TWO OBSERVATIONS ON THE PERTURBED WEDGE

FRED B. HOLT

ABSTRACT. Francisco Santos has described a new construction, perturbing apart a non-simple face, to offer a counterexample to the Hirsch Conjecture. We offer two observations about this *perturbed wedge* construction, regarding its effect on edge-paths. First, that an all-butsimple spindle of dimension d and length d + 1 is a counterexample to the nonrevisiting conjecture. Second, that there are conditions under which the perturbed wedge construction does not increase the diameter.

NOTE: These are simply working notes, offering two observations on the construction identified by Santos.

1. INTRODUCTION

Coincidence in high dimensions is a delicate issue. Santos has brought forward the perturbed wedge construction [San10], to produce a counterexample to the Hirsch conjecture. We start in dimension 5 with a non-simple counterexample to the nonrevisiting conjecture. If this counterexample were simple, then repeated wedging would produce a corresponding counterexample to the Hirsch conjecture. Since our counterexample to the nonrevisiting conjecture is not simple, we need an alternate method to produce the corresponding counterexample to the Hirsch conjecture, and the perturbed wedge provides this method.

A d-dimensional spindle (P, x, y) is a polytope with two distinguished vertices x and y such that every facet of P is incident to either x or y. The *length* of the spindle (P, x, y) is the distance $\delta_P(x, y)$. The spindle (P, x, y)is *all-but-simple* if every vertex of P other than x and y is a simple vertex.

Our first observation is that a *d*-dimensional all-but-simple spindle (P, x, y) of length d + 1 is a counterexample to the nonrevisiting conjecture.

Let P be a d-dimensional polytope with n facets. Let y be a nonsimple vertex of P incident to a facet G, and let F be a facet not incident to y. The *perturbed wedge* $\tilde{\omega}_{F,G}P$ of P is a wedge of P with foot F, which is a (d+1)-dimensional polytope with n+1 facets, followed by a perturbation of the image of the facet G in the coordinate for the new dimension.

Date: DRAFT 23 Feb 2013 - posted ArXiv 23 May 2013.

Our second observation is that for a spindle (P, x, y), with a facet F incident to x and a facet G incident to y, if there is a nonsimple edge in G from y to a vertex in F, then a short path from x to y along this edge is not increased in length in $\tilde{\omega}_{F,G}P$.

2. A Counterexample to the nonrevisiting conjecture

Lemma 2.1. Let (P, x, y) be a d-dimensional all-but-simple spindle of length d + 1. Then every path from x to y revisits at least one facet.

Proof. Let X be the n_1 facets incident to x, and let Y be the n_2 facets incident to y. Let $\rho = [x, u_1, \ldots, u_{k-1}, y]$ be a path from x to y of length k > d, with all of the u_i being simple vertices.

Then each u_i is incident to d facets. u_1 is incident to d-1 facets in X and one facet in Y, and u_k is incident to one facet in X and d-1 facets in Y. The incidence table for ρ looks like the following:

	X							Y					
x:	n_1 vertices	1	1	• • •	1	• • •	1						
$u_1:$	d vertices				1	•••	1	1					
						• • •			• • •				
$u_{k-1}:$	d vertices						1	1	• • •	1			
y:	n_2 vertices							1	•••	1	•••	1	1

Consider the facet-departures and facet-arrivals from u_1 to u_k (the simple part of the path). If any of the arrivals were back to a facet in X, this would be a revisit since these facets were all incident to the starting vertex x. So the arrivals must all be in Y.

Similarly, all of the departures must be from X. Any departure from a facet in Y would create a revisit since all the facets in Y are incident to the final vertex y.

There are too many arrivals and departures to prevent a revisit. The vertex u_1 is incident to only d-1 facets in X, and since each departure leaves a facet of X, u_j is incident to d-j facets in X. So u_d has completely departed from X. Since k > d, u_d occurs among the vertices u_1, \ldots, u_{k-1} , but since u_d is incident only to facets in Y, u_d must already be the vertex y, and ρ would have length at most d.

Santos [San10] has produced all-but-simple spindles in dimension 5 of length 6. Currently the smallest example has 25 facets. So in dimension 5 with 25 facets, we have a counterexample to the nonrevisiting conjecture, with length well below the Hirsch bound. Since the spindle is not simple, at x and y, the usual way of generating the corresponding counterexample [KW67, Hol03] to the Hirsch conjecture (through repeated wedging) does



FIGURE 1. This figure illustrates the Santos perturbed wedge construction and its dual construction. In the primal setting, we first perform a wedge of P over a facet F which is incident to x, followed by a vertical perturbation of a facet G incident to y.

not directly apply, and we need an alternate construction. The perturbed wedge accomplishes this.

3. The perturbed wedge

The perturbed wedge is constructed in two steps, first as a wedge over a facet, followed by a perturbation of a facet incident to a nonsimple vertex of the wedge.

Let P be a d-dimensional polytope with n facets and m vertices, and let F = F(u) be a facet incident to x in P. The wedge $W = \omega_F(P)$ is a (d+1)-dimensional polytope with n+1 facets and $2m - f_0(F)$ vertices. The wedge $\omega_F P$ over F in P, corresponds to the two-point suspension over u in P^* .

We now perturb a facet G = F(v) in W incident to the edge $[y^t, y_b]$. This is already interesting; we don't encounter non-simple edges until dimension 4. The perturbation of G is accomplished by introducing a small vertical

displacement to v; that is, instead of the last coordinate of the outwardpointing normal vector to G being 0, we perturb this coordinate to $\epsilon > 0$.

3.1. Embedded construction. As a canonical embedding for polytopes, we consider the vertices of P to be embedded in $1 \times \mathbb{R}^d$, with 0 in the interior of P. For the facets of P we take their outward-pointing normals. Since 0 is interior to P, we can assume that the first coordinate of each normal is -1. P is given by the embedding:

with

 $H_{n\times(d+1)}^T V_{(d+1)\times m} \le \langle 0 \rangle,$

$$\begin{bmatrix} -1 & h_i^T \end{bmatrix} \cdot \begin{bmatrix} 1 \\ x_j \end{bmatrix} = 0$$

iff vertex j is incident to facet (hyperplane) i. The facet-vertex incidence matrix for P is given by the $\{0, 1\}$ -matrix

$$M_{n \times m}(P) =$$
IsZero $(H^T V)$.

As a *d*-dimensional polytope, each vertex of P is incident to at least d facets, and each k-face of P is incident to at least k + 1 vertices. A simple vertex is incident to exactly d facets.

For a k-face of P, each incident facet contributes either to the affine space supporting this face or to its boundary [Hol04]. The space supporting this kface is the intersection of at least d-k facets of P, and we say that the space is *simple* iff this space is given by the coincident intersection of exactly d-kfacets of P. For k > 0, the boundary of the k-face is created by the various intersections of at least k + 1 other facets with the supporting space of the face. A face is simple iff its space is simple and all of its boundary elements are simple. A face can be nonsimple in a variety of ways or in multiple ways, through the nonsimplicity of its space or of its various boundary elements.

Let F be represented by the first outward-pointing normal in H^T , and G by the last outward-pointing normal. Then $H^T(\omega_F P)$ is given canonically by

$$H^{T}(\omega_{F}P) = \begin{bmatrix} -1 & h_{1}^{T} & 1 \\ -1 & h_{1}^{T} & -1 \\ -1 & h_{2}^{T} & 0 \\ \vdots & \vdots & \vdots \\ -1 & h_{n}^{T} & 0 \end{bmatrix}.$$

The facet F is replaced by two facets, the top and the base of the wedge. The top has final coordinate 1, and the base -1. Every other facet is replaced by a single *vertical* facet, meaning that the last coordinate (the new coordinate) is 0.

The vertices of $\omega_F P$ are given as follows. Rearrange the columns of V so that the vertices incident to F are given in the first block V_F and

the rest of the vertices occur in a second block V_{-} (denoted this way since $[-1 h_1^T] \cdot V_{-} < \langle 0 \rangle$).

$$\begin{split} V(P) &= \begin{bmatrix} V_F & V_- \end{bmatrix} .\\ \text{and} \\ V(\omega_F P) &= \begin{bmatrix} V_F & V_- & V_- \\ \langle 0 \rangle & -[-1 \ h_1^T] \cdot V_- & [-1 \ h_1^T] \cdot V_- \end{bmatrix} . \end{split}$$

In $V(\omega_F P)$, the vertices are now embedded in $1 \times \mathbb{R}^{d+1}$. The last coordinate for vertices in the foot F is 0, and for vertices not in the foot, there are two images, one in the top and one in the base.

Let the facet G have outward-pointing normal $[-1 h_n^T 0]$. We perturb the last coordinate to $\epsilon > 0$ to complete the construction of the perturbed wedge.

$$H^{T}(\tilde{\omega}_{F,G}P) = \begin{bmatrix} -1 & h_{1}^{T} & 1 \\ -1 & h_{1}^{T} & -1 \\ -1 & h_{2}^{T} & 0 \\ \vdots & \vdots & \vdots \\ -1 & h_{n-1}^{T} & 0 \\ -1 & h_{n}^{T} & \epsilon \end{bmatrix}.$$

Denote the outward-pointing normal for \tilde{G} by $h_{\tilde{G}}^T = [-1 \ h_n^T \ \epsilon]$.

To understand the effect of perturbing the facet G, we consider both Gand its perturbed image \tilde{G} . While we were able to write down the vertices of the wedge $\omega_F P$ explicitly, the effect of the perturbation is more complicated. The vertices incident to the facet G are of three types:

- Foot: $v \in G \cap T \cap B$ (or $v \in G \cap F$). For these vertices the last coordinate is 0, so $h_{\tilde{G}}^T v = 0$, and these vertices remain after the perturbation.
- Top: $v \in G \cap T \setminus B$. For these vertices, the last coordinate is positive, so $h_{\tilde{G}}^T v > 0$. If v consists combinatorially of a single edge terminated by G the case when v is a simple vertex but also when v consists of a single nonsimple edge terminated by G then the vertex v is perturbed back along this edge. If v consists combinatorially of more than one edge being terminated by G, then v is truncated away by the perturbation, and \tilde{G} introduces vertices along all of the edges incident to v but not lying in G.
- Base: $v \in G \cap B \setminus T$. For these vertices, the last coordinate is negative, so $h_{\tilde{G}}^T v < 0$. If v consists combinatorially of a single edge terminated by G the case when v is a simple vertex but also when v consists of a single nonsimple edge terminated by G then the vertex v is perturbed out along this edge. If v consists combinatorially of more than one edge being terminated by G, then v remains as a vertex

of $\tilde{\omega}_{F,G}P$, and the perturbation reveals new edges emanating from v and terminating in \tilde{G} in new vertices.

We now consider the effect of the perturbed wedge on vertex y and its natural images.

Under the wedge, y has two natural images y^t and y_b , in the top and base respectively. Since y is a nonsimple vertex, y^t and y_b are nonsimple vertices, and the edge $[y_b, y^t]$ between them is a nonsimple edge. The facet G is one of the facets supporting the space of this edge.

When G is perturbed to \tilde{G} , y_b is preserved as a vertex, y^t is truncated away, and \tilde{G} terminates the vertical edge at a new vertex y_0 whose last coordinate is 0. \tilde{G} introduces new vertices along the edges of $\omega_F P$ incident to y^t but not lying in G. \tilde{G} also introduces new vertices along the new edges emanating from y_b as revealed by \tilde{G} . That is, the collection of facets Y and the facet B intersect in edges that had lain beyond the facet G. \tilde{G} now introduces these edges as part of the boundary of $\tilde{\omega}_{F,G}P$ and terminates them in new vertices.

In considering the implications of the perturbed wedge construction on the Hirsch conjecture, we are interested in short paths from x to y in P and their tight natural images from x to y_0 in $\tilde{\omega}_{F,G}P$.

Claim: The perturbed wedge does not introduce revisits on tight natural images of short paths. The detailed study of this claim is beyond the scope of this note.

For the construction of the counterexample to the Hirsch conjecture, the revisits already exist in the initial 5-dimensional spindle. We see below that as a general construction, the perturbed wedge does not always increase the length of the input polytope by 1. However, the repeated application of the perturbed wedge to an all-but-simple spindle avoids the conditions of the following lemma.

Lemma 3.1. Let y be a nonsimple vertex of a d-dimensional polytope P. Let F be a facet of P not incident to y, and let G be a facet incident to y. If there is a nonsimple edge in G from y to a vertex w in $F \cap G$, then this edge remains after the perturbation.

Proof. As a nonsimple edge, the 1-dimensional space of [w, y] is defined by the coincidence of G and at least d-1 other facets \hat{Y} . See Figure 2. These facets $\hat{Y}G$ are incident to both w and y. The boundary of the edge at w is established by F and possibly more facets \hat{X} , none of which can be incident to y. The boundary of the edge at y is established by the facets $Y \setminus \hat{Y}$.

Under the wedge $\omega_F P$, the image of the edge is a nonsimple 2-dimensional face, the triangle with vertices w, y^t , and y_b .



FIGURE 2. Consider the facet incidences in the circumstance that y has a neighbor w along an edge of G that terminates in the facet F, which will be the foot of the wedge. Denote the facet incidences at the nonsimple vertex y as YG, in which Y is a set of at least d facets. The edge from y to wis the coincidence of $\hat{Y}G$ in which at least one facet of Y is omitted from \hat{Y} . The vertex w is coincident with the facets $\hat{Y}G$ and F and perhaps additional facets \hat{X} . \hat{X} may be empty.



FIGURE 3. Consider the action of the perturbed wedge on the facet-coincidences at y and w. Under the wedge, YGbecomes an edge and $\hat{Y}G$ becomes a 2-face. The vertex yhas two natural images, y^t which is incident to the facets YGT, and y_b which is incident to YGB. The facet F is replaced by two facets, the top T and the base B. Now we perturb the facet G, introducing a small positive value in the last coordinate of its outward normal. Since y was nonsimple, the vertex YB remains, but the vertex YT is truncated away. Instead, \tilde{G} now intersects the vertical edge Y in the plane $\{1\} \times \mathbb{R}^d \times \{0\}$. The 2-face $\hat{Y}G$ was nonsimple, and so the 2-face remains with its space supported by \hat{Y} , and \tilde{G} intersects it in an edge $[w, y_0]$.

Now, under the perturbation, the space of the triangular face is still defined by \hat{Y} . \tilde{G} intersects this face in the plane with last coordinate 0, creating an edge from w to y_0 .

In the dual setting, the wedge over the facet F(u) corresponds to a twopoint suspension $S_u(P^*)$ over the vertex u. See Figure 1. The perturbation of the facet G = F(v) corresponds to a vertical perturbation of the vertex v.

The observation in the previous lemma is that if the facet Y is adjacent to a facet W, such that $u, v \in W$ and that the ridge $Y \cap W$ is not simplicial, then after the two-point suspension over u and the perturbation of v, the new Y is still adjacent to W across the ridge, with v removed from the ridge.

Since the spindles used to seed the construction of the counterexample to the Hirsch conjecture are all-but-simple, the nonsimple edge does not occur on short paths between x and y. Although the natural images of xand y under this construction are not simple vertices, and although they are connected by nonsimple edges, the edges from y_0 run to natural images of y on the bases of various wedges, which should not be reused as the foot of the wedge. Thus the condition on $G \cap F$ does not occur for short paths.

4. Summary

Santos' construction of the first known counterexample to the Hirsch conjecture, for bounded polytopes, follows the strategy of first finding a counterexample to the nonrevisiting conjecture. Santos constructs a 5-dimensional all-but-simple spindle (P, x, y) of length 6, which is a counterexample to the nonrevisiting conjecture.

For simple polytopes, if we had a counterexample to the nonrevisiting conjecture, we would produce the corresponding counterexample to the Hirsch conjecture through repeated wedging, over all the facets not incident to x or y. However, Santos 5-dimensional spindle is not simple. Every facet is incident to either x or y, so we need an alternate method to produce the corresponding counterexample to the Hirsch conjecture. The perturbed wedge accomplishes this.

References

- [Hol03] F.B. Holt. Maximal nonrevisiting paths in simple polytopes. Discrete Mathematics, (1-3):105–128, 2003.
- [Hol04] F.B. Holt. Blending simple polytopes at faces. Discrete Mathematics, (1-3):141– 150, 2004.
- [San10] F. Santos. A counterexample to the Hirsch Conjecture. arXiv, 1006.2814v1:1–27, 2010.

5520 - 31st Ave NE; Seattle, WA 98105; fbholt@earthlink.net