

ON THE LEFT INVARIANT RANDERS AND MATSUMOTO METRICS OF BERWALD TYPE ON 3-DIMENSIONAL LIE GROUPS

H. R. SALIMI MOGHADDAM

ABSTRACT. In this paper we identify all simply connected 3-dimensional real Lie groups which admit Randers or Matsumoto metrics of Berwald type with a certain underlying left invariant Riemannian metric. Then we give their flag curvatures formulas explicitly.

1. Introduction

Let M be a smooth n -dimensional manifold and TM be its tangent bundle. A Finsler metric on M is a non-negative function $F : TM \rightarrow \mathbb{R}$ which has the following properties:

- (1) F is smooth on the slit tangent bundle $TM^0 := TM \setminus \{0\}$,
- (2) $F(x, \lambda y) = \lambda F(x, y)$ for any $x \in M$, $y \in T_x M$ and $\lambda > 0$,
- (3) the $n \times n$ Hessian matrix $[g_{ij}(x, y)] = [\frac{1}{2} \frac{\partial^2 F^2}{\partial y^i \partial y^j}]$ is positive definite at every point $(x, y) \in TM^0$.

An important family of Finsler metrics is the family of (α, β) -metrics. These metrics are introduced by M. Matsumoto (see [10]). Two families of interesting and important examples of (α, β) -metrics are Randers metrics $\alpha + \beta$, and Matsumoto metrics $\frac{\alpha^2}{\alpha - \beta}$, where $\alpha(x, y) = \sqrt{g_{ij}(x)y^i y^j}$ and $\beta(x, y) = b_i(x)y^i$ and g and β are a Riemannian metric and a 1-form respectively as follows:

$$(1.1) \quad g = g_{ij} dx^i \otimes dx^j$$

$$(1.2) \quad \beta = b_i dx^i.$$

These metrics occur naturally in physical applications. For example, in four-dimensional case, consider the above Randers metric as

$$(1.3) \quad F(x, \dot{x}) = (g_{ij}(x)\dot{x}^i \dot{x}^j)^{\frac{1}{2}} + b_i(x)\dot{x}^i,$$

where $\dot{x}^i = \frac{dx^i}{dq}$ and q is a parameter along a curve. Now suppose that $g_{ij}(x)$ is the (pseudo-)Riemannian metric tensor of space-time and

$$(1.4) \quad b_i(x) = \frac{e}{m_0 c^2} A_i(x),$$

where e and m_0 are the electric charge and the rest mass of a test particle respectively, c is the light velocity, and $A_i(x)$ denotes the electromagnetic vector potential. In this case the expression (1.3) is, up to a constant factor, the Lagrangian function of a test electric charge in the electromagnetic and gravitational fields described by the vector potential $A_i(x)$ and the Riemannian metric tensor $g_{ij}(x)$ (see [3]).

Key words and phrases. invariant metric, flag curvature, Berwald space, Randers metric, Matsumoto metric, 3-dimensional Lie group

AMS 2010 Mathematics Subject Classification: 22E60, 53C60.

Randers metrics were introduced by G. Randers in 1941 ([11]) when he was working on general relativity. On the other hand in the Matsumoto metric, the 1-form $\beta = b_i dx^i$ was originally to be induced by earths gravity (see [2] or [9].).

It has been shown that $\alpha + \beta$ (and in a similar way $\frac{\alpha^2}{\alpha - \beta}$) is a Finsler metric if and only if $\|\beta_x\|_\alpha = \sqrt{b_i(x)b^i(x)} < 1$, (for $\frac{\alpha^2}{\alpha - \beta}$, $\|\beta_x\|_\alpha = \sqrt{b_i(x)b^i(x)} < \frac{1}{2}$), where $b^i(x) = g^{ij}(x)b_j(x)$ and $[g^{ij}(x)]$ is the inverse matrix of $[g_{ij}(x)]$ (for more details see [1] and [5]).

The Riemannian metric g induces an inner product on any cotangent space T_x^*M such that $\langle dx^i(x), dx^j(x) \rangle = g^{ij}(x)$. The induced inner product on T_x^*M induces a linear isomorphism between T_x^*M and T_xM (see [6].). Then the 1-form β corresponds to a vector field \tilde{X} on M such that

$$(1.5) \quad g(y, \tilde{X}(x)) = \beta(x, y).$$

Therefore we can write the Randers and Matsumoto metrics as follows:

$$(1.6) \quad F(x, y) = \alpha(x, y) + g(\tilde{X}(x), y),$$

$$(1.7) \quad F(x, y) = \frac{\alpha(x, y)^2}{\alpha(x, y) - g(\tilde{X}(x), y)}.$$

One of geometric quantities which can be used in classification of Finsler metrics is flag curvature. This quantity is a generalization of the concept of sectional curvature in Riemannian geometry. Flag curvature is computed by the following formula:

$$(1.8) \quad K(P, Y) = \frac{g_Y(R(U, Y)Y, U)}{g_Y(Y, Y) \cdot g_Y(U, U) - g_Y^2(Y, U)},$$

where $g_Y(U, V) = \frac{1}{2} \frac{\partial^2}{\partial s \partial t} (F^2(Y + sU + tV))|_{s=t=0}$, $P = \text{span}\{U, Y\}$, $R(U, Y)Y = \nabla_U \nabla_Y Y - \nabla_Y \nabla_U Y - \nabla_{[U, Y]} Y$ and ∇ is the Chern connection induced by F (see [4] and [15].).

A Riemannian Metric g on the Lie group G is called left invariant if

$$(1.9) \quad g(x)(y, z) = g(e)(T_x l_{x^{-1}} y, T_x l_{x^{-1}} z) \quad \forall x \in G, \forall y, z \in T_x G,$$

where e is the unit element of G .

Similar to the Riemannian case, a Finsler metric is called left invariant if

$$(1.10) \quad F(x, y) = F(e, T_x l_{x^{-1}} y).$$

In our previous paper ([14]) we studied invariant Matsumoto metrics on homogeneous spaces and gave the flag curvature formula of them. In this paper we identify all simply connected 3-dimensional real Lie groups which admit Randers or Matsumoto metrics of Berwald type with a certain underlying left invariant Riemannian metric. Then we give their flag curvatures formulas explicitly.

2. Left invariant Randers and Matsumoto metrics of Berwald type on 3-dimensional Lie groups

The following proposition gives a method for constructing left invariant Randers or Matsumoto metrics on Lie groups.

Proposition 2.1. *Let \langle, \rangle be any left invariant Riemannian metric on a Lie group G . A function F of the form (1.6) or (1.7) is a left invariant Finsler metric (Randers or Matsumoto metric, respectively) if and only if the vector field \tilde{X} is a left invariant vector field and, $\|\tilde{X}(x)\|_\alpha = \sqrt{g(\tilde{X}(x), \tilde{X}(x))} < 1$ (for Randers metric) or $\|\tilde{X}(x)\|_\alpha < \frac{1}{2}$ (for Matsumoto metric), for any $x \in G$.*

Proof. With attention to proposition 3.6 of [13], it is clear. \square

Theorem 2.2. *Let G be a simply connected 3-dimensional real Lie group equipped with a left invariant Riemannian metric g and \mathfrak{g} denotes the Lie algebra of G . Then G admits a left invariant Randers or Matsumoto metrics of Berwald type with the underlying Riemannian metric g if and only if G , \mathfrak{g} and the associated matrix of g , with respect to the base $\{x, y, z\}$ of g and up to automorphism, be as follows:*

$$\text{i: } G = \mathbb{R}^3, [x, y] = 0, [x, z] = 0, [y, z] = 0, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$\text{ii: } G = \text{The solvable Lie group } \tilde{E}_0(2), [x, y] = 0, [x, z] = y, [y, z] = -x, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \nu \end{pmatrix},$$

where $\nu > 0$,

$$\text{iii: } G = \text{The non-unimodular Lie group } G_c, [x, y] = 0, [x, z] = -y, [y, z] = cx - 2y, \begin{pmatrix} 1 & \frac{1}{2} & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & 0 & \nu \end{pmatrix}, \text{ where } \nu > 0, c = 0.$$

Proof. We consider the classification of simply connected 3-dimensional real Lie groups given in [8]. Then, by some computations, we can compute the Levi-Civita connection of each case with respect to the base $\{x, y, z\}$ as table (1), where for case 15 we assumed that

$$(2.1) \quad A := \frac{\lambda^2(1 + \mu) + 1 - \mu}{2c^2\lambda^2},$$

$$(2.2) \quad B := \frac{1 - \mu}{2c\lambda^2},$$

$$(2.3) \quad D := \frac{1 - \mu}{2\lambda^2}.$$

We know that (see [4] and [2].) a Randers or Matsumoto metric is of Berwald type if and only if the vector field \tilde{X} (considered in (1.6) and (1.7)) is parallel with respect to the Levi-Civita connection of the Riemannian metric g and, $\|\tilde{X}\|_\alpha < 1$ (for Randers metric) or $\|\tilde{X}\|_\alpha < \frac{1}{2}$ (for Matsumoto metric). Therefore it is sufficient to find the cases which admit a parallel left invariant vector field \tilde{X} between these 15 cases. We can see in case 1 every left invariant vector field \tilde{X} is parallel with respect to the Levi-Civita connection of g , so for any left invariant vector field \tilde{X} with $\|\tilde{X}\|_\alpha < 1$ we have a Randers metric (and with $\|\tilde{X}\|_\alpha < \frac{1}{2}$ a Matsumoto metric). By a direct computation and using table (1) in case 5, G admits such left invariant vector field \tilde{X} if and only if $\mu = 1$. In this case the left invariant vector fields of the form $\tilde{X} = pz$, $p \in \mathbb{R}$, are parallel with respect to the Levi-Civita connection of g . Therefore, in Randers case, it is sufficient to consider $\|\tilde{X}\|_\alpha = \sqrt{g(pz, pz)} = |p|\sqrt{\nu} < 1$ or $|p| < \frac{1}{\sqrt{\nu}}$.

TABLE 1. The Levi-Civita connection of simply connected 3-dimensional real Lie groups

case	Lie algebra structure	Associated simply connected Lie group	Left Riemannian invariant metric	Conditions for parameters	$\nabla_x x$	$\nabla_x y$	$\nabla_x z$	$\nabla_y x$	$\nabla_y y$	$\nabla_y z$	$\nabla_z x$	$\nabla_z y$	$\nabla_z z$
case 1	$[x, y] = 0,$ $[x, z] = 0,$ $[y, z] = 0$	\mathbb{R}^3	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	-	0	0	0	0	0	0	0	0	0
case 2	$[x, y] = z,$ $[x, z] = 0,$ $[y, z] = 0$	The Heisenberg group Nil	$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\lambda > 0$	0	$\frac{1}{2}z$	$-\frac{1}{2\lambda}y$	$-\frac{1}{2}z$	0	$\frac{1}{2\lambda}x$	$-\frac{1}{2\lambda}y$	$\frac{1}{2\lambda}x$	0
case 3	$[x, y] = 0,$ $[x, z] = -x,$ $[y, z] = y$	The solvable Lie group Sol	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\nu > 0$	$\frac{z}{\nu}$	0	$-x$	0	$-\frac{z}{\nu}$	y	0	0	0
case 4	$[x, y] = 0,$ $[x, z] = -x,$ $[y, z] = y$	The solvable Lie group Sol	$\begin{pmatrix} 1 & 1 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\mu, \nu > 0$	$\frac{z}{\nu}$	0	$\frac{\mu}{1-\mu}x - \frac{\mu}{1-\mu}y$	0	$-\frac{\mu}{\nu}z$	$\frac{\mu}{1-\mu}x - \frac{\mu}{1-\mu}y$	$\frac{1}{1-\mu}x - \frac{\mu}{1-\mu}x$	$-\frac{\mu}{1-\mu}x - \frac{\mu}{1-\mu}y$	0
case 5	$[x, y] = 0,$ $[x, z] = y,$ $[y, z] = -x$	The solvable Lie group $E_0(2)$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$0 < \mu \leq 1, \nu > 0$	0	$\frac{1-\mu}{2\nu}z$	$\frac{\mu-1}{2\mu}y$	$\frac{1-\mu}{2\nu}z$	0	$\frac{\mu-1}{2}x$	$-\frac{1+\mu}{2\mu}y$	$\frac{1+\mu}{2}x$	0
case 6	$[x, y] = 2z,$ $[x, z] = -2y,$ $[y, z] = -2x$	The simple Lie group $\widetilde{PSL}(2, \mathbb{R})$	$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\mu \geq \nu > 0, \lambda > 0$	0	$\frac{\lambda+\mu+\nu}{\nu}z$	$-\frac{\lambda+\mu+\nu}{\mu}y$	$\frac{\lambda+\mu-\nu}{\nu}z$	0	$-\frac{\lambda-\mu+\nu}{\lambda}x$	$-\frac{\lambda+\mu-\nu}{\mu}y$	$\frac{\lambda-\mu+\nu}{\lambda}x$	0
case 7	$[x, y] = z,$ $[x, z] = -y,$ $[y, z] = x$	The simple Lie group $SU(2)$	$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\lambda \geq \mu \geq \nu > 0$	0	$-\frac{\lambda+\mu+\nu}{2\nu}z$	$\frac{\lambda-\mu-\nu}{2\mu}y$	$-\frac{\lambda+\mu-\nu}{2\nu}z$	0	$\frac{\lambda-\mu+\nu}{2\lambda}x$	$\frac{\lambda+\mu-\nu}{2\mu}y$	$-\frac{\lambda-\mu+\nu}{2\lambda}x$	0
case 8	$[x, y] = 0,$ $[x, z] = -x,$ $[y, z] = -y$	The non-unimodular Lie group G_I	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\nu > 0$	$\frac{z}{\nu}$	0	$-x$	0	$\frac{z}{\nu}$	$-y$	0	0	0
case 9	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$0 < \mu \leq c , \nu > 0$	0	$\frac{\mu-c}{2\nu}z$	$\frac{c-\mu}{2\mu}y$	$\frac{\mu-c}{2\nu}z$	$\frac{2\mu}{\nu}z$	$\frac{c-\mu}{2}x - 2y$	$\frac{c+\mu}{2\mu}y$	$-\frac{c+\mu}{2}x$	0
case 10	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\mu, \nu > 0, c = 0$	0	$\frac{\mu}{2\nu}z$	$-\frac{1}{2}y$	$\frac{\mu}{2\nu}z$	$\frac{2\mu}{\nu}z$	$-\frac{\mu}{2}x - 2y$	$\frac{1}{2}y$	$-\frac{\mu}{2}x$	0
case 11	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} 1 & \frac{1}{2} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\nu > 0, c = 0$	$\frac{z}{2\nu}$	$\frac{z}{\nu}$	$-y$	$\frac{z}{\nu}$	$\frac{2z}{\nu}$	$-2y$	0	0	0
case 12	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\nu > 0, c = \mu = 1, 0 < \mu \leq 1$	0	$\frac{\mu-1}{2\nu}z$	$\frac{1-\mu}{2\mu}y$	$\frac{\mu-1}{2\nu}z$	$\frac{2\mu}{\nu}z$	$\frac{1-\mu}{2}x - 2y$	$\frac{1+\mu}{2\mu}y$	$-\frac{1+\mu}{2}x$	0
case 13	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} \lambda & \lambda & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\nu > 0, c = 1, 0 < \mu \leq 1, 0 < \lambda < 1$	$\frac{\lambda}{\nu}z$	$\frac{\lambda}{\nu}z$	$-\frac{\lambda}{1+\lambda}(x+y)$	$\frac{\lambda}{\nu}z$	$\frac{2-\lambda}{\nu}z$	$\frac{\lambda}{1+\lambda}x - \frac{2+\lambda}{1+\lambda}y$	$-\frac{\lambda}{1+\lambda}x + \frac{1}{1+\lambda}y$	$-\frac{1}{1+\lambda}x + \frac{1}{1+\lambda}y$	0
case 14	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} 1 & 1 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$\nu > 0, c = 1, 1 < \mu \leq c$	$\frac{z}{\nu}$	$\frac{2+\mu-c}{2\nu}z$	$\frac{-2+\mu+c}{2(1-\mu)}x + \frac{\mu-c}{2(1-\mu)}y$	$\frac{2+\mu-c}{2\nu}z$	$\frac{2\mu-c}{\nu}z$	$\frac{-\mu(c+2)+2c+\mu^2}{2(1-\mu)}x + \frac{-2+3\mu-c}{2(1-\mu)}y$	$\frac{-2+\mu+c}{2(1-\mu)}x + \frac{2-\mu-c}{2(1-\mu)}y$	$\frac{\mu^2+\mu(c-2)}{2(1-\mu)}x + \frac{2-c-\mu}{2(1-\mu)}y$	0
case 15	$[x, y] = 0,$ $[x, z] = -y,$ $[y, z] = cx - 2y$	The non-unimodular Lie group G_c	$\begin{pmatrix} A & B & 0 \\ 0 & D & 0 \\ 0 & 0 & \nu \end{pmatrix}$	$0 \leq \mu < 1, \nu > 0, 0, \lambda = \sqrt{1-c}$	$\frac{1-\mu}{2c(1-c)\nu}z$	$\frac{\mu-c}{2c(c-1)\nu}z$	$\frac{-\mu}{1+\mu}x + \frac{c+c\mu-2\mu}{c(\mu^2-1)}y$	$\frac{\mu-c}{2c(c-1)\nu}z$	$\frac{\mu-1}{2(c-1)\nu}z$	$\frac{c}{1+\mu}x - \frac{2+\mu}{1+\mu}y$	$-\frac{\mu}{1+\mu}x + \frac{(-2+c+c\mu)\mu}{c(\mu^2-1)}y$	$-\frac{c\mu}{1+\mu}x + \frac{\mu}{1+\mu}y$	0

In a similar way, in Matsumoto case, it is sufficient to consider $\|\tilde{X}\|_\alpha < \frac{1}{2}$ or $|p| < \frac{1}{2\sqrt{\nu}}$. In case 11 the Lie group G admits a family of left invariant vector fields of the form $\tilde{X} = -2px + py$, $p \in \mathbb{R}$, which are parallel with respect to the Levi-Civita connection of g . Therefore, in this case, for constructing Randers metrics of Berwald type it is sufficient to let $\|\tilde{X}\|_\alpha = \sqrt{g(-2px + py, -2px + py)} = \sqrt{3}|p| < 1$ or $|p| < \frac{\sqrt{3}}{3}$ and for constructing Matsumoto metrics of Berwald type it is sufficient to consider $|p| < \frac{\sqrt{3}}{6}$. By using table (1) and a little complicated computation we can see in all other cases the Lie group G does not admit a left invariant vector field \tilde{X} such that it is parallel with respect to the Levi-Civita connection of g . Therefore proposition (2.1) guarantees that in other cases we have not any left invariant Randers or Matsumoto metric of Berwald type with the underlying left invariant Riemannian metric g . \square

Now we discuss about the flag curvatures of these metrics.

Remark 2.3. As I have shown in theorems 3.5 and 3.8 of [12] for Randers metrics (and with a similar way for Matsumoto metrics), the Finsler (Randers and Matsumoto) metrics considered in cases (i) and (ii) of theorem (2.2) are flat geodesically complete locally Minkowskian Finsler metrics.

Therefore it is sufficient to compute the flag curvature of Randers and Matsumoto metrics of case (iii).

Theorem 2.4. *Let $F = \alpha + \beta$ be the left invariant Randers metric derived from case (iii) of theorem (2.2). Suppose that $\{U = ax + by + cz, V = \tilde{a}x + \tilde{b}y + \tilde{c}z\}$ is an orthonormal set with respect to g . Then the flag curvature $K(P, U)$ of the flag $(P = \text{span}\{U, V\}, U)$ is given by*

$$(2.4) \quad K(P, U) = -4 \left(\frac{c(\tilde{a} + 2\tilde{b}) - \tilde{c}(a + 2b)}{3pa - 2} \right)^2.$$

Therefore F is of non-positive flag curvature.

Proof. F is of Berwald type so the curvature tensors of g and F coincide. By using table (1) we can compute the curvature tensor as follows

$$(2.5) \quad \begin{aligned} R(x, z)x &= -R(z, x)x = \frac{1}{2}R(x, z)y = -\frac{1}{2}R(z, x)y = \frac{1}{2}R(y, z)x = -\frac{1}{2}R(z, y)x \\ &= \frac{1}{4}R(y, z)y = -\frac{1}{4}R(z, y)y = \frac{z}{\nu}, \\ -\frac{1}{2}R(x, z)z &= \frac{1}{2}R(z, x)z = -\frac{1}{4}R(y, z)z = \frac{1}{4}R(z, y)z = y, \end{aligned}$$

$R = 0$, for other cases. Therefore we have

$$(2.6) \quad R(V, U)U = \frac{1}{\nu} \{ (a + 2b)(\tilde{a}c - \tilde{c}a + 2(\tilde{b}c - \tilde{c}b)) \} z - 2c \{ \tilde{a}c - \tilde{c}a + 2(\tilde{b}c - \tilde{c}b) \} y.$$

Let $\delta := \frac{1}{\nu}\{(a+2b)(\tilde{a}c - \tilde{c}a + 2(\tilde{b}c - \tilde{c}b))\}$ and $\sigma := -2c\{\tilde{a}c - \tilde{c}a + 2(\tilde{b}c - \tilde{c}b)\}$. Now by using formula of g_U given in [7] we have

$$(2.7) \quad g_U(R(V, U)U, V) = (\delta\tilde{c}\nu + \sigma(\tilde{b} + \frac{\tilde{a}}{2}))(-\frac{3}{2}pa + 1) - \frac{3}{2}p\tilde{a}(\delta c\nu + \sigma(b + \frac{a}{2}))$$

$$(2.8) \quad g_U(U, U) = (-\frac{3}{2}pa + 1)^2$$

$$(2.9) \quad g_U(V, V) = 1 - \frac{3}{2}pa + (-\frac{3}{2}p\tilde{a})^2$$

$$(2.10) \quad g_U(U, V) = -\frac{3}{2}p\tilde{a}(-\frac{3}{2}pa + 1).$$

Substitution of the last equations in (1.8) will completes the proof. \square

Theorem 2.5. *Suppose that $F = \frac{\alpha^2}{\alpha-\beta}$ is the left invariant Matsumoto metric derived from case (iii) of theorem (2.2). Then, with the assumptions of the previous theorem, the formula for the flag curvature of F is as follows*

$$(2.11) \quad K(P, U) = -\frac{(2+3ap)^3(1+3ap)(\tilde{c}a - \tilde{a}c + 2(\tilde{c}b - \tilde{b}c))^2}{2(4+18a^2p^2+18ap-27\tilde{a}^2p^2)}.$$

Proof. The proof is similar to the previous theorem except that for considered Matsumoto metric by a direct computation we have

$$(2.12) \quad g_U(R(V, U)U, V) = \frac{1}{(1+\frac{3}{2}ap)^4}\{(\sigma(\frac{\tilde{a}}{2} + \tilde{b}) + \tilde{c}\delta\nu)(1+3ap)(1+\frac{3}{2}ap) - \frac{3}{2}\tilde{a}p(\sigma(\frac{a}{2} + b) + c\delta\nu)(1+6ap)\}$$

$$(2.13) \quad g_U(U, U) = \frac{1}{(1+\frac{3}{2}ap)^2}$$

$$(2.14) \quad g_U(V, V) = \frac{1+\frac{9}{2}(a^2p^2 - \tilde{a}^2p^2 + ap)}{(1+\frac{3}{2}ap)^4}$$

$$(2.15) \quad g_U(U, V) = \frac{-3\tilde{a}p}{2(1+\frac{3}{2}ap)^3}.$$

\square

REFERENCES

1. H. An and S. Deng, *Invariant (α, β) -Metrics on Homogeneous Manifolds*, Monatsh. Math. **154** (2008), 89–102.
2. P. L. Antonelli, R. S. Ingarden, M. Matsumoto, *The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology*, Kluwer Academic Publishers, (1993).
3. G. S. Asanov, *Finsler Geometry, Relativity and Gauge Theories*, D. Reidel Publishing Company, Dordrecht, Holland, (1985).
4. D. Bao, S. S. Chern and Z. Shen, *An Introduction to Riemann-Finsler Geometry*, (Berlin: Springer) (2000).
5. S. S. Chern and Z. Shen, *Riemann-Finsler Geometry*, (World Scientific) (2005).
6. S. Deng, Z. Hou, *Invariant Randers Metrics on Homogeneous Riemannian Manifolds*, J. Phys. A: Math. Gen. **37** (2004), 4353–4360.
7. E. Esrafilian and H. R. Salimi Moghaddam, *Flag Curvature of Invariant Randers Metrics on Homogeneous Manifolds*, J. Phys. A: Math. Gen. **39** (2006) 3319–3324.

8. K. Y. Ha and J. B. Lee, *Left invariant metrics and curvatures on simply connected threedimensional Lie groups*, Math. Nachr. **282**(6), (2009), 868-898.
9. M. Matsumoto, *A slope of a mountain is a Finsler surface with respect to time measure*, J. Math. Kyoto Univ. **29** (1989), 17-25.
10. M. Matsumoto, *Theory of Finsler spaces with (α, β) -metric*, Rep. Math. Phys. **31** (1992), 43-83.
11. G. Randers, *On an Asymmetrical Metric in the Four-Space of General Relativity*, Phys. Rev. **59**(1941), 195-199.
12. H. R. Salimi Moghaddam, *Some Berwald Spaces of Non-positive flag Curvature*, Journal of Geometry and Physics **59** (2009) 969-975.
13. H. R. Salimi Moghaddam, *The Relation Between Automorphism Group and Isometry Group of Randers Lie Groups*, Results in Mathematics **61**, Issue 1 (2012), Page 137-142, DOI 10.1007/s00025-010-0081-x.
14. H. R. Salimi Moghaddam, *Invariant Matsumoto metrics on homogeneous spaces*, Osaka Journal of Mathematics, to appear.
15. Z. Shen, *Lectures on Finsler Geometry*, (World Scientific) (2001).

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF ISFAHAN, ISFAHAN, 81746-73441-IRAN.

E-mail address: salimi.moghaddam@gmail.com and hr.salimi@sci.ui.ac.ir