RIEMANN–SCHWARZ REFLECTION PRINCIPLE AND ASYMPTOTICS OF MODULES OF RECTANGULAR FRAMES

S.R. Nasyrov

23 of May, 2013

Abstract

We investigate asymptotical behavior of the conformal module of a doubly-connected domain which is the difference of two homothetic rectangles under stretching it along the abscissa axis. Thereby, we give the answer to a question put by Prof. M. Vuorinen.

1 Introduction

In recent years an interest increases to investigation of conformal modules of plane quadrilaterals, doubly-connected domains and capacities of condensers with polygonal boundaries. Special consideration is given to studying of behavior of the modules under various deformations of domains, their numerical calculation, and asymptotics at degeneration. In this regard we can note the review by R. Kühnau [1] and the papers [2]–[5].

At first, we recall some definitions. Consider a plane doubly-connected domain D with nondegenerated boundary components. One of its important characteristics is the conformal module m(D). There are several equivalent definitions of m(D); we give some of them.

If D is conformally equivalent to an annulus $\{r_1 < |z| < r_2\}$, then

$$m(D) := \frac{1}{2\pi} \ln \frac{r_2}{r_1}.$$

²⁰¹⁰ AMS Subject Classification. Primary 30C20.

KEY WORDS AND PHRASES. Conformal module, doubly-connected plane domain, polygon, quadrilateral, reflection principle, elliptic integrals, anharmonic ratio, convergence to a kernel, prime ends, asymptotics.

The work is supported by the grants No. 12-01-97015 and No. 11-01-00762 of the Russian Fund of Basic Research.

On the other hand,

$$m(D) := \lambda(\Gamma),$$

where $\lambda(\Gamma)$ is the extremal length of curve-family Γ consisting of all curves joining in D its boundary components. Besides,

$$m(D) := 1/\lambda(\Gamma'),$$

 Γ' being the family of all curves in D separating its boundary components. At last,

$$m(D) := 1/\operatorname{Cap}(C),$$

where $\operatorname{Cap}(C)$ is the conformal capacity of the condenser C defined by D.

Significant property of module is its invariance under conformal mappings. It is quasiinvariant under quasiconformal maps (see, e. g., [8]): if f is an Hquasiconformal mapping of D onto \widetilde{D} , then

$$\frac{1}{H}m(D) \le m(\widetilde{D}) \le Hm(D).$$

One of the simplest H-quasiconformal mappings is the stretching along the abscissa axis $f_H : x+iy \mapsto Hx+iy$. M. Vuorinen states the following problem: to investigate how the module m(D) is deformed under f_H for sufficiently large H. In particular, which is asymptotical behavior of m(D) if D is the difference of two homothetic squares?

The main result of the paper is

Theorem 1.1 If
$$D_1 = D_1^{\sigma} := [-1,1]^2 \setminus [-\sigma,\sigma]^2, \ \sigma \in (0,1), \ D_H = D_H^{\sigma} := f_H(D_1), \ then$$

$$m(D_H^{\sigma}) \sim \frac{1-\sigma}{4\sigma H}, \quad H \to \infty.$$
 (1)

In Section 2 we give a solution to the problem for $\sigma = 1/2$, in addition, we deduce an explicit formula for $m(D_H)$ via elliptic integrals. It should be noted that when H = 1 an explicit formula for $m(D_H)$ is well-known, see Remark 1.3 below. In Section 4 the general case is considered. The results of Sections 2 and 4 were announced in [7] and [9]. Preliminarily, in Section 3 we establish continuity of module of quadrilateral under kernel convergence in the sense of Carathéodory.

To prove our results we need to recall some more definitions and facts. A *quadrilateral* is a simply-connected domain D, conformally equivalent to a disk, with four marked distinct points (prime ends) A_k , $1 \le k \le 4$, on the boundary of D; increasing of k corresponds to the order in which the points occur when we bypass ∂D in the positive direction. We denote the quadrilateral as $D(A_1, A_2, A_3, A_4)$ or simply D if it is clear which points A_k are fixed. The parts of ∂D lying between A_1 and A_2 , A_3 and A_4 we call *horizontal sides* of D, the other two parts of the boundary are *vertical sides*. Let us map conformally

 $D(A_1, A_2, A_3, A_4)$ onto a rectangle $[0, a] \times [0, b]$ so that the horizontal sides are mapped onto the horizontal sides of the rectangular. The number

$$m(D) := \frac{a}{b}$$

is called the *module* of D. It is known (see, e. g., [8]) that m(D) is equal to the extremal length $\lambda(\Gamma)$ of the family Γ consisting of curves in D joining its vertical sides; therefore, it is invariant under conformal mappings and quasiinvariant under quasiconformal ones. Besides, $m(D) = 1/\lambda(\Gamma')$ where Γ' is the family of curves in D joining its horizontal sides.

Let $E := \{|z| < 1\}, U := \{\Im z > 0\}, E^+ := E \cap U\}, S_{\gamma\delta} := \{e^{i\varphi} \mid \gamma < \varphi < \delta\}, 0 < \delta - \gamma < 2\pi$. We denote by [a, b] the segment with endpoints $a, b \in \mathbb{C}$.

The elliptic integral of the first kind

$$K(r) := \int_0^1 \frac{d\xi}{\sqrt{(1-\xi^2)(1-r^2\xi^2)}} \, .$$

It is known (see, e. g., [10], [11]) that

$$\lim_{r \to 0} \left(K(r') - \ln \frac{4}{r} \right) = 0,$$
(2)

where as usual $r' = \sqrt{1 - r^2}$. From (2) it follows that $K(r) \sim \ln \frac{4}{r'}$ as $r \to 1$. Therefore,

$$K(r) \sim \frac{1}{2} \ln \frac{1}{1-r}, \quad \frac{K(r)}{K(r')} \sim \frac{1}{\pi} \ln \frac{1}{1-r}, \quad r \to 1.$$
 (3)

From (2) we also obtain that

$$\frac{K(r')}{K(r)} \sim \frac{2}{\pi} \ln \frac{1}{r}, \quad r \to 0.$$
(4)

Remark 1.2 The ring domain consisting of the unit disk minus a radial slit from 0 to r, 0 < r < 1, is usually called the Grötzsch ring and its modulus is often denoted

$$\mu(r) = \frac{\pi}{2} \frac{K(r')}{K(r)} \,,$$

see [1]. The asymptotic formula (4) can be refined by use of the results from [10], Theorem 5.13.

Remark 1.3 When H = 1 an explicit formula for $m(D_H^{\sigma})$ is well-known (see, e. g., [6]):

$$m(D_1^{\sigma}) = \mu\left(\left(\frac{l-l'}{l+l'}\right)^2\right), \quad l = \mu^{-1}\left(\frac{2}{\pi}\frac{1-\sigma}{1+\sigma}\right), \quad l' = \sqrt{1-l^2}.$$

2 The case $\sigma = 1/2$

Consider the part Q_H of D_H lying in the first quarter of the plane. It is the union of three rectangles of the same size. Let us map conformally one of the rectangles with vertices at the points (H + i)/2, H + i/2, H/2 + i, and H + i onto the quarter of the unit disk $U_1 := \{z \mid |z| < 1, \Re z > 0, \Im z > 0\}$ by the mapping f so that f((H + i)/2) = 0, f(H + i/2) = 1, and f(H/2 + i) = i. Let $e^{i\kappa} = f(H + i)$.

By the Riemann-Schwarz reflection principle f could be extended up to the conformal mapping of the rectangle $[0, H] \times [0, 1]$ onto the unit disk E; we will designate the extension also through f. Then f maps conformally Q_H onto the domain which is three quarters of the unit disk, besides, f(H/2) = -i, $f(H) = e^{-i\kappa}$, f(i/2) = -1, and $f(i) = -e^{-i\kappa}$.



Fig. 1

The function $g(\zeta) := (if(\zeta))^{2/3}$ maps conformally Q_H onto E^+ at an appropriate choice of a branch of the power function. We have g(H/2) = 1, $g(H) = e^{i\alpha}$, f(i/2) = -1, $f(i) = -e^{-i\beta}$ where

$$\alpha = (\pi - 2\kappa)/3, \quad \beta = \pi - 2\kappa/3.$$
 (5)

The module of D_H , by the symmetry principle for quasiconformal mappings (see, e. g., [8]) is equal

$$m(D_H) = 1/(4\lambda(\Gamma)) \tag{6}$$

where Γ is the family of all curves in Q_H which join [H/2, H] and [i/2, i]. Because of conformal invariance of the module we obtain

$$\lambda(\Gamma) = \lambda(\Gamma'),\tag{7}$$

where Γ' is a family of all curves in E^+ connecting $S_{0\alpha}$ to $S_{\beta\pi}$. By the symmetry principle,

$$\lambda(\Gamma') = 2\lambda(\Gamma) \tag{8}$$

where $\widetilde{\Gamma}$ is the family of all curves which join $S_{-\alpha,\alpha}$ and $S_{\beta,2\pi-\beta}$ in E^+ .

Let us map conformally the unit disk E onto the upper half-plane U so that the points $e^{i\beta}$, $e^{-i\beta}$, $e^{-i\alpha}$, and $e^{i\alpha}$ are mapped on -1/l, -1, 1, and 1/l, l > 1. Now we express l through α and β . Equating the anharmonic ratios

$$\frac{-1/l+1}{-1/l-1}\cdot\frac{1/l-1}{1/l+1} = \frac{e^{i\beta}-e^{-i\beta}}{e^{i\beta}-e^{-i\alpha}}\cdot\frac{e^{i\alpha}-e^{-i\alpha}}{e^{i\alpha}-e^{-i\beta}}$$

we obtain

$$l = \frac{\sqrt{1 - \cos(\alpha + \beta)} - \sqrt{2\sin\beta\sin\alpha}}{\sqrt{1 - \cos(\alpha + \beta)} + \sqrt{2\sin\beta\sin\alpha}}.$$
(9)

Besides,

$$\lambda(\widetilde{\Gamma}) = \frac{2K(l)}{K(l')}.$$
(10)

From (6), (7), (8), and (10) we have

$$m(D_H) = \frac{K(l')}{16K(l)}.$$
 (11)

Now we find the relation between κ and H. For this purpose we map conformally E onto the upper half-plane U by a function φ so that $-e^{-i\kappa}$, $-e^{i\kappa}$, $e^{-i\kappa}$, and $e^{i\kappa}$ are mapped on -1/k, -1, 1, and 1/k. We note that k satisfies the condition

$$\frac{2K(k)}{K(k')} = H \tag{12}$$

because the quadrilateral, which is the upper half-plane U with fixed points -1/k, -1, 1, and 1/k, is conformally equivalent to the rectangle of length H and height 1 under the mapping $\varphi \circ f$. From the equality of cross-ratios

$$\frac{-1/k+1}{-1/k-1} \cdot \frac{1/k-1}{1/k+1} = \frac{-e^{-i\kappa} + e^{i\kappa}}{-e^{-i\kappa} - e^{i\kappa}} \cdot \frac{e^{i\kappa} - e^{-i\kappa}}{e^{i\kappa} - e^{-i\kappa}}$$

we have

$$\kappa = \arcsin \frac{1-k}{1+k} \,. \tag{13}$$

Therefore, we prove

Theorem 2.1 For $\sigma = 1/2$ the module of $D_H = D_H^{\sigma}$ is defined by (11) where *l* is found from (9) taking into account (5), (12), and (13).

Corollary 2.2 We have

$$m(D_H) \sim \frac{1}{4H}, \quad H \to \infty.$$

Actually, from (13) it follows that $\kappa \sim (1-k)/2$ as $H \to \infty$. Now, taking into account (9) and (5), we obtain

$$1 - l \simeq \sqrt{\sin \beta} = \sqrt{\sin \frac{2\kappa}{3}} \simeq \sqrt{1 - k}.$$

Thus, using (11), (12), and (3), we have

$$m(D_H) = \frac{K(l')}{16K(l)} \sim \frac{\pi}{16\ln[(1-l)^{-1}]} \sim \frac{\pi}{8\ln[(1-k)^{-1}]} \sim \frac{1}{4H}, \quad H \to \infty.$$

3 Convergence of domains and their modules

At first, we recall some results of the theory of prime ends of a sequence of domains converging to a kernel [12].

Consider a sequence of simply-connected domains G_n on the Riemann sphere $\overline{\mathbb{C}}$ converging to a kernel G with respect to a fixed point $S_0 \in \overline{\mathbb{C}}$. We assume that the boundaries of G_n and G are nondegenerate, i. e., each of them contains more than one point. Further we provide G_n and G with metrics induced from the sphere; in the case when all G_n are contained in a fixed Euclidean disk it is possible to change the spherical metrics by the Euclidean one.

Consider a section γ of G, i. e., a Jordan arc in G with endpoints on ∂G . Let γ_n be a section of G_n . We say that the sequence (γ_n) is a section of the sequence $\widetilde{G} := (G_n)$ lying over γ if the following conditions are fulfilled:

- 1. For any neighborhood U of γ there exists n_0 such that γ_n lies in U for any $n \ge n_0$;
- 2. if points $p_1, p_2 \in G$ are separated by γ in D, then there exists n_1 such that p_1 and p_2 are separated by γ_n in G_n for any $n \ge n_1$.
- In [12] it is proved

Lemma 3.1 For any section γ of G there exists a section $\tilde{\gamma} := (\gamma_n)$ of \tilde{G} lying over γ .

Let (γ_m) be a chain of sections of G which defines a prime end P of G, and $\tilde{\gamma}_m := (\gamma_{mn})$ is a section of \tilde{G} lying over γ_m . It is possible to introduce a natural relation of equivalence on the set of all such sequences $(\tilde{\gamma}_m)$ (in more detail see [12]); the classes of equivalence \tilde{P} of such sequences are called *prime ends* of \tilde{G} .

If a sequence of sections (γ_m) defines a prime end P of G and prime end P of \widetilde{G} contains $(\widetilde{\gamma}_m)$ where $\widetilde{\gamma}_m$ is a section of \widetilde{G} lying over γ_m , then we will say that \widetilde{P} is the prime end of \widetilde{G} corresponding to prime end P of G. The described correspondence $\Phi : P \mapsto \widetilde{P}$ is a bijection between the set of prime ends of \widetilde{G} and the set of prime ends of \widetilde{G} .

Now consider a sequence (P_n) where P_n is a point or a boundary prime end of G_n . Let $\tilde{P} = \Phi(P)$ be the prime end of \tilde{G} corresponding to a prime end P of G and P is defined by a sequence (γ_m) . Let \tilde{P} is defined by $(\tilde{\gamma}_m)$, where $\tilde{\gamma}_m$ lies over γ_m for any m, and $\tilde{\gamma}_m := (\gamma_{mn})$.

We will say that the sequence (P_n) converges to \widetilde{P} if for any m there exists n_0 such that γ_{mn} separates P_n from S_0 in G_m for any $n \ge n_0$.

Let f_n and f be conformal mappings of E onto G_n and G, extended to \overline{E} up to homeomorphisms of prime ends. Let P_n be a sequence consisting of points or boundary prime ends of G_n , and let P be a boundary prime end of G. Denote $\zeta_n = f_n^{-1}(P_n), \, \zeta_0 = f^{-1}(P).$

Theorem 6, p. 75, from [12] implies

Theorem 3.2 A sequence (P_n) converges to the prime end \widetilde{P} of \widetilde{G} corresponding to P if and only if $\zeta_n \to \zeta_0$.

From Theorem 3.2 we deduce the following result on continuity of the module of quadrilateral under kernel convergence of domains.

Theorem 3.3 Let A_n , B_n , C_n , and D_n be distinct boundary prime ends of G_n . Let the sequences (A_n) , (B_n) , (C_n) , and (D_n) converge to the prime ends \widetilde{A} , \widetilde{B} , \widetilde{C} and \widetilde{D} of \widetilde{G} corresponding to distinct prime ends A, B, C, and D of \widetilde{G} . Then $m(G_n(A_n, B_n, C_n, D_n)) \to m(G(A, B, C, D))$.

Actually, taking into account conformal invariance of the module of quadrilateral, by Theorem 3.2 we have

$$m(G_n(A_n, B_n, C_n, D_n)) = m(E(a_n, b_n, c_n, d_n)) \rightarrow$$
$$\rightarrow E(a, b, c, d) = m(G(A, B, C, D))$$

where a_n , b_n , c_n , and d_n are preimages of A_n , B_n , C_n , and D_n under the map f_n , and a, b, c, and d are preimages of A, B, C, and D under the map f.

Remark 3.4 The statement of Theorem 3.3 is valid not only for univalent domains, it is true for p-valent domains (Riemann surfaces) as well (see, e. g., [13]).

4 General case

As in the case $\sigma = 1/2$, consider the part Q_H of D_H lying in the first quarter of the plane. Let us shift Q_H to the left on the value σH ; as a result we receive the domain \tilde{Q}_H . By the symmetry principle, the module of D_H is equal to

$$m(D_H) = \frac{1}{4\lambda(\Gamma)} \tag{14}$$

where Γ is the family of curves joining $[0, (1 - \sigma)H]$ and $[-\sigma H + i\sigma, -\sigma H + i]$ in the quadrilateral \widetilde{Q}_{H} . Now consider the domain

$$\widetilde{Q} := \bigcup_{H>0} \widetilde{Q}_H = (\mathbb{R} \times (0,1)) \setminus ((-\infty,0] \times (0,\sigma]).$$

Let us map conformally \widetilde{Q} onto the horizontal strip $\mathbb{R} \times (0,1)$ with keeping the infinite prime ends and the origin. For this purpose we map conformally the upper half-plane U onto \widetilde{Q} and $G = \{0 < \Im \omega < 1\}$ by the functions

$$z = C \int_{1}^{\zeta} \sqrt{\frac{\zeta - s}{\zeta - 1}} \frac{d\zeta}{\zeta}, \quad \omega = \frac{\ln \zeta}{\pi},$$

where C > 0, 0 < s < 1.

Near $\zeta = 0$ we have

$$z = C\sqrt{s}\ln\zeta + \sum_{k=0}^{\infty} \sigma_k \zeta^k.$$

Since the intersection of the domain \widetilde{Q} and the left half-plane is a half-strip of width $(1-\sigma)/\pi$, we have $C\sqrt{s} = (1-\sigma)/\pi$. Near $\zeta = \infty$

$$z = C \ln \zeta + \sum_{k=0}^{\infty} \frac{\beta_k}{\zeta^k},$$

therefore, similarly we obtain $C = 1/\pi$. Then $s = (1 - \sigma)^2$ and

$$z = \frac{1}{\pi} \int_{1}^{\zeta} \sqrt{\frac{\zeta - (1 - \sigma)^2}{\zeta - 1}}.$$

Considering it we conclude that for sufficiently large M > 0 we have

$$z = (1 - \sigma)\omega + \sum_{k=0}^{\infty} \sigma_k e^{k\pi\omega}.$$
(15)

in the half-plane $\Pi^{-}_{\mu} := \{ \Re \omega < -M, 0 < \Im \omega < 1 \}$, and

$$z = \omega + \sum_{k=0}^{\infty} \beta_k e^{-k\pi\omega}.$$
 (16)

in the half-plane $\Pi_{\mu}^{+} := \{ \Re \omega > M, 0 < \Im \omega < 1 \}.$ From rectilinearity of the boundary arcs of the domains and the Riemann-Schwarz reflection principle it follows that convergence of the series (15) and (16) is uniform in the closed half-planes $\overline{\Pi_{\mu}^{-}}$ and $\overline{\Pi_{\mu}^{+}}$.

From (15) and (16) we deduce that on the vertical segments in \widetilde{Q} , lying on the lines $\Re \omega = -\widetilde{\sigma} H$, where

$$\widetilde{\sigma} = \frac{\sigma}{1 - \sigma}$$

we have

$$\Re z(\omega) = -\widetilde{\sigma} H + O(1), \quad H \to \infty$$

In the same way, on the segments, lying on the lines $\Re \omega = (1 - \sigma)H$,

$$\Re z = (1 - \sigma)H + O(1), \quad H \to \infty,$$

Therefore,

$$\lambda(\Gamma) \sim m(\tilde{P}) \tag{17}$$

where \widetilde{P} is the quadrilateral which is the rectangle

$$P := [-\widetilde{\sigma} H, (1 - \sigma)H] \times [0, 1]$$

with the segments $[-\tilde{\sigma}H, -\tilde{\sigma}H+i]$ and $[0, (1-\sigma)H]$ as vertical sides.

Let us map U onto P by the function

$$z = C \int_0^{\zeta} \frac{d\xi}{\sqrt{(1-\xi^2)(1-k^2\xi^2)}} + C_1$$

where

$$C_1 = \frac{(1-\sigma)^2 - \sigma}{1-\sigma} H, \quad C = \frac{(1-\sigma)^2 + \sigma}{2K(k)(1-\sigma)},$$

and $k \in (0, 1)$ is defined from the relation

$$\frac{2K(k)}{K(k')} = \frac{(1-\sigma)^2 + \sigma}{1-\sigma} H.$$
 (18)

The mapping takes the points $-\tilde{\sigma} H + i, \tilde{\sigma} H, 0$, and $(1 - \sigma)H$, i. e., the vertices of quadrilateral P, into (-1/k), (-1), a, and 1, where

$$a = \operatorname{sn}\left[\frac{\sigma - (1 - \sigma)^2}{\sigma + (1 - \sigma)^2} K(k), k\right].$$
(19)

Here $\operatorname{sn}[\cdot, k]$ is the Jacobi elliptic sine corresponding to the parameter k (see, e. g., [11]).

Now we map the upper half-plane U onto itself conformally so that the points -1/k, -1, a, and 1 move into $-1/\nu$, -1, 1, and $1/\nu$ (0 < ν < 1). Then the module of the quadrilateral \tilde{P} is equal to

$$m(\tilde{P}) = \frac{2K(\nu)}{K(\nu')} \tag{20}$$

where ν is defined from the equality of the cross-ratios:

$$\frac{1/\nu - 1}{1/\nu + 1} \cdot \frac{-1/\nu + 1}{-1/\nu - 1} = \frac{1 - a}{1 + 1} \cdot \frac{-1/k + 1}{-1/k - a}$$
$$\frac{1 - \nu}{1 + 1} = \sqrt{\frac{1 - a}{1 + 1}} \cdot \sqrt{\frac{1 - k}{1 + 1}}.$$
(21)

or

$$\frac{1-\nu}{1+\nu} = \sqrt{\frac{1-a}{1+ka}} \cdot \sqrt{\frac{1-k}{2}}.$$
 (21)

Therefore, for finding the asymptotics of m(P) we need to know the asymptotic behavior of a as $H \to \infty$. It is possible to do using (19), but we prefer to apply geometric considerations which are based on rectilinearity of the boundary arcs and the reflection principle. Let us prove the following auxiliary statement.

Lemma 4.1 Let Q be a quadrilateral which is the square $[0,1]^2$ with vertices at the points c, 1, 1+i, and i, where $c \in (0,1)$. Denote $Q_H = f_H(Q)$. Then

$$m(Q_H) \sim (1-c)H, \quad H \to \infty.$$

Proof. Let $\tilde{Q}_H = (1/H)Q_H$. Since $m(\tilde{Q}_H)$ is a monotonic function of H, it is sufficient to consider the sequence $H_n = 2n$ and to prove that

$$m(Q_{H_n}) \sim (1-c)H_n, \quad n \to \infty.$$

For short we will write Q_n instead of Q_{H_n} .





Consider the domains $Q_n^1, Q_n^2, \ldots, Q_n^{2n}$, where

$$Q_n^k = [0,1] \times \left[\frac{k-1}{2n}, \frac{k}{2n}\right]$$

Let us glue Q_n^k and Q_n^{k+1} along $\{(x, y) \mid 0 \le x \le 1, y = j/(2n)\}$ for odd k, and along $\{(x, y) \mid c \le x \le 1, y = j/(2n)\}$ for even k. As a result, we obtain the domain G_n which is the unit square with (n-1) horizontal slits (Fig. 2).

We will consider G_n as a quadrilateral with vertices c, 1, c+i, and i. By the symmetry principle, $m(G_n) = m(Q_n)/(2n)$. The domains G_n converge to the rectangule $G := [c, 1] \times [0, 1]$ as $n \to \infty$, and the sequences of their vertices converge to four distinct prime ends of $\tilde{G} = (G_n)$ corresponding to the vertices of G. To show that we take (1/4) of concentric circles with radius $r_m \to 0$ as sections γ_m which define prime end being a vertex of G. Let γ_{mn} be a section of G_n which is the union of γ_m and, if it is necessary, a segment connecting one of its endpoint to one of the nearest points of ∂G_n . By Theorem 3.3, $m(G_n) \to m(G) = 1 - c$, and Lemma 4.1 is proved.

Remark 4.2 The quadrilateral Q_H could be considered as a generalized long quadrilateral. Asymptotics of the modules of long quadrilaterals was investigated in [16], [17], [15], [18], [19], and other papers where various methods for computing the modules were suggested.

Consider the quadrilateral P^* which is the rectangle P with the segments $[0, (1-\sigma)H]$ and $[-\tilde{\sigma} H+i, (1-\sigma)H+i]$ as horizontal sides. Taking into account conformal invariance of the module, by Lemma 4.1 we have

$$m(P^*) \sim (1 - \sigma)H, \quad H \to \infty.$$
 (22)

Now we can describe the behavior of a as $H \to \infty$. Let us map P^* conformally onto U such that the points $(1 - \sigma)H + i$, $-\tilde{\sigma}H + i$, 0, and $(1 - \sigma)H$ are mapped into $-1/\mu$, -1, 1, and $1/\mu$, $0 < \mu < 1$. Then

$$m(P^*) = \frac{K(\mu')}{2K(\mu)},$$
 (23)

We should note that because of $m(P^*) \to \infty$ as $H \to \infty$, by (23) and (4), we have $\mu \to 0, H \to \infty$. Comparing the anharmonic ratios of the points in ∂U , corresponding to each other under conformal automorphism, we obtain

$$\frac{1-1/\mu}{1+1} \cdot \frac{-1/\mu + 1}{-1/\mu - 1/\mu} = \frac{a-1}{a+1/k} \cdot \frac{1/k + 1/k}{1/k - 1}$$

 or

$$\frac{1-a}{1+ak} \cdot \frac{2k}{1-k} = \frac{(\mu-1)^2}{4\mu}.$$

Therefore, taking into account that $a, k \to 1$ as $H \to \infty$ we have

$$1 - a \sim \frac{1 - k}{\mu}.\tag{24}$$

By (22), (23), and (3),

$$\frac{1}{(1-\sigma)H} \sim \frac{2K(\mu)}{K(\mu')} \sim \frac{4}{\pi} \ln \frac{1}{\mu} \,. \tag{25}$$

Now we use the asymptotic behavior (3) of the elliptic integrals. With use of that and by (18)

$$\ln \frac{1}{1-k} \sim \pi \frac{K(k)}{K(k')} \sim \frac{\pi}{2} \cdot \frac{(1-\sigma)^2 + \sigma}{1-\sigma} H.$$

Now from (3), (20), (21), (24), and (25) we obtain

$$m(\tilde{P}) = \frac{2K(\nu)}{K(\nu')} \sim \frac{2}{\pi} \ln \frac{1}{1-\nu} \sim \frac{2}{\pi} \ln \frac{1+ak}{1-a} + \frac{1}{\pi} \ln \frac{1}{1-k}$$
$$\sim \frac{2}{\pi} \ln \frac{1}{1-k} - \frac{1}{\pi} \ln \frac{1}{\mu} \sim (1-\sigma+\tilde{\sigma})H - (1-\sigma)H = \tilde{\sigma}H.$$

Because of (14) and (17) it completes the proof of (1).

Acknowledgement. The author expresses his gratitude to Prof. M. Vuorinen for attraction attention to the problem and useful comments, and to E. V. Borisova for help in checking the argument in the proof of Theorem 2.1.

References

- R. Kühnau. The conformal module of quadrilaterals and of rings. In: Handbook of Complex Analysis: Geometric Function Theory, (ed. by R. Kühnau) Vol. 2. North Holland, Amsterdam: Elsevier, 2005, 99–129.
- [2] D. Betsakos, K. Samuelsson, and M. Vuorinen. The computation of capacity of planar condensers. Publications De L'institut Mathematique, 75(89), 233–252, (2004).
- [3] M. Vuorinen and X. Zhang. On exterior moduli of quadrilaterals and special functions. J. Fixed Point Theory Appl. (to appear), 16 pp.
- [4] V. N. Dubinin and M. Vuorinen. On conformal moduli of polygonal quadrilaterals. Israel Journal of Mathematics, 171(1), 111–125, (2009).
- [5] H. Hakula, A. Rasila, and M. Vuorinen. On moduli of rings and quadrilaterals: Algorithms and Experiments. Siam Journal on Scientific Computing, 33(1), 279–302, (2011).
- [6] F. Bowman. Introduction to Elliptic Functions with applications. English Universities Press Ltd., London, 1953.
- [7] E. V. Borisova and S.R. Nasyrov. Asymptotics of the module of doublyconnected domain which is the difference of two homothetic rectangles. Trudi Matem. Tzentr. N. I. Lobachevskogo, 2011, 44, 62–63. (in Russian)
- [8] L. V. Ahlfors. Lectures on quasiconformal mappings. D. Van Nostrand Company, Toronto-New York-London, 1966.
- [9] S. R. Nasyrov. Asymptotics of the module of a rectangular frame under stretching it along a coordinate axis. Complex Analysis and Appl. Materiali VI Petrozavodskoi Mezhd. Konf. (1–7 July, 2012, Petrozavodsk, Petr. Univ), 51–53. (in Russian)

- [10] G. D. Anderson, M. K. Vamanamurthy, M. Vuorinen. Conformal Invariants, Inequalities, and Quasiconformal Maps. New York: Wiley, 1997.
- [11] N. I. Akhiezer. Elements of the theory of elliptic functions. Moscow, Fizmatlit, 1970, 304 pp. Translated into English as AMS Translations of Mathematical Monographs. Vol. 79 (1990) AMS, Rhode Island.
- [12] G. D. Suvorov. Prime ends and sequences of plane mappings. Kiev: Naukova Dumka, 1986. 192 pp. (in Russian)
- [13] S. R. Nasyrov. Geometric problems of theory of ramified coverings of Riemann surfaces. Kazan, Magarif, 2008, 276 pp. (in Russian)
- [14] R. Kühnau. Zum konformen Modul eines Vierecks. Mitt. Math. Seminar Gieen, 211, 61–67, (1992). (in German)
- [15] R. Kühnau. Der konforme Modul schmaler Vierecke. Math. Nachr., 175, 193–198, (1995). (in German)
- [16] D. Gaier and W. K. Hayman. Moduli of long quadrilaterals and thick ring domains. Rend. Math. Appl., 10(7), 809–834, (1990).
- [17] D. Gaier and W. K. Hayman. On the computation of modules of long quadrilaterals. Constr. Approx., 7, 453–467, (1991).
- [18] R. Laugesen. Conformal mapping of long quadrilaterals and thick doubly connected domains. Constr. Approx., 1994, 10, 523-554, (1994).
- [19] M. I. Falcão, N. Papamichael, and N. S. Stylianopoulos. Approximating the conformal maps of elongated quadrilaterals by domain decomposition. Constr. Approx., 17, 589-617, (2001).

INSTITUTE OF MATHEMATICS AND MECHANICS, KAZAN (VOLGA REGION) FEDERAL UNIVERSITY, E-MAIL: snasyrov@kpfu.ru