

Nonabelian Dark Matter with Resonant Annihilation

Cheng-Wei Chiang,^{1,2,3} Takaaki Nomura,^{1,4} and Jusak Tandean⁵

¹*Department of Physics and Center for Mathematics and Theoretical Physics,
National Central University, Chungli 320, Taiwan*

²*Institute of Physics, Academia Sinica, Taipei 115, Taiwan*

³*Physics Division, National Center for Theoretical Sciences, Hsinchu 300, Taiwan*

⁴*Department of Physics, National Cheng-Kung University, Tainan 701, Taiwan*

⁵*Department of Physics and Center for Theoretical Sciences,
National Taiwan University, Taipei 106, Taiwan*

Abstract

We construct a model based on an extra gauge symmetry, $SU(2)_X \times U(1)_{B-L}$, which can provide gauge bosons to serve as weakly-interacting massive particle dark matter. The stability of the dark matter is naturally guaranteed by a discrete Z_2 symmetry that is a subgroup of $SU(2)_X$. The dark matter interacts with standard model fermions by exchanging gauge bosons which are linear combinations of $SU(2)_X \times U(1)_{B-L}$ gauge bosons. With the appropriate choice of representation for the new scalar multiplet whose vacuum expectation value spontaneously breaks the $SU(2)_X$ symmetry, the relation between the new gauge boson masses can naturally lead to resonant pair annihilation of the dark matter. After exploring the parameter space of the new gauge couplings subject to constraints from collider data and the observed relic density, we use the results to evaluate the cross section of the dark matter scattering off nucleons and compare it with data from the latest direct detection experiments. We find allowed parameter regions that can be probed by future direct searches for dark matter and LHC searches for new particles.

I. INTRODUCTION

The standard model (SM) of particle physics has been very successful in describing an enormous amount of experimental data at energies up to $\mathcal{O}(100)$ GeV. There are, however, questions remaining that require physics beyond the minimal SM to address. Among the outstanding issues are the explanations for the astronomical evidence of dark matter (DM) and for the numerous experimental indications of neutrino mass [1]. It is then of great interest to explore a new physics scenario in which the DM and neutrino sectors are intimately connected.

Previously, we have considered a simple model which provides not only DM of the popular weakly-interacting massive particle (WIMP) type, but also a means to endow neutrinos with mass [2]. The DM candidate belongs to a complex scalar singlet stabilized by a Z_2 symmetry that is not imposed in an *ad hoc* way, but instead emerges from an extra Abelian gauge group related to baryon number minus lepton number, $U(1)_{B-L}$, that is spontaneously broken by the nonzero vacuum expectation value (VEV) of a new scalar field, in the Krauss-Wilczek manner [3]. Light neutrino masses are produced via the well-known seesaw mechanism [4], which is triggered with the involvement of the same new scalar field after the addition of right-handed neutrinos. The DM relic density receives contributions mainly from diagrams mediated by the Higgs boson and also those mediated by the $U(1)_{B-L}$ gauge boson, Z' . It turns out that constraints from collider data and the observed relic density together imply that the Z' mass has to be in the resonance region of the Z' -mediated DM annihilation, namely about twice the DM mass. Furthermore, results from DM direct detection experiments and Higgs data from the LHC favor the dominance of the Z' -exchange contributions to the relic density. All this motivates us to look for a different possible scenario in which the resonance condition can be fulfilled naturally.

In this paper, we demonstrate that such a possibility can be realized in a model where the role of WIMP DM is played by massive gauge bosons associated with a nonabelian symmetry. Although most of the WIMP DM candidates proposed in the literature are either fermionic or spinless, those with spin one have also been considered before [5]. Here we construct a model based on the gauge group $G_{\text{SM}} \times SU(2)_X \times U(1)_{B-L}$, where G_{SM} refers to the SM group and the extra gauge symmetries offer gauge bosons which can act as WIMP candidates. The stability of the DM is naturally maintained by a discrete Z_2 symmetry which is a subgroup of the new nonabelian gauge group, $SU(2)_X$. This Z_2 symmetry appears after the spontaneous breaking of $SU(2)_X$ by the nonzero VEV of a new scalar multiplet, following the Krauss-Wilczek mechanism [3]. Then the Z_2 -odd gauge bosons associated with $SU(2)_X$ can serve as DM if they are lighter than other Z_2 -odd particles in the model. Since SM fermions are charged under $U(1)_{B-L}$, the DM can interact with SM fermions at tree level by exchanging gauge bosons which are obtained from the linear combinations of $SU(2)_X$ and $U(1)_{B-L}$ gauge fields. Thus, the new gauge interactions are responsible for both the relic abundance and the DM interactions with nucleons. Another interesting feature of the model is that, the DM being made up of $SU(2)_X$ gauge bosons, its mass is related to the masses of the mediating gauge bosons, implying that resonant pair annihilation can be naturally achieved by choosing suitable representations of the scalar fields involved in the breaking of the $SU(2)_X \times U(1)_{B-L}$ gauge symmetry and ensuring that their VEVs are sufficiently well separated. What's more, the presence of the $U(1)_{B-L}$ gauge symmetry requires the introduction of right-handed neutrinos for gauge-anomaly cancellation, which in turn participate in the type-I seesaw mechanism to generate

light neutrino masses [4], with the right-handed neutrino masses being connected to the $U(1)_{B-L}$ breaking scale. This model turns out to have sufficient parameter space that is consistent with current collider, relic density, and DM direct search data. Therefore, it can be probed further by ongoing or future DM direct detection experiments, and some of the new particles may be observable at the LHC with sufficient luminosities.

This paper is organized as follows. The next section contains the details of our model which possesses WIMP DM composed of the gauge bosons of an extra nonabelian gauge symmetry. We explain how the choices of the new particles and their quantum numbers can naturally translate into resonant annihilation of the DM. In Section III, we examine constraints on the new gauge couplings from collider data. In Section IV, we deal with the relic density of our DM candidates and extract the parameter values allowed by its observed value. In Section V, we use the results to predict the DM-nucleon scattering cross-section and compare it with current data from direct detection experiments. In Section VI, we comment on the collider phenomenology of the new particles in our model. We conclude in Section VII with the summary of our study and some more discussion.

II. A MODEL OF DARK MASSIVE GAUGE BOSON

Compared to the SM, the new model contains the additional gauge group $SU(2)_X \times U(1)_{B-L}$, where X refers to the massive gauge boson that serves as the DM, whereas B and L stand for baryon and lepton numbers, respectively. We denote the gauge fields associated with $SU(2)_X$ and $U(1)_{B-L}$ by C_k^μ and E^μ , respectively, $k = 1, 2, 3$, and their coupling constants g_X and g_{B-L} . The model also has new complex scalar fields S and Φ_5 as well as three extra fermions ν_{kR} , all of which are singlets under the SM gauge group, but carry nonzero $U(1)_{B-L}$ charges. Under $SU(2)_X$ transformations, S is a singlet, while Φ_5 is a five-plet represented by the column matrix $\Phi_5 = (\phi_2, \phi_1, \phi_0, \phi_{-1}, \phi_{-2})^T$, where ϕ_a corresponds to the eigenvalue $T_{3X} = a$ of the third generator of $SU(2)_X$. In Table I we collect the $SU(2)_X \times U(1)_{B-L}$ quantum number assignments for the fermions, scalars, and new gauge bosons in the model, with H being the usual scalar doublet.

The renormalizable Lagrangian for S and Φ_5 , with H included in the potential \mathcal{V} , is

$$\mathcal{L} = (\mathcal{D}^\mu S)^\dagger \mathcal{D}_\mu S + (\mathcal{D}^\mu \Phi_5)^\dagger \mathcal{D}_\mu \Phi_5 - \mathcal{V} , \quad (1)$$

where

$$\mathcal{D}^\mu S = \partial^\mu S + 2ig_{B-L} E^\mu S , \quad \mathcal{D}^\mu \Phi_5 = \partial^\mu \Phi_5 + ig_X C_k^\mu \mathcal{T}_k^{(5)} \Phi_5 + ig_{B-L} E^\mu \mathcal{Q}_{B-L}^{(5)} \Phi_5 , \quad (2)$$

$$\mathcal{V} = -\mu_\Phi^2 \Phi_5^\dagger \Phi_5 + (\lambda_S |S|^2 - \mu_S^2) |S|^2 + (\lambda_H H^\dagger H - \mu_H^2) H^\dagger H + (\text{other quartic terms}) . \quad (3)$$

| | f_{SM} | ν_R | H | S | ϕ_2 | ϕ_1 | ϕ_0 | ϕ_{-1} | ϕ_{-2} | X | X^\dagger | C_3 | E |
|------------------------|-----------------|----------|---------|---------|----------|----------|----------|-------------|-------------|---------|-------------|---------|---------|
| $SU(2)_X [U(1)_{B-L}]$ | $1 [B-L]$ | $1 [-1]$ | $1 [0]$ | $1 [2]$ | $5 [2]$ | $5 [2]$ | $5 [2]$ | $5 [2]$ | $5 [2]$ | $3 [0]$ | $3 [0]$ | $3 [0]$ | $1 [0]$ |
| T_{3X} | 0 | 0 | 0 | 0 | 2 | 1 | 0 | -1 | -2 | 1 | -1 | 0 | 0 |
| Z_2^X | + | + | + | + | + | - | + | - | + | - | - | + | + |

TABLE I: The charge assignments under $SU(2)_X \times U(1)_{B-L}$ and Z_2^X parity of the fermions, scalars and new gauge bosons in the model, with f_{SM} referring to SM fermions, $X = (C_1 - iC_2)/\sqrt{2}$, and T_{3X} denoting the eigenvalue of the third generator of $SU(2)_X$.

In $\mathcal{D}^\mu\Phi_5$ above, summation over $k = 1, 2, 3$ is implicit, and $\mathcal{T}_k^{(5)}$ and $\mathcal{Q}_{B-L}^{(5)}$ are matrices for the generators of $SU(2)_X$ and $U(1)_{B-L}$, respectively, acting on Φ_5 , where

$$\begin{aligned} \mathcal{T}_1^{(5)} &= \frac{1}{2} \begin{pmatrix} 0 & 2 & 0 & 0 & 0 \\ 2 & 0 & \sqrt{6} & 0 & 0 \\ 0 & \sqrt{6} & 0 & \sqrt{6} & 0 \\ 0 & 0 & \sqrt{6} & 0 & 2 \\ 0 & 0 & 0 & 2 & 0 \end{pmatrix}, & \mathcal{T}_2^{(5)} &= \frac{i}{2} \begin{pmatrix} 0 & -2 & 0 & 0 & 0 \\ 2 & 0 & -\sqrt{6} & 0 & 0 \\ 0 & \sqrt{6} & 0 & -\sqrt{6} & 0 \\ 0 & 0 & \sqrt{6} & 0 & -2 \\ 0 & 0 & 0 & 2 & 0 \end{pmatrix}, \\ \mathcal{T}_3^{(5)} &= \text{diag}(2, 1, 0, -1, -2), & \mathcal{Q}_{B-L}^{(5)} &= \text{diag}(2, 2, 2, 2, 2). \end{aligned} \quad (4)$$

In this paper, we consider the scenario in which the $SU(2)_X \times U(1)_{B-L}$ gauge symmetry is spontaneously broken according to

$$SU(2)_X \times U(1)_{B-L} \xrightarrow{\langle S \rangle} SU(2)_X \times Z_2^{B-L} \xrightarrow{\langle \Phi_5 \rangle} Z_2^X \times Z_2^{B-L}, \quad (5)$$

where $\langle S \rangle = v_S/\sqrt{2}$ and $\langle \Phi_5 \rangle = (v_\Phi, 0, 0, 0, 0)^T/\sqrt{2}$ are the VEVs of S and Φ_5 , with $v_S \gg v_\Phi > 0$. Since $\langle \Phi_5 \rangle \neq 0$ occurs via its $T_{3X} = 2$ component, $\langle \phi_2 \rangle \neq 0$, the Z_2^X symmetry emerges naturally as a subgroup of $SU(2)_X$ and the particles with even (odd) T_{3X} values will be Z_2^X even (odd), as Table I shows. On the other hand, Z_2^{B-L} is the remnant of $U(1)_{B-L}$ after $\langle S \rangle \neq 0$, as discussed in Ref. [2], but does not play a role in the stabilization of X . Thus, in this scenario the remaining Z_2^X guarantees the stability of the lightest Z_2^X -odd particle(s), which can therefore act as DM. Here we choose the gauge boson $X = (C_1 - iC_2)/\sqrt{2}$ and its conjugate X^\dagger to be the DM, hence tacitly taking the Z_2^X -odd scalar bosons to be more massive than X . It is worth mentioning that we would arrive at the same results below if $\langle \Phi_5 \rangle \neq 0$ through its $T_{3X} = -2$ component instead. As for H , its VEV is also nonvanishing and breaks the electroweak symmetry just as in the SM. We assume that the other parameters in the potential \mathcal{V} are such that the vacuum has the above desired properties, leaving a detailed analysis of \mathcal{V} for future work.

After $SU(2)_X \times U(1)_{B-L}$ spontaneously breaks into $Z_2^X \times Z_2^{B-L}$, the new gauge bosons acquire in \mathcal{L} the mass terms

$$\begin{aligned} \mathcal{L}_m &= \langle \Phi_5^\dagger \rangle \left[g_X C_k^\mu \mathcal{T}_k^{(5)} + g_{B-L} E^\mu \mathcal{Q}_{B-L}^{(5)} \right] \left[g_X C_{k'\mu} \mathcal{T}_{k'}^{(5)} + g_{B-L} E_\mu \mathcal{Q}_{B-L}^{(5)} \right] \langle \Phi_5 \rangle + 4g_{B-L}^2 E^2 \langle S \rangle^2 \\ &= g_X^2 v_\Phi^2 X_\mu^\dagger X^\mu + \frac{1}{2} (C_3^\mu \quad E^\mu) \begin{pmatrix} 4g_X^2 v_\Phi^2 & 4g_X g_{B-L} v_\Phi^2 \\ 4g_X g_{B-L} v_\Phi^2 & 4g_{B-L}^2 (v_\Phi^2 + v_S^2) \end{pmatrix} \begin{pmatrix} C_{3\mu} \\ E_\mu \end{pmatrix}. \end{aligned} \quad (6)$$

From the last line, upon diagonalizing the 2×2 matrix in the second term, we obtain the eigenvalues

$$m_X^2 = g_X^2 v_\Phi^2, \quad (7)$$

$$m_{Z_L, Z_H}^2 = 2g_X^2 v_\Phi^2 + 2g_{B-L}^2 (v_\Phi^2 + v_S^2) \mp 2\sqrt{[g_X^2 v_\Phi^2 - g_{B-L}^2 (v_\Phi^2 + v_S^2)]^2 + 4g_X^2 g_{B-L}^2 v_\Phi^4}, \quad (8)$$

assuming that $m_{Z_L} < m_{Z_H}$ for the mass eigenstates Z_L and Z_H which are given by

$$\begin{pmatrix} Z_L \\ Z_H \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} C_3 \\ E \end{pmatrix}, \quad (9)$$

$$\tan(2\theta) = \frac{2g_X g_{B-L} R_v}{g_X^2 R_v - g_{B-L}^2 (1 + R_v)}, \quad R_v = \frac{v_\Phi^2}{v_S^2}. \quad (10)$$

In this study, we focus on the case in which $v_S^2 \gg v_\Phi^2$ and $g_X \sim g_{B-L}$, implying that

$$|\theta| \simeq \frac{g_X}{g_{B-L}} R_v, \quad (11)$$

$$m_{Z_L}^2 \simeq 4m_X^2(1 - R_v), \quad (12)$$

$$m_{Z_H}^2 \simeq 4m_X^2 \frac{g_{B-L}^2}{g_X^2} R_v (1 + R_v). \quad (13)$$

Accordingly, with $R_v \ll 1$, we obtain the mass relation

$$m_{Z_L} \simeq 2m_X, \quad (14)$$

which naturally leads to resonant annihilation of the DM pair via the Z_L -mediated contribution.

It is worth noting that the five-plet Φ_5 is the minimal choice of $SU(2)_X$ representation that can result in the resonant relation in Eq. (14). In general, for an $SU(2)_X$ isospin value T_X and its third component T_{3X} , one would get $m_X^2/m_{Z_L}^2 \simeq [T_X(T_X + 1) - T_{3X}^2]/(2T_{3X}^2)$ assuming small mixing angle θ , in analogy to the ρ parameter in the electroweak sector [1].

The neutrino mass-generating sector is the same as that given in Ref. [2], the relevant Lagrangian having the form

$$\mathcal{L}_{m_\nu} = i\lambda_{kl} \bar{\nu}_{kR} H^T \tau_2 L_{lL} - \frac{1}{2} \lambda'_{kl} \bar{\nu}_{kR} (\nu_{lR})^c S^\dagger + \text{H.c.}, \quad (15)$$

where summation over $k, l = 1, 2, 3$ is implicit, $\lambda_{kl}^{(\prime)}$ are free parameters, τ_2 is the second Pauli matrix, L_{lL} represents a lepton doublet, and the superscript c indicates charge conjugation. The Dirac and Majorana mass matrices from these terms are $\mathcal{M}_D = \lambda v_H/\sqrt{2}$ and $\mathcal{M}_{\nu_R} = \lambda' v_S/\sqrt{2}$, respectively, where v_H is the VEV of H . Hence v_S sets the mass scale of the right-handed neutrinos, ν_{kR} . In our examples later on, we will see what values of v_S are compatible with the observed relic density and collider data.

Since X is our chosen candidate for DM and interacts with SM fermions by exchanging the $Z_{L,H}$ bosons at tree level, in the following two sections we evaluate the new gauge couplings subject to collider and relic density data. Subsequently, we use the allowed values of the couplings to predict the cross section of the DM-nucleon scattering and compare it with the existing results of DM direct detection experiments.

III. CONSTRAINTS FROM COLLIDER EXPERIMENTS

The gauge bosons Z_L and Z_H interact with SM fermions at tree level with coupling constants $g_{B-L} \sin \theta$ and $g_{B-L} \cos \theta$, respectively, according to the Feynman rules listed in Appendix A. It follows that measurements on processes mediated by Z_L and Z_H can offer constraints on these couplings. Significant restrictions may be available from the data on e^+e^- and hadron collisions into fermion pairs, which we treat in this section.

We first look at the constraints from $e^+e^- \rightarrow f\bar{f}$ scattering. In this work we assume that mixing between the Z boson and $Z_{L,H}$ is negligible, but we will comment on the impact of kinetic mixing between them later on and discuss it further in Appendix B. In the absence of the mixing, the new gauge couplings have no effects on the Z -pole observables at leading order. On the other hand,

the measurements of $e^+e^- \rightarrow f\bar{f}$ at LEP II with center-of-mass energies from 130 to 207 GeV are relevant [6]. We employ the data on the cross section and forward-backward asymmetry for $f = \mu, \tau$ and on the cross section for $f = \text{quark}$. To evaluate the limits on the new couplings, we include both the Z_L and Z_H contributions to the scattering amplitude, their couplings and masses satisfying the relations in Eqs. (11)-(13). Although Z_H is much heavier than Z_L , the fermionic couplings of the latter can be much smaller than those of the former to compensate for the suppression of the Z_H contribution to the amplitude due to its bigger mass. In the examples presented below, the Z_H contributions to $e^+e^- \rightarrow f\bar{f}$ turn out to dominate the Z_L ones.

For definiteness and simplicity, hereafter we set $g_X = g_{B-L} \geq 0$. Adopting the 90% confidence-level (CL) ranges of the LEP II measurements [6] and using the formulas given in Ref. [7], but with s -dependent Z and $Z_{L,H}$ widths [6], we then scan the m_X and R_ν space. To illustrate the results, we display in Figure 1 the upper limits on g_X versus m_X for $R_\nu = 10^{-2}$ (red dashed curve) and 10^{-3} (blue dashed curve) on the left and right sides, respectively. The horizontal, straight portions of the curves correspond to the perturbativity requirement, $g_X < \sqrt{4\pi}$.

The most recent data from the LHC on the cross-section of the Drell-Yan (DY) process in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$ with 4.5 fb^{-1} of integrated luminosity have revealed no discrepancy from the SM expectations and therefore no evidence of $Z_{L,H}$ bosons [8]. Consequently, we follow the same analysis as in Refs. [2, 9] to derive upper bounds on the coupling constants using the SM cross-section. In the present case, we can consider the Z_L and Z_H contributions separately because we focus on events with dilepton invariant mass around m_{Z_L} or m_{Z_H} where their effects are of different orders for small R_ν . Thus respective constraints are obtained for the pairs $(m_{Z_L}, g_{B-L} \sin \theta)$ and $(m_{Z_H}, g_{B-L} \cos \theta)$. To estimate the DY cross-section numerically, we utilize the CalcHEP package [10] by incorporating the new particles and Feynman rules of our model. Then we apply the one-bin log likelihood $LL = 2[N \ln(N/\nu) + \nu - N]$, where N (ν) is the number




FIG. 1: Upper limits on g_X versus m_X from LEP II and LHC data on $e^+e^- \rightarrow f\bar{f}$ and Drell-Yan scattering, respectively, for $R_\nu = 10^{-2}$ (left) and 10^{-3} (right) under the assumption that $g_X = g_{B-L}$, compared to the corresponding values of g_X (solid curves) consistent with the observed relic density. The horizontal, straight portions of the dashed and dotted curves correspond to the perturbativity condition, $g_X < \sqrt{4\pi}$.

of events predicted by the SM (SM plus the Z_L or Z_H boson) in the $\ell^+\ell^-$ invariant mass window of $\pm 20\%$ around the expected Z_L or Z_H mass, with $\sqrt{s} = 7 \text{ TeV}$ and 4.5 fb^{-1} of luminosity. The upper limit on the cross-section is obtained from the solved value of ν for each Z_L or Z_H mass, after adopting $LL = 2.7$ which corresponds to the 90% CL.

We find that the Z_H contribution to the DY process yields a stricter bound on g_{B-L} as a function of m_{Z_H} , as the Z_L contribution is strongly suppressed by the small $|\theta|$. We show the resulting upper-limits on $g_X (= g_{B-L})$ in Figure 1, where m_X is related to m_{Z_H} by Eq. (13), for $R_\nu = 10^{-2}$ (red dotted curve) and 10^{-3} (blue dotted curve) on the left and right, respectively. We notice that the limit in the $R_\nu = 10^{-2}$ case becomes large at $m_X \sim 5 \text{ GeV}$ corresponding to $m_{Z_H} \sim m_Z$ where the SM background is large.

IV. RESONANT DARK MATTER ANNIHILATION AND RELIC DENSITY

Now we estimate the relic density of the DM particle, X , in order to search for the model parameter space consistent with the observed relic density. The thermal relic abundance is found by solving the Boltzmann equation which describes the number density of the DM. We employ the approximate solution to the Boltzmann equation for the present-day relic density Ω , given by [11]¹

$$\begin{aligned} \Omega h^2 &= \frac{1.07 \times 10^9}{\sqrt{g_*} m_{\text{Pl}} J \text{ GeV}}, & J &= \int_{x_f}^{\infty} dx \frac{\langle \sigma v \rangle}{x^2}, \\ x_f &= \ln \frac{0.038 g m_X m_{\text{Pl}} \langle \sigma v \rangle}{\sqrt{g_*} x_f}, \end{aligned} \quad (16)$$

where h denotes the Hubble constant in units of 100 km/s/Mpc, g_* is the number of relativistic degrees of freedom below the freeze-out temperature $T_f = m_X/x_f$, $m_{\text{Pl}} = 1.22 \times 10^{19} \text{ GeV}$ is the Planck mass, $g = 3$ to account for X having spin-1, and $\langle \sigma v \rangle$ is the thermal average of the DM annihilation cross-section. More explicitly [12],

$$\langle \sigma v \rangle = \frac{x}{8m_X^5 K_2^2(x)} \int_{4m_X^2}^{\infty} ds \sqrt{s} (s - 4m_X^2) K_1(\sqrt{s} x/m_X) \sigma_{\text{ann}}, \quad (17)$$

where K_i is the modified Bessel function of the second kind of order i and σ_{ann} represents the cross section of $X^\dagger X$ annihilation into all possible final states.

Under the assumptions made in Section II, we find that the main contributions to σ_{ann} come from the s -channel transitions $X^\dagger X \rightarrow Z_L^* \rightarrow f_{\text{SM}} \bar{f}_{\text{SM}}$. Although Z_H -mediated diagrams also contribute, in this case they can be neglected because of the suppression due to $m_{Z_H} \gg m_{Z_L}$ and their lack of the resonance enhancement of the Z_L -mediated diagrams in the nonrelativistic region $\sqrt{s} \sim 2m_X$ due to $m_{Z_L} \simeq 2m_X$. Thus, with the Feynman rules in Appendix A, we arrive at

$$\begin{aligned} \sigma_{\text{ann}} &= \frac{g_X^2 g_{B-L}^2 \cos^2 \theta \sin^2 \theta}{432\pi} \sum_f \frac{\sqrt{(s - 4m_X^2)(s - 4m_f^2)}}{m_X^4 s} \frac{s^2 + 20 m_X^2 s + 12 m_X^4}{(s - m_{Z_L}^2)^2 + \Gamma_{Z_L}^2 m_{Z_L}^2} \\ &\quad \times \left[(s + 2m_f^2) |\hat{V}_f^{Z_L}|^2 + (s - 4m_f^2) |\hat{A}_f^{Z_L}|^2 \right] N_c^f, \end{aligned} \quad (18)$$

¹ For a more accurate approximation, see [27].

where the sum is over all fermions with masses $m_f < m_X$ and color factors N_c^f , the couplings $\hat{V}_f^{Z_L}$ and $\hat{A}_f^{Z_L}$ are given in Eq. (A2), and Γ_{Z_L} is the width of Z_L . Now, since $m_{Z_L}^2 = 4m_X^2(1 - R_v)$ and $s \geq 4m_X^2$ according to Eqs. (12) and (17), respectively, in the denominator of σ_{ann} above we have $(s - m_{Z_L}^2)^2 \geq 16m_X^4 R_v^2$. From the collider bounds on $g_{B-L} = g_X$ derived in the previous section, we find that for the mass range of interest $16m_X^4 R_v^2 \gg \Gamma_{Z_L}^2 m_{Z_L}^2$. Consequently, the Γ_{Z_L} term can be neglected in the calculation of Eq. (17).

With Eqs. (16)-(18), we can extract the (g_X, m_X) regions compatible with the observed Ω . Its most recent value has been determined by the Planck Collaboration from the Planck measurement and other data to be $\Omega h^2 = 0.1187 \pm 0.0017$ [13]. Accordingly, we require the relic density of X to satisfy the 90% CL (confidence level) range of its experimental value, $0.1159 \leq \Omega h^2 \leq 0.1215$. As mentioned in the preceding section, for simplicity we take $g_X = g_{B-L}$, implying that $|\theta| \simeq R_v$. The plots in Figure 1 display the resulting g_X values allowed by the relic data for $R_v = 10^{-2}$ (red solid curve) and 10^{-3} (blue solid curve) on the left and right panels, respectively. One can see that, although the $f\bar{f}Z_L$ couplings are suppressed by the small mixing angle, $|\theta| \ll 1$, the observed relic density can be reproduced with moderate-sized couplings $g_{B-L} = g_X = \mathcal{O}(0.1)\text{-}\mathcal{O}(1)$ over $m_X \leq 1000 \text{ GeV}$ due to the resonance enhancement. This can be partly understood from the fact that in the resonance region the denominator of σ_{ann} is dominated by the term $(4m_X^2 - m_{Z_L}^2)^2 \propto R_v^2$ which approximately cancels the R_v^2 factor in the numerator.

In Figure 1, we can also compare the coupling ranges that fulfill the requirements from both the collider and relic density data. Evidently, the constraints from LEP II data restrict the allowed masses to $m_X \gtrsim 400$ (220) GeV with couplings of $\mathcal{O}(1)$ for $R_v = 10^{-2}$ (10^{-3}). The cases with $R_v \lesssim 10^{-4}$ and $m_X \leq 1000 \text{ GeV}$ are excluded by the LEP II constraints.

Since we have the relation $m_X = g_X v_S \sqrt{R_v}$ from Eqs. (7) and (10), it is interesting to explore the values of v_S subject to the same experimental requirements. We illustrate this in Figure 2 obtained with the allowed g_X ranges in Figure 1. Hence v_S should be between about 5 and 10 TeV in order to satisfy both the collider and relic data. This suggests that our model is compatible with the TeV-scale type-I seesaw scenario.

V. DIRECT DETECTION OF DARK MATTER

The direct detection of X relies on its scattering off a nucleon N elastically, $XN \rightarrow XN$, which proceeds from $Z_{L,H}$ exchanges in the t channel. Since $m_{Z_H} \gg m_{Z_L}$, the Z_H contribution can be neglected. It follows that in the nonrelativistic limit the cross section of $XN \rightarrow XN$ is

$$\sigma_{\text{el}}^N = \frac{g_X^2 g_{B-L}^2 \cos^2 \theta \sin^2 \theta \mu_{XN}^2}{\pi m_{Z_L}^4} \simeq \frac{g_X^4 R_v^2 \mu_{XN}^2}{16\pi m_X^4}, \quad (19)$$

where $\mu_{XN} = m_X m_N / (m_X + m_N)$ and we have made use of $\langle N | \bar{u} \gamma^\alpha u + \bar{d} \gamma^\alpha d | N \rangle = 3\bar{N} \gamma^\alpha N$ [14], the other quarks having vanishing contributions. This indicates that σ_{el}^N is not sensitive to g_{B-L} for fixed $R_v \ll 1$.

In Figure 3 we plot σ_{el}^N as a function of m_X for the allowed parameter regions in Figure 1, the red and blue strips belonging to the $R_v = 10^{-2}$ and 10^{-3} cases, respectively. Also shown are the recent data from DM direct searches. Clearly, much of the σ_{el}^N prediction still escapes the existing




FIG. 2: Values of v_S versus m_X satisfying the requirements from both the collider and relic density data and corresponding to the allowed g_X regions in Figure 1.

constraints, including the strictest ones from XENON100 [17] and LUX [20], but it will be probed more stringently by future direct searches such as XENON1T [25].

Before moving on, we would like to make a few remarks regarding the potential implications of mixing between the SM and extra gauge bosons in our model. Since none of the scalar fields in the theory carries both the electroweak and new quantum numbers, there is no mass mixing between the SM and new gauge bosons. In contrast, as discussed in Appendix B, kinetic mixing between the $U(1)_Y$ and $U(1)_{B-L}$ gauge bosons can occur both at tree and loop levels. We find that the impact of this mixing is not significant on the results above for the allowed values of the new gauge couplings and Z_L mass. Especially, the relation $m_{Z_L} \simeq 2m_X$ is unaffected. We further find that, although the Z_H mass is sensitive to the kinetic mixing, being enhanced by it, the effect can be minimized if the mixing parameter has a magnitude below 0.5. Our rough estimate of the relevant loop diagram in Appendix B suggests that mixing size of order 0.5 is not atypical. Lastly, since the X annihilation and X -nucleon scattering processes are dominated by the Z_L contributions, the increased m_{Z_H} would not be important for them. It follows that it is reasonable to neglect the impact of the kinetic mixing.

VI. COMMENTS ON COLLIDER PHENOMENOLOGY

In this section, we briefly discuss how the extra scalar and gauge bosons in our model may be produced and detected at the LHC. The new scalar bosons coming from Φ_5 and S comprise twelve degrees of freedom in total. Four of them are “eaten” by the new gauge bosons, making them massive. The remaining extra scalar bosons can be expressed as ϕ_i^{even} and ϕ_j^{odd} , which are linear combinations of Z_2^X -even and -odd particles, respectively. Since two of the new massive gauge bosons are Z_2^X even and the other two Z_2^X odd, there are six ϕ^{even} 's and two ϕ^{odd} 's which are




FIG. 3: Cross-section σ_{el}^N of $XN \rightarrow XN$ scattering corresponding to the allowed parameter regions in Figure 1. The predicted cross-sections are compared to 90% CL upper-limits from XENON10 (green dashed-dotted curve) [15, 16], XENON100 (black short-dashed curve) [17], CDMS Ge (red long-dashed curves) [18], CDMS Si (blue solid curve) [19], and LUX (purple dashed-double-dotted curve) [20]. The prediction is also compared to the 90% CL (magenta) signal region suggested by CoGeNT [21], a gray patch compatible with the DAMA modulation signal at the 3σ level [22], two 2σ -confidence (light brown) areas representing CRESST-II data [23], and a cyan area for a possible signal at 90% CL from CDMS II [24]. Also plotted is the XENON1T projected sensitivity (brown dashed-triple-dotted curve) [25].

physical. In this study we do not specify the new scalars' mass spectrum, but one could obtain it by doing a detailed analysis of the scalar potential.

Taking into account the Z_2^X parities of the new particles, we find the decay patterns

$$Z_L \rightarrow f_{\text{SM}} \bar{f}_{\text{SM}}, \quad (20)$$

$$Z_H \rightarrow f_{\text{SM}} \bar{f}_{\text{SM}}, XX^\dagger, \phi_i^{\text{even}} \phi_j^{\text{even}}, \phi_i^{\text{odd}} \phi_j^{\text{odd}}, \quad (21)$$

$$\phi_i^{\text{even}} \rightarrow Z_{L,H} Z_{L,H}, XX^\dagger, X^{(\dagger)} \phi_i^{\text{odd}}, Z_{L,H} \phi_i^{\text{even}}, \phi_j^{\text{even}} \phi_k^{\text{even}}, \phi_j^{\text{odd}} \phi_k^{\text{odd}}, \quad (22)$$

$$\phi_i^{\text{odd}} \rightarrow Z_{L,H} X, \phi_j^{\text{even}} X, \phi_j^{\text{even}} \phi_k^{\text{odd}}, \quad (23)$$

where the particles on the right-hand sides may be off-shell depending on the masses involved. Throughout we have assumed that X is lighter than new scalar bosons, and so Z_L decays mostly to SM fermions. Since the couplings of $Z_{L,H}$ to the fermions are proportional to their $B - L$ numbers, $Z_{L,H}$ tend to decay into leptons rather than quarks, as the decay rates of $Z_{L,H}$ into a charged lepton pair and into a quark-antiquark pair, with relatively negligible masses, are related by $\Gamma_{Z_{L,H} \rightarrow \ell^+ \ell^-} : \Gamma_{Z_{L,H} \rightarrow q \bar{q}} \simeq 1 : 3(1/3)^2 = 3 : 1$. The decay branching fractions of the scalar bosons depend on their mass spectrum and couplings in the potential.

The DM pair, XX^\dagger , can be produced through $Z_{L,H}$ exchanges in the s -channel according to

$$pp \rightarrow Z_{L,H}^{(*)} + [\text{jet}(s), \text{photon}(s), \text{etc.}] \rightarrow XX^\dagger + [\text{jet}(s), \text{photon}(s), \text{etc.}] , \quad (24)$$

where we need particles other than XX^\dagger , such as jets (j 's) or photons (γ 's), for tagging. Their production signals are therefore jet(s) plus missing energy, photon(s) plus missing energy, etc. We note that the Z_L -mediated contributions dominate these processes because Z_H is much heavier than Z_L .

We now estimate the cross-sections of $pp \rightarrow XX^\dagger j$ and $pp \rightarrow XX^\dagger \gamma$ using the CalcHEP code package [10] incorporating the new Feynman rules in the model file. The cross sections are computed for the Tevatron and the LHC with different colliding energies employing two parameter sets taken from Figure 1. The results are listed in Table II. These cross sections are small compared to current search limits due to the suppression by the small $|\theta|$. For example, it is estimated that the upper limit of the cross section for jets plus missing energy in the squark-gluino-neutralino scenario of the minimal supersymmetric standard model is $\mathcal{O}(1)$ fb for the LHC at $\sqrt{s} = 7$ TeV with 4.7 fb^{-1} of luminosity [26], which is larger than our cross sections. However, at the LHC the cross-section of $pp \rightarrow XX^\dagger j$ can reach ~ 0.1 fb for $\sqrt{s} = 8$ TeV and ~ 0.5 fb for $\sqrt{s} = 14$ TeV, which are potentially testable with the appropriate amount of luminosity. The DM can also be produced singly in association with ϕ_i^{odd} , such as in

$$pp \rightarrow Z_{L,H}^{(*)} + \dots \rightarrow X^{(\dagger)} \phi_i^{\text{odd}} + \dots . \quad (25)$$

Since ϕ^{odd} would decay according to Eq. (23), the specific signal would be charged leptons plus missing energy: $X^{(\dagger)} \phi_i^{\text{odd}} \rightarrow XX^\dagger Z_L \rightarrow XX^\dagger \ell^+ \ell^-$ where ℓ is the electron or muon. The cross section of this channel is expected to be of similar order as that of XX^\dagger production since the coupling constants involved are the same.

The scalar bosons ϕ_i^{even} and ϕ_i^{odd} can be produced through $Z_{L,H}$ exchanges in the s channel,

$$pp \rightarrow Z_{L,H}^* \rightarrow Z_{L,H} \phi_i^{\text{even}} , \quad (26)$$

$$pp \rightarrow Z_{L,H}^{(*)} \rightarrow \phi_i^{\text{odd}} \phi_j^{\text{odd}} , \quad (27)$$

$$pp \rightarrow Z_{L,H}^{(*)} \rightarrow \phi_i^{\text{even}} \phi_j^{\text{even}} , \quad (28)$$

| | m_X (GeV) | g_X | $ \theta $ | $\sigma_{XX^\dagger j}$ (fb) | $\sigma_{XX^\dagger \gamma}$ (fb) |
|------------|-------------|-------|------------|------------------------------|-----------------------------------|
| Tevatron | 300 | 1.8 | 0.001 | 6.5×10^{-3} | 1.7×10^{-4} |
| | 600 | 1.2 | 0.01 | 6.0×10^{-5} | 1.8×10^{-6} |
| LHC 7 TeV | 300 | 1.8 | 0.001 | 8.7×10^{-2} | 1.6×10^{-3} |
| | 600 | 1.2 | 0.01 | 5.1×10^{-2} | 1.1×10^{-3} |
| LHC 8 TeV | 300 | 1.8 | 0.001 | 8.6×10^{-2} | 2.3×10^{-3} |
| | 600 | 1.2 | 0.01 | 0.12 | 1.9×10^{-3} |
| LHC 14 TeV | 300 | 1.8 | 0.001 | 0.46 | 8.2×10^{-3} |
| | 600 | 1.2 | 0.01 | 0.51 | 1.0×10^{-2} |

TABLE II: Estimated cross-sections of pp ($p\bar{p}$) $\rightarrow XX^\dagger j$ and pp ($p\bar{p}$) $\rightarrow XX^\dagger \gamma$ for two parameter sets taken from Figure 1 .

as well as the channels in Eq. (25). According to the decay patterns in Eqs. (22) and (23), the signals of ϕ_i^{even} and ϕ_i^{odd} would be two pairs of charged leptons and charged leptons plus missing energy, respectively. The production cross-sections of the Z_L -exchange processes are suppressed by $|\theta|^2$ because the $Z_L f \bar{f}$ coupling is proportional to $\sin\theta$, whereas the production cross-sections of Z_H -exchange processes are suppressed by the heavy Z_H mass. Thus high luminosities will be required to get a sufficient number of signal events. To carry out a detailed analysis, one would need to specify the self-couplings and thus the mass spectrum of scalar bosons. We leave such a study for future work.

VII. SUMMARY AND DISCUSSION

In this paper we have constructed a model possessing an extra gauge symmetry, $SU(2)_X \times U(1)_{B-L}$, which offers a massive gauge boson, X , associated with $SU(2)_X$ playing the role of WIMP DM. The new gauge bosons become massive after $SU(2)_X$ and $U(1)_{B-L}$ are spontaneously broken by scalar fields Φ_5 and S developing nonzero VEVs of v_Φ and v_S , respectively, with $v_S \gg v_\Phi$. The stability of the DM candidate is guaranteed by a residual Z_2 symmetry that is a subgroup of $SU(2)_X$. At tree level, the dark gauge boson X can interact with SM fermions by exchanging new gauge bosons $Z_{L,H}$ which arise from linear combinations of $SU(2)_X$ and $U(1)_{B-L}$ gauge fields. The XX^\dagger pair annihilates into SM fermions by exchanging $Z_{L,H}$ in the s channel. Since the DM is a gauge boson, its mass can be related to the masses of other gauge bosons in the model. The relation $m_{Z_L} \simeq 2m_X$ emerges when $SU(2)_X$ is broken by the VEV of the $SU(2)_X$ scalar five-plet Φ_5 , naturally leading to resonant pair annihilation of XX^\dagger via Z_L exchange. This model also supplies light neutrino masses with the aid of right-handed neutrinos whose mass terms are generated when $U(1)_{B-L}$ is spontaneously broken by the VEV of S , which activates the type-I seesaw mechanism.

We considered the values of the new gauge couplings g_X and g_{B-L} in the case where they are equal subject to constraints from collider data and the observed relic density. Assuming that $R_v = v_\Phi^2/v_S^2 \ll 1$ and $m_X \leq 1$ TeV, we obtain no parameter space which survives these experimental restrictions for $R_v \lesssim 10^{-4}$, but the $m_X \gtrsim 400$ (220) GeV region with $\mathcal{O}(1)$ couplings is still allowed for $R_v = 10^{-2}$ (10^{-3}). We also find that the corresponding values of v_S are between 5 and 10 TeV, implying that our model prefers the TeV-scale type-I seesaw scenario. Subsequently, we explored the X -nucleon elastic scattering cross-section, σ_{el}^N , for the surviving parameter regions and compared it with data from the latest DM direct detection experiments. The resulting σ_{el}^N largely evades the most recent XENON100 and LUX limits and will be tested more strictly by future DM direct searches such as XENON1T.

Finally, we schematically discussed some of the phenomenology of the new particles at the LHC. The DM particle can be produced as XX^\dagger or $X^{(\dagger)}\phi_i^{\text{odd}}$, where ϕ_i^{odd} is a Z_2^X -odd scalar boson. The signals of these production processes would be missing energy plus jets/photons and missing energy plus charged lepton pairs, respectively. The new scalar bosons can also be produced as $\phi^{\text{even(odd)}}\phi^{\text{even(odd)}}$ or $\phi^{\text{odd}}X$ ($\phi^{\text{even}}Z_{L,H}$), and the signals for ϕ^{even} and ϕ^{odd} would be two pairs of charged leptons and missing energy plus charged leptons, respectively. Their production cross-sections tend to be suppressed due to the small $|\theta|$ value and/or the heavy Z_H mass. Nevertheless, our model would be testable with an appropriate luminosity in the future. To perform a more

detailed study would require specifying the self-couplings in the scalar potential and thus the mass spectrum of the scalar bosons.

Acknowledgments

This research was supported in part by the National Science Council of Taiwan, R.O.C., under Grants No. NSC-100-2628-M-008-003-MY4 and No. NSC-100-2811-M-002-090 and by the MOE Academic Excellence Program under Grant No. 102R891505.

Appendix A: Feynman rules for new gauge interactions

The couplings of the fermion f in the model with the $Z_{L,H}$ bosons come from the $U(1)_{B-L}$ gauge interaction of f described by

$$\mathcal{L}' = -g_{B-L} \bar{f} \gamma^\mu \left(\hat{L}_f^E P_L + \hat{R}_f^E P_R \right) f E_\mu, \quad (\text{A1})$$

where $P_{L,R} = \frac{1}{2}(1 \mp \gamma_5)$ and \hat{L}_f^E (\hat{R}_f^E) is the $B-L$ value for $f_{L(R)}$. Since $E = Z_L \sin \theta + Z_H \cos \theta$, this leads to the Feynman rules

$$\bar{f} f Z_L^\mu : -ig_{B-L} \sin \theta \left(\hat{V}_f^E + \hat{A}_f^E \gamma_5 \right) \gamma^\mu, \quad (\text{A2})$$

$$\bar{f} f Z_H^\mu : -ig_{B-L} \cos \theta \left(\hat{V}_f^E + \hat{A}_f^E \gamma_5 \right) \gamma^\mu, \quad (\text{A3})$$

where $2\hat{V}_f^E = \hat{L}_f^E + \hat{R}_f^E$ and $2\hat{A}_f^E = \hat{L}_f^E - \hat{R}_f^E$. From the kinetic term of the C_k gauge bosons, $-\frac{1}{4}C_{k\mu\nu}C_k^{\mu\nu}$, where $C_k^{\mu\nu} = \partial^\mu C_k^\nu - \partial^\nu C_k^\mu - g_X \epsilon_{krs} C_r^\mu C_s^\nu$, with $C_3 = Z_L \cos \theta - Z_H \sin \theta$, we derive the Feynman rules

$$X^\mu X^{\nu\dagger} Z_L^\rho : -ig_X \cos \theta \left[(p_X^\rho - p_{X^\dagger}^\rho) g^{\mu\nu} + (p_{X^\dagger}^\mu - p_{Z_L}^\mu) g^{\nu\rho} + (p_{Z_L}^\nu - p_X^\nu) g^{\mu\rho} \right], \quad (\text{A4})$$

$$X^\mu X^{\nu\dagger} Z_H^\rho : -ig_X \sin \theta \left[(p_X^\rho - p_{X^\dagger}^\rho) g^{\mu\nu} + (p_{X^\dagger}^\mu - p_{Z_H}^\mu) g^{\nu\rho} + (p_{Z_H}^\nu - p_X^\nu) g^{\mu\rho} \right], \quad (\text{A5})$$

$$X_\mu X_\nu X_\rho^\dagger X_\sigma^\dagger : ig_X^2 (g_{\mu\nu} g_{\rho\sigma} - g_{\mu\rho} g_{\nu\sigma}), \quad (\text{A6})$$

$$X_\mu X_\nu^\dagger Z_{L\rho} Z_{L\sigma} : ig_X^2 \cos^2 \theta (g_{\mu\nu} g_{\rho\sigma} - g_{\mu\rho} g_{\nu\sigma}), \quad (\text{A7})$$

$$X_\mu X_\nu^\dagger Z_{H\rho} Z_{H\sigma} : ig_X^2 \sin^2 \theta (g_{\mu\nu} g_{\rho\sigma} - g_{\mu\rho} g_{\nu\sigma}), \quad (\text{A8})$$

$$X_\mu X_\nu^\dagger Z_{L\rho} Z_{H\sigma} : 2ig_X^2 \sin \theta \cos \theta (g_{\mu\nu} g_{\rho\sigma} - g_{\mu\rho} g_{\nu\sigma}), \quad (\text{A9})$$

where the momenta are all incoming.

Appendix B: Kinetic mixing between $U(1)_Y$ and $U(1)_{B-L}$

In the gauge $SU(2)_L \times U(1)_Y \times SU(2)_X \times U(1)_{B-L}$ sector of our model, the gauge bosons that may undergo mixing are W_3 , \mathcal{B} , C_3 , and E , respectively. We can express the Lagrangian for the kinetic

and mass terms of these particles after electroweak symmetry breaking as

$$\begin{aligned}\mathcal{L}_G &= -\frac{1}{4}W_3^{\alpha\omega}W_{3\alpha\omega} - \frac{1}{4}\mathcal{B}^{\alpha\omega}\mathcal{B}_{\alpha\omega} - \frac{1}{4}C_3^{\alpha\omega}C_{3\alpha\omega} - \frac{1}{4}E^{\alpha\omega}E_{\alpha\omega} - \frac{1}{2}\sin\chi\mathcal{B}^{\alpha\omega}E_{\alpha\omega} \\ &\quad + \frac{1}{2}m_W^2W_3^2 + \frac{1}{2}m_{\mathcal{B}}^2\mathcal{B}^2 - m_W m_{\mathcal{B}}W_3^\alpha\mathcal{B}_\alpha + \frac{1}{2}m_{C_3}^2C_3^2 + \frac{1}{2}m_E^2E^2 + m_{C_3}\mu_E C_3^\alpha E_\alpha \\ &= -\frac{1}{4}G_{\alpha\omega}^\top K G^{\alpha\omega} + \frac{1}{2}G_\alpha^\top M_G^2 G^\alpha,\end{aligned}\quad (\text{B1})$$

where $f_{\alpha\omega} = \partial_\alpha f_\omega - \partial_\omega f_\alpha$, the $\sin\chi$ term describes kinetic mixing between the hypercharge and B - L gauge bosons, \mathcal{B} and E , respectively,

$$m_W = \frac{1}{2}g_L v_H, \quad m_{\mathcal{B}} = \frac{1}{2}g_Y v_H, \quad (\text{B2})$$

$$m_{C_3} = 2g_X v_\Phi, \quad m_E^2 = 4g_{B-L}^2 v_S^2 + \mu_E^2, \quad \mu_E = 2g_{B-L} v_\Phi, \quad (\text{B3})$$

$$G = \begin{pmatrix} \mathcal{B} \\ W_3 \\ E \\ C_3 \end{pmatrix}, \quad K = \begin{pmatrix} 1 & 0 & s_\chi & 0 \\ 0 & 1 & 0 & 0 \\ s_\chi & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad M_G^2 = \begin{pmatrix} m_{\mathcal{B}}^2 & -m_{\mathcal{B}}m_W & 0 & 0 \\ -m_{\mathcal{B}}m_W & m_W^2 & 0 & 0 \\ 0 & 0 & m_E^2 & \mu_E m_{C_3} \\ 0 & 0 & \mu_E m_{C_3} & m_{C_3}^2 \end{pmatrix}, \quad (\text{B4})$$

with $s_\chi = \sin\chi$ and g_L and g_Y being the $SU(2)_L \times U(1)_Y$ gauge couplings, respectively. In \mathcal{L}_G we have included the s_χ term because it does not violate any of the symmetries in the theory, implying that in general s_χ can receive both tree- and loop-level contributions [28, 29], the latter being due to the SM fermions carrying both the $U(1)_Y$ and $U(1)_{B-L}$ charges.

It is straightforward to demonstrate that one can convert the kinetic part of \mathcal{L}_G into the canonical form, $-\frac{1}{4}\hat{G}_{\alpha\omega}^\top \hat{G}^{\alpha\omega}$, and diagonalize the M_G^2 matrix by making the transformation

$$G = \tilde{T} O_w \mathcal{O}_z \begin{pmatrix} A \\ Z \\ Z_H \\ Z_L \end{pmatrix}, \quad (\text{B5})$$

where A , Z , Z_H , and Z_L are the mass eigenstates, the photon A staying massless,

$$\tilde{T} = \begin{pmatrix} 1 & 0 & -t_\chi & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1/c_\chi & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad O_w = \begin{pmatrix} c_w & -s_w & 0 & 0 \\ s_w & c_w & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \mathcal{O}_z = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \mathcal{O}_{11} & \mathcal{O}_{12} & \mathcal{O}_{13} \\ 0 & \mathcal{O}_{21} & \mathcal{O}_{22} & \mathcal{O}_{23} \\ 0 & \mathcal{O}_{31} & \mathcal{O}_{32} & \mathcal{O}_{33} \end{pmatrix}, \quad (\text{B6})$$

$$c_\chi = \cos\chi, \quad t_\chi = \tan\chi, \quad c_w = \cos\theta_W, \quad s_w = \sin\theta_W = \frac{c_w m_{\mathcal{B}}}{m_W}. \quad (\text{B7})$$

The O_w and \mathcal{O}_z matrices are orthogonal, while \tilde{T} is not. The elements \mathcal{O}_{ij} encode the effect of the kinetic mixing, such that in its absence they are given by $\mathcal{O}_{11} = 1$, $\mathcal{O}_{12} = \mathcal{O}_{21} = \mathcal{O}_{13} = \mathcal{O}_{31} = 0$, $\mathcal{O}_{22} = \mathcal{O}_{33} = \cos\theta$, and $\mathcal{O}_{23} = -\mathcal{O}_{32} = \sin\theta$, which is the case treated in the main text.

It is also simple to see that the presence of kinetic mixing, $s_\chi \neq 0$, affects all of the couplings of Z and $Z_{L,H}$ to fermions. Since the fermions do not couple directly to the C_3 gauge boson, one can

write the Lagrangian for their interactions with the W_3 , \mathcal{B} , and E bosons in terms of the physical states A , Z , and $Z_{H,L}$ as

$$\begin{aligned}
\mathcal{L}' &= -g_L J_3^\lambda W_{3\lambda} - g_Y J_Y^\lambda \mathcal{B}_\lambda - g_{B-L} J_{B-L}^\lambda E_\lambda \\
&= -e J_{\text{em}}^\lambda A_\lambda - \left[(\mathcal{O}_{11} + \mathcal{O}_{21} t_\chi s_w) \hat{g}_Z \hat{J}_Z^\lambda + \frac{\mathcal{O}_{21}}{c_\chi} g_{B-L} J_{B-L}^\lambda - \mathcal{O}_{21} t_\chi c_w e J_{\text{em}}^\lambda \right] Z_\lambda \\
&\quad - \left[\frac{\mathcal{O}_{22}}{c_\chi} g_{B-L} J_{B-L}^\lambda + (\mathcal{O}_{12} + \mathcal{O}_{22} t_\chi s_w) \hat{g}_Z \hat{J}_Z^\lambda - \mathcal{O}_{22} t_\chi c_w e J_{\text{em}}^\lambda \right] Z_{H\lambda} \\
&\quad - \left[\frac{\mathcal{O}_{23}}{c_\chi} g_{B-L} J_{B-L}^\lambda + (\mathcal{O}_{13} + \mathcal{O}_{23} t_\chi s_w) \hat{g}_Z \hat{J}_Z^\lambda - \mathcal{O}_{23} t_\chi c_w e J_{\text{em}}^\lambda \right] Z_{L\lambda} , \tag{B8}
\end{aligned}$$

where $J_{3,Y,B-L}$ are the currents coupled to the respective fields and we have used the relations

$$e = g_L s_w = g_Y c_w, \quad J_{\text{em}} = J_3 + J_Y, \quad \hat{g}_Z \hat{J}_Z = c_w g_L J_3 - s_w g_Y J_Y. \tag{B9}$$

From the previous paragraphs, one can infer that the χ -dependent new terms translate into modifications to Z -pole observables and the $e^+e^- \rightarrow f\bar{f}$ cross-sections, as well as the Z and $Z_{H,L}$ masses. Consequently, such contributions must respect the pertinent experimental restrictions. After imposing them, we find that for the ranges of the new gauge couplings and Z_L mass satisfying the relic data requirement the kinetic-mixing effects are unimportant on the Z and Z_L masses, but for $\sin \chi$ not much less than 1 they could enlarge the Z_H mass substantially compared to that in the $\chi = 0$ case. Specifically, the increase in m_{Z_H} would be mild, no more than about 15%, if $|\sin \chi| \lesssim 0.5$.

To see if such mixing size is reasonable, we consider the two-point polarization diagram for the \mathcal{B} and E gauge bosons with fermions in the loop. Accordingly, we estimate the kinetic mixing parameter to be [28, 30]

$$\sin \chi \simeq - \sum_f \frac{g_{B-L} g_Y}{24\pi^2} (B_f - L_f) Y_f \ln \frac{|q^2|}{\Lambda^2}, \tag{B10}$$

where the sum is over the SM chiral fermions, $B_f - L_f$ and Y_f denote the $B - L$ and hypercharge values for fermion f , respectively, q is the renormalization scale which we take to be the $U(1)_{B-L}$ breaking scale, of order 1 TeV or greater, and we have applied the renormalization condition that at some higher scale Λ the sum of the loop and counterterm contributions vanishes. We note that one could evaluate $\sin \chi$ more precisely using the renormalization group equation to resum the large logarithms [30, 31], but the difference would amount to only a few percent for our scales of interest and therefore can be ignored. Thus, since $\sum_f (B_f - L_f) Y_f = 8$, taking $\Lambda^2 \sim 10^6 |q^2|$ in Eq. (B10) we get $\sin \chi \sim 0.16 g_{B-L}$. For the viable g_{B-L} values we have obtained, this result is compatible with the $|\sin \chi|$ number quoted in the last paragraph.

We remark that this is also consistent with the findings of a detailed analysis in Ref. [32] on the phenomenological constraints on a new massive Abelian gauge boson. The effects of such a particle can be compared to those of Z_H which contains mostly its $U(1)_{B-L}$ component E and has a mass of $\mathcal{O}(1-10)$ TeV in our study. For a new massive Abelian gauge boson in this mass range, the results of

Ref. [32] imply $|\sin \chi| \lesssim 0.6-1$. They also do not lead to additional restraints on the corresponding viable values of g_{B-L} .

-
- [1] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).
 - [2] C.W. Chiang, T. Nomura, and J. Tandean, Phys. Rev. D **87**, 073004 (2013) [arXiv:1205.6416 [hep-ph]].
 - [3] L.M. Krauss and F. Wilczek, Phys. Rev. Lett. **62**, 1221 (1989).
 - [4] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; Prog. Theor. Phys. **64**, 1103 (1980); M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979), p. 315; S.L. Glashow, in *Proceedings of the 1979 Cargese Summer Institute on Quarks and Leptons*, edited by M. Levy *et al.* (Plenum Press, New York, 1980), p. 687; R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980); J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); Phys. Rev. D **25**, 774 (1982).
 - [5] T. Hambye, JHEP **0901**, 028 (2009) [arXiv:0811.0172 [hep-ph]]; T. Hambye and M.H.G. Tytgat, Phys. Lett. B **683**, 39 (2010) [arXiv:0907.1007 [hep-ph]]; F. Chen, J.M. Cline, and A.R. Frey, Phys. Rev. D **80**, 083516 (2009) [arXiv:0907.4746 [hep-ph]]; J.L. Diaz-Cruz and E. Ma, Phys. Lett. B **695**, 264 (2011) [arXiv:1007.2631 [hep-ph]]; S. Bhattacharya, J.L. Diaz-Cruz, E. Ma, and D. Wegman, Phys. Rev. D **85**, 055008 (2012) [arXiv:1107.2093 [hep-ph]]; O. Lebedev, H.M. Lee, and Y. Mambrini, Phys. Lett. B **707**, 570 (2012) [arXiv:1111.4482 [hep-ph]]; Y. Farzan and A.R. Akbarieh, JCAP **1210**, 026 (2012) [arXiv:1207.4272 [hep-ph]]; T. Abe, M. Kakizaki, S. Matsumoto, and O. Seto, Phys. Lett. B **713**, 211 (2012) [arXiv:1202.5902 [hep-ph]]; F. D'Eramo, M. McCullough, and J. Thaler, JCAP **1304**, 030 (2013) [arXiv:1210.7817 [hep-ph]].
 - [6] ALEPH Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, and the LEP Electroweak Working Group (The LEP Collaborations), arXiv:hep-ex/0612034.
 - [7] C.W. Chiang, Y.F. Lin, and J. Tandean, JHEP **1111**, 083 (2011) [arXiv:1108.3969 [hep-ph]].
 - [8] CMS Collaboration, Report No. CMS-PAS-EWK-11-007, April 2012.
 - [9] C.W. Chiang, N.D. Christensen, G.J. Ding, and T. Han, Phys. Rev. D **85**, 015023 (2012) [arXiv:1107.5830 [hep-ph]].
 - [10] A. Pukhov *et al.*, arXiv:hep-ph/9908288.
 - [11] E.W. Kolb and M. Turner, *The Early Universe* (Westview Press, Boulder, 1990); K. Griest and D. Seckel, Phys. Rev. D **43**, 3191 (1991).
 - [12] P. Gondolo and G. Gelmini, Nucl. Phys. B **360**, 145 (1991).
 - [13] P.A.R. Ade *et al.* [Planck Collaboration], arXiv:1303.5062 [astro-ph.CO].
 - [14] D.B. Kaplan and A. Manohar, Nucl. Phys. B **310**, 527 (1988).
 - [15] J. Angle *et al.* [XENON10 Collaboration], Phys. Rev. Lett. **107**, 051301 (2011) [arXiv:1104.3088 [astro-ph.CO]]; **110**, 249901(E) (2013).
 - [16] M.T. Frandsen, F. Kahlhoefer, C. McCabe, S. Sarkar, and K. Schmidt-Hoberg, JCAP **1307**, 023 (2013) [arXiv:1304.6066 [hep-ph]].
 - [17] E. Aprile *et al.* [XENON100 Collaboration], Phys. Rev. Lett. **109**, 181301 (2012) [arXiv:1207.5988 [astro-ph.CO]].
 - [18] Z. Ahmed *et al.* [CDMS-II Collaboration], Science **327**, 1619 (2010) [arXiv:0912.3592 [astro-ph.CO]]; D.S. Akerib *et al.* [CDMS Collaboration], Phys. Rev. D **82**, 122004 (2010) [arXiv:1010.4290]

- [astro-ph.CO]]; Z. Ahmed *et al.* [CDMS-II Collaboration], Phys. Rev. Lett. **106**, 131302 (2011) [arXiv:1011.2482 [astro-ph.CO]].
- [19] R. Agnese *et al.* [CDMS Collaboration], Phys. Rev. D **88**, 031104 (2013) [arXiv:1304.3706 [astro-ph.CO]].
- [20] D.S. Akerib *et al.* [LUX Collaboration], arXiv:1310.8214 [astro-ph.CO].
- [21] C.E. Aalseth *et al.* [CoGeNT Collaboration], arXiv:1208.5737 [astro-ph.CO]; Phys. Rev. Lett. **106**, 131301 (2011) [arXiv:1002.4703 [astro-ph.CO]]; Phys. Rev. Lett. **107**, 141301 (2011) [arXiv:1106.0650 [astro-ph.CO]].
- [22] C. Savage, G. Gelmini, P. Gondolo, and K. Freese, JCAP **0904**, 010 (2009) [arXiv:0808.3607 [astro-ph]]; R. Bernabei *et al.* [DAMA and LIBRA Collaborations], Eur. Phys. J. C **67**, 39 (2010) [arXiv:1002.1028 [astro-ph.GA]].
- [23] G. Angloher *et al.* [CRESST Collaboration], Eur. Phys. J. C **72**, 1971 (2012) arXiv:1109.0702 [astro-ph.CO].
- [24] R. Agnese *et al.* [CDMS Collaboration], arXiv:1304.4279 [hep-ex].
- [25] E. Aprile, arXiv:1206.6288 [astro-ph.IM].
- [26] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. D **87**, 012008 (2013) [arXiv:1208.0949 [hep-ex]].
- [27] G. Steigman, B. Dasgupta, and J.F. Beacom, Phys. Rev. D **86**, 023506 (2012) [arXiv:1204.3622 [hep-ph]].
- [28] B. Holdom, Phys. Lett. B **166**, 196 (1986)
- [29] F. del Aguila, G.D. Coughlan, and M. Quiros, Nucl. Phys. B **307**, 633 (1988) [Erratum-ibid. B **312**, 751 (1989)]; R. Foot and X.G. He, Phys. Lett. B **267**, 509 (1991); R. Foot, X.G. He, H. Lew, and R.R. Volkas, Phys. Rev. D **50**, 4571 (1994) [hep-ph/9401250].
- [30] K.R. Dienes, C. Kolda, and J. March-Russell, Nucl. Phys. B **492**, 104 (1997).
- [31] K.S. Babu, C. Kolda, and J. March-Russell, Phys. Rev. D **54**, 4635 (1996),
- [32] M. Williams, C.P. Burgess, A. Maharana, and F. Quevedo, JHEP **1108**, 106 (2011) [arXiv:1103.4556 [hep-ph]].