

A Solution of Simultaneous Recurrences

Mark Korenblit
 Holon Institute of Technology, Israel
 korenblit@hit.ac.il

Vadim E. Levit
 Ariel University, Israel
 levitv@ariel.ac.il

1. Introduction

There exist problems whose solving requires a solution of simultaneous recurrences, such as, for example, the following ones:

$$\begin{cases} a_x = 2a_{x-1} + 4b_{x-1} + 1 \\ b_x = a_{x-1} + 3b_{x-1} + 2c_{x-1} + 1 \\ c_x = 2b_{x-1} + 4c_{x-1} + 1, \end{cases} \quad (1.1)$$

$$\begin{cases} a_x = \frac{37}{6}a_{x-1} - \frac{1}{6}b_{x-1} + 2 \\ b_x = \frac{15}{2}a_{x-1} - \frac{7}{2}b_{x-1} + 2c_{x-1} + 2 \\ c_x = \frac{16}{3}a_{x-1} - \frac{10}{3}b_{x-1} + 4c_{x-1} + 2 \end{cases} \quad (1.2)$$

Besides, the problem of solving simultaneous recurrences is of interest itself, regardless of any particular application. Section 2 discusses transformation of matrix recurrences to regular recurrences. Section 3 describes a process of solving special matrix recurrences of order three (as (1.1) and (1.2)) by their decomposition to matrix recurrences of order two.

2. Transformation of Matrix Recurrences to Regular Recurrences

Matrix recurrences look as

$$\begin{cases} y_{1x} = \alpha_{11}y_{1x-1} + \alpha_{12}y_{2x-1} + \alpha_{13}y_{3x-1} + \dots + \alpha_{1n}y_{nx-1} + \alpha_1 \\ y_{2x} = \alpha_{21}y_{1x-1} + \alpha_{22}y_{2x-1} + \alpha_{23}y_{3x-1} + \dots + \alpha_{2n}y_{nx-1} + \alpha_2 \\ y_{3x} = \alpha_{31}y_{1x-1} + \alpha_{32}y_{2x-1} + \alpha_{33}y_{3x-1} + \dots + \alpha_{3n}y_{nx-1} + \alpha_3 \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ y_{nx} = \alpha_{n1}y_{1x-1} + \alpha_{n2}y_{2x-1} + \alpha_{n3}y_{3x-1} + \dots + \alpha_{nn}y_{nx-1} + \alpha_n. \end{cases} \quad (2.1)$$

Here x is a current number of variables y_1, y_2, \dots, y_n (α followed by an index denotes a constant coefficient) in their sequences generated by means of (2.1) so that $y_{1_0}, y_{2_0}, \dots, y_{n_0}$ are initial values of y_1, y_2, \dots, y_n , respectively. We intend to separate variables of these recurrences so that each variable y_i is expressed in previous values of y_i only, i.e.

$$y_{ix} = \beta_1 y_{i_{x-1}} + \beta_2 y_{i_{x-2}} + \beta_3 y_{i_{x-3}} + \dots \quad (2.2)$$

where β_j ($j = 1, 2, 3, \dots$) is a constant coefficient. Recurrence (2.2) is a *regular recurrence*. Methods for solving regular recurrences, providing explicit formulae, are presented in [3].

First, we solve the problem for the special case when there are no absolute terms ($\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$). In this case (2.1) is reduced to

$$\begin{cases} y_{1x} = \alpha_{11}y_{1x-1} + \alpha_{12}y_{2x-1} + \alpha_{13}y_{3x-1} + \dots + \alpha_{1n}y_{nx-1} \\ y_{2x} = \alpha_{21}y_{1x-1} + \alpha_{22}y_{2x-1} + \alpha_{23}y_{3x-1} + \dots + \alpha_{2n}y_{nx-1} \\ y_{3x} = \alpha_{31}y_{1x-1} + \alpha_{32}y_{2x-1} + \alpha_{33}y_{3x-1} + \dots + \alpha_{3n}y_{nx-1} \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ y_{nx} = \alpha_{n1}y_{1x-1} + \alpha_{n2}y_{2x-1} + \alpha_{n3}y_{3x-1} + \dots + \alpha_{nn}y_{nx-1}. \end{cases} \quad (2.3)$$

We present equations (2.3) in the matrix-vectorial form as

$$Y_x = AY_{x-1} \quad (2.4)$$

where

$$A = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \dots & \alpha_{2n} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} & \dots & \alpha_{3n} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{n1} & \alpha_{n2} & \alpha_{n3} & \dots & \alpha_{nn} \end{pmatrix}$$

and

$$Y = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ \cdot \\ y_n \end{pmatrix}.$$

We will use the following basic notions.

The square matrix

$$I = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}$$

is called the *unit matrix*. If A is a square matrix of order n and I is a unit matrix of order n , the λ -matrix

$$A - \lambda I = \begin{pmatrix} \alpha_{11} - \lambda & \alpha_{12} & \alpha_{13} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} - \lambda & \alpha_{23} & \dots & \alpha_{2n} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} - \lambda & \dots & \alpha_{3n} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{n1} & \alpha_{n2} & \alpha_{n3} & \dots & \alpha_{nn} - \lambda \end{pmatrix}$$

is called the *characteristic matrix of A* and the determinant of A is called the *characteristic polynomial of A* .

By the *Hamilton-Cayley theorem* [1], [2], if A is a square matrix and $\phi(\lambda)$ its characteristic polynomial, then

$$\phi(A) = 0.$$

That is, if

$$\phi(\lambda) = u_0\lambda^n + u_1\lambda^{n-1} + u_2\lambda^{n-2} + \dots + u_{n-1}\lambda + u_n,$$

then

$$u_0A^n + u_1A^{n-1} + u_2A^{n-2} + \dots + u_{n-1}A + u_nI = 0.$$

As follows from the definition of the characteristic polynomial, $u_0 = (-1)^n$. Hence,

$$A^n = \beta_1A^{n-1} + \beta_2A^{n-2} + \dots + \beta_{n-1}A + \beta_nI \quad (2.5)$$

where $\beta_j = (-1)^{n+1} u_j$ ($j = 1, 2, \dots, n$). We multiply the left and the right parts of (3.1) by vector Y_{x-n} and get:

$$A^n Y_{x-n} = \beta_1 A^{n-1} Y_{x-n} + \beta_2 A^{n-2} Y_{x-n} + \dots + \beta_{n-1} A Y_{x-n} + \beta_n Y_{x-n}. \quad (2.6)$$

On the other hand, iterating recurrence (2.4) gives

$$Y_x = A Y_{x-1} = A^2 Y_{x-2} = \dots = A^{n-1} Y_{x-n+1} = A^n Y_{x-n}.$$

After substituting these results in (2.6) we have:

$$Y_x = \beta_1 Y_{x-1} + \beta_2 Y_{x-2} + \dots + \beta_{n-1} Y_{x-n+1} + \beta_n Y_{x-n}.$$

That is,

$$\begin{aligned} y_{1x} &= \beta_1 y_{1x-1} + \beta_2 y_{1x-2} + \dots + \beta_{n-1} y_{1x-n+1} + \beta_n y_{1x-n} \\ y_{2x} &= \beta_1 y_{2x-1} + \beta_2 y_{2x-2} + \dots + \beta_{n-1} y_{2x-n+1} + \beta_n y_{2x-n} \\ y_{3x} &= \beta_1 y_{3x-1} + \beta_2 y_{3x-2} + \dots + \beta_{n-1} y_{3x-n+1} + \beta_n y_{3x-n} \\ &\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ y_{nx} &= \beta_1 y_{nx-1} + \beta_2 y_{nx-2} + \dots + \beta_{n-1} y_{nx-n+1} + \beta_n y_{nx-n} \end{aligned} \quad (2.7)$$

where a coefficient β_j ($j = 1, 2, \dots, n$) equals a coefficient u_j near a term λ^{n-j} in a characteristic polynomial of matrix A for odd n and $-u_j$ for even n .

For example, for $n = 2$, when we have two the following recurrences:

$$\begin{cases} a_x = \alpha_{11} a_{x-1} + \alpha_{12} b_{x-1} \\ b_x = \alpha_{21} a_{x-1} + \alpha_{22} b_{x-1}, \end{cases}$$

the result is

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{22}) a_{x-1} + (\alpha_{12} \alpha_{21} - \alpha_{11} \alpha_{22}) a_{x-2} \\ b_x &= (\alpha_{11} + \alpha_{22}) b_{x-1} + (\alpha_{12} \alpha_{21} - \alpha_{11} \alpha_{22}) b_{x-2}. \end{aligned} \quad (2.8)$$

Now, move on to recurrences with absolute terms (2.1). There are $n + 1$ coefficients in each recurrence. We add the trivial $n + 1$ equation to (2.1) and get:

$$\begin{cases} y_{1x} = \alpha_{11} y_{1x-1} + \alpha_{12} y_{2x-1} + \alpha_{13} y_{3x-1} + \dots + \alpha_{1n} y_{nx-1} + \alpha_1 \\ y_{2x} = \alpha_{21} y_{1x-1} + \alpha_{22} y_{2x-1} + \alpha_{23} y_{3x-1} + \dots + \alpha_{2n} y_{nx-1} + \alpha_2 \\ y_{3x} = \alpha_{31} y_{1x-1} + \alpha_{32} y_{2x-1} + \alpha_{33} y_{3x-1} + \dots + \alpha_{3n} y_{nx-1} + \alpha_3 \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ y_{nx} = \alpha_{n1} y_{1x-1} + \alpha_{n2} y_{2x-1} + \alpha_{n3} y_{3x-1} + \dots + \alpha_{nn} y_{nx-1} + \alpha_n \\ 1 = 0 \quad + 0 \quad + 0 \quad + \dots + 0 \quad + 1. \end{cases} \quad (2.9)$$

Now, matrix A^* of coefficients looks as

$$A^* = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \dots & \alpha_{1n} & \alpha_1 \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \dots & \alpha_{2n} & \alpha_2 \\ \alpha_{31} & \alpha_{32} & \alpha_{33} & \dots & \alpha_{3n} & \alpha_3 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{n1} & \alpha_{n2} & \alpha_{n3} & \dots & \alpha_{nn} & \alpha_n \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}$$

and vector Y^* of variables is presented as

$$Y^* = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ \cdot \\ y_n \\ 1 \end{pmatrix}.$$

In such a case, equations (2.9) are presented in the matrix-vectorial form as

$$Y_x^* = A^* Y_{x-1}^*. \quad (2.10)$$

The characteristic polynomial of matrix A^* looks as

$$\begin{aligned} \phi^*(\lambda) &= \begin{vmatrix} \alpha_{11} - \lambda & \alpha_{12} & \alpha_{13} & \dots & \alpha_{1n} & \alpha_1 \\ \alpha_{21} & \alpha_{22} - \lambda & \alpha_{23} & \dots & \alpha_{2n} & \alpha_2 \\ \alpha_{31} & \alpha_{32} & \alpha_{33} - \lambda & \dots & \alpha_{3n} & \alpha_3 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{n1} & \alpha_{n2} & \alpha_{n3} & \dots & \alpha_{nn} - \lambda & \alpha_n \\ 0 & 0 & 0 & \dots & 0 & 1 - \lambda \end{vmatrix} \\ &= (-1)^{2n+2} (1 - \lambda) \begin{vmatrix} \alpha_{11} - \lambda & \alpha_{12} & \alpha_{13} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} - \lambda & \alpha_{23} & \dots & \alpha_{2n} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} - \lambda & \dots & \alpha_{3n} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{n1} & \alpha_{n2} & \alpha_{n3} & \dots & \alpha_{nn} - \lambda \end{vmatrix} \\ &= (1 - \lambda) \phi(\lambda) \end{aligned} \quad (2.11)$$

(we apply an *expansion of determinant by minors* [1]).

By the Hamilton-Cayley theorem , if

$$\phi^*(\lambda) = u_0^* \lambda^{n+1} + u_1^* \lambda^n + u_2^* \lambda^{n-1} + \dots + u_n^* \lambda + u_{n+1}^* ,$$

then

$$u_0^* (A^*)^{n+1} + u_1^* (A^*)^n + u_2^* (A^*)^{n-1} + \dots + u_n^* A^* + u_{n+1}^* I = 0.$$

As follows from 2.11, $u_0^* = (-1)^{n+1}$. Hence,

$$(A^*)^{n+1} = \beta_1^* (A^*)^n + \beta_2^* (A^*)^{n-1} + \dots + \beta_n^* A^* + \beta_{n+1}^* I \quad (2.12)$$

where $\beta_j^* = (-1)^n u_j^*$ ($j = 1, 2, \dots, n, n+1$). We multiply the left and the right parts of (2.12) by vector Y_{x-n-1}^* and get:

$$\begin{aligned} (A^*)^{n+1} Y_{x-n-1}^* &= \beta_1^* (A^*)^n Y_{x-n-1}^* + \beta_2^* (A^*)^{n-1} Y_{x-n-1}^* + \dots \\ &\quad + \beta_n^* A^* Y_{x-n-1}^* + \beta_{n+1}^* Y_{x-n-1}^*. \end{aligned} \quad (2.13)$$

Iterating recurrence 2.10 gives

$$Y_x^* = A^* Y_{x-1}^* = (A^*)^2 Y_{x-2}^* = \dots = (A^*)^n Y_{x-n}^* = (A^*)^{n+1} Y_{x-n-1}^*$$

and after substituting these results in (2.13) we have:

$$Y_x^* = \beta_1^* Y_{x-1}^* + \beta_2^* Y_{x-2}^* + \dots + \beta_n^* Y_{x-n}^* + \beta_{n+1}^* Y_{x-n-1}^*.$$

That is,

$$\begin{aligned} y_{1x} &= \beta_1^* y_{1x-1} + \beta_2^* y_{1x-2} + \dots + \beta_n^* y_{1x-n} + \beta_{n+1}^* y_{1x-n-1} \\ y_{2x} &= \beta_1^* y_{2x-1} + \beta_2^* y_{2x-2} + \dots + \beta_n^* y_{2x-n} + \beta_{n+1}^* y_{2x-n-1} \\ y_{3x} &= \beta_1^* y_{3x-1} + \beta_2^* y_{3x-2} + \dots + \beta_n^* y_{3x-n} + \beta_{n+1}^* y_{3x-n-1} \\ &\quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ y_{nx} &= \beta_1^* y_{nx-1} + \beta_2^* y_{nx-2} + \dots + \beta_n^* y_{nx-n} + \beta_{n+1}^* y_{nx-n-1} \\ 1 &= \beta_1^* + \beta_2^* + \dots + \beta_n^* + \beta_{n+1}^* \end{aligned} \quad (2.14)$$

where a coefficient β_j^* ($j = 1, 2, \dots, n, n+1$) equals a coefficient u_j^* near a term λ^{n+1-j} in a characteristic polynomial of matrix A^* for even n and $-u_j$ for odd n .

For example, for $n = 1$, we have a recurrence

$$a_x = \alpha_{11} a_{x-1} + \alpha_1$$

that can be presented as

$$a_x = (\alpha_{11} + 1)a_{x-1} - \alpha_{11}a_{x-2} ;$$

for $n = 2$, we have two the following recurrences:

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_1 \\ b_x = \alpha_{21}a_{x-1} + \alpha_{22}b_{x-1} + \alpha_2 \end{cases} \quad (2.15)$$

and the result is

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{22} + 1)a_{x-1} + (\alpha_{12}\alpha_{21} - \alpha_{11}\alpha_{22} - \alpha_{11} - \alpha_{22})a_{x-2} + \\ &\quad (\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21})a_{x-3} \\ b_x &= (\alpha_{11} + \alpha_{22} + 1)b_{x-1} + (\alpha_{12}\alpha_{21} - \alpha_{11}\alpha_{22} - \alpha_{11} - \alpha_{22})b_{x-2} + \\ &\quad (\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21})b_{x-3}. \end{aligned}$$

Thus, we expressed each variable y_i ($i = 1, 2, \dots, n$) from (2.1) in its $n + 1$ previous values. A coefficient β_j^* near a value y_{i_x-j} ($j = 1, 2, \dots, n, n + 1$) does not depend on i , i.e., all recurrences (2.14) have the same structure. Absolute terms $\alpha_1, \alpha_2, \dots, \alpha_n$ do not appear explicitly in (2.14). They are accounted for implicitly, by addition of a supplementary $n + 1$ previous value of y_i in the expressions (compare (2.14) with ((2.7))).

Nevertheless, it is also of interest to express variables of (2.1) directly using absolute terms. Such a presentation is more elegant since it contains explicitly all the information about initial recurrences. Specifically, it can be deduced for $n = 2$ and $n = 3$ through a number of transformations.

Consider recurrences (2.15). As follows from the first equation,

$$\begin{aligned} \alpha_{12}b_{x-1} &= a_x - \alpha_{11}a_{x-1} - \alpha_1 \Leftrightarrow \\ b_{x-1} &= \frac{a_x}{\alpha_{12}} - \frac{\alpha_{11}}{\alpha_{12}}a_{x-1} - \frac{\alpha_1}{\alpha_{12}} \Leftrightarrow \end{aligned} \quad (2.16)$$

$$b_x = \frac{a_{x+1}}{\alpha_{12}} - \frac{\alpha_{11}}{\alpha_{12}}a_x - \frac{\alpha_1}{\alpha_{12}}. \quad (2.17)$$

We substitute (2.16) and (2.17) into the second equation (2.15) and get:

$$\begin{aligned} \frac{a_{x+1}}{\alpha_{12}} - \frac{\alpha_{11}}{\alpha_{12}}a_x - \frac{\alpha_1}{\alpha_{12}} &= \alpha_{21}a_{x-1} + \alpha_{22} \left(\frac{a_x}{\alpha_{12}} - \frac{\alpha_{11}}{\alpha_{12}}a_{x-1} - \frac{\alpha_1}{\alpha_{12}} \right) + \alpha_2 \Leftrightarrow \\ \frac{a_x}{\alpha_{12}} - \frac{\alpha_{11}}{\alpha_{12}}a_{x-1} - \frac{\alpha_1}{\alpha_{12}} &= \alpha_{21}a_{x-2} + \alpha_{22} \left(\frac{a_{x-1}}{\alpha_{12}} - \frac{\alpha_{11}}{\alpha_{12}}a_{x-2} - \frac{\alpha_1}{\alpha_{12}} \right) + \alpha_2 \Leftrightarrow \end{aligned}$$

$$a_x = (\alpha_{11} + \alpha_{22})a_{x-1} + (\alpha_{12}\alpha_{21} - \alpha_{11}\alpha_{22})a_{x-2} + \alpha_1(1 - \alpha_{22}) + \alpha_2\alpha_{12}.$$

The expression for b_x is derived in the same way and, finally, we have:

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{22})a_{x-1} + (\alpha_{12}\alpha_{21} - \alpha_{11}\alpha_{22})a_{x-2} + \\ &\quad \alpha_1(1 - \alpha_{22}) + \alpha_2\alpha_{12} \\ b_x &= (\alpha_{11} + \alpha_{22})b_{x-1} + (\alpha_{12}\alpha_{21} - \alpha_{11}\alpha_{22})b_{x-2} + \\ &\quad \alpha_2(1 - \alpha_{11}) + \alpha_1\alpha_{21} \end{aligned} \tag{2.18}$$

(compare with (2.8)).

For $n = 3$, recurrences (2.1) can be presented as three following equations:

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_{13}c_{x-1} + \alpha_1 \\ b_x = \alpha_{21}a_{x-1} + \alpha_{22}b_{x-1} + \alpha_{23}c_{x-1} + \alpha_2 \\ c_x = \alpha_{31}a_{x-1} + \alpha_{32}b_{x-1} + \alpha_{33}c_{x-1} + \alpha_3. \end{cases} \tag{2.19}$$

The derivation is similar to the derivation for $n = 2$ and the final result is the following:

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{22} + \alpha_{33})a_{x-1} + \\ &\quad (\alpha_{12}\alpha_{21} + \alpha_{13}\alpha_{31} + \alpha_{23}\alpha_{32} - \alpha_{11}\alpha_{22} - \alpha_{11}\alpha_{33} - \alpha_{22}\alpha_{33})a_{x-2} + \\ &\quad (\alpha_{11}\alpha_{22}\alpha_{33} + \alpha_{12}\alpha_{23}\alpha_{31} + \alpha_{21}\alpha_{13}\alpha_{32} - \alpha_{12}\alpha_{21}\alpha_{33} - \alpha_{13}\alpha_{31}\alpha_{22} - \\ &\quad \alpha_{23}\alpha_{32}\alpha_{11})a_{x-3} + \\ &\quad \alpha_1(1 - \alpha_{22} - \alpha_{33}) + \alpha_2\alpha_{12} + \alpha_3\alpha_{13} + \alpha_1(\alpha_{22}\alpha_{33} - \alpha_{23}\alpha_{32}) + \\ &\quad \alpha_2(\alpha_{13}\alpha_{32} - \alpha_{33}\alpha_{12}) + \alpha_3(\alpha_{12}\alpha_{23} - \alpha_{22}\alpha_{13}) \\ b_x &= (\alpha_{11} + \alpha_{22} + \alpha_{33})b_{x-1} + \\ &\quad (\alpha_{12}\alpha_{21} + \alpha_{13}\alpha_{31} + \alpha_{23}\alpha_{32} - \alpha_{11}\alpha_{22} - \alpha_{11}\alpha_{33} - \alpha_{22}\alpha_{33})b_{x-2} + \\ &\quad (\alpha_{11}\alpha_{22}\alpha_{33} + \alpha_{12}\alpha_{23}\alpha_{31} + \alpha_{21}\alpha_{13}\alpha_{32} - \alpha_{12}\alpha_{21}\alpha_{33} - \alpha_{13}\alpha_{31}\alpha_{22} - \\ &\quad \alpha_{23}\alpha_{32}\alpha_{11})b_{x-3} + \\ &\quad \alpha_2(1 - \alpha_{11} - \alpha_{33}) + \alpha_1\alpha_{21} + \alpha_3\alpha_{23} + \alpha_2(\alpha_{11}\alpha_{33} - \alpha_{13}\alpha_{31}) + \\ &\quad \alpha_3(\alpha_{21}\alpha_{13} - \alpha_{11}\alpha_{23}) + \alpha_1(\alpha_{23}\alpha_{31} - \alpha_{33}\alpha_{21}) \\ c_x &= (\alpha_{11} + \alpha_{22} + \alpha_{33})c_{x-1} + \\ &\quad (\alpha_{12}\alpha_{21} + \alpha_{13}\alpha_{31} + \alpha_{23}\alpha_{32} - \alpha_{11}\alpha_{22} - \alpha_{11}\alpha_{33} - \alpha_{22}\alpha_{33})c_{x-2} + \\ &\quad (\alpha_{11}\alpha_{22}\alpha_{33} + \alpha_{12}\alpha_{23}\alpha_{31} + \alpha_{21}\alpha_{13}\alpha_{32} - \alpha_{12}\alpha_{21}\alpha_{33} - \alpha_{13}\alpha_{31}\alpha_{22} - \\ &\quad \alpha_{23}\alpha_{32}\alpha_{11})c_{x-3} + \\ &\quad \alpha_3(1 - \alpha_{11} - \alpha_{22}) + \alpha_1\alpha_{31} + \alpha_2\alpha_{32} + \alpha_3(\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21}) + \\ &\quad \alpha_1(\alpha_{32}\alpha_{21} - \alpha_{22}\alpha_{31}) + \alpha_2(\alpha_{31}\alpha_{12} - \alpha_{11}\alpha_{32}). \end{aligned} \tag{2.20}$$

Hence, each variable y_i ($i = 1, 2, \dots, n$) from (2.1) is expressed in its n previous values for $n = 2, 3$ if absolute terms appear explicitly in expressions. As follows from (2.18) and (2.20) the structure of absolute terms in these expressions depends on i .

3. A Solution of Matrix Recurrences of Order Three by their Decomposition to Matrix Recurrences of Order Two

We intend to solve three simultaneous recurrences (2.19), where x is a current number of variables a , b , and c in their sequences generated by means of (2.19) so that a_0 , b_0 , and c_0 are initial values of a , b , and c , respectively. That is, we intend to express a_x , b_x , and c_x of (2.19) directly in x .

It would be possible to solve this problem by the general method described in the previous section and then to apply the technique of solving regular recurrences from [3]. However, under certain conditions, there is a simpler way of arriving at a solution. This way is decomposition of (2.19) to matrix recurrences of order two.

Matrix recurrences of order two can also be solved by the general method. However, we use another way that seems to be more efficient under the special conditions.

Lemma 3.1. *If*

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_1 \\ b_x = \alpha_{21}a_{x-1} + \alpha_{22}b_{x-1} + \alpha_2 \end{cases}$$

and

$$\alpha_{11} + \alpha_{12} = \alpha_{21} + \alpha_{22} , \tag{3.1}$$

then

1. When $\alpha_{12} \neq -\alpha_{21}$, $\alpha_{11} + \alpha_{12} \neq 1$, $\alpha_{11} - \alpha_{21} \neq 1$

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{12})^x a_0 + \alpha_{12} (b_0 - a_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \\ &\quad \alpha_1 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{12} (\alpha_2 - \alpha_1)}{\alpha_{12} + \alpha_{21}} \times \\ &\quad \left(\frac{(\alpha_{11} + \alpha_{12})^x - \alpha_{11} - \alpha_{12}}{\alpha_{11} + \alpha_{12} - 1} - \frac{(\alpha_{11} - \alpha_{21})^x - \alpha_{11} + \alpha_{21}}{\alpha_{11} - \alpha_{21} - 1} \right) \\ b_x &= (\alpha_{11} + \alpha_{12})^x b_0 + \alpha_{21} (a_0 - b_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \end{aligned}$$

$$\alpha_2 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{21} (\alpha_1 - \alpha_2)}{\alpha_{12} + \alpha_{21}} \times \left(\frac{(\alpha_{11} + \alpha_{12})^x - \alpha_{11} - \alpha_{12}}{\alpha_{11} + \alpha_{12} - 1} - \frac{(\alpha_{11} - \alpha_{21})^x - \alpha_{11} + \alpha_{21}}{\alpha_{11} - \alpha_{21} - 1} \right).$$

2. When $\alpha_{12} \neq -\alpha_{21}$, $\alpha_{11} + \alpha_{12} = 1$ ($\alpha_{11} - \alpha_{21} \neq 1$)

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{12})^x a_0 + \alpha_{12} (b_0 - a_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \\ &\quad \alpha_1 x + \frac{\alpha_{12} (\alpha_2 - \alpha_1)}{\alpha_{11} - \alpha_{21} - 1} \left(\frac{(\alpha_{11} - \alpha_{21})^x - \alpha_{11} + \alpha_{21}}{\alpha_{11} - \alpha_{21} - 1} - x + 1 \right) \\ b_x &= (\alpha_{11} + \alpha_{12})^x b_0 + \alpha_{21} (a_0 - b_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \\ &\quad \alpha_2 x + \frac{\alpha_{21} (\alpha_1 - \alpha_2)}{\alpha_{11} - \alpha_{21} - 1} \left(\frac{(\alpha_{11} - \alpha_{21})^x - \alpha_{11} + \alpha_{21}}{\alpha_{11} - \alpha_{21} - 1} - x + 1 \right). \end{aligned}$$

3. When $\alpha_{12} \neq -\alpha_{21}$, $\alpha_{11} - \alpha_{21} = 1$ ($\alpha_{11} + \alpha_{12} \neq 1$)

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{12})^x a_0 + \alpha_{12} (b_0 - a_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \\ &\quad \alpha_1 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{12} (\alpha_2 - \alpha_1)}{\alpha_{11} + \alpha_{12} - 1} \left(\frac{(\alpha_{11} + \alpha_{12})^x - \alpha_{11} - \alpha_{12}}{\alpha_{11} + \alpha_{12} - 1} - x + 1 \right) \\ b_x &= (\alpha_{11} + \alpha_{12})^x b_0 + \alpha_{21} (a_0 - b_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \\ &\quad \alpha_2 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{21} (\alpha_1 - \alpha_2)}{\alpha_{11} + \alpha_{12} - 1} \left(\frac{(\alpha_{11} + \alpha_{12})^x - \alpha_{11} - \alpha_{12}}{\alpha_{11} + \alpha_{12} - 1} - x + 1 \right). \end{aligned}$$

4. When $\alpha_{12} = -\alpha_{21}$, $\alpha_{11} + \alpha_{12} \neq 1$ ($\alpha_{11} - \alpha_{21} \neq 1$)

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{12})^{x-1} ((\alpha_{11} + \alpha_{12}) a_0 + \alpha_{12} (b_0 - a_0) x) + \\ &\quad \alpha_1 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{12} (\alpha_2 - \alpha_1)}{(\alpha_{11} + \alpha_{12} - 1)^2} \times \\ &\quad ((x-1) (\alpha_{11} + \alpha_{12})^x - x (\alpha_{11} + \alpha_{12})^{x-1} + 1) \\ b_x &= (\alpha_{11} + \alpha_{12})^{x-1} ((\alpha_{11} + \alpha_{12}) b_0 + \alpha_{21} (a_0 - b_0) x) + \\ &\quad \alpha_2 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{21} (\alpha_1 - \alpha_2)}{(\alpha_{11} + \alpha_{12} - 1)^2} \times \end{aligned}$$

$$((x-1)(\alpha_{11} + \alpha_{12})^x - x(\alpha_{11} + \alpha_{12})^{x-1} + 1).$$

5. When $\alpha_{12} = -\alpha_{21}$, $\alpha_{11} + \alpha_{12} = 1$ ($\alpha_{11} - \alpha_{21} = 1$)

$$\begin{aligned} a_x &= (\alpha_{11} + \alpha_{12})^{x-1} ((\alpha_{11} + \alpha_{12})a_0 + \alpha_{12}(b_0 - a_0)x) + \\ &\quad \alpha_1 x + \alpha_{12}(\alpha_2 - \alpha_1) \frac{x(x-1)}{2} \\ b_x &= (\alpha_{11} + \alpha_{12})^{x-1} ((\alpha_{11} + \alpha_{12})b_0 + \alpha_{21}(a_0 - b_0)x) + \\ &\quad \alpha_2 x + \alpha_{21}(\alpha_1 - \alpha_2) \frac{x(x-1)}{2}. \end{aligned}$$

Proof. 1. Denote $\Delta_x = b_x - a_x$, $S = \alpha_{11} + \alpha_{12}$, $D = \alpha_{11} - \alpha_{21}$, $\delta = \alpha_2 - \alpha_1$. In such a case,

$$\begin{aligned} a_x &= \alpha_{11}a_{x-1} + \alpha_{12}(a_{x-1} + \Delta_{x-1}) + \alpha_1 \\ &= (\alpha_{11} + \alpha_{12})a_{x-1} + \alpha_{12}\Delta_{x-1} + \alpha_1 \\ &= Sa_{x-1} + \alpha_{12}\Delta_{x-1} + \alpha_1 \end{aligned}$$

As follows from (3.1)

$$\alpha_{11} - \alpha_{21} = \alpha_{22} - \alpha_{12}.$$

Therefore,

$$b_x - a_x = (\alpha_{11} - \alpha_{21})(b_{x-1} - a_{x-1}) + \alpha_2 - \alpha_1$$

or

$$\Delta_x = D\Delta_{x-1} + \delta.$$

Hence, we have two simultaneous recurrences:

$$\begin{cases} a_x = Sa_{x-1} + \alpha_{12}\Delta_{x-1} + \alpha_1 \\ \Delta_x = D\Delta_{x-1} + \delta. \end{cases} \quad (3.2)$$

Based on (3.2) we get:

$$\begin{aligned} a_x &= Sa_{x-1} + \alpha_{12}\Delta_{x-1} + \alpha_1 \\ &= S(Sa_{x-2} + \alpha_{12}\Delta_{x-2} + \alpha_1) + \alpha_{12}(D\Delta_{x-2} + \delta) + \alpha_1 \\ &= S^2a_{x-2} + \alpha_{12}(S+D)\Delta_{x-2} + (S+1)\alpha_1 + \alpha_{12}\delta \\ &= S^2(Sa_{x-3} + \alpha_{12}\Delta_{x-3} + \alpha_1) + \alpha_{12}(S+D)(D\Delta_{x-3} + \delta) + (S+1)\alpha_1 + \alpha_{12}\delta \\ &= S^3a_{x-3} + \alpha_{12}(S^2 + SD + D^2)\Delta_{x-3} + (S^2 + S + 1)\alpha_1 + \alpha_{12}(S + D + 1)\delta \end{aligned}$$

$$\begin{aligned}
&= S^4 a_{x-4} + \alpha_{12} (S^3 + S^2 D + S D^2 + D^3) \Delta_{x-4} + \\
&\quad (S^3 + S^2 + S + 1) \alpha_1 + \alpha_{12} (S^2 + S D + D^2 + S + D + 1) \delta \\
&= \dots = S^x a_0 + \alpha_{12} \Delta_0 \sum_{i=0}^{x-1} S^i D^{x-1-i} + \alpha_1 \sum_{i=0}^{x-1} S^i + \alpha_{12} \delta \sum_{j=0}^{x-2} \sum_{i=0}^j S^i D^{j-i} \\
&= S^x a_0 + \alpha_{12} \Delta_0 D^{x-1} \sum_{i=0}^{x-1} \left(\frac{S}{D}\right)^i + \alpha_1 \frac{S^x - 1}{S - 1} + \alpha_{12} \delta \sum_{j=0}^{x-2} D^j \sum_{i=0}^j \left(\frac{S}{D}\right)^i \\
&= S^x a_0 + \alpha_{12} \Delta_0 D^{x-1} \frac{\left(\frac{S}{D}\right)^x - 1}{\frac{S}{D} - 1} + \alpha_1 \frac{S^x - 1}{S - 1} + \alpha_{12} \delta \sum_{j=0}^{x-2} D^j \frac{\left(\frac{S}{D}\right)^{j+1} - 1}{\frac{S}{D} - 1} \\
&= S^x a_0 + \alpha_{12} \Delta_0 \frac{S^x - D^x}{S - D} + \alpha_1 \frac{S^x - 1}{S - 1} + \alpha_{12} \delta \sum_{j=0}^{x-2} \frac{S^{j+1} - D^{j+1}}{S - D} \\
&= S^x a_0 + \alpha_{12} \Delta_0 \frac{S^x - D^x}{S - D} + \alpha_1 \frac{S^x - 1}{S - 1} + \frac{\alpha_{12} \delta}{S - D} \left(\sum_{j=0}^{x-2} S^{j+1} - \sum_{j=0}^{x-2} D^{j+1} \right) \\
&= S^x a_0 + \alpha_{12} \Delta_0 \frac{S^x - D^x}{S - D} + \alpha_1 \frac{S^x - 1}{S - 1} + \frac{\alpha_{12} \delta}{S - D} \left(\frac{S(S^{x-1} - 1)}{S - 1} - \frac{D(D^{x-1} - 1)}{D - 1} \right) \\
&= S^x a_0 + \alpha_{12} \Delta_0 \frac{S^x - D^x}{S - D} + \alpha_1 \frac{S^x - 1}{S - 1} + \frac{\alpha_{12} \delta}{S - D} \left(\frac{S^x - S}{S - 1} - \frac{D^x - D}{D - 1} \right) \\
&= (\alpha_{11} + \alpha_{12})^x a_0 + \alpha_{12} (b_0 - a_0) \frac{(\alpha_{11} + \alpha_{12})^x - (\alpha_{11} - \alpha_{21})^x}{\alpha_{12} + \alpha_{21}} + \\
&\quad \alpha_1 \frac{(\alpha_{11} + \alpha_{12})^x - 1}{\alpha_{11} + \alpha_{12} - 1} + \frac{\alpha_{12} (\alpha_2 - \alpha_1)}{\alpha_{12} + \alpha_{21}} \times \\
&\quad \left(\frac{(\alpha_{11} + \alpha_{12})^x - \alpha_{11} - \alpha_{12}}{\alpha_{11} + \alpha_{12} - 1} - \frac{(\alpha_{11} - \alpha_{21})^x - \alpha_{11} + \alpha_{21}}{\alpha_{11} - \alpha_{21} - 1} \right).
\end{aligned}$$

We used that

$$S - D = \alpha_{11} + \alpha_{12} - \alpha_{11} + \alpha_{21} = \alpha_{12} + \alpha_{21}.$$

The result for b_x can be derived in the same way.

Statements 2, 3, 4, 5 are proven similarly. ■

Lemma 3.2. *If*

$$\begin{cases} a_x = \alpha_{11} a_{x-1} + \alpha_{12} b_{x-1} + \alpha_{13} c_{x-1} + \alpha_1 \\ b_x = \alpha_{21} a_{x-1} + \alpha_{22} b_{x-1} + \alpha_{23} c_{x-1} + \alpha_2 \\ c_x = \alpha_{31} a_{x-1} + \alpha_{32} b_{x-1} + \alpha_{33} c_{x-1} + \alpha_3 \end{cases} \quad (3.3)$$

and the following conditions hold:

1. $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{21} + \alpha_{22} + \alpha_{23} = \alpha_{31} + \alpha_{32} + \alpha_{33}$

2. $\exists w_1, w_2, w_1 + w_2 = 1$, that $\forall x, b_x = w_1 a_x + w_2 c_x$,

then three simultaneous recurrences (3.3) can be presented by means of representations of c through a and b , b through a and c , and a through b and c as the three pairs of the following simultaneous recurrences, respectively:

$$\begin{cases} a_x = \alpha'_{11} a_{x-1} + \alpha'_{12} b_{x-1} + \alpha_1 \\ b_x = \alpha'_{21} a_{x-1} + \alpha'_{22} b_{x-1} + \alpha_2 \end{cases} \quad (3.4)$$

where $\alpha'_{11} = \alpha_{11} - \frac{w_1}{w_2} \alpha_{13}$, $\alpha'_{12} = \alpha_{12} + \frac{1}{w_2} \alpha_{13}$, $\alpha'_{21} = \alpha_{21} - \frac{w_1}{w_2} \alpha_{23}$, $\alpha'_{22} = \alpha_{22} + \frac{1}{w_2} \alpha_{23}$;

$$\begin{cases} a_x = \alpha'_{11} a_{x-1} + \alpha'_{12} c_{x-1} + \alpha_1 \\ c_x = \alpha'_{21} a_{x-1} + \alpha'_{22} c_{x-1} + \alpha_3 \end{cases} \quad (3.5)$$

where $\alpha'_{11} = \alpha_{11} + w_1 \alpha_{12}$, $\alpha'_{12} = w_2 \alpha_{12} + \alpha_{13}$, $\alpha'_{21} = \alpha_{31} + w_1 \alpha_{32}$, $\alpha'_{22} = w_2 \alpha_{32} + \alpha_{33}$;

$$\begin{cases} b_x = \alpha'_{11} b_{x-1} + \alpha'_{12} c_{x-1} + \alpha_2 \\ c_x = \alpha'_{21} b_{x-1} + \alpha'_{22} c_{x-1} + \alpha_3 \end{cases} \quad (3.6)$$

where $\alpha'_{11} = \frac{1}{w_1} \alpha_{21} + \alpha_{22}$, $\alpha'_{12} = -\frac{w_2}{w_1} \alpha_{21} + \alpha_{23}$, $\alpha'_{21} = \frac{1}{w_1} \alpha_{31} + \alpha_{32}$, $\alpha'_{22} = -\frac{w_2}{w_1} \alpha_{31} + \alpha_{33}$ and for all these pairs of simultaneous recurrences

$$\alpha'_{11} + \alpha'_{12} = \alpha'_{21} + \alpha'_{22}.$$

Proof. Since $b_x = w_1 a_x + w_2 c_x$ then $c_x = -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x$. We substitute c in (3.3) and get:

$$\begin{aligned} a_x &= \alpha_{11} a_{x-1} + \alpha_{12} b_{x-1} + \alpha_{13} c_{x-1} + \alpha_1 \\ &= \alpha_{11} a_{x-1} + \alpha_{12} b_{x-1} + \alpha_{13} \left(\frac{1}{w_2} b_{x-1} - \frac{w_1}{w_2} a_{x-1} \right) + \alpha_1 \\ &= \left(\alpha_{11} - \frac{w_1}{w_2} \alpha_{13} \right) a_{x-1} + \left(\alpha_{12} + \frac{1}{w_2} \alpha_{13} \right) b_{x-1} + \alpha_1 \\ b_x &= \alpha_{21} a_{x-1} + \alpha_{22} b_{x-1} + \alpha_{23} c_{x-1} + \alpha_2 \\ &= \alpha_{21} a_{x-1} + \alpha_{22} b_{x-1} + \alpha_{23} \left(\frac{1}{w_2} b_{x-1} - \frac{w_1}{w_2} a_{x-1} \right) + \alpha_2 \\ &= \left(\alpha_{21} - \frac{w_1}{w_2} \alpha_{23} \right) a_{x-1} + \left(\alpha_{22} + \frac{1}{w_2} \alpha_{23} \right) b_{x-1} + \alpha_2. \end{aligned}$$

Here

$$\begin{aligned}
\alpha'_{11} + \alpha'_{12} &= \alpha_{11} - \frac{w_1}{w_2}\alpha_{13} + \alpha_{12} + \frac{1}{w_2}\alpha_{13} \\
&= \alpha_{11} + \alpha_{12} + \alpha_{13} \\
\alpha'_{21} + \alpha'_{22} &= \alpha_{21} - \frac{w_1}{w_2}\alpha_{23} + \alpha_{22} + \frac{1}{w_2}\alpha_{23} \\
&= \alpha_{21} + \alpha_{22} + \alpha_{23}.
\end{aligned}$$

Since $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{21} + \alpha_{22} + \alpha_{23}$ then $\alpha'_{11} + \alpha'_{12} = \alpha'_{21} + \alpha'_{22}$.

We substitute b in (3.3) and get:

$$\begin{aligned}
a_x &= \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_{13}c_{x-1} + \alpha_1 \\
&= \alpha_{11}a_{x-1} + \alpha_{12}(w_1a_{x-1} + w_2c_{x-1}) + \alpha_{13}c_{x-1} + \alpha_1 \\
&= (\alpha_{11} + w_1\alpha_{12})a_{x-1} + (w_2\alpha_{12} + \alpha_{13})c_{x-1} + \alpha_1 \\
c_x &= \alpha_{31}a_{x-1} + \alpha_{32}b_{x-1} + \alpha_{33}c_{x-1} + \alpha_3 \\
&= \alpha_{31}a_{x-1} + \alpha_{32}(w_1a_{x-1} + w_2c_{x-1}) + \alpha_{33}c_{x-1} + \alpha_3 \\
&= (\alpha_{31} + w_1\alpha_{32})a_{x-1} + (w_2\alpha_{32} + \alpha_{33})c_{x-1} + \alpha_3.
\end{aligned}$$

Here

$$\begin{aligned}
\alpha'_{11} + \alpha'_{12} &= \alpha_{11} + w_1\alpha_{12} + w_2\alpha_{12} + \alpha_{13} \\
&= \alpha_{11} + \alpha_{12} + \alpha_{13} \\
\alpha'_{21} + \alpha'_{22} &= \alpha_{31} + w_1\alpha_{32} + w_2\alpha_{32} + \alpha_{33} \\
&= \alpha_{31} + \alpha_{32} + \alpha_{33}.
\end{aligned}$$

Since $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{31} + \alpha_{32} + \alpha_{33}$ then $\alpha'_{11} + \alpha'_{12} = \alpha'_{21} + \alpha'_{22}$.

The pair of simultaneous recurrences for b and c and the equality $\alpha'_{11} + \alpha'_{12} = \alpha'_{21} + \alpha'_{22}$ for this case are derived in the same way. ■

Remark 3.3. Denote $w'_1 = -\frac{w_1}{w_2}$, $w'_2 = \frac{1}{w_2}$, $w''_1 = \frac{1}{w_1}$, $w''_2 = -\frac{w_2}{w_1}$. In such a case, $c = w'_1a + w'_2b$ and $a = w''_1b + w''_2c$, where $w'_1 + w'_2 = w''_1 + w''_2 = 1$. Hence, special conditions accompanying (3.3) hold for any combination of recurrences. For this reason, it would be enough to prove the equality $\alpha'_{11} + \alpha'_{12} = \alpha'_{21} + \alpha'_{22}$ for one pair of simultaneous recurrences in Lemma 3.2. The equality holds automatically for the other two pairs in this case.

Theorem 3.4. *If*

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_{13}c_{x-1} + \alpha_1 \\ b_x = \alpha_{21}a_{x-1} + \alpha_{22}b_{x-1} + \alpha_{23}c_{x-1} + \alpha_2 \\ c_x = \alpha_{31}a_{x-1} + \alpha_{32}b_{x-1} + \alpha_{33}c_{x-1} + \alpha_3 \end{cases}$$

and the following conditions hold:

1. $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{21} + \alpha_{22} + \alpha_{23} = \alpha_{31} + \alpha_{32} + \alpha_{33}$

2. $\exists w_1, w_2, w_1 + w_2 = 1$, that $\forall x, b_x = w_1a_x + w_2c_x$,

then for $C_1 = \alpha_{11} + \alpha_{12} + \alpha_{13}$, $C_2 = \alpha_{12} + \frac{1}{w_2}\alpha_{13}$, $C_3 = \alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23}$,
 $C_4 = \alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21} - \frac{w_1}{w_2}\alpha_{23}$, $C_5 = \alpha_{21} - \frac{w_1}{w_2}\alpha_{23}$:

1. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21} - \frac{w_1}{w_2}\alpha_{23} \neq 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} \neq 1$, $\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23} \neq 1$

$$\begin{aligned} a_x &= (C_1)^x a_0 + C_2 \frac{(C_1)^x - (C_3)^x}{C_4} (b_0 - a_0) + \alpha_1 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_2 - \alpha_1) C_2}{C_4} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - \frac{(C_3)^x - C_3}{C_3 - 1} \right) \\ b_x &= (C_1)^x b_0 + C_5 \frac{(C_1)^x - (C_3)^x}{C_4} (a_0 - b_0) + \alpha_2 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_1 - \alpha_2) C_5}{C_4} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - \frac{(C_3)^x - C_3}{C_3 - 1} \right) \\ c_x &= -\frac{w_1}{w_2}a_x + \frac{1}{w_2}b_x. \end{aligned}$$

2. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21} - \frac{w_1}{w_2}\alpha_{23} \neq 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} = 1$ ($\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23} \neq 1$)

$$\begin{aligned} a_x &= (C_1)^x a_0 + C_2 \frac{(C_1)^x - (C_3)^x}{C_4} (b_0 - a_0) + \alpha_1 x + \\ &\quad \frac{(\alpha_2 - \alpha_1) C_2}{C_3 - 1} \left(\frac{(C_3)^x - C_3}{C_3 - 1} - x + 1 \right) \\ b_x &= (C_1)^x b_0 + C_5 \frac{(C_1)^x - (C_3)^x}{C_4} (a_0 - b_0) + \alpha_2 x + \\ &\quad \frac{(\alpha_1 - \alpha_2) C_5}{C_3 - 1} \left(\frac{(C_3)^x - C_3}{C_3 - 1} - x + 1 \right) \\ c_x &= -\frac{w_1}{w_2}a_x + \frac{1}{w_2}b_x. \end{aligned}$$

3. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21} - \frac{w_1}{w_2}\alpha_{23} \neq 0$, $\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23} = 1$
 $(\alpha_{11} + \alpha_{12} + \alpha_{13} \neq 1)$

$$\begin{aligned} a_x &= (C_1)^x a_0 + C_2 \frac{(C_1)^x - (C_3)^x}{C_4} (b_0 - a_0) + \alpha_1 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_2 - \alpha_1) C_2}{C_1 - 1} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - x + 1 \right) \\ b_x &= (C_1)^x b_0 + C_5 \frac{(C_1)^x - (C_3)^x}{C_4} (a_0 - b_0) + \alpha_2 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_1 - \alpha_2) C_5}{C_1 - 1} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - x + 1 \right) \\ c_x &= -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x. \end{aligned}$$

4. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21} - \frac{w_1}{w_2}\alpha_{23} = 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} \neq 1$ ($\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23} \neq 1$)

$$\begin{aligned} a_x &= (C_1)^{x-1} (C_1 a_0 + C_2 (b_0 - a_0) x) + \alpha_1 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_2 - \alpha_1) C_2}{(C_1 - 1)^2} ((x - 1) (C_1)^x - x (C_1)^{x-1} + 1) \\ b_x &= (C_1)^{x-1} (C_1 b_0 + C_5 (a_0 - b_0) x) + \alpha_2 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_1 - \alpha_2) C_5}{(C_1 - 1)^2} ((x - 1) (C_1)^x - x (C_1)^{x-1} + 1) \\ c_x &= -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x. \end{aligned}$$

5. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21} - \frac{w_1}{w_2}\alpha_{23} = 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} = 1$ ($\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23} = 1$)

$$\begin{aligned} a_x &= (C_1)^{x-1} (C_1 a_0 + C_2 (b_0 - a_0) x) + \alpha_1 x + C_2 (\alpha_2 - \alpha_1) \frac{x(x-1)}{2} \\ b_x &= (C_1)^{x-1} (C_1 b_0 + C_5 (a_0 - b_0) x) + \alpha_2 x + C_5 (\alpha_1 - \alpha_2) \frac{x(x-1)}{2} \\ c_x &= -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x. \end{aligned}$$

Proof. As follows from Lemma 3.2, three simultaneous recurrences (3.3) can be reduced, specifically, to the pair of simultaneous recurrences (3.4). Based on Lemma 3.1 and after corresponding substitutions we obtain the expressions for a_x and b_x . The result for c_x is obtained automatically. ■

Remark 3.5. *It is not the only way to determine a_x , b_x , and c_x that is presented in Theorem 3.4. Each of recurrent variables a_x , b_x , and c_x can be determined through two of three pairs of simultaneous recurrences (3.4 – 3.6).*

Remark 3.6. *In principle, it could be sufficient to derive only one direct expression (for a_x , or b_x , or c_x). For example, if the formula for a_x is derived, then b_x and c_x can be expressed in a_x and in the difference between b_x and a_x . Actually, the difference $b_x - a_x$ was obtained implicitly in the course of the proof of Lemma 3.1. Based on the second recurrence in (3.2) we have:*

$$\Delta_x = D\Delta_{x-1} + \delta = D^x\Delta_0 + \delta\frac{D^{x-1} - 1}{D - 1}$$

for $D \neq 1$ and

$$\Delta_x = D\Delta_{x-1} + \delta = \Delta_0 + x\delta$$

for $D = 1$. Hence, for (3.3) and $C_3 = \alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21} + \frac{w_1}{w_2}\alpha_{23}$

$$b_x - a_x = (C_3)^x (b_0 - a_0) + (\alpha_2 - \alpha_1) \frac{(C_3)^{x-1} - 1}{C_3 - 1}$$

when $C_3 \neq 1$ and

$$b_x - a_x = b_0 - a_0 + x(\alpha_2 - \alpha_1)$$

when $C_3 = 1$. Relations between the differences are the following:

$$c_x - a_x = \frac{b_x - a_x}{w_2} = \frac{c_x - b_x}{w_1}.$$

The equality $b_x = w_1a_x + w_2c_x$ does not automatically mean the existence of the same proportion for coefficients ($\alpha_{21} = w_1\alpha_{11} + w_2\alpha_{31}$, $\alpha_{22} = w_1\alpha_{12} + w_2\alpha_{32}$, $\alpha_{23} = w_1\alpha_{13} + w_2\alpha_{33}$, $\alpha_2 = w_1\alpha_1 + w_3\alpha_3$, as in (1.1)). In the special case, the linear combination $b_x = w_1a_x + w_2c_x$ can be provided by the corresponding proportion for initial values of a , b , and c without observing the same proportion for coefficients. For example, it can be shown that in (1.2) $b_x = \frac{13}{19}a_x + \frac{6}{19}c_x$ for $a_0 = 41$, $b_0 = 47$, $c_0 = 60$ although this proportion does not take place for coefficients. Now, suppose that we have stronger conditions when also the proportion between coefficients takes place.

Lemma 3.7. *If*

1. $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{31} + \alpha_{32} + \alpha_{33}$

and

2. $\exists w_1, w_2, w_1 + w_2 = 1$, that $\alpha_{21} = w_1\alpha_{11} + w_2\alpha_{31}$, $\alpha_{22} = w_1\alpha_{12} + w_2\alpha_{32}$, $\alpha_{23} = w_1\alpha_{13} + w_2\alpha_{33}$,

then

$$\alpha_{21} + \alpha_{22} + \alpha_{23} = \alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{31} + \alpha_{32} + \alpha_{33}. \quad (3.7)$$

Proof.

$$\begin{aligned} \alpha_{21} + \alpha_{22} + \alpha_{23} &= w_1\alpha_{11} + w_2\alpha_{31} + w_1\alpha_{12} + w_2\alpha_{32} + w_1\alpha_{13} + w_2\alpha_{33} \\ &= w_1(\alpha_{11} + \alpha_{12} + \alpha_{13}) + w_2(\alpha_{31} + \alpha_{32} + \alpha_{33}) \\ &= (w_1 + w_2)(\alpha_{11} + \alpha_{12} + \alpha_{13}) \\ &= \alpha_{11} + \alpha_{12} + \alpha_{13}. \end{aligned}$$

■

Hence, the equality such as (3.7) is a redundant condition if the condition $b_x = w_1a_x + w_2c_x$ is provided by the corresponding proportion between coefficients. The equality of any two sums of coefficients from three ones in (3.7) is sufficient in this case.

Consider a situation with weaker conditions when it is only known that

$$\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{31} + \alpha_{32} + \alpha_{33}$$

and $b_x = w_1a_x + w_2c_x$, where $w_1 + w_2 = 1$. In this case, only one pair of simultaneous recurrences adduced in Lemma 3.2 can be generated. However, the solution of this pair of recurrences is sufficient for finding a_x , b_x , and c_x . On the other hand, two other pairs of simultaneous recurrences (not as in Lemma 3.2) can be generated in this case also. Indeed,

$$\begin{aligned} b_x &= w_1a_x + w_2c_x \\ &= w_1(\alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_{13}c_{x-1} + \alpha_1) + \\ &\quad w_2(\alpha_{31}a_{x-1} + \alpha_{32}b_{x-1} + \alpha_{33}c_{x-1} + \alpha_3) \\ &= (w_1\alpha_{11} + w_2\alpha_{31})a_{x-1} + (w_1\alpha_{12} + w_2\alpha_{32})b_{x-1} + (w_1\alpha_{13} + w_2\alpha_{33})c_{x-1} + \\ &\quad w_1\alpha_1 + w_2\alpha_3 \\ &= \alpha_{21}^*a_{x-1} + \alpha_{22}^*b_{x-1} + \alpha_{23}^*c_{x-1} + \alpha_2^*. \end{aligned}$$

where $\alpha_{21}^* = w_1\alpha_{11} + w_2\alpha_{31}$, $\alpha_{22}^* = w_1\alpha_{12} + w_2\alpha_{32}$, $\alpha_{23}^* = w_1\alpha_{13} + w_2\alpha_{33}$, $\alpha_2^* = w_1\alpha_1 + w_3\alpha_3$. Hence, the recurrence for b_x in (3.3) can be replaced so that we have the following three simultaneous recurrences:

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_{13}c_{x-1} + \alpha_1 \\ b_x = \alpha_{21}^*a_{x-1} + \alpha_{22}^*b_{x-1} + \alpha_{23}^*c_{x-1} + \alpha_2^* \\ c_x = \alpha_{31}a_{x-1} + \alpha_{32}b_{x-1} + \alpha_{33}c_{x-1} + \alpha_3 \end{cases} \quad (3.8)$$

By Lemma 3.7, these recurrences are of the same structure as (3.3). For this reason, by Lemma 3.2, (3.8) can be presented as three pairs of simultaneous recurrences. Thus, Theorem 3.4 can be generalized as follows.

Theorem 3.8. *If*

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_{13}c_{x-1} + \alpha_1 \\ b_x = \alpha_{21}a_{x-1} + \alpha_{22}b_{x-1} + \alpha_{23}c_{x-1} + \alpha_2 \\ c_x = \alpha_{31}a_{x-1} + \alpha_{32}b_{x-1} + \alpha_{33}c_{x-1} + \alpha_3 \end{cases}$$

and the following conditions hold:

1. $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{31} + \alpha_{32} + \alpha_{33}$

2. $\exists w_1, w_2, w_1 + w_2 = 1$, that $\forall x, b_x = w_1a_x + w_2c_x$,

then for $C_1 = \alpha_{11} + \alpha_{12} + \alpha_{13}$, $C_2 = \alpha_{12} + \frac{1}{w_2}\alpha_{13}$, $C_3 = \alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21}^* + \frac{w_1}{w_2}\alpha_{23}^*$, $C_4 = \alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^*$, $C_5 = \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^*$, $\alpha_{21}^* = w_1\alpha_{11} + w_2\alpha_{31}$, $\alpha_{23}^* = w_1\alpha_{13} + w_2\alpha_{33}$, $\alpha_2^* = w_1\alpha_1 + w_3\alpha_3$:

1. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^* \neq 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} \neq 1$, $\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21}^* + \frac{w_1}{w_2}\alpha_{23}^* \neq 1$

$$\begin{aligned} a_x &= (C_1)^x a_0 + C_2 \frac{(C_1)^x - (C_3)^x}{C_4} (b_0 - a_0) + \alpha_1 \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_2^* - \alpha_1) C_2}{C_4} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - \frac{(C_3)^x - C_3}{C_3 - 1} \right) \\ b_x &= (C_1)^x b_0 + C_5 \frac{(C_1)^x - (C_3)^x}{C_4} (a_0 - b_0) + \alpha_2^* \frac{(C_1)^x - 1}{C_1 - 1} + \\ &\quad \frac{(\alpha_1 - \alpha_2^*) C_5}{C_4} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - \frac{(C_3)^x - C_3}{C_3 - 1} \right) \\ c_x &= -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x. \end{aligned}$$

2. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^* \neq 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} = 1$ ($\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21}^* + \frac{w_1}{w_2}\alpha_{23}^* \neq 1$)

$$\begin{aligned}
a_x &= (C_1)^x a_0 + C_2 \frac{(C_1)^x - (C_3)^x}{C_4} (b_0 - a_0) + \alpha_1 x + \\
&\quad \frac{(\alpha_2^* - \alpha_1) C_2}{C_3 - 1} \left(\frac{(C_3)^x - C_3}{C_3 - 1} - x + 1 \right) \\
b_x &= (C_1)^x b_0 + C_5 \frac{(C_1)^x - (C_3)^x}{C_4} (a_0 - b_0) + \alpha_2^* x + \\
&\quad \frac{(\alpha_1 - \alpha_2^*) C_5}{C_3 - 1} \left(\frac{(C_3)^x - C_3}{C_3 - 1} - x + 1 \right) \\
c_x &= -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x.
\end{aligned}$$

3. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^* \neq 0$, $\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21}^* + \frac{w_1}{w_2}\alpha_{23}^* = 1$ ($\alpha_{11} + \alpha_{12} + \alpha_{13} \neq 1$)

$$\begin{aligned}
a_x &= (C_1)^x a_0 + C_2 \frac{(C_1)^x - (C_3)^x}{C_4} (b_0 - a_0) + \alpha_1 \frac{(C_1)^x - 1}{C_1 - 1} + \\
&\quad \frac{(\alpha_2^* - \alpha_1) C_2}{C_1 - 1} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - x + 1 \right) \\
b_x &= (C_1)^x b_0 + C_5 \frac{(C_1)^x - (C_3)^x}{C_4} (a_0 - b_0) + \alpha_2^* \frac{(C_1)^x - 1}{C_1 - 1} + \\
&\quad \frac{(\alpha_1 - \alpha_2^*) C_5}{C_1 - 1} \left(\frac{(C_1)^x - C_1}{C_1 - 1} - x + 1 \right) \\
c_x &= -\frac{w_1}{w_2} a_x + \frac{1}{w_2} b_x.
\end{aligned}$$

4. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^* = 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} \neq 1$ ($\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21}^* + \frac{w_1}{w_2}\alpha_{23}^* \neq 1$)

$$\begin{aligned}
a_x &= (C_1)^{x-1} (C_1 a_0 + C_2 (b_0 - a_0) x) + \alpha_1 \frac{(C_1)^x - 1}{C_1 - 1} + \\
&\quad \frac{(\alpha_2^* - \alpha_1) C_2}{(C_1 - 1)^2} ((x - 1) (C_1)^x - x (C_1)^{x-1} + 1) \\
b_x &= (C_1)^{x-1} (C_1 b_0 + C_5 (a_0 - b_0) x) + \alpha_2^* \frac{(C_1)^x - 1}{C_1 - 1} +
\end{aligned}$$

$$c_x = -\frac{w_1}{w_2}a_x + \frac{1}{w_2}b_x.$$

5. When $\alpha_{12} + \frac{1}{w_2}\alpha_{13} + \alpha_{21}^* - \frac{w_1}{w_2}\alpha_{23}^* = 0$, $\alpha_{11} + \alpha_{12} + \alpha_{13} = 1$ ($\alpha_{11} - \frac{w_1}{w_2}\alpha_{13} - \alpha_{21}^* + \frac{w_1}{w_2}\alpha_{23}^* = 1$)

$$\begin{aligned} a_x &= (C_1)^{x-1} (C_1 a_0 + C_2 (b_0 - a_0) x) + \alpha_1 x + C_2 (\alpha_2^* - \alpha_1) \frac{x(x-1)}{2} \\ b_x &= (C_1)^{x-1} (C_1 b_0 + C_5 (a_0 - b_0) x) + \alpha_2^* x + C_5 (\alpha_1 - \alpha_2^*) \frac{x(x-1)}{2} \\ c_x &= -\frac{w_1}{w_2}a_x + \frac{1}{w_2}b_x. \end{aligned}$$

In the special case, when (3.7) is true, Theorem 3.8 is reduced to Theorem 3.4.

4. Conclusions and Open Problems

We have considered the problem of finding a solution of simultaneous recurrences. Section 2 presents the general method of arriving at a solution. Section 3 describes a special method for solving matrix recurrences of order two and order three. This method works only under the special conditions (specifically, when the sums of coefficients of items in lines are equal) but seems to be more efficient. It might be noted that perhaps, this method can be generalized to matrix recurrences of order n .

It might also be of interest to investigate simultaneous recurrences with equal sums of coefficients of items in columns.

Lemma 4.1. *If*

$$\begin{cases} a_x = \alpha_{11}a_{x-1} + \alpha_{12}b_{x-1} + \alpha_1 \\ b_x = \alpha_{21}a_{x-1} + \alpha_{22}b_{x-1} + \alpha_2 \end{cases}$$

and

$$\alpha_{11} + \alpha_{21} = \alpha_{12} + \alpha_{22} ,$$

then

when $\alpha_{12} \neq -\alpha_{21}$, $\alpha_{11} - \alpha_{12} \neq 1$, $\alpha_{11} + \alpha_{21} \neq 1$

$$a_x = (\alpha_{11} - \alpha_{12})^x a_0 + \alpha_{12} (a_0 + b_0) \frac{(\alpha_{11} + \alpha_{21})^x - (\alpha_{11} - \alpha_{12})^x}{\alpha_{12} + \alpha_{21}} +$$

$$\begin{aligned}
& \alpha_1 \frac{(\alpha_{11} - \alpha_{12})^x - 1}{\alpha_{11} - \alpha_{12} - 1} + \frac{\alpha_{12}(\alpha_1 + \alpha_2)}{\alpha_{12} + \alpha_{21}} \times \\
& \left(\frac{(\alpha_{11} + \alpha_{21})^x - \alpha_{11} - \alpha_{21}}{\alpha_{11} + \alpha_{21} - 1} - \frac{(\alpha_{11} - \alpha_{12})^x - \alpha_{11} + \alpha_{12}}{\alpha_{11} - \alpha_{12} - 1} \right) \\
b_x = & (\alpha_{11} - \alpha_{12})^x b_0 + \alpha_{21} (a_0 + b_0) \frac{(\alpha_{11} + \alpha_{21})^x - (\alpha_{11} - \alpha_{12})^x}{\alpha_{12} + \alpha_{21}} + \\
& \alpha_2 \frac{(\alpha_{11} - \alpha_{12})^x - 1}{\alpha_{11} - \alpha_{12} - 1} + \frac{\alpha_{21}(\alpha_1 + \alpha_2)}{\alpha_{12} + \alpha_{21}} \times \\
& \left(\frac{(\alpha_{11} + \alpha_{21})^x - \alpha_{11} - \alpha_{21}}{\alpha_{11} + \alpha_{21} - 1} - \frac{(\alpha_{11} - \alpha_{12})^x - \alpha_{11} + \alpha_{12}}{\alpha_{11} - \alpha_{12} - 1} \right).
\end{aligned}$$

Proof. The proof is like that of Lemma 3.1. ■

We intend to generalize Lemma 4.1 to matrix recurrences of order n .

References

- [1] D. M. Bloom, *Linear Algebra and Geometry*, Cambridge University Press, Cambridge, London, 1979.
- [2] M. Bocher and E. P. R. Duval, *Introduction to Higher Algebra*, The Macmillan Company, New York, 1907.
- [3] K. H. Rosen (ed.), *Handbook of Discrete and Combinatorial Mathematics*, CRC Press, Boca Raton, 2000.