On free-group algorithms that sandwich a subgroup between free-product factors

Warren Dicks*

November 27, 2024

Abstract. Let F be a finite-rank free group and H be a finite-rank subgroup of F. We discuss proofs of two algorithms that sandwich H between an upper-layer free-product factor of F that contains H and a lower-layer free-product factor of F that is contained in H.

Richard Stong showed that the unique smallest-possible upper layer, denoted CL(H), is visible in the output of the polynomial-time cut-vertex algorithm of J. H. C. Whitehead. Stong's proof used bi-infinite paths in a Cayley tree and sub-surfaces of a three-manifold. We give a variant of his proof that uses edge-cuts of the Cayley tree induced by edge-cuts of a Bass-Serre tree.

A. Clifford and R. Z. Goldstein gave an exponential-time algorithm that determines whether or not the trivial subgroup is the only possible lower layer. Their proof used Whitehead's three-manifold techniques. We give a variant of their proof that uses Whitehead's cut-vertex results, and thereby obtain a somewhat simpler algorithm that yields a lower layer of maximum-possible rank.

2010 Mathematics Subject Classification. Primary: 20E05; Secondary: 20E36, 20E08. Key words. Sub-bases of free groups. Free-product factors. Cut-vertex algorithm. Cut-vertex lemma. Clifford-Goldstein algorithm.

1 Introduction

1.1 Definitions. For any set E, we let $\langle E | \rangle$ denote the free group on E. By a *basis* of $\langle E | \rangle$, we mean a free-generating set of $\langle E | \rangle$. By a *sub-basis* of $\langle E | \rangle$ we mean a subset of a basis of $\langle E | \rangle$. We let Aut $\langle E | \rangle$ denote the group of automorphisms of $\langle E | \rangle$ acting on the right as exponents.

For any subset Z of $\langle E | \rangle$, we let $\langle Z \rangle$ denote the subgroup of $\langle E | \rangle$ generated by Z. We let $\operatorname{supp}(Z \operatorname{rel} E)$ denote the \subseteq -smallest subset of E such that $Z \subseteq \langle \operatorname{supp}(Z \operatorname{rel} E) \rangle$. We let $\operatorname{CL}(Z)$ denote the intersection of all the free-product factors (generated by sub-bases) of $\langle E | \rangle$ that contain Z.

^{*}Partially supported by Spain's Ministerio de Ciencia e Innovación through Project MTM2011-25955.

1.2 Hypotheses. Throughout, let E be a finite set, let Z be a finite subset of $\langle E | \rangle$, and let E_Z denote supp(Z rel E).

1.3 History. Recall Hypotheses 1.2.

• In [8, publ. 1936], J. H. C. Whitehead gave his true-word and cyclic-word cut-vertex algorithms, and the former determines whether or not Z is a sub-basis of $\langle E | \rangle$. A little later, in [9, publ. 1936], he gave an exponential-time, general-purpose algorithm which has largely overshadowed the easier-to-prove, polynomial-time, limited-use algorithm. We wish to emphasize that the cut-vertex algorithm suffices to efficiently sandwich a subgroup between two free-product factors.

Whitehead defined a certain finite graph which we denote $W_{H_*}(Z \operatorname{rel} E_Z)$. He observed that if some vertex of $W_{H_*}(Z \operatorname{rel} E_Z)$ is what we call a Whitehead cut-vertex, then it is straightforward to construct an automorphism of $\langle E | \rangle$ that strictly reduces the total *E*-length of *Z*. Clearly, one then has an algorithm (with choices) which constructs some $\Psi \in \operatorname{Aut}\langle E | \rangle$ such that $W_{H_*}(Z^{\Psi} \operatorname{rel} E_{Z^{\Psi}})$ has no Whitehead cut-vertices. It then remains to extract information from Ψ and Z^{Ψ} . For example, it will transpire that the rank of $\operatorname{CL}(Z)$ is $|E_{Z^{\Psi}}|$. One reason this is interesting is that $\operatorname{Edward} C$. Turner [7, Theorem 1] showed that the rank of $\operatorname{CL}(Z)$ is |E| if and only if *Z* is a *test set* for injective endomorphisms of $\langle E | \rangle$ to be automorphisms, that is, each injective endomorphism of $\langle E | \rangle$ that maps $\langle Z \rangle$ onto itself is an automorphism.

Using a three-manifold model of $W_{H_*}(Z \text{ rel } E_Z)$, Whitehead proved a cutvertex lemma: If Z is a sub-basis of $\langle E | \rangle$, then $Z^{\Psi} \subseteq E^{\pm 1}$.

Hence, Z is a sub-basis of $\langle E | \rangle$ if and only if $Z \cap Z^{-1} = \emptyset$ and $Z^{\Psi} \subseteq E^{\pm 1}$; in this event, $Z \cup (E^{\Psi^{-1}} - Z^{\pm 1})$ is a basis of $\langle E | \rangle$.

Set $E' := E^{\Psi^{-1}}$ and $E'_Z := \operatorname{supp}(Z \operatorname{rel} E')$. Expressing the elements of Z^{Ψ} in terms of E is equivalent to expressing the elements of Z in terms of E'. The important point is that $\operatorname{WH}_*(Z \operatorname{rel} E'_Z)$ is isomorphic to $\operatorname{WH}_*(Z^{\Psi} \operatorname{rel} E_{Z^{\Psi}})$ and, hence, has no Whitehead cut-vertices.

In [6, publ. 1997], Richard Stong used bi-infinite paths in a Cayley tree and sub-surfaces homologous to an essential disk in a three-manifold to prove a more general cut-vertex lemma: The set E'_Z is a basis of $\operatorname{CL}(Z)$, and, for each free-product factorization $\operatorname{CL}(Z) = \underset{i \in I}{*} H_i$ such that $Z \subseteq \bigcup_{i \in I} H_i$, the set E'_Z contains a basis of each H_i .

Not only can a basis of $\operatorname{CL}(Z)$ be computed efficiently, but also there are only finitely many possibilities for the sets $\{H_i\}_{i \in I}$, and they can all be computed efficiently. To see how Stong's cut-vertex lemma generalizes Whitehead's, notice that if Z is a sub-basis of $\langle E | \rangle$, then $\operatorname{CL}(Z) = \langle Z \rangle = \underset{z \in Z}{*} \langle z \rangle$ and $Z \subseteq \bigcup_{z \in Z} \langle z \rangle$, and, here, for E'_Z to contain a basis of each $\langle z \rangle$, which is necessarily $\{z\}$ or $\{z^{-1}\}$, one must have $(E'_Z)^{\pm 1} \supseteq Z$, and, hence, $E^{\pm 1} \supseteq Z^{\Psi}$.

• In [1, publ. 2010], A. Clifford and R. Z. Goldstein revisited Whitehead's three-manifold techniques and constructed an ingenious exponential-time algo-

rithm which determines whether or not some element of $\langle Z \rangle$ lies in a basis of $\langle E \mid \rangle$, and, in the affirmative case, finds such an element.

1.4 Content. What we do in this article is formalize Whitehead's cut-vertex algorithm, give a Bass-Serre-theoretic proof of Stong's cut-vertex lemma, and give an algorithm that yields a basis E'' of $\langle E | \rangle$ that maximizes $|E'' \cap \langle Z \rangle|$.

In Section 2, for completeness and to develop the notation and basic results that will be used, we formalize part of Whitehead's discussion of cut-vertices and free-group automorphisms, including his true-word cut-vertex algorithm.

In Section 3, Stong's beautiful true-word cut-vertex lemma is proved using edge-cuts of a Cayley tree induced by edge-cuts of a Bass-Serre tree. At this stage, we will have given a detailed proof for the polynomial-time algorithm for computing a basis of CL(Z) that is more algebraic than Stong's proof.

In Section 4, we restructure the Clifford-Goldstein argument using Whitehead's cut-vertex results in place of the topology, and obtain a slightly faster, more powerful algorithm that yields a basis E'' of $\langle E | \rangle$ which maximizes $|E'' \cap \langle Z \rangle|$. In particular, $E'' \cap \langle Z \rangle \neq \emptyset$ if and only if some element of $\langle Z \rangle$ lies in a basis of $\langle E | \rangle$.

A formalized cut-vertex algorithm 2

This technical section gives elementary definitions and arguments that formalize part of Whitehead's discussion [8, pp.50–52] of cut-vertices and free-group automorphisms.

By a *graph*, we mean a set given as the disjoint union of two sets, called the vertex-set and the edge-set, together with an initial-vertex map and a terminal-vertex map, each of which maps the edge-set to the vertex-set. For any set S, we write $\mathbb{K}(S)$ to denote the graph which has vertex-set S and edge-set $S^{\times 2} := S \times S$, where an edge (x, y) has initial vertex x and terminal vertex y.

2.1 Notation. Recall Hypotheses 1.2.

• For $e \in E$, we write $\overline{e} \coloneqq e^{-1}$ and $e^{\pm 1} \coloneqq \{e, \overline{e}\}$. We write $E^{-1} \coloneqq \{\overline{e} \mid e \in E\}$ and $E^{\pm 1} := E \cup E^{-1}$. We shall be interested in the graph $\mathbb{K}(E^{\pm 1} \cup \{1\})$, which has basepoint 1 and an inversion map on the vertices.

• Consider any $z \in \langle E | \rangle$, and let $e_1 e_2 \cdots e_n$ represent the reduced $E^{\pm 1}$ -expression for z.

ession for z. • supp({z} rel E) = $\bigcup_{i=1}^{n} (E \cap e_i^{\pm 1})$ and supp(Z rel E) = $\bigcup_{z \in Z} \text{supp}(\{z\} rel E)$. • We set $||z||_E \coloneqq n$ and $||Z||_E \coloneqq \sum_{z \in Z} ||z||_E$.

• We say that a product xy has no $E^{\pm 1}$ -cancellation if $||xy||_E = ||x||_E + ||y||_E$, and then sometimes write xy as $x \cdot y$ for emphasis.

• Suppose that $z \neq 1$. We set $e_0 \coloneqq e_{n+1} \coloneqq 1$. For $i \in \{0, 1, \ldots, n\}$, we say that (e_i, e_{i+1}) occurs in the reduced $(E^{\pm 1} \cup \{1\})$ -expression for z, and note that there exist $g', g'' \in \langle E | \rangle$ such that $z = g' \cdot e_i \cdot e_{i+1} \cdot g''$ with no $E^{\pm 1}$ -cancellation, g' = 1 if $e_i = 1$, and g'' = 1 if $e_{i+1} = 1$. We set

 $\begin{aligned} & \operatorname{WH}_*(\{z\} \text{ rel } E) \coloneqq E^{\pm 1} \cup \{1\} \cup \{(\overline{e}_i, e_{i+1})\}_{i=0}^n \subseteq \mathbb{K}(E^{\pm 1} \cup \{1\}) \\ & \text{and } \operatorname{WH}_*(\{1\} \text{ rel } E) \coloneqq E^{\pm 1} \cup \{1\}. \quad \text{For example, for each } e \in E^{\pm 1}, \text{ we have } \\ & \operatorname{WH}_*(\{e\} \text{ rel } E) = E^{\pm 1} \cup \{1\} \cup \{(1, e), (\overline{e}, 1)\}. \text{ We also have the pentagonal example } \\ & \operatorname{WH}_*(\{x^2y^2\} \text{ rel } \{x, y\}) = \{x, y, \overline{x}, \overline{y}, 1, (1, x), (\overline{x}, x), (\overline{x}, y), (\overline{y}, y), (\overline{y}, 1)\}. \end{aligned}$

• Let S be a subset of $\langle E | \rangle$. If $S \neq \emptyset$, we set

$$WH_*(S \text{ rel } E) := \bigcup_{z \in S} WH_*(\{z\} \text{ rel } E) \subseteq \mathbb{K}(E^{\pm 1} \cup \{1\}),$$

and we set $W_{H_*}(\emptyset \text{ rel } E) := E^{\pm 1} \cup \{1\}$. In $W_{H_*}(S \text{ rel } E)$, a vertex e_* is said to be a *Whitehead cut-vertex* if removing e_* and all the edges incident to e_* leaves a basepointed graph that is not connected; this entails $e_* \neq 1$. If $W_{H_*}(S \text{ rel } E)$ is not connected, then each element of $E^{\pm 1}$ is a Whitehead cut-vertex, since the set of valence-zero vertices is closed under inversion.

• We let CUTS(E) denote the set of those ordered triples $({}_{0}D, {}_{1}D, e_{\star})$ such that ${}_{0}D \cup {}_{1}D = E^{\pm 1}, {}_{0}D \cap {}_{1}D = \{e_{\star}\}, \text{ and } {}_{1}D \neq \{e_{\star}\}.$ Clearly, ${}_{0}D \subseteq E^{\pm 1}, {}_{1}D \subseteq E^{\pm 1}, \text{ and } e_{\star} \in E^{\pm 1}.$ Suppose that $\mathbf{C} = ({}_{0}D, {}_{1}D, e_{\star}) \in \text{CUTS}(E).$

• For each $(\alpha, \beta) \in \{0, 1\}^{\times 2}$, we set $_{\alpha}D_{\beta} := _{\alpha}D \cap _{\beta}D^{-1}$ and $_{\alpha}E_{\beta} := E \cap _{\alpha}D_{\beta}$.

• Let $\chi: E^{\pm 1} \to \{0, 1\}, e \mapsto \chi(e) \coloneqq |\{e\} \cap {}_1D|$, be the characteristic map of ${}_1D$. We set $\eta_{\mathbf{C}} \coloneqq \chi(\overline{e}_{\star}) \in \{0, 1\}$ and $d_{\star} \coloneqq e_{\star}^{2\eta_{\mathbf{C}}-1} \in e_{\star}^{\pm 1}$, that is, $d_{\star} = \overline{e}_{\star}$ if $\overline{e}_{\star} \in {}_0D$, while $d_{\star} = e_{\star}$ if $\overline{e}_{\star} \in {}_1D$. We define $\varphi_{\mathbf{C}}$ to be the automorphism of $\langle E | \rangle$ that fixes d_{\star} and maps e to $d_{\star}^{\chi(e)} e \overline{d}_{\star} \chi(\overline{e})$ for each $e \in E - d_{\star}^{\pm 1}$.

• We define three subgraphs of $\mathbb{K}(E^{\pm 1} \cup \{1\})$: $_{0}WH(\mathbf{C}) \coloneqq \mathbb{K}(_{0}D \cup \{1\})$; $_{1}WH(\mathbf{C}) \coloneqq \mathbb{K}(_{1}D)$; and, $WH_{*}(\mathbf{C}) \coloneqq \mathbb{K}(_{0}D \cup \{1\}) \cup \mathbb{K}(_{1}D)$. We say that \mathbf{C} cuts each subgraph of $WH_{*}(\mathbf{C})$ with the full vertex-set, $E^{\pm 1} \cup \{1\}$.

If **C** cuts WH_{*}(Z rel E), then e_{\star} is a Whitehead cut-vertex of WH_{*}(Z rel E), since $_{0}D \cup \{1\}$ and $_{1}D$ have union $E^{\pm 1} \cup \{1\}$ and intersection $\{e_{\star}\}$, while $_{0}D \cup \{1\} \neq \{e_{\star}\} \neq _{1}D$.

2.2 Lemma. With Hypotheses 1.2, fix $\mathbf{C} = ({}_0D, {}_1D, e_{\star}) \in \text{CUTS}(E)$, and let $z \in \langle E | \rangle$. Then the following hold.

- (i) $\tilde{E} := \bigcup_{(\alpha,\beta)\in\{0,1\}^{\times 2}} (a^{\alpha}({}_{\alpha}E_{\beta})\overline{a}^{\beta})$ is a basis of $\langle E \cup \{a\} \mid \rangle$.
- (ii) $||z||_{\widetilde{E}} = ||z||_E$ if and only if **C** cuts WH_{*}({z} rel E).
- (iii) If **C** cuts WH_{*}({z} rel E), then $||z^{\overline{\varphi}_{\mathbf{C}}}||_E \leq ||z||_E$,
- (iv) If **C** cuts WH_{*}({z} rel E) and e_* has positive valence in the subgraph WH_{*}({z} rel E) $\cap_{1-\eta_{\mathbf{C}}}$ WH(**C**), then $||z^{\overline{\varphi}_{\mathbf{C}}}||_E < ||z||_E$.

Proof. Set $F \coloneqq \langle E | \rangle$, $\tilde{F} \coloneqq \langle E \cup \{a\} | \rangle$, $\eta \coloneqq \eta_{\mathbf{C}}$, and $\varphi \coloneqq \varphi_{\mathbf{C}}$. (i). Recall that $e_{\star} \in E^{\pm 1}$.

If $e_{\star} \in E$, then ${}_{0}E_{\eta} \cap {}_{1}E_{\eta} = \{e_{\star}\}$ and there are no other overlaps among the ${}_{\alpha}E_{\beta}$. Since $\{a^{0}e_{\star}\overline{a}^{\eta}, a^{1}e_{\star}\overline{a}^{\eta}\} \subseteq \tilde{E}$ and $(a^{1}e_{\star}\overline{a}^{\eta})(a^{0}e_{\star}\overline{a}^{\eta})^{-1} = a$, we see easily that \tilde{E} is a basis of \tilde{F} .

Warren Dicks

Similarly, if $e_{\star} \in E^{-1}$, then ${}_{\eta}E_0 \cap {}_{\eta}E_1 = \{\overline{e}_{\star}\}$ and there are no other overlaps among the ${}_{\alpha}E_{\beta}$. Again, \tilde{E} is a basis of \tilde{F} .

(ii). Let $e_1e_2\cdots e_n$ represent the reduced $E^{\pm 1}$ -expression for z. For any map $\{0,\ldots,n\} \to \{0,1\}, i \mapsto \chi_i$, the following three conditions are easily seen to be equivalent.

• the reduced $\tilde{E}^{\pm 1}$ -expression for z is $(a^{\chi_0}e_1\overline{a}^{\chi_1})(a^{\chi_1}e_2\overline{a}^{\chi_2})\cdots(a^{\chi_{n-1}}e_n\overline{a}^{\chi_n}).$

• $e_i \in \chi_{i-1} D_{\chi_i}$, i = 1, 2, ..., n, and $\chi_0 = \chi_n = 0$.

• $(\overline{e}_i, e_{i+1}) \in {}_{\chi_i} D^{\times 2}, i = 1, 2, ..., n-1, (\overline{e}_n, e_1) \in {}_0 D^{\times 2}, \text{ and } \chi_0 = \chi_n = 0.$ Now (ii) follows.

(iii). Let $\tilde{\varphi} \colon \tilde{F} \to F$ denote the retraction that carries a to $d_{\star} \coloneqq e_{\star}^{2\eta-1}$. We apply $\tilde{\varphi}$ to $\{a^{\eta}e_{\star} \overline{a}^{\eta}, a^{1-\eta}e_{\star} \overline{a}^{\eta}\} = \{a^{0}e_{\star} \overline{a}^{\eta}, a^{1}e_{\star} \overline{a}^{\eta}\} \subseteq \tilde{E}^{\pm 1}$. Here, we have $(a^{\eta}e_{\star} \overline{a}^{\eta})^{\tilde{\varphi}} = e_{\star} d_{\star}^{\eta-\eta} = e_{\star}$ and $(a^{1-\eta}e_{\star} \overline{a}^{\eta})^{\tilde{\varphi}} = e_{\star} d_{\star}^{1-\eta-\eta} = 1$. It follows that $\tilde{\varphi}$ carries \tilde{E} to $E^{\varphi} \cup \{1\}$. Since $z^{\tilde{\varphi}} = z$, we see that $||z||_{E^{\varphi}} \leq ||z||_{\tilde{E}}$. Now $||z^{\overline{\varphi}}||_{E} = ||z||_{E^{\varphi}} \leq ||z||_{\tilde{E}} = ||z||_{E}$, by (ii).

(iv). There exists some vertex e of $_{1-\eta}WH(\mathbf{C})$ such that $(\overline{e}, e_{\star})$ occurs in the reduced $(E^{\pm 1} \cup \{1\})$ -expression for z or \overline{z} . Necessarily, $e \neq e_{\star}$. Hence, $e \notin_{\eta}WH(\mathbf{C})$. As in (ii), the element $(a^{1-\eta}e_{\star}\overline{a}^{\eta}) \in \tilde{E}^{\pm 1}$ occurs in the reduced $\tilde{E}^{\pm 1}$ -expression for z or \overline{z} . Hence, $(a^{1-\eta}e_{\star}\overline{a}^{\eta})$ or $(a^{\eta}\overline{e_{\star}}\overline{a}^{1-\eta})$ occurs in the reduced $\tilde{E}^{\pm 1}$ -expression for z. As in (iii), each such term is mapped to 1 by $\tilde{\varphi}$. Thus, $||z^{\overline{\varphi}}||_{E} = ||z||_{E^{\varphi}} < ||z||_{\widetilde{E}} = ||z||_{E}$.

2.3 Algorithm. Recall Hypotheses 1.2. Whitehead's *cut-vertex subroutine* [8, p. 51] has the following structure.

INPUT: A Whitehead cut-vertex e_* of WH_{*}(Z rel E_Z).

OUTPUT: A $\mathbf{C} \in \text{CUTS}(E)$ with WH_{*}(Z rel E) \subseteq WH_{*}(\mathbf{C}) and $||Z^{\overline{\varphi}_{\mathbf{C}}}||_{E} < ||Z||_{E}$. PROCEDURE. We consider two cases.

CASE 1: $WH_*(Z \text{ rel } E_Z)$ is connected.

Deleting e_{\star} and its incident edges from WH_{*}(Z rel E_Z) leaves a subgraph that has a unique expression as the disjoint union of two nonempty subgraphs X_0 and X_1 such that X_0 is connected and contains {1}.

Set $_{0}D \coloneqq (X_{0} \cap E_{Z}^{\pm 1}) \cup \{e_{\star}\} \cup (E^{\pm 1} - E_{Z}^{\pm 1}), \ _{1}D \coloneqq (X_{1} \cap E_{Z}^{\pm 1}) \cup \{e_{\star}\},$ and $\mathbf{C} \coloneqq (_{0}D, _{1}D, e_{\star}) \in \text{CUTS}(E).$ Then $\text{WH}_{*}(Z \text{ rel } E) \subseteq \text{WH}_{*}(\mathbf{C}),$ and e_{\star} has positive valence in both $\text{WH}_{*}(Z \text{ rel } E) \cap _{0}\text{WH}(\mathbf{C})$ and $\text{WH}_{*}(Z \text{ rel } E) \cap _{1}\text{WH}(\mathbf{C}).$ Thus, e_{\star} has positive valence in $\text{WH}_{*}(Z \text{ rel } E) \cap _{1 - \eta_{\mathbf{C}}}\text{WH}(\mathbf{C}).$ It follows from Lemma 2.2(iii),(iv) that $||Z^{\overline{\varphi}_{\mathbf{C}}}||_{E} < ||Z||_{E}.$ We return \mathbf{C} and terminate the procedure.

CASE 2: $WH_*(Z \text{ rel } E_Z)$ is not connected.

Let X denote the component of $W_{H_*}(Z \operatorname{rel} E_Z)$ containing {1}, and let $D := X \cap E_Z^{\pm 1}$. If it were the case that $D^{-1} = D$, then it is not difficult to see that we would have $Z \subseteq \langle D \rangle$, $E_Z^{\pm 1} = D$, and $X = W_{H_*}(Z \operatorname{rel} E_Z)$, which would contradict the assumption that $W_{H_*}(Z \operatorname{rel} E_Z)$ is not connected. Thus, $D^{-1} \neq D$, $D \not\subseteq D^{-1}$, and $D - D^{-1} \neq \emptyset$.

Choose $e'_{\star} \in D - D^{-1}$, and set $_{0}D \coloneqq D \cup (E^{\pm 1} - E^{\pm 1}_{Z}), {}_{1}D \coloneqq (E^{\pm 1}_{Z} - D) \cup \{e'_{\star}\}$, and $\mathbf{C} \coloneqq ({}_{0}D, {}_{1}D, e'_{\star}) \in \text{CUTS}(E)$. It is clear that $\text{WH}_{*}(Z \text{ rel } E) \subseteq \text{WH}_{*}(\mathbf{C})$. Here, $\eta_{\mathbf{C}} = |\{(e'_{\star})^{-1}\} \cap {}_{1}D| = 1$. Also, $\text{WH}_{*}(Z \text{ rel } E) \cap {}_{0}\text{WH}(\mathbf{C}) \supseteq X$, the component of $\text{WH}_{*}(Z \text{ rel } E_{Z})$ that contains $\{e'_{\star}, 1\}$. Since e'_{\star} has positive valence in X, it follows from Lemma 2.2(iii),(iv) that $||Z^{\overline{\varphi}_{\mathbf{C}}}||_{E} < ||Z||_{E}$. We return \mathbf{C} and terminate the procedure.

2.4 Algorithm. Recall Hypotheses 1.2. Via the mock flow chart



Whitehead's cut-vertex algorithm [8, p. 51] returns a pair (Φ, Z') such that $\Phi \in \operatorname{Aut}\langle E | \rangle, Z' = Z^{\overline{\Phi}}$, and the isomorphic graphs $\operatorname{WH}_*(Z' \operatorname{rel supp}(Z' \operatorname{rel} E))$ and $\operatorname{WH}_*(Z \operatorname{rel supp}(Z \operatorname{rel} E^{\Phi}))$ have no Whitehead cut-vertices. It is then not difficult to find $\operatorname{supp}(Z' \operatorname{rel} E), E^{\Phi}$, and, hence, $\operatorname{supp}(Z \operatorname{rel} E^{\Phi})$.

Information about these will be given in Lemmas 3.4 and 3.7. For example, $|\operatorname{supp}(Z \operatorname{rel} E^{\Phi})|$ is smallest-possible over all bases of $\langle E| \rangle$, that is, $\operatorname{supp}(Z \operatorname{rel} E^{\Phi})$ is a basis of $\operatorname{CL}(Z)$. Also, Z is a sub-basis of $\langle E| \rangle$ if and only if $Z \cap Z^{-1} = \emptyset$ and $Z' \subseteq E^{\pm 1}$; in this event, $Z \cup (E^{\Phi} - Z^{\pm 1})$ is a basis of $\langle E| \rangle$. \Box

2.5 Notes. Although Whitehead did not mention it, it is possible to implement Algorithm 2.4 in such a way that it terminates in time that is polynomial (linear?) in $|E| + ||Z||_E$. Depth-first searches may be used to find the component X of WH_{*}(Z' rel $E_{Z'}$) that contains {1}, and to search for an element $e_* \in X \cap E_{Z'}^{\pm 1}$ such that either $\overline{e}_* \notin X$ or removing e_* and its incident edges from X leaves a graph that is not connected. If no such e_* exists then WH_{*}(Z' rel $E_{Z'}$) has no Whitehead cut-vertices, as was seen in Algorithm 2.3. If such an e_* exists, then it may be used to construct a φ such that $||Z'\overline{\varphi}||_E < ||Z'||_E$, as was also seen in Algorithm 2.3.

3 Bass-Serre proofs of cut-vertex lemmas

3.1 Review. Let F be a group.

• Let S be a subset of F. We let $\operatorname{Cayley}(F, S)$ denote the graph with vertex-set F and edge-set $F \times S$, where each edge $(g, s) \in F \times S$ has initial vertex g and terminal vertex gs; we shall sometimes write $\operatorname{edge}(g \xrightarrow{\bullet(s)} gs)$ to denote the pair (g, s) viewed as an edge. Then $\operatorname{Cayley}(F, S)$ is an F-graph. It is a tree when S is a basis of F. See, for example, [2, Theorem I.7.6].

• Let I be a set and $(H_i)_{i \in I}$ be a family of subgroups of F. We let BassSerre $(F, (H_i)_{i \in I})$ denote the graph whose vertex-set is the disjoint union of the set F together with the sets F/H_i , $i \in I$, and whose edge-set is $F \times I$, where each edge $(g, i) \in F \times I$ has initial vertex g and terminal vertex gH_i ; we shall sometimes write $edge(g \xrightarrow{\bullet(H_i)} gH_i)$ to denote the pair (g, i) viewed as an edge. Then BassSerre $(F, (H_i)_{i \in I})$ is an F-graph. It is a tree when $F = \underset{i \in I}{*} H_i$, by a result of H. Bass and J.-P. Serre. See, for example, [2, Theorem I.7.6].

Notice that if a subset S of $\langle E | \rangle$ contains E, then WH_{*}(S rel E) contains the basepointed star WH_{*}(E rel E), and therefore has no Whitehead cut-vertices. The following amazing partial converse can be extracted from the (1) \Rightarrow (3) part of [6, Theorem 10]. The case where each free-product factor is cyclic is essentially Whitehead's cut-vertex lemma [8, Lemma].

3.2 The Stong-Whitehead theorem. For each finite set E and free-product factorization $\langle E | \rangle = \underset{i \in I}{*} H_i$ such that $\bigcup_{i \in I} H_i \not\supseteq E$, the graph $\operatorname{WH}_*((\bigcup_{i \in I} H_i) \operatorname{rel} E)$ has a Whitehead cut-vertex.

Proof. Set $F := \langle E | \rangle = \underset{i \in I}{*} H_i$. Recall Review 3.1, and set T := Cayley(F, E)and $T^* := \text{BassSerre}(F, (H_i)_{i \in I})$. Thus, T and T^* are F-trees whose vertex-sets contain F.

We work first with T^* . We let $\lim_{T^*}(1)$ denote the set of T^* -edges incident to the T^* -vertex 1, and $\operatorname{star}_{T^*}(1)$ denote the set of components of the forest $T^* - \lim_{T^*}(1)$. For each T^* -vertex v, there exists a unique component $\chi(v) \in \operatorname{star}_{T^*}(1)$ such that $v \in \chi(v)$. For any T^* -vertices v and w, we let $T^*[v, w]$ denote the \subseteq -smallest subtree of T^* that contains $\{v, w\}$, and then $\chi(v) \neq \chi(w)$ if and only if $1 \in T^*[v, w]$ and $v \neq w$. Also, χ restricts to a map $F \to \operatorname{star}_{T^*}(1)$.

In T now, set $\delta := \{ \text{edge}(g \xrightarrow{\bullet(e)} ge) = (g, e) \in F \times E \subseteq T \mid \chi(g) \neq \chi(ge) \}$. Clearly, χ is constant on the vertex-set of each component of $T - \delta$. An element $(g, e) \in F \times E$ lies in δ if and only if $1 \in T^*[g, ge]$, or, equivalently, $\overline{g} \in T^*[1, e]$. Since E is nonempty and finite, it is clear that δ is nonempty and finite. Hence, there exists $(g_{\delta}, e_{\delta}) \in F \times E^{\pm 1}$ satisfying $||g_{\delta}e_{\delta}||_{E} = ||g_{\delta}||_{E} + 1$ and $\chi(g_{\delta}) \neq \chi(g_{\delta}e_{\delta})$ such that $||g_{\delta}||_{E}$ has the maximum possible value.

We shall now show that $g_{\delta} \neq 1$. By hypothesis, there exists $e_0 \in E - \bigcup_I H_i$. Since $e_0 \neq 1$, there exists some $T^*[1, e_0]$ -neighbour of 1, necessarily $1H_{i_0}$ for some $i_0 \in I$. Clearly $e_0 \neq 1H_{i_0}$; thus, there exists some $T^*[1, e_0]$ -neighbour of $1H_{i_0}$ other than 1, necessarily some $h_0 \in H_{i_0} - \{1\}$. Now $h_0 \in T^*[1, e_0]$, $1 \in T^*[\overline{h}_0, \overline{h}_0 e_0], \chi(\overline{h}_0) \neq \chi(\overline{h}_0 e_0)$, and $||g_\delta||_E \ge \min\{||\overline{h}_0||_E, ||\overline{h}_0 e_0||_E\}$. We know that $e_0 \in E - \{h_0\}$ and $h_0 \in H_{i_0} - \{1\}$. Hence, $1 \notin \{\overline{h}_0, \overline{h}_0 e_0\}$ and $g_\delta \neq 1$. There exists a unique $e_\star \in E^{\pm 1}$ such that $||g_\delta e_\star||_E = ||g_\delta||_E - 1$. Clearly, $e_\star \notin \{1, e_\delta\}$.

Let us review the graph of interest. In T, define $\lim_{T}(1)$ and $\operatorname{star}_{T}(1)$ as for T^* . For each $e \in E^{\pm 1} \cup \{1\}$, there exists a unique component $[e] \in \operatorname{star}_{T}(1)$ such that $e \in [e]$. Then the map $E^{\pm 1} \cup \{1\} \to \operatorname{star}_{T}(1), e \mapsto [e]$, is bijective. Fix an edge (e', e'') of $\operatorname{WH}_{*}(\bigcup_{I} H_{i} \operatorname{rel} E)$. Here, there exist $i \in I$, $h \in H_{i} - \{1\}$, and $g', g'' \in F$ such that $h = \overline{g'} \cdot \overline{e'} \cdot e'' \cdot g''$ with no $E^{\pm 1}$ -cancellation, g' = 1 if e' = 1, and g'' = 1 if e'' = 1. Thus, $e'g'h = e''g'', e'g'H_{i} = e''g''H_{i}, e' \cdot g' \in [e']$, and $e'' \cdot g'' \in [e'']$; it may happen that e' = 1 = g' and $[e'] = \{1\}$.

We now return to g_{δ} and e_{\star} . We see that $1 \in g_{\delta}[e_{\star}], \delta \subseteq g_{\delta} \operatorname{link}_{T}(1) \cup g_{\delta}[e_{\star}],$ and χ is constant on the vertex-set of each component of $T - (g_{\delta} \operatorname{link}_{T}(1) \cup g_{\delta}[e_{\star}]).$ We shall show that if $e_{\star} \notin \{e', e''\},$ then $\chi(g_{\delta}e') = \chi(g_{\delta}e'').$ As $e' \neq e_{\star}$, we see $1 \notin g_{\delta}[e']$ and χ maps the vertex-set of $g_{\delta}[e']$ to $\{\chi(g_{\delta}e')\}.$ As $g_{\delta}e'g' \in g_{\delta}[e'],$ we see $\operatorname{edge}(g_{\delta}e'g' \xrightarrow{\bullet(H_{i})} g_{\delta}e'g'H_{i}) \notin \operatorname{link}_{T^{*}}(1)$ and $\chi(g_{\delta}e'g') = \chi(g_{\delta}e').$ It follows that $\chi(g_{\delta}e') = \chi(g_{\delta}e'g') = \chi(g_{\delta}e'g'H_{i}) = \chi(g_{\delta}e''g''H_{i}) = \chi(g_{\delta}e''g'') = \chi(g_{\delta}e'').$

Let W denote the graph that is obtained from $WH_*(\bigcup_I H_i \text{ rel } E)$ by removing e_* and its incident edges. We have now proved that $\chi(g_{\delta}-)$ is constant on the vertex-sets of the components of W. Since 1 and e_{δ} are vertices of W such that $\chi(g_{\delta}1) \neq \chi(g_{\delta}e_{\delta})$, we see that W is not connected, and, hence, e_* is a Whitehead cut-vertex of $WH_*(\bigcup_I H_i \text{ rel } E)$.

3.3 Corollary. With Hypotheses 1.2, suppose that $W_{H_*}(Z \text{ rel } E)$ has no Whitehead cut-vertices. For each free-product factorization $\langle E | \rangle = \underset{i \in I}{*} H_i$ such that $Z \subseteq \bigcup_{i \in I} H_i$, the set E contains a basis of each H_i .

Proof. Let *i* range over *I*. Set $E_i := E \cap H_i$. Then the E_i are pairwise disjoint. As it contains $W_{H*}(Z \text{ rel } E)$, $W_{H*}(\bigcup_I H_i \text{ rel } E)$ has no Whitehead cut-vertices. By the contrapositive of Theorem 3.2, $E \subseteq \bigcup_I H_i$. Thus, $E = \bigcup_I E_i$. Hence, $\langle E | \rangle = \underset{i \in I}{*} \langle E_i \rangle \leq \underset{i \in I}{*} H_i = \langle E | \rangle$. It follows that $\langle E_i \rangle = H_i$ and, hence, E_i is a basis of H_i .

3.4 Whitehead's cut-vertex lemma. With Hypotheses 1.2, suppose that $W_{H_*}(Z \text{ rel } E_Z)$ has no Whitehead cut-vertices. If Z is a sub-basis of $\langle E | \rangle$, then $Z \subseteq E^{\pm 1}$. Hence, Z is a sub-basis of $\langle E | \rangle$ if and only if $Z \cap Z^{-1} = \emptyset$ and $Z \subseteq E^{\pm 1}$; in this event, $Z \cup (E - Z^{\pm 1})$ is a basis of $\langle E | \rangle$.

Proof. Let E' be a basis of $\langle E | \rangle$ that contains Z. A classic E'-length argument due to Nielsen shows that $E' \cap \langle E_Z \rangle$ is contained in some basis X of $\langle E_Z \rangle$; we shall mention Schreier's proof in Review 4.1. Now $\langle E_Z \rangle = \underset{x \in X}{*} \langle x \rangle$ and $Z \subseteq E' \cap \langle E_Z \rangle \subseteq X \subseteq \bigcup_{x \in X} \langle x \rangle$. By Corollary 3.3, E_Z contains a basis of each $\langle x \rangle$, necessarily $\{x\}$ or $\{\overline{x}\}$. Thus $E_Z^{\pm 1} \supseteq X \supseteq Z$.

We shall use the following strong form in the next section.

3.5 Corollary. If Z is a sub-basis of $\langle E | \rangle$ and $Z \not\subseteq E^{\pm 1}$, then there exists some $\mathbf{C} \in \text{CUTS}(E)$ such that $\text{WH}_*(Z \text{ rel } E) \subseteq \text{WH}_*(\mathbf{C})$ and $||Z^{\overline{\varphi}_{\mathbf{C}}}||_E < ||Z||_E$.

Proof. By the contrapositive of Lemma 3.4, $WH_*(Z \text{ rel } E_Z)$ has a Whitehead cut-vertex. The result now follows from Algorithm 2.3.

It remains to discuss free-product factors.

3.6 Review. • We now sketch a proof of a result of Kurosh: for any subgroups H and K of any group F, if K is a free-product factor of F, say F = K * L, then $H \cap K$ is a free-product factor of H.

We shall use Bass-Serre theory, although for our purposes the case $F = \langle E | \rangle$ and the graph-theoretic techniques of John R. Stallings [5] would suffice.

We may view BassSerre(F, (K, L)) as an H-tree, and then the vertex 1K can be extended to a fundamental H-transversal. The resulting graph of groups has $H \cap K$ as one of the vertex-groups and all the edge-groups are trivial. By another result of Bass and Serre, $H \cap K$ is a free-product factor of H. See, for example, [2, Theorem I.4.1].

It follows that, for any group, the set of all its free-product factors is closed under finite intersections.

• Recall Hypotheses 1.2 and set $F \coloneqq \langle E | \rangle$. Now |E| bounds the length of any strictly descending chain of free-product factors of F. Hence, the set of all the free-product factors of F is closed under arbitrary intersections.

In particular, CL(Z), the intersection of all the free-product factors of F containing Z, is the \subseteq -smallest free-product factor of F containing Z.

By Kurosh's result again, $\operatorname{CL}(Z) \cap \langle E_Z \rangle$ is a free-product factor of $\langle E_Z \rangle$. However, $\langle E_Z \rangle$ contains $\operatorname{CL}(Z)$, since $\langle E_Z \rangle$ is a free-product factor of F which contains Z. Thus, $\operatorname{CL}(Z)$ is a free-product factor of $\langle E_Z \rangle$. In particular, the bases of $\operatorname{CL}(Z)$ are the minimal-size supports of Z with respect to bases of F.

3.7 Stong's cut-vertex lemma. With Hypotheses 1.2, suppose that $W_{H_*}(Z \text{ rel } E_Z)$ has no Whitehead cut-vertices. Then E_Z is a basis of $C_L(Z)$, and, for each free-product factorization $C_L(Z) = \underset{i \in I}{*} H_i$ such that $Z \subseteq \bigcup_{i \in I} H_i$, the set E_Z contains a basis of each H_i .

Proof. We saw in Review 3.6 that there exists some free-product factorization $\langle E_Z \rangle = \operatorname{CL}(Z) * K$, and it is clear that $Z \subseteq \operatorname{CL}(Z) \cup K$. By Corollary 3.3, E_Z contains some basis E' of $\operatorname{CL}(Z)$. Since $Z \subseteq \operatorname{CL}(Z) = \langle E' \rangle$, we see that $\operatorname{supp}(Z \operatorname{rel} E) \subseteq E'$, that is, $E_Z \subseteq E'$. Hence, E_Z is a basis of $\operatorname{CL}(Z)$. The result now follows from Corollary 3.3.

4 A strengthened Clifford-Goldstein algorithm

Clifford and Goldstein [1] produced an ingenious algorithm which returns an element of $\langle Z \rangle$ that lies in a basis of $\langle E | \rangle$ or reports that no element of $\langle Z \rangle$ lies in a basis of $\langle E | \rangle$. They used Whitehead's three-manifold techniques to construct a sufficiently large finite set of finitely generated subgroups of $\langle E | \rangle$ whose elements of sufficiently bounded *E*-length give the desired information.

In this section, we restructure their argument, bypassing the topology and obtaining a less complicated, more powerful algorithm which yields as output a basis E'' of $\langle E | \rangle$ which maximizes $|E'' \cap \langle Z \rangle|$. In particular, $E'' \cap \langle Z \rangle = \emptyset$ if and only if no element of $\langle Z \rangle$ lies in a basis of $\langle E | \rangle$. We construct a smaller sufficiently large finite set of finitely generated subgroups of $\langle E | \rangle$ whose intersections with E give the desired information.

To fix notation, we sketch the proof of Schreier [4, publ. 1927] that subgroups of free groups are free. The finitely generated case had been proved by J. Nielsen [3, publ. 1921, in Danish].

4.1 Review. With Hypotheses 1.2, set $F := \langle E | \rangle$ and T := Cayley(F, E); see Review 3.1. Let H be a subgroup of F. The vertices of the *Schreier graph* $H \setminus T$ are the cosets Hg, $g \in F$, the basepoint is H1, and we write $\text{edge}(v \xrightarrow{\bullet(e)} ve) := (v, e) \in (H \setminus F) \times E$. The graph $H \setminus T$ is connected. Let $\pi(H \setminus T, H1)$ denote the fundamental group of $H \setminus T$ at the basepoint H1. Each (reduced) $H \setminus T$ -path from H1 to itself will be viewed as a (reduced) $E^{\pm 1}$ -expression for some element of H; for example, we would view

 $(H1 \xrightarrow{\bullet(e_1)} He_1 \xleftarrow{\bullet(e_2)} He_1\overline{e}_2 \xrightarrow{\bullet(e_3)} He_1\overline{e}_2e_3 = H1)$

as the $E^{\pm 1}$ -expression $e_1\overline{e_2}e_3$ for an element of H. Hence, we may identify $\pi(H\backslash T, H1)$ with H.

Choose a maximal subtree Y' of $H \setminus T$ and let Y'' denote the complement of Y' in $H \setminus T$; then Y'' is a set of edges. Each element y'' of Y'' determines the element of $\pi(H \setminus T, H1)$ that travels in Y' from H1 to the initial vertex of y'', travels along y'', and then travels in Y' from the terminal vertex of y'' to H1. By letting y'' range over Y'', we get a subset S of $\pi(H \setminus T, H1)$. By collapsing the tree Y' to a vertex, we find that S freely generates $\pi(H \setminus T, H1)$ (= H).

The vertices and edges involved in S form a connected basepointed subgraph of $H \setminus T$ denoted core_{*}(H rel E). An alternative description is that core_{*}(H rel E) consists of those vertices and edges that are involved in the reduced $H \setminus T$ -paths from H1 to itself. Thus, $\pi(\operatorname{core}_*(H \operatorname{rel} E), H1) = H$ and core_{*}(H rel E) is the \subseteq -smallest subgraph of $H \setminus T$ with this property.

For each $h \in E \cap H$, it is clear that $edge(H1 \xrightarrow{\bullet(h)} Hh = H1)$ is not in the tree Y', and, hence, $h \in S$. Thus, $E \cap H \subseteq S$. (I am indebted to Clifford and Goldstein for this paragraph.)

Warren Dicks

4.2 Algorithm. Stallings' *core algorithm* [5, Algorithm 5.4] has the following structure.

With Hypotheses 1.2, we shall suppress the information that the vertices of $\operatorname{core}_*(\langle Z \rangle \operatorname{rel} E)$ are certain cosets, and we shall build a basepointed *E*-labelled graph, denoted modelcore_{*}($\langle Z \rangle$ rel *E*), that has an abstract set as vertex-set and is isomorphic to $\operatorname{core}_*(\langle Z \rangle \operatorname{rel} E)$ as basepointed *E*-labelled graph.

For each $z \in Z$, we easily build modelcore_{*}($\langle z \rangle$ rel E) as a basepointed E-labelled lollipop graph, possibly trivial, using the reduced $E^{\pm 1}$ -expression for z.

We next amalgamate all these lollipop graphs at their basepoints. Throughout the construction, each edge will be assigned an expression of the form $\operatorname{edge}(v \xrightarrow{\bullet(e)} w)$ with v, w vertices and $e \in E$, but, for the moment, the expression need not determine the edge. While possible, we identify some distinct pair of edges with expressions $\operatorname{edge}(v \xrightarrow{\bullet(e)} w)$ and $\operatorname{edge}(v' \xrightarrow{\bullet(e)} w')$ where v = v' or w = w' or both; identifying the edges entails identifying w with w' or v with v'or neither, respectively. When no such pair of distinct edges is left, the procedure has yielded a basepointed E-labelled graph isomorphic to $\operatorname{core}_*(\langle Z \rangle \operatorname{rel} E)$; here, expressions $\operatorname{edge}(v \xrightarrow{\bullet(e)} w)$ do determine edges. \Box

Stallings gave the name *folding* to the foregoing edge-identifying process. The process itself had long been used unnamed, notably by Lyndon in his work on planar diagrams, where each nontrivial lollipop graph has a two-cell attached making a contractible CW-complex.

We now give the (strange) key construction of [1, Theorem 1].

4.3 Notation. With Hypotheses 1.2, fix $\mathbf{C} = ({}_0D, {}_1D, e_{\star}) \in \text{CUTS}(E)$, and set $F := \langle E | \rangle, \eta := \eta_{\mathbf{C}}, d_{\star} := e_{\star}^{2\eta-1}$, and $\varphi := \varphi_{\mathbf{C}}$; see Notation 2.1.

We first construct an F-map $\psi_{\mathbf{C}}$ from the edge-set of $T := \operatorname{Cayley}(F, E)$ to the edge-set of $T' := \operatorname{Cayley}(F, E^{\varphi})$. For any $\operatorname{edge}(g \xrightarrow{\bullet(e)} ge) \in F \times E$, there exists a unique $(\alpha, \beta) \in \{0, 1\}^{\times 2}$ such that $e \in {}_{\alpha}E_{\beta}$ and $e^{\varphi} = d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta}$; if $e^{\pm 1} \neq e_{\star}^{\pm 1}$, these two conditions are equivalent, while if $e^{\pm 1} = e_{\star}^{\pm 1}$, the two conditions together say that $\alpha = \beta = \eta$. We set $(\operatorname{edge}(g \xrightarrow{\bullet(e)} ge))^{\psi_{\mathbf{C}}} := \operatorname{edge}(g\overline{d}_{\star}^{\alpha} \xrightarrow{\bullet(e^{\varphi})} ge\overline{d}_{\star}^{\beta})$; we emphasize that no action of $\psi_{\mathbf{C}}$ on vertices is being defined. It is clear that $\psi_{\mathbf{C}}$ is an F-map.

Let H be a finitely generated subgroup of F. Then $\psi_{\mathbf{C}}$ induces a set map from the edge-set of $H \setminus T$ to the edge-set of $H \setminus T'$, and the image of the edge-set of core_{*}(H rel E) under this induced map is then the edge-set of a unique subgraph X of $H \setminus T'$ with the full vertex-set, $H \setminus F$. Let $K := \pi(X, H1) \leq \pi(H \setminus T', H1)$. We may identify the latter group with H, where $(H \setminus T')$ -paths are $(E^{\varphi})^{\pm 1}$ -expressions. We set $\partial_{\mathbf{C}} H := K^{\overline{\varphi}} \leq H^{\overline{\varphi}}$. Recall that modelcore_{*}(H rel E) was constructed in Algorithm 4.2; we shall be viewing $\partial_{\mathbf{C}}$ as a graph operation that converts modelcore_{*}(H rel E) into modelcore_{*}($\partial_{\mathbf{C}} H$ rel E).

- **4.4 Lemma.** With the foregoing notation, the following hold for $\partial_{\mathbf{C}} H \leq H^{\overline{\varphi}_{\mathbf{C}}}$. (i) modelcore_{*}($\partial_{\mathbf{C}} H$ rel E) may be constructed algorithmically.
- (ii) core_{*}(H rel E) has at least as many edges as core_{*}($\partial_{\mathbf{C}}H$ rel E).
- (iii) For each $z \in H$, if $WH_*(\{z\} \text{ rel } E) \subseteq WH_*(\mathbb{C})$, then $z^{\overline{\varphi}_{\mathbb{C}'}} \in \partial_{\mathbb{C}} H$.
- (iv) If Y is any sub-basis of $\langle E | \rangle$ such that $Y \subseteq H$ and $Y \not\subseteq E^{\pm 1}$, then there exists some $\mathbf{C}' \in \text{CUTS}(E)$ such that $Y^{\overline{\varphi}_{\mathbf{C}'}} \subseteq \partial_{\mathbf{C}'}H$ and $||Y^{\overline{\varphi}_{\mathbf{C}'}}||_E < ||Y||_E$.

Proof. (i). Since $K^{\overline{\varphi}} = \partial_{\mathbf{C}} H$, there is a natural graph isomorphism that maps core_{*}(K rel E^{φ}) to core_{*}($\partial_{\mathbf{C}} H$ rel E), changing each $Kg \xrightarrow{\bullet(e^{\varphi})} Kg(e^{\varphi})$ to $K^{\overline{\varphi}}g^{\overline{\varphi}} \xrightarrow{\bullet(e)} K^{\overline{\varphi}}g^{\overline{\varphi}}e$. Hence, there is a natural graph isomorphism that maps modelcore_{*}(K rel E^{φ}) to modelcore_{*}($\partial_{\mathbf{C}} H$ rel E), changing each $v \xrightarrow{\bullet(e^{\varphi})} w$ to $v \xrightarrow{\bullet(e)} w$; the labels on the non-basepoint vertices are irrelevant. Thus, it suffices to algorithmically construct modelcore_{*}(K rel E^{φ}) from modelcore_{*}(H rel E).

If $d_{\star} \in E$, resp. $\overline{d}_{\star} \in E$, we say that a vertex v of modelcore_{*} $(H \operatorname{rel} E)$ has a neighbour $v\overline{d}_{\star}$ if an edge of the form $\operatorname{edge}(w \xrightarrow{\bullet(d_{\star})} v)$, resp. $\operatorname{edge}(v \xrightarrow{\bullet(\overline{d}_{\star})} w)$, lies in modelcore_{*} $(H \operatorname{rel} E)$; in this event, we say that w is $v\overline{d}_{\star}$. We simultaneously add to modelcore_{*} $(H \operatorname{rel} E)$, for every vertex v that does not have a neighbour $v\overline{d}_{\star}$, a valence-zero vertex with label $v\overline{d}_{\star}$.

Next, in modelcore_{*}(*H* rel *E*) adorned with the valence-zero vertices, we simultaneously replace each edge($v \xrightarrow{\bullet(e)} w$) with edge($v\overline{d}_{\star}^{\alpha} \xrightarrow{\bullet(e^{\varphi})} w\overline{d}_{\star}^{\beta}$) for the unique $(\alpha, \beta) \in \{0, 1\}^{\times 2}$ such that $e \in {}_{\alpha}E_{\beta}$ and $e^{\varphi} = d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta}$. This particular operation alters incidence maps and edge labellings, but not the vertex-set or the edge-set.

In the resulting finite graph, we then keep only the component that has the basepoint. We next successively delete non-basepoint, valence-one vertices and their (unique) incident edges, while possible. When this is no longer possible, we have constructed modelcore_{*}(K rel E^{φ}) algorithmically.

(ii). It is clear from the constructions that $\operatorname{core}_*(H \operatorname{rel} E)$ has at least as many edges as $\operatorname{core}_*(K \operatorname{rel} E^{\varphi})$, which in turn has the same number of edges as $\operatorname{core}_*(\partial_{\mathbf{C}}H \operatorname{rel} E)$.

(iii). Consider any expression $Hg \xrightarrow{\bullet(e)} Hge$ corresponding to an edge or inverse edge in core_{*}(H rel E), and consider any $(\alpha, \beta) \in \{0, 1\}^{\times 2}$ such that $e \in {}_{\alpha}D_{\beta}$. Then $d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta} \in \{e^{\varphi}, 1\}$, for, if $d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta} \neq e^{\varphi}$, then either $e = e_{\star}$, $\alpha = 1-\eta$, $\beta = \eta$, $d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta} = d_{\star}^{1-\eta-\eta}e_{\star} = 1$, or $e = \overline{e}_{\star}$, $\alpha = \eta$, $\beta = 1-\eta$, $d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta} = d_{\star}^{\eta-1+\eta}\overline{e}_{\star} = 1$. This means that the expression $Hg\overline{d}_{\star}^{\alpha} \xrightarrow{\bullet(d_{\star}^{\alpha}e\overline{d}_{\star}^{\beta})} Hge\overline{d}_{\star}^{\beta}$ corresponds to an edge, inverse edge, or equality in the graph X of Notation 4.3.

Suppose that $z \in H$ and let $e_1 e_2 \cdots e_n$ represent the reduced $E^{\pm 1}$ -expression for z. We then have a corresponding reduced $H \setminus T$ -path from H1 to itself, which we may write in $H \setminus \text{Cayley}(F, E^{\pm 1})$ as

 $H1 \xrightarrow{\bullet(e_1)} He_1 \xrightarrow{\bullet(e_2)} He_1e_2 \xrightarrow{\bullet(e_3)} \cdots \xrightarrow{\bullet(e_n)} He_1e_2 \cdots e_n = Hz = H1.$

Warren Dicks

The $H \setminus T$ -path must then stay within the subgraph core_{*}(H rel E).

Suppose further that $W_{H_*}(\{z\} \text{ rel } E) \subseteq W_{H_*}(\mathbf{C})$. This means that there exists a (unique) set map $\{0, 1, \ldots, n\} \to \{0, 1\}, i \mapsto \chi_i$, such that $e_i \in \chi_{i-1}D_{\chi_i}$, $i = 1, 2, \ldots, n$, and $\chi_0 = \chi_n = 0$. In our core_{*}(H rel E)-path, let us change each vertex $He_1 \cdots e_i$ to $He_1 \cdots e_i \overline{d}_{\star}^{\chi_i}$ and each step $He_1 \cdots e_{i-1} \xrightarrow{\bullet(e_i)} He_1 \cdots e_{i-1}e_i$ to $He_1 \cdots e_{i-1} \overline{d}_{\star}^{\chi_{i-1}} \xrightarrow{\bullet(d_{\star}^{\chi_{i-1}}e_i \overline{d}_{\star}^{\chi_i})} He_1 \cdots e_{i-1}e_i \overline{d}_{\star}^{\chi_i}$, which we have seen corresponds to an edge, inverse edge, or equality in X. We thus obtain an X-path from H1 to itself that reads an $((E^{\varphi})^{\pm 1} \cup \{1\})$ -expression for z. This shows that $z \in \pi(X, H1) = K$, as desired.

(iv). By Corollary 3.5, there exists $\mathbf{C}' \in \text{CUTS}(E)$ such that $||Y^{\overline{\varphi}_{\mathbf{C}'}}||_E < ||Y||_E$ and WH_{*}(Y rel E) \subseteq WH_{*}(\mathbf{C}'). By (iii), $Y^{\overline{\varphi}_{\mathbf{C}'}} \subseteq \partial_{\mathbf{C}'}H$.

We now give a construction that is a somewhat less complicated variant of the algorithm of Clifford and Goldstein [1].

4.5 Notation. With Hypotheses 1.2, let \mathcal{F} denote the set of all finitely generated subgroups of $\langle E | \rangle$. Let Γ denote the graph whose vertex-set is \mathcal{F} and whose edge-set is $\mathcal{F} \times \text{CUTS}(E)$ where each edge $(H, \mathbf{C}) \in \mathcal{F} \times \text{CUTS}(E)$ has initial vertex H and terminal vertex $\partial_{\mathbf{C}} H$; see Notation 4.3.

Set $G \coloneqq \langle Z \rangle \in \mathcal{F}$. Let $(G \blacktriangleleft)$ denote the subgraph of Γ that radiates out from G, that is, $(G \blacktriangleleft)$ is the smallest subgraph of Γ that has G as a vertex and is closed in Γ under the operation of adding to each vertex H each outgoing edge (H, \mathbb{C}) and its terminal vertex $\partial_{\mathbb{C}} H$.

For each $n \ge 0$, each element $(\mathbf{C}_i)_{i=1}^n$ of $(\text{CUTS}(E))^{\times n}$ determines the oriented $(G\blacktriangleleft)$ -path with the edge-sequence $(H_i \xrightarrow{(H_i, \mathbf{C}_i)} H_{i+1})_{i=1}^n$ where $H_1 = G$ and $H_{i+1} = \partial_{\mathbf{C}_i} H_i$ for $i = 1, \ldots, n$. To simplify notation, we shall say that $(\mathbf{C}_i)_{i=1}^n$ itself is an oriented $(G\blacktriangleleft)$ -path with initial vertex G.

We usually think of a vertex H of $(G\blacktriangleleft)$ as the graph modelcore_{*}(H rel E), for ease of recognition. We shall see that we are interested in finding a vertex that maximizes the number of loops at the basepoint.

4.6 Theorem. With the foregoing notation, the following hold.

- (i) $(G \blacktriangleleft)$ is an algorithmically constructible finite graph whose vertices are viewed as finite, E-labelled, basepointed graphs.
- (ii) For each vertex H of $(G\blacktriangleleft)$, there is an algorithmically constructible oriented $(G\blacktriangleleft)$ -path $(\mathbf{C}_i)_{i=1}^n$ from G to H, $H = \partial_{\mathbf{C}_n} \cdots \partial_{\mathbf{C}_1} G \leqslant G^{\overline{\varphi}_{\mathbf{C}_1} \cdots \overline{\varphi}_{\mathbf{C}_n}}$, and $(E \cap H)^{\varphi_{\mathbf{C}_n} \cdots \varphi_{\mathbf{C}_1}} \subseteq E'' \cap G$, where $E'' := E^{\varphi_{\mathbf{C}_n} \cdots \varphi_{\mathbf{C}_1}}$.
- (iii) For each basis E'' of $\langle E | \rangle$, there exists some vertex H of $(G \blacktriangleleft)$ such that $|E \cap H| \ge |E'' \cap G|$.

Proof. (i). For each $H \in \mathcal{F}$, if *n* denotes the number of edges in core_{*}(*H* rel *E*), it is clear from Review 4.1 that *H* can be generated by *n*-or-less elements of $\langle E | \rangle$ of *E*-length 2*n*-or-less. By Lemma 4.4(ii), (*G* \triangleleft) is finite. By Lemma 4.4(i), we

may use a depth-first search to construct a maximal subtree of $(G \blacktriangleleft)$. We then add the missing edges of $(G \blacktriangleleft)$, although this is optional for our purposes.

(ii) is clear.

(iii). It follows from Lemma 4.4(iv) that there exists some $(\mathbf{C}_i)_{i=1}^n$ such that $(E'' \cap G)^{\overline{\varphi}_{\mathbf{C}_1}\overline{\varphi}_{\mathbf{C}_2}\cdots\overline{\varphi}_{\mathbf{C}_n}} \subseteq E^{\pm 1} \cap \partial_{\mathbf{C}_n}\cdots\partial_{\mathbf{C}_2}\partial_{\mathbf{C}_1}G.$

We now construct a basis E'' of $\langle E | \rangle$ which maximizes $|E'' \cap \langle Z \rangle|$.

4.7 Algorithm. Recall Hypotheses 1.2.

- Set $G := \langle Z \rangle$ and construct modelcore_{*}(G rel E); see Algorithm 4.2.
- Construct $(G \triangleleft)$ from modelcore_{*}(G rel E); see Theorem 4.6(i).

• In $(G \triangleleft)$, find a vertex H maximizing the number of loops at the basepoint of modelcore_{*}(H rel E), that is, maximizing $|E \cap H|$.

• Find an oriented $(G \triangleleft)$ -path $(\mathbf{C}_i)_{i=1}^n$ from G to H; see Theorem 4.6(ii).

• Return $E'' := E^{\varphi_{\mathbf{C}_n} \cdots \varphi_{\mathbf{C}_2} \varphi_{\mathbf{C}_1}}$, a basis of $\langle E | \rangle$ which maximizes $|E'' \cap \langle Z \rangle|$ by Theorem 4.6(ii),(iii).

References

- A. Clifford and R. Z. Goldstein, Subgroups of free groups and primitive elements, J. Group Theory 13 (2010), 601–611.
- Warren Dicks and M. J. Dunwoody, Groups acting on graphs, Camb. Stud. Adv. Math. 17, CUP, 1989. xvi+283 pp.

Errata at: http://mat.uab.cat/~dicks/DDerr.html

- J. Nielsen, Om Regnung med ikke kommutative Faktoren og dens Anvendelse i Gruppeteorien, Math. Tidsskrift B 107 (1921), 77–94.
- [4] O. Schreier, Die Untergruppen der freien Gruppen, Abh. Math. Univ. Hamburg 5 (1927), 161–183.
- [5] John R. Stallings, Topology of finite graphs, Invent. Math. 71 (1983), 551– 565.
- [6] Richard Stong, *Diskbusting elements of the free group*, Math. Res. Lett. 4 (1997), 201–210.
- [7] Edward C. Turner, Test words for automorphisms of free groups, Bull. London Math. Soc. 28 (1996), 255–263.
- [8] J. H. C. Whitehead, On certain sets of elements in a free group, Proc. London Math. Soc. 41 (1936), 48–56.
- J. H. C. Whitehead, On equivalent sets of elements in a free group, Ann. of Math. 37 (1936), 782–800.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA (BARCELONA), SPAIN email: dicks@mat.uab.cat URL: http://mat.uab.cat/~dicks/