ON SINGULARITY AS A FUNCTION OF TIME OF A CONDITIONAL DISTRIBUTION OF AN EXIT TIME

N.V. KRYLOV

ABSTRACT. We establish the singularity with respect to Lebesgue measure as a function of time of the conditional probability that the sum of two one-dimensional Brownian motions will exit from the unit interval before time t, given the trajectory of the second Brownian motion up to the same time. On the way of doing so we show that if one solves the one-dimensional heat equation with zero condition on a trajectory of a one-dimensional Brownian motion, which is the lateral boundary, then for each moment of time with probability one the normal derivative of the solution is zero, provided that the diffusion of the Brownian motion is sufficiently large.

1. MAIN RESULTS

Fix some constants $\sigma, \sigma_1 > 0$ and consider the equation

$$x_t = x_0 + \sigma_1 w_t + \sigma b_t, \quad y_t = b_t, \tag{1.1}$$

where w and b are independent one-dimensional standard Wiener processes, x_0 is independent of the couple (w, b) and has density $\pi_0 \in C_0^{\infty} = C_0^{\infty}(G)$ concentrated on G, where G = (0, 1). Define

$$\mathcal{F}_t^{b.} = \sigma(b_s : s \le t), \quad \tau = \inf\{t \ge 0 : x_t \notin G\},$$
$$A_t = P(\tau \le t \mid \mathcal{F}_t^{b.}).$$

Here is our main result.

Theorem 1.1. There is a continuous and nondecreasing modification of A_t which is singular with respect to Lebesgue measure, the latter provided that σ_1/σ is sufficiently small.

The process A_t in a more general framework arose in [6] as the main process governing the conditional distribution of a signal process x_t at the first time when it exists from a given domain. In [6] the observations y_t were only available until the first exit time of x_t from the domain. It turns out that in the setting of (1.1) the conditional and the so-called unnormalized conditional distributions of x_t before it exits from G given $y_s, 0 \le s \le t$,

²⁰⁰⁰ Mathematics Subject Classification. 60H15, 93E11.

Key words and phrases. Stochastic partial differential equations, heat equation in domains with irregular lateral boundaries, filtering of partially observable diffusion processes.

The author was partially supported by NSF Grant DNS-1160569.

coincide. These unnormalized conditional distributions are known to satisfy some linear stochastic partial differential equations and then the properties of A_t can be recovered from some properties of solutions of these equations.

To be more precise for $(t, x) \in (0, \infty) \times (0, 1)$ consider the following (filtering) equation

$$d\pi_t(x) = (1/2)aD^2\pi_t(x)\,dt - \sigma D\pi_t(x)\,db_t,$$
(1.2)

where $a = \sigma_1^2 + \sigma^2$, with initial condition $\pi_0(x)$ and zero lateral condition.

To explain in which sense we understand this equation, the initial condition, and the boundary condition, we need some notation. Introduce the space $W_2^1 = W_2^1(G)$ as the closure of the set of continuously differentiable functions in \overline{G} in the norm

$$||u||_{W_2^1} = ||u||_{L_2} + ||Du||_{L_2}$$

where Du is the derivative of u and $L_2 = L_2(G)$, and we introduce $\overset{0}{W_2^1} = \overset{0}{W_2^1}(G)$ as the closure of $C_0^{\infty} = C_0^{\infty}(G)$ in the above norm. Denote by \mathcal{P}^{b} the predictable σ -field in $\Omega \times (0, \infty)$ associated with the

Denote by \mathcal{P}^{b} the predictable σ -field in $\Omega \times (0, \infty)$ associated with the filtration $\{\mathcal{F}_t^{b}\}$. For $T \in (0, \infty)$ introduce

$$G_T = (0,T) \times G, \quad \overset{0}{W}{}_2^1(G_T) = L_2(\Omega \times (0,T), \mathcal{P}^{b}, \overset{0}{W}{}_2^1).$$

We are looking for a function $\pi_t(x)$ which is a generalized function on G for each (ω, t) such that $\pi \in \bigcap_T W_2^0(G_T)$ and for each $\zeta \in C_0^\infty$ with probability one for all $t \in [0, \infty)$ it holds that

$$(\pi_t, \zeta) = (\pi_0, \zeta) - \int_0^t (1/2) (aD\pi_s, D\zeta) \, ds - \int_0^t (\sigma D\pi_s, \zeta) \, db_s, \qquad (1.3)$$

where we use the notation

$$(f,g) = \int_G f(x)g(x) \, dx.$$

Observe that all expressions in (1.3) are well defined due to the fact that the coefficients of π and of $D\pi$ are constant and

$$\pi, D\pi \in \mathcal{L}_2(T) := L_2(\Omega \times (0, T), \mathcal{P}^{b}, L_2)$$

for any $T \in (0, \infty)$.

Recall that by assumption $\pi_0 \in C_0^{\infty}$.

Theorem 1.2. In the class $\bigcap_T \overset{\circ}{W}{}_2^1(G_T)$ there exists a unique solution π_t of equation (1.2) with initial condition π_0 . In addition, $\pi_t \ge 0$ for all $t \in [0, \infty)$ (a.s.). With probability one π_t is continuous in $L_1 = L_1(G)$ and in L_2 .

The existence, uniqueness, and the (a.s.) continuity in L_2 of π is a classical result proved in many places in a variety of settings (see, for instance, [7], [5], [8], and the references therein). That π_t is (a.s.) continuous as an L_1 function follows from its L_2 -continuity and the boundedness of G. The fact that $\pi \geq 0$ follows from the maximum principle (see, for instance, Theorem 1.1 of [3]) and the fact that, if $u \in \overset{0}{W_{2}^{1}}$, then $u^{+} \in \overset{0}{W_{2}^{1}}$.

The connection of A_t to π_t is established on the basis of Lemma 1.8 of [6], which our situation reads as follows.

Lemma 1.3. For any Borel bounded or nonnegative function ϕ on G and $t \in [0, \infty)$ we have (a.s.)

$$E\{I_{\tau>t}\phi(x_t) \mid \mathcal{F}_t^{b.}\} = (\pi_t, \phi).$$
(1.4)

In particular, for each $t \in [0, \infty)$ (a.s.)

$$P\{\tau > t \mid \mathcal{F}_t^{b.}\} = (\pi_t, 1).$$
(1.5)

Finally, (a.s.) we have $(\pi_t, 1) > 0$ for all $t \in [0, \infty)$.

By Lemma 1.3 for any $t \in [0, \infty)$

$$P(\tau \le t \mid \mathcal{F}_t^{b.}) = 1 - (\pi_t, 1)$$

(a.s.) and by Theorem 1.2 the right-hand side is continuous in t (a.s). Also it turns out (see [6]) that the process $(\pi_t, 1)$ is decreasing (a.s). Therefore, in Theorem 1.1 by the modification of A_t , which we identify with the original A_t , we mean $1 - (\pi_t, 1)$.

Observe that if in (1.3) we were allowed to first integrate by parts to replace

$$(D\pi_s, D\zeta)$$
 with $-(D^2\pi_s, \zeta)$

and then in the so modified version of (1.3) take $\zeta \equiv 1 \ (\notin C_0^{\infty})$, then we would formally obtain that

$$A_t = 1 - (\pi_t, 1) = (1/2)a \int_0^t [D\pi_s(0) - D\pi_s(1)] \, ds.$$
 (1.6)

This shows that A_t is related to the normal derivative of π_s on the boundary of G, investigating which is done on the basis of a different description of π_s .

We are going to state our second main result, which is about solutions of the heat equation in curvilinear cylinders whose lateral boundary is a trajectory of a Wiener process.

Theorem 1.4. For almost any ω there exists a unique function $u_t(x)$ defined, bounded, and continuous in the closure of

$$\Gamma(b_{\cdot}) = \{(t, x) : t \ge 0, \quad x \in (\sigma b_t, 1 + \sigma b_t)\}$$

such that it is infinitely differentiable with respect to (t, x) in $\Gamma(b)$, satisfies there the equation

$$\partial_t u_t = (1/2)\sigma_1^2 D^2 u \tag{1.7}$$

and satisfies the conditions $u_0(x) = \pi_0(x)$, $x \in [0,1]$, and $u_t(\sigma b_t) = u_t(1 + \sigma b_t) = 0$, $t \ge 0$. Furthermore, if σ_1/σ is sufficiently small, then for any

 $t \in [0,\infty)$

4

$$\lim_{x \downarrow \sigma b_t} \frac{u_t(x)}{x - \sigma b_t} = \lim_{x \uparrow 1 + \sigma b_t} \frac{u_t(x)}{1 + \sigma b_t - x} = 0$$
(1.8)

almost surely, so that the derivative of $u_t(x)$ on the boundary of $\Gamma(b_t)$ is zero (a.s.) for any fixed t.

Finally, with probability one

$$\int_{G} |u_t(x - \sigma b_t) - \pi_t(x)|^2 \, dx = 0 \tag{1.9}$$

for all $t \ge 0$, so that $u_t(x - \sigma b_t)$ is a modification of $\pi_t(x)$, and for this modification, for any $t \in [0, \infty)$,

$$\lim_{x \downarrow 0} \frac{\pi_t(x)}{x} = \lim_{x \uparrow 1} \frac{\pi_t(x)}{1 - x} = 0$$
(1.10)

almost surely if σ_1/σ is sufficiently small.

The last statement of Theorem 1.4 makes the representation (1.6) questionable, and, even though there is a limit procedure showing that (1.6) holds in a generalized sense similar to that of the local time for Brownian motion (see [6]), one would rightfully suspect that A_t is not absolutely continuous with respect to t.

The last statement of Theorem 1.4 should not make the reader over optimistic about the continuity properties of $\pi_t(x)$ in x near the boundary of G (see Remark 1.1).

Still the following theorem will be easily derived from known results. Take $\alpha \in (0, 1)$ and $c \in (0, \infty)$ and introduce

$$p = p(c) := P(\sup_{t \le 1} w_t - \inf_{t \le 1} w_t \ge c(\sqrt{2} - 1)/2), \quad r = r(\alpha, c) := \frac{\alpha(1 - p)}{p(1 - \alpha)},$$
$$\beta = \beta(\alpha, c) := 2\frac{(r - 1)p + 1}{r^{\alpha}} = 2\frac{1 - p}{1 - \alpha}r^{-\alpha}.$$

As is easy to see, for any $\alpha \in (0, 1)$, we have $p(c) \to 0$, $r(\alpha, c) \to \infty$, and $\beta(\alpha, c) \to 0$ as $c \to \infty$. It follows from [2] that there exists a function $\alpha(c)$, $c \in (0, \infty)$, with values in (0, 1) such that $\alpha(c) \to 0$ as $c \to \infty$ and $\alpha(c) \le \alpha$ for any α satisfying $\beta(\alpha, c) < 1$.

Next, take some constants $c\geq 0,\,d>0$ and for $x\in \mathbb{R}$ define

$$\tau_{d,x} = \inf\{t \ge 0 : d/\sqrt{2} + w_t = x\},\$$

$$\gamma(c,d,\delta) = P(\tau_{d,d} \land (\delta/2) < \tau_{d,-c}).$$

Theorem 1.5. Take the modification of π_t from Theorem 1.4. Then for any $t \in [0, \infty)$, c, d > 0, such that $\alpha(c\varepsilon) < 1$, where $\varepsilon = \sigma_1/\sigma$, and ν satisfying

$$0 \le \nu < \nu_0 := (1 - \alpha(c\varepsilon)) \log_2 \gamma^{-2}(c, d, 1),$$

we have that with probability one

$$\sup_{x \in (0,1)} \sup_{t \in [0,T]} \frac{\pi_t(x)}{x^{\nu}} < \infty.$$
(1.11)

Furthermore, there exists a constant $\nu \in (0,\infty)$ such that for any $T \in (0,\infty)$

$$E \sup_{x \in (0,1)} \sup_{t \in [0,T]} \frac{\pi_t(x)}{x^{\nu} (1-x)^{\nu}} < \infty.$$
(1.12)

Remark 1.1. The largest possible value of ν in (1.11) is unknown. However, Theorem 5.1 and Lemma 4.1 of [1] show that if we take a $\mu > 0$ and

$$\nu = (1+\mu)(2\pi\varepsilon^2)^{-1/2}e^{-1/(2\varepsilon^2)}$$

then for $\varepsilon = \sigma_1/\sigma$ small enough the left-hand side of (1.11) equals infinity with probability one. Therefore, the largest value of ν is extremely small if ε is small.

Remark 1.2. The fact that equation (1.8) holds (a.s.) does not contradict Remark 1.1, because (1.11) gives an estimate which is uniform with respect to t, and on almost each trajectory of b. there are points t such that $v_t(x)/x \to \infty$ as $x \downarrow 0$.

We prove Theorem 1.1 in Section 2 assuming that Theorems 1.4 and 1.5 are true. In Section 3 we prove Theorems 1.4 and 1.5. The first assertion of Theorem 1.4 and Itô's formula easily lead to the conclusion that $u_t(x - \sigma b_t)$ is a classical solution of (1.2) and the assertion concerning (1.9) is proved by showing that classical solutions coincide with generalized ones in a much more general situation in Section 4.

2. Proof of Theorem 1.1

We start by proving that for each $t_0 \in (0,\infty)$ and $t_n = t_0 + 1/n$ with probability one

$$E\left(\lim_{n \to \infty} \frac{1}{t_n - t_0} (A_{t_n} - A_{t_0}) \mid \mathcal{F}_{t_0}^{b.}\right) = 0.$$
 (2.1)

Observe that for any $\zeta \in C_0^\infty$ and $t>t_0$

$$(\pi_t, \zeta) = (\pi_{t_0}, \zeta) + \int_{t_0}^t (\pi_s, (1/2)aD^2\zeta) \, ds + \int_{t_0}^t (\pi_s, \sigma D\zeta) \, db_s, \qquad (2.2)$$

where and below we are dealing with the modification of π_t from Theorem 1.4. We multiply both parts of this equation by the indicator function of a set $F \in \mathcal{F}_{t_0}^{b}$ and then take the expectations of both parts. Then by denoting

$$\phi_t^F(x) = E\pi_t(x)I_F$$

we find

$$(\phi_t^F, \zeta) = (\phi_{t_0}^F, \zeta) + \int_{t_0}^t (\phi_s^F, (1/2)aD^2\zeta) \, ds.$$
(2.3)

Observe that $\phi_t^F(x)$ is continuous in \overline{G}_{∞} because such is π which is in addition bounded. Estimate (1.12) shows that $\phi_t^F(x) \to 0$ as $x \to \{0,1\}$,

 $x \in (0,1), t \geq 0$. Thus ϕ_t^F is a continuous in $[t_0,\infty) \times \overline{G}$ weak solution of the equation

$$\partial_t \eta_t = (1/2)aD^2 \eta_t. \tag{2.4}$$

By uniqueness of such solutions, ϕ^F_t is a classical solution of this equation with zero boundary data.

By the maximum principle we have $\phi_t^F \leq \psi_t^F$, $t \geq t_0$, where ψ_t^F is defined as a unique bounded classical solution of (2.4) for $t \geq t_0$, x > 0, with initial data $\psi_{t_0}^F(x) = \phi_{t_0}^F(x)I_{(0,1)}(x)$ and zero boundary condition. The following explicit representation for such solutions is well known:

$$\psi_t^F(x) = \frac{1}{\sqrt{2\pi a(t-t_0)}} \int_0^1 \phi_{t_0}^F(y) \exp\left[-\frac{(x-y)^2}{2a(t-t_0)}\right] - \exp\left[-\frac{(x+y)^2}{2a(t-t_0)}\right] dy$$

Furthermore,

$$E(A_t - A_{t_0})I_F = \int_0^1 [\phi_t^F(x) - \phi_{t_0}^F(x)] \, dx \le \int_0^\infty [\psi_t^F(x) - \phi_{t_0}^F(x)I_{(0,1)}(x)] \, dx$$

Observe that

$$\partial_t \int_0^\infty [\psi_t^F(x) - \phi_{t_0}^F(x)I_{(0,1)}(x)] \, dx = (1/2)a \int_0^\infty D^2 \psi_t^F(x) \, dx$$
$$= (1/2)aD\psi_t^F(0),$$

so that

$$E(A_t - A_{t_0})I_F \le \int_{t_0}^t \frac{1}{\sqrt{2\pi a(r - t_0)^3}} \int_0^1 y \phi_{t_0}^F(y) e^{-y^2/(2ar - 2at_0)} \, dy dr$$
$$= \int_{t_0}^t \frac{2a}{\sqrt{2\pi a(r - t_0)}} \int_0^{1/\sqrt{2ar - 2at_0}} \phi_{t_0}^F(x\sqrt{2ar - 2at_0}) x e^{-x^2} \, dx dr,$$

which after taking into account the arbitrariness of $F \in \mathcal{F}_{t_0}^{b}$ leads to

$$E(A_t - A_{t_0} \mid \mathcal{F}_{t_0}^{b})$$

$$\leq \int_{t_0}^t \frac{2a}{\sqrt{2\pi a(r - t_0)}} \int_0^\infty \pi_{t_0} (x\sqrt{2ar - 2at_0}) I_{x\sqrt{2ar - 2at_0} \le 1} x e^{-x^2} dx dr$$

almost surely for any $t > t_0$. By Theorem 1.4 with probability one $\pi_{t_0}(x) =$ $x\theta(x), x \in [0, 1]$, where θ is a bounded function of x tending to zero as $x \downarrow 0$. Hence

$$E(A_t - A_{t_0} \mid \mathcal{F}_{t_0}^{b.})$$

$$\leq \frac{2a}{\sqrt{\pi}} \int_{t_0}^t \int_0^\infty \theta(\sqrt{2arx - 2at_0x} \wedge 1) x^2 e^{-x^2} dx dr.$$

By the dominated convergence theorem (a.s.)

$$\lim_{r \downarrow t_0} \int_0^\infty \theta(\sqrt{2arx - 2at_0x} \wedge 1) x^2 e^{-x^2} \, dx = 0$$

implying that (a.s.)

$$\lim_{n \to \infty} \frac{1}{t_n - t_0} \int_{t_0}^{t_n} \int_0^{\infty} \theta(\sqrt{2arx - 2at_0x} \wedge 1) x^2 e^{-x^2} \, dx \, dr,$$
$$\overline{\lim_{n \to \infty} \frac{1}{t_n - t_0}} E(A_{t_n} - A_{t_0} \mid \mathcal{F}_{t_0}^{b.}) = 0,$$

which yields (2.1) by Fatou's lemma.

Thus, for any $t \geq 0$, for almost all ω

$$\lim_{t \to \infty} n(A_{t+1/n} - A_t) = 0.$$
(2.5)

By Fubini's theorem, for almost any ω , equation (2.5) holds for almost all t. It follows that the derivative of A_t is zero for almost all t and the theorem is proved.

3. Proof of Theorems 1.4 and 1.5

Proof of Theorem 1.4. On the space C of continuous functions on $[0, \infty)$ with Wiener measure W introduce the coordinate process $x_t(x_t) := x_t$, which is a Wiener process. For $t \ge 0$, $x \in \mathbb{R}$, and $x_t, y_t \in C$ such that $y_0 = 0$ define

$$\tau(t, x, x_{\cdot}, y_{\cdot}) = \inf\{s \ge 0 : x + \sigma_1 x_s \notin (\sigma y_{t-s}, \sigma y_{t-s} + 1)\},\$$

where $y_r := y_0$ for $r \leq 0$. Then the function

$$u_t(y_{\cdot}, x) := \int_C \pi_0(x_t) I_{\tau(t, x, x_{\cdot}, y_{\cdot}) \ge t} W(dx_{\cdot}).$$

is the probabilistic solution of the heat equation

$$\partial_t u_t = (1/2)\sigma_1^2 D^2 u_t$$

in $\Gamma(y)$ with boundary conditions

$$u_t(y_t) = u_t(y_t + 1) = 0, \quad t > 0,$$

 $u_0(x) = \pi_0(x), \quad 0 \le x \le 1.$

Due to interior estimates of derivatives of solutions to the heat equation, $u_t(y, x)$ is infinitely differentiable in $\Gamma(y)$. Its continuity up to $\{0\} \times [0, 1]$ easily follows from the fact that π_0 is a continuous function on [0, 1] vanishing at 0 and 1. The continuity of its derivatives up to $\{0\} \times (0, 1)$ follows from the fact that π_0 is infinitely differentiable. Next, as in the proof of Theorem 4.1 of [2] one shows that for constants ν (different in (3.1) and (3.2)) as in Theorem 1.5 we have that, for almost any trajectory of b.,

$$\sup_{x \in (0,1)} \sup_{t \in [0,T]} \frac{u_t(b,x)}{(b_t + x)^{\nu} (b_t + 1 - x)^{\nu}} < \infty,$$
(3.1)

$$E \sup_{x \in (0,1)} \sup_{t \in [0,T]} \frac{u_t(b,x)}{(b_t + x)^{\nu} (b_t + 1 - x)^{\nu}} < \infty.$$
(3.2)

In particular, with probability one $u_t(b, x)$ is continuous at the lateral boundary of $\Gamma(b)$.

Next we deal with (1.8) We organize the proof in the following way. For $t \ge 0, x \in \mathbb{R}$, and $x, y \in C$ such that $y_0 = 0$ define

$$\gamma(t, x, x, y) = \inf\{s \ge 0 : x + \sigma_1 x_s \le \sigma y_{t-s}\},\$$

where $y_r := y_0$ for $r \leq 0$. Also let

$$v_t(y_{\cdot}, x) := \int_C \pi_0 I_G(x_t) I_{\tau(t, x, x_{\cdot}, y_{\cdot}) \ge t} W(dx_{\cdot})$$

Lemma 3.1. Let B_t be a one-dimensional Wiener process and $\gamma \in (0, 1)$. Then with probability one there exists a sequence of integers $0 \le m_1 < m_2 < \ldots$ such that $B_{t_k} \ge \sqrt{t_k}$ for all k, where $t_k = \gamma^{m_k}$. Moreover, $m_k \le \beta k$ for all sufficiently large k, where β is any fixed number such that $\alpha\beta > 1$, $\alpha = P(B_1 \ge 1)$.

Proof. The sequence $I_{B_{\gamma^m} \geq \gamma^{m/2}}, m=0,1,...,$ is stationary, so that the limit

$$\lim_{n\to\infty}\frac{1}{m}\sum_{k=1}^m I_{B_{\gamma^k}\geq\gamma^{k/2}}$$

exists (a.s.). By the 0-1 law this limit is a constant and equals α (a.s.). Set

 $m_1 = \inf\{k \ge 1 : B_{\gamma^k} \ge \gamma^{k/2}\}, \quad m_{n+1} = \inf\{k > m_n : B_{\gamma^k} \ge \gamma^{k/2}\}.$

Then the number of $k \in \{1, 2, ...\}$ such that $m_k \leq m$ divided by m tends to α . It follows that the number of integers $i \in \{1, 2, ..., \beta k\}$ such that $m_i \leq \beta k$ for all large k is greater than $\beta' k \alpha$, where β' is any number such that $\beta' < \beta$ and $\beta' \alpha > 1$. On the other hand there are always exactly kvalues of $i \in \{1, 2, ..., m_k\}$ such that $B_{\gamma i} \geq \gamma^{i/2}$. Since $\beta' \alpha > 1$, it follows that for any sufficiently large k the inequality $m_k \geq \beta k$ is impossible. The lemma is proved.

Lemma 3.2. If σ_1/σ is sufficiently small, then for each $T \in (0, \infty)$

$$\lim_{x \downarrow 0} \frac{v_T(x, b_{\cdot})}{x - \sigma b_T} = 0$$
(3.3)

almost surely.

Proof. Fix a $T \in (0, \infty)$, set $\gamma = 1/2$, and take integers $0 \le m_1 < m_2 < ...$ such that $b_{T-t_k} - b_T \ge \sqrt{t_k}$ for all k, where $t_k = \gamma^{m_k}$ and $m_k \le \beta k$ for all large k. By Lemma 3.1 such a sequence exists with probability one. Then notice that the inequality $\tau(T, x, x, b) \ge T$ implies that

$$x + \sigma_1 x_{t_k} \ge \sigma b_{T-t_k} \ge \sigma b_T + \sigma \sqrt{t_k}$$

for all k such that $t_k \leq T$. Denote by k_0 the smallest k such that $t_k \leq T$. We also take into account that π_0 is a bounded function and conclude that for any integer $n \geq k_0$

$$v_T(x, b_{\cdot}) \le NW(z + \sigma_1 x_{t_k} - \sigma \sqrt{t_k} \ge 0, k = k_0, ..., n),$$

where $N = \sup \pi_0$ and $z = x - \sigma b_T$. It follows that to prove (3.3) it suffices to show that there exists an integer-valued function $n = n(x) \ge k_0$ such that

$$\lim_{x \downarrow 0} \frac{1}{x} P(x + w_{t_k} - K\sqrt{t_k} \ge 0, k = k_0, ..., n(x)) = 0,$$
(3.4)

where $K = \sigma / \sigma_1 = \varepsilon^{-1}$.

Observe that by Girsanov's theorem for any $n \ge k_0 + 1$

$$P(x + w_{t_k} - K\sqrt{t_k} \ge 0, k = k_0, k_0 + 1, ..., n)$$

= $EI_{\Gamma_n(x)} \exp\left(-\int_{t_n}^{t_{k_0}} f'(t) \, dw_t - (1/2) \int_{t_n}^{t_{k_0}} [f'(t)]^2 \, dt\right) =: I_n(x),$
where $f(t) = K\sqrt{t}$ for $t \ge t_n, f(t) = K\sqrt{t_n}$ for $t \in [0, t_n]$, and

$$\Gamma_n(x) = \{ \omega : x + w_{t_k} - K\sqrt{t_n} \ge 0, k = k_0, ..., n \}$$

Next, note that for bounded nonrandom functions g

$$E\{\exp\int_{s}^{t}g(r)\,dw_{r}\mid w_{s},w_{t}\}$$
$$=\exp\left(\frac{w_{t}-w_{s}}{t-s}\int_{s}^{t}g(u)\,du+(1/2)\int_{s}^{t}\left[g(r)-\frac{1}{t-s}\int_{s}^{t}g(u)\,du\right]^{2}\,dr\right)$$
cause

bee

$$\int_{s}^{t} \left[g(r) - \frac{1}{t-s} \int_{s}^{t} g(u) \, du \right] \, dw_{r}$$

is independent of w_s, w_t . Then we use the fact that

$$\int_{s}^{t} \left[g(r) - \frac{1}{t-s} \int_{s}^{t} g(u) \, du \right]^{2} dr = \int_{s}^{t} g^{2}(r) \, dr - \frac{1}{t-s} \left(\int_{s}^{t} g(u) \, du \right)^{2}$$

Hence, by applying this to g = -f' we find that

$$I_n(x) = EI_{\Gamma_n(x)} \exp\left(D_n - (1/2)C_n\right)$$

with

$$D_n := -K \sum_{k=k_0+1}^n s_k (w_{t_{k-1}} - w_{t_k}), \quad s_k = (\sqrt{t_{k-1}} + \sqrt{t_k})^{-1},$$
$$C_n := \sum_{k=k_0+1}^n \frac{1}{t_{k-1} - t_k} \left(\int_{t_k}^{t_{k-1}} f'(u) \, du\right)^2$$
$$= K^2 \sum_{k=k_0+1}^n \frac{\sqrt{t_{k-1}} - \sqrt{t_k}}{\sqrt{t_{k-1}} + \sqrt{t_k}} \ge K^2 \frac{1 - \sqrt{\gamma}}{1 + \sqrt{\gamma}} (n - k_0) =: 2K\kappa(n - k_0),$$

where the inequality follows from the fact that $\gamma t_{k-1} \ge t_k$ and κ is defined by the last equality.

Now we use summation by parts to see that

$$D_n = -Kw_{t_{k_0}}s_{k_0+1} + Kw_{t_n}s_n - K\sum_{k=k_0+1}^{n-1}w_{t_k}(s_{k+1}-s_k).$$

On the event $\Gamma_n(x)$ this quantity is smaller than

$$Kw_{t_n}s_n + K(x - K\sqrt{t_n}) \left(s_{k_0+1} + \sum_{k=k_0+1}^{n-1} (s_{k+1} - s_k) \right)$$

= $Kw_{t_n}s_n + K(x - K\sqrt{t_n})s_n.$

It follows that

$$I_n(x) \leq E I_{\Gamma_n(x)} \exp\left(K w_{t_n} s_n + K x s_n - K^2 \kappa (n - k_0)\right),$$

$$\leq E \exp\left(K w_{t_n} s_n + K x s_n - K^2 \kappa (n - k_0)\right)$$

$$= \exp\left((K^2/2) t_n s_n^2 + K x s_n - K^2 \kappa (n - k_0)\right).$$

Now it is time to choose n = n(x). We take n = n(x) so that $x^2 \in [t_n, t_{n-1}]$. Then $t_n s_n \leq \sqrt{t_n} \leq x$ and $x s_n \leq x/\sqrt{t_{n-1}} \leq 1$. Also $t_n s_n^2 \leq 1$ and since $t_n \geq \gamma^{\beta n}$, we have $x \geq \gamma^{\beta n/2}$, which implies that

$$\lim_{x \downarrow 0} x^{-1} I_{n(x)}(x)$$

$$\leq \lim_{n \to \infty} \exp\left(\frac{K^2}{2} + K - \frac{K^2 \kappa (n - k_0)}{(n - k_0)} - \frac{(1/2)\beta n \ln \gamma}{(1/2)\beta n \ln \gamma}\right) = 0$$

if $\beta |\ln \gamma| < 2K^2 \kappa$, which is true if σ_1/σ is small enough. This proves the lemma.

Corollary 3.3. If σ_1/σ is sufficiently small, then (1.8) holds (a.s.) for any fixed $t \ge 0$.

Indeed, the equality of the extreme terms in (1.8) follows from Lemma 3.2 since $u_t \leq v_t$. The remaining equality is proved similarly by replacing x with 1 - x.

It only remains to prove the last assertion of the theorem.

Observe that $\tau(t, x, x, y)$ is a lower semicontinuous function of its arguments. Therefore by Fubini's theorem $u_t(y, x)$ is a Borel function of (y, t, x). Furthermore, $u_t(y, x)$ will not change if we change y_r for r > t. Hence, $u_t(y, x)$ is \mathcal{N}_t -measurable, where $\mathcal{N}_t = \sigma(y_r : r \leq t, y \in C)$. Therefore,

$$v_t(x) := u_t(b_{\cdot}, x - \sigma b_t)$$

is \mathcal{F}_t -measurable for each $(t, x) \in \overline{G}_{\infty}$. After that in the same way the usual Itô's formula is proved on the basis of Taylor's formula and the fact that $u_t(y, x)$ is infinitely differentiable we obtain that for any $x \in G$ almost surely for all $t \geq 0$

$$v_t(x) = \pi_0(x) + (a/2) \int_0^t D^2 v_s(x) \, ds - \sigma \int_0^t D v_s(x) \, db_s.$$

The above properties of u_t and Theorem 4.1 now imply that (perhaps after modifying v_t on a set of probability zero) v_t satisfies (1.2) with zero boundary condition and initial condition π_0 in the sense explained below that formula and is such that

$$\int_0^T \|v_t\|_{W_2^1(G)}^2 \, dt < \infty$$

for each $T \ge 0$. Uniqueness of solutions of (1.2) in this class of functions is a classical result, and this proves the remaining assertions of the theorem.

Proof of Theorem 1.5. After (1.9) has been proved and the modification $u_t(x - \sigma b_t)$ of $\pi_t(x)$ has been chosen the assertions related to (1.11) and (1.12) follow directly from (3.1) and (3.2). This proves the theorem.

4. Appendix

Let w_t^1, w_t^2, \ldots be independent one-dimensional Wiener processes with respect to \mathcal{F}_t , and let \mathcal{P} denote the predictable σ -field related to $\{\mathcal{F}_t\}$. Assume that on $\Omega \times (0, \infty) \times \mathbb{R}^d$ we are given $a_t(x) = (a_t^{ij}(x))$ which is a $d \times d$ -symmetric matrix valued function, $b_t(x)$ which is an \mathbb{R}^d -valued function, real-valued $c_t(x)$, and $\sigma_t^{i}(x) = (\sigma_t^{i,1}(x), \sigma_t^{i,2}(x), \ldots), i = 1, \ldots, d$, and $\nu_t(x) = (\nu_t^1(x), \nu_t^2(x), \ldots)$, which are ℓ_2 -valued functions.

Assumption 4.1. (i) The functions a, b, c, σ, ν are bounded and measurable as functions of (ω, t, x) and are predictable as functions of (ω, t) for each x.

(ii) The functions $a_t(x)$ are Lipschitz continuous in x with constant K (independent of ω, t).

(ii) For any $\lambda, x \in \mathbb{R}^d, t \geq 0$, and ω

$$(2a_t^{ij} - \alpha_t^{ij})(x)\lambda^i \lambda^j \ge \delta_0 |\lambda|^2, \tag{4.1}$$

where δ_0 is a constant, $\delta_0 > 0$, $\alpha^{ij} := \sigma^{ik} \sigma^{jk}$.

Let $G \subset \mathbb{R}^d$ be an open set, $T \in [0, \infty)$, and let $u_t(x)$ be a real-valued function given on $\Omega \times [0, T] \times \overline{G}$. Also suppose that on $\Omega \times (0, T) \times G$ we are given functions $f_t(x)$ and $g_t(x) = (g_t^1(x), g_t^2(x), ...)$ with values in \mathbb{R} and ℓ_2 , respectively.

Define $D_i = \partial/\partial x^i$, $D_{ij} = D_i D_j$, Du the gradient of u, $D^2 u$ its Hessian.

Assumption 4.2. (i) For any ω the function u is continuous in $[0, T] \times \overline{G}$ and vanishes on $[0, T] \times \partial G$ and, moreover (if G is unbounded), for any $\delta > 0$ and ω there exists a compact set $\Gamma \subset G$ such that $|u_t(x)| \leq \delta$ for all $t \in [0, T]$ and $x \in G \setminus \Gamma$. For any x the function $u_t(x)$ is \mathcal{F}_t -adapted. For any ω (if Gis unbounded)

$$\int_G |u_0|^2 dx + \int_0^T \int_G |u_t|^2 dx dt < \infty.$$

(ii) For any ω , for almost any $t \in [0,T]$, the second-order derivatives $D^2 u_t(x)$ are continuous with respect to x and for any compact set $\Gamma \subset G$

$$\int_0^T \int_{\Gamma} |D^2 u_t| \, dx dt < \infty.$$

(iii) For each ω and compact set $\Gamma \subset G$

$$\int_0^T \int_{\Gamma} |Du_t|^2 \, dx \, dt < \infty.$$

(iii) The functions $f_t(x)$ and $g_t(x)$ are $\mathcal{P} \otimes \mathcal{B}(G)$ -measurable as functions of (ω, t, x) , where $\mathcal{B}(G)$ is the Borel σ -field on G. For each $x \in G$ and ω we have

$$\int_0^1 \left(|D^2 u_s(x)| + |D u_s(x)|^2 + |f_s(x)|^2 + |g_s(x)|_{\ell_2}^2 \right) ds < \infty.$$

(iv) For each ω

$$\int_G \int_0^T (|f_s(x)|^2 + |g_s(x)|^2_{\ell_2}) \, ds \, dx < \infty,$$

and for each $x \in G$ with probability one we have for all $t \in [0, T]$ that

$$u_t(x) = u_0(x) + \int_0^t \left[\sigma_s^{ik}(x)D_iu_s(x) + \nu_s^k(x)u_s(x) + g_s^k(x)\right] dw_s^k + \int_0^t \left[a_s^{ij}(x)D_{ij}u_s(x) + b_s^i(x)D_iu_s(x) + c_s(x)u_s(x) + f_s(x)\right] ds.$$

Theorem 4.1. Under the above assumptions

$$\int_0^T \int_G |Du_t(x)|^2 \, dx \, dt < \infty$$

(a.s.) and for any $\phi \in C_0^{\infty}(G)$ with probability one

$$(u_t, \phi) = (u_0, \phi) + \int_0^t (\phi, \sigma_s^{ik} D_i u_s + \nu_s^k u_s + g_s^k) \, dw_s^k + \int_0^t \left[(\phi, (b_s^i - D_j a^{ij}) D_i u_s + c_s u_s + f_s) - (D_j \phi, a_s^{ij} D_i u_s) \right] ds, t \in [0, T]$$

for all $t \in [0,T]$.

Proof. By the stochastic Fubini theorem (see, for instance, Lemma 2.7 of [4]), for any $\phi \in C_0^{\infty}(G)$ with probability one for all $t \ge 0$

$$(u_t^{\delta}, \phi) = (u_0^{\delta}, \phi) + \int_0^t (\phi, \sigma_s^{ik} D_i u_s + \nu_s^k u_s + g_s^k) \, dw_s^k + \int_0^t \left[(\phi, (b_s^i - D_j a^{ij}) D_i u_s + c_s u_s + f_s) - (D_j \phi, a_s^{ij} D_i u_s) \right] ds,$$

where $u^{\delta} = u - \delta$.

For $\varepsilon, \delta > 0$ define

$$K_{\varepsilon} = \{ x \in G : \operatorname{dist} (x, \partial D) \le \varepsilon \quad \text{or} \quad |x| \ge \varepsilon^{-1} \},\$$

$$\tau_{\varepsilon, \delta} = T \wedge \inf\{ t \ge 0 : \sup_{G \setminus K_{\varepsilon}} |u_t| \ge \delta \}.$$

Observe that as $\varepsilon \downarrow 0$ we have

$$\sup_{G\setminus K_{\varepsilon}} |u_t| \downarrow 0$$

uniformly with respect to $t \in [0,T]$ by assumption. Therefore $\tau_{\varepsilon,\delta} \to T$ as $\varepsilon \downarrow 0$ for any $\delta > 0$. Also notice that

$$|u_t(x)| \le \delta$$

in $G \setminus K_{\varepsilon}$ if $0 \le t < \tau_{\varepsilon,\delta}$.

By Lemma 2.5 of [3] for any $\phi \in C_0^{\infty}(G)$ (a.s.) for all $t \in [0,T]$

$$\|(\phi u_{t\wedge\tau_{\varepsilon,\delta}}^{\delta})^{+}\|_{L_{2}}^{2} = \|(\phi u_{0}^{\delta})^{+}\|_{L_{2}}^{2} + \int_{0}^{t} h_{s} \, ds + m_{t}, \tag{4.2}$$

where

$$h_s := 2((\phi u_s^{\delta})^+, \phi\{(b_s^i - D_j a^{ij})D_i u_s + c_t u_s + f_s\}$$
$$-a_t^{ij}(D_i u_s)D_j\phi)I_{s<\tau_{\varepsilon,\delta}} - 2(I_{\phi u_s^{\delta}>0}D_i(\phi u_s^{\delta}), \phi a_s^{ij}D_j u_s)I_{s<\tau_{\varepsilon,\delta}}$$
$$+\sum_k \left\|\phi\left[\sigma_s^{ik}D_i u_s + \nu_s^k u_s + g_s^k\right)\right]I_{\phi u_s^{\delta}>0}\right\|_{L_2}^2 I_{s<\tau_{\varepsilon,\delta}},$$
$$m_t := 2\int_0^t I_{s<\tau_{\varepsilon,\delta}}(\phi(\phi u^{\delta})_s^+, \sigma_s^{ik}D_i u_s + \nu_s^k u_s + g_s^k) dw_s^k.$$

We take $\phi \geq 0$ such that $\phi = 1$ on K_{ε} . Then for $0 \leq s < \tau_{\varepsilon,\delta}$

$$(\phi u_s^\delta)^+ = u_s^\delta I_{u_s > \delta}.$$

Indeed, if $u_s^{\delta}(x) \leq 0$, then both sides vanish. However, if $u_s^{\delta}(x) > 0$ and $0 \leq s < \tau_{\varepsilon,\delta}$, then $x \in K_{\varepsilon}$ and $\phi(x) = 1$. By also taking into account that

$$I_{u_s > \delta, s < \tau_{\varepsilon,\delta}} D_j \phi = 0$$

we transform (4.2) for such a ϕ into

$$\|(u_{t\wedge\tau_{\varepsilon,\delta}}^{\delta})^{+}\|_{L_{2}}^{2} = \|(u_{0}^{\delta})^{+}\|_{L_{2}}^{2} + \int_{0}^{t} h_{s} \, ds + m_{t}, \tag{4.3}$$

where

$$\begin{split} h_s &:= 2 \left(u_s^{\delta} I_{u_s > \delta}, (b_s^i - D_j a_{ij}^{ij}) I_{u_s > \delta} D_i u_s + c_t u_s + f_s \right) I_{s < \tau_{\varepsilon, \delta}} \\ &- 2 (I_{u_s > \delta} D_i u_s, a_t^{ij} D_j u_s) I_{s < \tau_{\varepsilon, \delta}} \\ &+ \sum_k \left\| \left[\sigma_s^{ik} D_i u_s^{\delta} + \nu_s^k u_s + g_s^k \right] \right] I_{u_s > \delta} \right\|_{L_2}^2 I_{s < \tau_{\varepsilon, \delta}}, \\ m_t &:= 2 \int_0^t I_{s < \tau_{\varepsilon, \delta}} ((u^{\delta})_s^+, \sigma_s^{ik} D_i u_s + \nu_s^k u_s^{\delta} + g_s^{k\delta}) \, dw_s^k. \end{split}$$

To proceed further we recall again the fact that the two conditions: $s < \tau_{\varepsilon,\delta}$ and $u_s(x) > \delta$, imply that $x \in K_{\varepsilon}$ so that

$$\int_0^{\tau_{\varepsilon,\delta}} \int_G |Du_t|^2 I_{u_t > \delta} \, dx dt \le \int_0^T \int_{K_\varepsilon} |Du_t|^2 \, dx dt < \infty.$$

After that we use (4.1) and the inequalities like $ab \leq \varepsilon a^2 + \varepsilon^{-1}b^2$. We also recall that $b_s^i - D_j a_s^{ij}, c_s$, and $|\nu_s|_{\ell_2}$ are bounded and then in an absolutely

standard way derive from (4.3) that there exists a constant $N \in (0, \infty)$, independent of $\omega, t, \varepsilon, \delta$, such that for any $t \in [0, T]$ and ω

$$N^{-1} \int_0^{t \wedge \tau_{\varepsilon,\delta}} \|I_{u_s > \delta} Du_s\|_{L_2}^2 \, ds \le N \|u_0\|_{L_2}^2 + N \int_0^{t \wedge \tau_{\varepsilon,\delta}} \|u_s\|_{L_2}^2 \, ds + m_t.$$

In particular, for any stopping time $\tau \leq T$

$$N^{-1} \int_0^{\tau \wedge \tau_{\varepsilon,\delta}} \|I_{u_s > \delta} Du_s\|_{L_2}^2 \, ds \le N \|u_0\|_{L_2}^2 I_{\tau > 0} + N \int_0^{\tau \wedge \tau_{\varepsilon,\delta}} \|u_s\|_{L_2}^2 \, ds + m_\tau.$$

If τ is a localizing time for the local martingale m_t starting at zero, then $Em_{\tau} = 0$ and

$$N^{-1}E \int_0^{\tau \wedge \tau_{\varepsilon,\delta}} \|I_{u_s > \delta} Du_s\|_{L_2}^2 \, ds \le NE \|u_0\|_{L_2}^2 I_{\tau > 0} + NE \int_0^{\tau \wedge \tau_{\varepsilon,\delta}} \|u_s\|_{L_2}^2 \, ds.$$

The last inequality, actually, holds for any stopping time $\tau \leq T$, which is easily proved by approximation.

Now we first let $\varepsilon \downarrow 0$ and then $\delta \downarrow 0$. Then by the monotone convergence theorem we obtain

$$N^{-1}E\int_0^\tau \|I_{u_s>0}Du_s\|_{L_2}^2 \, ds \le NE\|u_0\|_{L_2}^2 I_{\tau>0} + NE\int_0^\tau \|u_s\|_{L_2}^2 \, ds.$$

Similarly,

$$N^{-1}E\int_0^\tau \|I_{u_s<0}Du_s\|_{L_2}^2\,ds \le NE\|u_0\|_{L_2}^2I_{\tau>0} + NE\int_0^\tau \|u_s\|_{L_2}^2\,ds,$$

and since one knows that $I_{u=0}Du = 0$ (a.e.) for any function u of class W_2^1 we finally conclude that

$$N^{-1}E \int_0^\tau \|Du_s\|_{L_2}^2 \, ds \le NE \|u_0\|_{L_2}^2 I_{\tau>0} + NE \int_0^\tau \|u_s\|_{L_2}^2 \, ds. \tag{4.4}$$

For n > 0 and $\tau = \tau^n$, where

$$\tau_n = T \wedge \inf\{t \ge 0 : \|u_0\|_{L_2} + \int_0^t \|u_s\|_{L_2}^2 \, ds \ge n\},$$

the right-hand side of (4.4) is finite. Hence

$$\int_0^{\tau^n} \|Du_s\|_{L_2}^2 \, ds < \infty$$

(a.s.). To prove the first assertion of the theorem, it only remains to observe that, by assumption, for any ω , we have $\tau_n = T$ for all sufficiently large n. The second assertion follows from the stochastic Fubini theorem. The theorem is proved.

References

- N.V. Krylov, Brownian trajectory is a regular lateral boundary for the heat equation, SIAM J. Math. Anal., Vol. 34 (2003), No. 5, 1167-1182.
- [2] N.V. Krylov, One more square root law for Brownian motion and its application to SPDEs, Probab. Theory Relat. Fields, Vol. 127 (2003), 496-512.
- [3] N.V. Krylov, Maximum principle for SPDEs and its applications, pp. 311-338 in "Stochastic Differential Equations: Theory and Applications, A Volume in Honor of Professor Boris L. Rozovskii", P.H. Baxendale, S.V. Lototsky eds., Interdisciplinary Mathematical Sciences, Vol. 2, World Scientific, 2007.
- [4] N.V. Krylov, On the Itô-Wentzell formula for distribution-valued processes and related topics, Probability Theory and Related Fields, Vol. 150 (2011), No. 1-2, 295– 319.
- [5] N.V. Krylov and B.L. Rozovskii, *Stochastic evolution equations*, pp. 71–146 in "Itogy nauki i tekhniki", Vol. 14, VINITI, Moscow, 1979 in Russian; English translation in J. Soviet Math., Vol. 16, No. 4 (1981), 1233–1277.
- [6] N.V. Krylov and Teng Wang, Filtering partially observable diffusions up to the exit time from a domain, Stoch. Proc. Appl., Vol. 121 (2011), No.8, 1785–1815.
- [7] E. Pardoux, "Equations aux dérivées partielles stochastiques non linéaires monotones. Etude de solutions fortes de type Ito", (1975). Thèse Doct. Sci. Math. Univ. Paris Sud., http://www.cmi.univ-mrs.fr/~pardoux/Pardoux_these.pdf
- [8] B.L. Rozovskii, "Stochastic evolution systems", Kluwer, Dordrecht, 1990.
- *E-mail address*: krylov@math.umn.edu
- 127 VINCENT HALL, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN, 55455