

A NOTE ON ORTHOGONAL LIE ALGEBRAS IN DIMENSION 4 VIEWED AS CURRENT LIE ALGEBRAS

MARTIN CHAKTOURA AND FERNANDO SZECHTMAN

ABSTRACT. Orthogonal Lie algebras in dimension 4 are identified as current Lie algebras, thus producing a natural decomposition for them over any field.

Let $f : V \times V \rightarrow F$ be a non-degenerate symmetric bilinear form defined on a vector space V of dimension 4 over a field F . If $\text{char}(F) = 2$ we further assume that f is not alternating. Let D be the discriminant of f relative to a basis of V . Let $L(f)$ be the subalgebra of $\mathfrak{gl}(V)$ associated to f , formed by all $x \in \mathfrak{gl}(V)$ that are skew-adjoint relative to f . Let $M = [L(f), L(f)]$, which coincides with $L(f)$ if and only if $\text{char}(F) \neq 2$. In any case, M is a 6-dimensional orthogonal Lie algebra.

According to [B], Chapter 1, §6, Exercise 26, if $\text{char}(F) \neq 2$ then M is either the direct sum of two 3-dimensional simple ideals or simple, depending on whether D is a square in F or not. On the other hand, if $\text{char}(F) = 2$ then either $M = N \ltimes R$ or M is simple, depending, again, on whether D is a square in F or not; here N is a simple 3-dimensional simple subalgebra and R is the solvable radical of M .

We may view these orthogonal Lie algebras as current Lie algebras, and in this way explain all cases described above in a uniform manner.

Theorem. We have $M \cong [L(f|_W), L(f|_W)] \otimes F[X]/(X^2 - D)$, where W is an arbitrary 3-dimensional subspace of V such that $f|_W$ is non-degenerate.

Proof. It follows from [K], Theorems 4 and 20, that V admits an orthogonal basis $B = \{v_1, v_2, v_3, v_4\}$. Thus, the Gram matrix of f relative to B is diagonal with non-zero entries a, b, c, d and $D = abcd$. Let

$$\begin{aligned} f_1 &= be_{12} - ae_{21}, & f_2 &= ce_{23} - be_{32}, & f_3 &= ce_{13} - ae_{31}, \\ h_1 &= ab(de_{34} - ce_{43}), & h_2 &= bc(de_{14} - ae_{41}), & h_3 &= ac(be_{42} - de_{24}). \end{aligned}$$

Then $f_1, f_2, f_3, h_1, h_2, h_3$ is a basis of M , with multiplication table

$$\begin{aligned} [f_1, f_2] &= bf_3, & [f_2, f_3] &= cf_1, & [f_3, f_1] &= af_2, \\ [f_1, h_2] &= bh_3, & [f_2, h_3] &= ch_1, & [f_3, h_1] &= ah_2, \\ [f_2, h_1] &= -bh_3, & [f_3, h_2] &= -ch_1, & [f_1, h_3] &= -ah_2, \end{aligned}$$

and

$$[h_1, h_2] = Dbf_3, \quad [h_2, h_3] = Dcf_1, \quad [h_3, h_1] = Daf_2.$$

Thus M has the same multiplication table as $[L(f|_W), L(f|_W)] \otimes F[X]/(X^2 - D)$, where W is the span by v_1, v_2, v_3 . By [K], Theorems 4 and 20, W can be replaced by any 3-dimensional subspace of V where the restriction of f is non-degenerate. \square

2000 *Mathematics Subject Classification.* 17B05.

Key words and phrases. Classical Lie algebras, current Lie algebras.

The second author was supported in part by an NSERC discovery grant.

It is obvious from the Theorem that M decomposes exactly as prescribed above if D is a square. Suppose D is not a square. Then $K = F[X]/(X^2 - D)$ is a quadratic field extension of F . Since $[L(f|_W), L(f|_W)]$ is a perfect 3-dimensional Lie algebra over F , it follows that $[L(f|_W), L(f|_W)] \otimes K$ is a simple Lie algebra over K , and hence over F , as seen below. Thus, by the Theorem, M is a simple Lie algebra over F .

Lemma. (cf. [LP], Lemma 2.7) Let L be a simple Lie algebra over a field K . Suppose that F is a subfield of K . Then L is simple over F .

Proof. Let I be a non-zero ideal of L over F . The K -span of I is a non-zero ideal J of L over K , so $J = L = [L, L] = [L, J] = [L, I] \subseteq I$. \square

A simple Lie algebra need not remain simple, or even semisimple, upon field extension. For instance, let L be an absolutely simple Lie algebra over an imperfect field F of characteristic p , and let $s \in F \setminus F^p$ and $K = F[X]/(X^p - s)$, which has an element t satisfying $t^p = s$. Then $P = L \otimes K$ is simple over K and hence over F , but $P \otimes K \cong L \otimes K[X]/(X^p - s) \cong L \otimes K[X]/(X - t)^p$ is not semisimple over K .

Acknowledgment. We are grateful to A. Pianzola for pointing out reference [LP], which replaced a more convoluted argument.

REFERENCES

- [B] N. Bourbaki, *Lie groups and Lie algebras, Chapters 1-3*, Springer-Verlag, Berlin, 1989.
- [K] I. Kaplansky, *Linear algebra and geometry. A second course.*, Allyn and Bacon, Boston, 1969.
- [LP] M. Lau and A. Pianzola, *Maximal Ideals and Representations of Twisted Forms of Algebras*, to appear in Algebra and Number Theory.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, CANADA
E-mail address: martin_chaktoura@yahoo.com.ar

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, CANADA
E-mail address: fernando.szechtman@gmail.com