

RATIONAL CURVES ON QUOTIENTS OF ABELIAN VARIETIES BY FINITE GROUPS

BO-HAE IM AND MICHAEL LARSEN

ABSTRACT. In [3], it is proved that the quotient of an abelian variety A by a finite order automorphism g is uniruled if and only if some power of g satisfies a numerical condition $0 < \text{age}(g^k) < 1$. In this paper, we show that $\text{age}(g^k) = 1$ is enough to guarantee that $A/\langle g \rangle$ has at least one rational curve.

1. INTRODUCTION

Let G be a finite group of automorphisms of an abelian variety A/\mathbb{C} . It is a classical result [4, II. §1] that A itself cannot contain a rational curve. For $|G| > 1$, there may or may not be rational curves on A/G . For general abelian varieties, $\text{Aut}(A) = \pm 1$, and Pirola proved [7] that for A sufficiently general and of dimension at least three, $A/\pm 1$ has no rational curves. At the other extreme, regarding $A = E^n$ as the set of $n + 1$ -tuples of points on the elliptic curve E which sum to 0, A/S_{n+1} can be interpreted as the set of effective divisors linearly equivalent to $(n + 1)[0]$ and, as such, is just \mathbb{P}^n . More generally, Looijenga has shown [6] that the quotient of E^n by the Weyl group of a root system of rank n is a weighted projective space.

Rational curves on A/G over a field K are potentially a source of rational points over G -extensions of K . For instance, the method [5] for finding pairs $a, b \in \mathbb{Q}^\times$ such that the quadratic twists E_a , E_b , and E_{ab} all have positive rank amounts to finding a rational curve on $E^3/(\mathbb{Z}/2\mathbb{Z})^2$. Likewise, the theorem of Looijenga cited above gives for each elliptic curve E over a number field K and for each Weyl group W , a source of W -extensions L_i of K such that the representation of W on each $E(L_i) \otimes \mathbb{Q}$ contains the reflection representation. On the other hand, the result of Pirola cited above dims the hope of using geometric methods to show that every abelian variety over a number field K gains rank over infinitely many quadratic extensions of K . Thus, it is desirable from the viewpoint of arithmetic to understand when A/G can be expected to have a rational curve over a given field K ,

Date: September 17, 2018.

2000 Mathematics Subject Classification. 14K05.

Bo-Hae Im was partially supported by the National Research Foundation of Korea Grant funded by the Korean Government(MEST) (NRF-2011-0015557). Michael Larsen was partially supported by NSF grant DMS-1101424.

and to begin with, one would like to know when A/G has a rational curve over \mathbb{C} .

Any automorphism g of an abelian variety A defines an invertible linear transformation (also denoted g) on $\text{Lie}(A)$. If g is of finite order, there exists a unique sequence of rationals $0 \leq x_1 \leq x_2 \leq \dots \leq x_n < 1$ such that the eigenvalues of g are $e(x_1), \dots, e(x_n)$, where $e(x) := e^{2\pi i x}$. We say g is of *type* (x_1, \dots, x_n) . Following Kollár and Larsen [3], we write $\text{age}(g) = x_1 + \dots + x_n$. For instance, $\text{age}(g) = 1/2$ for every reflection g . The main result of [3] asserts that A/G is uniruled if and only if $0 < \text{age}(g) < 1$ for some $g \in G$. In this paper, we prove that to find a single rational curve in A/G , it suffices that $\text{age}(g) \leq 1$.

Since we need only consider the case $\text{age}(g) = 1$, we first classify all types of weight 1. This requires a combinatorial analysis, which we carried out using a computer algebra system to minimize the risk of an oversight. There are thirty-five cases (see Table 2 below), and our strategy for finding rational curves depends on case analysis. Abelian surfaces play a special role, since here we can use known results on K3 surfaces. The other key idea is to find a non-singular projective curve X on which G acts with quotient \mathbb{P}^1 and a G -equivariant map from X to A , or, equivalently, a G -homomorphism from the Jacobian variety of X to A .

We would like to thank Yuri Tschinkel and Alessio Corti for helpful comments on earlier versions of this paper.

2. CLASSIFYING TYPES

If $A = V/\Lambda$, then the Hodge decomposition $\Lambda \otimes \mathbb{C} \cong V \oplus \bar{V}$ respects the action of $\text{Aut}(A)$. Therefore, if g is of finite order with eigenvalues $e(x_1), \dots, e(x_n)$, then the multiset

$$(*) \quad \{e(x_1), \dots, e(x_n), e(-x_1), \dots, e(-x_n)\}$$

is $\text{Aut}(\mathbb{C})$ -stable. By a *type*, we mean a multiset $\{x_1, \dots, x_n\}$ with $x_i \in [0, 1)$ such that the multiset $(*)$ is $\text{Aut}(\mathbb{C})$ -stable. Equivalently, a type can be identified with a finitely supported function $f: \mathbb{Q}/\mathbb{Z} \rightarrow \mathbb{N}$ such that $f(x) + f(-x)$ depends only on the order of x in \mathbb{Q}/\mathbb{Z} . By the *weight* of $\{x_1, \dots, x_n\}$, we mean the sum $x_1 + \dots + x_n$, so that $\text{age}(g)$ is the weight of the type of g .

A type is *reduced* if 0 does not appear, and the reduced type of a given type is obtained by discarding all copies of 0. The *sum* of types is the union in the sense of multisets; at the level of associated functions on \mathbb{Q}/\mathbb{Z} it is the usual sum. A type which is not the sum of non-zero types is *primitive*. All the elements of a primitive type appear with multiplicity one, and they all have the same denominator. Every type can be realized (not necessarily uniquely) as a sum of primitive types; if the weight of the type is 1, each of the primitive types has weight ≤ 1 , so our first task is to classify primitive types with weight ≤ 1 .

A primitive type X of denominator $n \geq 2$ consists of fractions a_i/n where $0 < a_i < n$, and $(a_i, n) = 1$. Moreover, if $n \geq 3$, for each positive integer $a < n$ prime to n , exactly one of a/n and $1 - a/n$ belongs to X . If $a < b < n/2$ and $1 - a/n \in X$, then the weight of X exceeds 1 since either b/n or $1 - b/n$ belongs to X . Thus, if $0 < a < n/2$ and $1 - a/n \in X$, then a must be the largest integer in $(0, n/2)$ prime to n .

Lemma 1. *If $\phi(n) > 24$, then*

$$\sum_{x \in S_n} \min(x, n - x) > 2n,$$

where S_n is the set of positive integers $< n$ and prime to n . Moreover, the largest integer n such that $\phi(n) \leq 24$ is 90.

Proof. Note that

$$\min(x, n - x) > \frac{x(n - x)}{n} = \frac{n^2 - x^2 - (n - x)^2}{2n}.$$

In order to prove the first statement, we want to prove if $\phi(n) > 24$, then

$$\sum_{x \in S_n} \left(n^2 - x^2 - (n - x)^2 \right) > 4n^2,$$

or equivalently,

$$\phi(n)n^2 - 2 \sum_{x \in S_n} x^2 - 4n^2 > 0.$$

By Möbius inversion, one can prove that

$$\sum_{x \in S_n} x^2 = \frac{\phi(n)n^2}{3} + (-1)^{d_n} \frac{\phi(f(n))n}{6},$$

where $f(n)$ denotes the largest squarefree divisor of n and d_n is the number of distinct prime divisors of n . Thus, if $\phi(n) > 24$, then $\phi(n) > 24 \geq 12 \frac{n}{n-1}$, so $(n - 1)\phi(n) - 12n > 0$ and since $\phi(f(n)) \leq \phi(n)$,

$$\begin{aligned} \phi(n)n^2 - 2 \sum_{x \in S_n} x^2 - 4n^2 &\geq \left(\frac{\phi(n)}{3} - 4 \right) n^2 - \frac{\phi(n)n}{3} \\ &= \frac{n((n - 1)\phi(n) - 12n)}{3} > 0, \end{aligned}$$

which is the desired inequality.

For the second statement, if $\phi(n) \leq 24$ and p is a prime factor of n , then $\phi(p) = p - 1 \leq \phi(n) \leq 24$. Hence $p \leq 23$. Writing

$$n = 2^{n_2} 3^{n_3} 5^{n_5} 7^{n_7} 11^{n_{11}} 13^{n_{13}} 17^{n_{17}} 19^{n_{19}} 23^{n_{23}},$$

we have

$$0 \leq n_2 \leq 5, 0 \leq n_3 \leq 3, 0 \leq n_5 \leq 2,$$

and $0 \leq n_i \leq 1$ for $7 \leq i \leq 23$. Case analysis now shows $n \leq 90$. □

Proposition 2. *There are 28 primitive types with weight ≤ 1 :*

#	n	primitive types	weight
1	2	1/2	1/2
2	3	1/3	1/3
3		2/3	2/3
4	4	1/4	1/4
5		3/4	3/4
6	5	1/5, 2/5	3/5
7		1/5, 3/5	4/5
8	6	1/6	1/6
9		5/6	5/6
10	7	1/7, 2/7, 3/7	6/7
11		1/7, 2/7, 4/7	1
12	8	1/8, 3/8	1/2
13		1/8, 5/8	3/4
14	9	1/9, 2/9, 4/9	7/9
15		1/9, 2/9, 5/9	8/9
16	10	1/10, 3/10	2/5
17		1/10, 7/10	4/5
18	12	1/12, 5/12	1/2
19		1/12, 7/12	2/3
20	14	1/14, 3/14, 5/14	9/14
21		1/14, 3/14, 9/14	13/14
22	15	1/15, 2/15, 4/15, 7/15	14/15
23		1/15, 2/15, 4/15, 8/15	1
24	16	1/16, 3/16, 5/16, 7/16	1
25	18	1/18, 5/18, 7/18	13/18
26		1/18, 5/18, 11/18	17/18
27	20	1/20, 3/20, 7/20, 9/20	1
28	24	1/24, 5/24, 7/24, 11/24	1

Table 1

Proof. For $n \geq 3$, the weight of a primitive type of denominator n is at least

$$\sum_{\{x \in S_n \mid x < n/2\}} \frac{x}{n} \geq \frac{1}{2n} \sum_{x \in S_n} \min(x, n-x).$$

By Lemma 1, it suffices to carry out an exhaustive search up to $n = 90$. \square

Lemma 3. *There are 35 types with age 1 of automorphisms given in Table 2 below.*

#	n	types	notes
1	2	1/2, 1/2	Prop. 6
2	3	1/3, 1/3, 1/3	Th. 12
3		1/3, 2/3	Prop. 7
4	4	1/4, 1/4, 1/4, 1/4	Th. 12
5		1/4, 1/4, 2/4	$g^2 \rightarrow \#1$
6		1/4, 3/4	$g^2 \rightarrow \#1$
7	6	1/6, 1/6, 1/6, 1/6, 1/6, 1/6	Th. 12
8		1/6, 1/6, 1/6, 1/6, 2/6	Th. 12
9		1/6, 1/6, 1/6, 3/6	$g^2 \rightarrow \#2$
10		1/6, 1/6, 4/6	$g^3 \rightarrow \#1$
11		1/6, 5/6	$g^3 \rightarrow \#1$
12		1/6, 2/6, 3/6	$g^3 \rightarrow \#1$
13		1/6, 1/6, 2/6, 2/6	$g^3 \rightarrow \#1$
14	7	1/7, 2/7, 4/7	Cor. 10
15	8	1/8, 2/8, 5/8	$g^4 \rightarrow \#1$
16		1/8, 3/8, 4/8	$g^4 \rightarrow \#1$
17		1/8, 1/8, 3/8, 3/8	Th. 12
18		1/8, 2/8, 2/8, 3/8	$g^4 \rightarrow \#1$
19	10	1/10, 2/10, 3/10, 4/10	$g^5 \rightarrow \#1$
20	12	4/12, 2/12, 1/12, 5/12	$g^6 \rightarrow \#1$
21		4/12, 3/12, 3/12, 2/12	$g^6 \rightarrow \#1$
22		6/12, 1/12, 5/12	$g^6 \rightarrow \#1$
23		3/12, 3/12, 1/12, 5/12	$g^4 \rightarrow \#3$
24		2/12, 2/12, 2/12, 1/12, 5/12	$g^6 \rightarrow \#1$
25		1/12, 1/12, 5/12, 5/12	Th. 12
26		4/12, 1/12, 7/12	$g^6 \rightarrow \#1$
27		2/12, 2/12, 1/12, 7/12	$g^6 \rightarrow \#1$
28		3/12, 3/12, 2/12, 2/12, 2/12	$g^6 \rightarrow \#1$
29	15	1/15, 2/15, 4/15, 8/15	Cor. 11
30	16	1/16, 3/16, 5/16, 7/16	Cor. 9
31	20	1/20, 3/20, 7/20, 9/20	Cor. 9
32	24	1/24, 5/24, 7/24, 11/24	Cor. 9
33		8/24, 4/24, 3/24, 9/24	$g^{12} \rightarrow \#1$
34		3/24, 9/24, 2/24, 10/24	$g^{12} \rightarrow \#1$
35		4/24, 4/24, 4/24, 3/24, 9/24	$g^{12} \rightarrow \#1$

Table 2

Proof. Let $[a_i]$ be a formal variable representing the i th primitive type in Table 1, and let w_i denote the weight of the i th type. A monomial $\prod a_i^{m_i}$ stands for a sum of primitive types in which the i th type appears m_i times. The g.c.d. of the denominators of the w_i is 5040. Let $y = x^{1/5040}$, so $(1 - [a_i]x^{w_i})^{-1}$ is a power series in y for every i . By MAPLE 13, the coefficient

of y^{5040} in the product

$$\prod_{i=1}^{28} (1 - [a_i]x^{w_i})^{-1} = \prod_{i=1}^{28} (1 - [a_i]y^{w_i \cdot 5040})^{-1},$$

is $a1^2 + a2^3 + a2 a3 + a4 a5 + a8 a9 + a4 a13 + a6 a16 + a4^2 a1 + a28 + a27 + a23 + a24 + a11 + a19 a2 + a19 a8^2 + a18 a1 + a18 a4^2 + a18 a8^3 + a18 a12 + a18^2 + a12 a1 + a12 a4^2 + a12 a8^3 + a12^2 + a8^2 a2^2 + a8^2 a3 + a8^3 a1 + a8^3 a4^2 + a8^4 a2 + a8^6 + a4^4 + a18 a8 a2 + a12 a8 a2 + a8 a1 a2 + a8 a2 a4^2$.

Each monomial in this sum corresponds to an entry in Table 2. \square

3. RATIONAL CURVES IN $A/\langle g \rangle$

In this section we explain how to find rational curves on $A/\langle g \rangle$ in each case in Table 2.

Lemma 4. *If $A/\langle g^n \rangle$ has a rational curve for some positive integer n , then $A/\langle g \rangle$ has a rational curve.*

Proof. The morphism $A/\langle g^n \rangle \rightarrow A/\langle g \rangle$ is finite, so the image of a rational curve is again a rational curve. \square

Proposition 5. *Let A be an abelian variety and g an automorphism of finite order. Suppose that for every abelian variety B and finite-order automorphism $h \in \text{Aut}(B)$ whose type is the reduced type of g , $B/\langle h \rangle$ has a rational curve. Then $A/\langle g \rangle$ has a rational curve.*

Proof. Let B denote the image of $1 - g$ acting on A . Then B is an abelian subvariety of A , and g restricts to an automorphism h of B whose type is the reduced type of g . As $B/\langle h \rangle \subset A/\langle g \rangle$ has a rational curve, the same is true of $A/\langle g \rangle$. \square

The following proposition is well known.

Proposition 6. *If A is an abelian surface, then $A/\pm 1$ has a rational curve.*

Proof. Resolving the 16 singularities of $A/\pm 1$, we obtain a K3 surface with Picard number $\geq 16 \geq 5$. By work of Bogomolov and Tschinkel [1], any such surface is either elliptic or has infinite automorphism group and in either case has infinitely many rational curves, all but finitely many of which lie on $A/\pm 1$. \square

Note that Proposition 6 covers not only case #1 in Table 2 but twenty other cases as well, namely those (indicated in the “notes” column) for which the reduced type of some power of g is $(1/2, 1/2)$.

Proposition 7. *Let $A \cong V/\Lambda$ be an abelian surface with an automorphism g of type $(1/3, 2/3)$. Then $A/\langle g \rangle$ contains a rational curve.*

Proof. Let $G = \langle g \rangle$ and $X = A/G$. Regarding $1 - g$ as an isogeny of A , the number of fixed points of g is

$$\deg(1 - g) = \# \ker(1 - g) = \det(1 - g|\Lambda) = 3^2 = 9.$$

These are singularities of type A_2 , since under $(x, y) \mapsto (\omega x, \omega^2 y)$ where ω is a cube root of unity, the invariants are generated by $X = x^3, Y = y^3, Z = xy$, and so

$$\mathbb{C}[[x, y]]^G = \mathbb{C}[[X, Y, Z]]/(XY - Z^3).$$

This is isomorphic to $\mathbb{C}[[x, y, z]]/(x^2 + y^2 + z^3)$, which has a Du Val singularity of type A_2 (see [8, Ch.4, 4.2]).

Consider the minimal resolution $f : Y \rightarrow X$, for which the 9 exceptional divisors Y_i each consists of two projective lines $D_{i,1}$ and $D_{i,2}$ intersecting at one point. The canonical divisor of Y is $K_Y = f^*K_X = 0$. Hence Y is a K3-surface of Picard number ≥ 18 , and again by [1], we deduce that X has infinitely many rational curves.. \square

Theorem 8. *Let B be an abelian variety and $h \in \text{Aut}(B)$ an automorphism of finite order such that h and h^{-1} have disjoint types. Let A be an abelian variety with a finite order automorphism g whose type is contained in that of h . If there is a rational curve on $B/\langle h \rangle$ then there is a rational curve on $A/\langle g \rangle$.*

Proof. It suffices to prove that there exists a surjective homomorphism $p: B \rightarrow A$ such that the diagram

$$\begin{array}{ccc} B & \xrightarrow{p} & A \\ h \downarrow & & \downarrow g \\ B & \xrightarrow{p} & A \end{array}$$

commutes. Writing $B = \text{Lie}(B)/\Lambda_B$ and $A = \text{Lie}(A)/\Lambda_A$, the goal is to find a surjective $\mathbb{C}[t]$ -linear map $\phi: \text{Lie}(B) \rightarrow \text{Lie}(A)$ (where t acts as g on $\text{Lie}(A)$ and as h on $\text{Lie}(B)$) such that $\phi(\Lambda_B) \subset \Lambda_A$. If ψ is a surjective $\mathbb{C}[t]$ -linear map $\text{Lie}(B) \rightarrow \text{Lie}(A)$ such that $\psi(\Lambda_B \otimes \mathbb{Q}) = \Lambda_A \otimes \mathbb{Q}$, then we can define $\phi := n\psi$ for n a sufficiently divisible positive integer. It suffices, therefore, to find ψ with the desired properties.

As the type of A is a subset of the type of B , there exists a surjective $\mathbb{Q}[t]$ -linear map $T: \Lambda_B \otimes \mathbb{Q} \rightarrow \Lambda_A \otimes \mathbb{Q}$. Extending scalars to \mathbb{C} , $T \otimes 1$ maps $\Lambda_B \otimes \mathbb{C} = \overline{\text{Lie}(B)} \oplus \overline{\text{Lie}(B)}$ to $\Lambda_A \otimes \mathbb{C} = \text{Lie}(A) \oplus \overline{\text{Lie}(A)}$. The type of g acting on $\overline{\text{Lie}(A)}$ is the same as the type of g^{-1} acting on $\text{Lie}(A)$ and therefore disjoint from the type of g acting on $\text{Lie}(A)$, and the same is true for the type of h acting on $\text{Lie}(B)$. As $\text{Lie}(B)$ and $\text{Lie}(A)$ are direct sums of certain t -eigenspaces of $\Lambda_B \otimes \mathbb{C}$ and $\Lambda_A \otimes \mathbb{C}$ respectively and as the spectrum of t acting on $\text{Lie}(A)$ is the intersection of the spectra of t acting on $\text{Lie}(B)$ and on $\Lambda_A \otimes \mathbb{C}$, it follows that $T \otimes 1$ maps $\text{Lie}(B)$ to $\text{Lie}(A)$. The restriction of ψ to $\text{Lie}(B)$ is therefore the desired map. \square

Corollary 9. *Let $n \geq 3$ be a positive integer, and let $m = \lceil n/2 \rceil - 1$. If A is an abelian variety and $g \in \text{Aut}(A)$ is an automorphism of order n whose type is contained in $\{1/n, 2/n, \dots, m/n\}$, then $A/\langle g \rangle$ has a rational curve.*

Proof. Let X denote the non-singular projective hyperelliptic curve of genus m which contains the affine curve $y^2 = x^n - 1$. The order- n automorphism $h(x, y) = (e(1/n)x, y)$ extends to an automorphism of X and therefore defines an automorphism of $B := \text{Jac}(X)$. The Lie algebra $\text{Lie}(B)$ can be identified with the space $H^0(X, \Omega_X)$ of holomorphic differential forms on X , which has a basis

$$\left\{ \frac{dx}{y}, \frac{x dx}{y}, \dots, \frac{x^{m-1} dx}{y} \right\}.$$

Therefore, the type of h acting on B is $\{1/n, 2/n, \dots, m/n\}$, which is disjoint from the type of h^{-1} . On the other hand, $B/\langle h \rangle$ contains the rational curve $X/\langle h \rangle$. Thus, Theorem 8 applies. \square

Corollary 10. *If A is an abelian 3-fold and g is an automorphism of A of type $(1/7, 2/7, 4/7)$, then $A/\langle g \rangle$ has a rational curve.*

Proof. Let X denote the Klein quartic:

$$X : x^3y + y^3z + z^3x = 0$$

and B the Jacobian of X . The self-map $h(x, y, z) = (\zeta_7x, \zeta_7^4y, \zeta_7^2z)$ of X belongs to the automorphism group $\text{PSL}_2(\mathbb{F}_7)$ of X which acts non-trivially on the Jacobian variety B and therefore on $\text{Lie}(B) = H^0(X, \Omega_X)$. Conjugating h by the cyclic permutations of (x, y, z) , we see that h is conjugate to h^2 and h^4 in $\text{Aut}(X)$, and therefore the type of h is invariant under multiplication by 2 (mod 1). It is therefore $(1/7, 2/7, 4/7)$ or $(3/7, 5/7, 6/7)$, and replacing h by h^{-1} if necessary, we may assume that it is the former. \square

We remark that $B/\langle h \rangle$ in the proof of Corollary 10 has appeared in the literature; it is known to have a Calabi-Yau resolution [2, Example 6.3].

Corollary 11. *If A is an abelian variety and g an automorphism such that the type of A is contained in $(1/15, 2/15, 3/15, 4/15, 8/15, 9/15)$, then $A/\langle g \rangle$ has a rational curve.*

Proof. Let X be the non-singular projective curve which has a (singular, affine) model $X' : y^{15} = x^2(x - 1)$. This is singular only at $(0, 0)$, and the inverse image of this singularity under the normalization map $X \setminus \{P_\infty\} \rightarrow X'$ is a single point $P_0 \in X$. The automorphism $h : (x, y) \mapsto (x, e(1/15)y)$ of the affine curve induces an automorphism of X of order 15. As $15y^{14}dy = (3x^2 - 2x)dx$, any differential form $\frac{x^m y^n dy}{3x^2 - 2x}$, $m, n \geq 0$, is holomorphic except possibly at P_0 and P_∞ . One checks that

$$\frac{dy}{3x - 2}, \frac{ydy}{3x - 2}, \frac{y^2dy}{3x - 2}, \frac{y^3dy}{3x - 2}, \frac{y^7dy}{3x^2 - 2x}, \frac{y^8dy}{3x^2 - 2x}$$

are all holomorphic, and their eigenvalues under h are $e(1/15)$, $e(2/15)$, $e(3/15)$, $e(4/15)$, $e(8/15)$, $e(9/15)$ respectively. Applying the Riemann-Hurwitz theorem to the map $X \rightarrow \mathbb{P}^1$ given by y , we see that X is of genus 6, and therefore, that these differential forms form a basis of $\text{Lie}(\text{Jac}(B)) = H^0(X, \Omega_X)$. \square

Theorem 12. *If A is an abelian variety and g is an automorphism of finite order such that g and g^{-1} have disjoint types and the type of g is a sum of primitive types at least one of which has weight less than 1, then $A/\langle g \rangle$ has a rational curve.*

Proof. For every primitive type, there exists an abelian variety B_i with complex multiplication and an automorphism h_i of B_i with the given type. Indeed, the primitive types of denominator n are in natural correspondence with CM-types on $\mathbb{Q}(\zeta_n) = \mathbb{Q}(e(1/n))$. Any CM-type Φ on $\mathbb{Q}(\zeta_n)$, $n \geq 3$ defines an embedding $\mathbb{Q}(\zeta_n) \rightarrow \mathbb{C}^{\phi(n)/2}$. The image of $\mathbb{Z}[\zeta_n]$ defines a lattice $\Lambda \subset \mathbb{C}^{\phi(n)/2}$, and the quotient $\mathbb{C}^{\phi(n)/2}/\Lambda$ is a complex torus with a natural action of $\mathbb{Z}/n\mathbb{Z}$ of the type associated with Φ . The quotient $\mathbb{C}^{\phi(n)/2}/\Lambda$ admits a polarization [9, II 6 Theorem 4], so there exists a pair (B_i, h_i) as claimed. If $\text{age}(h_1) < 1$, then by [3], $B_1/\langle g_1 \rangle$ has a rational curve. If $A = A_1 \times \cdots \times A_m$, and $g = (g_1, \dots, g_m)$ is a finite order automorphism of A which stabilizes each factor, then $A_1/\langle g_1 \rangle \subset A/\langle g \rangle$, so $A/\langle g \rangle$ has a rational curve. The theorem now follows from Theorem 8. \square

To summarize, we have the following theorem:

Theorem 13. *Let A be an abelian variety with a nontrivial automorphism g of finite order such that $\text{age}(g) \leq 1$. Then $A/\langle g \rangle$ contains a rational curve.*

Corollary 14. *Let A be an abelian variety of dimension n with a nontrivial automorphism g of finite order. If $\dim(\ker(1 - g)) \geq n - 2$ (i.e. the codimension of the fixed subspace of A under g is less than or equal to 2), then the quotient $A/\langle g \rangle$ contains a uniruled hypersurface.*

Proof. Since $\dim(\ker(1 - g)) \geq n - 2$, $B := \text{im}(1 - g)$ is an abelian variety of dimension $n - \dim(\ker(1 - g)) \leq 2$. Let h denote the restriction of g to B . As $\text{age}(h) + \text{age}(h^{-1}) \geq 2$, we may assume without loss of generality that $\text{age}(h) \leq 1$, so $B/\langle h \rangle$ has a rational curve Z by Theorem 13. Let C denote the identity component of $\ker(1 - g)$, which is an abelian subvariety of dimension $\dim \ker(1 - g)$ on which g acts trivially. The addition morphism $B \times C \rightarrow A$ is an isogeny and respects the action of $\langle g \rangle$. We therefore obtain a finite morphism from

$$Z \times C \subset (B/\langle g \rangle) \times C \cong (B \times C)/\langle g \rangle$$

to $A/\langle g \rangle$. The image of an $n - 1$ -dimensional ruled variety under a finite morphism is a uniruled hypersurface. \square

Corollary 15. *Let E be an elliptic curve. If W is a Weyl group of simple roots of rank $n \geq 3$ acting on E^n and W^+ is an index 2-subgroup of W , then the quotient E^n/W^+ contains a rational curve.*

Proof. W is generated by reflections s_j of simple roots. Since W^+ is an index 2-subgroup of W , there exist two reflections s_1 and s_2 such that $s_1 s_2 \in W^+$. Then for each i , $\ker(1 - s_i)$ has codimension 1 and their intersection has codimension ≤ 2 . Since $\ker(1 - s_1 s_2)$ contains the intersection of $\ker(1 - s_1)$ and $\ker(1 - s_2)$, this follows from Corollary 14. \square

REFERENCES

- [1] Bogomolov, Fedor; Tschinkel, Yuri: Density of rational points on elliptic K3 surfaces, *Asian J. Math.* **4**, (2000), 351–368,
- [2] Bogomolov, Fedor; Tschinkel, Yuri: Rational curves and points on K3 surfaces. *Amer. J. Math.* **127** (2005), no. 4, 825–835.
- [3] Kollár, János; Larsen, Michael: Quotients of Calabi-Yau varieties. Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, 179–211, Progr. Math., 270, Birkhauser Boston, Inc., Boston, MA, 2009.
- [4] Lang, Serge: Abelian varieties, New York: Springer-Verlag, 1983.
- [5] Larsen, Michael: Rank of elliptic curves over almost algebraically closed fields. *Bull. London Math. Soc.* **35** (2003), no. 6, 817–820.
- [6] Looijenga, Eduard: Root systems and elliptic curves. *Invent. Math.* **38** (1976/77), no. 1, 17–32.
- [7] Pirola, Gian Pietro: Curves on generic Kummer varieties. *Duke Math. J.* **59** (1989), no. 3, 701–708.
- [8] Reid, Miles: Chapters on algebraic surfaces, Complex algebraic geometry (Park City, UT, 1993), 3–159, IAS/Park City Math. Ser., 3, Amer. Math. Soc., Providence, RI, 1997.
- [9] Shimura, Goro: Abelian Varieties with Complex Multiplication and Modular Functions. Princeton Mathematical Series, 46. Princeton University Press, Princeton, NJ, 1998.

DEPARTMENT OF MATHEMATICS, CHUNG-ANG UNIVERSITY, 221, HEUKSEOK-DONG,
DONGJAK-GU, SEOUL, 156-756, SOUTH KOREA
E-mail address: bohaeim@gmail.com, imbh@cau.ac.kr

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405,
USA
E-mail address: mjlarson@indiana.edu