

**ROOTS MULTIPLICITY AND SQUARE-FREE FACTORIZATION
OF POLYNOMIALS USING COMPANION MATRICES**

N. H. GUERSENZVAIG

Av. Corrientes 3985 6A, (1194) Buenos Aires, Argentina

email: nguersenz@fibertel.com.ar

and

FERNANDO SZECHTMAN¹

Department of Mathematics and Statistics, University of Regina, Saskatchewan, Canada

email: fernando.szechtman@gmail.com

Abstract

Given an arbitrary monic polynomial f over a field F of characteristic 0, we use companion matrices to construct a polynomial $M_f \in F[X]$ of minimum degree such that for each root α of f in the algebraic closure of F , $M_f(\alpha)$ is equal to the multiplicity $m(\alpha)$ of α as a root of f . As an application of M_f we give a new method to compute in $F[X]$ each component of the square-free factorization $f = P_1 P_2^2 \cdots P_m^m$, where P_k is the product of all $X - \alpha$ with $m(\alpha) = k$, for $k = 1, \dots, m = \max m(\alpha)$.

Keywords: Lagrange Interpolation Formula, square-free factorization, companion matrix.

2010 MSC: 12D05, 13A05, 15A24.

1. INTRODUCTION

We fix throughout the paper a field F of characteristic 0 and a monic polynomial $f \in F[X]$ of degree n . We say that f is square-free if for any monic polynomial $g \in F[X]$: $g^2 | f$ implies $g = 1$, or, equivalently, if $\gcd(f, f') = 1$. We assume henceforth that $n \geq 1$ and that f has prime factorization

$$(1.1) \quad f = f_1^{m_1} \cdots f_r^{m_r},$$

so f is square-free if and only if each $m_j = 1$. Set $m = \max_{1 \leq j \leq r} m_j$ and collect in P_k all monic irreducible factors f_j of f having multiplicity $m_j = k$ in (1.1), that is,

$$P_k = \prod_{\substack{1 \leq j \leq r \\ m_j = k}} f_j, \quad k = 1, \dots, m.$$

Each P_k is square-free, yielding the so-called square-free factorization of f , namely

$$(1.2) \quad f = P_1 P_2^2 \cdots P_m^m.$$

Efficient algorithms to compute (1.2) were developed about forty years ago by Tobey [7], Horowitz [4, 5], Musser [6] and Yun [8]. A more recent algorithm can be found in [1]. The square-free factorization (1.2) is usually utilized as a first step towards the computation of the full factorization (1.1) as well as to integrate rational functions, following a method due to Hermite [3], as explained in [1].

Let us review the Tobey-Horowitz algorithm, one of the simplest known methods to obtain (1.2). Let $g = f_j$, where $1 \leq j \leq r$, and set $k = m_j$ as well as $h = f/g^k$. We then have $f = g^k h$ with $\gcd(g, h) = 1$. Moreover,

$$f' = k g^{k-1} g' h + g^k h'.$$

¹Corresponding author

As g is relatively prime to $kg'h$, the multiplicity of g in $\gcd(f, f')$ is $k - 1$. Hence

$$\gcd(f, f') = P_2 P_3^2 \cdots P_m^{m-1}.$$

Setting

$$D_0 = f, D_1 = \gcd(D_0, D'_0), D_2 = \gcd(D_1, D'_1), \dots$$

a repeated application of the preceding discussion yields

$$D_k = \begin{cases} P_{k+1} P_{k+2}^2 \cdots P_m^{m-k} & k = 0, \dots, m-1, \\ 1 & k \geq m, \end{cases}$$

whence

$$(1.3) \quad \frac{D_{k-1}}{D_k} = P_k P_{k+1} \cdots P_m, \quad k = 1, \dots, m-1.$$

Therefore, not only m can be recognized as the smallest positive integer k such that $D_k = 1$, but we also we have

$$P_k = \frac{D_{k-1}}{D_k} : \frac{D_k}{D_{k+1}} = \frac{D_{k-1} D_{k+1}}{D_k^2}, \quad k = 1, \dots, m.$$

In this paper we furnish an entirely new practical method to compute (1.2) by means of Lagrange Interpolation Formula and companion matrices.

Let K be a splitting field of f over F and let $S(f)$ be the set of roots of f in K , whose size will be denoted by s . For each $\alpha \in S(f)$ let $m(\alpha)$ be the multiplicity of α as a root of f . Lagrange Interpolation Formula ensures the existence and uniqueness of a polynomial $M_f \in K[X]$ of degree less than s satisfying

$$(1.4) \quad M_f(\alpha) = m(\alpha), \quad \alpha \in S(f).$$

The polynomial $f_0 = f_1 \cdots f_r$ is called the square-free part of f . Clearly,

$$f_0 = \prod_{\alpha \in S(f)} (X - \alpha) = P_1 \cdots P_m.$$

Applying case $k = 1$ of (1.3) we obtain

$$f_0 = f / \gcd(f, f').$$

Combining the two preceding equations we obtain

$$(1.5) \quad P_k = \prod_{\substack{\alpha \in S(f) \\ m(\alpha)=k}} (X - \alpha) = \gcd(M_f - k, f / \gcd(f, f')), \quad k = 1, \dots, m.$$

The practical use of (1.5) seems to be limited due to the mysterious nature of M_f . A closer look reveals that

$$(1.6) \quad M_f = \sum_{\alpha \in S(f)} \frac{m(\alpha)}{f'_0(\alpha)} \frac{f_0}{X - \alpha},$$

which can be verified by evaluating both sides at each $\alpha \in S(f)$. This seems to confirm the ineffectiveness of (1.5), as (1.6) requires the use of all roots of f in K . However, applying the Galois group $G = \text{Gal}(K/F)$ to (1.4) we see that $M_f \in F[X]$. Indeed, let $\sigma \in G$. Then

$$M_f^\sigma(\alpha^\sigma) = M_f(\alpha)^\sigma = m(\alpha)^\sigma = m(\alpha) = m(\alpha^\sigma) = M_f(\alpha^\sigma),$$

so $M_f^\sigma = M_f$ by uniqueness, whence $M_f \in F[X]$, as claimed. The fact that $M_f \in F[X]$ suggests that there should be a rational procedure to obtain M_f from f . This is exactly what we do in this paper, by means of companion matrices.

The outcome is an efficient procedure to compute all components P_k of (1.2). Indeed, first compute $f_0 = f / \gcd(f, f')$, then M_f , as indicated in §2, and then proceed sequentially to find all P_1, P_2, \dots making use of (1.5), stopping at m , namely the smallest positive integer k satisfying $\deg(P_1) + 2 \deg(P_2) + \dots + k \deg(P_k) = n$.

2. USING COMPANION MATRICES TO FIND M_f FROM f

Given a monic polynomial $g = g_0 + g_1X + \dots + g_{s-1}X^{s-1} + X^s \in F[X]$ of positive degree s , its companion matrix $C_g \in M_s(F)$ is defined

$$C_g = \begin{pmatrix} 0 & 0 & \cdots & 0 & -g_0 \\ 1 & 0 & \cdots & 0 & -g_1 \\ 0 & 1 & \cdots & 0 & -g_2 \\ \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -g_{s-1} \end{pmatrix}.$$

We will use below the well-known fact that $g(C_g) = 0$. Let $F_s[X]$ be the subspace of $F[X]$ with basis $1, X, \dots, X^{s-1}$ and let $[R] \in F^s$ stand for the coordinates of $R \in F_s[X]$ relative to this basis. We will also use the following formula from [2]:

$$(2.1) \quad R(C_g) = ([R] \ C_g[R] \ C_g^2[R] \ \cdots \ C_g^{s-1}[R]).$$

Since the first column of a product of matrices, say AB , is equal to the product of A by the first column of B , it follows from (2.1) that for any $P, Q, T \in F_s[X]$,

$$(2.2) \quad P(C_g)Q(C_g) = T(C_g) \iff P(C_g)[Q] = [T].$$

Theorem 2.1. *Let F be a field of characteristic 0 and let $f \in F[X]$ be monic of degree $n \geq 1$. Set $f_0 = f / \gcd(f, f')$, $s = \deg(f_0)$ and $P = f' / \gcd(f, f')$. Let M_f , $S(f)$ and $m(\alpha)$ be as in Section 1. Then there exist unique $g \in F_s[X]$ and $h \in F[X]$ such that $f'_0g + f_0h = 1$. Moreover, for such P , f_0 and g ,*

$$[M_f] = P(C_{f_0})[g].$$

Proof. Definition of f_0 given in Section 1 guarantees $\gcd(f_0, f'_0) = 1$, which ensures the existence and uniqueness of g and h . Let R be the remainder of dividing $M_f f'_0$ by f_0 . By Lagrange Interpolation Formula, we have

$$R = \sum_{\alpha \in S(f)} \frac{R(\alpha)}{f'_0(\alpha)} \frac{f_0}{X - \alpha} = f_0 \sum_{\alpha \in S(f)} \frac{m(\alpha)}{X - \alpha} = f_0 \frac{f'}{f} = P.$$

Therefore,

$$M_f(C_{f_0})f'_0(C_{f_0}) = R(C_{f_0}) = P(C_{f_0}).$$

But $f'_0(C_{f_0})g(C_{f_0}) = I_s$, so

$$M_f(C_{f_0}) = P(C_{f_0})g(C_{f_0}),$$

which by (2.2) is equivalent to what we want to prove. \blacksquare

Example 2.2. Let $f = X^4 - 4X + 3 \in \mathbb{Q}[X]$. Then

$$f' = 4X^3 - 4 \text{ and } \gcd(f, f') = X - 1,$$

4

so

$$f_0 = X^3 + X^2 + X - 3, \quad f'_0 = 3X^2 + 2X + 1 \quad \text{and} \quad P = 4X^2 + 4X + 4.$$

We achieve $f'_0 g + f_0 h = 1$ by taking

$$g = (1/72)X^2 + (1/9)X + (1/24), \quad h = (-1/24)X - (23/72).$$

Using

$$(2.3) \quad C_{f_0} = \begin{pmatrix} 0 & 0 & 3 \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix},$$

we obtain

$$[M_f] = ([P] C_{f_0} [P] C_{f_0}^2 [P])[g] = \begin{pmatrix} 4 & 12 & 0 \\ 4 & 0 & 12 \\ 4 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1/24 \\ 1/9 \\ 1/72 \end{pmatrix} = \begin{pmatrix} 3/2 \\ 1/3 \\ 1/6 \end{pmatrix},$$

which means

$$(2.4) \quad M_f = \frac{1}{6}X^2 + \frac{1}{3}X + \frac{3}{2}.$$

Next we compute

$$P_1 = \gcd(M_f - 1, f_0) = X^2 + 2X + 3, \quad P_2 = \gcd(M_f - 2, f_0) = X - 1,$$

which yields the square-free factorization, $f = P_1 P_2^2$.

Remark 2.3. It is possible to find the degrees of the P_k 's before actually computing these polynomials. This can be done as follows. Suppose f_0 has roots $\alpha_1, \dots, \alpha_s$ in K . Then C_{f_0} is similar to the diagonal matrix $\text{Diag}(\alpha_1, \dots, \alpha_s)$, so $M_f(C_{f_0})$ is similar to $\text{Diag}(m(\alpha_1), \dots, m(\alpha_s))$ and has characteristic polynomial

$$\text{Char}_{M_f(C_{f_0})} = \prod_{1 \leq i \leq s} (X - m(\alpha_i)) = \prod_{1 \leq k \leq \max m(\alpha_i)} (X - k)^{\deg(P_k)}.$$

For instance, in Example 2.2 we can use (2.3) and (2.4) to obtain

$$M_f(C_{f_0}) = \begin{pmatrix} 3/2 & 1/2 & 1/2 \\ 1/3 & 4/3 & 1/3 \\ 1/6 & 1/6 & 7/6 \end{pmatrix}.$$

Therefore

$$\text{Char}_{M_f(C_{f_0})} = X^3 - 4X^2 + 5X - 2 = (X - 1)^2(X - 2),$$

which indicates the square-free factorization $f = P_1 P_2^2$, with $\deg(P_1) = 2$ and $\deg(P_2) = 1$, in complete agreement with what we found above.

Remark 2.4. Let R be an integrally closed domain of characteristic 0 with field of fractions F , and let $f \in R[X]$ be monic of degree $n \geq 1$. Then the square-free components of f , as found above, will have coefficients in R . Indeed, (1.2) shows that all α_i , and hence the coefficients of all P_k , are integral over R . But these coefficients are in F , and hence lie in R .

Acknowledgements. We thank V. Sergeichuk for valuable comments.

REFERENCES

- [1] J. Gerhard, Fast modular algorithms for squarefree factorization and Hermite integration, *Appl. Algebra Eng. Commun. Comput.* 11 (2001) 203–226.
- [2] N. H. Guersenzvaig and F. Szechtman, A closed formula for the product in simple integral extensions, *Linear Algebra Appl.* 430 (2009) 2464–2466.
- [3] C. Hermite, Sur l'integration des fonctions rationnelles, *Annales de Mathématique, 2^{ème} série* 11 (1872) 145–148
- [4] E. Horowitz, Algorithms for symbolic integration of rational functions, Ph.D. Thesis, University of Wisconsin, 1969.
- [5] E. Horowitz, Algorithms for partial fraction decomposition and rational function integration, *Proc. ACM Symp. on Symbolic and algebraic manipulation* (1971) 441–457
- [6] D. R. Musser, Algorithms for polynomial factorization, Ph.D. Thesis, University of Wisconsin, 1971.
- [7] R. G. Tobey, Algorithms for antidifferentiation of rational functions, Ph.D. Thesis, Harvard, 1967.
- [8] D.Y.Y. Yun, On square-free decomposition algorithms, *Symbolic and algebraic computation, Proc. 1976 ACM Symp.* (1976) 26–35.