

Intrinsic equations for a relaxed elastic line of second kind in Minkowski 3-space

Ergin Bayram and Emin Kasap
Ondokuz Mayıs University, Faculty of Arts and Sciences,
Mathematics Department, Samsun, Turkey

September 25, 2018

Abstract

Let α be an arc on a connected oriented surface S in Minkowski 3-space, parameterized by arc length s , with torsion τ and length l . The total square torsion H of α is defined by $H = \int_0^l \tau^2 ds$. The arc is called a relaxed elastic line of second kind if it is an extremal for the variational problem of minimizing the value of H within the family of all arcs of length l on S having the same initial point and initial direction as α . In this study, we obtain differential equation and boundary conditions for a relaxed elastic line of second kind on an oriented surface in Minkowski 3-space.

1 Preliminaries and Introduction

In this section, we give some fundamentals required for this paper.

Definition 1 E^n equipped with the metric

$$\langle u, w \rangle = - \sum_{i=1}^{\nu} u_i w_i + \sum_{j=\nu+1}^n u_j w_j, \quad u, w \in E^n, \quad 0 \leq \nu \leq n,$$

is called semi-Euclidean space and is denoted by E_ν^n , where ν is called the index of the metric. For $n = 3$, E_1^3 is called Minkowski 3-space [3].

Let $\alpha(s)$ denote an arc on a connected oriented surface S in E_1^3 , parameterized by arc length s , $0 \leq s \leq l$, with curvature $\kappa(s)$ and torsion $\tau(s)$. Let the energy density be given as some function of the curvature and torsion, $f(\kappa, \tau)$. Then

$$H = \int f(\kappa, \tau) ds \quad (1)$$

is an Hamiltonian for curves [1]. Thus the following integral can be taken as a special case of Hamiltonians for curves:

$$H = \int \tau^2 ds. \quad (2)$$

Ekici and Görgülü [5] handled the problem of minimization of the integral $\int \kappa^2 \tau ds$ in Minkowski 3-space. In [4] authors defined relaxed elastic line of second kind on an oriented surface in Minkowski space and for a curve of this kind lying on an oriented surface the Euler-Lagrange equations were derived. In this paper, we give intrinsic equations for a curve of this kind. Particularly, we obtain differential equation and boundary conditions for a curve to be a relaxed elastic line of second kind on an oriented surface.

Definition 2 *The arc α is called a relaxed elastic line of second kind in Minkowski 3-space if it is an extremal for the variational problem of minimizing the value of H within the family of all arcs of length l on S having the same initial point and initial direction as α in Minkowski 3-space.*

In this study, we would like to calculate the intrinsic equations for the curve α which is an extremal for (1). We shall require that the coordinate functions of S are smooth enough to have partial derivatives and coordinate functions of α , as functions of s , are smooth enough in these coordinates.

Definition 3 *A tangent vector v in E_1^3 is*
spacelike if $\langle v, v \rangle > 0$ or $v = 0$,
null if $\langle v, v \rangle = 0$ and $v \neq 0$,
timelike if $\langle v, v \rangle < 0$.

Definition 4 *At a point $\alpha(s)$ of α , let T denote the unit tangent vector to α , n the unit normal to S , and*

$$n \times T = \varepsilon Q(s), \quad \varepsilon = \pm 1, \quad (3)$$

respectively. Then $\{T, Q, n\}$ gives an orthonormal basis in E_1^3 . If S is a spacelike surface then $T \times Q = n$, $Q \times n = -T$, $n \times T = -Q$. Similarly if S is a timelike surface then $T \times Q = -n$, $Q \times n = \pm T$, $n \times T = \mp Q$ [2].

Theorem 5 *Let S be a surface in E_1^3 and α be a curve on S . The analogue of the Frenet-Serret formulas is given by*

$$\begin{pmatrix} T' \\ Q' \\ n' \end{pmatrix} = \begin{pmatrix} 0 & \varepsilon_2 \kappa_g & \varepsilon_3 \kappa_n \\ -\varepsilon_1 \kappa_g & 0 & \varepsilon_3 \tau_g \\ -\varepsilon_1 \kappa_n & -\varepsilon_2 \tau_g & 0 \end{pmatrix} \begin{pmatrix} T \\ Q \\ n \end{pmatrix}, \quad (4)$$

where $\varepsilon_1 = \langle T, T \rangle$, $\varepsilon_2 = \langle Q, Q \rangle$, $\varepsilon_3 = \langle n, n \rangle$. Here $k_g(s) = \langle T'(s), Q(s) \rangle$, $\tau_g(s) = \langle Q'(s), n(s) \rangle$ and $k_n(s) = \langle II(T(s), T(s)), n(s) \rangle$ are geodesic curvature, geodesic torsion and normal curvature of α , respectively.

Theorem 6 *Let α be any regular curve on a surface in E_1^3 . Then we have*

$$\kappa = \frac{\|\alpha' \times \alpha''\|}{\|\alpha'\|^3}, \quad \tau = \varepsilon_1 \varepsilon_2 \frac{\langle \alpha' \times \alpha'', \alpha''' \rangle}{\|\alpha' \times \alpha''\|^2}.$$

2 Derivation of equations

Suppose that α lies in a coordinate patch $(u, v) \rightarrow x(u, v)$ of a surface S , and let $x_u = \partial x / \partial u$, $x_v = \partial x / \partial v$. Then α is expressed as

$$\alpha(s) = x(u(s), v(s)), \quad 0 \leq s \leq l,$$

with

$$T(s) = \alpha'(s) = \frac{du}{ds} x_u + \frac{dv}{ds} x_v$$

and

$$Q(s) = p(s) x_u + q(s) x_v$$

for suitable scalar functions $p(s)$ and $q(s)$.

Now we will define variational fields for our problem. In order to obtain variational arcs of length l , we need to extend α to an arc $\alpha^*(s)$ defined for $0 \leq s \leq l^*$, with $l^* \geq l$ but sufficiently close to l so that α^* lies in the coordinate patch. Let $\mu(s)$, $0 \leq s \leq l^*$, be a scalar function of class C^2 , not vanishing identically. Define

$$\eta(s) = \mu(s) p^*(s), \quad \zeta(s) = \mu(s) q^*(s).$$

Then

$$\eta(s) x_u + \zeta(s) x_v = \mu(s) Q(s) \tag{5}$$

along α . Also assume that

$$\mu(0) = 0, \quad \mu'(0) = 0. \tag{6}$$

Now define

$$\beta(\sigma; t) = x(u(\sigma) + t\eta(\sigma), v(\sigma) + t\zeta(\sigma)), \tag{7}$$

for $0 \leq \sigma \leq l^*$. For $|t| < \varepsilon_1$ (where $\varepsilon_1 > 0$ depends upon the choice of α^* and

of μ), the point $\beta(\sigma; t)$ lies in the coordinate patch. For fixed t , $\beta(\sigma; t)$ gives an arc with the same initial point and initial direction as α , because of (6). For $t = 0$, $\beta(\sigma; 0)$ is the same as α^* and σ is arc length. For $t \neq 0$, the parameter σ is not arc length in general.

For fixed t , $|t| < \varepsilon_1$, let $L^*(t)$ denote the length of the arc $\beta(\sigma; t)$, $0 \leq \sigma \leq l^*$. Then

$$L^*(t) = \int_0^{l^*} \sqrt{\left| \left\langle \frac{\partial \beta}{\partial \sigma}, \frac{\partial \beta}{\partial \sigma} \right\rangle \right|} d\sigma \tag{8}$$

with

$$L^*(0) = l^* > l. \quad (9)$$

By (7) and (8) $L^*(t)$ is continuous and differentiable in t . Particularly, it follows from (9) that

$$L^*(t) > \frac{l+l^*}{2} > l \quad \text{for } |t| < \varepsilon \quad (10)$$

for a suitable ε satisfying $0 < \varepsilon \leq \varepsilon_1$. Because of (10) one can restrict $\beta(\sigma; t)$, $0 \leq |t| < \varepsilon$, to an arc of length l by restricting the parameter σ to an interval $0 \leq \sigma \leq \lambda(t) \leq l^*$ by requiring

$$\int_0^{\lambda(t)} \sqrt{\left| \left\langle \frac{\partial \beta}{\partial \sigma}, \frac{\partial \beta}{\partial \sigma} \right\rangle \right|} d\sigma = l. \quad (11)$$

Note that $\lambda(0) = l$. The function $\lambda(t)$ need not be determined explicitly, but we shall need

$$\left. \frac{d\lambda}{dt} \right|_{t=0} = \varepsilon_1 \int_0^l \mu \kappa_g ds. \quad (12)$$

The proof of (12) and of other results will depend on calculations from (7) such as

$$\left. \frac{\partial \beta}{\partial \sigma} \right|_{t=0} = T, \quad 0 \leq s \leq l, \quad (13)$$

which gives

$$\left. \frac{\partial^2 \beta}{\partial \sigma^2} \right|_{t=0} = T' = \varepsilon_2 \kappa_g Q + \varepsilon_3 \kappa_n n. \quad (14)$$

Also

$$\left. \frac{\partial \beta}{\partial t} \right|_{t=0} = \mu Q \quad (15)$$

because of (5). Further differentiation of (15) gives

$$\left. \frac{\partial^2 \beta}{\partial t \partial \sigma} \right|_{t=0} = \left. \frac{\partial^2 \beta}{\partial \sigma \partial t} \right|_{t=0} = \mu' Q + \mu Q' = -\varepsilon_1 \mu \kappa_g T + \mu' Q + \varepsilon_3 \mu \tau_g n \quad (16)$$

and using (4),

$$\begin{aligned} \left. \frac{\partial^3 \beta}{\partial t \partial \sigma^2} \right|_{t=0} &= (-2\varepsilon_1 \mu' \kappa_g - \varepsilon_1 \mu \kappa_g' - \varepsilon_1 \varepsilon_3 \mu \kappa_n \tau_g) T \\ &+ (\mu'' - \varepsilon_1 \varepsilon_2 \mu \kappa_g^2 - \varepsilon_1 \varepsilon_3 \mu \tau_g^2) Q \\ &+ (2\varepsilon_3 \mu' \tau_g - \varepsilon_1 \varepsilon_3 \mu \tau_g' + \varepsilon_1 \varepsilon_3 \mu \kappa_g \kappa_n) n. \end{aligned} \quad (17)$$

Also using (14) we have

$$\begin{aligned} \left. \frac{\partial^3 \beta}{\partial \sigma^3} \right|_{t=0} &= -(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) T + (\varepsilon_2 \kappa'_g - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g) Q \\ &\quad + (\varepsilon_3 \kappa'_n + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) n \end{aligned} \quad (18)$$

and by (15)

$$\begin{aligned} \left. \frac{\partial^4 \beta}{\partial t \partial \sigma^3} \right|_{t=0} &= (-3\varepsilon_1 \mu'' \kappa_g - 3\varepsilon_1 \mu' \kappa'_g - \varepsilon_1 \mu \kappa''_g - \varepsilon_1 \varepsilon_3 \mu \kappa'_n \tau_g - 2\varepsilon_1 \varepsilon_3 \mu \kappa_n \tau'_g) \\ &\quad - 3\varepsilon_1 \varepsilon_3 \mu' \kappa_n \tau_g + \varepsilon_3 \mu \kappa_g \kappa_n^2 + \varepsilon_2 \mu \kappa_g^3 + \varepsilon_1 \varepsilon_2 \varepsilon_3 \mu \kappa_g \tau_g^2) T \\ &\quad + (\mu''' - 3\varepsilon_1 \varepsilon_2 \mu' \kappa_g^2 - 3\varepsilon_2 \varepsilon_3 \mu' \tau_g^2 - 3\varepsilon_1 \varepsilon_2 \mu \kappa_g \kappa'_g - 3\varepsilon_2 \varepsilon_3 \mu \tau_g \tau'_g) Q \\ &\quad + (-3\varepsilon_1 \varepsilon_3 \mu' \kappa_g \kappa_n - 2\varepsilon_1 \varepsilon_3 \mu \kappa'_g \kappa_n - \varepsilon_1 \varepsilon_2 \varepsilon_3 \mu \kappa_g^2 \tau_g + 3\varepsilon_3 \mu'' \tau_g \\ &\quad + 3\varepsilon_3 \mu' \tau'_g + \varepsilon_3 \mu \tau''_g - \varepsilon_1 \varepsilon_3 \mu \kappa_g \kappa'_n - \varepsilon_2 \mu \tau_g^3 - \varepsilon_1 \mu \kappa_n^2 \tau_g) n. \end{aligned} \quad (19)$$

Now, let $H(t)$ denote the functional of a relaxed elastic line of second kind for the arc $\beta(\sigma; t)$, $0 \leq \sigma \leq \lambda(t)$, $|t| < \varepsilon$. Since, in general, σ is not the arc length for $t \neq 0$ functional (2) can be calculated as follows:

$$H(t) = \int_0^{\lambda(t)} \left(\frac{\left\langle \frac{\partial \beta}{\partial \sigma} \times \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial^3 \beta}{\partial \sigma^3} \right\rangle}{\left\langle \frac{\partial \beta}{\partial \sigma}, \frac{\partial \beta}{\partial \sigma} \right\rangle \left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial^2 \beta}{\partial \sigma^2} \right\rangle - \left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial \beta}{\partial \sigma} \right\rangle^2} \right)^2 d\sigma.$$

A necessary condition for α to be an extremal is that

$$\left. \frac{dH}{dt} \right|_{t=0} = 0$$

for arbitrary μ satisfying (6). We have

$$\begin{aligned} H'(t) &= \frac{d\lambda}{dt} \left\{ \left(\frac{\left\langle \frac{\partial \beta}{\partial \sigma} \times \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial^3 \beta}{\partial \sigma^3} \right\rangle}{\left\langle \frac{\partial \beta}{\partial \sigma}, \frac{\partial \beta}{\partial \sigma} \right\rangle \left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial^2 \beta}{\partial \sigma^2} \right\rangle - \left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial \beta}{\partial \sigma} \right\rangle^2} \right)^2 \right\}_{\sigma=\lambda(t)} \\ &\quad + 2 \int_0^{\lambda(t)} \frac{\partial}{\partial t} \left(\frac{\left\langle \frac{\partial \beta}{\partial \sigma} \times \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial^3 \beta}{\partial \sigma^3} \right\rangle}{\left\langle \frac{\partial \beta}{\partial \sigma}, \frac{\partial \beta}{\partial \sigma} \right\rangle \left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial^2 \beta}{\partial \sigma^2} \right\rangle - \left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial \beta}{\partial \sigma} \right\rangle^2} \right)^2 d\sigma. \end{aligned}$$

In calculating dH/dt ; we give explicitly only terms that do not vanish for $t = 0$. The omitted terms are those with factor

$$\left\langle \frac{\partial^2 \beta}{\partial \sigma^2}, \frac{\partial \beta}{\partial \sigma} \right\rangle$$

which vanishes at $t = 0$ because $\langle T, T' \rangle = 0$. Thus, using (11 – 14) and (16 – 19) we get

$$\begin{aligned}
H'(0) &= \varepsilon_1 f^2(l) \int_0^l \mu \kappa_g ds + 2 \int_0^l \frac{f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \{ (\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) [(-3\kappa_g^2 \kappa_n' \\
&\quad - 3\varepsilon_2 \kappa_g^3 \tau_g + 3\kappa_g \kappa_n' \kappa_n - 2\varepsilon_3 \kappa_g \kappa_n^2 \tau_g + \varepsilon_2 \kappa_g^4 + \varepsilon_3 \kappa_g^2 \kappa_n^2 - \varepsilon_2 \kappa_n' \tau_g^2 + \varepsilon_2 \varepsilon_3 \kappa_g' \tau_g' \\
&\quad + 4\kappa_n \tau_g \tau_g' - \varepsilon_2 \varepsilon_3 \kappa_g \tau_g'' + \varepsilon_1 \varepsilon_2 \kappa_g \kappa_n^2 \tau_g) + \mu' (-4\kappa_g^2 \kappa_n - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g \\
&\quad + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) + \mu'' (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
&\quad + \varepsilon_3 \mu''' \kappa_n - (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) (\mu (-4\varepsilon_2 \kappa_g^3 \\
&\quad - 4\varepsilon_3 \kappa_g \kappa_n^2 + 2\kappa_g \tau_g^2 + 2\varepsilon_1 \varepsilon_3 \kappa_n \tau_g') + 4\mu' \varepsilon_1 \varepsilon_3 \kappa_n \tau_g + 2\mu'' \varepsilon_1 \kappa_g) \} ds \\
&= \int_0^l \mu (\varepsilon_1 f^2(l) \kappa_g) ds + \int_0^l \mu \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-3\kappa_g^2 \kappa_n' \\
&\quad - 3\varepsilon_2 \kappa_g^3 \tau_g + 3\kappa_g \kappa_n' \kappa_n - 2\varepsilon_3 \kappa_g \kappa_n^2 \tau_g + \varepsilon_2 \kappa_g^4 + \varepsilon_3 \kappa_g^2 \kappa_n^2 - \varepsilon_2 \kappa_n' \tau_g^2 + \varepsilon_2 \varepsilon_3 \kappa_g' \tau_g' \\
&\quad + 4\kappa_n \tau_g \tau_g' - \varepsilon_2 \varepsilon_3 \kappa_g \tau_g'' + \varepsilon_1 \varepsilon_2 \kappa_g \kappa_n^2 \tau_g) - (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
&\quad \left. + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) (-4\varepsilon_2 \kappa_g^3 - 4\varepsilon_3 \kappa_g \kappa_n^2 + 2\kappa_g \tau_g^2 + 2\varepsilon_1 \varepsilon_3 \kappa_n \tau_g') \right] \} ds \\
&\quad + \int_0^l \mu' \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-4\kappa_g^2 \kappa_n - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g \right. \\
&\quad \left. + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) - 4\varepsilon_1 \varepsilon_3 \kappa_n \tau_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
&\quad \left. + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right] \} ds \\
&\quad + \int_0^l \mu'' \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
&\quad \left. - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right] \} ds \\
&\quad + \int_0^l \mu''' \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\} ds,
\end{aligned}$$

where

$$f(s) = \frac{-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)}{\varepsilon_1 (\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)}.$$

However, using integration by parts and (6) we get

$$\begin{aligned}
& \int_0^l \mu' \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-4\kappa_g^2 \kappa_n - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g) \right. \\
& + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) - 4\varepsilon_1 \varepsilon_3 \kappa_n \tau_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
& \left. + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\} ds = \mu(l) \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-4\kappa_g^2 \kappa_n \right. \\
& - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) \\
& \left. - 4\varepsilon_1 \varepsilon_3 \kappa_n \tau_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\}_{s=l} \\
& - \int_0^l \mu \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-4\kappa_g^2 \kappa_n - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g) \right. \\
& + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) - 4\varepsilon_1 \varepsilon_3 \kappa_n \tau_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
& \left. + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\}' ds
\end{aligned}$$

and

$$\begin{aligned}
& \int_0^l \mu'' \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
& \left. - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\} ds = \\
& \mu'(l) \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
& \left. - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\}_{s=l} \\
& - \mu(l) \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
& \left. - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\}'_{s=l} \\
& + \int_0^l \mu \left\{ \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} [(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \\
& \left. - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \right\}'' ds
\end{aligned}$$

and

$$\begin{aligned}
\int_0^l \mu''' \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\} ds &= \mu''(l) \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\}_{s=l} - \mu''(0) \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\}_{s=0} \\
&+ \mu'(l) \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\}'_{s=l} + \mu(l) \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\}''_{s=l} \\
&- \int_0^l \mu \left\{ \frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right\}''' ds.
\end{aligned}$$

Thus $H'(0)$ can be written as

$$\begin{aligned}
H'(0) = & \int_0^l \mu \left\{ \varepsilon_1 \kappa_g f^2(l) + \frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \left((\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) \left[(-3\kappa_g^2 \kappa_n' \right. \right. \right. \\
& - 3\varepsilon_2 \kappa_g^3 \tau_g + 3\kappa_g \kappa_g' \kappa_n - 2\varepsilon_3 \kappa_g \kappa_n^2 \tau_g + \varepsilon_2 \kappa_g^4 + \varepsilon_3 \kappa_g^2 \kappa_n^2 - \varepsilon_2 \kappa_n' \tau_g^2 + \varepsilon_2 \varepsilon_3 \kappa_g' \tau_g' \\
& + 4\kappa_n \tau_g \tau_g' - \varepsilon_2 \varepsilon_3 \kappa_g \tau_g'' + \varepsilon_1 \varepsilon_2 \kappa_g \kappa_n^2 \tau_g) - (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
& + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \left. \left. \left. (-4\varepsilon_2 \kappa_g^3 - 4\varepsilon_3 \kappa_g \kappa_n^2 + 2\kappa_g \tau_g^2 + 2\varepsilon_1 \varepsilon_3 \kappa_n \tau_g') \right] \right) \right. \\
& - \left[\frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \left[(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-4\kappa_g^2 \kappa_n - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g \right. \right. \\
& + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) - 4\varepsilon_1 \varepsilon_2 \kappa_n \tau_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
& + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \left. \left. \right] \right]' + \left[\frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \left[(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' \right. \right. \\
& - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \left. \left. \right] \right]'' \\
& - \left[\frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right]'''' \left. \right\} ds \\
& + \mu(l) \left\{ \frac{2f(l)}{(\varepsilon_2 \kappa_g^2(l) + \varepsilon_3 \kappa_n^2(l))^2} \left[(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-4\kappa_g^2 \kappa_n - \varepsilon_1 \kappa_n^3 + 2\varepsilon_2 \varepsilon_3 \kappa_g' \tau_g \right. \right. \\
& + 2\varepsilon_2 \kappa_n \tau_g^2 - 3\varepsilon_2 \varepsilon_3 \kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\varepsilon_2 \kappa_n \tau_g^2) - 4\varepsilon_1 \varepsilon_2 \kappa_n \tau_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
& + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \left. \left. \right] \right]_{s=l} - \left(\frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \left[(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) \right. \right. \\
& x (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \\
& + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \left. \left. \right] \right)'_{s=l} + \left(\frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right)''_{s=l} \left. \right\} \\
& + \mu'(l) \left\{ \left(\frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \left[(\varepsilon_1 \varepsilon_2 \kappa_g^2 + \varepsilon_1 \varepsilon_3 \kappa_n^2) (-\varepsilon_3 \kappa_n' - 4\varepsilon_2 \varepsilon_3 \kappa_g \tau_g) \right. \right. \right. \\
& - 2\varepsilon_1 \kappa_g (-\varepsilon_2 \kappa_g (\varepsilon_3 \kappa_n' + \varepsilon_2 \varepsilon_3 \kappa_g \tau_g) + \varepsilon_3 \kappa_n (\varepsilon_2 \kappa_g' - \varepsilon_2 \varepsilon_3 \kappa_n \tau_g)) \left. \left. \right] \right)_{s=l} \\
& - \left(\frac{2\varepsilon_3 \kappa_n f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right)'_{s=l} \left. \right\} + \mu''(l) \left(\frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right)_{s=l} \\
& - \mu''(0) \left(\frac{2f(s)}{(\varepsilon_2 \kappa_g^2 + \varepsilon_3 \kappa_n^2)^2} \right)_{s=0}.
\end{aligned}$$

2.1 Intrinsic equations for a generalized elastic line on a spacelike surface

On a spacelike surface n is timelike. So, we have T and Q are spacelike and $\varepsilon_1 = \langle T, T \rangle = 1$, $\varepsilon_2 = \langle Q, Q \rangle = 1$, $\varepsilon_3 = \langle n, n \rangle = -1$. Hence $H'(0)$ can be written as

$$\begin{aligned}
H'(0) = & \int_0^l \mu \left\{ \kappa_g f^2(l) + \frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} ((\kappa_g^2 - \kappa_n^2) [(-3\kappa_g^2 \kappa_n' \right. \\
& - 3\kappa_g^3 \tau_g + 3\kappa_g \kappa_n' \kappa_n + 2\kappa_g \kappa_n^2 \tau_g + \kappa_g^4 - \kappa_g^2 \kappa_n^2 - \kappa_n' \tau_g^2 - \kappa_g' \tau_g' \\
& + 4\kappa_n \tau_g \tau_g' + \kappa_g \tau_g'' + \kappa_g \kappa_n^2 \tau_g) - (\kappa_g (\kappa_n' + \kappa_g \tau_g) \\
& - \kappa_n (\kappa_g' + \kappa_n \tau_g)) (-4\kappa_g^3 + 4\kappa_g \kappa_n^2 + 2\kappa_g \tau_g^2 - \kappa_n \tau_g')] \\
& - \left[\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2 \kappa_n - \kappa_n^3 - 2\kappa_g' \tau_g \right. \\
& + 2\kappa_n \tau_g^2 + 3\kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\kappa_n \tau_g^2) - 4\kappa_n \tau_g (\kappa_g (\kappa_n' + \kappa_g \tau_g) \\
& - \kappa_n (\kappa_g' + \kappa_n \tau_g))] \Big] + \left[\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (\kappa_n' \right. \\
& + 4\kappa_g \tau_g) - 2\kappa_g (\kappa_g (\kappa_n' + \kappa_g \tau_g) - \kappa_n (\kappa_g' + \kappa_n \tau_g))] \Big]'' \\
& + \left. \left[\frac{2\kappa_n f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right]''' \right\} ds \\
& + \mu(l) \left\{ \frac{2f(l)}{(\kappa_g^2(l) - \kappa_n^2(l))^2} [(\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2 \kappa_n - \kappa_n^3 - 2\kappa_g' \tau_g \right. \\
& + 2\kappa_n \tau_g^2 + 3\kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\kappa_n \tau_g^2) - 4\kappa_n \tau_g (\kappa_g (\kappa_n' + \kappa_g \tau_g) \\
& - \kappa_n (\kappa_g' + \kappa_n \tau_g))] \Big]_{s=l} - \left(\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (\kappa_n' + 4\kappa_g \tau_g) \right. \\
& - 2\kappa_g (\kappa_g (\kappa_n' + \kappa_g \tau_g) - \kappa_n (\kappa_g' + \kappa_n \tau_g))] \Big]_{s=l}' - \left. \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right)'' \right)_{s=l} \\
& + \mu'(l) \left\{ \left(\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (\kappa_n' + 4\kappa_g \tau_g) \right. \right. \\
& - 2\kappa_g (\kappa_g (\kappa_n' + \kappa_g \tau_g) - \kappa_n (\kappa_g' + \kappa_n \tau_g))] \Big]_{s=l} \\
& + \left. \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right)' \right)_{s=l} \Big\} + \mu''(l) \left(\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right)_{s=l} \\
& - \mu''(0) \left(\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right)_{s=0} .
\end{aligned}$$

In order that $H'(0) = 0$ for all choices of the function $\mu(s)$ satisfying (6), with arbitrary values of $\mu(l)$, $\mu'(l)$ and $\mu''(l)$, spacelike arc α must satisfy four boundary conditions

$$\begin{aligned} & \frac{2f(l)}{(\kappa_g^2(l) - \kappa_n^2(l))^2} [(\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2\kappa_n - \kappa_n^3 - 2\kappa_g'\tau_g \\ & + 2\kappa_n\tau_g^2 + 3\kappa_g\tau_g' + 3\kappa_g^2\kappa_n + 3\kappa_n\tau_g^2) - 4\kappa_n\tau_g (\kappa_g (\kappa_n' + \kappa_g\tau_g) \\ & - \kappa_n (\kappa_g' + \kappa_n\tau_g))]_{s=l} - \left(\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (\kappa_n' + 4\kappa_g\tau_g) \right. \\ & \left. - 2\kappa_g (\kappa_g (\kappa_n' + \kappa_g\tau_g) - \kappa_n (\kappa_g' + \kappa_n\tau_g))] \right)'_{s=l} - \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right)'_{s=l} = 0, \end{aligned}$$

$$\begin{aligned} & \left(\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (\kappa_n' + 4\kappa_g\tau_g) \right. \\ & \left. - 2\kappa_g (\kappa_g (\kappa_n' + \kappa_g\tau_g) - \kappa_n (\kappa_g' + \kappa_n\tau_g))] \right)'_{s=l} \\ & + \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right)'_{s=l} = 0, \end{aligned}$$

$$f(l) = 0,$$

$$f(0) = 0,$$

and differential equation

$$\begin{aligned} & \kappa_g f^2(l) + \frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} ((\kappa_g^2 - \kappa_n^2) [(-3\kappa_g^2\kappa_n' - 3\kappa_g^3\tau_g \\ & + 3\kappa_g\kappa_g'\kappa_n + 2\kappa_g\kappa_n^2\tau_g + \kappa_g^4 - \kappa_g^2\kappa_n^2 - \kappa_n'\tau_g^2 - \kappa_g'\tau_g' \\ & + 4\kappa_n\tau_g\tau_g' + \kappa_g\tau_g'' + \kappa_g\kappa_n^2\tau_g) - (\kappa_g (\kappa_n' + \kappa_g\tau_g) \\ & - \kappa_n (\kappa_g' + \kappa_n\tau_g)) (-4\kappa_g^3 + 4\kappa_g\kappa_n^2 + 2\kappa_g\tau_g^2 - \kappa_n\tau_g')] \\ & - \left[\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2\kappa_n - \kappa_n^3 - 2\kappa_g'\tau_g \right. \\ & \left. + 2\kappa_n\tau_g^2 + 3\kappa_g\tau_g' + 3\kappa_g^2\kappa_n + 3\kappa_n\tau_g^2) - 4\kappa_n\tau_g (\kappa_g (\kappa_n' + \kappa_g\tau_g) \right. \\ & \left. - \kappa_n (\kappa_g' + \kappa_n\tau_g))] \right]' + \left[\frac{2f(s)}{(\kappa_g^2 - \kappa_n^2)^2} [(\kappa_g^2 - \kappa_n^2) (\kappa_n' \right. \\ & \left. + 4\kappa_g\tau_g) - 2\kappa_g (\kappa_g (\kappa_n' + \kappa_g\tau_g) - \kappa_n (\kappa_g' + \kappa_n\tau_g))] \right]' \\ & + \left[\frac{2\kappa_n f(s)}{(\kappa_g^2 - \kappa_n^2)^2} \right]''' = 0, \end{aligned}$$

where

$$f(s) = \frac{\kappa_g (\kappa'_n + \kappa_g \tau_g) - \kappa_n (\kappa'_g + \kappa_n \tau_g)}{\kappa_g^2 - \kappa_n^2}.$$

2.2 Intrinsic equations for a generalized elastic line on a timelike surface for timelike arc α

Since α is timelike T is timelike. So, we have Q and n are spacelike and $\varepsilon_1 = \langle T, T \rangle = -1$, $\varepsilon_2 = \langle Q, Q \rangle = 1$, $\varepsilon_3 = \langle n, n \rangle = 1$. Hence $H'(0)$ can be written as

$$\begin{aligned}
H'(0) = & \int_0^l \mu \left\{ -\kappa_g f^2(l) + \frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \left((-\kappa_g^2 - \kappa_n^2) \left[(-3\kappa_g^2 \kappa_n' \right. \right. \right. \\
& - 3\kappa_g^3 \tau_g + 3\kappa_g \kappa_g' \kappa_n - 2\kappa_g \kappa_n^2 \tau_g + \kappa_g^4 + \kappa_g^2 \kappa_n^2 - \kappa_n' \tau_g^2 + \kappa_g' \tau_g' \\
& + 4\kappa_n \tau_g \tau_g' - \kappa_g \tau_g'' - \kappa_g \kappa_n^2 \tau_g) - (-\kappa_g (\kappa_n' + \kappa_g \tau_g) \\
& + \kappa_n (\kappa_g' - \kappa_n \tau_g)) \left. \left. \left. (-4\kappa_g^3 - 4\kappa_g \kappa_n^2 + 2\kappa_g \tau_g^2 - 2\kappa_n \tau_g') \right] \right) \right. \\
& - \left[\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2 \kappa_n + \kappa_n^3 + 2\kappa_g' \tau_g \right. \right. \\
& + 2\kappa_n \tau_g^2 - 3\kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\kappa_n \tau_g^2) + 4\kappa_n \tau_g (-\kappa_g (\kappa_n' + \kappa_g \tau_g) \\
& + \kappa_n (\kappa_g' - \kappa_n \tau_g)) \left. \left. \right] \right]' + \left[\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 - \kappa_n^2) (-\kappa_n' \right. \right. \\
& - 4\kappa_g \tau_g) + 2\kappa_g (-\kappa_g (\kappa_n' + \kappa_g \tau_g) + \kappa_n (\kappa_g' - \kappa_n \tau_g)) \left. \left. \right] \right]'' \\
& - \left. \left[\frac{2\kappa_n f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right]''' \right\} ds \\
& + \mu(l) \left\{ \frac{2f(l)}{(\kappa_g^2(l) + \kappa_n^2(l))^2} \left[(-\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2 \kappa_n + \kappa_n^3 + 2\kappa_g' \tau_g \right. \right. \right. \\
& + 2\kappa_n \tau_g^2 - 3\kappa_g \tau_g' + 3\kappa_g^2 \kappa_n + 3\kappa_n \tau_g^2) + 4\kappa_n \tau_g (-\kappa_g (\kappa_n' + \kappa_g \tau_g) \\
& + \kappa_n (\kappa_g' - \kappa_n \tau_g)) \left. \left. \right] \right\}_{s=l} - \left(\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 - \kappa_n^2) (-\kappa_n' - 4\kappa_g \tau_g) \right. \right. \\
& + 2\kappa_g (-\kappa_g (\kappa_n' + \kappa_g \tau_g) + \kappa_n (\kappa_g' - \kappa_n \tau_g)) \left. \left. \right] \right)'_{s=l} + \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right)''_{s=l} \\
& + \mu'(l) \left\{ \left(\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 - \kappa_n^2) (-\kappa_n' - 4\kappa_g \tau_g) \right. \right. \right. \\
& + \kappa_g (-\kappa_g (\kappa_n' + \kappa_g \tau_g) + \kappa_n (\kappa_g' - \kappa_n \tau_g)) \left. \left. \right] \right) \right\}_{s=l} \\
& - \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right)'_{s=l} \left. \right\} + \mu''(l) \left(\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right)_{s=l} \\
& - \mu''(0) \left(\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right)_{s=0}.
\end{aligned}$$

In order that $H'(0) = 0$ for all choices of the function $\mu(s)$ satisfying (6), with arbitrary values of $\mu(l)$, $\mu'(l)$ and $\mu''(l)$, spacelike arc α must satisfy four boundary conditions

$$\begin{aligned}
& \frac{2f(l)}{(\kappa_g^2(l) + \kappa_n^2(l))^2} [(-\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2\kappa_n + \kappa_n^3 + 2\kappa_g'\tau_g \\
& + 2\kappa_n\tau_g^2 - 3\kappa_g\tau_g' + 3\kappa_g^2\kappa_n + 3\kappa_n\tau_g^2) + 4\kappa_n\tau_g (-\kappa_g (\kappa_n' + \kappa_g\tau_g) \\
& + \kappa_n (\kappa_g' - \kappa_n\tau_g))]_{s=l} - \left(\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} [(-\kappa_g^2 - \kappa_n^2) (-\kappa_n' - 4\kappa_g\tau_g) \right. \\
& \left. + 2\kappa_g (-\kappa_g (\kappa_n' + \kappa_g\tau_g) + \kappa_n (\kappa_g' - \kappa_n\tau_g))] \right)'_{s=l} + \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right)'_{s=l} = 0, \\
& \left(\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} [(-\kappa_g^2 - \kappa_n^2) (-\kappa_n' - 4\kappa_g\tau_g) + \kappa_g (-\kappa_g (\kappa_n' + \kappa_g\tau_g) + \kappa_n (\kappa_g' - \kappa_n\tau_g))] \right)'_{s=l} \\
& - \left(\frac{2\kappa_n f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right)'_{s=l} = 0, \\
& f(l) = 0, \\
& f(0) = 0,
\end{aligned}$$

and differential equation

$$\begin{aligned}
& -\kappa_g f^2(l) + \frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} ((-\kappa_g^2 - \kappa_n^2) [(-3\kappa_g^2\kappa_n' \\
& - 3\kappa_g^3\tau_g + 3\kappa_g\kappa_g'\kappa_n - 2\kappa_g\kappa_n^2\tau_g + \kappa_g^4 + \kappa_g^2\kappa_n^2 - \kappa_n'\tau_g^2 + \kappa_g'\tau_g' \\
& + 4\kappa_n\tau_g\tau_g' - \kappa_g\tau_g'' - \kappa_g\kappa_n^2\tau_g) - (-\kappa_g (\kappa_n' + \kappa_g\tau_g) \\
& + \kappa_n (\kappa_g' - \kappa_n\tau_g)) (-4\kappa_g^3 - 4\kappa_g\kappa_n^2 + 2\kappa_g\tau_g^2 - 2\kappa_n\tau_g')] \\
& - \left[\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} [(-\kappa_g^2 - \kappa_n^2) (-4\kappa_g^2\kappa_n + \kappa_n^3 + 2\kappa_g'\tau_g \right. \\
& \left. + 2\kappa_n\tau_g^2 - 3\kappa_g\tau_g' + 3\kappa_g^2\kappa_n + 3\kappa_n\tau_g^2) + 4\kappa_n\tau_g (-\kappa_g (\kappa_n' + \kappa_g\tau_g) \right. \\
& \left. + \kappa_n (\kappa_g' - \kappa_n\tau_g))] \right]' + \left[\frac{2f(s)}{(\kappa_g^2 + \kappa_n^2)^2} [(-\kappa_g^2 - \kappa_n^2) (-\kappa_n' \right. \\
& \left. - 4\kappa_g\tau_g) + 2\kappa_g (-\kappa_g (\kappa_n' + \kappa_g\tau_g) + \kappa_n (\kappa_g' - \kappa_n\tau_g))] \right]' \\
& - \left[\frac{2\kappa_n f(s)}{(\kappa_g^2 + \kappa_n^2)^2} \right]' = 0,
\end{aligned}$$

where

$$f(s) = \frac{\kappa_g (\kappa_n' + \kappa_g\tau_g) - \kappa_n (\kappa_g' - \kappa_n\tau_g)}{\kappa_g^2 + \kappa_n^2}.$$

2.3 Intrinsic equations for a generalized elastic line on a timelike surface for spacelike arc α

Now Q is timelike, T and n are spacelike. So we have $\varepsilon_1 = \langle T, T \rangle = 1$, $\varepsilon_2 = \langle Q, Q \rangle = -1$, $\varepsilon_3 = \langle n, n \rangle = 1$. Hence $H'(0)$ can be written as

$$\begin{aligned}
H'(0) = & \int_0^l \mu \left\{ \kappa_g f^2(l) + \frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \left((-\kappa_g^2 + \kappa_n^2) \left[(-3\kappa_g^2 \kappa_n' \right. \right. \right. \\
& + 3\kappa_g^3 \tau_g + 3\kappa_g \kappa_n' \kappa_n - 2\kappa_g \kappa_n^2 \tau_g - \kappa_g^4 + \kappa_g^2 \kappa_n^2 + \kappa_n' \tau_g^2 - \kappa_g' \tau_g' \\
& + 4\kappa_n \tau_g \tau_g' + \kappa_g \tau_g'' - \kappa_g \kappa_n^2 \tau_g) - (\kappa_g (\kappa_n' - \kappa_g \tau_g) \\
& \left. \left. \left. + \kappa_n (-\kappa_g' + \kappa_n \tau_g) \right) \right] (4\kappa_g^3 - 4\kappa_g \kappa_n^2 + 2\kappa_g \tau_g^2 + 2\kappa_n \tau_g') \right) \\
& - \left[\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 + \kappa_n^2) (-4\kappa_g^2 \kappa_n - \kappa_n^3 - 2\kappa_g' \tau_g \right. \right. \\
& - 2\kappa_n \tau_g^2 + 3\kappa_g \tau_g' + 3\kappa_g^2 \kappa_n - 3\kappa_n \tau_g^2) + 4\kappa_n \tau_g (\kappa_g (\kappa_n' - \kappa_g \tau_g) \\
& \left. \left. \left. + \kappa_n (-\kappa_g' + \kappa_n \tau_g) \right) \right] \right]' + \left[\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 + \kappa_n^2) (-\kappa_n' \right. \right. \\
& + 4\kappa_g \tau_g) - 2\kappa_g (\kappa_g (\kappa_n' - \kappa_g \tau_g) + \kappa_n (-\kappa_g' + \kappa_n \tau_g)) \right] \left. \right]'' \\
& - \left[\frac{2\kappa_n f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right]'' \left. \right\} ds \\
& + \mu(l) \left\{ \frac{2f(l)}{(-\kappa_g^2(l) + \kappa_n^2(l))^2} \left[(-\kappa_g^2 + \kappa_n^2) (-4\kappa_g^2 \kappa_n - \kappa_n^3 - 2\kappa_g' \tau_g \right. \right. \\
& - 2\kappa_n \tau_g^2 + 3\kappa_g \tau_g' + 3\kappa_g^2 \kappa_n - 3\kappa_n \tau_g^2) + 4\kappa_n \tau_g (\kappa_g (\kappa_n' - \kappa_g \tau_g) \\
& \left. \left. \left. + \kappa_n (-\kappa_g' + \kappa_n \tau_g) \right) \right]_{s=l} - \left(\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 + \kappa_n^2) (-\kappa_n' + 4\kappa_g \tau_g) \right. \right. \right. \\
& - 2\kappa_g (\kappa_g (\kappa_n' - \kappa_g \tau_g) \\
& \left. \left. \left. + \kappa_n (-\kappa_g' + \kappa_n \tau_g) \right) \right] \right)'_{s=l} + \left(\frac{2\kappa_n f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right)''_{s=l} \left. \right\} \\
& + \mu'(l) \left\{ \left(\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \left[(-\kappa_g^2 + \kappa_n^2) (-\kappa_n' + 4\kappa_g \tau_g) \right. \right. \right. \\
& - 2\kappa_g (\kappa_g (\kappa_n' - \kappa_g \tau_g) + \kappa_n (-\kappa_g' + \kappa_n \tau_g)) \left. \left. \left. \right] \right)_{s=l} \right. \\
& - \left(\frac{2\kappa_n f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right)'_{s=l} \left. \right\} + \mu''(l) \left(\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right)_{s=l} \\
& - \mu''(0) \left(\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right)_{s=0}.
\end{aligned}$$

In order that $H'(0) = 0$ for all choices of the function $\mu(s)$ satisfying (6), with arbitrary values of $\mu(l)$, $\mu'(l)$ and $\mu''(l)$, spacelike arc α must satisfy four boundary conditions

$$\begin{aligned} & \frac{2f(l)}{(-\kappa_g^2(l) + \kappa_n^2(l))^2} [(-\kappa_g^2 + \kappa_n^2) (-4\kappa_g^2\kappa_n - \kappa_n^3 - 2\kappa_g'\tau_g \\ & - 2\kappa_n\tau_g^2 + 3\kappa_g\tau_g' + 3\kappa_g^2\kappa_n - 3\kappa_n\tau_g^2) + 4\kappa_n\tau_g (\kappa_g (\kappa_n' - \kappa_g\tau_g) \\ & + \kappa_n (-\kappa_g' + \kappa_n\tau_g))]_{s=l} - \left(\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} [(-\kappa_g^2 + \kappa_n^2) (-\kappa_n' + 4\kappa_g\tau_g) \right. \\ & \left. - 2\kappa_g (\kappa_g (\kappa_n' - \kappa_g\tau_g) \right. \\ & \left. + \kappa_n (-\kappa_g' + \kappa_n\tau_g))]_{s=l} \right)' + \left(\frac{2\kappa_n f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right)''_{s=l} = 0, \end{aligned}$$

$$\begin{aligned} & \left(\frac{2f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} [(-\kappa_g^2 + \kappa_n^2) (-\kappa_n' + 4\kappa_g\tau_g) \right. \\ & \left. - 2\kappa_g (\kappa_g (\kappa_n' - \kappa_g\tau_g) + \kappa_n (-\kappa_g' + \kappa_n\tau_g))]_{s=l} \right. \\ & \left. - \left(\frac{2\kappa_n f(s)}{(-\kappa_g^2 + \kappa_n^2)^2} \right)'_{s=l} \right) = 0, \end{aligned}$$

$$f(l) = 0,$$

$$f(0) = 0,$$

where

$$f(s) = \frac{\kappa_g (\kappa_n' - \kappa_g\tau_g) - \kappa_n (\kappa_g' + \kappa_n\tau_g)}{-\kappa_g^2 + \kappa_n^2}.$$

References

- [1] Capovilla, R., Chryssomalakos, C. and Guven, J. Hamiltonians for curves, J. Phys. A: Math. Gen., 35, 6571–6587, (2002).
- [2] Uğurlu, H. and Kocayigit, H. The Frenet and Darboux instantaneous rotation vectors of curves on time-like surface, Mathematical & Computational Appl. Vol: 1, No: 2, 131-141 (1996).
- [3] O'Neill, B. Semi-Riemannian Geometry, London, Academic Press. New York 1983.

- [4] Tutar, A. and Sarıođlugil, A. Relaxed elastic lines of second kind on oriented surface in Minkowski space, Applied Mathematics and Mechanics (English Edition), Vol. 27 (11), 2006, 1481-1489.
- [5] Ekici, C. and G6rg6l6, A. Intrinsic equations for a generalized relaxed elastic line on an oriented surface in the Minkowski 3-space E_1^3 , Turk J. Math, 33 (2009) , 397 – 407.