

A curious relation between the flat cosmological model and the elliptic integral of the first kind (Research Note)

A. Mészáros¹ and J. Řípa²

¹ Charles University in Prague, Faculty of Mathematics and Physics, Astronomical Institute, V Holešovičkách 2, CZ 180 00 Prague 8, Czech Republic
e-mail: meszaros@cesnet.cz

² Institute of Basic Science, Natural Sciences Campus, Sungkyunkwan University Engineering Building 2, 2066 Seobu-ro, Jangan-gu, Suwon, Gyeonggi-do, 440-746, Korea
e-mail: ripa.jakub@gmail.com

Received June 14, 2013; accepted June 16, 2013

ABSTRACT

Context. The dependence of the luminosity distance on the redshift has a key importance in the cosmology. This dependence can well be given by standard functions for the zero cosmological constant.

Aims. The purpose of this article is to present such a relation also for the non-zero cosmological constant, if the universe is spatially flat.

Methods. A definite integral is used.

Results. The integration ends in the elliptic integral of the first kind.

Conclusions. The result shows that no numerical integration is needed for the non-zero cosmological constant, if the universe is spatially flat.

Key words. cosmology: theory

1. Introduction

The dependence of the luminosity distance $d_L(z)$ on the redshift z is a key formula in cosmology (Carroll et al. 1992). It is given by three independent cosmological parameters: by two omega-parameters Ω_M , Ω_Λ and by the Hubble constant H_0 . Its relation to the so called "proper-motion distance" is given by $d_{PM}(z)(1+z) = d_L(z)$ (Weinberg 1972).

One has (Carroll et al. 1992):

$$d_{PM}(z) = \frac{c}{H_0 \sqrt{|\Omega_k|}} \times$$

$$\text{sinn} \left\{ \sqrt{|\Omega_k|} \int_0^z \frac{dz'}{\sqrt{(1+z')^2(1+\Omega_M z') - z'(2+z')\Omega_\Lambda}} \right\}. \quad (1)$$

In this equation c is the speed of light in vacuum, and it holds $\Omega_k + \Omega_M + \Omega_\Lambda = 1$. The notation "sinn" means the standard function sinh for $\Omega_k > 0$, and sin for $\Omega_k < 0$, respectively. If $\Omega_k = 0$, then one simply has an integration:

$$\frac{H_0 d_{PM}(z)}{c} = \int_0^z \frac{dz'}{\sqrt{(1+z')^2(1+\Omega_M z') - z'(2+z')\Omega_\Lambda}}.$$

Send offprint requests to: A. Mészáros

(2)

In addition, from the physical point of view, in both equations it must be $\Omega_M > 0$. (The case $\Omega_M = 0$ can serve as a limit, but $\Omega_M < 0$ is fully unphysical.) On the other hand, Ω_Λ can have both signs, but the observations of the last two decades strongly disfavour negative values (see, e.g., Perrett et al. (2012) and the references therein).

In the special case of $\Omega_\Lambda = 0$ the integral in Eq.(1) can be given by the so-called Mattig-formula (Mattig 1958) for any $\Omega_M > 0$. The formula can then well be used in cosmological applications (see, e.g., Mészáros (2002)). For $\Omega_\Lambda \neq 0$ the integral in Eqs.(1,2) is usually solved numerically¹.

In this note we show that the integral on the right-hand-side of Eq.(2) can be solved analytically also for $\Omega_\Lambda \neq 0$.

2. The integration

We rewrite the right-hand side of Eq.(2) into the form

$$\frac{H_0 d_{PM}(z_0)}{c} = I(\Omega_M, z_0) = \int_0^{z_0} \frac{dz}{\sqrt{(1+z)^3 \Omega_M + \Omega_\Lambda}}, \quad (3)$$

where $z \geq 0$, $\Omega_M > 0$ and $\Omega_M + \Omega_\Lambda = 1$. Introducing the substitution $x = (1+z)^{-1}$, we obtain

$$I(\Omega_M, x_0) = \int_{x_0}^1 \frac{dx}{\sqrt{x} \sqrt{\Omega_M + \Omega_\Lambda x^3}}, \quad (4)$$

¹ See, cf., <http://www.astro.ucla.edu/~wright/cosmolog.htm>

where $0 < x_0 = (1 + z_0)^{-1} \leq 1$.

In this section we consider only the case $\Omega_M < 1$, i.e. $\Omega_\Lambda = 1 - \Omega_M > 0$. Introducing a further substitution $y = (\Omega_\Lambda/\Omega_M)^{1/3}x$, we obtain

$$I(\Omega_M, x_0) = \frac{1}{\Omega_M^{1/3}\Omega_\Lambda^{1/6}} \int_{y_1}^{y_2} \frac{dy}{\sqrt{y}\sqrt{1+y^3}}, \quad (5)$$

where $y_1 = (\Omega_\Lambda/\Omega_M)^{1/3}x_0$ and $y_2 = (\Omega_\Lambda/\Omega_M)^{1/3}$. The two limits are non-negative with $y_1 \leq y_2$.

We try to find the primitive function for

$$\int \frac{dy}{\sqrt{y}\sqrt{1+y^3}}, \quad (6)$$

where $y \geq 0$. The following substitution helps:

$$\alpha = \arccos\left(\frac{1 + (1 - \sqrt{3})y}{1 + (1 + \sqrt{3})y}\right). \quad (7)$$

There is a one-to-one correspondence between α and y . For $y = 0$ one has $\alpha = 0$, and for $y \rightarrow \infty$ one has $\alpha_\infty \rightarrow \arccos((1 - \sqrt{3})/(1 + \sqrt{3}))$, i.e. $\alpha_\infty = 105.54^\circ$. If y increases in the interval $[0, \infty)$, α is also increasing in the interval $[0, \alpha_\infty)$. Hence, this substitution is well-defined. Conversely, one obtains

$$y = \frac{1 - \cos \alpha}{(\sqrt{3} - 1) + (\sqrt{3} + 1) \cos \alpha} \quad (8)$$

and

$$dy = \frac{2\sqrt{3} \sin \alpha \, d\alpha}{[\sqrt{3}(1 + \cos \alpha) - (1 - \cos \alpha)]^2}. \quad (9)$$

Using these two formulas, we curiously obtain

$$\frac{dy}{\sqrt{y}\sqrt{1+y^3}} = \frac{d\alpha}{3^{1/4}\sqrt{1 - \frac{2+\sqrt{3}}{4}\sin^2 \alpha}}. \quad (10)$$

The right-hand side of Eq.(10) is the function in the elliptic integral of the first kind (Gradshteyn et al. 2007) with

$$m = \frac{2 + \sqrt{3}}{4}, \quad (11)$$

where $0 < m < 1$, as it should be in an elliptic integral.

In order to calculate the definite integral

$$\int_{y_1}^{y_2} \frac{dy}{\sqrt{y}\sqrt{1+y^3}} \quad (12)$$

in Eq.(5) for non-negative $y_1 \leq y_2$ one can write

$$\int_{y_1}^{y_2} \frac{dy}{\sqrt{y}\sqrt{1+y^3}} = \int_0^{y_2} \frac{dy}{\sqrt{y}\sqrt{1+y^3}} - \int_0^{y_1} \frac{dy}{\sqrt{y}\sqrt{1+y^3}}. \quad (13)$$

After this one should use the formula with α from Eq.(10) and determine the integration limits in variable α . The substitution from Eq.(7) gives for $y = 0$ the value $\alpha = 0$. This means that - using α - the lower limits in both definite integrals are zeros. The upper limits from y_1 and y_2 , respectively, are also calculable analytically and unambiguously

from Eq.(7) via the arccos function. One obtains α_1 and α_2 , respectively, as upper limits in the integrals. It must be $\alpha_2 > \alpha_1$. One should only precise that, of course, if it were α_2 from the interval $(\pi/2, \alpha_\infty)$, then the first elliptic integral itself should be given by a sum of two integrals: In one integral the limits should be 0 and $\pi/2$, and in the second one the limits should be $(\pi - \alpha_2)$ and $\pi/2$. Both integrals must give positive values. If it were also $\alpha_1 > \pi/2$, then one should proceed similarly in the second integral, too. In any case, the integral $I(\Omega_M, x_0)$ is well obtainable from standard elliptic integrals of the first kind.

3. Remarks

Integral in Eq.(5) is presented by Gradshteyn et al. (2007) (formula 3.166.22). It should also be noted that in Carroll et al. (1992) it is said that the integral of Eq.(2) can also be solved analytically. In Paál et al. (1992) similar efforts are done. But, on the other hand, we did not find in the literature any note about this non-numerical integration of Eq.(2) using the elliptic integrals. Therefore, we mean that the substitution given by Eqs.(7,8) is new and original.

For the sake of completeness it should still be added that integral of Eq.(4) can be solved also for $\Omega_\Lambda < 0$ and $\Omega_M > 1$. One obtains (up to a constant) the formula $dy/\sqrt{y(1-y^3)}$, which is also integrable (see the formula 3.166.23 of Gradshteyn et al. (2007)).

4. Conclusion

We have proven that the integral on the right-hand-side of Eq.(2) can be solved also analytically using the elliptic integral of the first kind.

Acknowledgements. We wish to thank M. Křížek for the useful discussions and comments on the manuscript. This study was supported by the OTKA Grant K77795, by the Grant Agency of the Czech Republic Grant P209/10/0734, by the Research Program MSM0021620860 of the Ministry of Education of the Czech Republic, and by Creative Research Initiatives (RCMST) of MEST/NRF.

References

- Carroll, S. M., Press, W. H., & Turner, E. L. 1992, ARA&A, 30, 499
- Gradshteyn, I. S., Ryzhik, I. M., Jeffrey, A., & Zwillinger, D. 2007, Table of Integrals, Series, and Products (Elsevier Academic Press)
- Mattig, W. 1958, Astronomische Nachrichten, 284, 109
- Mészáros, A. 2002, ApJ, 580, 12
- Paál, G., Horváth, I., & Lukács, B. 1992, Ap&SS, 191, 107
- Perrett, K., Sullivan, M., Conley, A., et al. 2012, AJ, 144, 59
- Weinberg, S. 1972, Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity (Wiley)