

NEW EXAMPLES OF NON-SYMMETRIC EINSTEIN SOLVMANIFOLDS OF NEGATIVE RICCI CURVATURE

MEGAN M. KERR

ABSTRACT. We obtain new examples of non-symmetric Einstein solvmanifolds by combining two techniques. In [T2], H. Tamaru constructs new *attached* solvmanifolds, which are submanifolds of the solvmanifolds corresponding to noncompact symmetric spaces, endowed with a natural metric. Extending this construction, we apply it to *associated* solvmanifolds, described in [GK], obtained by modifying the algebraic structure of the solvable Lie algebras corresponding to noncompact symmetric spaces. Our new examples are Einstein solvmanifolds with nilradicals of high nilpotency, which are geometrically distinct from noncompact symmetric spaces and their submanifolds.

1. INTRODUCTION

In this paper we construct new examples of solvmanifolds of constant negative Ricci curvature. These spaces provide many explicit new examples of homogeneous Einstein manifolds that are neither the solvmanifolds corresponding to noncompact symmetric spaces, nor their submanifolds. We obtain our examples by extending the method of H. Tamaru in [T2], in which, via parabolic subalgebras of semisimple Lie algebras, he builds solvable subalgebras by choosing a subset Λ' of the set Λ of simple roots. Tamaru proves that the solvable subalgebra of the restricted root system, $\mathfrak{g}_{\Lambda'}$, given a natural inner product, called an *attached* solvmanifold, is in fact an Einstein solvmanifold. We combine this with a method introduced by C. S. Gordon and the author in [GK] to construct Einstein solvmanifolds which are *associated* (but not isometric) to the Einstein solvable Lie groups corresponding to higher rank, irreducible symmetric spaces of noncompact type.

Tamaru's construction yields solvmanifolds which are naturally homogeneous submanifolds of symmetric spaces of noncompact type; however, they generally are *not* totally geodesic subalgebras. When Tamaru's method is extended to construct the Einstein solvmanifolds here, which are *attached* to *associated* solvmanifolds, we get completely new examples. We show, by isometry groups, that they cannot be submanifolds of symmetric spaces. Furthermore, we show that while our examples have constant negative Ricci curvature, they admit two-planes of positive sectional curvature.

The category of solvmanifolds is a natural place to look for Einstein manifolds of negative Ricci curvature. All known examples of homogeneous Einstein manifolds of negative Ricci curvature are Riemannian solvmanifolds. Indeed, in 1975 Alekseevskii conjectured that every noncompact homogeneous Einstein manifold is a solvmanifold [Al]. There has been much recent progress toward determining existence of Einstein manifolds of negative Ricci curvature, and more generally, Ricci solitons (e.g. [J1, J2, J3, L1, L2, L3, LW, N, P]).

It is a pleasure to thank Carolyn Gordon for helpful discussions.

Date: October 18, 2018.

2000 Mathematics Subject Classification. Primary: 53C30.

2. PRELIMINARIES

A solvable Lie group S together with a left-invariant metric g is called a *Riemannian solvmanifold*. A *metric Lie algebra* is a Lie algebra endowed with an inner product. The left-invariant metric g on a solvmanifold S defines an inner product $\langle \cdot, \cdot \rangle$ on the Lie algebra \mathfrak{s} of S . Conversely, an inner product on a Lie algebra \mathfrak{s} defines a unique left-invariant metric on S , the corresponding simply connected solvable Lie group. Given the one-to-one correspondence between simply connected Riemannian solvmanifolds and solvable metric Lie algebras, it makes sense to work at the Lie algebra level.

Definition. Let (S, g) be a simply connected Riemannian solvmanifold, and let $(\mathfrak{s}, \langle \cdot, \cdot \rangle)$ be the corresponding solvable metric Lie algebra. Write $\mathfrak{s} = \mathfrak{a} \oplus \mathfrak{n}$, where \mathfrak{n} is the nilradical of \mathfrak{s} . We say the metric Lie algebra $(\mathfrak{s}, \langle \cdot, \cdot \rangle)$ and the solvmanifold (S, g) are of *Iwasawa type* if

- (i) \mathfrak{a} is abelian,
- (ii) for all $A \in \mathfrak{a}$, $(\text{ad } A)$ is symmetric relative to the inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{s} ,
- (iii) if $A \neq 0$, $(\text{ad } A) \neq 0$, and
- (iv) for some $A_0 \in \mathfrak{a}$, the restriction $(\text{ad } A_0)|_{\mathfrak{n}}$ is positive definite on \mathfrak{n} .

Definition. We say two metric Lie algebras are *isomorphic* if there exists a Lie algebra isomorphism $\varphi : \mathfrak{s} \rightarrow \mathfrak{s}'$ which is also an inner product space isometry.

In what follows, all solvmanifolds will be of Iwasawa type, with real roots. This guarantees our solvmanifolds are in standard position within the identity component of the isometry group of S [GW]. In this case, the question of when two solvmanifolds are isometric is answered at the metric Lie algebra level.

Theorem 2.1. [GW, 5.2] *Let $(\mathfrak{s}, \langle \cdot, \cdot \rangle)$ and $(\mathfrak{s}', \langle \cdot, \cdot \rangle')$ be data in standard position for simply connected Riemannian solvmanifolds S and S' , respectively. Then S' is isometric to S if and only if their corresponding metric Lie algebras are isomorphic.*

We will use this to verify that our examples are distinct from previously described solvmanifolds.

2.1. Symmetric spaces and subalgebras. Some solvmanifolds correspond to symmetric spaces of non-compact type in the following way (see Helgason [Hel] for more details). Given a noncompact symmetric space M , let $G = \text{Isom}(M)$, the full isometry group. The isotropy subgroup K at a point of M is a maximal compact subgroup of G , so that $M \cong G/K$. The Lie algebra \mathfrak{g} of G admits an Iwasawa decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{a} + \mathfrak{n}$ where \mathfrak{k} is the Lie algebra of K , \mathfrak{a} is abelian, \mathfrak{n} is nilpotent. We note that the solvable Lie subgroup S of G , with Lie algebra $\mathfrak{s} := \mathfrak{a} + \mathfrak{n}$, acts simply transitively on M , so that we may identify M with the solvmanifold S .

Alternatively—equivalently—we may begin with a semisimple Lie algebra \mathfrak{g} and a Cartan involution σ on \mathfrak{g} . Via the Cartan involution σ , we obtain a decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$, where $\sigma|_{\mathfrak{k}} = \text{Id}$, $\sigma|_{\mathfrak{p}} = -\text{Id}$. Using the Killing form B , we define $B_\sigma(X, Y) := -B(X, \sigma(Y))$, an $\text{ad}_{\mathfrak{k}}$ -invariant inner product on \mathfrak{g} .

Let \mathfrak{a} be a maximal torus in \mathfrak{p} . We let Δ denote a root system of \mathfrak{g} , and we take the root space decomposition $\mathfrak{g} = \mathfrak{g}_0 + \sum_{\alpha \in \Delta} \mathfrak{g}_\alpha$, where $\mathfrak{g}_\alpha = \{X \in \mathfrak{g} \mid (\text{ad } A)X = \alpha(A)X \text{ for all } A \in \mathfrak{a}\}$, \mathfrak{g}_0 is the centralizer of \mathfrak{a} in \mathfrak{g} . Denote by Δ^+ the set of positive roots and take $\Lambda \subset \Delta^+$ to be a set of simple roots. Using Δ^+ , we define $\mathfrak{n} := \sum_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$, the nilradical of \mathfrak{g} . Define $\mathfrak{s} := \mathfrak{a} + \mathfrak{n}$, endowed with this inner product induced from B_σ : $\langle \cdot, \cdot \rangle = 2B_\sigma|_{\mathfrak{a} \times \mathfrak{a}} + B_\sigma|_{\mathfrak{n} \times \mathfrak{n}}$. It is straightforward to see that the corresponding simply connected solvmanifold is a symmetric space, and the *rank* of the symmetric space is $|\Lambda|$, the number of elements in Λ (cf. [Hel, p.532]).

We are ready to describe Tamaru's metric Lie subalgebras $\mathfrak{s}_{\Lambda'}$ [T2]. We begin with a description of the subalgebra construction, then we review his proof that the appropriate metric on $\mathfrak{s}_{\Lambda'}$ has constant Ricci curvature.

We have $\Lambda = \{\alpha_1, \dots, \alpha_r\}$, the set of simple roots. We take a dual basis $\{H^1, \dots, H^r\}$ in \mathfrak{a} , so that $\alpha_i(H^j) = \delta_{ij}$. Choose a *characteristic element*,

$$Z = c_{i_1}H^{i_1} + c_{i_2}H^{i_2} + \dots + c_{i_k}H^{i_k},$$

for positive integers c_{i_1}, \dots, c_{i_k} . Note that every eigenvalue of $\text{ad } Z$ is an integer [KA]. For each $j \in \mathbf{Z}$, define $\mathfrak{g}^j := \sum_{\alpha(Z)=j} \mathfrak{g}_\alpha$, summing over $\alpha \in \Delta$. For any choice of characteristic element Z , the eigenspace decomposition $\mathfrak{g} = \bigoplus \mathfrak{g}^j$ gives a grading of \mathfrak{g} : $[\mathfrak{g}^i, \mathfrak{g}^j] \subset \mathfrak{g}^{i+j}$ for all $i, j \in \mathbf{Z}$. Define $\Lambda' = \{\alpha_i \in \Lambda \mid \alpha_i(Z) = 0\}$. Equivalently, $\Lambda' = \Lambda \setminus \{\alpha_{i_1}, \dots, \alpha_{i_k}\}$.

Just as \mathfrak{g}_0 is the centralizer of \mathfrak{a} in \mathfrak{g} , we denote by \mathfrak{k}_0 the centralizer of \mathfrak{a} in \mathfrak{k} . The subalgebra $\mathfrak{k}_0 \oplus \mathfrak{a} \oplus \mathfrak{n}$ is called a *minimal parabolic subalgebra* of \mathfrak{g} . Any subalgebra $\mathfrak{q} \subset \mathfrak{g}$ which contains $\mathfrak{k}_0 \oplus \mathfrak{a} \oplus \mathfrak{n}$ (up to conjugation) is a *parabolic subalgebra*. Every parabolic subalgebra can be constructed from a subset $\Lambda' \subset \Lambda$ [Kn, 7.76]. In [T2, 3.6], Tamaru proves $\mathfrak{q}_{\Lambda'} := \sum_{k>0} \mathfrak{g}^k$ is a parabolic subalgebra of \mathfrak{g} . On $\mathfrak{q}_{\Lambda'}$, one has the following Langlands decomposition [T2, 3.8]:

- $\mathfrak{a}_{\Lambda'} = \text{span}\{H^{i_1}, \dots, H^{i_k}\}$,
- $\mathfrak{m}_{\Lambda'} = \mathfrak{g}^0 \ominus \mathfrak{a}_{\Lambda'}$,
- $\mathfrak{n}_{\Lambda'} = \sum_{k>0} \mathfrak{g}^k$.

Finally, our solvable metric Lie algebra is $\mathfrak{s}_{\Lambda'} := \mathfrak{a}_{\Lambda'} + \mathfrak{n}_{\Lambda'}$, endowed with the inner product inherited from that of \mathfrak{g} : $\langle \cdot, \cdot \rangle = 2B_\sigma|_{\mathfrak{a}_{\Lambda'} \times \mathfrak{a}_{\Lambda'}} + B_\sigma|_{\mathfrak{n}_{\Lambda'} \times \mathfrak{n}_{\Lambda'}}$. We say the corresponding simply connected solvmanifold $(S_{\Lambda'}, g)$ is an *attached solvmanifold*.

Remark. A subset $\Lambda' \subset \Lambda$ is said to be *trivial* if Λ' and $\Lambda \setminus \Lambda'$ are orthogonal. Provided $\Lambda' \subset \Lambda$ is nontrivial, the solvmanifold $(S_{\Lambda'}, g)$ will not be a totally geodesic submanifold of (S_\emptyset, g) [T2, 6.4].

Proposition 2.2. [T2, 4.4] *When Λ' is chosen to be the empty set, the attached solvmanifold $(\mathfrak{s}_\emptyset = \mathfrak{a}_\emptyset + \mathfrak{n}_\emptyset, \langle \cdot, \cdot \rangle)$ is a symmetric space and an Einstein manifold.*

Now, take any nonempty Λ' . Tamaru proves that $\mathfrak{s}_{\Lambda'} = \mathfrak{a}_{\Lambda'} + \mathfrak{n}_{\Lambda'}$ is a subalgebra of \mathfrak{s}_\emptyset , and the Ricci curvature of $\mathfrak{s}_{\Lambda'}$ is the same as that of \mathfrak{s}_\emptyset . Thus $(\mathfrak{s}_{\Lambda'}, \langle \cdot, \cdot \rangle)$ is an Einstein solvmanifold [T2, 5.3]. We outline Tamaru's proof below.

2.2. Ricci curvature. Recall the sectional and Ricci curvature formulae, viewed as bilinear forms on \mathfrak{s} (cf. [Bes]). Let $\{X_1, \dots, X_n\}$ be an orthonormal basis for \mathfrak{s} . The Levi-Civita connection $\nabla_X Y$ of (S, g) is given by

$$\nabla_X Y = \frac{1}{2}[X, Y] + U(X, Y)$$

where $U : \mathfrak{s} \times \mathfrak{s} \rightarrow \mathfrak{s}$ is the symmetric bilinear form satisfying

$$\langle U(X, Y), Z \rangle = \frac{1}{2}\langle [Z, X], Y \rangle + \frac{1}{2}\langle [Z, Y], X \rangle.$$

Let B denote the Killing form for \mathfrak{s} . Then for a pair of orthonormal vectors $X, Y \in \mathfrak{s}$, the sectional curvature of the two-plane spanned by X and Y is given by

$$K(X, Y) = -\frac{3}{4}\| [X, Y] \|^2 - \frac{1}{2}\langle [X, [X, Y]], Y \rangle - \frac{1}{2}\langle [Y, [Y, X]], X \rangle + |U(X, Y)|^2 - \langle U(X, X), U(Y, Y) \rangle.$$

The Ricci curvature is given by

$$\text{Ric}(X, Y) = -\frac{1}{2} \sum_i \langle [X, X_i], [Y, X_i] \rangle - \frac{1}{2}B(X, Y) + \frac{1}{4} \sum_{i,j} \langle [X_i, X_j], X \rangle \langle [X_i, X_j], Y \rangle - \langle U(X, Y), H_0 \rangle$$

where $H_0 = \sum U(X_i, X_i)$ is the mean curvature vector.

Since these solvable metric Lie algebras are of Iwasawa type, the Ricci curvature expression simplifies:

Lemma 2.3. [W, 1.4] *Let Ric denote the Ricci curvature tensor on $\mathfrak{s} = \mathfrak{a} \oplus \mathfrak{n}$, a solvable metric Lie algebra. Let Ricⁿ denote the Ricci curvature tensor on \mathfrak{n} . For $X \in \mathfrak{n}$, $A, A' \in \mathfrak{a}$,*

(i) $\text{Ric}(A, A') = -\text{tr}(\text{ad } A \circ \text{ad } A')$,

(ii) $\text{Ric}(A, X) = 0$,

(iii) $\text{Ric}(X) = \text{Ric}^n(X) - \langle [H_0, X], X \rangle$, where H_0 is the mean curvature vector.

The proof that Ricci curvature is unchanged relies on Wolter's lemma and these observations. (Recall, \mathfrak{g}^0 is the sum of the root spaces \mathfrak{g}_α for which $\alpha(Z) = 0$.)

- (1) Since $[\mathfrak{a}_{\Lambda'}, \mathfrak{g}^0] = 0$, we see $\text{tr}(\text{ad } A \circ \text{ad } A')|_{\mathfrak{n}_{\Lambda'}} = \text{tr}(\text{ad } A \circ \text{ad } A')|_{\mathfrak{n}_0}$ for all $A, A' \in \mathfrak{a}_{\Lambda'}$. Thus $\text{Ric}^{\mathfrak{s}_{\Lambda'}}(A, A') = \frac{1}{4}\langle A, A' \rangle = \text{Ric}^{\mathfrak{s}^0}(A, A')$.
- (2) For any $A \in \mathfrak{a}_{\Lambda'}$ and any $X \in \mathfrak{n}_{\Lambda'}$, $\text{Ric}^{\mathfrak{s}_{\Lambda'}}(A, X) = 0 = \text{Ric}^{\mathfrak{s}^0}(A, X)$.
- (3) Let $\{E'_i\}$ be an orthonormal basis for $\mathfrak{n}_{\Lambda'}$, and let $\{E_j^\perp\}$ be an orthonormal basis for $\mathfrak{n}_0 \ominus \mathfrak{n}_{\Lambda'}$. Define $H'_0 = \frac{1}{2} \sum [\sigma E'_i, E'_i]$ and $H^\perp = \frac{1}{2} \sum [\sigma E_j^\perp, E_j^\perp]$, so that $H_0 = H'_0 + H^\perp$. Hence $\text{Ric}^{\mathfrak{s}^0}(X, Y) = \langle \text{Ric}^{\mathfrak{n}^0}(X), Y \rangle - \langle [H_0, X], Y \rangle$, $\text{Ric}^{\mathfrak{s}_{\Lambda'}}(X, Y) = \langle \text{Ric}^{\mathfrak{n}_{\Lambda'}}(X), Y \rangle - \langle [H'_0, X], Y \rangle$.
- (4) We see $\text{Ric}^{\mathfrak{n}^0}(X) = -\frac{1}{4} \sum [E_i, [\sigma E_i, X]_{\mathfrak{n}_0}] + \frac{1}{2} \sum [\sigma E_i, [E_i, X]]_{\mathfrak{n}_0}$,
and $\text{Ric}^{\mathfrak{n}_{\Lambda'}}(X) = -\frac{1}{4} \sum [E'_j, [\sigma E'_j, X]_{\mathfrak{n}_{\Lambda'}}] + \frac{1}{2} \sum [\sigma E'_j, [E'_j, X]]_{\mathfrak{n}_{\Lambda'}}$.

Using these, Tamaru proves $\text{Ric}^{\mathfrak{n}^0}(X) - \text{Ric}^{\mathfrak{n}_{\Lambda'}}(X) = [H_0^\perp, X]$ for all $X \in \mathfrak{n}_{\Lambda'}$. Thus,

$$\begin{aligned} \text{Ric}^{\mathfrak{s}^0}(X, Y) - \text{Ric}^{\mathfrak{s}_{\Lambda'}}(X, Y) &= \langle \text{Ric}^{\mathfrak{n}^0}(X) - \text{Ric}^{\mathfrak{n}_{\Lambda'}}(X), Y \rangle - \langle [H_0 - H'_0, X], Y \rangle \\ &= \langle [H_0^\perp, X], Y \rangle - \langle [H_0^\perp, X], Y \rangle = 0. \end{aligned}$$

2.3. Associated solvmanifolds. We now review the method in [GK, 4.2] of modifying the algebraic structure of a solvmanifold while preserving the Ricci curvature. We begin with our real semisimple Lie algebra \mathfrak{g} . Let $\mathfrak{g}^{\mathbf{R}} = (\mathfrak{g})^{\mathbf{C}}$, the complexification of \mathfrak{g} , viewed as a real Lie algebra. View \mathfrak{g} , and also \mathfrak{s} , as subalgebras of $\mathfrak{g}^{\mathbf{R}}$. Recall that the root spaces \mathfrak{g}_α are mutually orthogonal relative to the inner product defined by the symmetric metric on S .

Definition. There exists an *adapted* orthonormal basis \mathcal{B} of the subspace \mathfrak{n} of \mathfrak{s} such that

- each basis vector lies in some \mathfrak{g}_β ,
- the bracket of any two basis vectors is a scalar multiple of another basis vector,
- for any X, Y , and U basis vectors with $Y \neq U$, then $[X, Y] \perp [X, U]$.

Definition. Let \mathcal{B} be an adapted orthonormal basis \mathcal{B} for \mathfrak{n} . Consider a new subspace of $\mathfrak{g}^{\mathbf{R}}$ with basis \mathcal{B}' , where we obtain \mathcal{B}' from \mathcal{B} by replacing some of the vectors $X \in \mathcal{B}$ by $\sqrt{-1}X$. Our new subspace \mathfrak{n}' will be a nilpotent algebra provided it is closed under the bracket operation in \mathfrak{g} . The brackets of the basis vectors in \mathfrak{n}' differ only by a sign from their corresponding brackets in \mathfrak{n} . Notice the subalgebra \mathfrak{a} of $\mathfrak{g}^{\mathbf{R}}$ normalizes \mathfrak{n}' . We choose the inner product on \mathfrak{n}' which makes \mathcal{B}' orthonormal, and choose the inner product on $\mathfrak{s}' := \mathfrak{a} \oplus \mathfrak{n}'$ as an orthogonal sum of inner product spaces. Then \mathfrak{s}' is a solvable metric Lie algebra. We say the corresponding Riemannian solvmanifold S' is *associated* to the symmetric space G/K .

Remark. The simply connected solvmanifold S' corresponding to \mathfrak{s}' is an Einstein manifold: the sign changes are cancelled out in the Ricci curvature equation [GK, 4.3]. It has since been proved that whenever \mathfrak{n} and \mathfrak{n}' are two real nilpotent Lie algebras whose complexifications are isomorphic, if \mathfrak{n} is an Einstein nilradical then \mathfrak{n}' is as well ([N, 6],[J3, 6.5]).

We note that one can first construct an associated solvmanifold S' (associated to S), then consider the solvmanifolds $S'_{\Lambda'}$ attached to S' . Alternatively, one can start from solvmanifold S , construct the solvmanifolds $S_{\Lambda'}$ attached to S , and from there, create their associated solvmanifolds $S_{\Lambda'}$. These two techniques can be combined in either order to obtain the same new Einstein solvmanifolds.

2.4. Isometries of solvmanifolds. We come to the issue of determining whether we have indeed created new solvmanifolds. We verify that these examples are distinct from Tamaru's. To do this, we note that every attached solvmanifold $S_{\Lambda'}$ will inherit isometries from S_{\emptyset} .

Proposition 2.4. *Consider $\text{Isom}(S_{\Lambda'})$, the group of isometries of our solvmanifold. Let $M_{\Lambda'}$ denote the compact component of $\text{Isom}(S_{\Lambda'})$. Then $M_{\emptyset} \subseteq M_{\Lambda'}$.*

Proof. We know from the work of Gordon and Wilson that the compact component M of the isometry group of a solvmanifold S is the normalizer of A in K , the maximal compact subgroup of $G = \text{Isom}(S)$ [GW]. That is, M_{\emptyset} is the normalizer of A in K (here $S_{\emptyset} = G/K$, and $G = \text{Isom}(S_{\emptyset}) = KAN$). The isometries of a solvmanifold S must preserve root spaces; hence all isometries of S_{\emptyset} take $S_{\Lambda'}$ to itself. Our conclusion, $M_{\emptyset} \subseteq M_{\Lambda'}$, follows directly [GW, La. 6.4]. \square

Now, we contrast the attached solvmanifold case with the case of associated solvmanifolds.

Proposition 2.5. [GK, 4.4] *If S' and S'' are two Riemannian solvmanifolds associated with G/K , then S' is isometric to S'' if and only if there exists an automorphism ϕ of the root system (i.e., a Weyl group element) and an isomorphism $\tau : \mathfrak{n}' \rightarrow \mathfrak{n}''$ such that $\tau(\mathfrak{n}'_{\alpha}) = \mathfrak{n}''_{\phi(\alpha)}$ for all roots α .*

Proposition 2.6. *Let S' be a solvmanifold associated to the symmetric space G/K . Consider $\text{Isom}(S')$, the group of isometries of our solvmanifold S' . Let M denote the compact component of $\text{Isom}(G/K)$ and M' denote the compact component of $\text{Isom}(S')$. Then $M' \subset M$.*

Proof. Recall, $M \subset \text{Isom}(G/K)$ is the normalizer of A in K . Since the isometries in M' must preserve root spaces, we see that when a selection of basis vectors X are replaced by $\sqrt{-1}X$ to construct \mathcal{B}' (and hence S'), some isometries may be lost. \square

In what follows, in each case we will verify that for our associated solvmanifolds S' , $M' \subsetneq M$. That is, the compact component M' of $\text{Isom}(S')$ is a proper subset of M .

3. NEW EXAMPLES

3.1. $SO(p, q)/SO(p)SO(q)$, $SU(p, q)/S(U(p)U(q))$, and $Sp(p, q)/Sp(p)Sp(q)$. We start with the noncompact symmetric spaces of the form $SO(p, q)/S(O(p)O(q))$, $SU(p, q)/S(U(p)U(q))$ or $Sp(p, q)/Sp(p)Sp(q)$. Without loss of generality, we may assume that $q \geq p$, so that the rank of these symmetric spaces is p . Then

$$\mathfrak{g} = \{A \in \mathfrak{gl}(p+q, \mathbf{F}) \mid AM = -MA^t\} \quad \text{where} \quad M = \begin{pmatrix} \text{Id}_p & 0 \\ 0 & -\text{Id}_q \end{pmatrix},$$

where $\mathbf{F} = \mathbf{R}$ in the case of $\mathfrak{so}(p, q)$, $\mathbf{F} = \mathbf{C}$ in the case of $\mathfrak{su}(p, q)$, and $\mathbf{F} = \mathbf{H}$ in the case of $\mathfrak{sp}(p, q)$.

Elements of \mathfrak{g} may be written in matrix block form this way, $\begin{pmatrix} A & B & C \\ B^t & D & E \\ C^t & -E^t & F \end{pmatrix}$, such that blocks

A, B, D are $p \times p$, blocks C, E are $p \times (q - p)$, and block F is $(q - p) \times (q - p)$, and furthermore, $A = -A^t$, $D = -D^t$, $F = -F^t$, while B, C, E are arbitrary.

We consider the maximal torus \mathfrak{a} consisting of the matrices in \mathfrak{g} for which B is diagonal, and all other blocks are zero. Let $\{\omega_1, \dots, \omega_p\}$ denote the basis of duals to the standard basis for the set of diagonal elements of B . Then we may choose the set of positive roots

$$\Delta^+ = \{\omega_i \mid 1 \leq i \leq p\} \cup \{\omega_j \pm \omega_i \mid 1 \leq i < j \leq p\} \text{ (orthogonal)}$$

$$\Delta^+ = \{\omega_i \mid 1 \leq i \leq p\} \cup \{\omega_j \pm \omega_i \mid 1 \leq i < j \leq p\} \cup \{2\omega_i \mid 1 \leq i \leq p\} \text{ (unitary, symplectic).}$$

We may take $\Lambda = \{\omega_1, \omega_2 - \omega_1, \omega_3 - \omega_2 \dots, \omega_p - \omega_{p-1}\}$ as the fundamental system. We give bases of root vectors below. In what follows, E_{ij} is the skew-symmetric matrix with 1 in the (i, j) entry, -1 in the (j, i) entry and zeroes elsewhere; F_{ij} is the symmetric matrix with 1 in the (i, j) and (j, i) entries and zeroes elsewhere. (While the basis elements for Λ above are not orthogonal, the basis vectors below are orthonormal.)

$$\begin{aligned} \mathfrak{n}_{\omega_k} &= \text{span}\{U_{km} = (F_{km} + E_{p+k,m}), \epsilon V_{k,m} = \epsilon(E_{km} + F_{p+k,m}) \mid 2p+1 \leq m \leq p+q\} \\ \mathfrak{n}_{\omega_j - \omega_i} &= \text{span}\{Y_{jk}^- = \frac{1}{\sqrt{2}}(E_{jk} + E_{p+j,p+k} - F_{j,p+k} - F_{k,p+j}), \\ &\quad \epsilon Z_{jk}^- = \frac{\epsilon}{\sqrt{2}}(F_{jk} + F_{p+j,p+k} - E_{j,p+k} + E_{k,p+j})\} \\ \mathfrak{n}_{\omega_j + \omega_i} &= \text{span}\{Y_{jk}^+ = \frac{1}{\sqrt{2}}(E_{jk} - E_{p+j,p+k} - F_{j,p+k} + F_{k,p+j}), \\ &\quad \epsilon Z_{jk}^+ = \frac{\epsilon}{\sqrt{2}}(F_{jk} - F_{p+j,p+k} - E_{j,p+k} - E_{k,p+j})\} \\ \mathfrak{n}_{2\omega_k} &= \text{span}\{\epsilon W_k = \epsilon(F_{kk} - F_{p+k,p+k} - E_{k,p+k})\} \end{aligned}$$

The basis vectors of type V_{ij} , Z_{ij}^\pm and W_j occur only in the unitary and symplectic cases, and $\epsilon^2 = -1$. In the unitary case, $\epsilon = i$. In the symplectic case, $\epsilon \in \{i, j, k\}$.

Notice that for each k , the root space \mathfrak{n}_{ω_k} lies in $W = \left\{ \begin{pmatrix} 0 & 0 & C \\ 0 & 0 & C \\ C^t & -C^t & 0 \end{pmatrix} \right\}$. In particular, \mathfrak{n}_{ω_k}

is the space for which all but the k^{th} row of C is zero. Meanwhile, the root spaces $\mathfrak{n}_{\omega_j - \omega_i}$, $\mathfrak{n}_{\omega_j + \omega_i}$,

and $\mathfrak{n}_{2\omega_i}$ lie in $\begin{pmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & 0 \end{pmatrix}$. These observations will be useful in what follows.

In the case that our space has $q - p \geq 2$, we obtain an *associated* solvmanifold as follows. For any integer a with $1 \leq a \leq q - p$ and for b such that $a + b = q - p$, we obtain a decomposition $W = W_a \oplus W_b$. In W_a , the last b columns of C are all zero. In W_b , the first a columns of C are all zero.

We construct \mathfrak{s}_a by replacing each X in W_b by $X' = \sqrt{-1}X$ in $\mathfrak{s}^{\mathbf{C}}$, leaving W_a unchanged. With the following observations it is clear that \mathfrak{s}_a is closed under the bracket in $\mathfrak{s}^{\mathbf{C}}$:

- (i) W commutes with all root spaces of type $\mathfrak{n}_{2\omega_i}$, $\mathfrak{n}_{\omega_j + \omega_i}$
- (ii) W_b commutes with W_a
- (iii) The maximal torus \mathfrak{a} and the root spaces $\mathfrak{n}_{\omega_j - \omega_i}$ normalize each of W_a and W_b .

The associated simply connected solvmanifold S_a corresponding to \mathfrak{s}_a is Einstein. Yet S_a , unlike the symmetric manifold, admits some positive sectional curvature. Thus S_a cannot be isometric to its corresponding symmetric associate.

To see a two-plane in S_a admitting positive curvature, let i and j be integers with $1 \leq i \leq a < j \leq a + b$. For $1 \leq k < l \leq p$, let X_k be the element of \mathfrak{n}_{ω_k} for which C has (k, i) and (k, j) entries equal to $\frac{1}{\sqrt{2}}$ and all other entries zero, and let X_l in \mathfrak{n}_{ω_l} be defined analogously. Let X'_k and X'_l be the corresponding unit vectors in \mathfrak{s}_a . Then we have $[X'_k, X'_l] = 0$ and $\langle U(X'_k, X'_k), U(X'_l, X'_l) \rangle = 0$. On the other hand, $U(X'_k, X'_l)$ is a nonzero element of $\mathfrak{n}_{\omega_l - \omega_k}$. Thus $K(X'_k, X'_l) > 0$ [GK].

Remark. Two solvmanifolds S_a and $S_{a'}$ both associated to S will be isometric to each other if and only if either $a = a'$ or $a = (q - p) - a'$, by Prop. 2.5.

In the construction of our solvable Lie group \mathfrak{s}_a out of \mathfrak{s} , the maximal torus \mathfrak{a} , the root system Δ and the fundamental system Λ are unchanged. Thus we have a dual basis for \mathfrak{a} , $\{H^1, H^2, \dots, H^p\}$, such that H^i is dual to the i^{th} element of Λ . Each choice of characteristic element $Z = c_{i_1} H^{i_1} + \dots + c_{i_k} H^{i_k}$ (with $c_{i_j} \in \mathbf{Z}_+$), determines $\Lambda' \subset \Lambda$ and, in turn, Λ' determines a solvable metric Lie subalgebra $(\mathfrak{s}_a)_{\Lambda'} = \mathfrak{a}_{\Lambda'} + (\mathfrak{n}_a)_{\Lambda'} \subset \mathfrak{s}_a$. This proves the statement below.

Proposition 3.1. *Let S_a be a solvmanifold associated to a solvmanifold corresponding to a noncompact symmetric space of type $\mathrm{SO}(p, q)/\mathrm{S}(\mathrm{O}(p)\mathrm{O}(q))$, $\mathrm{SU}(p, q)/\mathrm{S}(\mathrm{U}(p)\mathrm{U}(q))$ or $\mathrm{Sp}(p, q)/\mathrm{Sp}(p)\mathrm{Sp}(q)$, in the sense of [GK]. Let $(S_a)_{\Lambda'}$ be an attached solvmanifold to S_a . Then $((S_a)_{\Lambda'}, g')$ is an Einstein solvmanifold.*

For each such symmetric space, there are $\lfloor \frac{q-p}{2} \rfloor$ associated Einstein solvmanifolds (without loss of generality, $q \geq p$). For each associated solvmanifold, there are approximately $2^p - 1$ attached solvmanifolds (depending on the size of the Weyl group).

Remark. The compact subgroup of isometries of G/K is $M = \mathrm{SO}(q-p)$, which corresponds exactly to the block labeled F in our matrix diagram. The subspace W is invariant under the elements of M . For the associated solvmanifold S_a , the compact subgroup of isometries is $M_a = \mathrm{SO}(a)\mathrm{SO}(b) \subset \mathrm{SO}(q-p)$, elements for which now each of the separate components W_a and W_b are invariant (if $a = b$, M_a also has a $\mathbf{Z}/2\mathbf{Z}$ factor, interchanging W_a and W_b).

Analogous statements hold for the unitary and symplectic symmetric spaces.

Example. We will consider the specific example of $S \cong S_\emptyset$, the solvmanifold which corresponds to the symmetric space $\mathrm{SO}(3, 5)/\mathrm{SO}(3)\mathrm{SO}(5)$. From there we construct an associated solvmanifold S_1 , still Einstein but no longer symmetric; for example, our solvmanifold is no longer negatively curved. Finally, we will choose a nontrivial subset $\Lambda' \subset \Lambda$ of fundamental roots, to study an attached submanifold $(S_1)_{\Lambda'} \subset S_1$.

On the Lie algebra level, we start with $\mathfrak{g} = \mathfrak{so}(3, 5)$, $\mathfrak{k} = \mathfrak{so}(3)\mathfrak{so}(5)$, and $\mathfrak{}$. As described above, \mathfrak{a} is made up of the matrices in \mathfrak{g} for which B is diagonal and all other blocks are zero. The basic roots are given by $\Lambda = \{\omega_1, \omega_2 - \omega_1, \omega_3 - \omega_2\}$. The set of positive roots is

$$\Delta^+ = \Lambda \cup \{\omega_2, \omega_3, \omega_2 + \omega_1, \omega_3 + \omega_2, \omega_3 \pm \omega_1\}.$$

The vectors listed here give an adapted basis for S :

$$\begin{aligned}\mathfrak{a} &= \text{span}\{F_{14}, F_{25}, F_{36}\}, \\ \mathfrak{n}_{\omega_k} &= \text{span}\{U_{km} = F_{km} + E_{3+k,m} \mid m = 7, 8\}, \quad k = 1, 2, 3, \\ \mathfrak{n}_{\omega_j \pm \omega_i} &= \text{span}\{Y_{ij}^\pm = \frac{1}{\sqrt{2}}(E_{ij} - F_{i,3+j} \pm F_{j,3+i} \mp E_{3+i,3+j})\}, \quad 1 \leq i < j \leq 3.\end{aligned}$$

In this example, there is only one non-isomorphic associated solvmanifold, S_1 : take $a = b = 1$ in the description above. Then $W_a = \text{span}\{U_{17}, U_{27}, U_{37}\}$ and $W_b = \text{span}\{U_{18}, U_{28}, U_{38}\}$. To construct our associated solvmanifold S_1 , we replace U_{i7} with $\sqrt{-1}U_{i7}$, for $i = 1, 2, 3$. Here we note that S_1 is different from S ; we show we have introduced positive sectional curvature. Let $X = \frac{1}{\sqrt{2}}(U_{27} + U_{28})$ and $Y = \frac{1}{\sqrt{2}}(U_{37} + U_{38})$. Then $[X, Y] = 0$. Meanwhile, $U(X, Y)$ is a nonzero element of $\mathfrak{n}_{\omega_3 - \omega_2}$, whereas $\langle U(X, X), U(Y, Y) \rangle = 0$. Thus we get $K(X, Y) = |U(X, Y)|^2 > 0$.

Next, we consider a characteristic element Z in \mathfrak{a} : we take $Z = H^1 + H^2 = F_{14} + 2F_{25} + 2F_{36}$. With this choice of Z , we find that $\Lambda' = \{\alpha \in \Lambda \mid \alpha(Z) = 0\} = \{\omega_3 - \omega_2\}$. Now $\mathfrak{n}_{\omega_3 - \omega_2}$ is removed. We consider the same section as above, $X = \frac{1}{\sqrt{2}}(U_{27} + U_{28})$ and $Y = \frac{1}{\sqrt{2}}(U_{37} + U_{38})$. One sees that we still have $[X, Y] = 0$, but now $U_{\Lambda'}(X, Y) = 0$. And because we must project from \mathfrak{a}_θ to $\mathfrak{a}_{\Lambda'}$, we see $\langle U_{\Lambda'}(X, X), U_{\Lambda'}(Y, Y) \rangle > 0$. Thus the section which has positive sectional curvature in S_1 has negative sectional curvature in $(S_1)_{\Lambda'}$.

3.2. $SO(n, \mathbf{H})/U(n)$. We now consider the noncompact symmetric space $SO(n, \mathbf{H})/U(n)$, of rank $m = \lfloor \frac{n}{2} \rfloor$. We have

$$\mathfrak{g} = \mathfrak{so}(n, \mathbf{H}) = \left\{ \begin{pmatrix} X & Y \\ -\bar{Y} & X \end{pmatrix} \in \mathfrak{gl}(2n, \mathbf{C}) \mid X = -X^t, Y = \bar{Y}^t \right\}.$$

Within \mathfrak{g} we consider the maximal torus $\mathfrak{a} = \{H_j = \frac{i}{\sqrt{2}}(E_{2j-1,2j} - E_{n+2j-1,n+2j}) \mid j = 1, \dots, m\}$. We choose the set of positive roots

$$\begin{aligned}\Delta^+ &= \{\omega_j \pm \omega_k \mid 1 \leq j < k \leq m\} \cup \{2\omega_j \mid j = 1, \dots, m\} \quad (\text{if } n \text{ even}) \\ \Delta^+ &= \{\omega_j \pm \omega_k \mid 1 \leq j < k \leq m\} \cup \{2\omega_j \mid j = 1, \dots, m\} \cup \{\omega_j \mid j = 1, \dots, m\} \quad (\text{if } n \text{ odd}).\end{aligned}$$

For completeness, we describe the root spaces below (see [GK, 4.2]):

$$\begin{aligned}\mathfrak{n}_{\omega_j \pm \omega_k} &= \text{span}\{A_{jk}^\pm = \frac{1}{2}((E_{2j-1,2k-1} \mp E_{2j,2k} + E_{n+2j-1,n+2k-1} \mp E_{n+2j,n+2k}) \\ &\quad + i(\mp E_{2j-1,2k} - E_{2j,2k-1} \pm E_{n+2j-1,n+2k} + E_{n+2j,n+2k-1))), \\ B_{jk}^\pm &= \frac{1}{2}((E_{2j-1,2k} \pm E_{2j,2k-1} + E_{n+2j-1,n+2k} \pm E_{n+2j,n+2k-1}) \\ &\quad + i(\pm E_{2j-1,2k-1} - E_{2j,2k} \mp E_{n+2j-1,n+2k-1} + E_{n+2j,n+2k})), \\ C_{jk}^\pm &= \frac{1}{2}((E_{2j-1,n+2k} \mp E_{2j,n+2k-1} \mp E_{2k-1,n+2j} + E_{2k,n+2j-1}) \\ &\quad + i(\mp E_{2j-1,n+2k-1} - E_{2j,n+2k} \pm E_{2k-1,n+2j-1} + E_{2k,n+2j})), \\ D_{jk}^\pm &= \frac{1}{2}((E_{2j-1,n+2k-1} + E_{2k-1,n+2j-1} \pm E_{2j,n+2k} \pm E_{2k,n+2j}) \\ &\quad + i(\pm E_{2j-1,n+2k} - E_{2j,n+2k-1} + E_{2k-1,n+2j} \mp E_{2k,n+2j-1})); \\ \mathfrak{n}_{2\omega_k} &= \text{span}\{G_k = \frac{1}{\sqrt{2}}((E_{2k-1,n+2k-1} + E_{2k,n+2k}) + i(E_{2k-1,n+2k} - E_{2k,n+2k-1})); \\ (\text{and if } n \text{ odd}) \quad \mathfrak{n}_{\omega_k} &= \text{span}\{X_k = \frac{1}{\sqrt{2}}((E_{2k,n} + E_{n+2k,2n}) + i(E_{2k-1,n} - E_{n+2k-1,2n})),\end{aligned}$$

$$\begin{aligned}
Y_k &= \frac{1}{\sqrt{2}}((E_{2k-1,n} + E_{n+2k-1,2n}) - i(E_{2k,n} - E_{n+2k,2n})), \\
Z_k &= \frac{1}{\sqrt{2}}((E_{2k,2n} + E_{n,n+2k}) + i(E_{2k-1,2n} - E_{n,n+2k-1})), \\
W_k &= \frac{1}{\sqrt{2}}((E_{2k-1,2n} + E_{n,n+2k-1}) - i(E_{2k,2n} - E_{n,n+2k})).
\end{aligned}$$

In the case that n is even, we can modify \mathfrak{s} by replacing each of the basis elements $B_{jk}^-, C_{jk}^-, A_{jk}^+, D_{jk}^+$ and G_k by its respective product with $\sqrt{-1}$, an element of $\mathfrak{s}^{\mathbf{C}}$, the complexified Lie algebra. In the case that n is odd, instead we modify \mathfrak{s} by replacing each of the basis elements $X_k, Z_k, B_{jk}^+, C_{jk}^+, B_{jk}^-, C_{jk}^-$ by its respective product with $\sqrt{-1}$. In either case, with this changed basis, we obtain a new Lie algebra \mathfrak{s}' associated to \mathfrak{s} , and the corresponding simply-connected Riemannian solvmanifold S' is Einstein.

As with the previous examples, these new solvmanifolds are not isomorphic to their associated symmetric solvmanifolds. To see this, we show here that S' admits some positive sectional curvature [GK]. To construct a two-plane of positive curvature, we set

$$X = \frac{1}{\sqrt{2}}((A_{j,j+1}^-)' + (B_{j,j+1}^-)') \quad \text{and} \quad Y = \frac{1}{\sqrt{2}}((C_{j+1,j+2}^-)' + (D_{j+1,j+2}^-)').$$

Then one sees that while $[X, Y] = 0$ and $U(X, Y) = 0$, we have $U(X, X) = \frac{1}{\sqrt{2}}(H_j - H_{j+1})$ and $U(Y, Y) = \frac{1}{\sqrt{2}}(H_{j+1} - H_{j+2})$. Thus

$$K(X, Y) = -\langle U(X, X), U(Y, Y) \rangle = -\langle \frac{1}{\sqrt{2}}(H_j - H_{j+1}), \frac{1}{\sqrt{2}}(H_{j+1} - H_{j+2}) \rangle = \frac{1}{2}\|H_{j+1}\| > 0.$$

Proposition 3.2. *Let S' be the solvmanifold associated to a solvmanifold corresponding to a noncompact symmetric space of type $\mathrm{SO}(n, \mathbf{H})/\mathrm{U}(n)$, in the sense of [GK]. Let $m = \lfloor \frac{n}{2} \rfloor$. Let $S'_{\Lambda'}$ be an attached solvmanifold to S' in the sense of [T2]. Then $(S'_{\Lambda'}, g')$ is an Einstein solvmanifold.*

Proof. In the construction of our solvable Lie group \mathfrak{s}' , the maximal torus \mathfrak{a} , the root system Δ , and the fundamental system Λ are unchanged from \mathfrak{s} . We have a dual basis for \mathfrak{a} , $\{H^1, H^2, \dots, H^m\}$ such that H^i is dual to the i^{th} element of Λ . Each choice of characteristic element $Z = c_{i_1}H^{i_1} + \dots + c_{i_k}H^{i_k}$ (each $c_{i_j} \in \mathbf{Z}_+$), determines $\Lambda' \subset \Lambda$ and, in turn, Λ' determines an attached solvable metric Lie subalgebra $\mathfrak{s}'_{\Lambda'} = \mathfrak{a}_{\Lambda'} + \mathfrak{n}'_{\Lambda'} \subset \mathfrak{s}'$. \square

For each n , there is one associated Einstein solvmanifold, and for each associated solvmanifold, there are approximately $2^m - 1$ attached solvmanifolds (depending on the number of symmetries of the Dynkin diagram).

Remark. The compact subgroup of isometries of G/K is $M = \mathrm{SO}(4)$ (where the action permutes $A_{jk}^{\pm}, B_{jk}^{\pm}, C_{jk}^{\pm}$ and D_{jk}^{\pm} , preserving the plus or minus type). For the associated solvmanifold S' , the analogous compact subgroup of isometries is $M' = \mathrm{SO}(2)\mathrm{SO}(2)$.

3.3. $\mathrm{SL}(n, \mathbf{H})/\mathrm{Sp}(n)$. Finally, we consider the noncompact symmetric space $\mathrm{SL}(n, \mathbf{H})/\mathrm{Sp}(n)$, of rank $n - 1$. Here

$$\mathfrak{g} = \mathfrak{sl}(n, \mathbf{H}) = \left\{ \begin{pmatrix} X & -\bar{Y} \\ Y & \bar{X} \end{pmatrix} \mid X, Y \in \mathfrak{gl}(n, \mathbf{C}), \mathrm{tr}(X + \bar{X}) = 0 \right\}.$$

Within \mathfrak{g} we have $\mathfrak{a} = \{\mathrm{diag}(a_1, \dots, a_n, a_1, \dots, a_n) \in \mathfrak{gl}(2n, \mathbf{R}) \mid \sum_{i=1}^n a_i = 0\}$. For an arbitrary element $H = \mathrm{diag}(a_1, \dots, a_n, a_1, \dots, a_n) \in \mathfrak{a}$, define $\omega_j \in \mathfrak{a}^*$ by $\omega_j(H) = a_j$. The positive roots are

$\Delta^+ = \{\omega_k - \omega_j \mid 1 \leq j < k \leq n\}$. The root spaces are described below (see [GK, 4.3]):

$$\begin{aligned} \mathfrak{n}_{\omega_j - \omega_k} = \text{span}\{ & A_{jk} = i\sqrt{2}(F_{jk} - F_{n+j, n+k}), \quad B_{jk} = i\sqrt{2}(F_{j, n+k} + F_{n+j, k}), \\ & C_{jk} = \sqrt{2}(F_{j, n+k} - F_{n+j, k}), \quad D_{jk} = \sqrt{2}(F_{jk} + F_{n+j, n+k})\}. \end{aligned}$$

We modify \mathfrak{s} by replacing each A_{jk} and C_{jk} by their respective products with $\sqrt{-1}$, to get new elements of $\mathfrak{s}^{\mathbf{C}}$ for every $1 \leq j < k \leq n$. With these changes, we obtain a new Lie algebra \mathfrak{s}' associated to \mathfrak{s} , and the corresponding simply-connected Riemannian solvmanifold S' is Einstein.

We show that, since S' admits some positive sectional curvature, it cannot be isomorphic to S , the symmetric solvmanifold [GK]. Let $X = \frac{1}{\sqrt{2}}(A'_{ij} + B'_{ij})$ and $Y = \frac{1}{\sqrt{2}}(C'_{jk} + D'_{jk})$. Then we see $[X, Y] = 0$ and $U(X, Y) = 0$, while $U(X, X) = H_{ij}$ and $U(Y, Y) = H_{jk}$ (here $H_{ij} = X_{ii} - X_{jj} + X_{n+i, n+i} - X_{n+j, n+j}$ in \mathfrak{a}). Thus

$$K(X, Y) = -\langle U(X, X), U(Y, Y) \rangle = -\langle H_{ij}, H_{jk} \rangle > 0.$$

Proposition 3.3. *Let S' be the solvmanifold associated to a solvmanifold corresponding to a non-compact symmetric space of type $\mathbf{SL}(n, \mathbf{H})/\mathbf{Sp}(n)$, in the sense of [GK]. Let $S'_{\Lambda'}$ be an attached solvmanifold to S' in the sense of [T2]. Then $(S'_{\Lambda'}, g')$ is an Einstein solvmanifold.*

For each such symmetric space, there is one associated Einstein solvmanifold, and for each associated solvmanifold, there are approximately $2^{n-1} - 1$ attached solvmanifolds (depending on the number of symmetries of the Dynkin diagram).

Remark. As in the previous family of examples, the compact subgroup of isometries of G/K is $M = \mathbf{SO}(4)$ (where the action permutes A_{jk} , B_{jk} , C_{jk} and D_{jk}). For the associated solvmanifold S' , the analogous compact subgroup of isometries is $M' = \mathbf{SO}(2)\mathbf{SO}(2)$.

REFERENCES

- [Al] D. V. Alekseevskii, *Homogeneous Riemannian spaces of negative curvature*, Mat. Sb. **25** (1975), 87-109; English transl, Math. USSR-Sb. **96** (1975), 93-117.
- [AW1] R. Azencott and E. N. Wilson, *Homogeneous Manifolds with Negative Curvature, I*, Transactions, A.M.S. **215** (1976), 323-362.
- [AW2] R. Azencott and E. N. Wilson, *Homogeneous Manifolds with Negative Curvature, II*, Memoirs, A.M.S. **8** (1976), no. 178.
- [Bes] A. Besse, *Einstein Manifolds*, Springer, 1987.
- [GK] C. S. Gordon and M. M. Kerr, *New homogeneous Einstein metrics of negative Ricci curvature*, Ann. Global Anal. Geom. **19** (2001), no. 1, 75-101.
- [GW] C. S. Gordon and E. N. Wilson, *Isometry Groups of Riemannian Solvmanifolds*, Transactions A.M.S. **307** (1988) no. 1, 245-269.
- [Hb] J. Heber, *Noncompact homogeneous Einstein spaces*, Invent. Math. **133** (1998), 279-352.
- [Hel] S. Helgason, *Differential geometry, Lie groups and symmetric spaces*. Academic Press (1978)
- [J1] M. Jablonski, *Moduli of Einstein and non-Einstein nilradicals*, Geom. Dedicata **152** (2011), 63-84.
- [J2] M. Jablonski, *Concerning the existence of Einstein and Ricci soliton metrics on solvable Lie groups*, Geom. Topol. **15** (2011), no. 2, 735-764.
- [J3] M. Jablonski, *Distinguished orbits of reductive groups*, Rocky Mountain J. Math. **42** (2012), no. 5, 1521-1549.
- [KA] S. Kaneyuki and H. Asano, *Graded Lie algebras and generalized Jordan triple systems*, Nagoya Math. J. **112** (1988), 81-115.
- [Kn] A. Knapp, *Lie Groups: Beyond an Introduction*, Birkhäuser, 1996.
- [L1] J. Lauret, *Einstein solvmanifolds and nilsolitons*, New developments in Lie theory and geometry, 135, Contemp. Math. **491**, Amer. Math. Soc., Providence, RI, 2009.
- [L2] J. Lauret, *Einstein solvmanifolds are standard*, Ann. of Math. (2) **172** (2010), no. 3, 1859-1877.

- [L3] J. Lauret, *Ricci soliton solvmanifolds*, J. Reine Angew. Math. **650** (2011), 1-21.
- [LW] J. Lauret and C. Will, *Einstein solvmanifolds: existence and non-existence questions*, Math. Ann. **350** (2011), no. 1, 199-225.
- [N] Y. Nikolayevsky, *Einstein solvmanifolds and the pre-Einstein derivation*, Trans. Amer. Math. Soc. **363** (2011), no. 8, 3935-3958.
- [P] T. Payne, *The existence of soliton metrics for nilpotent Lie groups*, Geom. Dedicata **145** (2010), 71-88.
- [T1] H. Tamaru, *Homogeneous submanifolds in noncompact symmetric spaces*, Proceedings of the 14th International Workshop on Differential Geometry, **14** (2010), 111-144.
- [T2] H. Tamaru, *Parabolic subgroups of semisimple Lie groups and Einstein solvmanifolds*, Math. Ann. **351** (2011), no. 1, 51-66.
- [T3] H. Tamaru, *Parabolic subgroups and submanifold geometry of noncompact symmetric spaces*, Proceedings of the 15th International Workshop on Differential Geometry, **15** (2011), 29-38.
- [W] T. H. Wolter, *Einstein Metrics on Solvable Groups*, Math. Z. **206** (1991), 457-471.

DEPARTMENT OF MATHEMATICS, WELLESLEY COLLEGE, 106 CENTRAL ST., WELLESLEY, MA 02481
E-mail address: mkerr@wellesley.edu