

On Poisson structures on \mathbb{R}^4

Rubén Flores-Espinoza
Departamento de Matemáticas,
Universidad de Sonora, México

Diciembre 2013

Abstract

This paper is devoted to the study of Poisson structures on the Euclidean four dimensional space \mathbb{R}^4 . By using the properties of the trace operator associated to a volumen form and the elementary vector calculus operations in \mathbb{R}^3 , we give explicit formulas for the main geometric objects associated to the Poisson structures in \mathbb{R}^4 , including its characteristic foliation, the Hamiltonian and Poisson vector fields, normal forms and some useful decomposition formulae for Poisson tensors. We also discuss the class of unimodular Poisson structures and give two results about the existence of Poisson structures having as its characteristic foliation a given arbitrary regular foliation.

1 Introduction

In the last fifty years, Poisson geometry has been an important field of research in differential geometry and mathematical physics for its relevance in the mathematical foundations of classical and quantum mechanics. These geometric structures are associated to the existence of contravariant antisymmetric 2-tensors satisfying the so called Jacobi equation. The Poisson tensors, or solutions of the Jacobi identity, define a generalized integrable vector field distribution in the sense of Sussman [17] decomposing the total space into a family of symplectic manifolds fitting in a smooth manner.

For low dimensional vector spaces, one can find in the literature several papers devoted to the study of Jacobi equations, Poisson geometry and Hamiltonian dynamics. For the tridimensional case, see for example, [5], [7], [8], [13], [21]. In dimension four, the classification of linear and quadratic Poisson tensors has been done in [10] and [16]. In this article, we study from a general point of view the Poisson structures on the four dimensional euclidean space \mathbb{R}^4 . We give explicit formulas for the different quantities and structures associated to a Poisson tensor on \mathbb{R}^4 by using the trace operator properties and the elementary operations of vector calculus. In particular, we study constant rank Poisson structures and those preserving a given volume form. We discuss the automorphisms and infinitesimal symmetries for Poisson tensors, focusing on the

modular Poisson vector field and its role on the preserving volume properties of Hamiltonian vector fields. We include various useful decomposition formulas for Poisson tensors extending some given results in the literature used to classify linear and quadratic Poisson structures. Finally, we show that any regular distribution in the sense of Frobenius corresponds to the symplectic foliation of some Poisson structure. In particular, if the foliation is given by level sets of two independent smooth functions, then the Poisson structure has a maximal basis of infinitesimal automorphisms, transversal to the symplectic foliation.

The paper is organized as follows: In Section 2, we introduce the trace operator on orientable manifolds, discussing its main properties and its relation with the Schouten calculus for contravariant antisymmetric tensor fields. We give explicit formulas for its action on the different contravariant antisymmetric tensors in \mathbb{R}^4 . In Section 3 we introduce the fundamental concepts on Poisson structures and its particular expressions on the vector space \mathbb{R}^4 . We also derive some explicit formulas for the different quantities associated to Poisson tensors on \mathbb{R}^4 , by using the elementary vector calculus operations for three dimensional vector functions. Besides that, we present some important classes of Poisson tensors. In Section 4, we study the group of symmetries of a Poisson tensor and its infinitesimal generators, introducing the modular vector field and the class of unimodular Poisson tensors. In Section 5 we give some interesting decomposition results for Poisson tensors in \mathbb{R}^4 by applying the particular properties of the trace operator on orientable manifolds. Finally, in Section 6, we study the regular Poisson structures on \mathbb{R}^4 and show that any regular 2-dimensional distribution on \mathbb{R}^4 , in the sense of Frobenius, is the characteristic foliation of a regular Poisson structure. Furthermore, in the particular case of foliations given by level sets of independent global smooth functions, the Poisson structure possesses a maximal algebra of transversal Poisson vector fields.

Acknowledgements. This research was suggested by Professor Yu. Vorobiev to whom I express my gratitude. I am grateful also to G. Omelianov, J. Vallejo R., E. Velasco B., J.C. Ruiz P. and G. Dávila-Rascón for support and helpful discussions and comments on this work. This paper was written under grant CB-178690 by Consejo Nacional de Ciencia y Tecnología (México).

2 The trace operator

Let M be a smooth, orientable m -manifold and Ω a volume form on M . For $k = 0, 1, \dots, m$ denote by $V^k(M)$ the space of antisymmetric contravariant k -tensor fields on M . The *trace operator* \mathbf{D} relative to the volume form Ω , is an operator defined on the space of contravariant tensor fields on M . For a k -tensor field $A \in V^k(M)$, $\mathbf{D}(A)$ is the unique $(k - 1)$ -tensor field defined by

$$\mathbf{d}_A \Omega = \mathbf{i}_{\mathbf{D}(A)} \Omega. \quad (1)$$

Here, \mathbf{i}_A is the usual interior product operator on tensor fields. The trace operator was introduced by Koszul in [12] and it is also called the *curl operator* or *divergence operator* [2], [5].

We have the following formulae for the different contravariant tensor fields on M :

- (a) If X is a vector field on M , then $\mathbf{D}(X)$ is the smooth function defined by the relation

$$\mathbf{D}(X)\Omega = \text{di}_X\Omega = L_X\Omega.$$

- (b) For any vector fields X, Y on M , we have

$$\mathbf{D}(X \wedge Y) = [Y, X] + \mathbf{D}(Y)X - \mathbf{D}(X)Y \quad (2)$$

- (c) If $A \in V^p(M)$ and $B \in V^q(M)$, then

$$\mathbf{D}(A \wedge B) = (-1)^q \mathbf{D}(A) \wedge B + A \wedge \mathbf{D}(B) - (-1)^{q+p} [A, B] \quad (3)$$

where $[A, B]$ denotes the Schouten bracket between contravariant anti-symmetric tensors. Recall that for undecomposable p and q -vector fields on M , the Schouten bracket is defined by

$$\begin{aligned} [X_1 \wedge \cdots \wedge X_p, Y_1 \wedge \cdots \wedge Y_q] = & (-1)^{p+1} \sum_{i,j} (-1)^{i+j} [X_i, Y_j] \wedge X_1 \wedge \cdots \\ & \cdots \wedge \widehat{X}_i \wedge \cdots \wedge X_p \wedge Y_1 \wedge \cdots \wedge \widehat{Y}_j \wedge \cdots \wedge Y_q, \end{aligned}$$

where \widehat{X}_i and \widehat{Y}_j are deleted in each summand, see [18].

- (d) For $A \in V^p(M)$ and $B \in V^q(M)$

$$\mathbf{D}([A, B]) = (-1)^q [\mathbf{D}(A), B] + [A, \mathbf{D}(B)]. \quad (4)$$

- (e) The trace operator \mathbf{D} is a cohomology operator

$$\mathbf{D}^2 = 0. \quad (5)$$

If a p -tensor A is in the kernel of the trace operator, we say that A is a *zero trace tensor*. In particular, the flow of a zero-trace vector field preserves the volume form Ω . In the case of a zero trace 2-tensor Λ , the vector field $[f, \Lambda]$ is a zero-trace vector field for each smooth function f . Moreover, the trace of the Schouten bracket of zero-trace tensors is, again, a zero trace tensor.

If we consider another volume form $\widetilde{\Omega} = f\Omega$ with $f \neq 0$, then the corresponding trace operator $\widetilde{\mathbf{D}}$ is related with \mathbf{D} through the following relation

$$\widetilde{\mathbf{D}}(A) = \mathbf{D}(A) - (-1)^p [\ln(|f|), A], \quad \forall A \in V^p(M). \quad (6)$$

For more information about the trace operator see [2], [5], [11].

Remark 1 If $L = \sum_{i=1}^n x^i \frac{\partial}{\partial x^i}$ is the Euler vector field on \mathbb{R}^n and A is a k -homogeneous 2-tensor A on \mathbb{R}^n , then, $[A, L] = (k - 2)A$, and the following decomposition formula holds

$$A = \frac{1}{n + k - 2} (\mathbf{D}(A \wedge L) + \mathbf{D}(A) \wedge L). \quad (7)$$

Decomposition formula (7) has been used in [10] and [16], to classify quadratic and linear Poisson tensors on \mathbb{R}^4 , respectively.

Now, consider \mathbb{R}^4 with global coordinates $(\mathbf{x}, y) = (x_1, x_2, x_3, y)$. In what follows, in order to write down the different geometric quantities on \mathbb{R}^4 , we will use the operations $\times, \cdot, \nabla, \text{div}, \text{rot}$ of elementary vector calculus on the three variables $\mathbf{x} = (x_1, x_2, x_3)$ and for any smooth vector function $X = (X_1, X_2, X_3)$ on \mathbb{R}^3 , we denote by $X \frac{\partial}{\partial \mathbf{x}}$ the vector field

$$X \frac{\partial}{\partial \mathbf{x}} = X_1 \frac{\partial}{\partial x_1} + X_2 \frac{\partial}{\partial x_2} + X_3 \frac{\partial}{\partial x_3},$$

and by $X \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}}$ the 2-tensor

$$X \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} = X_1 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} + X_2 \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial x_1} + X_3 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2},$$

With this notation we give the following formulae for the value of the trace operator \mathbf{D} on \mathbb{R}^4 relative to the canonical volume form

$$\Omega = d\mathbf{x} \wedge dy = dx_1 \wedge dx_2 \wedge dx_3 \wedge dy.$$

(a) For a 4-tensor $A = f \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial y}$,

$$\mathbf{D}A = \frac{\partial f}{\partial y} \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} - \nabla f \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}$$

(b) For a 3-tensor $A = g \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} + \Sigma \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}$, we have

$$\mathbf{D}A = \left(\nabla g + \frac{\partial \Sigma}{\partial y} \right) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} - \text{rot}(\Sigma) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}$$

(c) For a 2-tensor $A = \Psi \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + \Phi \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}$, we have

$$\mathbf{D}(A) = \left(\text{rot}(\Psi) + \frac{\partial \Phi}{\partial y} \right) \frac{\partial}{\partial \mathbf{x}} - \text{div}(\Phi) \frac{\partial}{\partial y}$$

(d) For a vector field $X = W \frac{\partial}{\partial \mathbf{x}} + b \frac{\partial}{\partial y}$

$$\mathbf{D}(X) = \text{div}(W) + \frac{\partial b}{\partial y}$$

3 Poisson structures on \mathbb{R}^4

On \mathbb{R}^4 with global coordinates $(\mathbf{x}, y) = (x_1, x_2, x_3, y)$, any antisymmetric contravariant 2-tensor Λ takes the form

$$\begin{aligned} \Lambda = & \Psi_1 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} + \Psi_2 \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial x_1} + \Psi_3 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} + \\ & + \Phi_1 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial y} + \Phi_2 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial y} + \Phi_3 \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial y}, \end{aligned} \quad (8)$$

for smooth functions Ψ_i, Φ_i on \mathbb{R}^4 and $i = 1, 2, 3$.

If we consider the vector functions $\Psi = (\Psi_1, \Psi_2, \Psi_3)$ and $\Phi = (\Phi_1, \Phi_2, \Phi_3)$, we can identify the 2-tensor Λ with the pair (Ψ, Φ) and write (8) in the simplified notation

$$\Lambda = \Psi \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + \Phi \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}. \quad (9)$$

Each antisymmetric contravariant 2-tensor Λ defines a morphism

$$\Lambda^\# : \bigwedge^1(\mathbb{R}^4) \rightarrow V^1(\mathbb{R}^4)$$

between the space of differential 1-forms and the space of vector fields on \mathbb{R}^4 . The value of $\Lambda^\#$ on a 1-form $\alpha = A dx + b dy$ on \mathbb{R}^4 is the vector field

$$\Lambda^\#(\alpha) = (\Psi \times A - b\Phi) \frac{\partial}{\partial \mathbf{x}} + A \cdot \Phi \frac{\partial}{\partial y} \quad (10)$$

Any contravariant 2-tensor Λ define a bilinear operation on the space of smooth functions on \mathbb{R}^4 with values on the same space of smooth functions given by the bracket

$$\{f, g\} = dg(\Lambda^\#(df)). \quad (11)$$

In terms of the vector functions Ψ and Φ we have

$$\{f, g\} = \Psi \cdot \nabla f \times \nabla g + \Phi \cdot \left(\frac{\partial f}{\partial y} \nabla g - \frac{\partial g}{\partial y} \nabla f \right). \quad (12)$$

A contravariant antisymmetric 2-tensor Λ is called a *Poisson tensor* or a *Poisson structure* on \mathbb{R}^4 if the bracket (11) satisfies the Jacobi identity

$$\{f, \{g, h\}\} + \{h, \{f, g\}\} + \{g, \{h, f\}\} = 0, \quad (13)$$

for any smooth functions f, g, h on \mathbb{R}^4 . In this case, the bracket (11) is called the *Poisson bracket* associated to the Poisson tensor Λ . A 2-tensor Λ satisfies the Jacobi identity (13), if and only if the vector functions (Ψ, Φ) are solutions of the equation

$$\Psi \cdot \left(\text{rot } \Psi + \frac{\partial \Phi}{\partial y} \right) = \frac{\partial}{\partial y}(\Psi \cdot \Phi), \quad (14)$$

$$\Phi \times \left(\text{rot } \Psi + \frac{\partial \Phi}{\partial y} \right) + (\text{div } \Phi) \Psi = \nabla(\Psi \cdot \Phi). \quad (15)$$

Therefore, we can identify the Poisson tensors on \mathbb{R}^4 with solutions of (14), (15).

In a general smooth manifold M , the Jacobi identity (13), is equivalent to the equation

$$[\Lambda, \Lambda] = 0. \quad (16)$$

For orientable manifolds we have, from (3), that a 2-tensor Λ is a Poisson tensor if and only if

$$\mathbf{D}(\Lambda \wedge \Lambda) = 2\Lambda \wedge \mathbf{D}(\Lambda). \quad (17)$$

Moreover, if Λ is a 2-tensor on M and we consider the $(m-2)$ -form ω given by

$$\omega = \mathbf{i}_\Lambda \Omega, \quad (18)$$

then Λ is a Poisson tensor if and only if

$$\mathbf{d}(\mathbf{i}_\Lambda \omega) = 2\mathbf{i}_\Lambda \mathbf{d}\omega. \quad (19)$$

Given a Poisson tensor Λ on \mathbb{R}^4 and a smooth function H , the vector field

$$X_H = \Lambda^\#(\mathbf{d}H) = \left(\Psi \times \nabla H - \frac{\partial H}{\partial y} \Phi \right) \frac{\partial}{\partial \mathbf{x}} + \nabla H \cdot \Phi \frac{\partial}{\partial y}, \quad (20)$$

is called the *Hamiltonian vector field* with *Hamiltonian function* H . Using the properties (3) of the trace operator, the Hamiltonian vector field X_H satisfies

$$\mathbf{i}_{X_H} \Omega = -\mathbf{d}H \wedge \mathbf{i}_\Lambda \Omega.$$

In particular for Poisson structures on \mathbb{R}^4 , the Hamiltonian vector fields with Hamiltonian functions $H_1 = -y$ and $H_2 = \Phi \cdot \Psi$ are, respectively,

$$X_{H_1} = \Phi \frac{\partial}{\partial x} \quad \text{and} \quad X_{H_2} = (\Psi \cdot \Phi) \left(\left(\text{rot}(\Psi) + \frac{\partial \Phi}{\partial y} \right) \frac{\partial}{\partial \mathbf{x}} - \text{div}(\Phi) \frac{\partial}{\partial y} \right).$$

In the case when $\text{div}(\Phi) = 0$, the Hamiltonian vector fields X_{-y} and $X_{\Psi \cdot \Phi}$ commute.

The Jacobi identity for the Poisson bracket (11) endows the space of smooth functions with a Poisson algebra structure, which is homomorphic to the Lie subalgebra of Hamiltonian vector fields

$$[X_f, X_g] = X_{\{f, g\}}, \quad \forall f, g \in C^\infty(\mathbb{R}^4). \quad (21)$$

Thus, the space of Hamiltonian vector fields define an integrable generalized distribution of vector fields in the sense of Sussman [17] and the total space \mathbb{R}^4 is foliated by leaves of (possible different) even dimension. This foliation is called the *characteristic foliation* of Λ and each leaf \mathfrak{L} is a submanifold equipped with a symplectic structure given by the 2-form

$$\omega_{\mathfrak{L}}(X_f, X_g) = \{f, g\} \Big|_{\mathfrak{L}}. \quad (22)$$

The characteristic foliation is also called the *symplectic foliation* of the Poisson structure.

Relative to a given Poisson tensor Λ on \mathbb{R}^4 , a smooth function k is called a *Casimir function* if $\Lambda^\#(dk) = 0$. Moreover, k is a Casimir function if and only if

$$\left(\Psi \times \nabla k - \frac{\partial k}{\partial y} \Phi \right) = 0 \quad \text{and} \quad \nabla k \cdot \Phi = 0. \quad (23)$$

The Poisson bracket of a Casimir function with any other smooth function vanishes. Therefore any Casimir function is a first integral of any vector field $\Lambda^\#(\omega)$, where ω is an arbitrary 1-form in \mathbb{R}^4 .

In terms of the associated 2-form ω (18) k is a Casimir function if and only if

$$dk \wedge \omega = 0. \quad (24)$$

The rank of the Poisson tensor $\Lambda = (\Psi, \Phi)$ is constant on the points of each symplectic leaf. The classification of the symplectic leaves, according to its dimension, is given by the following

Proposition 2 *If $p = (\mathbf{x}, y) \in \mathbb{R}^4$, the rank of a Poisson tensor $\Lambda = (\Psi, \Phi)$ at p , given by (9), takes the following values*

$$\text{rank } \Lambda(p) = \begin{cases} 0 & \text{if and only if } (\Phi^2 + \Psi^2)(p) = 0 \\ 2 & \text{if and only if } (\Phi^2 + \Psi^2)(p) \neq 0 \text{ and } (\Phi \cdot \Psi)(p) = 0 \\ 4 & \text{if and only if } (\Phi \cdot \Psi)(p) \neq 0 \end{cases}$$

Moreover, the characteristic foliation of Λ has the open sets

$$S^+ = \{(\mathbf{x}, u) \in \mathbb{R}^4 \mid \Phi \cdot \Psi > 0\}$$

and

$$S^- = \{(\mathbf{x}, u) \in \mathbb{R}^4 \mid \Phi \cdot \Psi < 0\},$$

as its 4-dimensional symplectic leaves and its boundary

$$\partial S = \{(\mathbf{x}, u) \in \mathbb{R}^4 \mid \Phi \cdot \Psi = 0\},$$

is foliated by 2-dimensional and 0-dimensional symplectic leaves. Notice that a point p belongs to a singular symplectic leaf if $(\Phi \cdot \Psi)(p) = 0$ and $d(\Phi \cdot \Psi)(p) \neq 0$.

Example 3 *The solutions of Jacobi equation with $\Phi = 0$ define a parametrized family of three dimensional Poisson structures where the parameter y can be considered as new variable commuting with all functions.*

Example 4 *The linear solutions (Ψ, Φ) of the Jacobi equation (14), (15) take the general form*

$$\Psi(\mathbf{x}, y) = M\mathbf{x} + p \times \mathbf{x} + y\alpha, \quad (25)$$

$$\Phi(\mathbf{x}, y) = N\mathbf{x} + q \times \mathbf{x} + y\beta, \quad (26)$$

where M and N are symmetric 3×3 matrices and $p, q, \alpha, \beta \in \mathbb{R}^3$ satisfy the equations

$$\begin{aligned} 2Mp &= N\alpha - q \times \alpha \\ 2\alpha \cdot p &= \alpha \cdot \beta \\ (2M + \Lambda \circ \beta)(N + \Lambda \circ q) + (N - \Lambda \circ q)(2M - \Lambda \circ \beta) &= 2 \operatorname{Tr}(N)M \\ \operatorname{Tr}(N)\beta &= N(2p + \beta) + q \times (2p + \beta) \\ (\operatorname{Tr}N)\alpha - M\beta - p \times \beta &= N\alpha - q \times \alpha. \end{aligned}$$

Example 5 *The characteristic foliation of the linear Poisson bracket*

$$\Lambda = 2x_1 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} + \frac{1}{2}x_1 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial y} + \frac{1}{4}x_2 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial y} + \frac{1}{4}x_3 \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial y}$$

consists of:

- (a) *Two 4-dimensional symplectic leaves: $\mathcal{L}_1 = \{(\mathbf{x}, y) \mid x_1 > 0\}$ and $\mathcal{L}_2 = \{(\mathbf{x}, y) \mid x_1 < 0\}$.*
- (b) *The boundary $\{(\mathbf{x}, y) \mid x_1 = 0\}$ is a submanifold foliated by 2-dimensional symplectic leaves of the form $\{x_1 = 0, \quad bx_2 - cx_3 = 0, \quad bc \neq 0\}$, and zero dimensional leaves given by the points $(\mathbf{0}, y)$.*

In particular, the 2-dimensional symplectic leaves are given by intersection of the level set of $k = \frac{x_2}{x_3}$ with $x_1 = 0$.

The Hamiltonian vector fields take the form

$$\begin{aligned} X_f &= \left(-2x_1 \frac{\partial f}{\partial x_3} + \frac{1}{4}x_2 \frac{\partial f}{\partial y} \right) \frac{\partial}{\partial x_2} + \left(2x_1 \frac{\partial f}{\partial x_2} + \frac{1}{4}x_3 \frac{\partial f}{\partial y} \right) \frac{\partial}{\partial x_3} \\ &\quad + \left(\frac{1}{2}x_1 \frac{\partial f}{\partial x_1} + \frac{1}{4}x_2 \frac{\partial f}{\partial x_2} + \frac{1}{4}x_3 \frac{\partial f}{\partial x_3} \right) \frac{\partial}{\partial y}. \end{aligned}$$

If Λ is a Poisson tensor on \mathbb{R}^4 has a smooth global Casimir function $k : \mathbb{R}^4 \rightarrow \mathbb{R}$, we have

$$dk \wedge \mathbf{i}_\Lambda \Omega = -\mathbf{i}_{[k, \Lambda]} \Omega = 0,$$

and there exists a 1-form θ on \mathbb{R}^4 with

$$dk \wedge \theta = \mathbf{i}_\Lambda \Omega. \tag{27}$$

In this case, the Jacobi identity becomes

$$d\theta \wedge \theta \wedge dk = 0. \tag{28}$$

Moreover, if we consider the 3-tensor T given by

$$\theta = \mathbf{i}_T \Omega,$$

then Λ takes the form

$$\Lambda = [k, T]. \quad (29)$$

In global coordinates if $\theta = -A dx + f dy$, with $A = A(\mathbf{x}, y)$ a smooth vector function and f a smooth real function, then the Poisson 2-tensor Λ in (29) takes the form

$$\Lambda = \left(f \nabla k + \frac{\partial k}{\partial y} A \right) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + (A \times \nabla k) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}, \quad (30)$$

and the condition (28) becomes

$$\left(\left(\nabla f + \frac{\partial A}{\partial y} \right) \times A - f \operatorname{rot}(A) \right) \cdot \nabla k - \frac{\partial k}{\partial y} (A \cdot \operatorname{rot}(A)) = 0. \quad (31)$$

Similarly, for those Poisson structures Λ having 2 independent global Casimir functions k_1, k_2 , we have

$$f dk_1 \wedge dk_2 = \mathbf{i}_\Lambda \Omega$$

for some smooth function f . In this case, the Poisson tensor Λ has the form

$$\Lambda = f \left(\frac{\partial k_2}{\partial y} \nabla k_1 - \frac{\partial k_1}{\partial y} \nabla k_2 \right) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + f (\nabla k_1 \times \nabla k_2) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}. \quad (32)$$

Example 6 *The linear Poisson tensors Λ having as Casimir function the homogeneous polynomial of degree two $k_1 = \frac{1}{2} \mathbf{x}^T P \mathbf{x} + (\alpha \cdot \mathbf{x}) y + \frac{1}{2} b y^2$, with $P = P^T$, $\alpha \in \mathbb{R}^3$ and $b \in \mathbb{R}$, take the form*

$$\Lambda = (c(P \mathbf{x} + y \alpha) + (\alpha \cdot \mathbf{x} + b y) A) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + (A \times (P \mathbf{x} + y \alpha)) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}$$

where $A \in \mathbb{R}^3$, $c \in \mathbb{R}$. In this case, we have another global Casimir function k_2 given by the linear function

$$k_2 = A \cdot \mathbf{x} - c y$$

For the general theory on Poisson structures, the reader can consult the papers by A. Lichnerowicz [14] and A. Weinstein [19], or the books [5], [9], [13], [18].

4 Automorphisms of Poisson structures on \mathbb{R}^4

Consider a Poisson tensor $\Lambda = (\Psi, \Phi)$ on \mathbb{R}^4 . A diffeomorphism $F : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ with $F(\mathbf{x}, y) = (S(\mathbf{x}, y), h(\mathbf{x}, y))$, where $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ and $h : \mathbb{R}^4 \rightarrow \mathbb{R}$ are smooth functions, is called a *Poisson map* if it preserves the Poisson tensor or, equivalently, preserves the Poisson bracket

$$\{f, g\} \circ F = \{f \circ F, g \circ F\}, \quad \forall f, g \in C^\infty(\mathbb{R}^4). \quad (33)$$

In terms of the vector functions (Ψ, Φ) associated to the Poisson tensor Λ , the condition (33) is written as

$$\Psi(S(\mathbf{x}, y), h(\mathbf{x}, y)) = \det(D_{\mathbf{x}}S)(D_{\mathbf{x}}^{-1}S)^T(\Psi) + D_{\mathbf{x}}S(\Phi) \times \frac{\partial S}{\partial y}, \quad (34)$$

$$\Phi(S(\mathbf{x}, y), h(\mathbf{x}, y)) = -D_{\mathbf{x}}S(\Psi \times \nabla_{\mathbf{x}}h) + \frac{\partial h}{\partial y}D_{\mathbf{x}}S(\Phi) - (\Phi \cdot \nabla_{\mathbf{x}}h)\frac{\partial S}{\partial y}. \quad (35)$$

Here, $D_{\mathbf{x}}S$ denotes the differential of S with respect to the variable $\mathbf{x} = (x_1, x_2, x_3)$.

Remark that expression (35) is equivalent to

$$\Phi(S(\mathbf{x}, y), h(\mathbf{x}, y)) = -DF(X_h), \quad (36)$$

where X_h is the Hamiltonian vector field corresponding to function $h(\mathbf{x}, y)$ relative to the Poisson structure (Ψ, Φ) . Notice that the function $\Phi \cdot \Psi$ is transformed into

$$(\Phi \cdot \Psi) \circ F = (\det DF)(\Phi \cdot \Psi).$$

In general, applying the transformation formulae (34), (35), the Poisson tensors having a global Casimir function can be transformed to a normal form, as stated by the following result.

Proposition 7 *Any Poisson tensor $\Lambda = (\Psi, \Phi)$ on \mathbb{R}^4 having a global Casimir function k with $\frac{\partial k}{\partial y} \neq 0$ is transformed, under the diffeomorphism $F : \mathbb{R}^4 \rightarrow \mathbb{R}^4$, with $F(\mathbf{x}, y) = (\mathbf{x}, k(\mathbf{x}, y))$, into the normal form*

$$\tilde{\Lambda} = \tilde{\Psi}(\mathbf{x}, y)\frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}}. \quad (37)$$

The constructions of Poisson brackets on smooth manifolds with prescribed Casimir functions has been studied in [3].

Given a Poisson tensor Λ , a vector field $X = W\frac{\partial}{\partial \mathbf{x}} + b\frac{\partial}{\partial y}$ preserving the Poisson structure is called an *infinitesimal automorphism* of Λ or *Poisson vector field*. In terms of the Poisson bracket, X is a Poisson vector field if

$$L_X \{f, g\} = \{L_X f, g\} + \{f, L_X g\}, \quad (38)$$

for any smooth functions $f, g \in C^\infty(\mathbb{R})$, or equivalently,

$$[X, \Lambda] = 0. \quad (39)$$

In terms of the trace operator, condition (39) reads

$$\mathbf{D}(\Lambda \wedge X) + \mathbf{D}(\Lambda) \wedge X - \mathbf{D}(X) \wedge \Lambda = 0. \quad (40)$$

Moreover, if Λ is a Poisson tensor and $\omega = \mathbf{i}_\Lambda \Omega$, as in (18), then a vector field X is a Poisson vector field if and only if

$$L_X \omega = \mathbf{D}(X)\omega. \quad (41)$$

In \mathbb{R}^4 with global coordinates (\mathbf{x}, y) , the vector field $X = W \frac{\partial}{\partial \mathbf{x}} + b \frac{\partial}{\partial y}$ is a Poisson vector field if and only if

$$\nabla(\Psi \cdot W) - (\operatorname{div} W)\Psi - W \times \operatorname{rot} \Psi + b \frac{\partial \Psi}{\partial y} + \frac{\partial W}{\partial y} \times \Phi = 0, \quad (42)$$

$$\operatorname{rot}(\Phi \times W) - \operatorname{div}(W)\Phi + \operatorname{div}(\Phi)W + \Psi \times \nabla b - \frac{\partial b}{\partial y}\Phi + b \frac{\partial \Phi}{\partial y} = 0. \quad (43)$$

The space of Poisson vector fields is a Lie algebra of vector fields.

For each Poisson structure Λ on an oriented manifold (M, Ω) , the vector field $Z_\Lambda = \mathbf{D}(\Lambda)$ satisfies

$$[Z_\Lambda, \Lambda] = 0. \quad (44)$$

Thus Z_Λ is a Poisson vector field called the *modular vector field*. If we take on M another volume form $\tilde{\Omega} = f \Omega_0$ with $f \neq 0$, the corresponding modular vector field \tilde{Z}_Λ differs from Z_Λ by a Hamiltonian vector field,

$$\tilde{Z}_\Lambda - Z_\Lambda = X_{-\ln(|f|)}.$$

If Λ is a Poisson tensor on \mathbb{R}^4 (8), the modular vector field Z_Λ takes the form

$$Z_\Lambda = \left(\operatorname{rot}(\Psi) + \frac{\partial \Phi}{\partial y} \right) \frac{\partial}{\partial \mathbf{x}} - \operatorname{div} \Phi \frac{\partial}{\partial y}. \quad (45)$$

Proposition 8 *The modular tensor field $Z_\Lambda = \mathbf{D}(\Lambda)$, associated to a Poisson tensor Λ on an orientable manifold M , has the following properties:*

1. *For each Hamiltonian vector field X_H we have:*

$$\mathbf{D}(X_H) = L_{Z_\Lambda} H, \quad (46)$$

$$[Z_\Lambda, X_H,] = X_{\mathbf{D}(X_H)} \quad (47)$$

where $L_{Z_\Lambda} H$ denotes the Lie derivative of H along Z_Λ .

2. *The Lie bracket of a Poisson vector field W with the modular vector field Z_Λ is a Hamiltonian vector field with Hamiltonian function given by the trace of W ,*

$$[W, Z_\Lambda] = X_{\mathbf{D}(W)}. \quad (48)$$

From Proposition 8 we deduce the following facts:

- (a) If the modular vector field Z_Λ vanishes, then any Hamiltonian vector field has zero trace. Reciprocally, if all Hamiltonian vector fields have zero-trace, then the modular vector field vanishes.
- (b) From (48), the Lie bracket of Z_Λ with any other Poisson vector field W is a Hamiltonian vector field. The trace of any Poisson vector field commuting with the modular vector field Z_Λ is a Casimir function.

For Poisson tensors on \mathbb{R}^4 , the modular vector field Z_Λ has the following additional properties:

- (a) Z_Λ is tangent to the level sets of $\Phi \cdot \Psi$.
- (b) The formula $\Lambda^\#(d(\Phi \cdot \Psi)) = (\Phi \cdot \Psi)Z_\Lambda$ holds and for each Hamiltonian vector field with Hamiltonian function H , we have

$$(\Phi \cdot \Psi)L_{Z_\Lambda}H = -L_{X_H}(\Phi \cdot \Psi). \quad (49)$$

Therefore, if H is a first integral of the modular vector field Z_Λ , the Hamiltonian vector field X_H is tangent to the level set of the function $\Phi \cdot \Psi$. On the contrary, at any point $p \in \mathbb{R}^4$ where $(\Phi \cdot \Psi)(p) \neq 0$ and $L_{Z_\Lambda}H(p) \neq 0$, the Hamiltonian vector field X_H is transversal to the level set of $\Phi \cdot \Psi$.

On a general orientable manifold M , a Poisson tensor Λ is called *unimodular* if for some volume form Ω , the modular vector field Z_Λ vanishes. The unimodular Poisson tensors on \mathbb{R}^4 take the form

$$\Lambda = \mathbf{D}(T) \quad (50)$$

where T is a 3-contravariant tensor

$$T = f \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_2} + \Sigma \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}, \quad (51)$$

where Σ is a smooth vector function and f a smooth function satisfying

$$\left(\nabla f + \frac{\partial \Sigma}{\partial y} \right) \cdot \text{rot}(\Sigma) = c, \quad (52)$$

with c a constant.

Considering the 1-form θ defined by

$$\theta = \mathbf{i}_T \Omega = -\Sigma d\mathbf{x} + f dy, \quad (53)$$

we have

$$d\theta = \mathbf{i}_\Lambda \Omega,$$

and condition (52) is equivalent to

$$d\theta \wedge d\theta = c\Omega.$$

If the constant c is non-zero, the unimodular Poisson tensor is symplectic. If $c = 0$, then Λ is a rank 2, regular Poisson tensor. The unimodular Poisson tensors Λ on \mathbb{R}^4 take the form

$$\Lambda = \left(\nabla f + \frac{\partial \Sigma}{\partial y} \right) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} - \text{rot}(\Sigma) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}. \quad (54)$$

The unimodular Poisson tensors Λ on \mathbb{R}^4 have the following properties:

- (a) The 2-form $\omega = \mathbf{i}_\Lambda \Omega$ is exact.
- (b) The Poisson vector fields tangent to the symplectic foliation have zero trace.
- (c) The trace of a Poisson vector field transversal to symplectic leaves is a Casimir function.
- (d) For any 2-tensor Θ commuting with the unimodular Poisson tensor Λ , the vector field $\mathbf{D}(\Theta)$ is a Poisson vector field.

An example of unimodular Poisson tensors are those in \mathbb{R}^4 of the form

$$\Lambda = \nabla G(\mathbf{x}, y) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}}$$

Here, we have two global Casimir functions $k_1(\mathbf{x}, y) = y$ and $k_2 = G(\mathbf{x}, y)$.

Example 9 In \mathbb{R}^4 with the canonical volume form, the quadratic Poisson tensor [10]

$$\Lambda = -x_1 x_2 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_3} + x_1 x_3 \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial x_1} - y x_1 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial y} - y x_2 \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial y} + y x_3 \frac{\partial}{\partial x_3} \wedge \frac{\partial}{\partial y}$$

has a modular Poisson vector field given by

$$Z_\Lambda = -2x_1 \frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_2} + (2x_3 + x_1) \frac{\partial}{\partial x_3} + y \frac{\partial}{\partial y}. \quad (55)$$

In this case, we have two open 4-dimensional symplectic leaves given by $\mathcal{L} = \{(\mathbf{x}, y) \mid y x_1 x_2 (x_1 - x_3) > 0 (< 0)\}$. The set of rank-two points (\mathbf{x}, y) is

$$\begin{aligned} & \{(\mathbf{x}, y) \mid y x_1 = 0, y^2 + x_1^2 \neq 0, x_2 x_3 \neq 0\} \cup \{(\mathbf{x}, y) \mid y x_1 \neq 0, x_2 = 0\} \\ & \cup \{(\mathbf{x}, y) \mid y x_1 \neq 0, x_1 = x_3\}, \end{aligned}$$

and the zero dimensional leaves are given by the points $\{(\mathbf{x}, y) \mid y^2 + x_1^2 = 0\} \cup \{(\mathbf{x}, y) \mid \mathbf{x} = \mathbf{0}\} \cup \{(\mathbf{x}, y) \mid x_2 = x_3 = y = 0\}$.

The unimodular Poisson structures has been studied by A. Weinstein in [20].

5 Decomposition of Poisson structures

In this section, we present some useful results on the decomposition of Poisson brackets on \mathbb{R}^4 as a sum of some special 2-tensors. We recall that a pair of contravariant antisymmetric 2-tensors A and B is called a *Poisson pair* if both are commuting Poisson tensors, that is $[A, A] = [B, B] = 0$ and $[A, B] = 0$. In this case, for any scalars $\lambda, \beta \in \mathbb{R}$, the 2-tensor $\lambda A + \beta B$ is again a Poisson tensor.

Proposition 10 Let $\Lambda = (\Psi, \Phi)$ be a Poisson tensor on \mathbb{R}^4 and $\alpha = A dx + a dy$ and $\beta = B dx + b dy$ two independent 1-forms such that $d\alpha \wedge \alpha \wedge \beta = d\beta \wedge \alpha \wedge \beta = 0$ and $\beta(\Lambda^\#(\alpha)) = 1$. Then Λ has the following decomposition

$$\Lambda = \Lambda^\#(\alpha) \wedge \Lambda^\#(\beta) - (\Psi \cdot \Phi) S_{(\alpha, \beta)}, \quad (56)$$

where $S_{(\alpha, \beta)}$ is the rank-two Poisson tensor given by

$$S_{(\alpha, \beta)} = (bA - aB) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + (A \times B) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}. \quad (57)$$

In particular, if there exists independent smooth functions f, g with $\{f, g\} \neq 0$, then we have the decomposition formula

$$\Lambda = \frac{1}{\{f, g\}} X_f \wedge X_g - \frac{\Psi \cdot \Phi}{\{f, g\}} S_{(df, dg)}, \quad (58)$$

where

$$S_{(df, dg)} = \left(\frac{\partial g}{\partial y} \nabla f - \frac{\partial f}{\partial y} \nabla g \right) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + (\nabla f \times \nabla g) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}. \quad (59)$$

Remark 11 Notice that in (58), the 2-tensor $S_{(df, dg)}$ is also a Poisson tensor and $S_{(df, dg)}(df) = S_{(df, dg)}(dg) = 0$. Moreover, if a Hamiltonian vector field X_f is non-zero in some point p , then there exists a function h with $\{f, h\} = 1$, and $[X_f, X_h] = 0$ in some neighborhood of p . In this case, we have in the decomposition formula (58), that the tensor $X_f \wedge X_h$ is a Poisson tensor. Thus, locally, we have a decomposition of Λ as a sum of a Poisson pair in the form

$$\Lambda = X_f \wedge X_h - (\Psi \cdot \Phi) S_{(df, dh)}. \quad (60)$$

and we recover the classical local decomposition theorem by Weinstein [19].

Another type of decompositions of Poisson tensors on \mathbb{R}^4 are the following:

Proposition 12 For each Poisson tensor $\Lambda = (\Psi, \Phi)$ on \mathbb{R}^4 , as in (8), with $(\operatorname{div} \Phi) \neq 0$, we have the decomposition

$$\Lambda = Z_\Lambda \wedge \left(\frac{\Phi}{\operatorname{div}(\Phi)} \right) \frac{\partial}{\partial \mathbf{x}} + \Lambda_0, \quad (61)$$

where

$$\Lambda_0 = \frac{1}{\operatorname{div}(\Phi)} \nabla(\Phi \cdot \Psi) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}}$$

is a Poisson tensor of constant rank equal to 2.

Corollary 13 If the Poisson tensor is linear with (Ψ, Φ) given by (25), (26) and $\operatorname{div} \Phi \neq 0$, then the decomposition formula (61) can be rewritten as

$$\Lambda = Z_\Lambda \wedge \Sigma \frac{\partial}{\partial \mathbf{x}} + \nabla(\Sigma \cdot \Psi) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}}, \quad (62)$$

with $\Sigma = \frac{1}{\operatorname{div} \Phi} \Phi$. In this case, the two 2-tensors $A = Z_\Lambda \wedge \Sigma \frac{\partial}{\partial \mathbf{x}}$ and $B = \nabla(\Sigma \cdot \Psi) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}}$ are a Poisson pair.

For linear and quadratic Poisson tensors on \mathbb{R}^4 , we recall the following results given in [16] and [15], [21], respectively:

For linear Poisson tensor on a finite-dimensional vector space, applying formula (3) we have

$$\Lambda = \frac{1}{3}(\mathbf{D}(\Lambda \wedge L) + Z_\Lambda \wedge L)$$

where L denotes the *Euler vector field* $L = \sum_{i=1}^n x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y}$.

If the Poisson tensor Λ is quadratic, then $[\Lambda, L] = 0$ and applying directly formula (3) we obtain

$$\Lambda = \frac{1}{4}(\mathbf{D}(\Lambda \wedge L) + Z_\Lambda \wedge L). \quad (63)$$

In the quadratic case L and Z_Λ are commuting Poisson vector fields of Λ . Moreover, $Z_\Lambda \wedge L$ and $\mathbf{D}(\Lambda \wedge L)$ are commuting Poisson tensors and then, we have a decomposition of Λ as a sum of a Poisson pair, one of them with zero trace. Applying the decomposition formula (63), the quadratic Poisson tensors on \mathbb{R}^4 take the form

$$\Lambda = \mathbf{D}(T) + \frac{1}{4}X \wedge L,$$

where X is a linear vector field and T is cubic 3-tensor on \mathbb{R}^4 satisfying the conditions

$$[\mathbf{D}(T), \mathbf{D}(T)] = 0, \quad [\mathbf{D}(T), X] = 0, \quad \mathbf{D}(X) = 0. \quad (64)$$

If we consider the 1-form $\theta = \mathbf{i}_T \Omega$, conditions (64) can be written in the equivalent form

$$d\theta \wedge d\theta = 0, \quad d(L_X \theta) = 0, \quad \mathbf{D}(X) = 0, \quad (65)$$

Therefore, θ and X , satisfying (65), parametrize the quadratic Poisson tensors on \mathbb{R}^4 .

6 Regular Poisson structures on \mathbb{R}^4

A Poisson tensor is called *regular* if it has a constant rank. It means that its characteristic foliation is a regular foliation in the sense of Frobenius. For regular Poisson tensors on \mathbb{R}^4 we have two cases: one, when the rank of the Poisson tensor is 4 and the foliation consists of only one leaf, and the other one, when each symplectic leaf is two-dimensional.

The first case correspond to symplectic Poisson tensors. From Jacobi conditions (14) and (15), a 2-contravariant tensor $\Lambda = (\Psi, \Phi)$ defines a symplectic

structure on \mathbb{R}^4 if and only if

$$(\Phi \cdot \Psi) \neq 0, \quad (66)$$

$$\operatorname{div} \left(\frac{\Phi}{\Phi \cdot \Psi} \right) = 0, \quad (67)$$

$$\frac{\partial}{\partial y} \left(\frac{\Phi}{\Phi \cdot \Psi} \right) + \operatorname{rot} \left(\frac{\Psi}{\Phi \cdot \Psi} \right) = 0. \quad (68)$$

Thus, from conditions (66)-(68), the contravariant 2-tensor $\Lambda = (\Psi, \Phi)$ defines a symplectic structure on \mathbb{R}^4 if and only if $\Phi \cdot \Psi \neq 0$. In this case, the 2-tensor $\frac{1}{\Phi \cdot \Psi} \Lambda$ has zero trace and, on simply connected domains, tensor Λ takes the form $\Lambda = (\Phi \cdot \Psi) \mathbf{D}(T)$, where T is some contravariant 3-tensor. Here, the modular vector field Z_Λ is the Hamiltonian vector field with Hamiltonian function $H = \ln(|\Phi \cdot \Psi|)$.

Now, let us consider the Poisson structures Λ on \mathbb{R}^4 of rank equal 2. In this case, we have $\Lambda(\mathbf{x}, y) \neq 0$, for all $(\mathbf{x}, y) \in \mathbb{R}^4$ and

$$\begin{aligned} \Lambda \wedge \Lambda &= 0, \\ \Lambda \wedge \mathbf{D}(\Lambda) &= 0. \end{aligned}$$

In global coordinates (\mathbf{x}, y) , the regular Poisson tensors Λ of rank 2 takes the form (8) with $\Psi \cdot \Phi = 0$ and $\Psi^2 + \Phi^2 \neq 0$. The Jacobi identity reduces to equations

$$\begin{aligned} \Psi \cdot \left(\operatorname{rot} \Psi + \frac{\partial \Phi}{\partial y} \right) &= 0, \\ \Phi \times \left(\operatorname{rot}(\Psi) + \frac{\partial \Phi}{\partial y} \right) + (\operatorname{div} \Phi) \Psi &= 0. \end{aligned}$$

Notice that if Λ is a regular Poisson structure of rank 2, then for any smooth function $\lambda \neq 0$, the tensor $\lambda \Lambda$ is also a regular Poisson structure having the same characteristic foliation. Moreover, for any smooth functions f, g with $\{f, g\} \neq 0$ we have the following representation,

$$\Lambda = \frac{1}{\{f, g\}} X_f \wedge X_g. \quad (69)$$

In particular, in the open set $\Phi \cdot \mathbf{x} \neq 0$ we have

$$\Lambda = \left(\frac{-1}{\Phi \cdot \mathbf{x}} \right) X_{(\frac{1}{2}\mathbf{x}^2)} \wedge X_y. \quad (70)$$

In the open domain where $\Phi \neq 0$, the regular Poisson tensor Λ takes the form

$$\begin{aligned} \Lambda &= (\Phi \times \Sigma) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + \Phi \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y} \\ &= \Phi \frac{\partial}{\partial \mathbf{x}} \wedge \left(\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y} \right), \end{aligned} \quad (71)$$

for some vector function Σ and the Jacobi identity reduces to the single equation

$$\Phi \frac{\partial}{\partial \mathbf{x}} \wedge \left[\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y}, \Phi \frac{\partial}{\partial \mathbf{x}} \right] = 0. \quad (72)$$

The Hamiltonian vector field X_H , at each point, belongs to the distribution generated by the vector fields $\Phi \frac{\partial}{\partial \mathbf{x}}$ and $\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y}$ and takes the form

$$X_H = - \left(L_{\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y}} H \right) \Phi \frac{\partial}{\partial \mathbf{x}} + \left(L_{\Phi \frac{\partial}{\partial \mathbf{x}}} H \right) \left(\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y} \right). \quad (73)$$

The foliated symplectic form ω is

$$\omega = \frac{1}{\Phi^2 + (\Phi \times \Sigma)^2} (\Phi d\mathbf{x}) \wedge (\Sigma d\mathbf{x} + dy), \quad (74)$$

and the modular vector field Z_Λ (45) is given by

$$Z_\Lambda = \left(\left[\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y}, \Phi \frac{\partial}{\partial \mathbf{x}} \right] - \operatorname{div}(\Phi) \left(\Sigma \frac{\partial}{\partial \mathbf{x}} + \frac{\partial}{\partial y} \right) + \operatorname{div}(\Sigma) \Phi \frac{\partial}{\partial \mathbf{x}} \right). \quad (75)$$

Taking into account (75) we verify that Z_Λ is always a tangent Poisson vector field to the symplectic leaves. Notice that, on the domain where $\operatorname{div}(\Phi) \neq 0$, formula (75) allow us to write down the Poisson tensor (71) in the form

$$\Lambda = \frac{1}{\operatorname{div}(\Phi)} Z_\Lambda \wedge \left(\Phi \frac{\partial}{\partial \mathbf{x}} \right). \quad (76)$$

Any rank two regular foliation on \mathbb{R}^4 is generated by two independent 1-forms α, β , with $\alpha \wedge \beta \neq 0$, and satisfying the integrability conditions

$$d\alpha \wedge \alpha \wedge \beta = 0, \quad (77)$$

$$d\beta \wedge \alpha \wedge \beta = 0. \quad (78)$$

In this case, the 2-tensor Λ defined by the the relation

$$\alpha \wedge \beta = \mathbf{i}_\Lambda \Omega, \quad (79)$$

is a regular Poisson tensor.

Furthermore, using the local expressions for α, β given by $\alpha = A d\mathbf{x} + a dy$ and $\beta = B d\mathbf{x} + b dy$, the 2-tensor field Λ in (79) takes the form

$$\Lambda = (bA - aB) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial \mathbf{x}} + (A \times B) \frac{\partial}{\partial \mathbf{x}} \wedge \frac{\partial}{\partial y}, \quad (80)$$

If $\alpha \wedge \beta$ is a closed 2-form, then the modular vector $\mathbf{D}(\Lambda)$ vanishes and the Poisson structure Λ is unimodular.

We summarize the previous discussion in the following result.

Proposition 14 *Given a 2-dimensional integrable distribution \mathfrak{F} on \mathbb{R}^4 generated by two independent 1-forms α and β satisfying (77)-(78), there exists a Poisson structure Λ having \mathfrak{F} as its characteristic foliation.*

For a regular foliation defined by the level sets of two independent functions f and g , we can improve Proposition 14 constructing a Poisson structure which has the additional property of possessing a basis of vector fields, transversal to the foliation. This kind of structures are called *transversally constant Poisson structures* [18].

Given two independent smooth functions f, g , we are going to construct a Poisson structure with the above properties, in the following way: Take a symplectic Poisson tensor $\Lambda = (\Psi, \Phi)$ on \mathbb{R}^4 with $\{f, g\} \neq 0$; then, consider the 2-tensor Δ given by

$$\Delta = \Lambda - \frac{1}{\{f, g\}} X_f \wedge X_g. \quad (81)$$

The 2-tensor Δ in (81) is a regular Poisson having f and g as its Casimir functions and the vector fields

$$W_1 = \frac{1}{\{f, g\}} X_f, \quad W_2 = \frac{1}{\{f, g\}} X_g, \quad (82)$$

as independent Poisson vector fields, transversal to its symplectic leaves. The tensor Δ in (81) is called the *Dirac-Poisson tensor associated to Λ and constraint functions f and g* .

Taking into account the decomposition (58), the Poisson tensor Δ takes the form

$$\Delta = -\frac{\Psi \cdot \Phi}{\{f, g\}} S_{(df, dg)},$$

and then, Δ is a regular Poisson structures having the desired properties.

From the above discussion, given any regular foliation defined by the level sets of two independent smooth functions f and g we can take any symplectic structure Λ and multiply the 2-tensor $S_{(df, dg)}$ by the factor $\lambda = -\frac{\Psi \cdot \Phi}{\{f, g\}}$ to have a transversally maximal Poisson structure whose characteristic foliation coincides with the level sets of f and g . Notice that, in general, the constructed tensor Δ is not unimodular.

Proposition 15 *For any regular foliation \mathfrak{F} of rank 2, defined as the level sets of two independent smooth functions f and g , there exists a transversally constant regular Poisson structure, having \mathfrak{F} as its characteristic foliation.*

References

- [1] Ay A, Gurses M, and Zheltukin K., Hamiltonian equations in \mathbb{R}^3 , *J. Math. Phys.* **44**, (2003) 5688-5705
- [2] Cariñena J.F., Ibort, A., Marmo G., Perelomov A. On the geometry of Lie algebras and Poisson tensors., *J. Phys. A: Math. Gen.* **27** (1994) 7425-7449.

- [3] Daminou A. P, and Petalidou F. Poisson brackets with prescribed Casimirs.,arXiv:1103.0849v1 [Math. DG] 4 Mar 2011
- [4] Dufour J.P. and Haraki A. Rotationnels et structures de Poisson quadratiques, C.R. Acad. Sci. Paris Ser. I, Math. **312** (1991), no.1, 137-140
- [5] Dufour J.-P. and Zung N.T. *Poisson structures and their normal forms*, Progress in Mathematics 242, Birkhauser Verlag, Basel (2005)
- [6] Gumral H., Nutku Y., Poisson structure of dynamical systems with three degree of freedom. *J. Math. Phys.* **34**, 5691 (1993)
- [7] Hernandez-Bermejo B. New solutions of the Jacobi equations for three dimensional Poisson structures, *J. Math. Phys.* **42**, (2001) 4984-4996
- [8] Hernandez-Bermejo B., New solutions family of the Jacobi equations: Characterization, invariants, and global Darboux analysis, *J. Math. Phys.* **48** (2007) 022903-022914
- [9] Karasev, M. V. and Maslov, V. P., *Nonlinear Poisson brackets: Geometry and Quantization*, Translations of Mathematical Monographs, V. 119, Amer. Math. Soc., Providence, 1993.
- [10] Klinker F., Quadratic Poisson structures in dimension four, www.Mathematik.uni-dortmund.de/~klinker/Paper/PoissonQuadraticFour_Klinker.pdf
- [11] Klinker F., Polynomial Poly-vector fields, arXiv:math/0409157v4 [math. DG] Feb. 2008
- [12] Koszul, J.L., Crochets de Schouten-Nijenhuis et cohomologie, *Asterisque, Soc. Math. de France, hors serie, (1985), 257-271*
- [13] Laurent-Gengoux C., Pichereau A. and Vanhaeaecke P. *Poisson Structures*, Springer Verlag Berlin Heidelberg 2013.
- [14] Lichnerowicz, A., Les varietes de Poisson et leur algebres associes. *J. Diff. Geometry.* **12** (1977) 253-300
- [15] Lin, Q., Liu, Z.-J., and Sheng, Y.-H. Quadratic deformations of Lie-Poisson structures, *Lett. Math. Phys.* (2008) 83:217-299
- [16] Sheng. Y., Linear Poisson structures on \mathbb{R}^4 . *Journal of Geometry and Physics* 57 (11) 2398-2410 (2007)
- [17] Sussmann, H.J., Orbits of families of vector fields and integrability of distributions. Transactions Amer. Math. Soc. 170 (1973), 171-188
- [18] Vaismann, I., *Lectures on the Geometry of Poisson Manifolds*, Birkhauser, Basel , 1994.
- [19] Weinstein, A., The local structure of Poisson manifolds, *J. Diff. Geom.* **18** (1983), 523-557

- [20] Weinstein, A., The modular automorphism group of a Poisson manifold, *J. Geom. Phys.* **26** (1997), 379-394
- [21] Zhang Ju Liu and Xu Ping., On quadratic Poisson structures. *Lett. Math. Phys.* **26** (1) (1992), 33-42