

# Primitive Central Idempotents of Rational Group Algebras

Geoffrey Janssens

## Abstract

We give a description of the primitive central idempotents of the rational group algebra  $\mathbb{Q}G$  of a finite group  $G$ . Such a description is already investigated by Jespers, Olteanu and del Río, but some unknown scalars are involved. Our description also gives answers to their questions.

Let  $G$  be a finite group. The complex group algebra  $\mathbb{C}G$  is semisimple and a description of its primitive central idempotents is well known. These are the elements  $e(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(1)\chi(g^{-1})g$ , where  $\chi$  runs through the irreducible characters of  $G$ . Using Galois descent one obtains that the primitive central idempotents of the semisimple rational group algebra  $\mathbb{Q}G$  are the elements  $e_{\mathbb{Q}}(\chi) = \sum_{\sigma \in G_{\chi}} \sigma(e(\chi))$ , with  $G_{\chi} = \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})$ . Rather recently, Olivieri et al. [3] obtained a character free method for describing the primitive central idempotents of  $\mathbb{Q}G$  provided that  $G$  is a finite monomial group. Their method relies on a theorem of Shoda on pairs of subgroups  $(H, K)$  of  $G$  with  $K$  normal in  $H$ ,  $H/K$  abelian and so that an irreducible linear character of  $H$  with kernel  $K$  induces an irreducible character of  $G$ . Such pairs are called Shoda pairs. The main ingredient in this theory are the elements  $\epsilon(H, K)$  of  $\mathbb{Q}H$  with  $K \triangleleft H \leq G$ . These are defined as  $\prod_{M/K \in \mathcal{M}(H/K)} (\tilde{K} - \tilde{M})$  if  $H \neq K$  and as  $\tilde{H}$  if  $H = K$ , where  $\mathcal{M}(H/K)$  denotes the set of minimal non-trivial normal subgroups of  $H/K$  and  $\tilde{K} = \frac{1}{|K|} \sum_{k \in K} k$ . Furthermore,  $e(G, H, K)$  denotes the sum of the different  $G$ -conjugates of  $\epsilon(H, K)$ .

For arbitrary finite groups, Jespers, Olteanu and del Río obtained a description of  $e_{\mathbb{Q}}(\chi)$  in [2]. It is shown that  $e_{\mathbb{Q}}(\chi)$  is a  $\mathbb{Q}$ -linear combination of the elements  $e(G, H_i, K_i)$ , with  $(H_i, K_i)$  Shoda pairs in some subgroups of  $G$ . They posed the question ([2, Remark 3.4]) whether one could determine the scalars and the Shoda pairs involved. In this paper we answer both questions by giving a full description of the primitive central idempotents of  $\mathbb{Q}G$ , for  $G$  a finite group.

Throughout  $G$  is a finite group. For  $\chi$  an arbitrary complex character of  $G$  we put:  $e(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(1)\chi(g^{-1})g$  and  $e_{\mathbb{Q}}(\chi) = \sum_{\sigma \in G_{\chi}} \sigma(e(\chi))$ . Note that in general these elements do not have to be idempotents. Recall that the Möbius  $\mu$ -function,  $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ , is the map defined by  $\mu(1) = 1$ ,  $\mu(n) = 0$  if  $a^2 | n$  with  $a > 1$  and  $\mu(n) = (-1)^r$  if  $n = p_1 p_2 \dots p_r$  for different primes  $p_1, \dots, p_r$ . The induction of a character  $\phi$  of a subgroup  $H$  to  $G$  is defined as  $\phi_H^G(g) = \frac{1}{|H|} \sum_{y \in G} \phi(y^{-1}gy)$ , where  $\phi(x) = \phi(x)$  if  $x \in H$  and  $\phi(x) = 0$  otherwise. By  $1_G$  we denote the trivial character of  $G$ .

To prove our result we make use of the Artin Induction Theorem. Although this is probably well known, we state and prove it in the following specific form. Recall that for a rational valued character  $\chi$  of a group  $G$ ,  $\chi(g) = \chi(g^i)$  for  $(i, o(g)) = 1$ .

**Proposition 1** (Artin) *If  $\psi$  is a rational valued character of  $G$ , then*

$$\psi = \sum_{i=1}^r d_{C_i} 1_{C_i}^G,$$

where the sum runs through a set  $\{C_1, \dots, C_r\}$  of representatives of conjugacy classes of cyclic subgroups of  $G$ . Furthermore, if  $C_i = \langle c_i \rangle$  then

$$d_{C_i} = \frac{[G : \text{Cen}_G(c_i)]}{[G : C_i]} \sum_{C_i^* \geq C_i} \mu([C_i^* : C_i]) \psi(z^*),$$

where the sum runs through all the cyclic subgroups  $C_i^*$  of  $G$  containing  $C_i$  and  $C_i^* = \langle z^* \rangle$ .

**Proof.** For every cyclic subgroup  $C = \langle c \rangle$  of  $G$ , there exists exactly one  $i \in \{1, \dots, r\}$  such that  $C$  is  $G$ -conjugated to  $C_i$ . Say,  $C = C_i^{g^{-1}}$ . Set  $a_C = \frac{|\text{Cen}_G(c)|}{|C|} d_C$ . First we prove that  $a_C = a_{C_i}$  and  $1_C^G = 1_{C_i}^G$ . To prove the second equality, note that  $1_C^G(g) = \frac{1}{|C|} \sum_{y \in G} \dot{1}_C(y^{-1}gy)$ , where the function  $\dot{1}_C(y^{-1}gy)$  is defined as 1 if  $y^{-1}gy \in C$  and 0 otherwise. This combined with the facts that conjugation preserves the order of subgroups and that it is an automorphism of  $G$  we easily see that  $1_C^G = 1_{C_i}^G$ .

Now we prove that  $a_C = a_{C_i}$ . Define the sets  $(C_i) \uparrow^{\geq} = \{K \mid C_i \leq K \leq G\}$  and  $(C) \uparrow^{\geq} = \{K \mid C \leq K \leq G\}$ . There is a bijective correspondance between these sets. A map from  $(C_i) \uparrow^{\geq}$  to  $(C) \uparrow^{\geq}$  is given by conjugation with  $g^{-1}$  and the invers map is conjugation by  $g$ . Along with the fact that  $C^g = \langle c^g \rangle$  if  $C = \langle c \rangle$  and  $|C| = |C^g|$ , we see immediatly that  $a_C = a_{C_i}$ .

All this yields,  $\sum_C a_C 1_C^G = \sum_{i=1}^r k_i a_{C_i} 1_{C_i}^G = \sum_{i=1}^r d_{C_i} 1_{C_i}^G$ , where  $k_i = |\mathcal{C}_G(c_i)| = \frac{|G|}{|\text{Cen}_G(c_i)|}$  (with  $\mathcal{C}_G(c_i)$  the conjugacy class of  $c_i$  in  $G$ ). The result now follows from Artin's Induction Theorem, [1, page 489], which says that every rational valued character of  $G$  is of the form  $\sum_C a_C 1_C^G$ , with  $a_C$  as above and the sum runs over all cyclic subgroups  $C$  of  $G$ .  $\square$

Recall that by  $\tilde{C}_i$  we denote  $\epsilon(C_i, C_i) = \frac{1}{|C_i|} \sum_{c_i \in C_i} c_i$ .

**Theorem 2** *Let  $G$  be a finite group and  $\chi$  an irreducible complex character of  $G$ . Let  $C_i = \langle c_i \rangle$ , then we denote*

$$b_{C_i} = \frac{[G : \text{Cen}_G(c_i)]}{[G : C_i]} \sum_{C_i^* \geq C_i} \mu([C_i^* : C_i]) \left( \sum_{\sigma \in G_\chi} \sigma(\chi) \right) (z^*)$$

where the sum runs through all the cyclic subgroups  $C_i^*$  of  $G$  which contain  $C_i$  and  $z^*$  is a generator of  $C_i^*$ . Then

$$e_{\mathbb{Q}}(\chi) = \sum_{i=1}^r \frac{b_{C_i} \chi(1)}{[G : \text{Cen}_G(\tilde{C}_i)]} e(G, C_i, C_i) = \sum_{i=1}^r \frac{b_{C_i} \chi(1)}{[G : C_i]} \left( \sum_{k=1}^{m_i} \tilde{C}_i^{g_{ik}} \right),$$

where the first sum runs through a set  $\{C_1, \dots, C_r\}$  of representatives of conjugacy classes of cyclic subgroups of  $G$  and  $T_i = \{g_{i1}, \dots, g_{im_i}\}$  a right transversal of  $C_i$  in  $G$ .

**Proof.** Let  $\chi$  be an irreducible complex character of  $G$ . First we suppose that  $\chi(G) \subseteq \mathbb{Q}$ . Then  $G_\chi = \{1\}$  and then by Proposition 1,  $\chi = \sum_{i=1}^r b_{C_i} 1_{C_i}^G$ .

We get

$$\begin{aligned}
e_{\mathbb{Q}}(\chi) = e(\chi) &= \frac{\chi(1)}{|G|} \sum_{g \in G} (\sum_{i=1}^r b_{C_i} 1_{C_i}^G(g^{-1}))g \\
&= \frac{\chi(1)}{|G|} \sum_{i=1}^r \frac{b_{C_i}}{|C_i^G(1)|} \sum_{g \in G} 1_{C_i}^G(1) 1_{C_i}^G(g^{-1})g \\
&= \frac{\chi(1)}{|G|} \sum_{i=1}^r \frac{b_{C_i}}{|C_i^G(1)|} |G| e(1_{C_i}^G) \\
&= \sum_{i=1}^r \frac{b_{C_i} \chi(1)}{|G : C_i|} e(1_{C_i}^G)
\end{aligned}$$

Let  $T_i = \{g_{i1}, \dots, g_{im_i}\}$  be a right transversal of  $C_i$  in  $G$ . Then

$$\begin{aligned}
e(1_{C_i}^G) &= \frac{1}{|G|} \sum_{g \in G} 1_{C_i}^G(1) 1_{C_i}^G(g^{-1})g \\
&= \frac{1}{|G|} \sum_{g \in G} \frac{|G|}{|C_i|} 1_{C_i}(1) 1_{C_i}^G(g^{-1})g \\
&= \sum_{g \in G} \frac{1}{|C_i|} (\frac{1}{|C_i|} \sum_{y \in G} 1_{C_i}(yg^{-1}y^{-1}))g \\
&= \sum_{g \in G} \frac{1}{|C_i|} (\sum_{j=1}^{m_i} 1_{C_i}(g_{ij}g^{-1}g_{ij}^{-1}))g \\
&= \frac{1}{|C_i|} \sum_{j=1}^{m_i} \sum_{h \in C_i} 1_{C_i}(h^{-1})g_{ij}^{-1}hg_{ij} \\
&= \sum_{j=1}^{m_i} \tilde{C}_i^{g_{ij}}
\end{aligned}$$

With this expression for  $e(1_{C_i}^G)$  we obtain one of the equalities in the statement of the result.

Obviously, the sum  $\sum_{k=1}^{m_i} \tilde{C}_i^{g_{ik}}$  adds the elements of the  $G$ -orbit of  $\tilde{C}_i = \epsilon(C_i, C_i)$  and each of them  $[Cen_G(\tilde{C}_i) : C_i]$  times. So

$$\sum_{k=1}^{m_i} \tilde{C}_i^{g_{ik}} = [Cen_G(\tilde{C}_i) : C_i] e(G, C_i, C_i).$$

A simple substitution in the earlier found expression for  $e_{\mathbb{Q}}(\chi)$  yields the theorem.

Assume now that  $\chi$  is an arbitrary irreducible complex character of  $G$ . Then it is clear and well known that  $\sum_{\sigma \in G_{\chi}} \sigma \circ \chi$  is a rational valued character of  $G$ . Hence, by the first part we get

$$\begin{aligned}
e(\sum_{\sigma \in G_{\chi}} \sigma(\chi)) &= \frac{1}{|G|} \sum_{g \in G} (\sum_{\sigma \in G_{\chi}} \sigma(\chi(1))) (\sum_{\sigma \in G_{\chi}} \sigma(\chi(g^{-1})))g \\
&= \frac{|G_{\chi}| \cdot \chi(1)}{|G|} \sum_{\sigma \in G_{\chi}} \sum_{g \in G} \sigma(\chi(g^{-1}))g \\
&= \frac{|G_{\chi}| \cdot \chi(1)}{|G|} \sum_{\sigma \in G_{\chi}} \sigma(\sum_{g \in G} \chi(g^{-1})g) \\
&= |G_{\chi}| e_{\mathbb{Q}}(\chi)
\end{aligned}$$

Since  $\sum_{\sigma \in G_{\chi}} \sigma(\chi(1)) = |G_{\chi}| \chi(1)$ , the rational case yields the theorem.  $\square$

We finish with some remarks. First note that the elements  $e(G, C_i, C_i)$  are not necessarily idempotens. Second, the definition of  $b_{C_i}$  is not character-free. However one easily obtains a character free upperbound:

$$b_{C_i} \leq \frac{[G : Cen_G(c_i)]}{[G : C_i]} \sum_{C_i^* \geq C_i} \mu([C_i^* : C_i]) \phi(n) \chi(1) \leq \frac{[G : Cen_G(c_i)][G : Z(G)]}{[G : C_i]} \sum_{C_i^* \geq C_i} \mu([C_i^* : C_i]) \phi(n),$$

where  $\phi$  denotes the  $\phi$ -Euler function. Hence, we can obtain a finite algorithm, that easily can be implemented in for example GAP, to compute all primitive central idempotents of  $\mathbb{Q}G$ . This answers one of the questions posed in [2, Remark 3.4]. Also the description of the idempotents only makes use of pairs of subgroups  $(C_i, C_i)$ , with  $C_i$  cyclic. This answers the second question posed in [2, Remark 3.4].

## References

- [1] Bertram Huppert, Character Theory of Finite Groups, De Gruyter expositions in mathematics; 25, 1998
- [2] Eric Jespers, Gabriela Olteanu and Ángel del Río, Rational Group Algebras of Finite Groups: from Idempotents to Units of Integral Group Rings, to appear in, Algebras and Representation Theory
- [3] A.Olivieri, Á.del Río, J.J. Simón, On monomial characters and central idempotents of rational group algebras, Communications in Algebra 32 (2004), 1531-1550.

G.Janssens

Departement of Mathematics, Vrije Universiteit Brussel, Pleinlaan 2, 1050, Brussels, Belgium  
e-mail: geofjans@vub.ac.be