Hamilton cycles in almost distance-hereditary graphs

Bing Chen^{a*} and Bo Ning^{b†}

^aDepartment of Applied Mathematics, School of Science,

Xi'an University of Technology, Xi'an, Shaanxi 710048, P.R. China ^bDepartment of Applied Mathematics, School of Science,

Northwestern Polytechnical University, Xi'an, Shaanxi 710072, P.R. China

Abstract

Let G be a graph on $n \geq 3$ vertices. A graph G is almost distance-hereditary if each connected induced subgraph H of G has the property $d_H(x, y) \leq d_G(x, y) + 1$ for any pair of vertices $x, y \in V(H)$. A graph G is called 1-heavy (2-heavy) if at least one (two) of the end vertices of each induced subgraph of G isomorphic to $K_{1,3}$ (a claw) has (have) degree at least n/2, and called claw-heavy if each claw of G has a pair of end vertices with degree sum at least n. Thus every 2-heavy graph is claw-heavy. In this paper we prove the following two results: (1) Every 2-connected, claw-heavy and almost distance-hereditary graph is Hamiltonian. (2) Every 3-connected, 1-heavy and almost distance-hereditary graph is Hamiltonian. In particular, the first result improves a previous theorem of Feng and Guo. Both results are sharp in some sense.

Keywords: Hamilton cycle; Almost distance-hereditary graph; Claw-free graph; 1-heavy graph; 2-heavy graph; Claw-heavy graph

AMS Subject Classification (2000): 05C38, 05C45

1 Introduction

We use Bondy and Murty [4] for terminology and notation not defined here and consider simple graphs only.

Let G be a graph. For a vertex v and a subgraph H of G, we use $N_H(v)$ to denote the set, and $d_H(v)$ the number, of neighbors of v in H, respectively. We call $d_H(v)$ the degree

^{*}Supported by NSFC (No. 11271300) and the Scientific Research Program of Shaanxi Provincial Education Department (No. 2013JK0580). E-mail address: cbing_2004@163.com (B. Chen).

[†]Corresponding author. Supported by NSFC (No. 11271300) and the Doctorate Foundation of Northwestern Polytechnical University (No. cx201326). E-mail address: ningbo_math84@mail.nwpu.edu.cn (B. Ning).

of v in H. For $x, y \in V(G)$, an (x, y)-path is a path connecting x and y. If $x, y \in V(H)$, the distance between x and y in H, denoted by $d_H(x, y)$, is the length of a shortest (x, y)-path in H. When there is no danger of ambiguity, $N_G(v)$, $d_G(v)$ and $d_G(x, y)$ are abbreviated to N(v), d(v) and d(x, y), respectively.

A graph is called Hamiltonian if it contains a Hamilton cycle, i.e., a cycle passing through all the vertices of the graph. The study of cycles, especially Hamilton cycles, maybe one of the most important and most studied areas of graph theory. It is well-known that to determine whether a given graph contains a Hamilton cycle is \mathcal{NP} -complete, shown by R.M. Karp [15]. However, if we only consider some restricted graph classes, then the situation is completely changed. A graph G is called distance-hereditary if each connected induced subgraph H has the property that $d_H(x, y) = d_G(x, y)$ for any pair of vertices x, y in H. This concept was introduced by Howorka [13] and a complete characterization of distance-hereditary graphs can be found in [13]. In 2002, Hsieh, Ho, Hsu and Ko [14] obtained an O(|V| + |E|)-time algorithm to solve the Hamiltonian problem on distancehereditary graphs. Some other optimization problems can also be solved in linear time for distance-hereditary graphs although they are proved to be \mathcal{NP} -hard for more general graphs. For references in this direction, we refer to [2, 8].

A graph G is called *almost distance-hereditary* if each connected induced subgraph H of G has the property $d_H(x, y) \leq d_G(x, y) + 1$ for any pair of vertices $x, y \in V(H)$. For some properties and a characterization of almost-distance hereditary graphs, we refer to [1].

Let G be a graph. An induced subgraph of G isomorphic to $K_{1,3}$ is called a *claw*, the vertex of degree 3 in it is called its *center* and the other vertices are its *end vertices*. G is called *claw-free* if G contains no claw. Throughout this paper, whenever the vertices of a claw are listed, its center is always the first one.

The class of claw-free graphs is important in graph theory. Maybe one big reason is due to Matthews and Sumner's conjecture [17] which states that every 4-connected claw-free graph is Hamiltonian. Many results about the existence of Hamilton cycles in claw-free graphs have been obtained. For surveys on Matthews and Sumner's conjecture and on claw-free graphs, we refer the reader to [5] and [9], respectively.

In particular, Feng and Guo [10] gave the following result on Hamiltonicity of almost distance-hereditary claw-free graphs.

Theorem 1 (Feng and Guo [10]). Let G be a 2-connected claw-free graph. If G is almost distance-hereditary, then G is Hamiltonian.

Let G be a graph. A vertex v of G on n vertices is called *heavy* if $d(v) \ge n/2$. Broersma

et al. [6] introduced the concepts of 1-heavy graph and 2-heavy graph, and the concept of claw-heavy graph was introduced by Fujisawa [12]. Following [6, 12], we say that a claw in G is 1-heavy (2-heavy) if at least one (two) of its end vertices is (are) heavy. G is called 1-heavy (2-heavy) if every claw of it is 1-heavy (2-heavy), and called *claw-heavy* if every claw of it has two end vertices with degree sum at least n. It is easily seen that every claw-free graph is 1-heavy (2-heavy, claw-heavy), every 2-heavy graph is claw-heavy but not every claw-heavy graph is 2-heavy, and every claw-heavy graph is 1-heavy but not every 1-heavy graph is claw-heavy.

Broersma et al. [6] proved some sufficient conditions for Hamiltonicity of 1-heavy graphs and 2-heavy graphs. Motivated by the works of Broersma et al. [6], Feng and Guo [11] extended Theorem 1 to a larger graph class of 2-heavy graphs.

Theorem 2 (Feng and Guo [11]). Let G be a 2-connected 2-heavy graph. If G is almost distance-hereditary, then G is Hamiltonian.

Later, Chen et al. [7] extended two results of Broersma et al. [6] to the more larger graph class of claw-heavy graphs. Motivated by Chen et al.' previous works [7], in this paper we obtain the following two theorems which extend Theorem 1 and Theorem 2. In particular, Theorem 3 improves Theorem 2.

Theorem 3. Let G be a 2-connected claw-heavy graph. If G is almost distance-hereditary, then G is Hamiltonian.

Theorem 4. Let G be a 3-connected 1-heavy graph. If G is almost distance-hereditary, then G is Hamiltonian.

Remark 1. The graph in Fig.1 shows that the result in Theorem 3 indeed strengthen that in Theorem 2. As shown in [7, Fig.2], let $n \ge 10$ be an even integer and $K_{n/2} + K_{n/2-3}$ denote the join of two complete graphs $K_{n/2}$ and $K_{n/2-3}$. Choose a vertex $y \in V(K_{n/2})$ and construct a graph G with $V(G) = V(K_{n/2} + K_{n/2-3}) \cup \{v, u, x\}$ and $E(G) = E(K_{n/2} + K_{n/2-3}) \cup \{uv, uy, ux\} \cup \{vw, xw : w \in V(K_{n/2-3})\}$. It is easy to see that G is a Hamiltonian graph satisfying the condition of Theorem 3, but not the condition of Theorem 2.




Fig.1

Remark 2. In [11], Feng and Guo constructed a graph which is 2-connected almost distance-hereditary 1-heavy graph but not Hamiltonian. Hence the connectivity condition in our Theorem 4 is sharp.

We postpone the proofs of Theorem 3 and 4 to the next section.

2 Proofs of Theorems 3 and 4

Before the proofs, we first introduce some additional terminology and notation.

Let H be a path (a cycle) with a given orientation. We denote H by the same graph as H but with the reverse orientation. When $v \in V(H)$, v_H^+ and v_H^- denote the successor (if it exists) and the predecessor (if it exists) of v on H, respectively. If $S \subseteq V(H)$, then define $S_H^+ = \{s_H^+ : s \in S\}$ and $S_H^- = \{s_H^- : s \in S\}$. If there is no danger of ambiguity, then we denote v_H^+ , v_H^- , S_H^+ and S_H^- by v^+ , v^- , S^+ and S^- , respectively. For two vertices $u, v \in V(H)$, we use H[u, v] to denote the segment of H from u to v, and by H[u, v), H(u, v] and H(u, v), we mean the path $H[u, v] - \{v\}$, $H[u, v] - \{u\}$ and $H[u, v] - \{u, v\}$, respectively.

Let G be a graph on n vertices and $k \ge 3$ be an integer. Recall that a vertex of degree at least n/2 in G is a heavy vertex; otherwise it is light. A claw in G is called a light claw if all its end vertices are light, and called an *o*-light claw if any pair of end vertices has degree sum less than n. A heavy cycle in G is a cycle containing all heavy vertices in G. Following [16], we use $\tilde{E}(G)$ to denote the set $\{xy : xy \in E(G) \text{ or } d(x) + d(y) \ge n, x, y \in V(G)\}$. A sequence of vertices $C = u_1 u_2 \dots u_k u_1$ is called an *Ore-cycle* or briefly, *o*-cycle of G, if $u_i u_{i+1} \in \tilde{E}(G)$ for every $i \in [1, k]$, where $u_1 = u_{k+1}$.

To prove Theorems 3 and 4, the following three lemmas are needed.

Lemma 1 (Bollobás and Brightwell [3], Shi [18]). Every 2-connected graph contains a heavy cycle.

Lemma 2 (Li, Ryjáček, Wang and Zhang [16]). Let G be graph on n vertices and C' be an o-cycle of G. Then there exists a cycle C of G such that $V(C') \subseteq V(C)$.

Lemma 3. Let G be a non-Hamiltonian graph, C be a longest heavy cycle (a longest cycle) of G, R a component of G - V(C), and $A = \{v_1, v_2, \ldots, v_k\}$ the set of neighbors of R on C. Let $u \in V(R)$, $v_i, v_j \in A$ and P be a (v_i, v_j) -path with all internal vertices in R. Then there hold

(a) $uv_i^- \notin \widetilde{E}(G), uv_i^+ \notin \widetilde{E}(G);$ (b) $v_i^- v_i^- \notin \widetilde{E}(G), v_i^+ v_i^+ \notin \widetilde{E}(G);$

- (c) If $v_i^- v_i^+ \in \widetilde{E}(G)$, then $v_i v_j^- \notin \widetilde{E}(G)$, $v_i v_j^+ \notin \widetilde{E}(G)$;
- (d) Let $l \in V(C[v_i, v_i^-]) \cap N(v_i) \cap N(v_i^-)$. If $v_i^- v_i^+ \in \widetilde{E}(G)$, then $l^-l^+ \notin \widetilde{E}(G)$, $v_i l^+ \notin \widetilde{E}(G)$;
- (e) Let $l \in V(C[v_i, v_i^-]) \cap N(v_i)$. If $v_i^- v_i^+ \in \widetilde{E}(G)$, then $l^- v_i^- \notin \widetilde{E}(G)$, $v_i^+ l^+ \notin \widetilde{E}(G)$;
- $(f) \ Let \ l \in V(C[v_i, v_j^-]). \ If \ v_i^- l \in \widetilde{E}(G), \ then \ l^-v_j^- \notin \widetilde{E}(G), \ l^-v_j^+ \notin \widetilde{E}(G), \ l^+v_j^+ \notin \widetilde{E}(G);$

Furthermore, if G is a 2-connected claw-heavy graph, then

- $(g) \ v_i^- v_i^+ \in \widetilde{E}(G), \ v_j^- v_j^+ \in \widetilde{E}(G);$
- (h) $v_i^- v_i^+ \in E(G) \text{ or } v_j^- v_j^+ \in E(G).$

Proof. (a), (b), (c) The proof of (a) is similar as the proof of Claim 1, and the proof of (b), (c) is similar as the proof of Claim 3, in [16, Theorem 8], respectively.

(d) Suppose that $l^-l^+ \in \widetilde{E}(G)$. Then $C' = PC[v_j, v_i^-]v_i^-v_i^+C[v_i^+, l^-]l^-l^+C[l^+, v_j^-]$ $v_j^-lv_i$ is an o-cycle such that $V(C) \subset V(C')$. By Lemma 2, there is a longer cycle C'' containing all vertices in C, contradicting the choice of C. Suppose that $v_i l^+ \in \widetilde{E}(G)$. Then $C' = PC[v_j, v_i^-]v_i^-v_i^+C[v_i^+, l]lv_j^-\overleftarrow{C}[v_j^-, l^+]l^+v_i$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction.

(e) Suppose that $l^-v_j^- \in \widetilde{E}(G)$. Then $C' = \overleftarrow{P}v_i lC[l, v_j^-]v_j^-l^-\overleftarrow{C}[l^-, v_i^+]v_i^+v_i^-\overleftarrow{C}[v_i^-, v_j]$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction. Suppose that $v_j^+l^+ \in \widetilde{E}(G)$. Then $C' = P\overleftarrow{C}[v_j, l^+]l^+v_j^+C[v_j^+, v_i^-]v_i^-v_i^+C[v_i^+, l]lv_i$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction.

(f) Suppose that $l^-v_j^- \in \widetilde{E}(G)$. Then $C' = PC[v_j, v_i^-]v_i^-lC[l, v_j^-]v_j^-l^-\overleftarrow{C}[l^-, v_i]$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction. Suppose that $l^-v_j^+ \in \widetilde{E}(G)$. Then $C' = \overleftarrow{P}C[v_i, l^-]l^-v_j^+C[v_j^+, v_i^-]v_i^-lC[l, v_j]$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction. Suppose that $l^+v_j^+ \in \widetilde{E}(G)$. Then $C' = P\overleftarrow{C}[v_j, l^+]l^+v_j^+C[v_j^+, v_i^-]v_i^-l\overleftarrow{C}[l, v_i]$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction. Suppose that $l^+v_j^+ \in \widetilde{E}(G)$. Then $C' = P\overleftarrow{C}[v_j, l^+]l^+v_j^+C[v_j^+, v_i^-]v_i^-l\overleftarrow{C}[l, v_i]$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction.

(g), (h) The proof of (g) and (h) is similar as the proof of Claim 2, 4 in [16, Theorem 8], respectively. \Box

Proof of Theorem 3

Let G be a graph satisfying the condition of Theorem 3. Let C be a longest cycle of G and assign an orientation to it. Suppose G is not Hamiltonian. Then $V(G)\setminus V(C) \neq \emptyset$. Let R be a component of G - C, and $A = \{v_1, v_2, \ldots, v_k\}$ be the set of neighbors of R on C. Since G is 2-connected, there exists a (v_i, v_j) -path $P = v_i u_1 \ldots u_r v_j$ with all internal vertices in R, and $v_i, v_j \in A$. Choose P such that:

- (1) $|V(C(v_i, v_j))|$ is as small as possible;
- (2) |V(P)| is as small as possible subject to (1).

By the choice of P, the first claim is obvious.

Claim 1. For any two vertices $u_s, u_t \in V(P) \setminus \{v_i, v_j\}$ and $v \in V(C(v_i, v_j))$, there hold $u_s u_t \notin E(G)$ if $d_P(u_s, u_t) \ge 2$, $u_s v \notin E(G)$, $v_i u_s \notin E(G)$ and $v_j u_s \notin E(G)$.

Claim 2. There is no *o*-cycle C' in G such that $V(C) \subset V(C')$.

Proof. Otherwise, C' is an *o*-cycle such that $V(C) \subset V(C')$. By Lemma 2, there exists a cycle containing all vertices in C' and longer than C, contradicting the choice of C.

By Lemma 3 (h), without loss of generality, assume that $v_i^- v_i^+ \in E(G)$.

Claim 3. r = 1, that is, $V(P) = \{v_i, u_1, v_j\}$.

Proof. Suppose that $r \ge 2$. Consider $H = G[V(P) \cup V(C[v_i, v_j])] - v_j$. By Lemma 3 (g), $v_i^- v_i^+ \in \widetilde{E}(G)$. Since $v_i^- v_i^+ \in \widetilde{E}(G)$, $v_i v_j^- \notin E(G)$ by Lemma 3 (c). Thus $d_H(v_j^-, v_i) \ge 2$. By the choice of P, we have $d_P(v_i, u_r) \ge 2$. Now by Claim 1, $d_H(v_j^-, u_r) = d_H(v_j^-, v_i) + d_P(v_i, u_r) \ge 4$, which yields a contradiction to the fact G is almost distance-hereditary and $d_G(v_j^-, u_r) = 2$. Hence $V(P) = \{v_i, u_1, v_j\}$.

Claim 4. $|V(C[v_i, v_j])| \ge 5.$

Proof. Suppose $|V(C[v_i, v_j])| = 4$ or $|V(C[v_i, v_j])| = 3$. This means $C[v_i, v_j] = v_i v_i^+ v_j^- v_j$ or $C[v_i, v_j] = v_i v_i^+ v_j$. Let $C' = v_i u_1 v_j v_j^- v_j^+ C[v_j^+, v_i^-] v_i^- v_i^+ v_i$ or $C' = v_i u_1 v_j v_j^- v_j^+ C[v_j^+, v_i]$. Then C' is an o-cycle such that $V(C) \subset V(C')$, contradicting Claim 2.

Let $H = G[\{u_1, v_i^-\} \cup V(C[v_i, v_j])] - v_i$. Since $d_G(v_i^-, u_1) = 2$ and G is almost distancehereditary, $d_H(v_i^-, u_1) \leq 3$. By Lemma 3 (c) and Claim 1, we have $v_i^-v_j \notin E(G)$ and $u_1v_s \notin E(G)$, where $v_s \in C[v_i^+, v_j^-]$. It follows that $d_H(v_i^-, u_1) = 3$ and $d_H(v_i^-, v_j) = 2$. By Lemma 3 (b) and (c), $v_i^-v_j^- \notin E(G)$ and $v_i^+v_j \notin E(G)$. Thus there exits a vertex $w \in C(v_i^+, v_j^-)$ such that $v_i^-w \in E(G)$ and $wv_j \in E(G)$. Note that w is well-defined by Claim 4.

Claim 5. $v_i^- w^+ \in \widetilde{E}(G)$.

Proof. Suppose that $v_i^-w^+ \notin \widetilde{E}(G)$. By Lemma 3 (e) and symmetry, we have $v_jw^+ \notin \widetilde{E}(G)$. Note that $v_jv_i^- \notin \widetilde{E}(G)$ by Lemma 3 (c). Thus $\{w, w^+, v_j, v_i^-\}$ induces an o-light claw in G, a contradiction.

Claim 6. $wv_i^+ \in \widetilde{E}(G)$.

Proof. First we will show that $w^-v_i^- \notin \widetilde{E}(G)$. Since $v_j^-v_j^+ \in \widetilde{E}(G)$ and $v_jw \in E(G)$, we have $w^-v_i^- \notin \widetilde{E}(G)$ by Lemma 3 (e) and symmetry.

Next we will show that $w^-v_j \in \widetilde{E}(G)$. Suppose not. Consider the subgraph induced by $\{w, w^-, v_j, v_i^-\}$. Note that $v_j v_i^- \notin \widetilde{E}(G)$ by Lemma 3 (c) and $w^-v_i^- \notin \widetilde{E}(G)$. Then $\{w, w^-, v_j, v_i^-\}$ induces an o-light claw, a contradiction.

Next we will show that $u_1w \notin \widetilde{E}(G)$. Otherwise, $C' = u_1wC[w, v_j^-]v_j^-v_j^+C[v_j^+, w^-]w^$ v_ju_1 (note that $w^-v_j \in \widetilde{E}(G)$) is an *o*-cycle such that $V(C) \subset V(C')$, contradicting Claim 2.

Now we will show that $wv_j^+ \in \widetilde{E}(G)$. Consider the subgraph induced by $\{v_j, u_1, w, v_j^+\}$. Note that $u_1w \notin \widetilde{E}(G)$ by the analysis above and $u_1v_j^+ \notin \widetilde{E}(G)$ by Lemma 3 (a). Since G is claw-heavy, we have $wv_j^+ \in \widetilde{E}(G)$.

By Claim 5, we have $v_i^- w^+ \in \widetilde{E}(G)$. By Lemma 3 (f), we have $wv_j^+ \notin \widetilde{E}(G)$, contradicting Claim 6. The proof of Theorem 3 is complete.

Proof of Theorem 4

Let G be a graph satisfying the condition of Theorem 4. By Lemma 1, there exists a heavy cycle in G. Now choose a longest heavy cycle C of G and assign an orientation to it. Suppose G is not Hamiltonian. Then $V(G)\setminus V(C) \neq \emptyset$. Let R be a component of G - Cand $A = \{w_1, w_2, \ldots, w_k\}$ be the set of neighbors of R on C. Since G is 3-connected, for any vertex u of R, there exists a (u,C)-fan F such that $F = (u;Q_1,Q_2,Q_3)$, where $Q_1 = uu_1 \ldots u_{r_1} w_i, Q_2 = us_1 \ldots s_{r_2} w_j$ and $Q_3 = uy_1 \ldots y_{r_3} w_k$ are three internally disjoint paths, $V(Q_1) \cap V(C) = w_i, V(Q_2) \cap V(C) = w_j, V(Q_3) \cap V(C) = w_k$, and w_i, w_j, w_k are in the order of the orientation of C.

By the choice of C, all internal vertices on F are not heavy. By Lemma 3 (b), there is at most one heavy vertex in $N_C^+(R)$ and at most one heavy vertex in $N_C^-(R)$. Now assume that w_i^-, w_i^+ are light. Hence $w_i^- w_i^+ \in E(G)$, otherwise $\{w_i, w_i^-, w_i^+, u_{r_1}\}$ induces a light claw, contradicting G is 1-heavy.

Claim 1. There exists a (u,C)-fan F such that $V(F) = \{u, w_i, w_j, w_k\}$.

Proof. Now we choose the fan F in such a way that:

- (1) $|V(Q_1)| = 2$ and $w_i \in V(Q_1);$
- (2) $|V(C[w_i, w_j])|$ is as small as possible subject to (1);
- (3) $|V(Q_2)|$ is as small as possible subject to (1), (2);
- (4) $|V(C[w_k, w_i])|$ is as small as possible subject to (1), (2) and (3);
- (5) $|V(Q_3)|$ is as small as possible subject to (1), (2), (3) and (4)

Since G is 3-connected, this choice condition is well-defined. Without loss of generality, assume $Q_1 = uw_i$. Note that $w_i^- w_i^+ \in E(G)$.

First we show that $V(Q_2) = \{u, w_j\}$. Suppose there exists a vertex of $V(Q_2) \setminus \{u, w_j\}$. Without loss of generality, set $x = s_{r_2}$. Let $H = G[V(Q_1) \cup V(Q_2) \cup V(C[w_i, w_j])] - w_j$. By Lemma 3 (c), it is easy to see that $w_i w_j^- \notin E(G)$, so $d_H(w_j^-, w_i) \ge 2$, and the choice condition (2) implies that $N_{C(w_i,w_j)}(Q_2 \setminus \{u, w_j\}) = \emptyset$. This means that $d_H(w_i, x) = d_F(w_i, x)$. If $xw_i \notin E(G)$, then $d_F(w_i, x) \ge 2$. Since $|V(Q_2)| > 2$, we have $d_H(w_j^-, x) = d_H(w_j^-, w_i) + d_F(w_i, x) \ge 4$. It yields a contradiction to the fact G is almost distance-hereditary and $d_G(w_j^-, x) = 2$. Thus, $xw_i \in E(G)$. Let $F = (x, xw_i, xw_j, Q_2[x, u]Q_3[u, w_j])$. Then F is a (x, C)-fan satisfying (1), (2) and $|\{x, w_j\}| = 2$, contradicting the choice condition (3), a contradiction. Hence $V(Q_2) = \{u, w_j\}$.

Next we show that $V(Q_3) = \{u, w_k\}$. Suppose there exists a vertex of $V(Q_3) \setminus \{u, w_k\}$. Without loss of generality, set $x = y_{r_3}$. If $xw_i \notin E(G)$, let $H = G[V(Q_1) \cup V(Q_3) \cup V(C[w_k, w_i])] - w_k$. By Lemma 3 (c), we have $w_k^+ w_i \notin E(G)$. This means $d_H(w_k^+, w_i) \ge 2$. By the choice condition (4) (5), we have $N(Q_3 \setminus \{w_k\}) \cap V(C(w_k, w_i)) = \emptyset$. Since $|V(Q_3)| \ge 3$, $d_H(w_i, x) \ge 2$ and we have $d_H(w_k^+, x) = d_H(w_k^+, w_i) + d_F(w_i, x) \ge 4$. It yields a contradiction to the fact G is almost distance-hereditary and $d_G(w_k^+, x) = 2$. Thus, $xw_i \in E(G)$. Since $d_G(x, w_i^+) = 2$ and G is almost distance-hereditary, let $H = G[V(C[w_i, w_j]) \cup V(Q_3[u, x])] - \{w_i\}$, we have $d_H(w_i^+, x) = 3$. Note that $N_C(x) \cap C(w_i, w_j] = \emptyset$ and $N_C(u) \cap C(w_i, w_j) = \emptyset$. It follows that $w_i^+ w_j \in E(G)$, and $w_j^- w_j^+ \notin \widetilde{E}(G)$ by Lemma 3 (c). Consider the subgraph induced by $\{w_j, w_i^+, w_j^+, u\}$. We know that w_j^+ is heavy. Now let $H = G[\{w_i^-\} \cup V(C[w_i, w_j]) \cup V(Q_3[u, x])] - \{w_i\}$. Since $d_G(w_i^-, x) = 2$, we have $d_H(w_i^-, x) = 3$, and $w_i^- w_j \in E(G)$. Consider the subgraph induced by $\{w_j, w_i^-, w_j^-, u\}$. We can see that w_j^- is heavy. Now $w_j^- w_j^+ \in \widetilde{E}(G)$, a contradiction. The proof is complete.

By Claim 1, there exists a (u, C)-fan F such that $V(F) \setminus V(C) = \{u\}$. Suppose that $N_C(u) = \{v_1, v_2, \ldots, v_r\}$ $(r \ge 3)$ and v_1, v_2, \ldots, v_r are in the order of the orientation of C. In the following, all subscripts of v are taken modulo r, and $v_0 = v_r$.

Claim 2. For any vertex $v_i \in N_C(u)$ such that $v_i^- v_i^+ \in \widetilde{E}(G)$, there exists a vertex $l_i \in C[v_i^+, v_{i+1}^-]$ such that $v_{i+1}^- l \in E(G)$ and $lv_i \in E(G)$; there exists a vertex $l_i' \in C(v_{i-1}^+, v_i^-]$ such that $v_{i-1}^+ s \in E(G)$ and $sv_i \in E(G)$.

Proof. Let $H = G[\{u\} \cup V(C[v_i, v_{i+1}])] - v_{i+1}$. Since $d_G(v_{i+1}^-, u) = 2$ and G is almost distance-hereditary, we have $d_H(v_{i+1}^-, u) \leq 3$. By Lemma 3 (c), we have $v_i v_{i+1}^- \notin E(G)$ and $d_H(v_{i+1}^-, u) = 3$. It follows that $d_H(v_{i+1}^-, v_i) = 2$. So there exists a vertex $l_i \in C[v_i^+, v_{i+1}^-)$ such that $v_{i+1}^- l_i \in E(G)$ and $l_i v_i \in E(G)$. The other assertion can be proved similarly. \Box

By Lemma 3 (b), there is at most one heavy vertex in $N_C^+(u)$ and at most one heavy vertex in $N_C^-(u)$. Since $r \ge 3$, we know that there exits $v_j \in N_C(u)$, such that $v_j^- v_j^+ \in$ E(G). Without loss of generality, assume that $v_1^-v_1^+ \in E(G)$. We divide the proof into two cases.

Case 1. $v_2^-v_2^+ \notin E(G)$ and $v_r^-v_r^+ \notin E(G)$.

Both $\{v_2, v_2^-, v_2^+, u\}$ and $\{v_r, v_r^-, v_r^+, u\}$ induce claws. By Lemma 3 (b) and the fact G is 1-heavy, v_2^-, v_r^+ are heavy or v_2^+ and v_r^- are heavy vertices.

By Claim 2, there exists a vertex $l_1 \in C[v_1^+, v_2^-)$ such that $v_2^- l_1 \in E(G)$ and $l_1 v_1 \in E(G)$, and there exists a vertex $l_1' \in C(v_r^+, v_1^-)$ such that $v_r^+ l_1' \in E(G)$ and $l_1' v_1 \in E(G)$.

Claim 3. (1) $v_1, v_1^-, v_1^+, l_1^+, l_1^-, l_1'^-, l_1'^+$ are light vertices;

- (2) $v_1 l_1^- \in E(G);$
- (3) l_1, l_1' are light vertices;
- (4) $v_1^- v_2 \notin E(G);$
- (5) $v_1^- l_1 \in E(G)$ and $l_1 v_2 \in E(G)$.

Proof. (1) By Lemma 3 (b), (c) and (e), it is obvious that $v_1, v_1^-, v_1^+, l_1^+, l_1^-, l_1'^-, l_1'^+$ are light vertices.

(2) Suppose that $v_1l_1^- \notin E(G)$. Note that $v_1l_1^+ \notin \widetilde{E}(G)$ and $l_1^-l_1^+ \notin \widetilde{E}(G)$ by Lemma 3 (d). Since l_1^+, v_1, l_1^- are light, $\{l_1, l_1^+, v_1, l_1^-\}$ induces a light claw, a contradiction.

(3) Since $v_1 l_1^- \in E(G)$, we have $v_2^+ l_1 \notin \widetilde{E}(G)$ and $v_r^+ l_1 \notin \widetilde{E}(G)$ by Lemma 3 (e). Note that either v_r^+ or v_2^+ is a heavy vertex. This implies l_1 is a light vertex. The other assertion can be proved similarly.

(4) Suppose that $v_1^-v_2 \in E(G)$ and v_2^- , v_3^+ are heavy. Let $C' = v_1^-v_2C[v_2, v_r]v_ruv_1$ $C[v_1, v_2^-]v_2^-v_r^+C[v_r^+, v_1^-]$. Then C' is an o-cycle such that $V(C) \subset V(C')$, a contradiction.

Suppose that $v_1^-v_2 \in E(G)$ and v_2^+ , v_r^- are heavy. Now $\{v_2, v_2^-, u, v_1^-\}$ induces a light claw, a contradiction.

(5) Suppose that $v_1^- l_1 \notin E(G)$. Note that $uv_1^- \notin \widetilde{E}(G)$ by Lemma 3 (a) and $ul_1 \notin \widetilde{E}(G)$ by Lemma 3 (c) and $v_1 l_1^- \in E(G)$. Now $\{v_1, l_1, u, v_1^-\}$ induces a light claw, a contradiction.

Suppose that $l_1v_2 \notin E(G)$. Let $H = G[\{v_1^-, l_1, v_2^-, v_2, u\}]$. By Lemma 3, we have $uv_2^- \notin E(G)$, $ul_1 \notin E(G)$, $uv_1^- \notin E(G)$, $v_2v_1^- \notin E(G)$ and $v_1^-v_2^- \notin E(G)$. Now $G[\{v_1^-, l_1, v_2^-, v_2, u\}]$ is an induced path of length 4 in G. It follows that $d_H(v_1^-, u) = 4$, contradicting the fact that $d_G(v_1^-, u) = 2$ and G is almost distance-hereditary. Hence we have $l_1v_2 \in E(G)$.

Now we consider the following two subcases.

Subcase 1.1. v_2^-, v_r^+ are heavy vertices.

By Lemma 3 (c) and $v_1 l_1^- \in E(G)$, $ul_1 \notin \widetilde{E}(G)$. By Lemma 3 (a) and (e), we have $uv_2^+ \notin \widetilde{E}(G)$ and $l_1v_2^+ \notin \widetilde{E}(G)$. By Claim 3 (1) and (3), $\{v_2, l_1, u, v_2^+\}$ induces a light claw, a contradiction.

Subcase 1.2. v_2^+ and v_r^- are heavy.

Claim 4. $G[\{v_r^+, l_1', l_1, v_2, u\}]$ is an induced path in G.

Proof. It is easily to check that $v_r^+ l_1' \in E(G)$, $l_1 v_2 \in E(G)$ (by Claim 3 (5)) and $v_2 u \in E(G)$. By Lemma 3 (a), $v_r^+ u \notin E(G)$. From the proof of Claim 3 (3), we know that $v_r^+ l_1 \notin E(G)$. Since $v_1 l_1^- \in E(G)$, $ul_1 \notin \widetilde{E}(G)$ by Lemma 3 (c). By symmetry, we have $ul_1' \notin \widetilde{E}(G)$ since $ul_1 \notin \widetilde{E}(G)$. Next we only need show that $l_1' l_1 \in E(G)$, $v_r^+ v_2 \notin E(G)$ and $l_1' v_2 \notin E(G)$.

Claim 4.1. $l_1' l_1 \in E(G)$.

Proof. Otherwise, $\{v_1, l_1, l_1', u\}$ induces a light claw, contradicting G is 1-heavy.

Claim 4.2. $v_r^+v_2 \notin E(G)$ and $l_1'v_2 \notin E(G)$.

Proof. Suppose that $v_r^+v_2 \in E(G)$. Consider the subgraph induced by $\{v_2, v_2^-, v_r^+, u\}$. By Lemma 3 (a), we have $uv_2^- \notin E(G)$ and $uv_r^+ \notin E(G)$. Since v_2^-, v_r^+, u are light and G is 1heavy, $v_r^+v_2^- \in E(G)$. Now $C' = v_1l_1^-\overleftarrow{C}[l_1^-, v_1^+]v_1^+v_1^-\overleftarrow{C}[v_1^-, v_r^+]v_r^+v_2^-\overleftarrow{C}[v_2^-, l_1]l_1v_2C[v_2, v_r]$ v_ruv_1 is an o-cycle such that $V(C) \subset V(C')$, a contradiction.

Suppose that $l_1'v_2 \in E(G)$. Consider the subgraph induced by $\{v_2, v_2^-, l_1', u\}$. Since v_2^-, l_1', u are light and G is 1-heavy, $l_1'v_2^- \in E(G)$. Now $C' = v_1uv_2C[v_2, l_1']l_1'v_2^ \overleftarrow{C}[v_2^-, v_1^+]v_1^+l_1'^+C[l_1'^+, v_1]$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction. (Note that $v_1^+l_1'^+ \in E(G)$ since otherwise $\{l_1', v_r^+, v_1^+, l_1'^+\}$ induces a light claw, a contradiction.)

Now it is proved that $G[\{v_r^+, l_1', l_1, v_2, u\}]$ is an induced path in G.

By Claim 4, $G[\{v_r^+, l_1', l_1, v_2, u\}]$ is an induced path of length 4 in G. It follows that $d_H(v_r^+, u) = 4$, contradicting the fact that $d_G(v_r^+, u) = 2$ and G is almost distance-hereditary.

Case 2. $v_2^- v_2 \in E(G)$ or $v_r^- v_r^+ \in E(G)$.

Without loss of generality, by symmetry, assume that $v_2^-v_2^+ \in E(G)$.

By Claim 2, for any vertex v_i such that $v_i^- v_i^+ \in E(G)$, there exists a vertex $l_i \in C[v_i^+, v_{i+1}^-)$ such that $l_i v_i \in E(G)$ and $v_{i+1}^- l \in E(G)$.

Claim 5. $v_{i+1}^+ l_i \in E(G)$.

Proof. Suppose that $v_{i+1}^+ l_i \notin E(G)$. Let $H = G[\{v_{i+1}^+, v_{i+1}^-, l_i, v_i, u\}]$. By Lemma 3 (c), we have $v_i v_{i+1}^- \notin E(G)$ and $v_i v_{i+1}^+ \notin E(G)$. We can see that H is an induced path of length 4 in G. Hence $d_H(v_{i+1}^+, u) = 4$, contradicting the fact that $d_G(v_{i+1}^+, u) = 2$ and G is almost distance-hereditary.

By Claim 5 and Lemma 3 (b), we have $l_i \neq v_i^+$.

Claim 6. $\{l_i, l_i^-, v_i, v_{i+1}^-\}$ induces a claw.

Proof. By Lemma 3 (c) and (e), we have $v_i v_{i+1}^- \notin \widetilde{E}(G)$ and $l_i^- v_{i+1}^- \notin \widetilde{E}(G)$. By Claim 5 and Lemma 3 (e), we have $v_i l_i^- \notin E(G)$. So $\{l_i, l_i^-, v_i, v_{i+1}^-\}$ induces a claw.

Subcase 2.1. $v_r^- v_r^+ \notin E(G)$.

Claim 7. $\{l_1, l_1^+, v_1, v_2^+\}$ induces a light claw.

Proof. By Lemma 3 (a), we have $uv_r^- \notin E(G)$ and $uv_r^+ \notin E(G)$. Now $\{v_r, v_r^-, v_r^+, u\}$ induces a claw. Since G is 1-heavy and u is light, v_r^- is heavy or v_r^+ is heavy.

Claim 7.1. v_r^+ is heavy.

Proof. Suppose that v_r^- is heavy. By Lemma 3 (b), (c) and (e), v_2^- , v_1 and l_1^- are light. By Claim 6, $\{l_1, l_1^-, v_1, v_2^-\}$ induces a light claw, a contradiction.

Claim 7.2. v_1^+, v_2^+, v_1 and l_1^+ are light.

Proof. By Claim 7.1, v_r^+ is heavy. By Lemma 3 (b), (c) and (e), v_1^+, v_2^+, v_1 and l_1^+ are light.

Claim 7.3. $v_1 l_1^+ \notin E(G)$.

Proof. If $v_1l_1^+ \in E(G)$, then consider the subgraph induced by $\{v_1, u, v_1^+, l_1^+\}$. It is obvious that $uv_1^+ \notin \widetilde{E}(G)$ and $ul_1^+ \notin \widetilde{E}(G)$. By Claim 5 and Lemma 3 (f), we have $v_1^+l_1^+ \notin E(G)$. By Claim 7.2, $\{v_1, u, v_1^+, l_1^+\}$ induces a light claw, a contradiction.

Now consider the subgraph induced by $\{l_1, l_1^+, v_1, v_2^+\}$. By Claim 5 and Lemma 3 (c), (e), we have $v_1v_2^+ \notin \widetilde{E}(G)$ and $l_1^+v_2^+ \notin \widetilde{E}(G)$. By Claim 7.3, we have $v_1l_1^+ \notin E(G)$. By Claim 7.2, $\{l_1, l_1^+, v_1, v_2^+\}$ induces a light claw.

By Claim 7, a contradiction.

Subcase 2.2. $v_r^- v_r^+ \in E(G)$.

Claim 8. v_1 is heavy.

Proof. By Claim 2, there exists a vertex $l_1 \in C[v_1^+, v_2^-)$ such that $v_2^-l_1 \in E(G)$ and $l_1v_1 \in E(G)$. By Claims 5 and 6, $v_2^+l_1 \in E(G)$, $\{l_r, l_r^-, v_r, v_1^-\}$ and $\{l_1, l_1^-, v_1, v_2^-\}$ induce claws.

Suppose that v_2^- is heavy. By Lemma 3 (b), (c) and (e), we have v_1^- , v_r and l_r^- are light. Now $\{l_r, l_r^-, v_r, v_1^-\}$ induces a light claw, contradicting that G is 1-heavy.

Suppose that l_1^- is heavy. By Claim 8, Lemma 3 (e) and (f), since $v_2^+ l_1 \in E(G)$, we have $v_r l_1^- \notin \widetilde{E}(G)$ and $v_1^- l_1^- \notin \widetilde{E}(G)$. This implies that v_r, v_1^- are light. At the same time, we can prove that l_r^- is light (otherwise, $C' = v_r u v_2 \overleftarrow{C} [v_2, l_1] l_1 v_2^+ C[v_2^+, v_r^-] v_r^- v_r^+ C[v_r^+, l_r^-] l_r^- l_1^- \overleftarrow{C} [l_1^-, l_r] l v_r$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction). Now $\{l_r, l_r^-, v_r, v_1^-\}$ induces a light claw, contradicting that G is 1-heavy.

Note that $\{l_1, l_1^-, v_1, v_2^-\}$ induces a claw and v_2^-, l_1^- are light. We can see v_1 is heavy since G is 1-heavy.

By Claim 2, there exists a vertex $l_2' \in C(v_1^+, v_2^-]$ such that $v_1^+ l_2' \in E(G)$ and $l_2' v_2 \in E(G)$.

Claim 9. $l_r^-, l_r^+, {l_2'}^-, {l_2'}^+$ are light.

Proof. If l_r^+ is heavy, then $C' = v_r u v_1 l_r^+ C[l_r^+, v_1^-] v_1^- v_1^+ C[v_1^+, v_r^-] v_r^- v_r^+ C[v_r^+, l_r] l_r v_r$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction. If l_r^- is heavy, then $C' = v_r l_r C[l_r, v_1^-] v_1^- v_1^+ C[v_1^+, v_r^-] v_r^- v_r^+ C[v_r^+, l_r^-] l_r^- v_1 u v_r$ is an o-cycle such that $V(C) \subset V(C')$, a contradiction.

Similarly, by symmetry, we can prove that $l_2'^-, l_2'^+$ are light.

Claim 10. v_r is heavy.

Proof. Consider the subgraph induced by $\{l_r, l_r^-, l_r^+, v_r\}$. By Claim 5 and Lemma 3 (e), we have $v_r l_r^- \notin \widetilde{E}(G)$. By Lemma 3 (d), we have $l_r^- l_r^+ \notin \widetilde{E}(G)$ and $v_r l_r^+ \notin \widetilde{E}(G)$. Since G is 1-heavy, v_r is heavy by Claim 9.

Claim 11. $\{v_1^-, v_1^+, l_r^+, l_2^{\prime+}\}$ induces a light claw.

Proof. By Claim 9, $l_r^+, l_2'^+$ are light. Note that $v_1^-v_1^+ \in E(G)$. Now we suffice to check the following facts: $\{v_1^-l_r^+, v_1^-l_2'^+\} \subset E(G)$ and $\{v_1^+l_r^+, v_1^+l_2'^+, l_r^+l_2'^+\} \cap E(G) = \emptyset$.

Claim 11.1. $v_1^- l_r^+ \in E(G)$ and $v_1^- l_2'^+ \in E(G)$.

Proof. Suppose that $v_1^- l_r^+ \notin E(G)$. By Lemma 3 (d) and (e), we have $l_r^- l_r^+ \notin \widetilde{E}(G)$ and $v_1^- l_r^- \notin \widetilde{E}(G)$. Now $\{l_r, l_r^-, l_r^+, v_1^-\}$ induces a light claw, a contradiction.

By Claim 5 and symmetry, we have $v_1^- l_2' \in E(G)$. Suppose that $v_1^- l_2'^+ \notin E(G)$. By Lemma 3 (d) and (e), we have $l_2'^- l_2'^+ \notin \widetilde{E}(G)$ and $v_1^- l_2'^- \notin \widetilde{E}(G)$. Now $\{l_2', l_2'^-, l_2'^+, v_1^-\}$ induces a light claw, a contradiction.

Claim 11.2. $\{v_1^+l_r^+, v_1^+l_2'^+, l_r^+l_2'^+\} \cap E(G) = \emptyset.$

Proof. By Lemma 3 (e), since $v_r l_r \in E(G)$ and $v_2 l_2' \in E(G)$, we have $v_1^+ l_r^+ \notin \widetilde{E}(G)$ and $v_1^+ l_2'^+ \notin \widetilde{E}(G)$. At the same time, we can prove that $l_r^+ l_2'^+ \notin E(G)$, since otherwise, $C' = v_r u v_2 l_2' \overleftarrow{C}[l_2', l_r^+] l_r^+ l_2'^+ C[l_2'^+, v_2^-] v_2^- v_2^+ C[v_2^+, v_r^-] v_r^- v_r^+ C[v_r^+, l_r] l_r v_r$ is an ocycle such that $V(C) \subset V(C')$, a contradiction.

It is proved that $\{v_1^-, v_1^+, l_r^+, l_2^{\prime+}\}$ induces a light claw.

By Claim 11, a contradiction. The proof of Theorem 4 is complete. \Box

References

- [1] M. Aider, Almost distance-hereditary graphs, Discrete Math. 242 (2002) 1-16.
- [2] D. Alessandro and M. Marina, Distance-hereditary graphs, Steiner trees, and connected domination, SIAM J. Comput. 17 (3) (1988) 521-538.
- [3] B. Bollobás and G. Brightwell, Cycles through specified vertices, Combinatorica 13 (1993) 147-155.
- [4] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications, Macmillan London and Elsevier, New York (1976).
- [5] H.J. Broersma, Z. Ryjáček and P. Vrana, How many conjectures can you stand? A survey, *Graphs and Combin.* 28 (2012) 57-75.
- [6] H.J. Broersma, Z. Ryjáček and I. Schiermeyer, Dirac's minimum degree condition restricted to claws, *Discrete Math.* 167/168 (1997) 155-166.
- B. Chen, S. Zhang and S. Qiao, Hamilton cycles in claw-heavy graphs, *Discrete Math.* 309 (2009) 2015-2019.
- [8] O. Cogis and E. Thierry, Computing maximum stable sets for distance-hereditary graphs, *Discrete Optimization* **2** (2) (2005) 185-188.
- [9] R.J. Faudree, E. Flandrin and Z. Ryjáček, Claw-free graphs- A survey, *Discrete Math.* 164 (1997) 87-147.
- [10] J. Feng and Y. Guo, Hamiltonian problem on claw-free and almost distance-hereditary graphs, *Discrete Math.* **308** (24) (2008) 6558-6563.

- [11] J. Feng and Y. Guo, Hamiltonian cycle in almost distance-hereditary graphs with degree condition restricted to claws, *Optimazation* 57 (1) (2008) 135-141.
- [12] J. Fujisawa and T. Yamashita, Degree conditions on claws and modified claws for Hamiltonicity of graphs, *Discrete Math.* 9 (308) (2008) 1612-1619.
- [13] E. Howorka, A characterization of distance-hereditary graphs, Quart. J. Math. Oxford Ser. 28 (112) (1977) 417-420.
- [14] S. Hsieh, C. Ho, T. Hsu, and M. Ko, Efficient Algorithms for the Hamiltonian Problem on Distance-Hereditary Graphs, *Computing and Combinatorics*, Lecture Notes in Computer Science 2387 (2002) 77-86.
- [15] R.M. Karp, Reducibility among combinatorial problems, In: R.E. Miler et al: Complexity of Computer Computations, New York, Plenum Press (1972), pp. 85-103.
- [16] B. Li, Z. Ryjáček, Y. Wang and S. Zhang, Pairs of heavy subgraphs for Hamiltonicity of 2-connected graphs, SIAM J. Discrete Math. 26 (3) (2012) 1088-1103.
- [17] M.M Matthews and D.P. Sumner, Hamiltonian results in $K_{1,3}$ -free graphs, J. Graph Theory 8 (1984) 139-146.
- [18] R. Shi, 2-Neighborhoods and hamiltonian conditions, J. Graph Theory 16 (1992) 267-271.