ON THE DIFFERENTIABLE VECTORS FOR CONTRAGREDIENT REPRESENTATIONS

INGRID BELTIŢĂ AND DANIEL BELTIŢĂ

ABSTRACT. We establish a few simple results on contragredient representations of Lie groups, with a view toward applications to the abstract characterization of some spaces of pseudo-differential operators. In particular, this method provides an abstract approach to J. Nourrigat's recent description of the norm closure of the pseudo-differential operators of order zero.

1. Introduction

In this note we study the abstract characterization of some spaces of pseudo-differential operators by using a few simple results on the contragredients of Banach space representations of Lie groups. The applicability of the method based on a contragredient representation is due to the fact that such a representation may be discontinuous even if the original representation is continuous; see for instance the representation (3.1) below, which is discontinuous if $r = \infty$. In particular, we provide an abstract approach to J. Nourrigat's recent description [No12] of the norm closure of the pseudo-differential operators of order zero (see Example 3.3 below) and we also bring additional information on some results from the earlier literature.

Preliminaries. For any complex Banach space \mathcal{Y} we denote by \mathcal{Y}^* its topological dual and by $\mathcal{B}(\mathcal{Y})^{\times}$ the group of invertible elements in the Banach algebra $\mathcal{B}(\mathcal{Y})$ of all bounded linear operators. If G is any group, then a Banach space representation of G is a group homomorphism $\pi\colon G\to \mathcal{B}(\mathcal{Y}_{\pi})^{\times}$, where \mathcal{Y}_{π} is a complex Banach space. The contragredient representation of π is the representation

$$\pi^*: G \to \mathcal{B}(\mathcal{Y}_{\pi}^*)^{\times}, \quad \pi^*(g) := \pi(g^{-1})^*,$$

so that $\mathcal{Y}_{\pi^*} := \mathcal{Y}_{\pi}^*$. If $\sup_{g \in G} \|\pi(g)\| < \infty$, then we say that π is uniformly bounded, and in this case also π^* is uniformly bounded.

Now assume that G is a topological group and for the uniformly bounded representation $\pi\colon G\to \mathcal{B}(\mathcal{Y}_\pi)^\times$ define $\mathcal{Y}_{\pi_0}:=\{x\in\mathcal{Y}_\pi\mid \pi(\cdot)x\in\mathcal{C}(G,\mathcal{Y}_\pi)\}$, where \mathcal{C} indicates the space of continuous mappings. Then \mathcal{Y}_{π_0} is a closed linear subspace of \mathcal{Y} since π is uniformly bounded, and moreover \mathcal{Y}_{π_0} is invariant under π . Hence we obtain a strongly continuous representation $\pi_0\colon G\to \mathcal{B}(\mathcal{Y}_{\pi_0}), \, \pi_0(g):=\pi(g)|_{\mathcal{Y}_{\pi_0}}$. By using this construction for the contragredient representation, we define

$$\mathcal{Y}_{\pi_0^*} := \{ \xi \in \mathcal{Y}_\pi^* \mid \pi^*(\cdot) \xi \in \mathcal{C}(G, \mathcal{Y}_\pi^*) \} = \{ \xi \in \mathcal{Y}_\pi^* \mid \lim_{g \to 1} \| \pi^*(g) \xi - \xi \| = 0 \}$$

Date: June 22, 2013.

²⁰¹⁰ Mathematics Subject Classification. Primary 22E45; Secondary 47G30, 47B10. Key words and phrases. contragredient representation; differentiable vector; commutator.

and

$$\pi_0^* : G \to \mathcal{B}(\mathcal{Y}_{\pi_0^*})^{\times}, \quad \pi_0^*(g) := \pi^*(g)|_{\mathcal{Y}_{\pi_0^*}}.$$

If moreover G is a Lie group, then we also define $\mathcal{Y}_{\pi}^{k} := \{y \in \mathcal{Y} \mid \pi(\cdot)y \in \mathcal{C}^{k}(G,\mathcal{Y})\}$ for every integer $k \geq 0$, so in particular $\mathcal{Y}_{\pi}^{0} = \mathcal{Y}_{\pi_{0}}$. Moreover, if the representation π is strongly continuous, that is, $\mathcal{Y} = \mathcal{Y}_{\pi_{0}}$, then for every $k \geq 1$ and every basis $\{X_{1}, \ldots, X_{m}\}$ in the Lie algebra \mathfrak{g} of G we have

$$\mathcal{Y}_{\pi}^{k} = \bigcap_{1 \leq j_{1}, \dots, j_{k} \leq m} \mathcal{D}(\mathrm{d}\pi(X_{j_{1}}) \cdots \mathrm{d}\pi(X_{j_{k}}))$$

$$(1.1)$$

(see for instance [Ne10, Th. 9.4]). Here and in what follows we denote by $\mathcal{D}(T)$ the domain of any unbounded operator T.

2. The main abstract results

The following theorem can be regarded as a version of (1.1) for some discontinuous representations of Lie groups, namely for the contragredient of any uniformly bounded and strongly continuous representation.

Theorem 2.1. Let G be a Lie group with a strongly continuous representation $\pi: G \to \mathcal{B}(\mathcal{Y})$ which is also assumed to be uniformly bounded. If $\{X_1, \ldots, X_m\}$ is any basis in the Lie algebra \mathfrak{g} of G, then

$$\mathcal{Y}_{\pi^*}^k \subseteq \bigcap_{1 \le j_1, \dots, j_k \le m} \mathcal{D}(\mathrm{d}\pi(X_{j_1})^* \cdots \mathrm{d}\pi(X_{j_k})^*) \subseteq \mathcal{Y}_{\pi^*}^{k-1}$$

for every integer $k \geq 1$, and the above inclusions could simultaneously be strict.

For proving the theorem it will be convenient to use the notation

$$\mathcal{C}^k(\pi^*) := \bigcap_{1 \le j_1, \dots, j_k \le m} \mathcal{D}(\mathrm{d}\pi(X_{j_1})^* \cdots \mathrm{d}\pi(X_{j_k})^*)$$

for an arbitrary integer $k \geq 1$. It is clear that $C^1(\pi^*) \supseteq C^2(\pi^*) \supseteq \cdots$.

The proof will be based on the following auxiliary result, which should be thought of as an embedding lemma on abstract Sobolev spaces.

Lemma 2.2. We have $C^1(\pi^*) \subseteq \mathcal{Y}_{\pi_0^*}$.

Proof. For every $X \in \mathfrak{g}$ let us denote $\gamma_X : \mathbb{R} \to G$, $\gamma_X(t) := \exp_G(tX)$. It follows by [vNe92, Th. 1.3.1] that

$$\mathcal{D}(\mathrm{d}\pi(X)^*) \subseteq \mathcal{Y}_{\pi^* \circ \gamma_X} = \{ \xi \in \mathcal{Y}^* \mid \pi^*(\gamma_X(\cdot))\xi \in \mathcal{C}(\mathbb{R}, \mathcal{Y}^*) \}$$
 (2.1)

for arbitrary $X \in \mathfrak{g}$. On the other hand, we have

$$\mathcal{Y}_{\pi^*} = \bigcap_{i=1}^m \mathcal{Y}_{\pi^* \circ \gamma_{X_j}} \tag{2.2}$$

since the inclusion \subseteq is obvious while the inclusion \supseteq holds true for the following reason. For all $t_1, \ldots, t_m \in \mathbb{R}$ and $\xi \in \mathcal{Y}^*$ we have

$$\|\pi^*(\gamma_{X_1}(t_1)\cdots\gamma_{X_m}(t_m))\xi - \xi\|$$

$$\leq \sum_{j=1}^m \|\pi^*(\gamma_{X_1}(t_1)\cdots\gamma_{X_{j-1}}(t_{j-1}))(\pi^*(\gamma_{X_j}(t_j))\xi - \xi)\|$$

$$\leq M\sum_{j=1}^m \|\pi^*(\gamma_{X_j}(t_j))\xi - \xi\|$$

where $M:=\sup_{g\in G}\|\pi(g)\|$. Since $\{X_1,\ldots,X_m\}$ is a basis in the Lie algebra \mathfrak{g} , it follows that the mapping $\mathbb{R}^m\to G,\ (t_1,\ldots,t_m)\mapsto \gamma_{X_1}(t_1)\cdots\gamma_{X_m}(t_m)$, is a local diffeomorphism at $0\in\mathbb{R}^m$, and then the above estimate shows that for every $\xi\in\bigcap_{j=1}^m\mathcal{Y}_{\pi^*\circ\gamma_{X_j}}$ we have $\lim_{g\to 1}\|\pi^*(g)\xi-\xi\|=0$, hence $\xi\in\mathcal{Y}_{\pi^*}$. This completes the proof of (2.2).

Now, since $\mathcal{D}(d\pi(X_1)^*) \cap \cdots \cap \mathcal{D}(d\pi(X_m)^*) = \mathcal{C}^1(\pi^*)$, the assertion follows by (2.1) and (2.2).

Proof of Theorem 2.1. By using Lemma 2.2 and [Po72, Lemma 1.1] we obtain

$$\mathcal{C}^{k}(\pi^{*}) \subseteq \bigcap_{1 \le j_{1}, \dots, j_{k-1} \le m} \mathcal{D}(\mathrm{d}\pi_{0}^{*}(X_{j_{1}}) \cdots \mathrm{d}\pi_{0}^{*}(X_{j_{k-1}})) = \mathcal{Y}_{\pi^{*}}^{k-1}$$

where the latter equality follows by using (1.1) for the strongly continuous representation π_0^* . The inclusion $\mathcal{Y}_{\pi^*}^k \subseteq \mathcal{C}^k(\pi^*)$ can be easily proved by using (1.1) and the fact that for every $X \in \mathfrak{g}$ we have $\mathcal{D}(\mathrm{d}\pi_0^*(X)) \subset \mathcal{D}(\mathrm{d}\pi(X)^*)$ and $\mathrm{d}\pi(X)^*|_{\mathcal{D}(\mathrm{d}\pi_0^*(X))} = \mathrm{d}\pi_0^*(X)$.

We now prove by example that the inclusion in the statement can be strict for k=1. To this end let $G=\mathbb{R}$, \mathcal{Y} be the space of trace-class operators on $L^2(\mathbb{R})$, and consider the regular representation $\rho \colon \mathbb{R} \to \mathcal{B}(L^2(\mathbb{R}))$, $\rho(t)f = f(\cdot + t)$. Then define $\pi \colon \mathbb{R} \to \mathcal{B}(\mathcal{Y})$, $\pi(t)A = \rho(t)A\rho(t)^{-1}$ and for every $\phi \in L^{\infty}(\mathbb{R})$ denote by $\phi(Q)$ the multiplication-by- ϕ operator on $L^2(\mathbb{R})$, so that $\phi(Q) \in \mathcal{B}(L^2(\mathbb{R})) \simeq \mathcal{Y}^*$. It was noted in [ABG96, Ex. 6.2.7] that $\phi(Q) \in \mathcal{Y}^k_{\pi^*}$ if and only the first k derivatives of ϕ exist, are bounded, and the k-th derivative is also uniformly continuous on \mathbb{R} .

On the other hand, if we denote by $P=-\mathrm{i}\frac{\mathrm{d}}{\mathrm{d}t}$ the infinitesimal generator of ρ , then it is easily checked that $\phi(Q)\in\mathcal{C}^1(\pi^*)$ if and only if the commutator $[\phi(Q),P]$ belongs to $\mathcal{B}(L^2(\mathbb{R}))$, hence by using also [ABG96, Prop. 5.1.2(b)] and again [ABG96, Ex. 6.2.7] we see that the latter commutator condition is equivalent to the fact that ϕ is bounded and satisfies the Lipschitz condition globally on \mathbb{R} . Therefore there exist $\phi,\psi\in L^\infty(\mathbb{R})$ such that $\phi(Q)\in\mathcal{C}^1(\pi^*)\setminus\mathcal{Y}^1_{\pi^*}$ and $\psi(Q)\in\mathcal{Y}^2_{\pi^*}\setminus\mathcal{C}^1(\pi^*)$. This completes the proof.

Corollary 2.3. In the setting of Theorem 2.1, the linear subspace

$$\bigcap_{k\geq 1}\bigcap_{1\leq j_1,\ldots,j_k\leq m}\mathcal{D}(\mathrm{d}\pi(X_{j_1})^*\cdots\mathrm{d}\pi(X_{j_k})^*)$$

is dense in $\mathcal{Y}_{\pi_0^*}$.

Proof. It follows by Theorem 2.1 that this linear subspace is equal to the space of smooth vectors for the strongly continuous representation π_0^* , hence it is dense in the representation space $\mathcal{Y}_{\pi_0^*}$ (see [Gå47]).

3. Applications

In this section we will develop a more general version of the example used in the proof of Theorem 2.1. Let G be a Lie group with a continuous unitary representation $\rho \colon G \to \mathcal{B}(\mathcal{H})$. If $1 \leq p < \infty$, denote by $\mathfrak{S}_p(\mathcal{H})$ the p-th Schatten ideal, and let $\mathfrak{S}_{\infty}(\mathcal{H}) := \mathcal{B}(\mathcal{H})$ and $\mathfrak{S}_0(\mathcal{H})$ be the ideal of all compact operators on \mathcal{H} . It is well known that if $p, q \in \{0\} \cup [1, \infty]$ with $\frac{1}{p} + \frac{1}{q} = 1$ and $p \neq \infty$, then there exists an isometric linear isomorphism $\mathfrak{S}_p(\mathcal{H})^* \simeq \mathfrak{S}_q(\mathcal{H})$ defined by the duality pairing

$$\langle \cdot, \cdot \rangle \colon \mathfrak{S}_q(\mathcal{H}) \times \mathfrak{S}_p(\mathcal{H}) \to \mathbb{C}, \quad \langle Y, V \rangle := \operatorname{Tr}(YV).$$

The representation $\rho^{(q)}$ can thus be regarded as the contragredient representation of the *strongly continuous* representation $\rho^{(p)}$, where

$$(\forall r \in \{0\} \cup [1, \infty]) \quad \rho^{(r)} \colon G \to \mathcal{B}(\mathfrak{S}_r(\mathcal{H})), \quad \rho^{(r)}(g)Y = \rho(g)Y\rho(g)^{-1} \qquad (3.1)$$
 (see also [BB10]).

Here is a consequence of the results from the previous section. In the special case of the Heisenberg group, this establishes a direct relationship between the classical characterizations of pseudo-differential operators from [Be77] and [Co79].

Corollary 3.1. In the above setting, pick any basis $\{X_1, \ldots, X_m\}$ in the Lie algebra \mathfrak{g} of G. Assume $1 \leq q \leq \infty$ and denote

$$\Psi_q(\rho) := \{ Y \in \mathfrak{S}_p(\mathcal{H}) \mid \rho^{(q)}(\cdot)Y \in \mathcal{C}^{\infty}(G, \mathfrak{S}_p(\mathcal{H})) \}.$$

Then the following assertions hold:

(1) The linear subspace $\Psi_q(\rho)$ is precisely the set of all $Y \in \mathfrak{S}_q(\mathcal{H})$ for which

$$[d\rho(X_{j_1}),\ldots,[d\rho(X_{j_k}),Y]\ldots]\in\mathfrak{S}_q(\mathcal{H})$$

for arbitrary $k \geq 1$ and $j_1, \ldots, j_k \in \{1, \ldots, m\}$.

(2) If $1 \leq q < \infty$, then $\Psi_q(\rho)$ is dense in $\mathfrak{S}_q(\mathcal{H})$. If $q = \infty$, then $\Psi_\infty(\rho)$ contains the ideal of compact operators on \mathcal{H} and is dense in the norm-closed subspace $\{Y \in \mathcal{B}(\mathcal{H}) \mid \rho^{(\infty)}(\cdot)Y \in \mathcal{C}(G,\mathcal{B}(\mathcal{H}))\}$ of $\mathcal{B}(\mathcal{H})$.

Proof. We have that

$$\mathcal{C}^{1}(\rho^{(q)}) = \{ Y \in \mathfrak{S}_{q}(\mathcal{H}) \mid [\mathrm{d}\rho(X_{j}), Y] \in \mathfrak{S}_{q}(\mathcal{H}) \text{ for } j = 1, \dots, m \}.$$

Then both assertions are special cases of Theorem 2.1 and Corollary 2.3. $\hfill\Box$

We can now prove a corollary which shows that the first two conditions in [Me00, Th. 1] are equivalent irrespective of the unitary representation involved therein. This also shows that the \mathcal{C}^{∞} part of the relation between differentiability and existence of commutators suggested after [Co95, Eq. (8.4)] holds true although the \mathcal{C}^1 part of that suggestion fails to be true, since the following corollary would be false with the class \mathcal{C}^{∞} replaced by \mathcal{C}^k for any $k < \infty$. In fact, recall from the proof of Theorem 2.1 that the corresponding inclusions are strict in a special instance of the present setting, which is precisely the special instance referred to in [Co95].

Corollary 3.2. If $Y \in \mathcal{B}(\mathcal{H})$ then the above mapping $\rho^{(\infty)}(\cdot)Y \colon G \to \mathcal{B}(\mathcal{H})$ is of class \mathcal{C}^{∞} with respect to the norm operator topology on $\mathcal{B}(\mathcal{H})$ if and only if it is \mathcal{C}^{∞} with respect to the strong operator topology.

Proof. The mapping $\rho^{(\infty)}(\cdot)Y \colon G \to \mathcal{B}(\mathcal{H})$ is smooth with respect to any topology on $\mathcal{B}(\mathcal{H})$ if and only if it is smooth on any neighborhood of $\mathbf{1} \in G$. On the other hand, just as in the proof of [ABG96, Prop. 5.1.2(b)], one can see that this mapping is smooth with respect to the strong operator topology on $\mathcal{B}(\mathcal{H})$ if and only if the iterated commutator condition in Corollary 3.1(1) is satisfied, hence the conclusion follows by Corollary 3.1(1), where the smoothness of $\rho^{(\infty)}(\cdot)Y$ is understood with respect to the norm operator topology on $\mathfrak{S}_{\infty}(\mathcal{H}) = \mathcal{B}(\mathcal{H})$.

Example 3.3. Let $G = \mathbb{H}_{2n+1}$ be the (2n+1)-dimensional Heisenberg group with the Schrödinger representation $\rho \colon G \to \mathcal{B}(\mathcal{H})$. As recalled in [No12] for $1 \leq p \leq \infty$, the set $\Psi_p(\rho)$ of the above Corollary 3.1 is precisely the set of pseudo-differential operators on $L^2(\mathbb{R}^n)$ corresponding to the space of symbols

$$\{a \in \mathcal{C}^{\infty}(\mathbb{R}^{2n}) \mid (\forall \alpha \in \mathbb{N}^{2n}) \ \partial^{\alpha} a \in L^p(\mathbb{R}^{2n})\}$$

(see also [BB12] for similar results on more general nilpotent Lie groups). Thus our Corollary 3.1 leads to the main results of [No12].

Example 3.4. The above Corollary 3.1 also provides additional information in the setting of pseudo-differential operators on a compact manifold acted on by a Lie group, as studied for instance in [Ta97] and [Me00]. Thus, it follows that the notions of U-smoothness and \mathfrak{A} -smoothness from [Ta97, Sect. 2] actually coincide.

Remark 3.5. It would be interesting to extend the above result of Ex. 3.3 to the setting of the magnetic Weyl calculus of [IMP10]. Such an extension is likely to require infinite-dimensional Lie groups.

Acknowledgment. This research has been partially supported by the Grant of the Romanian National Authority for Scientific Research, CNCS-UEFISCDI, project number PN-II-ID-PCE-2011-3-0131.

References

- [ABG96] W.O. AMREIN, A. BOUTET DE MONVEL, V. GEORGESCU, Co-groups, commutator methods and spectral theory of N-body Hamiltonians. Progress in Mathematics, 135. Birkhäuser Verlag, Basel, 1996.
- [Be77] R. BEALS, Characterization of pseudodifferential operators and applications. Duke Math. J. 44 (1977), no. 1, 45–57.
- [BB10] I. BELTIŢĂ, D. BELTIŢĂ, Smooth vectors and Weyl-Pedersen calculus for representations of nilpotent Lie groups. An. Univ. Bucureşti Mat. 58 (2010), no.1, 17–46.
- [BB12] I. Beltiţă, D. Beltiţă, Boundedness for Weyl-Pedersen calculus on flat coadjoint orbits. Preprint arXiv:1203.0974v1 [math.AP], 2012.
- [Co79] H.O. CORDES, On pseudo-differential operators and smoothness of special Lie-group representations. manuscr. math. 28 (1979), nos. 1-3, 51-69.
- [Co95] H.O. CORDES, The technique of pseudodifferential operators. London Mathematical Society Lecture Note Series, 202. Cambridge University Press, Cambridge, 1995.
- [Gå47] L. GÅRDING, Note on continuous representations of Lie groups. Proc. Nat. Acad. Sci. U.S.A. 33 (1947), 331–332.
- [IMP10] V. IFTIMIE, M. MĂNTOIU, R. PURICE, Commutator criteria for magnetic pseudodifferential operators. Comm. Partial Differential Equations 35 (2010), no. 6, 1058–1094.
- [Me00] S.T. Melo, Smooth operators for the regular representation on homogeneous spaces. Studia Math. 142 (2000), no. 2, 149–157.
- [Ne10] K.-H. NEEB, On differentiable vectors for representations of infinite dimensional Lie groups. J. Funct. Anal. 259 (2010), no. 11, 2814–2855.
- [vNe92] J. VAN NEERVEN, The adjoint of a semigroup of linear operators. Lecture Notes in Mathematics, 1529. Springer-Verlag, Berlin, 1992.

- [No12] J. NOURRIGAT, Closure of the set of pseudodifferential operators. C. R. Math. Acad. Sci. Paris 350 (2012), no. 7-8, 355–358.
- [Po72] N.S. POULSEN, On C^{∞} -vectors and intertwining bilinear forms for representations of Lie groups. J. Functional Analysis 9 (1972), 87–120.
- [Ta97] M.E. TAYLOR, Beals-Cordes-type characterizations of pseudodifferential operators. Proc. Amer. Math. Soc. 125 (1997), no. 6, 1711–1716.

Institute of Mathematics "Simion Stoilow" of the Romanian Academy, P.O. Box 1-764, Bucharest, Romania

 $E\text{-}mail\ address: \texttt{Ingrid.Beltita@imar.ro, ingrid.beltita@gmail.com}$

 $E\text{-}mail\ address: \texttt{Daniel.Beltita@imar.ro, beltita@gmail.com}$