SU(2)-IRREDUCIBLY COVARIANT AND EPOSIC CHANNELS

Muneerah Al Nuwairan

Department of Mathematics

University of Ottawa, Ottawa, ON.

E-mail: malnu009@uottawa

ABSTRACT. In this paper we introduce EPOSIC channels, a class of SU(2)-covariant quantum channels. We give their definition, a Kraus representation of them, and compute their Choi matrices. We show that these channels form the set of extreme points of all SU(2)-irreducibly covariant channels. We also compute their complementary channels, and their dual maps. As application of these channel, we get a new example of positive map that is not completely positive.

INTRODUCTION

Given two representations of a compact group G on Hilbert spaces K, E, one can construct G-covariant quantum channels. Indeed, the G-space $K \otimes E$ decomposes into a direct sum of G-irreducible invariant subspaces $\{H_i\}_{i \in I}$. For each $i \in I$, the Gequivariant inclusion $\alpha_i : H_i \longrightarrow K \otimes E$ provides (by a standard construction) two G-covariant channels $\Phi_i^1 : End(H_i) \longrightarrow End(K)$ and $\Phi_i^2 : End(H_i) \longrightarrow End(E)$. Unless we make further assumptions on the G-spaces K, E these channels remain difficult to study. To simplify the study of these channels, we choose K and E to be Girreducible spaces. In this case, the channels Φ_i^1, Φ_i^2 are called G-irreducibly covariant channels [6]. If G = SU(n) then Clebsch-Gordan formula [14, ch.5] describes the decomposition of the SU(n)-space $K \otimes E$ into SU(n)-irreducible subspaces and gives their multiplicity. Studying SU(n)-equivariant maps between spaces of the form $K \otimes E$, is in general difficult, due to the fact that the multiplicity of SU(n)-irreducible subspaces of $K \otimes E$ might be greater than one. However, in case of SU(2), the multiplicity of any SU(2)-irreducible subspace in $K \otimes E$ is always one, simplifying the study of such maps.

In this paper we introduce EPOSIC channels, a class of SU(2)-covariant quantum channels. We show that if H, K are SU(2)-irreducible spaces then EPOSIC channels from End(H) into End(K) forms the extreme points of the convex set of all SU(2)irreducibly covariant channels form End(H) into End(K). The first section contains definitions, lemmas and propositions from both representation theory and quantum information theory. In Section 2, we introduce our examples of quantum channels and exhibit some of their properties. In section 3, we get Kraus operators and the Choi matrix of EPOSIC channel. Section 4 shows that the set of all SU(2)-irreducibly covariant channels forms a simplex and that its extreme points are nothing but EPOSIC channels. In section 5 we compute a complementary channel and the dual map for EPOSIC channel. The paper ends with an application of the EPOSIC channels where we get an example of positive map that is not completely positive.

This work was done under the supervision of professors B. Collins and T. Giordano as a part of a PhD thesis of the author. I greatly appreciate their patient advice and help.

1. BACKGROUND DEFINITION AND RESULTS

In this section, we recall standard notions and fix the notations we will then use. We assume all vector spaces are finite dimensional. A Hilbert space H is a complex vector space endowed with an inner product $\langle | \rangle_{H}$. If H, K are Hilbert spaces, then End(H, K) denotes the vector space of linear maps from H to K. It is a Hilbert space endowed with Hilbert-Schmidt inner product given by $\langle A | B \rangle_{End(H,K)} = tr(A^*B)$ for $A, B \in End(H, K)$. For $y \in H$, let y^* denote the linear form given by $y^*(z) = \langle y | z \rangle_{H}$. If $x \in K$ then $xy^* \in End(H, K)$ denotes the map $xy^*(z) = \langle y | z \rangle_{H} x$, for any $z \in H$. Recall that $\{xy^* : x \in K, y \in H\}$ forms a set of generators of End(H, K). As usual, we write End(H) for End(H, H) and I_H for the identity map on H. To a Hilbert space H, we associate the conjugate space \overline{H} , the space \overline{H} is a vector space with the same underlying abelian group as H and with scalar multiplication $(\lambda, v) \longrightarrow \overline{\lambda} v$. The inner product on \overline{H} is then defined by $\langle h_1 | h_2 \rangle_{\overline{H}} = \langle h_2 | h_1 \rangle_H$. One can easily verify that $End(\overline{H}) = \overline{End(H)}$ and the inner products of the two spaces coincide.

1.1. The *G*-equivariant maps.

Definition 1.1. Let G be a topological group. A representation of G on a Hilbert space H is a continues homomorphism $\pi_H : G \longrightarrow GL(H)$ that makes H a G-space. If $\pi_{H(g)}$ is a unitary operator for each $g \in G$ then π_H is called unitary representation. We restrict the use of the symbol ρ_H to an irreducible representation on H.

If W is a subspace of H such that $\pi_{H(g)}(w) \in W$ for any $g \in G$. Then π_H can be made into a representation of G on W, called subrepresentation. In such case, W is called G-invariant subspace of H.

There are standard methods for constructing a new representation from given ones. For example if π_H, π_K are two representations of G on H, K, then the maps given by $g \mapsto \pi_H(g) \otimes \pi_K(g)$ and $g \longrightarrow l(g)$ where $l(g)(A) = \pi_K(g)A\pi_H(g^{-1})$, for $A \in End(H, K)$, are representations of G on the Hilbert spaces $H \otimes K$ and End(H, K) respectively.

If $\pi : G \longrightarrow GL(H)$ is a representation of G then $\check{\pi} : G \longrightarrow GL(\overline{H})$ defined by $\check{\pi}_{(g)} = \pi^t_{(g^{-1})}$ for each $g \in G$ (t denotes the transpose map), and $\overline{\pi} : G \longrightarrow GL(\overline{H})$ defined by $\overline{\pi}_{(g)} = \overline{\pi}_{(g)}$ for each $g \in G$, are representations of G on the Hilbert space \overline{H} . These representations are known as the contragredient and the conjugate representation of G respectively. These two representations on \overline{H} coincide if π is unitary representation.

Definition 1.2. Let G be a group and $\pi_H : G \longrightarrow GL(H)$ is a representation of G on the Hilbert space H, the commutant of $\pi_H(G)$, denoted by $\pi_H(G)'$, is the set $\{T \in End(H) : T\pi_H(g) = \pi_H(g)T \; \forall g \in G\}.$

Remark 1.3. If π_H is a representation of G on a Hilbert space H, and W is a subspace of H, then W is a G-invariant subspace of H if and only if q_W , the orthogonal projection on W, belongs to $\pi_H(G)'$.

Definition 1.4. Let G be a group, and π_H, π_K be representations of G on the Hilbert spaces H, K. A linear map $\alpha : H \longrightarrow K$ is said to be G-equivariant if $\pi_{K(g)}\alpha = \alpha \pi_{H(g)}$ for all $g \in G$. We denote the set of G-equivariant maps from $H \longrightarrow K$ by $End(H, K)^G$.

Example 1.5. Let G be a compact group, and π_H , π_K be representations of G on the Hilbert spaces H, K, then the following are standard constructions of G-equivariant maps.

- (1) If $\alpha: H \longrightarrow K$ is a *G*-equivariant then so is $\alpha^*: K \longrightarrow H$.
- (2) The map $flip_{K}^{H}: H \otimes K \longrightarrow K \otimes H$ given by linearly extending $h \otimes k \longrightarrow k \otimes h$ is *G*-equivariant map. It is a unitary with adjoint $(flip_{K}^{H})^{*} = flip_{H}^{K}$ and satisfy $Tr_{K}(flip_{K}^{H}Aflip_{H}^{K}) = Tr_{K}(A)$ for any $A \in End(H \otimes K)$. The map Tr_{K} denotes the partial trace [18, p.19], and it is defined on the generators set of $End(H) \otimes End(K)$ by $A_{1} \otimes A_{2} \longmapsto tr(A_{2})A_{1}$.
- (3) The natural isomorphism

$$T: End(K) \otimes End(\overline{H}) \longrightarrow End(K \otimes \overline{H})$$

defined by the map taking $A \otimes B$ to $T(A \otimes B)(k \otimes h) = A(k) \otimes B(h)$ and extending linearly, is a *G*-equivariant map.

(4) The map Vec : End(H, K) → K ⊗ H [18, p.23] defined to be the unique linear extension of Vec(xy*) = x ⊗ y, is an example of G-equivariant map. It represents any element in End(H, K) as a vector in the tensor product space K ⊗ H. The map Vec is a unitary in the sense that

$$\langle A | B \rangle_{End(H,K)} = \langle \operatorname{Vec}(A) | \operatorname{Vec}(B) \rangle_{K \otimes \overline{H}}$$

for any $A, B \in End(H, K)$.

(5) Let $B^* : End(H) \longrightarrow \mathbb{C}$ denote the map $B^*X = \langle B | X \rangle_{End(H)}$. Then the Choi-Jamiolkowski map

$$C: End(End(H), End(K)) \longrightarrow End(K \otimes \overline{H})$$

defined by taking AB^* to $A \otimes \overline{B}$, for $A \in End(K)$, $B \in End(H)$ and extending linearly, is a unitary *G*-equivariant map. The map *C* assigns for each Φ a unique matrix $C(\Phi)$, known as Choi matrix of Φ , that can be computed using the formula $C(\Phi) = \sum_{ij} \Phi(E_{ij}) \otimes E_{ij}$ where E_{ij} are the standard basis for End(H) [4].

Remark 1.6. The Choi-Jamiolkowski map C is the composition of the map

$$\operatorname{Vec}: End(End(H), End(K)) \longrightarrow End(K) \otimes \overline{End(H)} = End(K) \otimes End(\overline{H})$$

and the natural isomorphism $T: End(K) \otimes End(\overline{H}) \longrightarrow End(K \otimes \overline{H}).$

Proposition 1.7. Let G be a group, and π_H, π_K be representations of G on the Hilbert spaces H, K. The set of G-equivariant maps $End(H, K)^G$ forms a subspace of End(H, K). In particular, it is a convex set.

1.2. G-Covariant quantum channels.

Let H, K be Hilbert spaces. Recall that quantum channel $\Phi : End(H) \longrightarrow End(K)$ is a completely positive trace preserving map [18, p54], and that such a channel has several equivalent representations:

Proposition 1.8. [18, p.54] Let H, K be Hilbert spaces and $\Phi : End(H) \longrightarrow End(K)$ be a quantum channel, then

 A Stinespring representation (dilation) of Φ is a pair (E, α) consisting of a Hilbert space E (an environment space), and an isometry α : H → K ⊗ E such that $\Phi(A) = Tr_E(\alpha A \alpha^*)$ for any $A \in End(H)$. The map Tr_E denotes the partial trace over E.

- (2) A Kraus representation of Φ is a set of operators $T_j \in End(H, K)$, called Kraus operators, that satisfy $\Phi(A) = \sum_{j=1}^{k} T_j A T_j^*$ and $\sum_{j=1}^{k} T_j^* T_j = I_H$.
- (3) Choi representation of Φ , it states that Φ is channel if and only if its Choi matrix $C(\Phi)$ is a positive matrix satisfies $Tr_K(C(\Phi)) = I_{\overline{H}}$.

For any quantum channel a Stinespring and a Kraus representation always exist [18, p.54], but neither is unique. However, the Choi representation gives a unique characterization of a quantum channel. The following proposition [18, p.51-p.54] gives some relations between these different representations of a quantum channel.

Proposition 1.9. Let H, K be Hilbert spaces, and $\Phi : End(H) \longrightarrow End(K)$ be a quantum channel. Then

- (1) If (E, α) is a Stinespring representation of Φ , and $\{e_j : 1 \leq j \leq d_E\}$ is an orthonormal basis of E then the set of maps $T_j : H \longrightarrow K$, $1 \leq j \leq d_E$ given by $T_j = (I_K \otimes e_j^*) \alpha$ forms a Kraus operators of Φ .
- (2) If $\{T_j : 1 \le j \le k\}$ is a Kraus operators of Φ , then the Choi matrix of Φ is given by $C(\Phi) = \sum_{j=1}^k \operatorname{Vec}(T_j) \operatorname{Vec}(T_j)^*$.

Remark 1.10. As a corollary to the last proposition and Theorem 5.3 in [18, p.51], the rank of the Choi matrix of Φ gives an achievable lower bound for both the number of any Kraus operators, and of the dimension of any environment space.

Remark 1.11. The three representations of a quantum channel are exist in general for any completely positive map, see [11, ch.4] and [3]. In the case of quantum channels, more conditions come up as a consequence of being trace preserving map.

Definition 1.12. Let G be a group, and π_H, π_K be representations of G on the Hilbert spaces H, K respectively. A quantum channel $\Phi : End(H) \longrightarrow End(K)$ is

said to be G-covariant if it is a G-equivariant map from End(H) into End(K). i.e

(1.1)
$$\Phi(\pi_H(g)A\pi_H^*(g)) = \pi_K(g)\Phi(A)\pi_K^*(g)$$

for each $A \in End(H)$ and $g \in G$. If both π_H, π_K are irreducible representations, then Φ is called *G*-irreducibly covariant channel.

The set of all G-covariant channels from End(H) to End(K) is denoted by $QC(\pi_H, \pi_K)$.

Proposition 1.13. Let G be a group, and π_H, π_K be representations of G on the Hilbert spaces H, K. Then the set $QC(\pi_H, \pi_K)$ is a convex set.

Proposition 1.14. Let G be a group, and π_H, π_K be representations of G on the Hilbert spaces H, K. Let $\Phi : End(H) \longrightarrow End(K)$ be a quantum channel given by a Stinespring representation (E, α) . If $\alpha : H \longrightarrow K \otimes E$ is a G-equivariant map then Φ is a G-covariant channel.

Recall that for a Hilbert space H, the set of all density operators on H, denoted by D(H), is $\{\varrho \in End(H) : \varrho \ge 0, tr(\varrho) = 1\}$.

Proposition 1.15. [4] If G is a group, and π_H, π_K are representations of G on the Hilbert spaces H, K. Let $\Phi : End(H) \longrightarrow End(K)$ be a linear map. Then

- (1) The map Φ is G-equivariant map if and only if $C(\Phi) \in (\pi_{\kappa} \otimes \check{\pi}_{H}(G))'$.
- (2) If π_H is an irreducible representation, then Φ is G-covariant channel if and only if $\frac{1}{d_H}C(\Phi) \in (\pi_K \otimes \check{\pi}_H(G))' \bigcap D(K \otimes \overline{H}).$

For further information on these results, we refer the reader to [1],[18],[11],[6], [4] and [13].

2. EPOSIC CHANNELS

The goal of this section is to introduce our example of a class of SU(2)-irreducibly covariant channels. The first two subsections contain all the results that are needed to construct the channels.

2.1. SU(2)-irreducible representations and SU(2)-equivariant maps.

2.1.1. The irreducible representations of SU(2). For $m \in \mathbb{N}$, let P_m denote the space of homogeneous polynomials of degree m in the two variables x_1, x_2 . It is a complex vector space of dimension m + 1 with a basis consists of $\{x_1^i x_2^{m-i} : 0 \leq i \leq m\}$. The space P_{-1} will denote the zero vector space. For any $m \in \mathbb{N}$, the compact group $SU(2) = \left\{ \begin{bmatrix} a & b \\ -\bar{b} & \bar{a} \end{bmatrix} : a, b \in \mathbb{C}, |a|^2 + |b|^2 = 1 \right\}$ has a representation ρ_m on P_m given by

(2.1)
$$(\rho_m(g)f)(x_1,x_2) = f((x_1,x_2)g) = f(ax_1 - \bar{b}x_2, bx_1 + \bar{a}x_2)$$

for $f \in P_m$ and $g \in SU(2)$.

Proposition 2.1. [14, p.181], [2, p.85-p.86], and [16, p.276-p.279].

(1) For $m \in \mathbb{N}$, ρ_m is a unitary representation with respect to the inner product on P_m given by

(2.2)
$$\langle x_1^l x_2^{m-l}, x_1^k x_2^{m-k} \rangle_{P_m} = l! (m-l)! \,\delta_{lk}$$

(2) The set $\{\rho_m : m \in \mathbb{N}\}$ constitutes the full list of the irreducible representations of SU(2).

To facilitate the computations, we choose the orthonormal basis of P_m given by the functions $\{f_l^m = a_m^l x_1^l x_2^{m-l} : 0 \le l \le m\}$, where $a_m^l = \frac{1}{\sqrt{l!(m-l)!}}$. This basis is called the canonical basis of the SU(2)-irreducible space P_m . The corresponding standard basis of $End(P_m)$ will be $\{E_{lk} = f_{l-1}^m f_{k-1}^{m^*} : 1 \le l, k \le m+1\}$.

It follows directly from the definition that the action of $g = \begin{bmatrix} a & b \\ -\bar{b} & \bar{a} \end{bmatrix} \in SU(2)$ on the canonical basis for P_m will be given by

(2.3)
$$\rho_m(g)(f_l^m) = a_m^l (ax_1 - \bar{b}x_2)^l (bx_1 + \bar{a}x_2)^{m-l}$$

In particular, for $g_0 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$, we have $\rho_m(g_0)(f_l^m) = (-1)^l f_{m-l}^m$ for any $0 \le l \le m$. The element g_0 will play a special role in constructing an SU(2)-equivariant unitary map of P_m onto \overline{P}_m . We fix our notation for g_0 to be $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.

Definition 2.2. For $m \in \mathbb{N}$, define the endomorphisms

(1) $\Theta_m : P_m \longrightarrow \overline{P}_m$ by $\Theta_m \left(\sum_{l=0}^m \lambda_l f_l^m \right) = \sum_{l=0}^m \lambda_l \cdot f_l^m$, where \cdot is the multiplication in \overline{P}_m . (2) $J_m : P_m \longrightarrow \overline{P}_m$ by $J_m = \Theta_m \rho_m(g_0)$.

Proposition 2.3. For $m \in \mathbb{N}$,

- (1) Θ_m is a unitary isomorphism that satisfies $\overline{\rho_m(g)}\Theta_m = \Theta_m\rho_m(\overline{g})$ for any $g \in SU(2)$.
- (2) J_m is an SU(2)-equivariant unitary isomorphism from P_m onto \overline{P}_m .

Proof.

The first assertion is straightforward. For the second one, note that J_m is a composition of unitary maps, so is unitary. As $g_0g = \bar{g}g_0$ for any $g \in SU(2)$, we have

$$J_m \rho_m(g) = \Theta_m \rho_m(g_0) \rho_m(g) = \Theta_m \rho_m(\overline{g}) \rho_m(g_0) = \overline{\rho_m(g)} \Theta_m \rho_m(g_0) = \overline{\rho_m(g)} J_m.$$

Note that on a basis element f_l^m of $P_m,\, 0\leq l\leq m$, we have

(2.4)
$$J_m(f_l^m) = (-1)^l f_{m-l}^m , \quad J_m^*(f_l^m) = (-1)^{m-l} f_{m-l}^m$$

Recall the definition of the *flip* map in Example 1.5. By direct computations on the basic elements $f_l^m \otimes f_j^n$ of $P_m \otimes P_n$, we obtain

Proposition 2.4. For $m, n \in \mathbb{N}$, the map $flip_{P_n}^{\overline{P}_m}(J_m \otimes I_{P_n}) : P_m \otimes P_n \longrightarrow P_n \otimes \overline{P}_m$ is an SU(2)-equivariant isomorphism that satisfies

$$flip_{P_n}^{\overline{P}_m}(J_m \otimes I_{P_n}) = (I_{P_n} \otimes J_m)flip_{P_n}^{P_m}$$

2.1.2. Clebsch-Gordan decomposition and SU(2)-equivariant maps on $P_m \otimes P_n$.

For $m, n \in \mathbb{N}$, let ρ_m, ρ_n be the corresponding irreducible representations of SU(2)on P_m and P_n respectively. The new constructed representation by taking the tensor product of ρ_m and ρ_n is not necessarily irreducible. Clebsch Gordan Decomposition formula [2, p.87], gives the decomposition of $\rho_m \otimes \rho_n$ into irreducible representations, namely $\rho_m \otimes \rho_n = \bigoplus_{h=0}^{\min\{m,n\}} \rho_{m+n-2h}$. Note that the corresponding decomposition for $P_m \otimes P_n$ will be given by the formula

(2.5)
$$P_m \otimes P_n \cong \bigoplus_{h=0}^{\min\{m,n\}} P_{m+n-2h}$$

Since the representations ρ_{m+n-2h} are irreducible, by Schur Lemma [13, p.13], the direct sum in the last formula will be an orthogonal direct sum.

To obtain a concrete representation of $P_m \otimes P_n$, let $x := (x_1, x_2), y := (y_1, y_2),$ $P_m := P_m(x)$, and $P_n := P_n(y)$. We embed the tensor product $P_m(x) \otimes P_n(y)$ into $\mathbb{C}[x, y]$ as follows. Define the map $: P_m(x) \times P_n(y) \longrightarrow \mathbb{C}[x, y]$ by $(f(x), g(y)) \longmapsto$ f(x)g(y), it is a bilinear map hence extends to a linear $T : P_m(x) \otimes P_n(y) \longrightarrow \mathbb{C}[x, y]$ taking $f(x) \otimes g(y)$ to f(x)g(y). Let $P_{m,n}$ denote the vector space of polynomials in xand y of bi-degree (m, n) (homogeneous polynomials of degree m in $x = (x_1, x_2)$ and of degree n in $y = (y_1, y_2)$). The space $P_{m,n}$ has a basis consist of $\{x_1^s x_2^{m-s} y_1^t y_2^{n-t} = \frac{1}{a_s^m a_t^n} T(f_s^m \otimes f_t^n) : 0 \le s \le m, 0 \le t \le n\}$. Since the map T takes the basis of $P_m \otimes P_n$ to a basis in $P_{m,n}$, it is an isomorphism. Henceforth, we will use $P_{m,n}$ as a concrete representation of $P_m \otimes P_n$. Using this identification, we define the following polynomial maps

$$\Delta_{xy} : P_m \otimes P_n \longrightarrow P_{m+1} \otimes P_{n-1} \qquad \Delta_{yx} : P_m \otimes P_n \longrightarrow P_{m-1} \otimes P_{n+1} \\ \Delta_{xy}(f(x,y)) = \left(x_1 \frac{\partial}{\partial y_1} + x_2 \frac{\partial}{\partial y_2}\right) f(x,y) \qquad \Delta_{yx}(f(x,y)) = \left(y_1 \frac{\partial}{\partial x_1} + y_2 \frac{\partial}{\partial x_2}\right) f(x,y)$$

$$\Gamma_{xy}: P_m \otimes P_n \longrightarrow P_{m+1} \otimes P_{n+1} \qquad \qquad \Omega_{xy}: P_m \otimes P_n \longrightarrow P_{m-1} \otimes P_{n-1} \\ \Gamma_{xy}f(x,y) = (x_1y_2 - y_1x_2) f(x,y) \qquad \qquad \Omega_{xy}f(x,y) = \left(\frac{\partial}{\partial x_1}\frac{\partial}{\partial y_2} - \frac{\partial}{\partial x_2}\frac{\partial}{\partial y_1}\right) f(x,y)$$

for $f(x,y) \in P_m \otimes P_n$.

Remark 2.5. Let $f(x,y) \in P_m \otimes P_n$, then by direct computations, we have:

(1) For
$$g(x, y) \in P_{m+1} \otimes P_{n-1}$$
,
 $\langle \Delta_{xy} \left(f(x, y) \right) | g(x, y) \rangle_{P_{m+1} \otimes P_{n-1}} = \langle f(x, y) | \Delta_{yx} \left(g(x, y) \right) \rangle_{P_m \otimes P_n}$.
(2) For $g(x, y) \in P_{m+1} \otimes P_{n+1}$,
 $\langle \Gamma_{xy} \left(f(x, y) \right) | g(x, y) \rangle_{P_{m+1} \otimes P_{n+1}} = \langle f(x, y) | \Omega_{xy} \left(g(x, y) \right) \rangle_{P_m \otimes P_n}$.

By the remark above and [12, p.47], we have:

Proposition 2.6. The operators Δ_{xy} , Δ_{yx} , Γ_{xy} and Ω_{xy} are SU(2)-equivariant maps that satisfy $\Delta_{xy}^* = \Delta_{yx}$ and $\Gamma_{xy}^* = \Omega_{xy}$.

Theorem 2.7. [12, chp.3] For a polynomial f(x, y) of bi-degree (m, n) we have

$$f(x,y) = \sum_{h=0}^{\min\{m,n\}} c_{m,n,h} \Gamma^h_{xy} \Delta^{n-h}_{yx} \Delta^{n-h}_{xy} \Omega^h_{xy}(f(x,y))$$

where the coefficients $c_{m,n,h}$ are determined by induction as follows: $c_{m,0,0} = 1$ and for $n \ge 1$

$$c_{m,n,h} = \begin{cases} \frac{1}{(m+1)n} c_{m+1,n-1,h} & h=0\\ \frac{1}{(m+1)n} [c_{m-1,n-1,h-1} + c_{m+1,n-1,h}] & 0 < h < n\\ \frac{1}{(m+1)n} c_{m-1,n-1,h-1} & h=n \end{cases}$$

2.2. Forming the SU(2)-equivariant isometry $\alpha_{m,n,h}$.

We now define an isometry $\alpha_{m,n,h}$ that will be used in constructing our examples of SU(2)-covariant channels. We also find a computational formula for this isometry.

2.2.1. The isometry $\alpha_{m,n,h}$.

Definition 2.8. For $m, n, h \in \mathbb{N}$, with $0 \le h \le \min\{n, m\}$, let $\alpha_{m,n,h} : P_{m+n-2h} \longrightarrow P_m \otimes P_n$ be the map defined by

$$\alpha_{m,n,h}(f(x_1, x_2)) = \sqrt{c_{m,n,h}} \Gamma^h_{xy} \Delta^{n-h}_{yx}(f(x_1, x_2))$$

where $f(x_1, x_2)$ is a homogeneous polynomial in x_1, x_2 of degree m + n - 2h.

As a direct result of Proposition 2.6 and Theorem 2.7, we have the following lemma:

Lemma 2.9. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{n, m\}$,

(1) The adjoint map of $\alpha_{m,n,h}$ is given by $\alpha_{m,n,h}^*: P_m \otimes P_n \longrightarrow P_{m+n-2h}$ where

$$\alpha^*_{m,n,h}(f(x,y)) = \sqrt{c_{m,n,h}} \ \Delta^{n-h}_{xy} \Omega^h_{xy}(f(x,y))$$

(2) $\int_{h=0}^{\min\{m,n\}} \alpha_{m,n,h} \alpha_{m,n,h}^* = I_{P_m \otimes P_n}.$

Proposition 2.10. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, the map $\alpha_{m,n,h}$ is an SU(2)-equivariant isometry.

Proof.

By Proposition 2.6, the maps Γ_{xy}^{h} , Δ_{yx}^{n-h} are SU(2)-equivariant, so $\alpha_{m,n,h}$ is. To show that $\alpha_{m,n,h}$ is an isometry, let $0 \leq h, s \leq \min\{m,n\}$. the map $\alpha_{m,n,h}^{*}\alpha_{m,n,s}$: $P_{m+n-2s} \longrightarrow P_{m+n-2h}$ is an SU(2)-equivariant map, thus by Schur Lemma [13, p.13], we have

(2.6)
$$\alpha_{m,n,h}^* \alpha_{m,n,s} \equiv \begin{cases} 0 & if \quad h \neq s \\ \lambda I_{P_{m+n-2h}} & if \quad h = s \end{cases}$$

It remains to show that $\lambda = 1$. By Lemma 2.9 we have: $\sum_{s=0}^{\min\{m,n\}} \alpha_{m,n,s} \alpha_{m,n,s}^* = I_{P_m \otimes P_n}.$ Thus, for any $0 \le h \le \min\{m,n\}$ $\min\{m,n\}$

$$\alpha_{m,n,h}^* = \alpha_{m,n,h}^* I_{P_m \otimes P_n} = \alpha_{m,n,h}^* \sum_{s=0}^{\min\{m,n\}} \alpha_{m,n,s} \alpha_{m,n,s}^* = \sum_{s=0}^{\min\{m,n\}} \alpha_{m,n,h}^* \alpha_{m,n,s} \alpha_{m,n,s}^*$$

By Equation (2.6) above, we get

$$\alpha_{m,n,h}^* = \alpha_{m,n,h}^* \alpha_{m,n,h} \alpha_{m,n,h}^* = \lambda \alpha_{m,n,h}^*$$

13

Since $\alpha_{m,n,h} \neq 0$ $(\alpha_{m,n,h}(x_1^{m+n-2h}) \neq 0)$, we have $\alpha_{m,n,h}^* \neq 0$ and $\lambda = 1$.

The following straightforward corollary describes the SU(2)-equivariant projections of $P_m \otimes P_n$ onto the SU(2)- subspaces $W_{m+n-2h} = \alpha_{m,n,h}(P_{m+n-2h})$. Note that by the equivariance of $\alpha_{m,n,h}$ and by Schur Lemma [13, p.13], the space W_{m+n-2h} is an irreducible.

Corollary 2.11. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, and $\alpha_{m,n,h}$ be as in Definition 2.8. Then

(1) α_{m,n,h}α^{*}_{m,n,h} is the orthogonal projection of P_m⊗P_n onto the SU(2)-irreducible subspace W_{m+n-2h} = α_{m,n,h}(P_{m+n-2h}) ≅ P_{m+n-2h}.
 (2) P_m⊗P_n = ⊕ W_{m+n-2h} W_{m+n-2h}, where the subspaces W_{m+n-2h} are mutually orthog-

A similar result can be proved for the SU(2)-space $P_m \otimes \overline{P}_n$. We start by constructing isometric embedding in the following lemma. Recall that \overline{P}_n is an SU(2)irreducible space under the contragredient representation ρ_n .

Lemma 2.12. For $m, n \in \mathbb{N}$,

onal.

- (1) The map $I_{P_m} \otimes J_n : P_m \otimes P_n \longrightarrow P_m \otimes \overline{P}_n$ is an SU(2)-equivariant unitary isomorphism whose inverse is $I_{P_m} \otimes J_n^*$.
- (2) The map $\eta_{m,n,h} = (I_{P_m} \otimes J_n) \alpha_{m,n,h}$ is an SU(2)-equivariant isometry from P_{m+n-2h} into $P_m \otimes \overline{P}_n$.

Using the embedding above, we obtain the decomposition of $P_m \otimes \overline{P}_n$ as follows:

Corollary 2.13. Let $m, n, h \in \mathbb{N}$ with $0 \leq h \leq \min\{m, n\}$. Let $\eta_{m,n,h}$ be defined as in Lemma 2.12. Then $\eta_{m,n,h}\eta_{m,n,h}^*$ is the orthogonal projection of $P_m \otimes \overline{P}_n$ onto the SU(2)-subspace $V_{m+n-2h} = \eta_{m,n,h}(P_{m+n-2h}) \cong P_{m+n-2h}$. Moreover, $P_m \otimes \overline{P}_n = \bigoplus_{h=0}^{\min\{m,n\}} V_{m+n-2h}$, where V_{m+n-2h} , $0 \leq h \leq \min\{m,n\}$ are mutually orthogonal SU(2)irreducible subspaces.

2.2.2. A computational formula for $\alpha_{m,n,h}$.

In this subsection, we derive a computational formula for the isometry $\alpha_{m,n,h}$, and prove some related corollaries. For the rest of this paper, we will systematically use the following notations without further mention.

For $m, n, h, i, j \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}, 0 \le i \le m + n - 2h$, and $0 \le j \le n$. Let

- $k_1(i) := \max\{0, -m+i+h\}.$
- $k_2(i) := \min\{i, n-h\}.$
- $B(i) := \{j : k_1(i) \le j \le k_2(i) + h\}.$
- $l_{ij} := i j + h$.
- $c_{m,n,h}$ is as defined in Theorem 2.7.

Lemma 2.14. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let r = m + n - 2h, and $\{f_i^r : 0 \le i \le r\}$ be the canonical basis for P_r . Then for $0 \le i \le r$,

$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sum_{s=0}^{h} \sum_{j=k_{1}(i)+s}^{k_{2}(i)+s} \beta_{i,s,j}^{m,n,h} f_{l_{ij}}^{m} \otimes f_{j}^{n}$$

where $\beta_{i,s,j}^{m,n,h} = (-1)^{s} \sqrt{\frac{c_{m,n,h} r! m! n!}{\binom{r}{(i)} \binom{m}{(i-j+h)}\binom{n}{j}} \frac{\binom{h}{s} \binom{n-h}{j-s} \binom{m-h}{(i-j+s)}}{(m-h)!}}{(m-h)!}.$

Proof.

Recall that the canonical basis element of P_r is $f_i^r = a_r^i x_1^i x_2^{r-i}$ where $a_r^i = \frac{1}{\sqrt{i!(r-i)!}}$. As

$$\Delta_{yx}^{n-h} = \left(y_1 \frac{\partial}{\partial x_1} + y_2 \frac{\partial}{\partial x_2}\right)^{n-h} = \sum_{t=0}^{n-h} \binom{n-h}{t} y_1^t y_2^{n-h-t} \frac{\partial^t}{\partial x_1^t} \frac{\partial^{n-h-t}}{\partial x_2^{n-h-t}},$$

and

$$\Gamma^{h}_{xy} = (x_1y_2 - y_1x_2)^{h} = \sum_{s=0}^{h} (-1)^{s} {h \choose s} (x_1y_2)^{h-s} (x_2y_1)^{s}.$$

$SU(2)\mbox{-}\mbox{iRREDUCIBLY COVARIANT AND EPOSIC CHANNELS}$

We have by the definition of $\alpha_{m,n,h}$ (see Definition 2.8), that for $0 \leq i \leq r$

$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sqrt{c_{m,n,h}}a_{r}^{i}\sum_{s=0}^{h}\sum_{t=0}^{n-h}(-1)^{s}\binom{h}{s}\binom{n-h}{t}x_{1}^{h-s}x_{2}^{s}y_{1}^{(t+s)}y_{2}^{n-(s+t)}\frac{\partial^{t}}{\partial x_{1}^{t}}x_{1}^{i}\frac{\partial^{n-h-t}}{\partial x_{2}^{n-h-t}}x_{2}^{r-i}$$

But

$$\frac{\partial^{t}}{\partial x_{1}^{t}}x_{1}^{i}\frac{\partial^{n-h-t}}{\partial x_{2}^{n-h-t}}x_{2}^{r-i} = \begin{cases} \frac{i!(r-i)!}{(i-t)!(m-h-i+t)!}x_{1}^{i-t}x_{2}^{m+t-h-i} & if & -m+i+h \le t \le i \\ 0 & 0 & otherwise \end{cases}$$

Thus, we can rewrite the above sum as

$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sqrt{c_{m,n,h}}a_{r}^{i}\sum_{s=0}^{h}\sum_{t=\max\{0,-m+i+h\}}^{\min\{i,n-h\}} (-1)^{s}\binom{h}{s}\binom{n-h}{t}\frac{i!(r-i)!}{(i-t)!(m-h-i+t)!}x_{1}^{h-(s+t)+i}x_{2}^{m+(s+t)-h-i}y_{1}^{(t+s)}y_{2}^{n-(s+t)}$$

Changing the summation variable in the inner sum to j = s + t, we obtain

$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sqrt{c_{m,n,h}} a_{r}^{i} \sum_{s=0}^{h} \sum_{j=k_{1(i)}+s}^{k_{2}(i)+s} (-1)^{s} \binom{h}{s} \binom{n-h}{j-s} \frac{i!(r-i)!}{(i-j+s)!(m-h-i+j-s)!} x_{1}^{l_{ij}} x_{2}^{m-l_{ij}} y_{1}^{j} y_{2}^{n-j}$$

Finally, as
$$\frac{a_r^i}{a_m^{l_{ij}}a_n^j} = \sqrt{\frac{j!l_{ij}!(m-l_{ij})!(n-j)!}{i!(r-i)!}}$$
, we have

$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sqrt{c_{m,n,h}} \sum_{s=0}^{h} \sum_{j=k_{1}(i)+s}^{k_{2}(i)+s} (-1)^{s} \frac{\binom{h}{s}\binom{n-h}{j-s}\binom{m-h}{i-j+s}}{(m-h)!} \sqrt{\frac{r!\ m!\ n!}{\binom{n}{i}\binom{n}{j}\binom{n}{j}}} f_{lij}^{m} \otimes f_{j}^{n}$$
$$= \sum_{s=0}^{h} \sum_{j=k_{1}(i)+s}^{k_{2}(i)+s} \beta_{i,s,j}^{m,n,h} f_{lij}^{m} \otimes f_{j}^{n}$$

The following corollary gives a more compact form of the above formula for $\alpha_{m,n,h}(f_i^r)$.

Corollary 2.15. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let r = m + n - 2h. Then

$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sum_{j \in B(i)} \varepsilon_{i}^{j}{}_{(m,n,h)} f_{l_{ij}}^{m} \otimes f_{j}^{n}$$

where $\varepsilon_i^j = \sum_{s=\max\{0,j-i,j+h-n\}}^{\min\{h,j,j+m-i-h\}} \beta_{i,s,j}^{m,n,h}$.

Proof.

Let
$$A = \{(s, j) : 0 \le s \le h, k_1(i) + s \le j \le k_2(i) + s\}$$
, and $A_j = \{s : (s, j) \in A\}$.
Then $A_j = \{s : 0 \le s \le h, j - k_2(i) \le s \le j - k_1(i)\}$
 $= \{s : \max\{0, j - k_2(i)\} \le s \le \min\{h, j - k_1(i)\}\}$
 $= \{s : \max\{0, j - i, j + h - n\} \le s \le \min\{h, j, j + m - i - h\}.$

Observe that $A = \bigcup_{j} \{(s, j) : s \in A_j\}$, where the index j range from $k_1(i)$ at s = 0 to $k_2(i) + h$ at s = h. Thus, by Lemma 2.14, we have

$$\alpha_{m,n,h}(f_i^r) = \sum_{(s,j)\in A} \beta_{i,s,j}^{m,n,h} f_{l_{ij}}^m \otimes f_j^n = \sum_{j=k_1(i)}^{k_2(i)+h} \sum_{s\in A_j} \beta_{i,s,j}^{m,n,h} f_{l_{ij}}^m \otimes f_j^n$$
$$= \sum_{j=k_1(i)}^{k_2(i)+h} \sum_{s=\max\{0,j-i,j+h-n\}}^{\min\{h,j,j+m-i-h\}} \beta_{i,s,j}^{m,n,h} f_{l_{ij}}^m \otimes f_j^n = \sum_{j\in B(i)} \varepsilon_i^j(m,n,h) f_{l_{ij}}^m \otimes f_j^n$$

We now state and prove identities for the coefficients $\varepsilon_{i(m,n,h)}^{j}$. When m, n, h are clear from the context, we write $\varepsilon_{i(m,n,h)}^{j}$ as ε_{i}^{j} .

Lemma 2.16. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let r = m + n - 2h. Then, the coefficients $\varepsilon_i^j := \varepsilon_i^{j}{}_{(m,n,h)}$ satisfy

$$\begin{array}{ll} (1) \ \varepsilon_{i}^{j} = (-1)^{h} \varepsilon_{r-i}^{n-j}, & for \ 0 \leq i \leq r \ and \ j \in B(i). \\ (2) \ \varepsilon_{i}^{i+h} = \beta_{i,h,h+i}^{m,n,h}, & for \ any \ i \leq n-h. \\ (3) \ For \ n-h \leq i \leq r, \ we \ have \ \varepsilon_{i}^{n} = \beta_{i,h,n}^{m,n,h} = (-1)^{h} \left| \beta_{i,h,n}^{m,n,h} \right| \neq 0. \\ (4) \ For \ j \in B(0), \ we \ have \ \varepsilon_{0}^{j} = \beta_{0,j,j}^{m,n,h} = (-1)^{j} \left| \beta_{0,j,j}^{m,n,h} \right| \neq 0. \end{array}$$

Proof.

(1) Let
$$g_0 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$
. By the $SU(2)$ -equivariance of $\alpha_{m,n,h}$, we have
 $(\rho_m(g_0) \otimes \rho_n(g_0)) \alpha_{m,n,h} \rho_r^*(g_0)(f_i^r) = \alpha_{m,n,h}(f_i^r)$

for $0 \leq i \leq r$. As $\rho_r^*(g_0)(f_i^r) = (-1)^{r-i}f_{r-i}^r$, $\rho_m(g_0)(f_l^m) = (-1)^l f_{m-l}^m$, and $\rho_n(g_0)(f_j^n) = (-1)^j f_{n-j}^n$, by Corollary 2.15, the above equation can be written as

$$(-1)^{h} \sum_{s=k_{1}(r-i)}^{k_{2}(r-i)+h} \varepsilon_{r-i}^{s} f_{l_{i}(n-s)}^{m} \otimes f_{n-s}^{n} = \sum_{j=k_{1}(i)}^{k_{2}(i)+h} \varepsilon_{i}^{j} f_{l_{ij}}^{m} \otimes f_{j}^{n}$$

Let t = n - s. Since $h + k_2(r - i) = n - k_1(i)$ and $h + k_1(r - i) = n - k_2(i)$, we have

$$(-1)^{h} \sum_{t=k_{1}(i)}^{k_{2}(i)+h} \varepsilon_{r-i}^{n-t} f_{lit}^{m} \otimes f_{t}^{n} = \sum_{j=k_{1}(i)}^{k_{2}(i)+h} \varepsilon_{i}^{j} f_{lij}^{m} \otimes f_{j}^{n}$$

Since the vectors $f_l^m \otimes f_j^n$ are linearly independent, the last equality implies that $(-1)^h \varepsilon_{r-i}^{n-j} = \varepsilon_i^j$ for each $k_1(i) \leq j \leq k_2(i) + h$.

The claims (2),(3), and (4) follow by direct computations.

Using the fact that $\alpha_{m,n,h}$ is an isometry, one can prove that the coefficients $c_{m,n,h}$ in Theorem 2.7 are given by

$$c_{m,n,h} = \frac{((m-h)!)^2}{(m+n-2h)! \ m! \ n! \left(\sum_{k=0}^h \frac{\binom{h}{k}^2}{\binom{m}{h-k}\binom{n}{k}}\right)}$$

Consequently, we have the following corollary

Corollary 2.17. Let $m, n, h \in \mathbb{N}$ be such that $0 \le h \le \min\{m, n\}$. Then

(1) The coefficients $c_{m,n,h}$ in Theorem 2.7 satisfy $c_{m,n,h} = \frac{((m-h)!)^2}{((n-h)!)^2} c_{n,m,h}$.

(2) For
$$0 \le i \le m + n - 2h$$
 and $j \in B(i)$, we have
(a) $\beta_{i,h-s,l_{ij}}^{n,m,h} = (-1)^h \beta_{i,s,j}^{m,n,h}$, for any $0 \le s \le h$.
(b) $\varepsilon_i^j(m,n,h) = (-1)^h \varepsilon_i^{l_{ij}}(n,m,h)$.

2.3. EPOSIC channels.

The SU(2)-equivariant isometry $\alpha_{m,n,h}$ in Proposition 2.10, will induce an SU(2)covariant quantum channel $\Phi_{m,n,h} : End(P_{m+n-2h}) \to End(P_m)$ that has a Stinespring representation $(P_n, \alpha_{m,n,h})$, see Proposition 1.8. According to Definition 1.12, $\Phi_{m,n,h}$ is SU(2)-irreducibly covariant channel. The following proposition records this result.

Proposition 2.18. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let r = m + n - 2hand $\alpha_{m,n,h}$ be as in Definition 2.8. The map $\Phi_{m,n,h} : End(P_r) \to End(P_m)$ defined by

$$\Phi_{m,n,h}(A) = Tr_{P_n}(\alpha_{m,n,h}A\alpha_{m,n,h}^*) \qquad A \in End(P_r)$$

is an SU(2)-irreducibly covariant channel.

Definition 2.19. Let $r, m \in \mathbb{N}$, and $\mathcal{E}(r, m) = \{(n, h) \in \mathbb{N}^2 : r = m + n - 2h, 0 \le h \le \min\{m, n\}\}.$ For $(n, h) \in \mathcal{E}(r, m)$, We call the quantum channel $\Phi_{m,n,h} : End(P_r) \longrightarrow End(P_m)$, defined in Proposition 2.18, an EPOSIC channel.

Recall that by Proposition 1.13, the set $QC_{SU(2)}(\rho_r, \rho_m)$ of all SU(2)-irreducibly covariant channels from $End(P_r)$ into $End(P_m)$, is a convex set. As we will show in Section 4, the set of all EPOSIC channels form $End(P_r)$ into $End(P_m)$ forms the extreme points of $QC_{SU(2)}(\rho_r, \rho_m)$, justifying the nomenclature EPOSIC.

We denote by EC(r, m) the set of all EPOSIC channels form $End(P_r)$ into $End(P_m)$, and abbreviate EC(m, m) to EC(m).

Lemma 2.20. Let $r, m \in \mathbb{N}$, then

$$\mathcal{E}(r,m) = \{ (r+m-2l,m-l) \in \mathbb{N}^2 : 0 \le l \le \min\{r,m\} \}$$

Proof.

Let $\mathcal{B} = \{(r+m-2l, m-l) \in \mathbb{N}^2 : 0 \le l \le \min\{r, m\}\}$. If $(r+m-2l, m-l) \in \mathcal{B}$ for some $0 \le l \le \min\{r, m\}$, set $n_0 = r+m-2l$, and $h_0 = m-l$. Then (n_0, h_0) satisfies $r = n_0 + m - 2h_0$ and $0 \le h_0 \le \min\{m, n_0\}$. (note that $h_0 \le n_0$ since $h_0 \le h_0 + (r - l) = m - l + r - l = m + r - 2l = n_0$). Thus, $(r + m - 2l, m - l) = (n_0, h_0) \in \mathcal{E}(r, m)$. Conversely, if $(n, h) \in \mathcal{E}(r, m)$, set $l_0 = m - h$. Then, since r = n + m - 2h, we have $n = r - m + 2h = r + m - 2l_0$. As $0 \le h \le \min\{m, n\}$, we have $0 \le l_0 \le m$ and $l_0 \le l_0 + (n - h) = (m - h) + (n - h) = m + n - 2h = r$. Hence, $0 \le l_0 \le \min\{r, m\}$, and $(n, h) = (r + m - 2l_0, m - l_0) \in \mathcal{B}$.

As a consequence of the above lemma, we have:

Proposition 2.21. Let $r, m \in \mathbb{N}$, then

$$EC(r,m) = \{\Phi_{m,r+m-2l,m-l}, \ 0 \le l \le \min\{r,m\}\}\$$

Definition 2.22. [5, p.125] Let H, K be two finite dimensional spaces. Then a quantum channel $\Phi : End(H) \longrightarrow End(K)$ is said to be unital if it satisfies that $\Phi(I_H) = \frac{d_H}{d_K} I_K$.

Remark 2.23.

- (1) According to a result that is due to A. HOLEVO [6], any *G*-irreducibly covariant channel is unital. Thus, EPOSIC channels are unital.
- (2) Since the coefficient c_{m,0,0} in Theorem 2.7 is equal to 1, then using the natural isomorphism to identify the spaces P_m ⊗ P₀ and P_m in Definition 2.8, we have that α_{m,0,0} is the identity map on P_m. Thus, Φ_{m,0,0} is the identity map on End(P_m).

3. A KRAUS REPRESENTATION AND THE CHOI MATRIX OF EPOSIC CHANNELS

In this section, we use the same notation introduced in Section 2.2.2. Recall that for $k \in \mathbb{N}$ the set $\{f_s^k : 0 \le s \le k\}$ denotes the canonical basis for P_k .

3.1. A Kraus representation of EPOSIC channel.

Definition 3.1. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$. For $0 \le j \le n$, we define $T_j : P_{m+n-2h} \longrightarrow P_m$ by

$$T_j = (I_{P_m} \otimes f_j^{n^*}) \alpha_{m,n,h}$$

By Proposition 1.9, the set $\{T_j : 0 \le j \le n\}$ is a Kraus representation of $\Phi_{m,n,h}$. We call the Kraus operators defined above, EPOSIC Kraus operators. By direct computations using Corollary 2.15 and Definition 3.1, we have:

Proposition 3.2. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let $\{T_j : 0 \le j \le n\}$ be the EPOSIC Kraus operators of $\Phi_{m,n,h}$. Then for $0 \le j \le n$

$$T_j(f_i^r) = \begin{cases} \varepsilon_i^j f_{l_{ij}}^m & \text{if } j \in B(i) \\ 0 & \text{otherwise} \end{cases}$$

Corollary 3.3. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let r = m + n - 2h. For $0 \le i_1, i_2 \le r$, we have

$$\Phi_{m,n,h}(f_{i_1}^r f_{i_2}^{r^*}) = \sum_{j \in B(i_1) \cap B(i_2)} \varepsilon_{i_1}^j \varepsilon_{i_2}^j f_{l_{i_1j}}^m f_{l_{i_2j}}^{m^*}$$

Corollary 3.4. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, and r = m + n - 2h. Let D_k denote the set of all diagonal operators of $End(P_k)$. Then $\Phi_{m,n,h}(D_r) \subseteq D_m$.

The SU(2) equivariance property of $\alpha_{m,n,h}$ implies the symmetry relations for the T_j 's. This relation is given in the next proposition whose proof follows easily by the following lemma.

Lemma 3.5. Let $m, n, h \in \mathbb{N}$ with $0 \leq h \leq \min\{m, n\}$, r = m + n - 2h, and $\{T_j : 0 \leq j \leq n\}$ be the EPOSIC Kraus operators of $\Phi_{m,n,h}$. Then for any $0 \leq i \leq r$, $0 \leq l \leq m$, and $0 \leq j \leq n$, we have

$$\left\langle f_l^m \left| T_j(f_i^r) \right\rangle_{P_m} = (-1)^h \left\langle f_{m-l}^m \left| T_{n-j}(f_{r-i}^r) \right\rangle_{P_m} \right.$$

where $\{f_s^k : 0 \le s \le k\}$ is the canonical basis for $P_k, k \in \{r, m\}$.

Proof.

Fix $j \in \mathbb{N}$ such that $0 \leq j \leq n$, then we have one of the following cases:

• If $j \in B(i)$ then by Lemma 2.16, we have $\varepsilon_i^j = (-1)^h \varepsilon_{r-i}^{n-j}$, and since $l = l_{ij}$ if and only if $m - l = l_{(r-i)(n-j)}$, it follows that

$$\left\langle f_{l}^{m} \left| T_{j}(f_{i}^{r}) \right\rangle_{P_{m}} = \varepsilon_{i}^{j} \delta_{ll_{ij}} = (-1)^{h} \varepsilon_{r-i}^{n-j} \delta_{(m-l)l_{(r-i)(n-j)}} = \left\langle f_{m-l}^{m} \left| T_{n-j}(f_{r-i}^{r}) \right\rangle_{P_{m}} \right\rangle_{P_{m}}$$

• If $j \notin B(i)$, then $n - j \notin n - B(i) = B(r - i)$ and both of $T_j(f_i^r)$, $T_{n-j}(f_{r-i}^r)$ are zero. This implies that the identity also holds in this case.

Proposition 3.6. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, and $\{T_j : 0 \le j \le n\}$ be the EPOSIC Kraus operators of $\Phi_{m,n,h}$. Then for each $0 \le j \le n$, we have

$$\rho_m(g_0)T_j\rho_r^*(g_0) = (-1)^j T_{n-j}$$

where $g_0 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.

The relation in Proposition 3.6 can be translated into a symmetry relation for the vectors representing the operators T_j in $P_m \otimes \overline{P}_r$, see Example 1.5, to get

(3.1)
$$\operatorname{Vec}(T_{n-j}) = (-1)^{j} \left(\rho_{m}(g_{0}) \otimes \check{\rho}_{r}(g_{0}) \right) \operatorname{Vec}(T_{j})$$

It is worth noticing that even though Φ is a *G*-covariant channel, its Kraus operators are not necessary *G*-equivariant.

Notations: Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, and r = m = n - 2h. For any $0 \le j \le n$, let $l_e(j) = \max\{0, h - j\}$, $l_f(j) = \min\{r - j + h, m\}$, and $R(j) = \{l : l_e(j) \le l \le l_f(j)\}$.

Remark 3.7. Using the same notation above, we have:

- (1) For fixed j such $0 \le j \le n$, and for $i, 0 \le i \le r$, we have $j \in B(i) \iff l_{ij} \in R(j)$.
- (2) By Proposition 3.2 and (1) above, we have that the Kraus operator T_j can be written as $T_j = \sum_{l \in R(j)} \varepsilon_{l+j-h}^j f_l^m f_{l+j-h}^{r^*}$.

(3) Using (2) above, we have
$$\operatorname{Vec}(T_j) = \sum_{l \in R(j)} \varepsilon_{l+j-h}^j f_l^m \otimes f_{l+j-h}^r$$
.

The first part of the following proposition can by proved by taking the adjoint of both sides of the equation in Remark 3.7 (2), while the second part is direct computations.

Proposition 3.8. Let $m, n, h \in \mathbb{N}$ with $0 \leq h \leq \min\{m, n\}$ and r = m + n - 2h. Let $\{T_j : 0 \leq j \leq n\}$ be the EPOSIC Kraus operators for $\Phi_{m,n,h}$ and T_j^* denotes the adjoint map of T_j . Then for $0 \leq j \leq n$ we have

(1)
$$T_{j}^{*} = \sum_{l \in R(j)} \varepsilon_{l+j-h}^{j} f_{l+j-h}^{r} f_{l}^{m^{*}} i.e$$

$$T_{j}^{*}\left(f_{l}^{m}\right) = \begin{cases} \varepsilon_{l+j-h}^{j} f_{l+j-h}^{r} & if \quad l \in R(j) \\ 0 & else \end{cases}$$

(2)
$$flip_{P_r}^{\overline{P}_m} \left(J_m \otimes J_r^* \right) \left(Vec(T_{n-j}) \right) = (-1)^{m+j} Vec(T_j^*)$$

where $\{f_s^k : 0 \le s \le k\}$, is the canonical basis for P_k , and J_k as in Definition 2.2, $k \in \{r, m\}$.

3.2. The Choi matrix of $\Phi_{m,n,h}$.

Recall that by Corollary 2.13, the space $P_m \otimes \overline{P}_r$ decomposes into an orthogonal direct sum of SU(2)-irreducible subspaces V_{m+r-2l} , $0 \leq l \leq \min\{m,r\}$. The maps $\eta_{m,r,l}: V_{m+r-2l} \longrightarrow P_m \otimes \overline{P}_r$, $0 \leq l \leq \min\{m,r\}$ are isometries such that the final supports $q_{m,r,l} = \eta_{m,r,l}\eta_{m,r,l}^*$ are the mutually orthogonal projections onto V_{m+r-2l} , $0 \leq l \leq \min\{m,r\}$. These projections satisfy $\sum_{l=0}^{\min\{m,r\}} q_{m,r,l} = I_{P_m \otimes \overline{P}_r}$. By Schur Lemma [13, p.13], we get: **Proposition 3.9.** [17] Let ρ_m , ρ_r be the irreducible representation of SU(2) on P_m , P_r respectively. Then $(\rho_m \otimes \check{\rho}_r (SU(2)))'$ is an abelian algebra that is generated by the projections on the SU(2)-subspaces V_{m+r-2l} of $P_m \otimes \overline{P}_r$ where $0 \leq l \leq \min\{m, r\}$.

By the last Proposition and Proposition 1.15, we have:

Corollary 3.10. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, and r = m + n - 2h, the Choi matrix of $\Phi_{m,n,h}$ is generated by $\{q_{m,r,l} = \eta_{m,r,l}\eta_{m,r,l}^* : 0 \le l \le \min\{m,r\}\}$.

Remark 3.11. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, if r = m + n - 2h then since $0 \le m - h \le \min\{m, r\}$, the set $\{q_{m,r,l} : 0 \le l \le \min\{m, r\}\}$ always contains the projection $q_{m,r,m-h} = \eta_{m,r,m-h} \eta^*_{m,r,m-h}$.

The following lemma will be used to prove Proposition 3.13.

Lemma 3.12. Let $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, and r = m + n - 2h. Then the operator $C(\Phi_{m,n,h})\eta_{m,r,m-h}$ is non zero.

Proof.

Let $\{f_s^k : 0 \le s \le k\}$ denote the canonical basis of P_k where $k \in \{r, n, m\}$. To show that $C(\Phi_{m,n,h})\eta_{m,r,m-h}$ is non zero, it is enough to show that $C(\Phi_{m,n,h})(\eta_{m,r,m-h}(f_0^n)) \ne 0$. By Corollary 3.3, we have

$$C(\Phi_{m,n,h}) = \sum_{i_1,i_2=0}^r \Phi_{m,n,h}(f_{i_1}^r f_{i_2}^{r^*}) \otimes f_{i_1}^r f_{i_2}^{r^*} = \sum_{i_1,i_2=0}^r \sum_{j \in B(i_1) \cap B(i_2)} \varepsilon_{i_1}^j \varepsilon_{i_2}^j f_{l_{i_1j}}^m f_{l_{i_2j}}^{m^*} \otimes f_{i_1}^r f_{i_2}^{r^*}.$$

By Lemma 2.12, and Corollary 2.16, we have

$$\eta_{m,r,m-h}(f_0^n) = (I_{P_m} \otimes J_r) \alpha_{m,r,m-h}(f_0^n) = (I_{P_m} \otimes J_r) \left(\sum_{t \in B(0)} \varepsilon_{0}^t{}_{(m,r,m-h)} f_{l_{0t}}^m \otimes f_t^r \right)$$
$$= \sum_{t \in B(0)} (-1)^t \varepsilon_{0}^t{}_{(m,r,m-h)} f_{l_{0t}}^m \otimes f_{r-t}^r$$
$$= \sum_{t \in B(0)} \lambda_t f_{m-(t+h)}^m \otimes f_{r-t}^r \quad \text{where } \lambda_t > 0 .$$

Hence, we get:

$$C(\Phi_{m,n,h})(\eta_{m,r,m-h}(f_0^n)) = \sum_{i_1,i_2=0}^r \sum_{j \in B(i_1) \cap B(i_2)} \varepsilon_{i_1}^j \varepsilon_{i_2}^j f_{l_{i_1j}}^m f_{l_{i_2j}}^{m^*} \otimes f_{i_1}^r f_{i_2}^{r^*} \left(\sum_{t \in B(0)} \lambda_t f_{m-(t+h)}^m \otimes f_{r-t}^r\right)$$
$$= \sum_{i_1,i_2=0}^r \sum_{j \in B(i_1) \cap B(i_2)} \sum_{t \in B(0)} \lambda_t \varepsilon_{i_1}^j \varepsilon_{i_2}^j \left(f_{l_{i_1j}}^m f_{l_{i_2j}}^{m^*}\right) f_{m-(t+h)}^m \otimes \left(f_{i_1}^r f_{i_2}^{r^*}\right) f_{r-t}^r$$

But, for $0 \leq i_1, i_2 \leq r, j \in B(i_1) \cap B(i_2)$ and $t \in B(0)$, we have:

$$f_{l_{i_{1}j}}^{m}f_{l_{i_{2}j}}^{m^{*}}f_{m-(t+h)}^{m} \otimes f_{i_{1}}^{r}f_{i_{2}}^{r^{*}}f_{r-t}^{r} = \begin{cases} f_{l_{i_{1}n}}^{m} \otimes f_{i_{1}}^{r} & if \quad i_{2} = r-t, j = n \\ \\ 0 & otherwise \end{cases}$$

As for $t \in B(0)$, $n \in B(i_1) \cap B(r-t)$ if and only if $n-h \le i_1 \le r$, we have

$$C(\Phi_{m,n,h})\left(\eta_{m,r,m-h}(f_0^n)\right) = \sum_{i_1=n-h}^r \sum_{t\in B(0)} \lambda_t \varepsilon_{i_1}^n \varepsilon_{r-t}^n f_{i_1-n+h}^m \otimes f_{i_1}^r$$

Assume that

$$0 = C(\Phi_{m,n,h})(\eta_{m,r,m-h}(f_0^n)) = \sum_{i_1=n-h}^r \sum_{t \in B(0)} \lambda_t \varepsilon_{i_1}^n \varepsilon_{r-t}^n f_{i_1-n+h}^m \otimes f_{i_1}^r$$

then by the linearly independence of $\{f_i^m \otimes f_i^r : 0 \le l \le m, 0 \le i \le r\}$, we would have that $\sum_{t \in B(0)} \lambda_t \varepsilon_{i_1}^n \varepsilon_{r-t}^n = 0$ for $n - h \le i_1 \le r$.

As by Corollary 2.16, $\varepsilon_{i_1}^n \neq 0$ for $n-h \leq i_1 \leq r$, we would have $\sum_{t \in B(0)} \lambda_t \varepsilon_{r-t}^n = 0$. But by Corollary 2.16, we have $\varepsilon_{r-t}^n = (-1)^h \theta_t$ with $\theta_t > 0$ holds for $t \in B(0) = \{t : 0 \leq t \leq m-h\}$. Therefore, $\sum_{t \in B(0)} \lambda_t \varepsilon_{r-t}^n = (-1)^h \sum_{t=0}^{m-h} \lambda_t \theta_t \neq 0$ which is a contradiction. \Box

Proposition 3.13. Let $m, n, h \in \mathbb{N}$ with $0 \leq h \leq \min\{m, n\}$, and r = m + n - 2h. Then the Choi matrix of the EPOSIC channel $\Phi_{m,n,h}$ is equal to $\frac{r+1}{n+1}q_{m,r,m-h}$ where $q_{m,r,m-h}$ is the projection of $P_m \otimes \overline{P}_r$ onto the SU(2)-invariant subspace of dimension n + 1. Proof.

By Corollary 3.10, we have $C(\Phi_{m,n,h}) = \sum_{l=0}^{\min\{m,r\}} \lambda_l q_{m,r,l}$ where $q_{m,r,l} = \eta_{m,r,l} \eta_{m,r,l}^*, 0 \le l \le \min\{m,r\}$ are the mutually orthogonal projections onto V_{m+r-2l} . Consequently, $rank(C(\Phi_{m,n,h})) = \sum_{l=0,\lambda_l\neq 0}^{\min\{m,r\}} rank(q_{m,r,l}).$

By Lemma 3.12, we have $\lambda_{m-h} \neq 0$, hence $rank(C(\Phi_{m,n,h})) \geq rank(q_{m,r,m-h}) = dim(V_n) = n + 1$. As by Remark 1.10, $rank(C(\Phi_{m,n,h})) \leq n + 1$, we obtain that $C(\Phi_{m,n,h}) = \lambda_{m-h}q_{m,r,m-h}$. Taking the trace of both side of the equation, we get

$$r + 1 = tr(C(\Phi_{m,n,h})) = \lambda_{m-h}tr(q_{m,r,m-h}) = \lambda_{m-h}n + 1$$

and $\lambda_{m-h} = \frac{r+1}{n+1}$.

Remark 3.14. Note that the above proposition establishes a one to one corresponding between EC(r,m) and the projections on the SU(2)-subspaces of $P_m \otimes \overline{P}_r$ $(\Phi_{m,m+r-2l,m-l} \longleftrightarrow \frac{r+1}{m+r-2l+1}q_{m,r,l}).$

Corollary 3.15. For $r, m \in \mathbb{N}$, there are exactly $\min\{r, m\}+1$ elements in EC(r, m).

Recall that for $r, m \in \mathbb{N}$, the set $End(End(P_r), End(P_m))^{SU(2)}$ denotes the vector space of SU(2)-equivariant maps Φ from $End(P_r)$ to $End(P_m)$. The following corollaries are direct consequence of Proposition 3.13.

Corollary 3.16. Let $r, m \in \mathbb{N}$, let ρ_r and ρ_m be the irreducible representations of SU(2) on P_r and P_m . Then EC(r, m) is a spanning set of $End(End(P_r), End(P_m))^{SU(2)}$.

Proof.

By Proposition 3.9, and Proposition 3.13, the commutant

$$\left(\rho_m \otimes \check{\rho}_r \left(SU(2)\right)\right)' = \left\{\sum_{l=0}^{\min\{r,m\}} \lambda_l q_{m,r,l} : \lambda_l \in \mathbb{C}\right\}$$

$$= \left\{ \sum_{l=0}^{\min\{r,m\}} \mu_l C(\Phi_{m,m+r-2l,m-l}) : \mu_l \in \mathbb{C} \right\}$$
$$= Span \left\{ C \left(EC(r,m) \right) \right\}$$

The result now follows from Proposition 1.15.

Corollary 3.17. Let $r, m \in \mathbb{N}$. Then $QC_{SU(2)}(\rho_r, \rho_m)$ is the convex hull of EC(r, m).

Proof.

Let $\Phi \in QC_{SU(2)}(\rho_r, \rho_m)$. Since Φ is SU(2)-equivariant map then by Corollary 3.16, we have

(3.2)
$$\Phi = \sum_{l=0}^{\min\{r,m\}} \lambda_l \Phi_{m,m+r-2l,m-l} \qquad \lambda_l \in \mathbb{C}$$

It remains to show that $0 \leq \lambda_l$ and $\sum_{l=0}^{\min\{r,m\}} \lambda_l = 1$. By Remark 3.14, we have $C(\Phi) = \sum_{l=0}^{\min\{r,m\}} \frac{r+1}{m+r-2l+1} \lambda_l q_{m,r,l}$ where $q_{m,r,l}$, $0 \leq l \leq \min\{r,m\}$ are mutually orthogonal projections of $P_m \otimes \overline{P}_r$. By the orthogonality of $q_{m,r,l}$,s and the positivity of $C(\Phi)$ (see Proposition 1.8), we have $\lambda_l \geq 0$ for $0 \leq l \leq \min\{m,r\}$. Since both Φ and $\Phi_{m,m+r-2l,m-l}$ are trace preserving, choosing any state $\varrho \in D(P_r)$, we have $1 = tr(\varrho) = tr(\Phi(\varrho)) = \sum_{l=0}^{\min\{r,m\}} \lambda_l tr(\Phi_{m,m+r-2l,m-l}(\varrho)) = \sum_{l=0}^{\min\{r,m\}} \lambda_l$.

4. The extreme points of SU(2)-irreducibly covariant channels

In this section, we show that EC(r, m), the set of all EPOSIC channels from $End(P_r)$ to $End(P_m)$, forms the set of the extreme points of $QC_{SU(2)}(\rho_r, \rho_m)$. We also show that any completely positive SU(2)-equivariant map $\Phi : End(P_r) \longrightarrow End(P_m)$ is a multiple of SU(2)-covariant channel. Recall that $EC(r, m) = \{\Phi_{m,r+m-2l,m-l}, 0 \le l \le \min\{r, m\}\}$.

Proposition 4.1. For $r, m \in \mathbb{N}$, the set of extreme points in $QC(\rho_r, \rho_m)$ is EC(r, m).

Proof.

Since by Corollary 3.17, we have $QC(\rho_r, \rho_m) = Conv(EC(r, m))$, it is enough to show that any element in $QC_{SU(2)}(\rho_r, \rho_m)$ is uniquely written as linear combination of elements of EC(r, m). Let $\Psi \in QC_{SU(2)}(\rho_r, \rho_m)$ such that $\sum_{l=0}^{\min\{r,m\}} \lambda_l \Phi_l = \Psi = \sum_{l=0}^{\min\{r,m\}} \mu_l \Phi_l$ where $\Phi_l = \Phi_{m,m+r-2l,m-l}$, then by Proposition 3.13, and the orthogonality of $q_{m,r,l}$, $0 \leq l \leq \min\{m, r\}$, we have

$$\frac{r+1}{m+r-2l+1}\lambda_l q_{m,r,l} = q_{m,r,l}C(\Psi) = \frac{r+1}{m+r-2l+1}\mu_l q_{m,r,l}$$

Thus, $\lambda_l = \mu_l$, and EC(r, m) are extreme points of $QC_{SU(2)}(\rho_r, \rho_m)$. To complete the proof, note that since any extreme point of $QC_{SU(2)}(\rho_r, \rho_m)$ can not be written as a linear combination of elements of $QC_{SU(2)}(\rho_r, \rho_m)$ other than itself, then any extreme point of $QC_{SU(2)}(\rho_r, \rho_m)$ must be in EC(r, m).

As EC(r, m) is a spanning set for both the SU(2)-irreducibly equivariant maps and the SU(2)-irreducibly covariant channels, we have the following corollary:

Corollary 4.2. For $r, m \in \mathbb{N}$, any completely positive SU(2)-equivariant map Φ : $End(P_r) \longrightarrow End(P_m)$ is a multiple of an SU(2)-irreducibly covariant channel.

Proof.

By Corollary 3.16, we have $\Phi = \sum_{l=0}^{\min\{r,m\}} \lambda_l \Phi_{m,r+m-2l,m-l}$ for some $\lambda_l \in \mathbb{C}$. Since Φ is completely positive the coefficients λ_l are non-negative (otherwise, $C(\Phi) = \sum_{l=0}^{\min\{r,m\}} \frac{r+1}{m+r-2l+1} \lambda_l q_{m,r,l}$ will have a negative eigenvalue). Let $\lambda = \sum_{l=0}^{\min\{r,m\}} \lambda_l$. If $\lambda = 0$ then $\lambda_l = 0$ for all $0 \leq l \leq \min\{r,m\}$, and $\Phi = 0$ is a multiple of any SU(2)-irreducibly covariant channel. If $\lambda \neq 0$, then $\Psi = \sum_{l=0}^{\min\{m,r\}} \frac{\lambda_l}{\lambda} \Phi_{m,m+r-2l,m-l}$ is a convex combination of EPOSIC channels. Thus, by corollary 3.17, Ψ is an SU(2)-irreducibly covariant channel, and $\Phi = \lambda \Psi$.

5. Some maps related to eposic channel

In this section, given an EPOSIC channel $\Phi_{m,n,h}$, we construct a complementary channel $\tilde{\Phi}_{m,n,h}$. We also give the condition for the dual map $\Phi_{m,n,h}^*$ to be a quantum channel.

5.1. A complementary channel of $\Phi_{m,n,h}$.

Let us first recall the notion of complementary channels [8]. Given three Hilbert spaces H, K, E and a linear isometry $\alpha : H \longrightarrow K \otimes E$ one associates two quantum channels

$$\Phi:End(H)\longrightarrow End(K)\qquad \text{ and }\quad \Psi:End(H)\longrightarrow End(E)$$
 defined for $A\in End(H)$ by

$$\Phi(A) = Tr_E(\alpha A \alpha^*)$$
 and $\Psi(A) = Tr_K(\alpha A \alpha^*)$

The maps Φ and Ψ are called mutually complementary. For any quantum channel, a complementary channel always exists, see Proposition 1.8. However, due to the fact that Stinespring representation (dilation) is not unique, there can be many candidates for "the" complementary channel. In [8], Holevo clarifies in what sense the complementary map is unique. He showed that if (E, α) and (E', α') are two Stinespring representations (dilations) of $\Phi : End(H) \longrightarrow End(K)$ then there exist a partial isometry $J : E \longrightarrow E'$ such that $\alpha' = (I_K \otimes J)\alpha$, and $\alpha = (I_K \otimes J^*)\alpha'$. It follows that if $\Phi_E, \Phi_{E'}$ are complementary channels of Φ , then they are equivalent in the sense that there exist a partial isometry $J : E \longrightarrow E'$ such that $\Phi_E(\varrho) =$ $J^*\Phi_{E'}(\varrho)J$, and $\Phi_{E'}(\varrho) = J\Phi_E(\varrho)J'$ for any $\varrho \in D(H)$. Stinespring representations with minimal dimensionality of the space E are called minimal dilation, and any two minimal dilations are isometric. By Remark 1.10, the Stinespring representation with an environment space that satisfies $dim(E) = rank(C(\Phi))$ is a minimal dilation. Remark 5.1. [8, p.96] If G is a group, and π_H, π_K are representations of G on the Hilbert spaces H, K. Then $\Phi : End(H) \longrightarrow End(K)$ is G-covariant channel if and only if any complementary channel of Φ is G-covariant.

The following proposition will be used below to construct a complementary channel $\tilde{\Phi}_{m,n,h}$ of $\Phi_{m,n,h}$. Recall the notations in Section 2.2.2.

Proposition 5.2. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$

$$flip_{P_n}^{P_m}\alpha_{m,n,h} = (-1)^h \alpha_{n,m,h}$$

where $\alpha_{m,n,h}$ is the isometry in Definition 2.8.

Proof.

Let $\{f_s^k : 0 \leq s \leq k\}$ be the canonical bases for P_k , $k \in \{r, m, n\}$. Let $B^{m,n,h}(i) = \{j : \max\{0, -m+i+h\} \leq j \leq \min\{i+h, n\}\}$ (the set B(i) associated to $\alpha_{m,n,h}$), and $B^{n,m,h}(i) = \{j : \max\{0, -n+i+h\} \leq j \leq \min\{i+h, m\}\}$ (the set B(i) associated to $\alpha_{n,m,h}$), see Corollary 2.15). By Corollary 2.17, and since $j \in B^{m,n,h}(i)$ if and only if $l_{ij} \in B^{n,m,h}(i)$, we have for $0 \leq i \leq r$,

$$flip_{P_{n}}^{P_{m}}\alpha_{m,n,h}(f_{i}^{r}) = \sum_{j \in B^{m,n,h}(i)} \varepsilon_{i}^{j}(m,n,h) f_{j}^{n} \otimes f_{l_{ij}}^{m} = (-1)^{h} \sum_{l_{ij} \in B^{n,m,h}(i)} \varepsilon_{i}^{l_{ij}}(n,m,h) f_{j}^{n} \otimes f_{l_{ij}}^{m}$$

By taking the sum over $l = l_{ij}$, we get

$$flip_{P_n}^{P_m}\alpha_{m,n,h}(f_i^r) = (-1)^h \sum_{l \in B(i)} \varepsilon_{i^{(n,m,h)}}^l f_{i-l+h}^n \otimes f_l^m = (-1)^h \sum_{l \in B(i)} \varepsilon_{i^{(n,m,h)}}^l f_{l_{il}}^n \otimes f_l^m$$

Using the proposition above and the equation $Tr_{\kappa}(flip_{\kappa}^{H}Aflip_{H}^{\kappa}) = Tr_{\kappa}(A)$ for any $A \in End(H \otimes K)$, the following corollary becomes straightforward

Corollary 5.3. The channel $\Phi_{n,m,h}$ is a complementary channel for $\Phi_{m,n,h}$.

The following corollary to Proposition 5.2 will be used in the proof of Corollary 5.6. Recall that by Proposition 2.3, $J_m : P_m \longrightarrow \overline{P}_m$ is a unitary map.

Corollary 5.4. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$,

$$flip_{P_n}^{\overline{P}_m}(J_m\otimes J_n^*)\eta_{m,n,h} = (-1)^h \eta_{n,m,h}$$

Proof.

By Lemma 2.12, we have

$$flip_{P_n}^{\overline{P}_m}(J_m \otimes J_n^*)\eta_{m,n,h} = flip_{P_n}^{\overline{P}_m}(J_m \otimes J_n^*)(I_{P_m} \otimes J_n)\alpha_{m,n,h}$$
$$= flip_{P_n}^{\overline{P}_m}(J_m \otimes I_{P_n})\alpha_{m,n,h}$$

Thus, by Propositions 2.4 and 5.2, we have

$$flip_{P_n}^{\overline{P}_m}(J_m \otimes J_n^*)\eta_{m,n,h} = (I_{P_n} \otimes J_m)flip_{P_n}^{P_m}\alpha_{m,n,h} = (-1)^h(I_{P_n} \otimes J_m)\alpha_{n,m,h}$$
$$= (-1)^h\eta_{n,m,h}.$$

5.2. The dual map of $\Phi_{m,n,h}$.

For Hilbert spaces H, K and a linear map $\Phi : End(H) \longrightarrow End(K)$, the dual map of Φ is defined to be the unique map $\Phi^* : End(K) \longrightarrow End(H)$ such that $\langle B | \Phi(A) \rangle_{End(K)} = \langle \Phi^*(B) | A \rangle_{End(H)}$ for all $A \in End(H), B \in End(K)$. One can easily check that if $\Phi : End(H) \longrightarrow End(K)$ is a quantum channel, then

• Φ^* is a quantum channel if and only if $\Phi(I_H) = I_K$.

- If {T_j : 1 ≤ j ≤ d} are Kraus operators for Φ, then {T_j^{*} : 1 ≤ j ≤ d} are Kraus operators for Φ^{*}.
- If G is a group and π_H, π_K are representations of G on the Hilbert spaces H, K, then Φ is G-equivariant map if and only if Φ^* is G-equivariant.

To obtain a relation between $\Phi_{m,n,h}$ and $\Phi_{m,n,h}^*$, we examine their Choi matrices. Recall that by Proposition 3.8, we have

$$flip_{P_r}^{\overline{P}_m}\left(J_m \otimes J_r^*\right)\left(Vec(T_{n-j})\right) = (-1)^{m-j}Vec(T_j^*)$$

Proposition 5.5. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let $\mathcal{T}_{m,r} = flip_{P_r}^{\overline{P}_m} (J_m \otimes J_r^*)$. Then $C(\Phi_{m,n,h}^*) = \mathcal{T}_{m,r}C(\Phi_{m,n,h})\mathcal{T}_{m,r}^*$.

Proof.

Let $\{T_j : 0 \le j \le n\}$ be the EPOSIC Kraus operators for $\Phi_{m,n,h}$. As $\{T_j^* : 0 \le j \le n\}$ are Kraus operators for $\Phi_{m,n,h}^*$, then by Propositions 1.9 and 3.8, we have

$$C(\Phi_{m,n,h}^{*}) = \sum_{j=0}^{n} Vec(T_{j}^{*}) \left(VecT_{j}^{*} \right)^{*} = \sum_{j=0}^{n} \mathcal{T}_{m,r} Vec(T_{n-j}) \left(\mathcal{T}_{m,r} Vec(T_{n-j}) \right)^{*}$$

$$= \sum_{j=0}^{n} \mathcal{T}_{m,r} Vec(T_{n-j}) \left(Vec(T_{n-j}) \right)^{*} \mathcal{T}_{m,r}^{*}$$

$$= \mathcal{T}_{m,r} \left(\sum_{j=0}^{n} Vec(T_{n-j}) \left(Vec(T_{n-j}) \right)^{*} \right) \mathcal{T}_{m,r}^{*}$$

$$= \mathcal{T}_{m,r} \left(\sum_{j=0}^{n} Vec(T_{j}) \left(Vec(T_{j}) \right)^{*} \right) \mathcal{T}_{m,r}^{*}$$

$$= \mathcal{T}_{m,r} C(\Phi_{m,n,h}) \mathcal{T}_{m,r}^{*}.$$

Corollary 5.6. For $m, n, h \in \mathbb{N}$ with $0 \le h \le \min\{m, n\}$, let r = m + n - 2h. Then

$$\Phi_{m,n,h}^* = \frac{r+1}{m+1} \Phi_{r,n,n-h}$$

Proof.

It suffices to show that $C(\Phi_{m,n,h}^*) = \frac{r+1}{m+1}C(\Phi_{r,n,n-h})$. By Proposition 5.5, this is equivalent to $\mathcal{T}_{m,r}C(\Phi_{m,n,h})\mathcal{T}_{m,r}^* = \frac{r+1}{m+1}C(\Phi_{r,n,n-h})$, and by Proposition 3.13, it is

equivalent to $\mathcal{T}_{m,r}q_{m,r,m-h}\mathcal{T}_{m,r}^* = q_{r,m,m-h}$ where $q_{m,r,m-h}$, $q_{r,m,m-h}$ are the projections on the SU(2)-irreducible subspaces of $P_m \otimes \overline{P}_r$ and $P_r \otimes \overline{P}_m$ respectively.

By Corollary 5.4, one has

$$\mathcal{T}_{m,r}q_{m,r,m-h}\mathcal{T}_{m,r}^{*} = \mathcal{T}_{m,r}\eta_{m,r,m-h}\eta_{m,r,m-h}^{*}\mathcal{T}_{m,r}^{*}$$

$$= flip_{P_{r}}^{\overline{P}_{m}} \left(J_{m} \otimes J_{r}^{*}\right)\eta_{m,r,m-h} \left(flip_{P_{r}}^{\overline{P}_{m}} \left(J_{m} \otimes J_{r}^{*}\right)\eta_{m,r,m-h}\right)^{*}$$

$$= \eta_{r,m,m-h}\eta_{r,m,m-h}^{*} = q_{r,m,m-h}.$$

Remark 5.7. The dual map for $\Phi_{m,n,h}$ is a channel if and only if n = 2h, in which case $\Phi_{m,2h,h}^*$ is $\Phi_{m,2h,h}$.

6. Application in operator algebra: An example of positive non-completely positive map

In this section, using EPOSIC channels, we derive a new example of positive map that is not completely positive. We begin with reviewing notions we need.

Definition 6.1. Let H and K be Hilbert spaces, a linear map $\Phi : End(H) \longrightarrow End(K)$ is said to be

- (1) positive map, if $\Phi(A) \ge 0$ for any positive matrix $A \in End(H)$.
- (2) *n*-positive map, if $\Phi \otimes I_n$ is positive where $\Phi \otimes I_n : End(H) \otimes M_n \longrightarrow End(K) \otimes M_n$ given by $A \otimes B \longmapsto \Phi(A) \otimes B$ and extended by linearity.
- (3) completely positive map, if it is *n*-positive for each $n \ge 1$.

Clearly, any completely positive map is automatically positive, but the converse is not true. In fact there are some examples of positive, non completely positive maps. Here, we use the EPOSIC channels EC(1,m), $m \in \mathbb{N} \setminus \{0\}$ to derive a new example of these maps. Recall that for $m \in \mathbb{N} \setminus \{0\}$, the EC(1,m) consists of two EPOSIC channels from $End(P_1)$ to $End(P_m)$, namely $EC(1,m) = \{\Phi_{m,m+1,m}, \Phi_{m,m-1,m-1}\}$. Recall also that P_1 has the canonical basis $\{f_0^1, f_1^1\}$ where $f_0^1(x_1, x_2) = x_2$, and $f_1^1(x_1, x_2) = x_1$. **Lemma 6.2.** Let $h \in P_1$ with ||h|| = 1. Then

- (1) There exist $g_h \in SU(2)$ such that $\rho_{1(g_h)}(f_0^1) = h$.
- (2) If $\Phi : End(P_1) \longrightarrow End(P_m)$ is an SU(2)-equivariant map then the matrices $\Phi(hh^*)$ and $\Phi(E_{11})$ have the same eigenvalues.

Proof.

- (1) Since *h* is a unit element in P_1 then *h* can be written as $u_0 f_0^1 + u_1 f_1^1$ for some $u_0, u_1 \in \mathbb{C}$ that satisfy $u_0^2 + u_1^2 = 1$. Let $g_h = \begin{bmatrix} \overline{u}_0 & u_1 \\ -\overline{u}_1 & u_0 \end{bmatrix} \in SU(2)$. Then by Equation 2.1, we have $(\rho_1(g_h)f_0^1)(x_1,x_2) = f_0^1(\overline{u}_0x_1 - \overline{u}_1x_2, u_1x_1 + u_0x_2) = u_1x_1 + u_0x_2 = u_0f_0^1(x_1,x_2) + u_1f_1^1(x_1,x_2) = h(x_1,x_2)$.
- (2) By item (1), $hh^* = \rho_1(g_h) f_0^{1} f_0^{1^*} \rho_1^*(g_h) = \rho_1(g_h) E_{11} \rho_1^*(g_h)$, and by equivarince property of Φ we have that $\Phi(hh^*) = \Phi(\rho_1(g_h) E_{11} \rho_1^*(g_h)) = \rho_1(g_h) \Phi(E_{11}) \rho_1^*(g_h)$ which gives the result.

By direct computations using the formula of ε_i^j (Corollary 2.15), and the equation in Corollary 3.3, one can show:

Lemma 6.3. For $m \in \mathbb{N} \setminus \{0\}$

(1)
$$\Phi_{m,m+1,m}(E_{11}) = \sum_{j=0}^{m} \frac{2(m-j+1)}{(m+1)(m+2)} E_{m-j+1,m-j+1}$$

(2) $\Phi_{m,m-1,m-1}(E_{11}) = \sum_{j=0}^{m-1} \frac{2(j+1)}{m(m+1)} E_{m-j,m-j}.$

Proposition 6.4. For $m \in \mathbb{N} \setminus \{0\}$ and $\alpha \in \mathbb{R}$, the map $\Phi_{m,m+1,m} - \alpha \Phi_{m,m-1,m-1}$ is a positive map if and only if $\alpha \leq \frac{1}{m+2}$.

Proof.

Let A be a positive matrix in $End(P_1)$. By the spectral theorem there exist an orthonormal basis $\{x_1, x_2\}$ of P_1 , and non-negative numbers λ_1, λ_2 such that

 $A = \sum_{i=1}^{2} \lambda_i x_i x_i^*$. To show that $\Phi := \Phi_{m,m+1,m} - \alpha \Phi_{m,m-1,m-1}$ is positive, it suffices to show the positivity of $\Phi(x_i x_i^*)$, and by Lemma 6.2, this is equivalent to check the positivity of $\Phi(E_{11})$. Note that by Lemma 6.3, we have

$$\Phi(E_{11}) = \Phi_{m,m+1,m}(E_{11}) - \alpha \Phi_{m,m-1,m-1}(E_{11}) \ge 0$$

if and only if

$$\sum_{j=0}^{m} \frac{2(m-j+1)}{(m+1)(m+2)} E_{m-j+1,m-j+1} - \alpha \sum_{j=0}^{m-1} \frac{2(j+1)}{m(m+1)} E_{m-j,m-j} \ge 0$$

if and only if

$$\frac{2}{m+2}E_{m+1,m+1} + \sum_{j=0}^{m-1} \left[\frac{2(m-j)}{(m+1)(m+2)} - \alpha \frac{2(j+1)}{m(m+1)}\right] E_{m-j,m-j} \ge 0$$

Thus, $\Phi(E_{11}) \ge 0$ if and only if $\alpha \le \min\{\frac{m(m-j)}{(m+2)(j+1)} : 0 \le j \le m-1\}$. Since the map $f(t) = \frac{(m-t)}{t+1}$ is decreasing map for $0 \le t \le m-1$, then $\min\{\frac{m(m-j)}{(m+2)(j+1)} : 0 \le j \le m-1\} = \frac{1}{m+2}$. Consequently, $\Phi(E_{11}) \ge 0$ if and only if $\alpha \le \frac{1}{m+2}$.

Using the formula for ε_i^j in Corollary 2.15, we get:

Lemma 6.5. For $m \in \mathbb{N} \setminus \{0\}$, we have

(1)
$$\varepsilon_1^{0}{}_{(m,1,0)} = \sqrt{\frac{m}{m+1}}, \ \varepsilon_1^{1}{}_{(m,1,0)} = \sqrt{\frac{1}{m+1}}, \ \varepsilon_0^{0}{}_{(m,1,1)} = \sqrt{\frac{1}{m+1}}, \ \text{and} \ \varepsilon_0^{1}{}_{(m,1,1)} = -\sqrt{\frac{m}{m+1}}.$$

Proposition 6.6. For $m \in \mathbb{N} \setminus \{0\}$, and $\alpha > 0$, the map $\Phi_{m,m+1,m} - \alpha \Phi_{m,m-1,m-1}$ is not completely positive.

Proof.

Let $\Phi = \Phi_{m,m+1,m} - \alpha \Phi_{m,m-1,m-1}$. We show that $-\frac{2\alpha}{m}$ is an eigenvalue of $C(\Phi)$ with a corresponding eigenvectors $v = \sqrt{m}(f_0^m \otimes f_0^1) + f_1^m \otimes f_1^1$. By Proposition 3.13, $C(\Phi_{m,m+1,m}) = \frac{2}{m+2}\eta_{m,1,0}\eta_{m,1,0}^*$, and $C(\Phi_{m,m-1,m-1}) = \frac{2}{m}\eta_{m,1,1}\eta_{m,1,1}^*$. Thus, $C(\Phi_{m,m+1,m})(v) = \frac{2}{m+2}\eta_{m,1,0}\eta_{m,1,0}^*(\sqrt{m}(f_0^m \otimes f_0^1) + f_1^m \otimes f_1^1)$

$$= \frac{2}{m+2} \eta_{m,1,0} \left[\left(-\sqrt{m} \varepsilon_1^{1}{}_{(m,1,0)} + \varepsilon_1^{0}{}_{(m,1,0)} \right) f_1^{m+1} \right]$$

$$= \frac{2}{m+2} \eta_{m,1,0} \left[\left(-\sqrt{\frac{m}{m+1}} + \sqrt{\frac{m}{m+1}} \right) f_1^{m+1} \right] \text{ (Lemma 6.5)}$$

$$= \frac{2}{m+2} \eta_{m,1,0} \left[0 \times f_1^{m+1} \right] = 0.$$

Similarly,

$$C(\Phi_{m,m-1,m-1})(v) = \frac{2}{m}\eta_{m,1,1}\eta_{m,1,1}^{*}\left(\sqrt{m}(f_{0}^{m}\otimes f_{0}^{1}) + f_{1}^{m}\otimes f_{1}^{1}\right)$$

$$= \frac{2}{m+2}\eta_{m,1,1}\left[\left(-\sqrt{m}\varepsilon_{0}^{1}(m,1,1) + \varepsilon_{0}^{0}(m,1,1)\right)f_{0}^{m-1}\right]$$

$$= \frac{2}{m}\left(\frac{m+1}{\sqrt{m+1}}\right)\eta_{m,1,1}(f_{0}^{m-1}) \text{ (Lemma 6.5)}$$

$$= \frac{2\sqrt{m+1}}{m}\left[\sum_{j=0}^{1}(-1)^{j}\varepsilon_{0}^{j}f_{1-j}^{m}\otimes f_{1-j}^{1}\right]$$

$$= \frac{2\sqrt{m+1}}{m}\left(\frac{1}{\sqrt{m+1}}f_{1}^{m}\otimes f_{1}^{1} + \sqrt{\frac{m}{m+1}}f_{0}^{m}\otimes f_{0}^{1}\right)$$

$$= \frac{2}{m}(f_{1}^{m}\otimes f_{1}^{1} + \sqrt{m}f_{0}^{m}\otimes f_{0}^{1}) = \frac{2}{m}v.$$

Thus, we have $C(\Phi)(v) = C(\Phi_{m,m+1,m})(v) - \alpha C(\Phi_{m,m-1,m-1})(v) = -\frac{2\alpha}{m}v$. Hence, $-\frac{2\alpha}{m}$ is a negative eigenvalue of $C(\Phi)$ for any $\alpha > 0$.

It is straightforward to show that if Φ is *n*-positive then it is *s*-positive for $1 \le s \le n$. Thus, combining the result of the proposition above and CHOI result [11, p.35] about the *n*-positivity, we get the following:

Corollary 6.7. For $0 < \alpha$ and $m \in \mathbb{N} \setminus \{0\}$ the map $\Phi_{m,m+1,m} - \alpha \Phi_{m,m-1,m-1}$ is not n-positive for any n > 1.

References

- C.H. Bennett and P.W. Shor, Quantum Information Theory, IEEE Trans. Info. Theory Vol 44 Issue 6 (1998) 2724–2748.
- [2] T. Brocker and T. Dieck, Representations of Compact Lie Groups, Springer- Verlag New York Inc 1985.
- [3] M. Choi, Completely positive linear maps on complex matrices, Lin. Alg. Appl. 10 (1975) 285-290.
- [4] M. Fannes, B. Haegeman, M. Mosonyi and D. Vanpeteghem, Additivity of minimal entropy output for a class of covariant channels, arXiv: quant-ph/040195v1.
- [5] M. Hayashi, Quantum Information An Introduction, Springer-Verlag Berlin Heidelberg, 2006.

- [6] A. Holevo, Additivity conjecture and covariant channel, Int. J. Quant.Inform. 3 (1) (2005) 41-48.
- [7] A. Holevo, A note on covariant dynamical semigroup, Rep. Math. Phys. 32 (1993) 211-216.
- [8] A. Holevo, Complementary channel and the additivity problem, Theory Probab. Appl. Vol 51. No 1, pp.92-100.
- [9] A. Holevo, The additivity problem in quantum information theory, ICM 2006 Madrid, Section Lectures, Mathematical Aspect of Computer Science.
- [10] M. Nielsen, I. Chuang, Quantum Computation and Quantum Information, Cambridge university press 2000.
- [11] V. Paulson, Completely bounded maps and operator algebras, Cambridge university press, UK (2002).
- [12] C. Procesi, Lie Groups An Approach through Invariants and Representations, Springer Science+ Business Media, LLC, 2007.
- [13] J.-P.Serre, Linear Representations of Finite Groups, Springer-Verlag New York Inc, 1977.
- [14] S. STERNBERG, Group Theory And Physics, Cambridge University Press 1994.
- [15] F. Verstraeteab, H. Verscheldea, On Quantum Channels, arXiv: quant-ph/0202124v2
- [16] N.Ja. Vilenkin, A.U. Klimyk, Representation of Lie Groups and Special Functions, Volume 1:Simplest Lie Groups, Special Functions and Integral Transforms. Kluwer academic publishers. Dordrecht/Boston/London (1991).
- [17] K.Vollbrecht, R. Werner, Entanglement Meausers under symmetry, arXiv: quant-ph/0010095.
- [18] J. Watrous, CS 766/QIC Theory of Quantum Information, Institute for Quantum Computing, University of Waterloo, Fall 2011.

LIST OF EQUATIONS THAT ARE USED IN THE COMPUTATIONS.

For any $m, n \in \mathbb{N}$ and $0 \le h \le \min\{m, n\}$, let r = m + n - 2h then

•
$$c_{m,n,h} = \frac{((m-h)!)^2}{(m+n-2h)! \ m! \ n! \left(\sum_{k=0}^h \frac{\binom{h}{k}^2}{\binom{m}{h-k}\binom{n}{k}}\right)}.$$

• $\beta_{i,s,j}^{m,n,h} = (-1)^s \sqrt{\frac{c_{m,n,h} \ r! \ m! \ n!}{\binom{r}{i}\binom{m}{(i-j+h)}\binom{n}{j}}} \frac{\binom{h}{s}\binom{n-h}{j-s}\binom{m-h}{i-j+s}}{(m-h)!}.$

• $k_1(i) = \max\{0, -m+i+h\}, k_2(i) = \min\{i, n-h\}, \text{ and } l_{ij} = h+i-j.$

•
$$\varepsilon_{i^{(m,n,h)}}^{j} = \sum_{s=\max\{0,j-i,j+h-n\}}^{\min\{h,j,j+m-i-h\}} \beta_{i,s,j}^{m,n,h}.$$

•
$$B(i) = \{j : k_1(i) \le j \le k_2(i) + h\}.$$

•
$$\{f_l^m = a_m^l x_1^l x_2^{m-l} : 0 \le l \le m\}$$
 where $a_m^l = \frac{1}{\sqrt{l!(m-l)!}}$

•
$$J_m(f_l^m) = (-1)^l f_{m-l}^m$$
 and $J_m^*(f_l^m) = (-1)^{m-l} f_{m-l}^m$.

•
$$P_m \otimes P_n \cong \bigoplus_{h=0}^{\min\{m,n\}} P_{m+n-2h}.$$

•
$$\alpha_{m,n,h}\left(f_{i}^{r}\right) = \sum_{j \in B(i)} \varepsilon_{i}^{j}{}^{(m,n,h)} f_{lij}^{m} \otimes f_{j}^{n}.$$

• $\eta_{m,n,h} = (I_{P_m} \otimes J_n) \alpha_{m,n,h} : P_{m+n-2h} \longrightarrow P_m \otimes \overline{P}_n.$

•
$$C(\Phi_{m,n,h}) = \frac{r+1}{n+1}q_{m,r,m-h}$$
.