

Explicit formulae for primes in arithmetic progressions, I^{*†}

Tomohiro Yamada

Abstract

We shall give an explicit formula for $\psi(x, q, a)$ with an error term of the form $C/\log^\alpha x$ under the condition that $q < \log^{\alpha_1} x$ is nonexceptional, for various values of α and α_1 . We shall also give an explicit formula for $\psi(x, q, a)$ with error terms $C/\log^A x$ working whether q is exceptional or nonexceptional, but under the condition that $\frac{0.4923A}{\pi}q^{1/2}\log^2 q < \log x/\log\log x$. Moreover, we shall give an explicit form of Bombieri-Vinogradov theorem over non-exceptional moduli.

1 Introduction

The prime number theorem for arithmetic progressions states that $\psi(x, q, a) = \sum_{p^e \leq x, p^e \equiv a \pmod{Q}} \log p$ is $x/\varphi(k) + o(x)$ as $x \rightarrow \infty$.

Moreover, Siegel-Walfisz theorem states that for any $\alpha > 0$, the error term can be bounded by $O(x \exp -\frac{c \log^{3/5} x}{(\log\log x)^{1/5}})$ uniformly for $q < \log^\alpha x$.

Prime number theorem for arithmetic progressions has many applications in number theory. Naturally we need have explicit estimates for the error term for the ordinary prime number theorem to give explicit results in such applications.

The classical results are Rosser[12] and Rosser and Schoenfeld[13][14], giving explicit estimates for the ordinary prime number theorem (i.e. the case $q = 1, 2$). It is after 1980's that various explicit formulae for primes

^{*}2010 Mathematics Subject Classification: 11N13.

[†]Key words and phrases: Primes in arithmetic progressions.

in arithmetic progression have been obtained by several authors such as McCurley[7, 8, 9], Ramaré and Rumely[11], Dusart[4], Liu and Wang[5] and Kadiri[6].

The purpose of this paper is to give an explicit formula with an error term of the form $C/\log^{\alpha_2} x$ under the condition $q < \log^{\alpha_1} x$ for various values of α_1 and α_2 . Our main results are the following two theorems.

Theorem 1.1. *Let $\alpha_1, \alpha_2, Y_0 = \log \log X_0, C$ be constants given in some column of Table 5. Let q be a modulus $\leq \log^{\alpha_1} x$. Let $E_0 = 1$ and β_0 denote the Siegel zero modulo q if it exists and $E_0 = 0$ otherwise. If $(a, q) = 1$ and $x \geq X_0$, then*

$$-1 + x^{-1} \sum_{\chi \pmod{q}} |\psi(x, \chi)| < \frac{C}{\log^{\alpha_2} x} + E_0 \frac{x^{\beta_0 - 1}}{\beta_0} \quad (1)$$

and

$$\frac{\varphi(q)}{x} \left| \psi(x; q, a) - \frac{x}{\varphi(q)} \right| < \frac{C}{\log^{\alpha_2} x} + E_0 \frac{x^{\beta_0 - 1}}{\beta_0}. \quad (2)$$

Theorem 1.2. *Let $A, Y_0 = \log \log X_0, C$ be constants given in some column of Table 5 with $(\alpha_1, \alpha_2) = (1, A)$ and $C' = C + 1/(1 - AY_0/e^{Y_0})$. If $x > X_0$ and $\frac{0.4923A}{\pi} q^{1/2} \log^2 q < \log x / \log \log x$, then we have*

$$-1 + x^{-1} \sum_{\chi \pmod{q}} |\psi(x, \chi)| < \frac{C'}{\log^A x} \quad (3)$$

and

$$\frac{\varphi(q)}{x} \left| \psi(x; q, a) - \frac{x}{\varphi(q)} \right| < \frac{C'}{\log^A x}. \quad (4)$$

One of applications of our formula is obtaining (almost) completely explicit Bombieri-Vinogradov theorem.

The well-known Bombieri-Vinogradov theorem, first proved by Bombieri[2], states that for any constant $A > 0$, there exists some constant B such that

$$\sum_{\substack{q \leq \frac{x^{\frac{1}{2}}}{\log^B x} \\ (a, q)=1}} \max_{2 \leq y \leq x} \left| \psi(y; q, a) - \frac{\psi(y)}{\varphi(q)} \right| = O \left(\frac{x}{\log^A x} \right), \quad (5)$$

where $\pi(y; q, a)$ denotes the number of primes $\leq y$ congruent to $a \pmod{q}$. Akbary and Hamrook[1] gave an explicit formula with $B = A + \frac{9}{2}$, with moduli limited to those which is free of small prime factors. Our explicit formula enables us to give (almost) completely explicit version of Bombieri-Vinogradov theorem.

Before stating our version of Bombieri-Vinogradov theorem, we introduce a result concerning zeros of Dirichlet L-functions.

Lemma 1.3. *Define $\Pi(s, q) = \prod_{\chi \pmod{q}} L(s, \chi)$, $R_0 = 6.3970$ and $R_1 = 2.0452$. Then the function $\Pi(s, q)$ has at most one zero $\rho = \beta + it$ in the region $0 \leq \beta \leq 1 - 1/R_0 \log \max\{q, q|t|\}$. If such a zero exists, then it must be real and simple and must correspond to a nonprincipal real character $\chi \pmod{q}$. Moreover, for any given Q_1 , such a zero must satisfy $\beta < 1 - 1/2R_1 \log Q_1$ except possibly one modulus $\leq Q_1$.*

This result easily follows from Theorems 1.1 and 1.3 of [6].

For a given Q_1 , we call a modulus $q_0 \leq Q_1$ to be exceptional up to Q_1 if $\Pi(s, q)$ has a zero $\rho = \beta + it$ with $\beta \geq 1 - 1/2R_1 \log Q_1$. Now we shall state our explicit version of Bombieri-Vinogradov theorem.

Theorem 1.4. *Let A be an integer with $6 \leq A \leq 10$ and $Y_0 = 10.1(A = 6), 10.5(A = 7), 10.9(A = 8), 11.2(A = 9)$ and $11.5(A = 10)$, C be the constant in Table 5 corresponding to the column $(\alpha_1, \alpha, Y) = (A, A-3, Y_0)$, c_0, c_1 be the constants defined by*

$$c_0 = \frac{2^{\frac{13}{2}}}{9\pi \log 2} \left(\frac{1}{3} + \frac{3}{2 \log 2} \right) \left(\frac{2 + \log(\log 2 / \log(4/3))}{\log 2} \right) \sqrt{\frac{\psi(113)}{113}} \quad (6)$$

and

$$c_1 = \prod_p \left(1 + \frac{1}{p(p-1)} \right) = \frac{\zeta(2)\zeta(3)}{\zeta(6)}. \quad (7)$$

We note that $c_0 < 48.833$ and $c_1 < 1.9436$.

Let $Q = \frac{x^{1/2}}{\log^A x}$ and $1 \leq Q_1 \leq \log^A x$. Let q_0 denote the exceptional modulus up to Q_1 if it exists. If $\log \log x > X_0$, then the inequality

$$\begin{aligned} & \sum_{q \leq Q, q_0 \nmid q} \max_a \left| \psi(x, q, a) - \frac{\psi(x)}{\varphi(q)} \right| \\ & < x^{\frac{1}{2}} + \frac{c_1 c_0 (2 + e^{-800}) x}{\log^{A-\frac{9}{2}} x} + \frac{2c_0 c_1 x \log^{\frac{9}{2}} x}{Q_1} + \frac{c_1^2 (C_0 + e^{-17}) x (1 + A \log \log x)}{2 \log^{A-4} x}. \end{aligned} \quad (8)$$

holds.

Our explicit Bombieri-Vinogradov theorem is not completely explicit in two senses: First, $\max_{2 \leq y \leq x}$ should be removed. But in many applications, this would not cause a serious problem. Second, we must avoid exceptional

Table 1: constants C_1, C_2

C_1	C_2
0.247	9.359
0.264	8.049
0.281	7.323
0.298	6.828
0.315	6.455
0.332	6.156
0.349	5.907
0.365	5.694
0.382	5.506
0.399	5.338

moduli. However, our version would still have several applications. Indeed, combined with the author's explicit linear sieve [18] and a technique which discharges us from large exceptional moduli, the author obtained explicit version of Chen's celebrated theorem that any sufficiently large even numbers can be written as the sum of a prime and a product of at most two primes[19]; the author showed that $\exp \exp 36$ suffices.

For calculations of constants, we used PARI-GP. Our script is available from <http://tyamada1093.web.fc2.com/math/files/prim0003pari.txt> and can be used to calculate constants for arbitrary values of α and α_1 .

2 Preliminary Lemmas

We combine the method of McCurley in [8] and results of Kadiri concerning the distribution of zeros of Dirichlet L-functions in [6]. We begin by the following zero-free region.

Let $T = \log^\alpha x$, where $\alpha = \alpha_1 + \alpha_2 + 3 \geq 5$. We let two constants C_1, C_2 be given in any column of Table 1 and put

$$F(T) = \frac{T}{\pi} \log \frac{qT}{2\pi e}, R(T) = C_1 \log qT + C_2. \quad (9)$$

Then, Theorem 1 of [17] gives that $N(\chi, T) \leq F(T) + R(T)$.

For a Dirichlet character χ modulo q , we use $z(\chi)$ to represent the set of zeros $\rho = \beta + i\gamma$ of $L(s, \chi)$ with $\beta \geq 0$ and $\rho \neq 0$. Moreover, for $H > 1$ and

$R > 0$, we use $z_0(\chi, H, R)$ to represent the set of zeros $\rho = \beta + i\gamma$ of $L(s, \chi)$ with $\frac{1}{2} < \beta \leq 1 - 1/R \log qH$, $|\gamma| \leq H$ and $z_1(\chi, H, R)$ to represent the set of zeros $\rho = \beta + i\gamma$ of $L(s, \chi)$ with $1 - 1/R \log qH < \beta \leq 1 - 1/R_0 \log qH$, $|\gamma| \leq H$. Of course, GRH implies that these sets would be empty.

We set

$$\Sigma = \sum_{\chi} \sum_{\rho \in z(\chi), |\gamma| \leq T} \frac{x^{\beta-1}}{|\rho|}. \quad (10)$$

We can estimate error terms in the prime number theorem in arithmetic progressions by these sums.

Similarly to Theorem 8 of [5], we have the following lemma.

Lemma 2.1. *If $t > \exp \exp 6.8$, then we have*

$$|\psi(t, \chi) - \delta(\chi)t| \leq \sum_{\rho \in z(\chi), |\gamma| \leq T} \frac{x^{\beta}}{|\rho|} + 1.3833 \frac{t \log^2 x}{T}, \quad (11)$$

where $\delta(\chi) = 1$ if χ is principal and $\delta(\chi) = 0$ otherwise.

Proof. We shall mainly consider the case χ is primitive and show that (11) holds with 1.3833 replaced by 1.38329. If $\chi \pmod{q}$ is induced by the primitive character $\chi_1 \pmod{q_1}$, then $|\psi(t, \chi) - \psi(t, \chi_1)| \leq \log q / \log 2 \leq A \log L / \log 2 \leq e^{-20} t L^2 / T$ and (11) holds. If $\chi = \chi_0$ is the principal character, then the following argument can be proceeded with $\psi(t, \chi)$ replaced by $\psi(t, \chi) - t$.

We begin by seeing that Lemma 8 of [3] gives $\sum_{|\gamma-T| \leq 1} 1 \leq 3.5(0.5 \log q(T+2) + 0.59773) \leq C_3 \log(qT) - 1$, where $C_3 < 1.83$, so that there exists a real T_0 such that $|T_0 - T| \leq 1$ and

$$\frac{1}{|\gamma - T_0|} \leq C_3 \log(qT). \quad (12)$$

We can confirm that L and T are enough large for us to obtain

$$\psi(t, \chi) = \frac{1}{2\pi i} \int_{b-iT_0} b + iT_0 - \frac{L'}{L}(s, \chi) \frac{t_0^s}{s} ds + R_6, \quad (13)$$

where $t_0 = \lfloor t + 0.5 \rfloor$, and

$$\psi(2.5, \chi) = \frac{1}{2\pi i} \int_{b-iT_0} b + iT_0 - \frac{L'}{L}(s, \chi) \frac{2.5^s}{s} ds + R_7, \quad (14)$$

with $|R_6| \leq 1.38305tL^2/T$ and $|R_7| \leq 2L/(T-1) \leq e^{-1000}tL^2/T$. Thus (4.5) in [5] holds with $|R_8| \leq |R_6| + |R_7| + \log 2 < 1.38306tL^2/T$ and (4.6) in [5] holds with $|R_9| \leq |R_8| + \exp(\log(t+1)/L)TL/(2\pi) \leq 1.38307tL^2/T$.

Using Lemmas 9 and 9' of [3], we have

$$\left| \frac{L'}{L}(\sigma + iT, \chi) \right| \leq 4.6 \log(qT) + C_3^2 \log^2(qT) \quad (15)$$

for $-0.5 \leq \sigma \leq b$, and

$$\left| \frac{L'}{L}(-0.5 + it, \chi) \right| \leq 1.5 \log q(T+2) + 1.79319 + \frac{35}{8} \log(T+2) + 13.4. \quad (16)$$

Thus we obtain

$$\left| \frac{1}{2\pi i} \int_{-0.5 \pm iT_0} b \pm iT_0 - \frac{L'}{L}(s, \chi) \frac{t_0^s - 2.5^s}{s} ds \right| \leq \frac{0.00003tL^2}{T} \quad (17)$$

and

$$\left| \frac{1}{2\pi i} \int_{-0.5 - iT_0} -0.5 + iT_0 - \frac{L'}{L}(s, \chi) \frac{t_0^s - 2.5^s}{s} ds \right| \leq 1.31 \log T \log(qT) \quad (18)$$

Hence (4.8) in [5] holds with $|R_{10}| \leq 1.3831tL^2/T$. Since $3.5(0.5 \log q(T+2) + 0.59773) < 0.000092(T-1)L^2/T$, (4.9) in [5] holds with the constant 1.38039 replaced by 1.3832.

By Theorem 1 of [17], we have $N(\chi, 1) \leq 0.399 \log q + 5.338$ and therefore

$$\begin{aligned} \sum_{|\gamma| \leq 1} \frac{1}{\Re \rho} &\leq \frac{0.4923q^{\frac{1}{2}} \log^2 q}{\pi} + R_0(\log q)N(\chi, 1) \\ &\leq 0.33q^{\frac{1}{2}} \log^2 q \leq 0.39\alpha_1^2 L^{\frac{\alpha_1}{2}} \log^2 L \\ &\leq e^{-1000} \frac{tL^2}{T}. \end{aligned} \quad (19)$$

We see that, using Theorem 6 of [5], or Theorem 1 of [17],

$$\sum_{1 \leq |\gamma| \leq T+1} \frac{1}{|\rho|} \leq \frac{N(\chi, T+1)}{T+1} + \int_1^{T+1} \frac{N(\chi, y)}{y^2} dy \leq \frac{\log^2(T+1)}{2\pi} \leq e^{-1000} \frac{tL^2}{T}. \quad (20)$$

Hence $\sum_{|\gamma| \leq T+1} |2.5^\rho / \rho| \leq e^{-999} tL^2/T$.

Similarly, observing that $t^{1-\beta}1 - \beta$ with $1 - \pi/(0.4934q^{1/2} \log^2 q) \leq \beta \leq 1/2$ takes the maximum at $\beta = 1/2$, we obtain

$$\begin{aligned} \sum_{|\gamma| \leq 1, \beta \leq 1/2} \frac{t^{\frac{1}{2}}}{\Re \rho} &\leq t^{\frac{1}{2}}(2 + R_0(\log q)N(\chi, 1)) \\ &\leq 10t^{\frac{1}{2}} \log^2 q \leq 10\alpha_1^2 t^{\frac{1}{2}} \log^2 L \\ &\leq e^{-1000} \frac{tL^2}{T} \end{aligned} \quad (21)$$

and

$$\sum_{1 \leq |\gamma| \leq T+1} \frac{1}{|\rho|} \leq \frac{N(\chi, T+1)}{T+1} + \int_1^{T+1} \frac{N(\chi, y)}{y^2} dy \leq \frac{\log^2(T+1)}{2\pi} < e^{-500} \frac{tL^2}{T}. \quad (22)$$

This proves (11) with 1.3833 replaced by 1.38329. In the imprimitive case, (11) follows from the argument in the beginning of the proof and the proof is complete. \square

So that, it suffices to obtain upper bounds for Σ . The following lemmas, combined with Lemma 1.3, enable us to obtain these upper bounds. However, an upper bound for Σ obtained by Lemma 1.3 would be large. We can improve an upper bound by dividing the sum Σ as follows.

We additionally set the following two sums

$$\begin{aligned} \mu_0 &= \sum_{\chi} \sum_{\rho \in z_0(\chi, T, R)} \frac{x^{\beta-1}}{|\rho|}, \\ \mu_1 &= \sum_{\chi} \sum_{\rho \in z_1(\chi, T, R)} \frac{x^{\beta-1}}{|\rho|}. \end{aligned} \quad (23)$$

Then we can easily see that

$$\Sigma \leq \mu_0 + \mu_1 + E_0 \frac{x^{\beta_0-1}}{\beta_0}. \quad (24)$$

Now results in [5] concerning the number of zeros near the line $\sigma = 1$ enable us to improve an upper bound for Σ by choosing R larger than R_0 .

We can easily see that

$$\sum_{\rho \in z_0(\chi, T, R)} \frac{x^{\beta-1}}{|\rho|} \leq \frac{1}{2} x^{-1/R \log q T} \left(2N(\chi, 1) + \int_1^T \frac{dN(\chi, t)}{t} \right), \quad (25)$$

Using to (3.30) in [8, p. 278], we obtain

$$2N(\chi, 1) + \int_1^T \frac{dN(\chi, t)}{t} \leq \frac{N(\chi, T)}{T} + N(\chi, 1) + \frac{\log^2 T}{2\pi} + F(1) \log T + R(1) + C_1. \quad (26)$$

Since $N(T) \leq F(T) + R(T)$ and $N(1) \leq F(1) + R(1)$, this is at most

$$\leq S := \frac{\alpha^2 + 2\alpha\alpha_1 l^2 + 2\alpha_1 - \alpha \log(2\pi)}{2\pi} l + \frac{R(T)}{T} + 2R(1) - \frac{2(1 + \log(2\pi))}{\pi} + C_1, \quad (27)$$

where $l = \log L = \log \log x$.

Hence we obtain $\mu_0 \leq (\varphi(q)S)x^{-1/R \log qT}/2$ and using (11), we conclude that

$$\sum_{\chi \pmod{q}} |\psi(t, \chi) - \delta(\chi)t| \leq \mu_1 + \frac{qS}{2}x^{-1/R \log qT} + 1.38 \frac{x}{\log^{\alpha_2+1} x}. \quad (28)$$

In order to estimate the remaining term μ_1 , we use the following inequality.

Lemma 2.2. *Let $\rho = \beta + it$ be a zero of $L(s, \chi)$ with $\beta < 1 - 1/R_0 \log qT$, $|t| \leq T$ and $\rho \neq 0$. Let λ be such that $\beta = 1 - \lambda/\log qT$. Then we have*

$$\left| \frac{x^{\rho-1}}{\rho} \right| < \begin{cases} \frac{x^{-1/R_0 \log q}}{1 - \frac{1}{R_0 \log q}} & \text{if } |t| \leq 1, \\ q \exp -2\sqrt{\frac{\log x}{R_0}} & \text{if } |t| > 1, \exp \sqrt{\frac{x}{R_0}} < (qT)^{1/R_0 \lambda}, \\ q \frac{x^{-\lambda/\log qT}}{(qT)^{1/R_0 \lambda}} & \text{otherwise.} \end{cases} \quad (29)$$

Proof. We may assume that $t > 0$ and see that $\left| \frac{x^{\rho-1}}{\rho} \right| < \frac{x^{\beta-1}}{t}$. In the case $|t| < 1$,

$$\left| \frac{x^{\rho-1}}{\rho} \right| < \frac{x^{\beta-1}}{\beta}. \quad (30)$$

Since $\beta > \frac{1}{R_0 \log q} \geq \frac{1}{\log x}$, we have (29) in this case.

If $t \geq 1$, then we have

$$\frac{x^{\beta-1}}{t} < \frac{x^{-1/R_0 \log qt}}{t} \leq q \exp -2\sqrt{\frac{\log x}{R_0}} \quad (31)$$

since

$$\frac{1}{R_0 \log qt} + \frac{\log t}{\log x} = \frac{1}{R_0 \log qt} + \frac{\log qt}{\log x} - \frac{\log q}{\log x} \geq 2\sqrt{\frac{1}{R_0 \log x}} - \frac{\log q}{\log x}. \quad (32)$$

Let $t_0 = \frac{1}{q} \exp \sqrt{\frac{x}{R_0}}$ and $t_1 = q^{\frac{1}{R_0\lambda}-1} T^{\frac{1}{R_0\lambda}}$. Now we may assume that $t_0 \geq t_1$, that is, $qt_0 = \exp \sqrt{\frac{x}{R_0}} \geq (qT)^{1/R_0\lambda} = qt_1$. In the case $t \leq t_1$ we have

$$\frac{x^{-1/R_0 \log qt}}{t} \leq \frac{x^{-1/R_0 \log qt_1}}{t_1} = q \frac{x^{-\lambda/\log qT}}{(qT)^{1/R_0\lambda}} \quad (33)$$

since $\frac{x^{-1/R_0 \log qt}}{t}$ is increasing below t_0 . Moreover, in the other case we have

$$\frac{x^{-\lambda/\log qT}}{t} \leq \frac{x^{-\lambda/\log qT}}{t_1} = q \frac{x^{-\lambda/\log qT}}{(qT)^{1/R_0\lambda}}. \quad (34)$$

□

3 Proof of main results

Now we shall estimate μ_1 , the sum of $\left| \frac{x^{\rho-1}}{\rho} \right|$ over zeros ρ of $L(s, \chi)$ with $1 - R < \log qT < \beta_i < 1 - R_0/\log qT, 1 \leq |t_i| \leq T$.

Let $\rho_1 = \beta_1 + it_1, \rho_2 = \beta_2 + it_2, \dots$ be all zeros of $\Pi(s, q)$ with $\beta_i = 1 - \lambda_i/\log qT, 1 \leq t_i \leq T$ and $1/R_0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$

We shall summarize some results of [5] concerning the distribution of zeros of $\Pi(s, q)$.

Lemma 3.1. *Assume that $qT \geq 8 \times 10^9$. Then $\lambda_1 \leq \eta_i$ implies $\lambda_2 > \xi_i$ for each $i = 1, 2, \dots, 12$, where η_i 's and ξ_i 's are given as in Table 2, and $\lambda_3 > 0.26213$. Moreover, if $qT \geq 10^{11}$, then for each n of Table 3, λ_n must be greater than the corresponding value,*

Proof. Theorem 2 and Table 1 of [5, p.p. 269–270] give the result for λ_1 and λ_2 . The result for λ_3 is just Theorem 1 of [5, p. 265]. The remaining results follow from Table 3-5 of [5, p.p. 277–279]. □

By Lemma 2.2, we have $\left| \frac{x^{\rho-1}}{\rho} \right| < \mu(\lambda)$, where $\mu(\lambda)$ is the function represented by the right-hand side of (29). Now Lemma 3.1 gives an upper bound for μ_1 .

Corollary 3.2. *Assume that $qT \geq 10^{11}$. Then, for some $i = 1, 2, \dots, 11$ we have*

$$\mu_1 < 2 \min_{0 \leq J \leq 12} \left(\mu(\eta_i) + \mu(\xi_{i+1}) + \sum_{j=1}^J M_j \mu(\max\{\xi_{i+1}, \nu_j\}) \right) \quad (35)$$

Table 2: constants η_i and ξ_i

i	η_i	ξ_i
1	0.10367089	0.3534
2	0.12	0.3221
3	0.13	0.3050
4	0.14	0.2891
5	0.15	0.2743
6	0.16	0.2605
7	0.17	0.2477
8	0.18	0.2356
9	0.19	0.2242
10	0.20	0.2135
11	0.206	0.2074
12	0.2067	0.2067

Table 3: constants related to each modulus

n	4	5	6	7	10	18	45
$\lambda_n >$	0.28	0.31	0.32	0.33	0.36	0.39	0.42
n	91	146	332	834	7000		
$\lambda_n >$	0.45	0.46	0.47	0.475	0.478		

Table 4: constants ν_j and M_j in Corollary 3.2

j	1	2	3	4	5	6	7	8
ν_j	0.26213	0.27	0.30	0.32	0.33	0.36	0.39	0.42
M_j	1	1	1	1	4	7	27	57

j	9	10	11	12	13
ν_j	0.45	0.46	0.47	0.475	0.478
M_j	55	186	502	6166	—

with $R = R_{i,J} = \max\{\xi_{i+1}, \nu_{J+1}\}$, where M_j 's and ν_j 's are defined in Table 4.

So that the sum $\varphi(q)Sx^{-1/R \log qT} + \mu_1$ in (28) is at most

$$\max_{1 \leq i \leq 11} \min_{0 \leq J \leq 12} qSx^{-1/R_{i,J} \log qT} + 2 \left(\mu(\eta_i) + \mu(\xi_{i+1}) + \sum_{j=1}^J M_j \mu(\max\{\xi_{i+1}, \nu_j\}) \right). \quad (36)$$

Now we shall prove Theorem 1.1. We shall use (36); we have $qT \geq 10^{11}$ for any value of α_2 in Table 5. For each pair (α_1, α_2) in Table 5, we calculated an upper bound for C in appropriate ranges for each X_0 such that $10 \log \log X_0$ is an integer and we obtained the desired results using PARI-GP (The output is available from <http://tyamada1093.web.fc2.com/math/files/prim0003pariout>). This completes the proof of Theorem 1.1.

Next we shall prove Theorem 1.2. Since our assumption implies that $q < \log^2 x$, we have by (28) with $\alpha_1 = 2$,

$$-1 + x^{-1} \sum_{\chi \pmod{q}} |\psi(x, \chi)| < \frac{C \log \log x}{\log x} + E_0 \frac{x^{\beta_0 - 1}}{\beta_0}. \quad (37)$$

Now assume that $E_0 = 1$ and let $\beta_0 > \frac{1}{2}$ be the Siegel zero of $L(s, \chi)$. Then, Theorem 3 of [5] gives $\beta_0 \leq 1 - \frac{\pi}{0.4923q^{1/2} \log^2 q}$.

Since we have assumed that $\frac{0.4923A}{\pi} q^{1/2} \log^2 q < \log x / \log \log x$, we have

$$\frac{x^{\beta_0 - 1}}{\beta_0} < \frac{x^{-\frac{\pi}{0.4923q^{1/2} \log^2 q}}}{1 - \frac{\pi}{0.4923q^{1/2} \log^2 q}} < \frac{1}{\left(1 - \frac{A \log \log x}{\log x}\right) \log^A x}. \quad (38)$$

Combining (37) and (38), we have

$$\frac{\varphi(q)}{x} \left| \psi(x; q, a) - \frac{x}{\varphi(q)} \right| < \frac{C'}{\log^A x} \quad (39)$$

for $x \geq X_0$ and $\frac{0.4923A}{\pi} q^{1/2} \log^2 q < \log x / \log \log x$.

4 Proof of Theorem 1.4

We see that

$$|\psi(x, \chi) - \psi(x, \chi^*)| \leq \sum_{p^k \leq x, p|q} \log p < \omega(q)(\log x) \leq \log^2(qx), \quad (40)$$

where $\omega(q)$ denotes the number of distinct prime factors of q . Noting that $\psi(x, \chi_0^*) = \psi(x)$, we have

$$\left| \psi(x, q, a) - \frac{\psi(x)}{\varphi(q)} \right| \leq \frac{1}{\varphi(q)} \left| \sum_{\chi \pmod{q}, \chi \neq \chi_0} \psi(x, \chi^*) \right| + \log^2(qx). \quad (41)$$

Hence we obtain

$$\begin{aligned} \sum_{q \leq Q} \max_a \left| \psi(x, q, a) - \frac{\psi(x)}{\varphi(q)} \right| &\leq Q \log^2(Qx) + \sum_{1 < q \leq Q, q_0 \nmid q} \frac{1}{\varphi(q)} \left| \sum_{\chi \pmod{q}, \chi \neq \chi_0} \psi(x, \chi^*) \right| \\ &< x^{\frac{1}{2}} + \sum_{1 < q \leq Q, q_0 \nmid q} \frac{1}{\varphi(q)} \left| \sum_{\chi \pmod{q}, \chi \neq \chi_0} \psi(x, \chi^*) \right|. \end{aligned} \quad (42)$$

Since each character $\chi \pmod{q}$ is induced from a certain primitive character $\chi^* \pmod{q^*}$ with $q^* \mid q$, the last sum is at most

$$\begin{aligned} &\sum_{\substack{m \geq 1, q^* > 1, \\ q_0 \nmid q^*, \\ mq^* \leq Q}} \frac{1}{\varphi(mq^*)} \left| \sum_{\chi \pmod{q^*}}^* \psi(x, \chi) \right| \\ &\leq \left(\sum_{1 \leq m \leq Q} \frac{1}{\varphi(m)} \right) \sum_{1 < q \leq Q, q_0 \nmid q} \left| \sum_{\chi \pmod{q}}^* \psi(x, \chi) \right|. \end{aligned} \quad (43)$$

Using Theorem 1.2 of [1], we have

$$\begin{aligned}
\sum_{Q_1 < q \leq Q} \frac{1}{\varphi(q)} \left| \sum_{\chi \pmod{q}}^* \psi(x, \chi) \right| &\leq c_0 \left(\frac{4x}{Q_1} + 4x^{\frac{1}{2}}Q + 18x^{\frac{2}{3}}Q^{\frac{1}{2}} + \frac{5}{2}x^{\frac{5}{6}} \log x \right) \log^{\frac{7}{2}} x \\
&\leq c_0 \left(\frac{4x}{Q_1} + \frac{4x}{\log^A x} + \frac{18x^{\frac{11}{12}}}{\log^{\frac{A}{2}} x} + \frac{5}{2}x^{\frac{5}{6}} \log x \right) \log^{\frac{7}{2}} x \\
&\leq \frac{4c_0 x}{Q_1} + \frac{c_0 x}{\log^{A-\frac{7}{2}} x} \left(4 + \frac{18 \log^{\frac{A}{2}} x}{x^{\frac{1}{12}}} + \frac{5 \log^{A+1} x}{2x^{\frac{1}{6}}} \right) \\
&< \frac{4c_0 x}{Q_1} + \frac{c_0(4 + e^{-800})x}{\log^{A_1 - \frac{7}{2}} x},
\end{aligned} \tag{44}$$

where

$$\begin{aligned}
c_0 &= \frac{2^{\frac{13}{2}}}{9\pi \log 2} \left(\frac{1}{3} + \frac{3}{2 \log 2} \right) \left(\frac{2 + \log(\log 2 / \log(4/3))}{\log 2} \right) \sqrt{\frac{\psi(113)}{113}} \\
&< 48.833.
\end{aligned} \tag{45}$$

Now we consider the sum $\sum_{\chi \pmod{q}}^* \psi(x, \chi)$ for moduli q with $q \leq Q_1$, $q_0 \nmid q$.

The proof of Theorem 3.6 of [8] shows that the left-hand side quantity $\frac{\varphi(q)}{x} \left| \psi(x; q, a) - \frac{x}{\varphi(q)} \right|$ in this theorem can be replaced by $-1 + x^{-1} \sum_{\chi \pmod{q}} |\psi(x, \chi)|$. This also applies to Theorem 1.1.

Now, provided that $x \geq x_0$, Theorem 1.1 gives

$$\left| -1 + \frac{1}{x} \sum_{\chi \pmod{q}} |\psi(x, \chi)| \right| < \frac{C_0}{\log^{A+1} x} + E_0 \frac{x^{1-\beta_0}}{\beta_0}, \tag{46}$$

where $E_0 = 1$ and β_0 denote the Siegel zero modulo q if it exists and $E_0 = 0$ otherwise.

We see that

$$\begin{aligned}
\sum_{\chi \pmod{q}}^* |\psi(x, \chi)| &\leq \left| -1 + \frac{1}{x} \sum_{\chi \pmod{q}} |\psi(x, \chi)| \right| + \left| \frac{\psi(x, \chi_0)}{x} - 1 \right| \\
&\leq \left| -1 + \frac{1}{x} \sum_{\chi \pmod{q}} |\psi(x, \chi)| \right| + \left| \frac{\psi(x)}{x} - 1 \right| + \frac{\log q}{x \log 2} \\
&\leq \left| -1 + \frac{1}{x} \sum_{\chi \pmod{q}} |\psi(x, \chi)| \right| + \frac{e^{-18}x}{\log^{A-3} x},
\end{aligned} \tag{47}$$

using the inequality (Theorem 11 of [16])

$$\left| \frac{\psi(x)}{x} - 1 \right| < \sqrt{\frac{8}{17\pi}} \left(\frac{\log x}{R_2} \right)^{\frac{1}{4}} \exp - \sqrt{\frac{\log x}{R_2}} \tag{48}$$

for $x \geq 101$, where $R_2 = 9.645908801$. Therefore (46) becomes

$$\sum_{\chi \pmod{q}}^* |\psi(x, \chi)| < \frac{(C_0 + e^{-18})x}{\log^{A-3} x} + E_0 \frac{x^{1-\beta_0}}{1-\beta_0}, \tag{49}$$

where $E_0 = 1$ and β_0 denote the Siegel zero modulo q if it exists and $E_0 = 0$ otherwise.

Since $q \leq Q_1$ is non-exceptional, the right-hand side of (49) is at most

$$\frac{(C_0 + e^{-18})x}{\log^{A-3} x} + \frac{x^{1-\frac{1}{2AR_1 \log \log x}}}{1 - \frac{1}{2AR_1 \log \log x}}. \tag{50}$$

We can easily confirm that, provided that $x \geq x_0$,

$$\frac{x^{1-\frac{1}{2AR_1 \log \log x}}}{1 - \frac{1}{2AR_1 \log \log x}} < e^{-50} \frac{x}{\log^{A-3} x} \tag{51}$$

and therefore

$$\sum_{\chi \pmod{q}}^* |\psi(x, \chi)| < \frac{(C_0 + e^{-17})x}{\log^{A-3} x}. \tag{52}$$

We see that $\sum_{1 \leq m \leq Q_1} \frac{1}{\varphi(m)} < c_1(1 + \log Q_1)$ from the argument in the proof of Theorem A. 17 in [10, p. 316]. Thus we obtain

$$\sum_{q \leq Q_1, q_0 \nmid q} \frac{1}{\varphi(q)} \sum_{\chi \pmod{q}}^* |\psi(x, \chi)| \leq \frac{c_1(C_0 + e^{-17})x(1 + A \log \log x)}{\log^{A-3} x}. \tag{53}$$

Observing that $\sum_{m \leq Q} \frac{1}{\varphi(m)} < c_1(1 + \log Q) < \frac{c_1}{2} \log x$ and substituting (44), (53) into (43), we see that the last sum of (42) is at most

$$\frac{c_1 c_0 (2 + e^{-800})x}{\log^{A-\frac{9}{2}} x} + \frac{2c_0 c_1 x \log^{\frac{9}{2}} x}{Q_1} + \frac{c_1^2 (C_0 + e^{-17})x(1 + A \log \log x)}{2 \log^{A-4} x}. \quad (54)$$

This proves Theorem 1.4.

References

- [1] Amir Akbary and Kyle Hambrook, A variant of the Bombieri-Vinogradov theorem with explicit constants and applications, *Math. Comp.* (to appear).
- [2] E. Bombieri, On the large sieve, *Mathematika* **12** (1965), 201–225.
- [3] Chen Jing-Run and Wang Tian-Ze, On distribution of primes in an arithmetical progression, *Sci. Sinica Ser. A* **33** (1990), 397–408.
- [4] P. Dusart, Estimates of $\theta(x; k, l)$ for large values of x , *Math. Comp.* **71** (2002), 1137–1166.
- [5] Ming-Chit Liu and Tianze Wang, Distribution of zeros of Dirichlet L -functions and an explicit formula for $\psi(t, \chi)$, *Acta Arith.* **102** (2002), 261–293.
- [6] Habiba Kadiri, An explicit zero-free region for the Dirichlet L-functions, <http://arxiv.org/abs/0510570>.
- [7] Kevin S. McCurley, Explicit zero-free regions for Dirichlet L -functions, *J. Number Theory* **19** (1984), 7–32.
- [8] Kevin S. McCurley, Explicit estimates for the error term in the prime number theorem for arithmetic progressions, *Math. Comp.* **42** (1985), 265–285.
- [9] Kevin S. McCurley, Explicit estimates for $\theta(x; 3, 1)$ and $\psi(x; 3, 1)$, *Math. Comp.* **42** (1985), 287–296.
- [10] Melvyn B. Nathanson, *Additive Number Theory: The Classical Bases*, GTM 164, Springer-Verlag, New York, 1996.
- [11] O. Ramaré and R. Rumely, Primes in arithmetic progressions, *Math. Comp.* **65** (1996), 397–425.

- [12] Barkley Rosser, Explicit bounds for some functions of prime numbers, *Amer. J. Math.* **63** (1941), 211–232.
- [13] J. Barkley Rosser and Lowell Schoenfeld, Approximate formulas for some functions of prime numbers, *Illinois J. Math.* **6** (1962), 64–94.
- [14] J. Barkley Rosser and Lowell Schoenfeld, Sharper Bounds for the Chebyshev Functions $\theta(x)$ and $\psi(x)$, *Math. Comp.* **29** (1975), 243–269.
- [15] R. Rumely, Numerical computations concerning the ERH, *Math. Comp.* **61** (1993), 415–440.
- [16] Lowell Schoenfeld, Sharper Bounds for the Chebyshev Functions $\theta(x)$ and $\psi(x)$ II, *Math. Comp.* **30** (1976), 337–360.
- [17] T. S. Trudgian, An improved upper bound for the error in the zero-counting formulae for Dirichlet L -functions and Dedekind zeta-functions, (submitted).
- [18] Tomohiro Yamada, An explicit formula for the linear sieve, (in preparation).
- [19] Tomohiro Yamada, Explicit Chen’s theorem, (in preparation).

Tomohiro Yamada
Center for Japanese language and culture
Osaka University
562-8558
8-1-1, Aomatanihigashi, Minoo, Osaka
Japan
e-mail: tyamada1093@gmail.com

Table 5: constants related to Theorem 1

α_1	α	Y_0	C	α_1	α	Y_0	C
1	1	68	341.37279	1	7	98	224.55734
1	1	69	139.24784	1	7	99	0.2773700
1	1	70	51.198745	1	7	100	0.0002301
1	1	71	16.805224	1	7	101	0.0000569
1	1	72	4.9039512	1	8	101	8.1198690
1	1	73	1.2429184	1	8	102	0.0024266
1	1	74	0.2720959	1	8	103	0.0000469
1	1	75	0.0511499	1	9	103	70.055386
1	1	76	0.0085318	1	9	104	0.0085547
1	1	77	0.0016318	1	9	105	0.0000385
1	1	78	0.0006712	1	10	105	145.53327
1	1	79	0.0005215	1	10	106	0.0062008
1	2	77	287.04559	1	10	107	0.0000313
1	2	78	49.708220	1	10	108	0.0000001
1	2	79	7.0904806	2	1	77	366.59584
1	2	80	0.8182991	2	1	78	63.466310
1	2	81	0.0751978	2	1	79	9.0503268
1	2	82	0.0056819	2	1	80	1.0500668
1	2	83	0.0006284	2	1	81	0.0967331
1	2	84	0.0003224	2	1	82	0.0072533
1	2	85	0.0002818	2	1	83	0.0007180
1	3	84	27.074957	2	1	84	0.0003264
1	3	85	1.4777161	2	1	85	0.0002820
1	3	86	0.0591888	2	2	84	33.655774
1	3	87	0.0018985	2	2	85	1.8367608
1	3	88	0.0002410	2	2	86	0.0735210
1	3	89	0.0001891	2	2	87	0.0023052
1	4	88	169.98979	2	2	88	0.0002489
1	4	89	4.3948140	2	2	89	0.0001892
1	4	90	0.0767410	2	3	88	206.74646
1	4	91	0.0010204	2	3	89	5.3442833
1	4	92	0.0001459	2	3	90	0.0932719
1	5	92	94.371270	2	3	91	0.0012073
1	5	93	0.8513896	2	3	92	0.0001472
1	5	94	0.0047470	2	4	92	112.75512
1	5	95	0.0001181	2	4	93	1.0170960
1	5	96	0.0000938	2	4	94	0.0056481
1	6	95	411.32655	2	4	95	0.0001209
1	6	96	1.5600920	2	4	96	0.0000938
1	6	97	0.0033364				
1	6	98	0.0000803				

α_1	α	Y_0	C	α_1	α	Y_0	C
2	5	95	484.26872	3	4	95	557.01746
2	5	96	1.8365549	3	4	96	2.1123032
2	5	97	0.0039123	3	4	97	0.0044867
2	5	98	0.0000809	3	4	98	0.0000815
2	6	98	261.09757	3	5	98	297.55254
2	6	99	0.3224670	3	5	99	0.3674601
2	6	100	0.0002573	3	5	100	0.0002844
2	6	101	0.0000569	3	5	101	0.0000569
2	7	101	9.3404567	3	6	101	10.558512
2	7	102	0.0027835	3	6	102	0.0031396
2	7	103	0.0000469	3	6	103	0.0000470
2	8	103	79.844778	3	7	103	89.615782
2	8	104	0.0097436	3	7	104	0.0109304
2	8	105	0.0000386	3	7	105	0.0000386
2	9	105	164.53130	3	8	105	183.49683
2	9	106	0.0070055	3	8	106	0.0078087
2	9	107	0.0000313	3	8	107	0.0000313
2	10	107	63.493001	3	9	107	70.410044
2	10	108	0.0007963	3	9	108	0.0008799
2	10	109	0.0000256	3	9	109	0.0000256
2	10	110	0.0000001	3	10	109	3.7235338
3	1	84	40.187937	3	10	110	0.0000334
3	1	85	2.1926074	3	10	111	0.0000003
3	1	86	0.0890408	3	10	112	0.0000001
3	1	87	0.0028106	4	1	88	279.89541
3	1	88	0.0002627	4	1	89	7.2338934
3	1	89	0.0001896	4	1	90	0.1298417
3	2	88	243.40542	4	1	91	0.0017720
3	2	89	6.2933477	4	1	92	0.0001558
3	2	90	0.1099001	4	2	92	149.37762
3	2	91	0.0013993	4	2	93	1.3500825
3	2	92	0.0001487	4	2	94	0.0075153
3	3	92	131.08327	4	2	95	0.0001277
3	3	93	1.1823671	4	2	96	0.0000938
3	3	94	0.0065480	4	3	95	629.69393
3	3	95	0.0001237	4	3	96	2.3881358
3	3	96	0.0000938	4	3	97	0.0050654
				4	3	98	0.0000822

α_1	α	Y_0	C	α_1	α	Y_0	C
4	4	98	333.97042	5	4	101	12.992382
4	4	99	0.4124195	5	4	102	0.0038532
4	4	100	0.0003115	5	4	103	0.0000470
4	4	101	0.0000569	5	5	103	109.13817
4	5	101	11.775583	5	5	104	0.0133020
4	5	102	0.0034955	5	5	105	0.0000388
4	5	103	0.0000470	5	6	105	221.39416
4	6	103	99.380273	5	6	106	0.0094139
4	6	104	0.0121164	5	6	107	0.0000314
4	6	105	0.0000387	5	7	107	84.233287
4	7	105	202.45206	5	7	108	0.0010471
4	7	106	0.0086116	5	7	109	0.0000256
4	7	107	0.0000314	5	8	109	4.4201216
4	8	107	77.323788	5	8	110	0.0000353
4	8	108	0.0009635	5	9	110	11419.253
4	8	109	0.0000256	5	9	111	0.0221440
4	9	109	4.0719188	5	9	112	0.0000190
4	9	110	0.0000343	5	10	112	22.759465
4	10	110	10553.818	5	10	113	0.0000240
4	10	111	0.0204677	5	10	114	0.0000001
4	10	112	0.0000190	6	1	95	774.84324
4	10	113	0.0000001	6	1	96	2.9379648
5	1	92	167.66332	6	1	97	0.0062067
5	1	93	1.5120638	6	1	98	0.0000955
5	1	94	0.0100915	6	2	98	406.74243
5	1	95	0.0001509	6	2	99	0.5177958
5	1	96	0.0000939	6	2	100	0.0005074
5	2	95	702.27252	6	2	101	0.0000572
5	2	96	2.6767555	6	3	101	14.230994
5	2	97	0.0058542	6	3	102	0.0043119
5	2	98	0.0000850	6	3	103	0.0000473
5	3	98	370.41678	6	4	103	118.90823
5	3	99	0.4578772	6	4	104	0.0145133
5	3	100	0.0003417	6	4	105	0.0000389
5	3	101	0.0000570				

α_1	α	Y_0	C	α_1	α	Y_0	C
6	5	105	240.33821	7	7	110	13150.105
6	5	106	0.0102188	7	7	111	0.0254970
6	5	107	0.0000314	7	7	112	0.0000190
6	6	107	91.142884	7	8	112	26.082288
6	6	108	0.0011307	7	8	113	0.0000250
6	6	109	0.0000256	7	9	113	15059.773
6	7	109	4.7683260	7	9	114	0.0026411
6	7	110	0.0000362	7	10	115	0.4569526
6	8	110	12284.687	7	10	116	0.0000127
6	8	111	0.0238203	7	10	117	0.0000001
6	8	112	0.0000190	8	1	101	16.639402
6	9	112	24.420945	8	1	102	0.0049176
6	9	113	0.0000245	8	1	103	0.0000478
6	10	113	14130.221	8	2	103	138.40436
6	10	114	0.0024791	8	2	104	0.0168563
6	10	115	0.0000044	8	2	105	0.0000514
6	10	116	0.0000001	8	3	105	278.20999
7	1	98	443.12389	8	3	106	0.0176009
7	1	99	0.5471351	8	3	107	0.0000329
7	1	100	0.0003927	8	4	107	105.25995
7	1	101	0.0000574	8	4	108	0.0015069
7	2	101	15.423819	8	4	109	0.0000256
7	2	102	0.0082841	8	5	109	5.4707339
7	2	103	0.0000503	8	5	110	0.0000394
7	3	103	129.05908	8	6	110	14015.192
7	3	104	0.0169840	8	6	111	0.0271902
7	3	105	0.0000402	8	6	112	0.0000190
7	4	105	259.38714	8	7	112	27.743089
7	4	106	0.0111522	8	7	113	0.0000255
7	4	107	0.0000315	8	8	113	15988.974
7	5	107	98.059267	8	8	114	0.0028032
7	5	108	0.0012172	8	9	115	0.4842973
7	5	109	0.0000256	8	9	116	0.0000127
7	6	109	5.1166158	8	10	116	42.515883
7	6	110	0.0000372	8	10	117	0.0000119
				8	10	118	0.0000001

α_1	α	Y_0	C
9	1	103	148.15370
9	1	104	0.0180405
9	1	105	0.0000390
9	2	105	297.13835
9	2	106	0.0126220
9	2	107	0.0000332
9	3	107	111.86299
9	3	108	0.0043396
9	3	109	0.0000263
9	4	109	6.0873692
9	4	110	0.0000896
9	5	110	14886.409
9	5	111	0.0302231
9	5	112	0.0000190
9	6	112	29.418754
9	6	113	0.0000267
9	7	113	16918.289
9	7	114	0.0029683
9	8	115	0.5116813
9	8	116	0.0000127
9	9	116	44.850504
9	9	117	0.0000119
9	10	117	1923.3233
9	10	118	0.0000167
9	10	119	0.0000001
α_1	α	Y_0	C
10	1	105	316.06672
10	1	106	0.0134237
10	1	107	0.0000314
10	2	107	118.76799
10	2	108	0.0014647
10	2	109	0.0000257
10	3	109	6.1605138
10	3	110	0.0002801
10	4	110	15744.953
10	4	111	0.0946919
10	4	112	0.0000209
10	5	112	31.907392
10	5	113	0.0000663
10	6	113	17854.358
10	6	114	0.0034270
10	7	115	0.5399268
10	7	116	0.0000128
10	8	116	47.186782
10	8	117	0.0000120
10	9	117	2020.8916
10	9	118	0.0000171
10	10	118	40634.957
10	10	119	0.0000610
10	10	120	0.0000001