

AN APPLICATION OF COHN'S RULE TO CONVOLUTIONS
OF UNIVALENT HARMONIC MAPPINGS

Raj Kumar^{a *}, Sushma Gupta^a, Sukhjit Singh^a and Michael Dorff^{b †}

^a *Sant Longowal Institute of Engineering and Technology, Longowal-148106 (Punjab), India.*

^b *Department of Mathematics, Brigham Young University, Provo, Utah, 84602, USA.*

Abstract

Dorff et al. [4], proved that the harmonic convolution of right half-plane mapping with dilatation $-z$ and mapping $f_\beta = h_\beta + \bar{g}_\beta$, where f_β is obtained by shearing of analytic vertical strip mapping, with dilatation $e^{i\theta}z^n$, $n = 1, 2$, $\theta \in \mathbb{R}$, is in S_H^0 and is convex in the direction of the real axis. In this paper, by using Cohn's rule, we generalize this result by considering dilatations $(a - z)/(1 - az)$ and $e^{i\theta}z^n$ ($n \in \mathbb{N}$, $\theta \in \mathbb{R}$) of right half-plane mapping and f_β , respectively.

Key Words : Cohn's rule, univalent harmonic mappings, harmonic convolutions.

AMS Subject Classification: 30C45.

1 Introduction

Let S_H denotes the class of all harmonic, sense-preserving and univalent mappings $f = h + \bar{g}$ in the unit disk $E = \{z : |z| < 1\}$, which are normalized by the conditions $f(0) = 0$ and $f_z(0) = 1$. Therefore a function $f = h + \bar{g}$ in the class S_H has the representation,

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} \bar{b}_n \bar{z}^n, \quad (1)$$

for all z in E . Denote by S_H^0 the subclass of S_H whose functions satisfy an additional condition of normalization $f_{\bar{z}}(0) = 0$. Lewy's Theorem asserts that a harmonic mapping $f = h + \bar{g}$ defined in E , is locally univalent and sense-preserving if and only if the Jacobian of the mapping, defined by $J_f = |h'|^2 - |g'|^2$, is positive or equivalently, if and only if $h' \neq 0$ in E and the dilatation function ω of f , defined by $\omega = g'/h'$, satisfies $|\omega(z)| < 1$, for all $z \in E$. Further, let $K_H(K_H^0)$ be

*rajgarg2012@yahoo.co.in

†mdorff@math.byu.edu

the subclass of $S_H(S_H^0)$ consisting of functions which map the unit disk E onto convex domains. A domain Ω is said to be convex in a direction ϕ , $0 \leq \phi < \pi$, if every line parallel to the line joining 0 and $e^{i\phi}$ has a connected intersection with Ω .

Kumar et al. [6], defined the harmonic mappings in the right half-plane $F_a = H_a + \overline{G}_a \in K_H$, given by $H_a(z) + G_a(z) = z/(1-z)$ with dilatation $\omega_a(z) = (a-z)/(1-az)$, $a \in (-1, 1)$. Then, by using the shearing technique due to Clunie and Sheil-Small [1], we get,

$$H_a(z) = \frac{\frac{1}{1+a}z - \frac{1}{2}z^2}{(1-z)^2} \quad \text{and} \quad G_a(z) = \frac{\frac{a}{1+a}z - \frac{1}{2}z^2}{(1-z)^2}. \quad (2)$$

Let $f_\beta = h_\beta + \overline{g}_\beta$ be the collection of harmonic mappings obtained by shearing of analytic vertical strip mapping

$$h_\beta(z) + g_\beta(z) = \frac{1}{2i \sin \beta} \log \left(\frac{1 + ze^{i\beta}}{1 + ze^{-i\beta}} \right), \quad 0 < \beta < \pi. \quad (3)$$

For more details about this map see Dorff [3].

Harmonic Convolution (Hadamard product) of two harmonic functions $F(z) = H(z) + \overline{G}(z) = z + \sum_{n=2}^{\infty} A_n z^n + \sum_{n=1}^{\infty} \overline{B}_n \overline{z}^n$ and $f(z) = h(z) + \overline{g}(z) = z + \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} \overline{b}_n \overline{z}^n$ in S_H is defined as,

$$\begin{aligned} (F * f)(z) &= (H * h)(z) + \overline{(G * g)}(z) \\ &= z + \sum_{n=2}^{\infty} a_n A_n z^n + \sum_{n=1}^{\infty} \overline{b}_n \overline{B}_n \overline{z}^n. \end{aligned}$$

It is well known that unlike the case of conformal mappings, harmonic convolution of two mappings from the class K_H^0 may not be in K_H^0 , may not be even in S_H^0 . Also, if $f_1 \in K_H^0$ and $f_2 \in S_H^0$, then $f_1 * f_2$ is not necessarily in S_H^0 . These facts generated a lot of interest in the study of harmonic convolutions of univalent harmonic mappings and a good number of papers recently appeared in the literature on this topic (for example see [2, 4, 5, 6, 7, 8]). In particular, Dorff [2] proved the following result:

Theorem A. *Let $f = h + \overline{g}$ with $h(z) + g(z) = z/(1-z)$ be the right half-plane mapping and $f_\beta = h_\beta + \overline{g}_\beta \in K_H^0$ be given by (3) with $\pi/2 \leq \beta < \pi$. Then $f * f_\beta \in S_H^0$ and is convex in the direction of the real axis, provided $f * f_\beta$ is locally univalent and sense-preserving.*

Recently, Dorff et al. [4] obtained the following result which shows that the condition of convolution being locally univalent and sense-preserving as required in Theorem A, can be dropped in some special cases but not in general.

Theorem B. *If $f = h + \overline{g}$, where $h(z) + g(z) = z/(1-z)$ is a right half-plane mapping with dilatation $\omega(z) = -z$ and $f_\beta = h_\beta + \overline{g}_\beta$ is the harmonic map as defined in Theorem A above with dilatation $\omega_n(z) = e^{i\theta} z^n$; then for $n = 1, 2$, $f * f_\beta \in S_H^0$ and is convex in the direction of the real axis.*

Proceeding as in Remark 1 in [4] one can verify that Theorem B does not hold true for $n \geq 3$. Aim of the present paper is to study convolutions of more general harmonic mappings $F_a = H_a + \overline{G}_a$, where H_a and G_a are defined by (2) and the mappings $f_\beta = h_\beta + \overline{g}_\beta$, where h_β and g_β are the solutions of (3), with dilatation $\omega_n(z) = g'_\beta(z)/h'_\beta(z) = e^{i\theta}z^n$, $n \in \mathbb{N}$. We shall find the range of real constant a such that $F_a * f_\beta \in S_H^0$ and is convex in the direction of the real axis even if $n \geq 3$. Cohn's rule stated below shall play a central role in the proofs of our results in this paper.

Lemma A. (Cohn's Rule [9, p.375]) *Given a polynomial*

$$t(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$$

of degree n , let

$$t^*(z) = z^n \overline{\left(\frac{1}{z}\right)} = \overline{a_n} + \overline{a_{n-1}}z + \overline{a_{n-2}}z^2 + \dots + \overline{a_0}z^n.$$

Denote by r and s the number of zeros of t inside and on the unit circle $|z| = 1$, respectively. If $|a_0| < |a_n|$, then

$$t_1(z) = \frac{\overline{a_n}t(z) - a_0t^*(z)}{z}$$

is of degree $n - 1$ and has $r_1 = r - 1$ and $s_1 = s$ number of zeros inside and on the unit circle $|z| = 1$, respectively.

2 Main results

In addition to Cohn's rule stated above, we begin by proving following two results which are also essential for the proofs of our main results.

Lemma 2.1. *Let $f_\beta = h_\beta + \overline{g}_\beta \in K_H^0$ be given by (3) with dilatation $\omega = g'_\beta/h'_\beta$ and $F_a = H_a + \overline{G}_a$ be a mapping in the right half-plane defined by (2). Then $\tilde{\omega}$, the dilatation of $F_a * f_\beta$, is given by*

$$\tilde{\omega}(z) = \left[\frac{2\omega(1+\omega)(a + az\cos\beta + z\cos\beta + z^2) - z\omega'(1-a)(1 + 2z\cos\beta + z^2)}{2(1 + z\cos\beta + az\cos\beta + az^2)(1+\omega) - z\omega'(1-a)(1 + 2z\cos\beta + z^2)} \right]. \quad (4)$$

Proof. Since

$$F_a * f_\beta = H_a * h_\beta + \overline{G_a * g_\beta} = H + \overline{G} \text{ (say),}$$

so

$$H(z) = \frac{1}{2} \left[h_\beta + \frac{(1-a)}{(1+a)} zh'_\beta \right]$$

and

$$G(z) = \frac{1}{2} \left[g_\beta - \frac{(1-a)}{(1+a)} zg'_\beta \right].$$

If $\tilde{\omega}$ is the dilatation of $F_a * f_\beta$, then

$$\tilde{\omega}(z) = \frac{G'(z)}{H'(z)} = \left[\frac{2ag'_\beta - (1-a)zg''_\beta}{2h'_\beta + (1-a)zh''_\beta} \right]. \quad (5)$$

Since

$$g'_\beta = \omega h'_\beta, \quad h'_\beta(z) = \frac{1}{(1+\omega)(1+ze^{i\beta})(1+ze^{-i\beta})} \quad \text{and}$$

$$h''_\beta(z) = -\frac{2(\cos\beta + z)(1+\omega) + \omega'(1+2z\cos\beta + z^2)}{(1+\omega)^2(1+ze^{i\beta})^2(1+ze^{-i\beta})^2},$$

therefore, from (5) we get

$$\tilde{\omega}(z) = \left[\frac{2a\omega h'_\beta - (1-a)z(\omega h''_\beta + \omega' h'_\beta)}{2h'_\beta + (1-a)zh''_\beta} \right].$$

Substituting the values of h'_β and h''_β and simplifying, we get (4). \square

Lemma 2.2. *Following inequalities hold true:*

$$(a) \left| -2\cos\beta + 4e^{-i\theta}\cos^2\beta - 3e^{-i\theta} + e^{i\theta} \right| \leq \left| 4 - 2\cos^2\beta - 2\cos\beta\cos\theta \right| \text{ for } \beta \in (0, \pi) \text{ and } \theta \in \mathbb{R}.$$

$$(b) \left| \cos\beta(e^{-i\theta} - 5) \right| < \left| 6 - \cos^2\beta + 2e^{-i\theta} - 3e^{-i\theta}\cos^2\beta \right| \text{ for } \beta \in (0, \pi) \text{ and } \theta \in \mathbb{R}.$$

$$(c) \left| 2(1 + e^{-i\theta}) - 3e^{-i\theta}\cos^2\beta \right| < \left| 4 - \cos^2\beta \right| \text{ for } \beta \in (0, \pi/2) \cup (\pi/2, \pi) \text{ and } \theta \neq 2n\pi, n \in \mathbb{Z}.$$

Proof. Let $x = \cos\beta$ and $y = \cos\theta$. Then

$$\begin{aligned} (a) & \left| -2\cos\beta + 4e^{-i\theta}\cos^2\beta - 3e^{-i\theta} + e^{i\theta} \right|^2 - \left| 4 - 2\cos^2\beta - 2\cos\beta\cos\theta \right|^2 \\ &= 12x^4 - 12x^2 - 24x^3y + 24xy + 12x^2y^2 - 12y^2 \\ &= -12(1-x^2)(x-y)^2 \\ &\leq 0, \text{ as } x \in (-1, 1) \text{ and } y \in [-1, 1]. \end{aligned}$$

$$\begin{aligned} (b) & \left| \cos\beta(e^{-i\theta} - 5) \right|^2 - \left| 6 - \cos^2\beta + 2e^{-i\theta} - 3e^{-i\theta}\cos^2\beta \right|^2 \\ &= -2(5x^4 + 3x^4y - 15x^2y - 25x^2 + 12y + 20) \\ &= -2(x^2 - 4)(x^2 - 1)(5 + 3y) \\ &< 0, \text{ for } x \in (-1, 1) \text{ and } y \in [-1, 1]. \end{aligned}$$

$$\begin{aligned} (c) & \left| 2(1 + e^{-i\theta}) - 3e^{-i\theta}\cos^2\beta \right|^2 - \left| 4 - \cos^2\beta \right|^2 \\ &= 4(2x^4 - x^2 - 3yx^2 + 2y - 2) \\ &= 4[2(x^2 - 1)(x^2 + 1 - y) - x^2(y + 1)] \\ &< 0, \text{ for } x \in (-1, 0) \cup (0, 1) \text{ and } y \in [-1, 1]. \end{aligned} \quad \square$$

We are now in a position to prove our main results.

Theorem 2.3. *Let $f_\beta = h_\beta + \bar{g}_\beta \in K_H^0$ be a mapping given by (3) with dilatation $\omega_1(z) = e^{i\theta}z$. If F_a defined by (2) is a mapping in the right half-plane, then $F_a * f_\beta \in S_H^0$ and is convex in the direction of the real axis for $a \in [-1/3, 1)$.*

Proof. Let $\tilde{\omega}$ be the dilatation of $F_a * f_\beta$. We claim that $|\tilde{\omega}(z)| < 1$ for all $z \in E$ i.e., $F_a * f_\beta$ is locally univalent and sense-preserving. Our result shall then follow from Theorem A. By setting $\omega(z) = \omega_1(z) = e^{i\theta}z$ in (4) we get,

$$\begin{aligned}\tilde{\omega}(z) &= z \left[\frac{e^{i\theta}z^3 + (\frac{1}{2} + ae^{i\theta} \cos \beta + e^{i\theta} \cos \beta + \frac{a}{2})z^2 + a(2 \cos \beta + e^{i\theta})z + \frac{(3a-1)}{2}}{e^{-i\theta} + (e^{-i\theta} \cos \beta + ae^{-i\theta} \cos \beta + \frac{1}{2} + \frac{a}{2})z + a(e^{-i\theta} + 2 \cos \beta)z^2 + \frac{(3a-1)}{2}z^3} \right] \\ &= z \frac{p(z)}{p^*(z)},\end{aligned}$$

where,

$$\begin{aligned}p(z) &= e^{i\theta}z^3 + (\frac{1}{2} + ae^{i\theta} \cos \beta + e^{i\theta} \cos \beta + \frac{a}{2})z^2 + a(2 \cos \beta + e^{i\theta})z + \frac{(3a-1)}{2} \\ &= a_3z^3 + a_2z^2 + a_1z + a_0\end{aligned}$$

and

$$p^*(z) = e^{-i\theta} + (e^{-i\theta} \cos \beta + ae^{-i\theta} \cos \beta + \frac{1}{2} + \frac{a}{2})z + a(e^{-i\theta} + 2 \cos \beta)z^2 + \frac{(3a-1)}{2}z^3 = z^3 \overline{p(\frac{1}{z})}.$$

Thus if $z_0, z_0 \neq 0$, is a zero of p then $1/\bar{z}_0$ is a zero of p^* . Therefore we can write

$$\tilde{\omega}(z) = z \frac{(z+A)(z+B)(z+C)}{(1+\bar{A}z)(1+\bar{B}z)(1+\bar{C}z)}.$$

In order to prove that $|\tilde{\omega}(z)| < 1$, it suffices to show that $|A| \leq 1$, $|B| \leq 1$ and $|C| \leq 1$ or equivalently, all the zeros, $-A, -B$ and $-C$, of the polynomial p lie in or on the unit circle $|z| = 1$ for $a \in [-1/3, 1)$. When $a = -1/3$, then $|\tilde{\omega}(z)| = |-ze^{i\theta}| < 1$ for all $z \in E$, so we take $a \in (-1/3, 1/3) \cup (1/3, 1)$ and the case when $a = 1/3$ will be settled separately.

Let

$$\begin{aligned}p_1(z) &= \frac{\bar{a}_3p(z) - a_0p^*(z)}{z} \\ &= \frac{1}{4}(1 + 2a - 3a^2)[3z^2 + 2(2 \cos \beta + e^{-i\theta})z + (2 \cos \beta e^{-i\theta} + 1)] \\ &= b_2z^2 + b_1z + b_0.\end{aligned}\tag{6}$$

As $|a_0| < |a_3|$ and $1 + 2a - 3a^2 \neq 0$ for $a \in (-1/3, 1/3) \cup (1/3, 1)$, therefore by Cohn's rule polynomial p_1 has one less number of zeros inside $|z| = 1$ than p and same number of zeros on $|z| = 1$ as p . Again, let

$$p_2(z) = \frac{\bar{b}_2p_1(z) - b_0p_1^*(z)}{z}$$

$$= \frac{1}{16}(1 + 2a - 3a^2)^2[(9 - |2 \cos \beta + e^{i\theta}|^2)z + 6(2 \cos \beta + e^{-i\theta}) - 2e^{-i\theta}(2 \cos \beta + e^{i\theta})^2],$$

where $p_1^*(z) = z^2 \overline{p_1\left(\frac{1}{z}\right)}$. We can again apply Cohn's rule on p_1 because

$|b_0| = |(1 + 2a - 3a^2)\frac{1}{4}(2 \cos \beta e^{-i\theta} + 1)| < |(1 + 2a - 3a^2)\frac{3}{4}| = |b_2|$. If the zero of p_2 is denoted by z_0 , then

$$z_0 = \frac{-2 \cos \beta + 4e^{-i\theta} \cos^2 \beta - 3e^{-i\theta} + e^{i\theta}}{4 - 2 \cos^2 \beta - 2 \cos \beta \cos \theta}$$

and in view of Lemma 2.2 (a), $|z_0| \leq 1$. Thus we have proved that all the zeros of p_1 and therefore of p lie inside or on the unit circle $|z| = 1$ for $a \in (-1/3, 1/3) \cup (1/3, 1)$. In case when $a = 1/3$, then

$$p(z) = \frac{1}{3}e^{i\theta}z \left[3z^2 + 2(2 \cos \beta + e^{-i\theta})z + (2 \cos \beta e^{-i\theta} + 1) \right].$$

In view of (6) it can be easily verified that all the zeros of p lie inside or on the circle $|z| = 1$ for $a = 1/3$. This completes the proof. \square

Remark 2.4. By setting $a = 0$ in Theorem 2.3, we get Theorem B (for $n = 1$), stated in Section 1.

Theorem 2.5. *If $f_\beta \in K_H^0$ is a mapping with dilatation $\omega_2(z) = e^{i\theta}z^2$, then $F_a * f_\beta \in S_H^0$ and is convex in the direction of the real axis for $a \in [0, 1)$, where F_a is defined by (2).*

Proof. The case when $a = 0$ has already been proved by Dorff et al. [4], so we take $a \in (0, 1)$. Now, if $\tilde{\omega}$ is the dilatation of $F_a * f_\beta$, then letting $\omega(z) = \omega_2(z) = e^{i\theta}z^2$ in (4) we get,

$$\tilde{\omega}(z) = z^2 e^{2i\theta} \left[\frac{z^4 + \cos \beta(a+1)z^3 + a(1 + e^{-i\theta})z^2 + e^{-i\theta} \cos \beta(3a-1)z + e^{-i\theta}(2a-1)}{1 + \cos \beta(a+1)z + a(1 + e^{i\theta})z^2 + e^{i\theta} \cos \beta(3a-1)z^3 + e^{i\theta}(2a-1)z^4} \right].$$

In view of Theorem A, we need only to show that $F_a * f_\beta$ is locally univalent and sense-preserving, i.e., $|\tilde{\omega}| < 1$ in E . Consider

$$\begin{aligned} q(z) &= z^4 + \cos \beta(a+1)z^3 + a(1 + e^{-i\theta})z^2 + e^{-i\theta} \cos \beta(3a-1)z + e^{-i\theta}(2a-1) \\ &= a_4 z^4 + a_3 z^3 + a_2 z^2 + a_1 z + a_0, \end{aligned}$$

and

$$\begin{aligned} q^*(z) &= z^4 \overline{q\left(\frac{1}{z}\right)} \\ &= 1 + \cos \beta(a+1)z + a(1 + e^{i\theta})z^2 + e^{i\theta} \cos \beta(3a-1)z^3 + e^{i\theta}(2a-1)z^4. \end{aligned}$$

$$\text{Then } \tilde{\omega}(z) = z^2 e^{2i\theta} \frac{q(z)}{q^*(z)} = z^2 e^{2i\theta} \frac{(z+A)(z+B)(z+C)(z+D)}{(1+\overline{A}z)(1+\overline{B}z)(1+\overline{C}z)(1+\overline{D}z)},$$

where $-A, -B, -C$ and $-D$ are the zeros of q . We shall show that these zeros lie inside or on the unit circle $|z| = 1$ for $a \in (0, 1)$. First we take $a \in (0, 1/2) \cup (1/2, 1)$ and the case when $a = 1/2$ will be dealt separately. For $a \in (0, 1/2) \cup (1/2, 1)$, we have $|a_0| = |2a - 1| < 1 = |a_4|$,

so we can apply Cohn's rule on q . Let

$$\begin{aligned} q_1(z) &= \frac{\bar{a}_4 q(z) - a_0 q^*(z)}{z} \\ &= 2a(1-a) \left(2z^3 + 3 \cos \beta z^2 + (1 + e^{-i\theta})z + e^{-i\theta} \cos \beta \right). \\ &= b_3 z^3 + b_2 z^2 + b_1 z + b_0. \end{aligned} \tag{7}$$

It is easy to verify that Cohn's rule is applicable to q_1 also. So, let

$$\begin{aligned} q_2(z) &= \frac{\bar{b}_3 q_1(z) - b_0 q_1^*(z)}{z} \\ &= 4(a(1-a))^2 \left[(4 - \cos^2 \beta)z^2 + \cos \beta(5 - e^{-i\theta})z + 2(1 + e^{-i\theta}) - 3e^{-i\theta} \cos^2 \beta \right] \\ &= c_2 z^2 + c_1 z + c_0. \end{aligned}$$

To apply Cohn's rule on q_2 we need $|2(1 + e^{-i\theta}) - 3e^{-i\theta} \cos^2 \beta| < |4 - \cos^2 \beta|$, which is true in view of Lemma 2.2 (c) provided $\beta \neq \pi/2$ and $\theta \neq 2n\pi, n \in \mathbb{Z}$. Therefore by applying Lemma A again on q_2 , we have

$$\begin{aligned} q_3(z) &= \frac{\bar{c}_2 q_2(z) - c_0 q_2^*(z)}{z} \\ &= 16(a(1-a))^4 \{ [(4 - \cos^2 \beta)^2 - (2(1 + e^{-i\theta}) - 3e^{-i\theta} \cos^2 \beta)^2] z + \\ &\quad \cos \beta(5 - e^{-i\theta}) [(4 - \cos^2 \beta) - 2(1 + e^{-i\theta}) + 3e^{-i\theta} \cos^2 \beta] \}. \end{aligned}$$

The only zero of q_3 shall lie inside or on the circle $|z| = 1$, provided

$$|\cos \beta(e^{-i\theta} - 5)| < |6 - \cos^2 \beta + 2e^{-i\theta} - 3e^{-i\theta} \cos^2 \beta|,$$

which is true in view of Lemma 2.2(b). Thus by Cohn's rule all zeros of q_2 , q_1 and therefore of q lie in or on the unit circle $|z| = 1$ for $\beta \neq \pi/2$ and $\theta \neq 2n\pi, n \in \mathbb{Z}$.

In case $\beta = \pi/2$ and $\theta = 2n\pi, n \in \mathbb{Z}$, we have $q_2(z) = 16(a(1-a))^2(z^2 + 1)$ and obviously, zeros of q_2 lie on $|z| = 1$ because $a \in (0, 1/2) \cup (1/2, 1)$. Therefoer by Cohn's rule all zeros of q_1 and of q lie in or on the unit circle $|z| = 1$ in this case also.

Now when $a = 1/2$, then we have

$$q(z) = \frac{z}{2} \left(2z^3 + 3 \cos \beta z^2 + (1 + e^{-i\theta})z + e^{-i\theta} \cos \beta \right).$$

Keeping (7) in view, we can easily verify that in this case all zeros of q lie in or on the unit circle $|z| = 1$. Hence $|\tilde{\omega}| < 1$ for $a \in [0, 1)$. \square

As pointed out in Section 1, Theorem B is not true for $n \geq 3$, however following theorem, which we state without proof (as proof runs on the same lines as of Theorem 2.5), asserts that the conclusion of Theorem 2.5 remains valid when dilatation of f_β is taken as $e^{i\theta} z^3$ and value of real constant a is restricted in the interval $[1/5, 1)$.

Theorem 2.6. Let $f_\beta \in K_H^0$ be a harmonic map with dilatation $\omega_3(z) = e^{i\theta} z^3$. Then $F_a * f_\beta \in S_H^0$ and is convex in the direction of the real axis for $a \in [1/5, 1)$, where F_a is defined by (2).

Remark 2.7. It is expected that, keeping $\omega_n(z) = e^{i\theta} z^n$ ($\theta \in \mathbb{R}, n \in \mathbb{N}$) as the dilatation of f_β , the convolution $F_a * f_\beta \in S_H^0$ and is convex in the direction of the real axis for $a \in [\frac{n-2}{n+2}, 1)$. The authors have verified the case when $n = 4$ but when n increases calculations become extremely cumbersome.

Acknowledgement: The first author is thankful to the Council of Scientific and Industrial Research, New Delhi, for financial support vide grant no. 09/797/0006/2010 EMR-1.

References

- [1] J. Clunie and T. Sheil-Small, *Harmonic univalent functions*, Ann. Acad. Sci. Fenn. Ser. A. I Math., **9** (1984), 3-25.
- [2] M. Dorff, *Convolutions of planar harmonic convex mappings*, Comp. Var. Theory Appl. **45**(3) (2001), 263-271.
- [3] M. Dorff, *Harmonic mappings onto asymmetric vertical strips*, Computational Methods and Function Theory 1997, (N.Papamichael, St. Ruscheweyh and E.B.Saff, eds), 171-175, World Sci. Publishing, River Edge, NJ,1999.
- [4] M. Dorff, M. Nowak and M. Woloszkiwicz, *Convolutions of harmonic convex mappings*, Comp. Var. Elliptic Eqn., **57**(5) (2012), 489-503.
- [5] M.Goodloe, *Hadamard Products of Harmonic Mappings*, Complex Var. Theory Appl., **47**(2) (2002), 81-92.
- [6] R. Kumar, M. Dorff, S.Gupta and S.Singh, *Convolution properties of some harmonic mappings in the right half-plane*, see arXiv:1206.4364.
- [7] L. Li and S. Ponnusamy, *Solution to an open problem on convolutions of harmonic mappings*, Comp. Var. Elliptic Eqn. (2012), accepted.
- [8] L. Li and S. Ponnusamy, *Convolution of slanted right half-plane harmonic mappings*, Analysis (2012) accepted (or see arXiv:1206.4364).
- [9] Q.I. Rahman and G. Schmeisser, *Analytic Theory of Polynomials*, London Mathematical Society Monographs New Series, Vol. 26, Oxford University Press, Oxford, 2002.