Thresholds of Random Quasi-Abelian Codes

Yun Fan, Liren Lin

School of Mathematics and Statistics Central China Normal University, Wuhan 430079, China

Abstract

For a random quasi-abelian code of rate r, it is shown that the GV-bound is a threshold point: if r is less than the GV-bound at δ , then the probability of the relative distance of the random code being greater than δ is almost 1; whereas, if r is bigger than the GV-bound at δ , then the probability is almost 0. As a consequence, there exist many asymptotically good quasi-abelian codes with any parameters attaining the GV-bound.

Key words: Random quasi-abelian code, threshold, GV-bound, balanced code, cumulative weight enumerator.

1 Introduction

Random codes play an important role in Informatics, Statistical Physics and Coding Theory; for example, see [1], [17]. For a random linear code of rate r over a finite field F with q elements, Varshamov [22] and Pierce [19] showed in fact that the GV-bound (see (1.1) below) is a threshold point: if r is less than the GV-bound at δ where $0 < \delta < 1 - q^{-1}$, then the probability of the relative distance of the random linear code being greater than δ is almost 1; whereas, if r is bigger than the GV-bound at δ , then the probability is almost 0. Recently, in [8] the cumulative distance enumerators of random codes are introduced and their thresholds are investigated; as a consequence, the above threshold of random linear codes is redescribed explicitly with the parameters r and δ .

By means of random codes, [3] showed that, if 2 is primitive for infinitely many primes (this is a so-called *Artin's conjecture*), then the asymptotically good binary quasi-cyclic codes exist. Later, [4] and [12] made big improvements from different points of view and proved that, without the Artin's conjecture, the asymptotically good binary quasi-cyclic codes exist.

For a finite group G of order m, any element $\sum_{z\in G} a_z z$ (with $a_z\in F$) of the group algebra FG over the finite field F can be viewed as a word $(a_z)_{z\in G}$

Email address: yfan@mail.ccnu.edu.cn (Yun Fan); L_R_Lin@163.com (Liren Lin).

of length m over F. By extension, any element of the free module $(FG)^n$ of rank n can be viewed as a word of length mn. Any FG-submodule C of $(FG)^n$ is called a *quasi-group code* of index n. The code C is just the so-called *group code* if n = 1; whereas it is just the usual quasi-cyclic code of index n if G is cyclic. And, C is called a *quasi-abelian code* if G is abelian; see [6], [23].

In 2006, Bazzi and Mitter [2] constructed a class of random binary quasiabelian codes and a class of random binary dihedral group codes, and showed that the probability of the parameters of the random codes of any one of the two classes attaining GV-bound is large; as a consequence, within the two classes the asymptotically good codes exist. Soon after, with the similar random method Martínez-Pérez and Willems [14] proved that self-dual doubly-even binary dihedral group codes are asymptotically good.

We are interested in general random quasi-abelian codes and their thresholds. Modifying the random linear code ensemble in Shannon's Information Theory (cf. [17, ch.6]), in Section 2 we construct the general random quasi-abelian code ensemble, and state our main theorem, see Theorem 2.1 below, which asserts that the GV-bound is still a threshold point, i.e. the probability of the relative distance of the random code of the ensemble being greater than a given δ is almost 1 if the parameters are below the GV-bound; whereas, the probability is almost 0 if the parameters are beyond the GV-bound. The Varshamov-Pierce's threshold for random linear codes mentioned above is the special case of our main theorem by taking the finite group to be trivial.

The proof of the main theorem consists of three parts. In Section 3, we extend a result on weights of so-called *balanced codes*; this result appeared in [15], [20] and [21] in a binary version, which played a key role in [2] and [14]. We generalize it to any q-ary version, see Theorem 3.3 below, so that we can treat any q-ary codes. Theorem 3.3 has independent significance; for example, from it quite a part of [2] can be extended to any q-ary case.

In Section 4, a threshold of the expectation of the cumulative weight enumerator of the random code of the ensemble is obtained in Theorem 4.1 below, from which the first part ("below the GV-bound") of the main theorem follows immediately.

In Section 5, we prove the second part ("beyond the GV-bound") of the main theorem by estimating the second moment of the cumulative weight enumerator of the random code of the ensemble.

From the random quasi-abelian code ensemble and the main theorem, in Section 6, we draw the random quasi-abelian codes of given rate r and describe their thresholds; in particular, for any finite abelian group, for any r and δ attaining the GV-bound, there is a series of quasi-abelian codes such that the limit of their rates and the limit of their relative distances are equal to r and δ respectively.

In this paper, $h_q(x) = x \log_q(q-1) - x \log_q x - (1-x) \log_q(1-x)$ with the convention that $0 \log_q 0 = 0$, the function $h_q(x)$ is called the *q-ary entropy*

(different from the entropy with base q in Informatics, see [5, §2.1]); and let

$$g_q(x) = 1 - h_q(x) = 1 - x \log_q(q - 1) + x \log_q x + (1 - x) \log_q(1 - x),$$
 (1.1)

which is the q-ary asymptotic Gilbert-Varshamov bound, or GV-bound in short; note that $g_q(x)$ for $x \in [0,1]$ is a convex function and has a unique zero point at $x = 1 - q^{-1}$, hence $g_q(x)$ is a strictly decreasing function for $x \in [0, 1 - q^{-1}]$; see [10, §2.10.6]. About fundamentals on coding theory and group theory, please refer to [10] and [11] respectively.

2 Random quasi-abelian code ensembles

In this paper we always assume that F is a finite field with cardinality $|F| = q = p^e$ where p is a prime, and G is a finite abelian group of order |G| = m.

By $FG = \left\{ \sum_{z \in G} a_z z \mid a_z \in F \right\}$ we denote the group algebra of G over F. Each element $a = \sum_{z \in G} a_z z$ of FG is viewed as a word $(a_z)_{z \in G}$ of length m over F, and $\mathbf{w}(a) = \mathbf{w}((a_z)_{z \in G})$ stands for the usual Hamming weight of the word $(a_z)_{z \in G}$. In this way, $a = \sum_{z \in G} a_z z \in FG$ and word $(a_z)_{z \in G} \in F^m$ are identified with each other; but note that for $a, b \in FG$ we have the product ab in the algebra FG.

Let n be any positive integer. We consider the free FG-module of rank n:

$$(FG)^n = \{ \mathbf{a} = (a_1, \dots, a_n) \mid a_i \in FG, \ i = 1, \dots, n \}.$$

Each element $\mathbf{a} = (a_1, \dots, a_n) \in (FG)^n$ is identified with a concatenated word $((a_{1z})_{z \in G}, \dots, (a_{nz})_{z \in G})$ of length mn over F, thus the Hamming weight $\mathbf{w}(\mathbf{a}) = \mathbf{w}(a_1, \dots, a_n) = \mathbf{w}(a_1) + \dots + \mathbf{w}(a_n)$. As mentioned in Introduction, any submodule C of the FG-module $(FG)^n$ is said to be a quasi-abelian code of G over F (or quasi-FG code more precisely) with index n. In particular, it is just the usual abelian code if n = 1; whereas, it is just the usual quasi-cyclic code with index n if G is cyclic.

We always take the following parameters:

$$r \in (0,1), \quad \delta \in (0,\delta_0) \text{ where } \delta_0 = 1 - q^{-1},$$
 (2.1)

and set k = [rn], the integer nearest to rn. We consider the set of $k \times n$ matrices over FG:

$$(FG)^{k \times n} = \left\{ A = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \cdots & \cdots & \cdots \\ a_{k1} & \cdots & a_{kn} \end{pmatrix} \middle| a_{ij} \in FG \right\},$$
 (2.2)

which is viewed as a probability space with equiprobability. Following a notation in Shannon's information theory, we call this probability space the random quasi-abelian code ensemble. In particular, if G = 1 is trivial then $(FG)^{k \times n} = F^{k \times n}$ is just the usual random linear code ensemble; cf. [17, ch.6].

Take $A = (a_{ij})_{k \times n} \in (FG)^{k \times n}$, i.e. A is a random $k \times n$ matrix over FG. We write $A = (A_1, \dots, A_n)$ with $A_j = (a_{1j}, \dots, a_{kj})^T$ being the j'th column of the matrix A, where the superscript "T" stands for the transpose. Then we have a random quasi-abelian code C_A of index n as follows:

$$C_A = \left\{ \mathbf{b}A = \left(\mathbf{b}A_1, \dots, \mathbf{b}A_n \right) \mid \mathbf{b} = (b_1, \dots, b_k) \in (FG)^k \right\}, \tag{2.3}$$

where $\mathbf{b}A_j = b_1 a_{ij} + \dots + b_k a_{kj} \in FG$. Note that the rate $R(C_A) = \frac{\dim C_A}{mn}$. It is obvious that $R(C_A) \leq \frac{k}{n} \approx r$, and $R(C_A) = \frac{k}{n}$ if and only if the FG-rank of A is equal to k; so, we can get the random quasi-abelian codes of rate r from the ensemble, see Section 6 below. About the rank of a matrix over a ring, please see $[7, \S 2]$, or related refs such as [16].

By $\Delta(C_A)$ we denote the relative distance of the random quasi-abelian code C_A , i.e. $\Delta(C_A) = \frac{\mathrm{w}(C_A)}{mn}$, where $\mathrm{w}(C_A)$ denotes the minimum weight of C_A . Then $\Delta(C_A)$ is a random variable over the probability space $(FG)^{k \times n}$. We consider the asymptotic property (with $n \to \infty$) of $\mathrm{Pr}(\Delta(C_A) > \delta)$ which stands for the probability that $\Delta(C_A) > \delta$, and state our main theorem.

Theorem 2.1. Let notations be as in (2.1), (2.2) and (2.3). Then

$$\lim_{n \to \infty} \Pr\left(\Delta(C_A) > \delta\right) = \begin{cases} 1, & \text{if } r < g_q(\delta); \\ 0, & \text{if } r > g_q(\delta); \end{cases}$$

and both the limits converge exponentially.

If G = 1 is trivial, then FG = F is just the finite field F and the theorem exhibits just the threshold of random linear codes obtained by Vasharmov [22] and Pierce [19] (cf. [8, Corollary 3.2]), as mentioned in Introduction.

The key idea for the proof of the theorem is to estimate the first moment (i.e. the expectation) and the second moment of the cumulative weight enumerator of the random code C_A , so that we can bound $\Pr\left(\Delta(C_A) > \delta\right)$ suitably; for estimating the moments we need a result on weights of balanced codes which appeared in references, as we've seen so far, only in binary version, so we extend it to q-ary version first. Thus, as we mentioned in Introduction, the proof of the main theorem will be completed in Sections 3, 4 and 5.

3 The weights of balanced codes

Let $I = \{1, 2, \dots, n\}$ be an index set; let $F^I = F^n$ be the set of all words over F of length n. For any subset $I' = \{i_1, \dots, i_d\}$ of I with $1 \le i_1 < \dots < i_d \le n$, we have a projection ρ' from F^I to $F^{I'}$ as follows: $\rho'(\mathbf{a}) = (a_{i_1}, \dots, a_{i_d}) \in F^{I'}$ for any $\mathbf{a} = (a_1, \dots, a_n) \in F^I$.

Definition 3.1. Let $C \subseteq F^n = F^I$. If there are subsets I_1, \dots, I_s (with repetition allowed) of the index set I with every cardinality $|I_j| = d$ and an integer t such that

- (i) for any index $i \in I$, the number of the subscripts j satisfying that $i \in I_j$ is equal to t;
- (ii) for any $j=1,\dots,s$, the projection $\rho_j:F^I\to F^{I_j}$ maps C bijectively onto F^{I_j} ;

then we say that C is a balanced code of F^n with information length d, and I_1, \dots, I_s form a balanced system of information index sets of C.

Remark 3.2. For example, any group code C (i.e. any ideal) of the group algebra FG is a balanced code, see [2]; similarly, any coset a + C for $a \in FG$ is a balanced code too.

For any word $\mathbf{a} = (a_1, \dots, a_n) \in F^n$, the fraction $\mathbf{w}(\mathbf{a})/n$ is called the relative weight of \mathbf{a} . The following is a generalization of a result in [15], [20] and [21], where only the binary case is considered.

Theorem 3.3. Let C be a balanced code of F^n with information length d and B be a non-empty subset of C, and let $\omega = \frac{\sum_{\mathbf{b} \in B} \mathbf{w}(\mathbf{b})}{n|B|}$ (the average relative weight of B). If $0 \le \omega \le 1 - q^{-1}$, then

$$|B| \le q^{dh_q(\omega)}. (3.1)$$

Before proving the theorem, we show two corollaries.

Corollary 3.4. Let C be a balanced code of F^n with information length d, let $C^{\leq \delta}$ be the set of the codewords of C which relative weight are at most δ . If $0 \leq \delta \leq 1 - q^{-1}$, then $|C^{\leq \delta}| \leq q^{dh_q(\delta)}$.

Proof. The average relative weight of $C^{\leq \delta}$ is at most δ , and $h_q(x)$ is an increasing function in $[0, 1-q^{-1}]$.

For $C \subseteq F^n$, the Cartesian product of n' copies of C in $(F^n)^{n'}$ is as follows:

$$C^{n'} = \{ (\mathbf{c}_1, \dots, \mathbf{c}_{n'}) \mid \mathbf{c}_i \in C, \ i = 1, \dots, n' \}.$$

$$(3.2)$$

Corollary 3.5. Let C be a balanced code of F^n with information length d. Then the product code $C^{n'}$ is a balanced code of $F^{nn'}$ with information length dn'; in particular, if $0 \le \delta \le 1 - q^{-1}$ then $|(C^{n'})^{\le \delta}| \le q^{dn'h_q(\delta)}$.

Proof. Assume that the subsets I_1, \dots, I_s of the index set $I = \{1, \dots, n\}$ form a balanced system of information index sets of C. We write the index set of the product code $C^{n'}$ as:

$$I^{n'} = \{1^{(1)}, \cdots, n^{(1)}, \cdots, 1^{(n')}, \cdots, n^{(n')}\}.$$

For each $I_j = \{j_1, \dots, j_d\}$, we can form a subset $I_j^{n'}$ of $I^{n'}$ by concatenating n' copies of I_j as follows:

$$I_i^{n'} = \{j_1^{(1)}, \cdots, j_d^{(1)}, \cdots, j_1^{(n')}, \cdots, j_d^{(n')}\}.$$

Then it is easy to check that $I_1^{n'},\ \cdots,\ I_s^{n'}$ form a balanced system of information index sets of the product code $C^{n'}$.

The rest of this section contributes to the proof of the theorem.

Proof of Theorem 3.3. First we assume that d = n, i.e. $C = F^n$ which is of course balanced (with s = 1, $I_1 = I$ and t = 1), and prove the inequality (3.1); this is a key step of the proof.

Set M = |B|. Consider B as a probability space with equiprobability. Each $\mathbf{b} \in B$ is an n-tuple: $\mathbf{b} = (b_1, \dots, b_n)$. For each index $i, 1 \le i \le n$, we have a random variable X_i defined over the probability space B and taking values in F as follows: $X_i(\mathbf{b}) = b_i$; hence we have a discrete distribution function $p_i(a) = \Pr(X_i = a)$ for $a \in F$; we write the distribution as:

$$p_i = (p_i(a))_{a \in F}, \qquad i = 1, \dots, n.$$

Set

$$p = \frac{p_1 + \dots + p_n}{n};$$

then p is a distribution function. Denote $F^* = F \setminus \{0\}$ (which denotes the difference set). It is obvious that

$$\omega = \sum_{a \in F^*} p(a) = \sum_{a \in F^*} \sum_{i=1}^n \frac{p_i(a)}{n};$$
 (3.3)

hence we also have that

$$1 - \omega = p(0) = \sum_{i=1}^{n} \frac{p_i(0)}{n}.$$
 (3.4)

Consider the random *n*-tuple $\mathbf{X} = (X_1, \dots, X_n)$ and its entropy with base q:

$$H_q(\mathbf{X}) = H_q(X_1, \dots, X_n) = \sum_{\mathbf{a} \in F^n} -\Pr(\mathbf{X} = \mathbf{a}) \log_q \Pr(\mathbf{X} = \mathbf{a}).$$

For any $\mathbf{a} = (a_1, \dots, a_n) \in F^n$, by the definition of the random variables X_i 's, we have

$$\Pr(\mathbf{X} = \mathbf{a}) = \begin{cases} \frac{1}{M}, & \mathbf{a} \in B; \\ 0, & \mathbf{a} \notin B. \end{cases}$$

So we get

$$H_q(\mathbf{X}) = H_q(X_1, \dots, X_n) = \log_q M. \tag{3.5}$$

On the other hand, by an inequality for entropy of joint distribution (see [5, Theorem 2.6.6]), we have

$$H_q(X_1, \dots, X_n) \le H_q(X_1) + \dots + H_q(X_n) = \sum_{i=1}^n \sum_{a \in F} -p_i(a) \log_q p_i(a);$$

so

$$H_q(\mathbf{X}) \le \left(\sum_{i=1}^n -p_i(0)\log_q p_i(0)\right) + \left(\sum_{i=1}^n \sum_{a \in F^*} -p_i(a)\log_q p_i(a)\right).$$

Since $-x \log_q x$ is a concave function, for the second bracket of the right hand side of the above inequality we get (with the help of Eqn (3.3))

$$\frac{\sum_{i=1}^{n} \sum_{a \in F^{*}} -p_{i}(a) \log_{q} p_{i}(a)}{n(q-1)} \leq -\frac{\sum_{i=1}^{n} \sum_{a \in F^{*}} p_{i}(a)}{n(q-1)} \log_{q} \frac{\sum_{i=1}^{n} \sum_{a \in F^{*}} p_{i}(a)}{n(q-1)} \\
= -\frac{\omega}{q-1} \log_{q} \frac{\omega}{q-1};$$

that is

$$\sum_{i=1}^{n} \sum_{a \in F^*} -p_i(a) \log_q p_i(a) \le n \left(\omega \log_q (q-1) - \omega \log_q \omega\right).$$

Similarly, with the help of Eqn (3.4) we can obtain

$$\sum_{i=1}^{n} -p_i(0) \log_q p_i(0) \le -n(1-\omega) \log_q (1-\omega).$$

Thus we get

$$H_q(\mathbf{X}) \le n \Big(\omega \log_q(q-1) - \omega \log_q \omega - (1-\omega) \log_q(1-\omega) \Big) = nh_q(\omega).$$

Combining it with Eqn (3.5), we obtain that

$$\log_q |B| = \log_q M \le nh_q(\omega).$$

which is just the inequality (3.1) since we have assumed that d = n.

Next we turn to the general case. That is, there are subsets I_1, \dots, I_s of the index set $I = \{1, 2, \dots, n\}$ with each $|I_j| = d$ such that any index $i \in I$ appears in exactly t members of the s subsets I_1, \dots, I_s ; in particular, we have

$$tn = sd. (3.6)$$

Set |B| = M again. For each I_j , by ρ_j we denote the projection from $F^n = F^I$ onto F^{I_j} ; then $|\rho_j(B)| = M$.

Let \hat{I} be the disjoint union of I_1, \dots, I_s (though they may be not disjoint), so $|\hat{I}| = sd$, and $F^{\hat{I}} = F^{I_1} \times \dots \times F^{I_s}$ is the product of F^{I_j} for $j = 1, \dots, s$, i.e. the words of $F^{\hat{I}}$ are the concatenations of the words of F^{I_j} for $j = 1, \dots, s$:

$$F^{\hat{I}} = \left\{ (\mathbf{a}_1, \cdots, \mathbf{a}_s) \mid \mathbf{a}_j \in F^{I_j}, \ j = 1, \cdots, s \right\}.$$

Consider the following subset of $F^{\hat{I}}$:

$$\hat{B} = \rho_1(B) \times \dots \times \rho_s(B) = \left\{ \left(\rho_1(\mathbf{b}_1), \dots, \rho_s(\mathbf{b}_s) \right) \mid \mathbf{b}_1, \dots, \mathbf{b}_s \in B \right\}.$$

Since $|\rho_i(B)| = M$ for $j = 1, \dots, s$, we see that

$$|\hat{B}| = M^s. \tag{3.7}$$

Set $\widehat{\mathbf{w}(\hat{B})} = \sum_{\hat{\mathbf{b}} \in \hat{B}} \mathbf{w}(\hat{\mathbf{b}})$, which can be computed as follows:

$$\widehat{\mathbf{w}}(\widehat{B}) = \sum_{\mathbf{b}_1, \dots, \mathbf{b}_s \in B} \mathbf{w}(\rho_1(\mathbf{b}_1), \dots, \rho_s(\mathbf{b}_s))$$

$$= \sum_{\mathbf{b}_1, \dots, \mathbf{b}_s \in B} \sum_{j=1}^s \mathbf{w}(\rho_j(\mathbf{b}_j)) = \sum_{j=1}^s \sum_{\mathbf{b}_1, \dots, \mathbf{b}_s \in B} \mathbf{w}(\rho_j(\mathbf{b}_j)).$$

For j = 1 we have that

$$\sum_{\mathbf{b}_{1},\cdots,\mathbf{b}_{s}\in B} \mathbf{w}(\rho_{1}(\mathbf{b}_{1})) = \sum_{\mathbf{b}_{1}\in B} \sum_{\mathbf{b}_{2},\cdots,\mathbf{b}_{s}\in B} \mathbf{w}(\rho_{1}(\mathbf{b}_{1}))$$

$$= \sum_{\mathbf{b}_{1}\in B} M^{s-1} \mathbf{w}(\rho_{1}(\mathbf{b}_{1})) = M^{s-1} \sum_{\mathbf{b}\in B} \mathbf{w}(\rho_{1}(\mathbf{b})).$$

Similarly, $\sum_{\mathbf{b}_1,\dots,\mathbf{b}_s\in B} w(\rho_j(\mathbf{b}_j)) = M^{s-1} \sum_{\mathbf{b}\in B} w(\rho_j(\mathbf{b}))$. So

$$\widehat{\mathbf{w}(\hat{B})} = \sum_{j=1}^{s} M^{s-1} \sum_{\mathbf{b} \in B} \mathbf{w} (\rho_j(\mathbf{b})) = M^{s-1} \sum_{\mathbf{b} \in B} \sum_{j=1}^{s} \mathbf{w} (\rho_j(\mathbf{b})).$$

By (i) of Definition 3.1, we have

$$\sum_{j=1}^{s} w(\rho_{j}(\mathbf{b})) = w(\rho_{1}(\mathbf{b}), \cdots, \rho_{s}(\mathbf{b})) = tw(\mathbf{b}).$$

Recalling that $\omega = \frac{\sum_{\mathbf{b} \in B} w(\mathbf{b})}{nM}$, we obtain that

$$\widehat{\mathbf{w}(\hat{B})} = M^{s-1}t \sum_{\mathbf{b} \in B} \mathbf{w}(\mathbf{b}) = M^{s-1}t\omega nM = \omega t nM^s.$$

By Eqns (3.6) and (3.7), we compute the average relative weight of \hat{B} as follows:

$$\widehat{\mathbf{w}}(\widehat{B}) / sdM^s = \omega tnM^s / sdM^s = \omega.$$

Applying the conclusion proved in the first step (i.e. the case "d=n") to the subset \hat{B} of $F^{\hat{I}}$, we obtain that $M^s=|\hat{B}|\leq q^{sdh_q(\omega)}$; in other words,

$$|B| = M \le q^{dh_q(\omega)}.$$

Theorem 3.3 is proved.

4 Cumulative weight enumerators of C_A

We keep the notations in (2.1), (2.2) and (2.3), and further set

$$\hat{\mathcal{N}}_{C_A}(\delta) = \left| \left\{ \mathbf{b} \in (FG)^k \mid 1 \le \mathbf{w}(\mathbf{b}A) \le mn\delta \right\} \right|, \tag{4.1}$$

which is a non-negative integral random variable defined over the probability space $(FG)^{k\times n}$. Obviously, $\hat{\mathcal{N}}_{C_A}(\delta)$ stands for the number of such elements **b** of $(FG)^k$ that **b**A is a non-zero codewords of C_A with relative weights at most δ ; so we call it the *cumulative weight enumerator* of the random code C_A ; in particular (cf. [8, §3]),

$$\hat{\mathcal{N}}_{C_A}(\delta) \ge 1 \iff \Delta(C_A) \le \delta.$$
 (4.2)

We are concerned with the asymptotic behavior of the expectation $E(\hat{\mathcal{N}}_{C_A}(\delta))$. The following is the main result of this section.

Theorem 4.1. Let notation be as in (2.1), (2.2), (2.3) and (4.1). Then

$$\lim_{n \to \infty} E(\hat{\mathcal{N}}_{C_A}(\delta)) = \begin{cases} 0, & r < g_q(\delta); \\ \infty, & r > g_q(\delta); \end{cases}$$

and both the limits converge exponentially.

Before proving the theorem, we show that the first part of Theorem 2.1 is an immediate consequence of the first part of the above theorem.

Corollary 4.2. If $r < g_q(\delta)$ then $\lim_{n \to \infty} \Pr(\Delta(C_A) > \delta) = 1$ and the convergence speed is exponential.

Proof. By Eqn (4.2), Markov's inequality (see [18, Theorem 3.1]) and the first part of Theorem 4.1, we have

$$\lim_{n \to \infty} \Pr\left(\Delta(C_A) \le \delta\right) = \lim_{n \to \infty} \Pr\left(\hat{\mathcal{N}}_{C_A}(\delta) \ge 1\right) \le \lim_{n \to \infty} \mathbb{E}\left(\hat{\mathcal{N}}_{C_A}(\delta)\right) = 0. \quad \Box$$

To prove Theorem 4.1 (and Theorem 2.1 also), a key step is to write $\hat{\mathcal{N}}_{C_A}(\delta)$ as a sum of Bernoulli random variables.

For every $\mathbf{b} \in (FG)^k$ we define a Bernoulli random variable over the probability space $(FG)^{k \times n}$:

$$X_{\mathbf{b}} = \begin{cases} 1, & \text{if } 1 \leq \mathbf{w}(\mathbf{b}A) \leq mn\delta; \\ 0, & \text{otherwise.} \end{cases}$$

Set $X = \sum_{\mathbf{b} \in (FG)^k} X_{\mathbf{b}}$. It is obvious that $X_{\mathbf{0}} = 0$ and

$$\hat{\mathcal{N}}_{C_A}(\delta) = \sum_{\mathbf{b} \in (FG)^k} X_{\mathbf{b}} = X. \tag{4.3}$$

Fixing any $\mathbf{b} = (b_1, \dots, b_k) \in (FG)^k$, we have an FG-homomorphism induced by \mathbf{b} as follows:

$$\beta_{\mathbf{b}}: (FG)^{k \times n} \longrightarrow (FG)^{n}, A \longmapsto \mathbf{b}A = (\mathbf{b}A_{1}, \cdots, \mathbf{b}A_{n}).$$

$$(4.4)$$

For each j, $\mathbf{b}A_j = b_1 a_{1j} + \cdots + b_k a_{kj}$; so the set of $\mathbf{b}A_j$ with A_j running over $(FG)^k$ is an ideal of FG generated by b_1, \dots, b_k , we denote it by $I_{\mathbf{b}}$:

$$I_{\mathbf{b}} = FGb_1 + \dots + FGb_k$$
, for $\mathbf{b} = (b_1, \dots, b_k) \in (FG)^k$;

and denote $d_{\mathbf{b}} = \dim I_{\mathbf{b}}$. Thus, the image of $\beta_{\mathbf{b}}$ is the product code $I_{\mathbf{b}}^n \subseteq (FG)^n$, and $\dim I_{\mathbf{b}}^n = d_{\mathbf{b}}n$.

Since $\beta_{\mathbf{b}}$ is an FG-homomorphism, the number of the pre-images in $(FG)^{k \times n}$ of every $\mathbf{a} \in I_{\mathbf{b}}^{n}$ is equal to $\frac{q^{mkn}}{q^{d_{\mathbf{b}}n}}$, which is independent of the choice of \mathbf{a} . And, by Remark 3.2 and Corollary 3.5, we have $|(I_{\mathbf{b}}^{n})^{\leq \delta}| \leq q^{d_{\mathbf{b}}nh_{q}(\delta)}$. So

$$E(X_{\mathbf{b}}) = \Pr\left(1 \le w(\mathbf{b}A) \le mn\delta\right) \le \frac{q^{d_{\mathbf{b}}nh_q(\delta)} - 1}{q^{d_{\mathbf{b}}n}};$$

that is

$$E(X_{\mathbf{b}}) \le q^{-d_{\mathbf{b}}ng_q(\delta)} - q^{-d_{\mathbf{b}}n}, \qquad \forall \ \mathbf{b} \in (FG)^k. \tag{4.5}$$

For any ideal I of FG, we denote $d_I = \dim I$ and set

$$I^{k*} = \{ \mathbf{b} \in I^k \mid I_{\mathbf{b}} = I \}; \tag{4.6}$$

in particular, $(FG)^{k*} = \{\mathbf{b} \in (FG)^k \mid I_{\mathbf{b}} = FG\}$. Obviously, we have a disjoint union $(FG)^k = \bigcup_{I \leq FG} I^{k*}$, where the subscript " $I \leq FG$ " means that I runs over the ideals of FG. Thus, by the linearity of expectation, we get

$$E(X) = E\left(\sum_{\mathbf{b} \in (FG)^k} X_{\mathbf{b}}\right) = \sum_{0 \neq I \leq FG} \sum_{\mathbf{b} \in I^{k*}} E(X_{\mathbf{b}}). \tag{4.7}$$

To get a lower bound of $E(X_b)$ for $b \in (FG)^{k*}$, we recall an estimation of a partial sum of binomials:

$$q^{nh_q(k/n) - \frac{1}{2}\log_q n} \le \sum_{i=1}^k \binom{n}{i} (q-1)^i \le q^{nh_q(k/n)}, \tag{4.8}$$

see [8, Eqn(2.3)]. One can also check the upper bound of (4.8) from Corollary 3.4 (by taking $C = F^n$, i.e. k = n in the corollary), and check the lower bound by the argument in [18, Lemma 9.2].

Let $\mathbf{b_1} \in (FG)^{k*}$, i.e. $I_{\mathbf{b_1}} = FG$; then the image of $\beta_{\mathbf{b_1}}$ in (4.4) is just the whole space $(FG)^n \cong F^{mn}$, so

$$E(X_{\mathbf{b_1}}) = \Pr\left(1 \le w(\mathbf{b_1}A) \le mn\delta\right) = \frac{\left|(F^{mn})^{\le \delta}\right| - 1}{\left|F^{mn}\right|};\tag{4.9}$$

in particular,

$$E(X_{\mathbf{b_1}}) = E(X_{\mathbf{b_2}}), \quad \forall \ \mathbf{b_1}, \mathbf{b_2} \in (FG)^{k*}.$$
 (4.10)

Further, since $|(F^{mn})^{\leq \delta}| = \sum_{i=1}^{mn\delta} {mn \choose i} (q-1)^i$ and $|F^{mn}| = q^{mn}$; by the inequality (4.8) we get that

$$\mathrm{E}(X_{\mathbf{b_1}}) \geq q^{-mng_q(\delta) - \frac{1}{2}\log_q(mn)} - q^{-mn}, \qquad \forall \ \mathbf{b_1} \in (FG)^{k*}.$$

Moreover, since $|(FG)^{k*}| = \frac{|(FG)^{k*}|}{|(FG)^k|} \cdot q^{mk}$, for $\mathbf{b_1} \in (FG)^{k*}$ we have

$$|(FG)^{k*}| \cdot \mathrm{E}(X_{\mathbf{b_1}}) \ge \frac{|(FG)^{k*}|}{|(FG)^k|} \left(q^{mn\left(\frac{k}{n} - g_q(\delta)\right) - \frac{1}{2}\log_q(mn)} - q^{-mn(1 - \frac{k}{n})} \right). \tag{4.11}$$

Recalling from (4.3) that $\hat{\mathcal{N}}_{C_A}(\delta) = X$, we show a proof of Theorem 4.1.

Proof of Theorem 4.1.

Now we assume that $r < g_q(\delta)$. Since k = [rn], there is a positive number γ such that for large enough n we have $\frac{k}{n} - g_q(\delta) < -\gamma$. For any $\mathbf{b} \in I^{k*}$ as above, since $d_{\mathbf{b}} = d_I$, from Eqn (4.5) we have $\mathrm{E}(X_{\mathbf{b}}) \leq q^{-nd_Ig_q(\delta)}$. Further, because $|I^{k*}| \leq |I^k| = q^{d_Ik}$, we get that

$$\sum_{\mathbf{b}\in I^{k*}} \mathrm{E}(X_{\mathbf{b}}) \le q^{d_I k} q^{-nd_I g_q(\delta)} = q^{nd_I \left(\frac{k}{n} - g_q(\delta)\right)} < q^{-\gamma d_I n},$$

and the right hand side is exponentially convergent to 0 as $n \to \infty$. Note that FG has only finitely many ideals, by Eqn (4.7) we obtain that

$$\lim_{n \to \infty} \mathrm{E}(\hat{\mathcal{N}}_{C_A}(\delta)) = \lim_{n \to \infty} \mathrm{E}(X) = \sum_{0 \neq I \leq FG} \lim_{n \to \infty} \sum_{\mathbf{b} \in I^{k*}} \mathrm{E}(X_{\mathbf{b}}) = 0.$$

In the following we assume that $r>g_q(\delta)$. Since k=[rn], there is a positive number γ such that for large enough n we have $\frac{k}{n}-g_q(\delta)>\gamma$ and $1-\frac{k}{n}>\gamma$. Fixing a $\mathbf{b_1}\in (FG)^{k*}$, from Eqn (4.7) and the Eqn (4.10) we have:

$$\mathrm{E}(X) \ge \sum_{\mathbf{b} \in (FG)^{k*}} \mathrm{E}(X_{\mathbf{b}}) = |(FG)^{k*}| \cdot \mathrm{E}(X_{\mathbf{b_1}}).$$

Since $k \to \infty$ as $n \to \infty$, by Lemma 4.3 below, we have $\lim_{n \to \infty} \frac{|(FG)^{k*}|}{|(FG)^k|} = 1$; so, by the inequality (4.11), we obtain the following exponentially convergent limit:

$$\lim_{n\to\infty} \mathrm{E}(X) > \lim_{n\to\infty} \frac{|(FG)^{k*}|}{|(FG)^k|} \Big(q^{mn\gamma - \frac{1}{2}\log_q(mn)} - q^{-mn\gamma} \Big) = \infty.$$

The proof of Theorem 4.1 is finished.

Lemma 4.3. Assume that $m = |G| = p^{\mu}m'$ with m' coprime to p, G has h irreducible characters over F with degree d_1, \dots, d_h respectively, and $(FG)^{k*}$ is defined as in (4.6). Then the cardinality

$$|(FG)^{k*}| = \prod_{j=1}^{h} q^{(p^{\mu}-1)d_jk} (q^{d_jk} - 1); \tag{4.12}$$

and

$$\frac{|(FG)^{k*}|}{|(FG)^k|} = \prod_{j=1}^h (1 - q^{-d_j k}) \xrightarrow[k \to \infty]{} 1$$
 (4.13)

with exponential convergence speed.

Proof. By the assumptions, the abelian group G has a subgroup G' of order m' and a subgroup G'' of order p^{μ} such that $G = G'' \times G'$; hence we can assume that the group algebra FG' has h irreducible ideals E_j over F and denote $d_j = \dim_F E_j$ for $j = 1, \dots, h$. Then each E_j is a field extension of F and

$$FG' = E_1 \oplus E_2 \oplus \cdots \oplus E_h$$
.

Since $FG \cong FG'' \otimes_F FG'$ and FG'' is a local ring with head $FG''/J(FG'') \cong F$ where J(FG'') denotes the Jacobson radical, we have

$$FG \cong R_1 \oplus \cdots \oplus R_h$$
, (4.14)

where $R_j = FG'' \otimes_F E_j$ for $j = 1, \dots, h$ is a local algebra with

$$R_j/J(R_j) \cong E_j$$
, $\dim_F R_j = p^{\mu}d_j$ and $\dim_F J(R_j) = p^{\mu}d_j - d_j$.

Thus we get

$$(FG)^k \cong R_1^k \oplus \cdots \oplus R_h^k$$

A vector $\mathbf{b}_j = (b_{j1}, \cdots, b_{jk})$ of R_j^k generates R_j (i.e. $R_j b_{j1} + \cdots + R_j b_{jk} = R_j$) if and only if the image of \mathbf{b}_j in the residue $R_j^k/J(R_j)^k \cong E_j^k$ is non-zero, i.e. $R_j^{k*} = R_j^k \backslash J(R_j)^k$ (the difference set). So we get

$$|R_j^{k*}| = q^{p^{\mu}d_jk} - q^{(p^{\mu}-1)d_jk} = q^{(p^{\mu}-1)d_jk}(q^{d_jk} - 1).$$

It is easy to check that $(FG)^{k*} = R_1^{k*} \times \cdots \times R_h^{k*}$. We obtain that

$$|(FG)^{k*}| = \prod_{j=1}^{h} q^{(p^{\mu}-1)d_jk} (q^{d_jk} - 1);$$

hence

$$\frac{|(FG)^{k*}|}{|(FG)^k|} = \prod_{i=1}^h (1 - q^{-d_j k}),$$

which converges, as $k \to \infty$, exponentially to 1.

5 Second moment method for the main theorem

In this section we keep the notations in Theorem 2.1 and Eqn (4.3).

In this section we always assume that $r > g_q(\delta)$ and prove that

$$\lim_{n \to \infty} \Pr(X \ge 1) = 1 \quad \text{with exponential convergence speed;}$$
 (5.1)

which completes the proof of Theorem 2.1, since $\Pr\left(\Delta(C_A) \leq \delta\right) = \Pr(X \geq 1)$, see (4.2), hence Eqn (5.1) implies that $\lim_{n \to \infty} \Pr\left(\Delta(C_A) > \delta\right) = 0$.

By a known inequality, see [18, Theorem 6.10], we have that

$$\Pr(X \ge 1) \ge \sum_{\mathbf{b} \in (FG)^k} \frac{\mathrm{E}(X_{\mathbf{b}})}{\mathrm{E}(X|X_{\mathbf{b}} = 1)},$$

where $E(X|X_b=1)$ denotes the conditional expectation, which is essentially involved in the second moment of X. Such a way to investigate phase transitions (thresholds) by means of second moments is usually named the *second moment method*; e.g. see [9, Appendix].

Since $(FG)^{k*}$ is a part of $(FG)^k$, see Eqn (4.6), we have

$$\Pr(X \ge 1) \ge \sum_{\mathbf{b} \in (FG)^{k_*}} \frac{E(X_{\mathbf{b}})}{E(X|X_{\mathbf{b}} = 1)}.$$
 (5.2)

By the linearity of expectations, for $\mathbf{b_1} \in (FG)^k$ we have

$$E(X|X_{\mathbf{b_1}} = 1) = E\left(\sum_{\mathbf{b} \in (FG)^k} X_{\mathbf{b}} \middle| X_{\mathbf{b_1}} = 1\right) = \sum_{\mathbf{b} \in (FG)^k} E(X_{\mathbf{b}} | X_{\mathbf{b_1}} = 1).$$

By the conditional probability formula (and noting that $X_{\mathbf{b}}$'s are 0-1 variables), we further have

$$E(X_{\mathbf{b}}|X_{\mathbf{b_1}}=1) = \frac{\Pr(X_{\mathbf{b}}=1 \& X_{\mathbf{b_1}}=1)}{\Pr(X_{\mathbf{b_1}}=1)} = \frac{E(X_{\mathbf{b}}X_{\mathbf{b_1}})}{E(X_{\mathbf{b_1}})}.$$

Set

$$\mathcal{A}(\mathbf{b}, \mathbf{b_1}) = \left\{ A \in (FG)^{k \times n} \, \middle| \, 1 \leq \mathrm{w}(\mathbf{b}A), \mathrm{w}(\mathbf{b_1}A) \leq mn\delta \right\},$$

then

$$\mathrm{E}(X_{\mathbf{b}}X_{\mathbf{b_1}}) = \mathrm{Pr}\left(X_{\mathbf{b}} = 1 \& X_{\mathbf{b_1}} = 1\right) = \frac{|\mathcal{A}(\mathbf{b}, \mathbf{b_1})|}{|(FG)^{k \times n}|} = \frac{|\mathcal{A}(\mathbf{b}, \mathbf{b_1})|}{q^{mnk}}.$$

Thus we get that

$$E(X_{\mathbf{b}}|X_{\mathbf{b_1}} = 1) = \frac{|\mathcal{A}(\mathbf{b}, \mathbf{b_1})|}{q^{mnk}E(X_{\mathbf{b_1}})}.$$
(5.3)

For any invertible $k \times k$ matrix Q over FG,

$$\begin{split} \mathcal{A}(\mathbf{b}Q,\mathbf{b_1}Q) &= & \left\{A \in (FG)^{k \times n} \,\middle|\, 1 \leq \mathbf{w}(\mathbf{b}QA), \mathbf{w}(\mathbf{b_1}QA) \leq mn\delta\right\} \\ &= & \left\{A \in (FG)^{k \times n} \,\middle|\, QA \in \mathcal{A}(\mathbf{b},\mathbf{b_1})\right\} = \left\{Q^{-1}A \,\middle|\, A \in \mathcal{A}(\mathbf{b},\mathbf{b_1})\right\}; \end{split}$$

in particular, we have that $|\mathcal{A}(\mathbf{b}Q, \mathbf{b_1}Q)| = |\mathcal{A}(\mathbf{b}, \mathbf{b_1})|$.

Now we can show that

$$E(X|X_{\mathbf{b_1}} = 1) = E(X|X_{\mathbf{b_2}} = 1), \quad \forall \ \mathbf{b_1}, \mathbf{b_2} \in (FG)^{k*}.$$
 (5.4)

To see it, by [7, Proposition 2.11] we can take an invertible $k \times k$ matrix Q over FG such that $\mathbf{b_2} = \mathbf{b_1}Q$; then, by Eqns (5.3) and (4.10), we have

$$\begin{split} \mathrm{E}(X_{\mathbf{b}}|X_{\mathbf{b_1}} = 1) &= \frac{|\mathcal{A}(\mathbf{b}, \mathbf{b_1})|}{q^{mnk}\mathrm{E}(X_{\mathbf{b_1}})} = \frac{|\mathcal{A}(\mathbf{b}Q, \mathbf{b_1}Q)|}{q^{mnk}\mathrm{E}(X_{\mathbf{b_1}})} \\ &= \frac{|\mathcal{A}(\mathbf{b}Q, \mathbf{b_2})|}{q^{mnk}\mathrm{E}(X_{\mathbf{b_2}})} = \mathrm{E}(X_{\mathbf{b}Q}|X_{\mathbf{b_2}} = 1); \end{split}$$

hence

$$\mathrm{E}(X|X_{\mathbf{b_1}}=1) \quad = \quad \sum_{\mathbf{b} \in (FG)^k} \mathrm{E}(X_{\mathbf{b}}|X_{\mathbf{b_1}}=1) = \sum_{\mathbf{b} \in (FG)^k} \mathrm{E}(X_{\mathbf{b}Q}|X_{\mathbf{b_2}}=1);$$

noting that $\mathbf{b}Q$ runs over $(FG)^k$ when \mathbf{b} runs over $(FG)^k$, we obtain that

$$E(X|X_{\mathbf{b_1}} = 1) = \sum_{\mathbf{b} \in (FG)^k} E(X_{\mathbf{b}}|X_{\mathbf{b_2}} = 1) = E(X|X_{\mathbf{b_2}} = 1),$$

which is just Eqn (5.4).

From now on to the end of this section we fix $\mathbf{b_1} = (0, \dots, 0, 1)$, which belongs obviously to $(FG)^{k*}$. By Eqns (5.2), (4.10) and (5.4), we have

$$\Pr(X \ge 1) \ge \frac{|FG^{k*}| \cdot E(X_{\mathbf{b_1}})}{E(X|X_{\mathbf{b_1}} = 1)}.$$

Thus, to prove Eqn (5.1), it is enough to prove that

$$\lim_{n \to \infty} \frac{E(X|X_{\mathbf{b_1}} = 1)}{|FG^{k*}| \cdot E(X_{\mathbf{b_1}})} = \lim_{n \to \infty} \frac{\sum_{\mathbf{b} \in (FG)^k} E(X_{\mathbf{b}}|X_{\mathbf{b_1}} = 1)}{|FG^{k*}| \cdot E(X_{\mathbf{b_1}})} = 1$$
(5.5)

and it converges exponentially.

For any $A \in (FG)^{k \times n}$, by \underline{A}_i we denote the *i*'th row of A. To compute $\mathcal{A}(\mathbf{b}, \mathbf{b_1})$, we set

$$\mathcal{A}(\mathbf{b_1}) = \{ A \in (FG)^{k \times n} \mid 1 \le w(\mathbf{b_1}A) \le mn\delta \};$$

since $\mathbf{b} = (0, \dots, 0, 1)$, it is clear that

$$\mathcal{A}(\mathbf{b_1}) = \left\{ A \in (FG)^{k \times n} \mid \mathbf{0} \neq \underline{A}_k \in \left((FG)^n \right)^{\leq \delta} \right\},\,$$

that is, $\mathcal{A}(\mathbf{b_1})$ is the set of the $k \times n$ matrices A over FG such that the k'th row $\underline{A}_k \neq \mathbf{0}$ and $\mathbf{w}(\underline{A}_k) \leq mn\delta$; see the notation in Corollary 3.4.

Given any non-zero $(a_{k1}, \dots, a_{kn}) \in ((FG)^n)^{\leq \delta}$, we denote

$$\mathcal{A}(\mathbf{b}_1)_{(a_{k1},\cdots,a_{kn})} = \left\{ A \in (FG)^{k \times n} \mid \underline{A}_k = (a_{k1},\cdots,a_{kn}) \right\}.$$

Then any $\mathbf{b} = (b_1, \dots, b_{k-1}, b_k) \in (FG)^k$ induces a map:

$$\bar{\beta}_{\mathbf{b}}: \quad \mathcal{A}(\mathbf{b_1})_{(a_{k1}, \dots . a_{kn})} \quad \longrightarrow \quad (FG)^n,$$

$$A \quad \longmapsto \quad \mathbf{b}A = b_1 \underline{A}_1 + \dots + b_{k-1} \underline{A}_{k-1} + b_k \underline{A}_k;$$
 (5.6)

Set $\bar{\mathbf{b}} = (b_1, \dots, b_{k-1}), \ I_{\bar{\mathbf{b}}} = FGb_1 + \dots + FGb_{k-1}$ which is the ideal of FG generated by b_1, \dots, b_{k-1} , and set $d_{\bar{\mathbf{b}}} = \dim I_{\bar{\mathbf{b}}}$. It is easy to see that the image of the map $\bar{\beta}_{\mathbf{b}}$ is a coset of $I_{\bar{\mathbf{b}}}^n \subseteq (FG)^n$ as follows

$$I^n_{\bar{\mathbf{b}}} + b_k \underline{A}_k, \qquad \text{with cardinality} \ |I^n_{\bar{\mathbf{b}}} + b_k \underline{A}_k| = |I^n_{\bar{\mathbf{b}}}| = q^{d_{\bar{\mathbf{b}}}n};$$

and the number of the pre-images in $\mathcal{A}(\mathbf{b_1})_{(a_{k1},\dots a_{kn})}$ of any $\mathbf{a} \in I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k$ is equal to $\frac{q^{mn(k-1)}}{q^{d_{\bar{\mathbf{b}}}^n}} = q^{mn(k-1)-d_{\bar{\mathbf{b}}}n}$, which is independent of the choices of \mathbf{a} and (a_{k1},\dots,a_{kn}) . Thus the cardinality of the pre-image in $\mathcal{A}(\mathbf{b_1})_{(a_{k1},\dots,a_{kn})}$ of the set $(I_{\bar{\mathbf{b}}} + b_k \underline{A}_k)^{\leq \delta} \setminus \{\mathbf{0}\}$ is

$$(|(I_{\bar{\mathbf{h}}}^n + b_k \underline{A}_k)^{\leq \delta}| - \lambda)q^{mn(k-1)-d_{\bar{\mathbf{h}}}n}$$

with

$$\lambda = \begin{cases} 1, & \text{if } \mathbf{0} \in I_{\bar{\mathbf{b}}} + b_k \underline{A}_k; \\ 0, & \text{otherwise;} \end{cases}$$
 (5.7)

hence

$$|\mathcal{A}(\mathbf{b}, \mathbf{b_1})| = \sum_{\mathbf{0} \neq (a_{k1}, \dots, a_{kn}) \in ((FG)^n) \leq \delta} \left(|(I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k)^{\leq \delta}| - \lambda \right) q^{mn(k-1) - d_{\bar{\mathbf{b}}} n};$$

that is

$$|\mathcal{A}(\mathbf{b}, \mathbf{b_1})| = \left(|((FG)^n)^{\leq \delta}| - 1 \right) \left(|(I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k)^{\leq \delta}| - \lambda \right) \cdot q^{mn(k-1) - d_{\bar{\mathbf{b}}}n}.$$

But $|((FG)^n)^{\leq \delta}| - 1 = q^{mn} E(X_{\mathbf{b_1}})$, see Eqn (4.9). By Eqn (5.3), we get that

$$E(X_{\mathbf{b}}|X_{\mathbf{b_1}} = 1) = \left(|(I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k)^{\leq \delta}| - \lambda \right) \cdot q^{-d_{\bar{\mathbf{b}}}n}. \tag{5.8}$$

By the disjoint union $(FG)^{k-1}=\bigcup_{I\leq FG}I^{(k-1)*}$ again, cf. Eqn (4.7) (but this time we consider $\bar{\mathbf{b}}$ which has length k-1), we have

$$\sum_{\mathbf{b} \in (FG)^k} \mathrm{E}\big(X_{\mathbf{b}} \big| X_{\mathbf{b_1}} = 1\big) = \sum_{I \leq FG} \sum_{\bar{\mathbf{b}} \in I^{(k-1)*}} \sum_{b_k \in FG} \mathrm{E}\big(X_{\mathbf{b}} \big| X_{\mathbf{b_1}} = 1\big),$$

where $\bar{\mathbf{b}} = (b_1, \dots, b_{k-1})$ and $\mathbf{b} = (b_1, \dots, b_{k-1}, b_k)$. Thus

$$\frac{\sum_{\mathbf{b} \in (FG)^k} E(X_{\mathbf{b}} | X_{\mathbf{b_1}} = 1)}{|(FG)^{k*}| \cdot E(X_{\mathbf{b_1}})} = \sum_{0 \neq I \leq FG} S_I$$
 (5.9)

with

$$S_I = \frac{\sum_{\bar{\mathbf{b}} \in I^{(k-1)*}} \sum_{b_k \in FG} \mathrm{E}(X_{\mathbf{b}} | X_{\mathbf{b_1}} = 1)}{|(FG)^{k*}| \cdot \mathrm{E}(X_{\mathbf{b_1}})}.$$

We compute the S_I 's for ideals I of FG into two cases.

Case 1. $0 \neq I \neq FG$; note that there are only finitely many such ideals of FG. Let $d_I = \dim I$; then $d_I < m$. By Remark 3.2 and Corollary 3.5, we see that $I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k$ is a balanced code and

$$|(I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k)^{\leq \delta}| - \lambda \leq |(I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k)^{\leq \delta}| \leq q^{d_{\bar{\mathbf{b}}} n h_q(\delta)}.$$

By Eqn (5.8) and the above inequality, we obtain that

$$E(X_{\mathbf{b}}|X_{\mathbf{b}_1}=1) \le q^{-d_{\bar{\mathbf{b}}}ng_q(\delta)}.$$

Note that $d_{\bar{\mathbf{b}}} = d_I$ for any $\bar{\mathbf{b}} \in I^{(k-1)*}$, $|I^{(k-1)*}| \leq |I^{(k-1)}| = q^{d_I(k-1)}$ and $|FG| = q^m$. By the inequality (4.11) and the above inequality, we have

$$S_{I} \leq \frac{q^{d_{I}(k-1)}q^{m} \cdot q^{-d_{I}ng_{q}(\delta)}}{\frac{|(FG)^{k*}|}{|(FG)^{k}|} \left(q^{mn(\frac{k}{n}-g_{q}(\delta))-\frac{1}{2}\log_{q}(mn)} - q^{-mn(1-\frac{k}{n})}\right)}$$

$$= \frac{|(FG)^{k}|}{|(FG)^{k*}|} \cdot \frac{q^{d_{I}n(\frac{k}{n}-g_{q}(\delta))-d_{I}+m}}{q^{mn(\frac{k}{n}-g_{q}(\delta))-\frac{1}{2}\log_{q}(mn)} - q^{-mn(1-\frac{k}{n})}}.$$

Since k=[rn] and $1>r>g_q(\delta)$, there is a real number $\gamma>0$ such that for large enough n we have $\frac{k}{n}-g_q(\delta)>\gamma$ and $1-\frac{k}{n}>\gamma$; hence

$$\frac{1}{S_I} \ge \frac{|(FG)^{k*}|}{|(FG)^k|} \left(q^{\gamma(m-d_I)n + \frac{1}{2}\log_q(mn) + d_I - m} - O(q^{-\gamma n}) \right),$$

where $O(q^{-\gamma n})$ stands for a quantity bounded from above by a multiple of $q^{-\gamma n}$. Recalling that $d_I < m$ and $\lim_{n \to \infty} \frac{|(FG)^k|}{|(FG)^{k*}|} = 1$ (see Eqn (4.13)), we get that

$$\lim_{n \to \infty} S_I = 0, \quad \text{if } I \neq FG, \tag{5.10}$$

and the limit converges exponentially.

Case 2. I=FG, and $\bar{\mathbf{b}}\in I^{(k-1)*}$. Then $d_{\bar{\mathbf{b}}}=m$, and $I^n_{\bar{\mathbf{b}}}=(FG)^n$, hence $I^n_{\bar{\mathbf{b}}}+b_k\underline{A}_k=(FG)^n$; in particular, $\lambda=1$ in Eqn (5.7). By Eqn (4.9), we get

$$|(I_{\bar{\mathbf{b}}}^n + b_k \underline{A}_k)^{\leq \delta}| - \lambda = |((FG)^n)^{\leq \delta}| - 1 = q^{mn} \mathbf{E}(X_{\mathbf{b_1}}).$$

By Eqn (5.8) we can compute

$$S_{I} = \frac{|(FG)^{(k-1)*}| \cdot q^{m} \cdot q^{mn} E(X_{\mathbf{b_{1}}}) \cdot q^{-mn}}{|(FG)^{k*}| \cdot E(X_{\mathbf{b_{1}}})}$$

$$= \frac{|(FG)^{(k-1)*}| \cdot q^{m}}{|(FG)^{k*}|} = \frac{|(FG)^{(k-1)*}|}{|(FG)^{k-1}|} \cdot \frac{|(FG)^{k}|}{|(FG)^{k*}|}.$$

By the exponential convergence $\lim_{n\to\infty}\frac{|(FG)^k|}{|(FG)^{k*}|}=1$ (see Eqn (4.13)) again, we get the following exponential convergent limit:

$$\lim_{n \to \infty} S_I = 1, \quad \text{if } I = FG. \tag{5.11}$$

Finally, by Eqn (5.9), Eqn (5.10) and Eqn (5.11), we obtain that

$$\lim_{n \to \infty} \frac{\sum_{\mathbf{b} \in (FG)^k} E(X_{\mathbf{b}} | X_{\mathbf{b_1}} = 1)}{|(FG)^{k*}| \cdot E(X_{\mathbf{b_1}})} = 1$$

and it converges exponentially; this is just what Eqn (5.5) requires.

6 Random quasi-abelian codes

Keep notations in (2.1), (2.2) and (2.3).

Recall that for $A \in (FG)^{k \times n}$ the rate $R(C_A) = \frac{k}{n}$ if and only if the FG-rank of A is equal to k; at that case we say that A is full-rank.

In order to get random quasi-abelian codes of rate $\frac{k}{n} \approx r$, we consider the probability space \mathcal{F} , which sample space is $\{A \in (FG)^{k \times n} \mid A \text{ is full-rank}\}$ and probability function is equiprobability. Take $\mathcal{A} \in \mathcal{F}$, construct $C_{\mathcal{A}}$ the same as in (2.3). By $\Pr_{\mathcal{F}} (\Delta(C_{\mathcal{A}}) > \delta)$ we emphasize that the probability is computed over the probability space \mathcal{F} .

Corollary 6.1. Let notation be as above. Then

$$\lim_{n \to \infty} \Pr_{\mathcal{F}} (\Delta(C_{\mathcal{A}}) > \delta) = \begin{cases} 1, & r < g_q(\delta); \\ 0, & r > g_q(\delta); \end{cases}$$

and both the limits converge exponentially.

Proof. Let $A \in (FG)^{k \times n}$. By the total probability formula we have

$$\Pr(\Delta(C_A) > \delta) = \Pr\left(\Delta(C_A) > \delta \mid A \text{ is full-rank}\right) \cdot \Pr(A \text{ is full-rank}) + \Pr\left(\Delta(C_A) > \delta \mid A \text{ is not full-rank}\right) \cdot \Pr(A \text{ is not full-rank}).$$

Noting that

$$\Pr\left(\Delta(C_A) > \delta \mid A \text{ is full-rank}\right) = \Pr_{\mathcal{F}}\left(\Delta(C_A) > \delta\right);$$

and by Lemma 6.3 below,

$$\lim_{n\to\infty}\Pr(A\text{ is not full-rank})=0,\quad \lim_{n\to\infty}\Pr(A\text{ is full-rank})=1\,;$$

so we get

$$\lim_{n \to \infty} \Pr\left(\Delta(C_A) > \delta\right) = \lim_{n \to \infty} \Pr_{\mathcal{F}}\left(\Delta(C_A) > \delta\right).$$

Then the corollary follows from Theorem 2.1 at once.

From the first part (the case $r < g_q(\delta)$) of the above corollary we obtain the following result immediately.

Corollary 6.2. For any $(r, \delta) \in (0, 1) \times (0, 1 - q^{-1})$ satisfying that $r < g_q(\delta)$, there exists a series of quasi-FG codes C_1, C_2, \cdots such that:

- (i) the length of C_i goes to infinity;
- (ii) $\lim_{i \to \infty} R(C_i) = r;$ (iii) $\lim_{i \to \infty} \Delta(C_i) \ge \delta.$

Lemma 6.3. Let $A \in (FG)^{k \times n}$ where k = [rn] and 0 < r < 1. Then $\lim_{n\to\infty} \Pr(A \text{ is full-rank}) = 1$, or equivalently, $\lim_{n\to\infty} \Pr(A \text{ is not full-rank}) = 0$; and both the limits converge exponentially.

Proof. Let $A = (a_{\alpha\beta})_{k \times n}$ with $a_{\alpha\beta} \in FG$. We quote the decomposition of FG in (4.14) and adopt their notations. Each $a_{\alpha\beta}$ can be written as

$$a_{\alpha\beta} = (a_{\alpha\beta}^{(1)}, \cdots, a_{\alpha\beta}^{(h)}), \qquad a_{\alpha\beta}^{(j)} \in R_j;$$

hence the matrix A can be rewritten as $A = (A^{(1)}, \dots, A^{(h)})$ with $A^{(j)} =$ $(a_{\alpha\beta}^{(j)})_{k\times n}$ being $k\times n$ matrix over the local algebra R_j (cf. [7, Eqn (2.2)]), and A is full-rank if and only if every $A^{(j)}$ is full-rank over the field E_j for $j=1,\cdots,h,$ cf. [7, Lemma 2.2]. For $1\leq i\neq j\leq h,$ it is clear that $A^{(i)}\in R_i^{k\times n}$ and $A^{(j)} \in \mathbb{R}_{j}^{k \times n}$ are randomly independent of each other. So we have

$$\Pr\left(A \text{ is full-rank}\right) = \prod_{j=1}^{h} \Pr\left(A^{(j)} \text{ is full-rank}\right). \tag{6.1}$$

Let j with $1 \le j \le h$ be given. First we claim that

$$A^{(j)}$$
 is full-rank $\iff \bar{A}^{(j)}$ is full-rank, (6.2)

where $\bar{A}^{(j)} = (\bar{a}_{\alpha\beta}^{(j)})_{k\times n}$ is the image of $A^{(j)}$ in $E_j^{k\times n}$, i.e. each $\bar{a}_{\alpha\beta}^{(j)}$ is the image of the element $a_{\alpha\beta}^{(j)}$ in the residue filed $E_j = R_j/J(R_j)$. To see it, we remark that $A^{(j)}$ is full-rank if and only if it is right invertible, cf. [7, Lemma 2.6]. Suppose that $A^{(j)}$ is full-rank, then $A^{(j)}B = I_{k \times k}$ for a $B \in R_j^{n \times k}$, where $I_{k \times k}$ stands for the identity $k \times k$ matrix; mapping them to matrices over E_j , we get that $\bar{A}^{(j)}\bar{B} = \bar{I}_{k\times k}$, which implies that $\bar{A}^{(j)}$ is full-rank over E_j . Conversely, if $\bar{A}^{(j)}$ is full-rank over E_j , then $\bar{A}^{(j)}\bar{B} = \bar{I}_{k\times k}$ for a $B \in R_j^{n\times k}$, hence

$$A^{(j)}B = I_{k \times k} + C$$
, with $C \in J(R_i)^{k \times k}$;

since C is a nilpotent matrix, $I_{k\times k}+C$ is an invertible matrix; hence $A^{(j)}$ is full-rank over R_j .

Next we claim that

$$\Pr\left(A^{(j)} \text{ is not full-rank}\right) \le q^{d_j(k-n)} \approx q^{d_j(r-1)n}. \tag{6.3}$$

To see it, we note three points: $\bar{A}^{(j)}$ is not full-rank if and only if there a (k-1)-dimensional subspace of E_j^k which contains all the columns of $\bar{A}^{(j)}$; the probability that a (k-1)-dimensional subspace of E_j^k contains all the columns of $\bar{A}^{(j)}$ is $1/q^{d_j n}$ (recall that the cardinality $|E_j| = q^{d_j}$); the number of the (k-1)-dimensional subspaces of E_j^k is $\frac{q^{d_j k}-1}{q^{d_j-1}} \leq q^{d_j k}$; thus

$$\Pr\left(\bar{A}^{(j)} \text{ is not full-rank}\right) \leq q^{d_j k} \cdot \frac{1}{q^{d_j n}} = q^{d_j (k-n)}.$$

Each matrix in $E_j^{k \times n}$ has exactly $|J(R_j)|^{kn}$ inverse images in $R_j^{k \times n}$. So the claim (6.3) follows from the above inequality and the conclusion (6.2).

Finally, since r-1 < 0, from the inequality (6.3) we obtain

$$\lim_{n\to\infty} \Pr\left(A^{(j)} \text{ is not full-rank}\right) \leq \lim_{n\to\infty} q^{d_j(r-1)n} = 0.$$

By Eqn (6.1), we are done for the lemma.

Acknowledgements

The research of the authors is supported by NSFC with grant numbers 11171370 and 11271005.

References

- [1] A. Barg, G. D. Forney, "Random codes: Minimum distances and error exponents", *IEEE Trans. Inform. Theory*, vol.48, pp.2568-2573, 2002.
- [2] L. M. J. Bazzi, S. K. Mitter, "Some randomized code constructions from group actions", *IEEE Trans. Inform. Theory*, vol.52, pp.3210-3219, 2006.
- [3] C.L. Chen, W.W. Peterson, E.J. Weldon, "Some results on quasi-cyclic codes", Information and Control, vol.15, pp407-423, 1969.
- [4] V. Chepyzhov, "New lower bounds for minimum distance of linear quasi-cyclic and almost linear quasi-cyclic codes", *Problemy Peredachi Inform.*, vol.28, pp33-44, 1992.

- [5] Thomas M. Cover, Joy A. Thomas, *Elements of Information Theory*, John Wiley & Sons, Inc., 1991.
- [6] B. K. Dey, B. S. Rajan "Codes Closed Under Arbitrary Abelian Group of Permutations", Siam J. Discrete Math., vol.18, pp1-18, 2004.
- [7] Yun Fan, San Ling, Hongwei Liu, "Matrix product codes over finite commutative Frobenius rings", Des. Codes Cryptogr., published online: July 2012. DOI 10.1007/s10623-012-9726-y.
- [8] Yun Fan, San Ling, Hongwei Liu, Jing Shen and Chaoping Xing, "Cumulative distance enumerators of random codes and their thresholds", arXiv:1212.5679, 2012.
- [9] Yun Fan, Jing Shen, "On the phase transitions of random k-constraint satisfaction problems", Artificial Intelligence, vol.175, pp.914-927, 2011.
- [10] W. C. Huffman, V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, 2003.
- [11] B. Huppert, N. Blackburn, Finite Groups II, Springer Verlag, Berlin, 1982.
- [12] T. Kasami, "A Gilbert-Varshamov bound for quasi-cyclic codes of rate 1/2", IEEE Trans. Inform. Theory, vol.20, p679, 1974.
- [13] San Ling, P. Solé, "Good self-dual quasi-cyclic codes exist", IEEE Trans. Inform. Theory, vol.49, pp.1052-1053, 2003.
- [14] C. Martínez-Pérez, W. Willems, "Self-dual double-even 2-quasi-cyclic transitive codes are asymptotically good", *IEEE Trans. Inform. Theory*, vol.53, pp.4302-4308, 2007.
- [15] J. L. Massey, "On the fractional weight of distinct binary n-tuples", IEEE Trans. Inform. Theory, vol.20, p.130, 1974.
- [16] B.R. McDonald, Finite Rings with Identity, New York: Marcel Dekker, 1974.
- [17] M. Mézard, A. Montanari, Information, Physics and Computation, Oxford University Press, 2009.
- [18] M. Mitzenmacher, E. Upfal, Probability and Computing: Randomized Algorithm and Probabilistic Analysis, Cambridge Univ. Press, Cambridge, 2005.
- [19] J. N. Pierce, "Limit distribution of the minimum distance of random linear codes", *IEEE Trans. Inform. Theory*, vol.13, pp.595-599, 1967.
- [20] P. H. Piret, "An upper bound on the weight distribution of some codes", *IEEE Trans. Inform. Theory*, vol.31, pp.520-521, 1985.
- [21] I. E. Shparlinsky, "On weight enumerators of some codes", Problemy Peredechi Inform., vol.2, pp.43-48, 1986.
- [22] R. R. Varshamov, "Estimate of the number of signals in error-correcting codes" (in Russian), Dokl. Acad. Nauk, vol.117, pp.739-741, 1957.
- [23] S. K. Wasan, "Quasi Abelian codes", Publ. Inst. Math. (Beograd) N.S., vol.21, pp201-206, 1977.