

Quantization of Drinfeld Zastava in type C

Michael Finkelberg and Leonid Rybnikov

To Vladimir Drinfeld on his 60th birthday

ABSTRACT

Drinfeld zastava is a certain closure of the moduli space of maps from the projective line to the Kashiwara flag scheme of an affine Lie algebra $\hat{\mathfrak{g}}$. In case \mathfrak{g} is the symplectic Lie algebra \mathfrak{sp}_N , we introduce an affine, reduced, irreducible, normal quiver variety Z which maps to the zastava space isomorphically in characteristic 0. The natural Poisson structure on the zastava space Z can be described in terms of Hamiltonian reduction of a certain Poisson subvariety of the dual space of a (nonsemisimple) Lie algebra. The quantum Hamiltonian reduction of the corresponding quotient of its universal enveloping algebra produces a quantization Y of the coordinate ring of Z . The same quantization was obtained in the finite (as opposed to the affine) case generically in the work of Gerasimov-Kharchev-Lebedev-Oblezin (2005). We prove that Y is a quotient of the affine Borel Yangian. The analogous results for $\mathfrak{g} = \mathfrak{sl}_N$ were obtained in our previous work.

1. Introduction

1.1

This note is a continuation of [4] where we have studied the Drinfeld zastava spaces $Z^d(\widehat{\mathfrak{sl}}_N)$ from the Invariant Theory viewpoint. Recall that given a collection of complex vector spaces $(V_l)_{l \in \mathbb{Z}/N\mathbb{Z}}$, $\underline{\dim}(V_l)_{l \in \mathbb{Z}/N\mathbb{Z}} = (d_l)_{l \in \mathbb{Z}/N\mathbb{Z}} = \underline{d}$ along with $(W_l)_{l \in \mathbb{Z}/N\mathbb{Z}}$, $\underline{\dim}(W_l)_{l \in \mathbb{Z}/N\mathbb{Z}} = (1, \dots, 1)$, we consider the space $M_{\underline{d}} = \{(A_l, B_l, p_l, q_l)_{l \in \mathbb{Z}/N\mathbb{Z}}\} =$


$$\bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} \text{End}(V_l) \oplus \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} \text{Hom}(V_l, V_{l+1}) \oplus \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} \text{Hom}(W_{l-1}, V_l) \oplus \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} \text{Hom}(V_l, W_l)$$

2010 *Mathematics Subject Classification* 19E08, (22E65, 37K10)

Keywords: chainsaw quiver, quadratic spaces, affine Yangian, hamiltonian reduction, quantization.

Both authors were partially supported by the RFBR grants 12-01-00944, 12-01-33101, 13-01-12401/13, the National Research University Higher School of Economics' Academic Fund award No.12-09-0062 and the AG Laboratory HSE, RF government grant, ag. 11.G34.31.0023. This study comprises research findings from the "Representation Theory in Geometry and in Mathematical Physics" carried out within The National Research University Higher School of Economics' Academic Fund Program in 2012, grant No 12-05-0014. L.R. was supported by the RFBR grant 11-01-93105-CNRSL-a. This study was carried out within The National Research University Higher School of Economics Academic Fund Program in 2013-2014, research grant No. 12-01-0065.

of representations of the following chainsaw quiver:



Furthermore, we consider the closed subscheme $M_{\underline{d}} \subset M_{\underline{d}}$ cut out by the equations $A_{l+1}B_l - B_lA_l + p_{l+1}q_l = 0 \forall l$, and two open subschemes $M_{\underline{d}}^s \subset M_{\underline{d}}$ (resp. $M_{\underline{d}}^c \subset M_{\underline{d}}$) formed by all $\{(A_l, B_l, p_l, q_l)_{l \in \mathbb{Z}/N\mathbb{Z}}\}$ such that for any $\mathbb{Z}/N\mathbb{Z}$ -graded subspace $V'_l \subset V_l$ with $A_lV'_l \subset V'_l$, and $B_lV'_l \subset V'_{l+1} \forall l$, if $p_l(W_{l-1}) \subset V'_l \forall l$, then $V'_l = V_l$. (resp. if $V'_l \subset \text{Ker } q_l \forall l$, then $V'_l = 0$). The group $G(V_{\bullet}) = \prod_{l \in \mathbb{Z}/N\mathbb{Z}} GL(V_l)$ acts naturally on $M_{\underline{d}}$ preserving the subschemes $M_{\underline{d}}, M_{\underline{d}}^s, M_{\underline{d}}^c$.

According to [4], [3], the action of $G(V_{\bullet})$ on $M_{\underline{d}} \cap M_{\underline{d}}^s \cap M_{\underline{d}}^c$ is free, and the quotient $(M_{\underline{d}} \cap M_{\underline{d}}^s \cap M_{\underline{d}}^c)/G(V_{\bullet})$ is naturally isomorphic to the moduli space $\overset{\circ}{Z}^{\underline{d}}$ of based maps of degree \underline{d} from the projective line to the Kashiwara flag scheme of the affine Lie algebra $\widehat{\mathfrak{sl}}_N$. Moreover, the categorical quotient $M_{\underline{d}}//G(V_{\bullet})$ is naturally isomorphic to the Drinfeld zastava closure $Z^{\underline{d}}(\widehat{\mathfrak{sl}}_N) \supset \overset{\circ}{Z}^{\underline{d}}$. Furthermore, the scheme $M_{\underline{d}}//G(V_{\bullet}) \simeq Z^{\underline{d}}(\widehat{\mathfrak{sl}}_N)$ is reduced, irreducible, normal. One of the crucial points in proving this consists in checking that $(M_{\underline{d}} \cap M_{\underline{d}}^s \cap M_{\underline{d}}^c) \subset M_{\underline{d}}$ is dense, while $M_{\underline{d}} \subset M_{\underline{d}}$ is a complete intersection.

1.2

One of the goals of this note is to extend the above results to the case of zastava spaces for the affine *symplectic* Lie algebra $\widehat{\mathfrak{sp}}_N$ (in case N is even). We prove that any zastava scheme for $\widehat{\mathfrak{sp}}_N$ is reduced, irreducible, normal (note that these properties of zastava schemes were established in [3] for all *finite dimensional* simple Lie algebras).

To this end we again invoke Invariant Theory. Following [1], we equip $W := \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} W_l$ with a symplectic form such that W_l and W_k are orthogonal unless $l + k = N - 1$. We equip $V := \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} V_l$ with a nondegenerate *symmetric* bilinear form such that V_l and V_k are orthogonal unless $l + k = 0$. In particular, we must have $d_{-l} = d_l \forall l$, so that the collection \underline{d} is encoded by $\bar{d} := (d_0, d_1, \dots, d_{N/2})$. We denote by $O(V_{\bullet})$ the Levi subgroup of the orthogonal group $O(V)$ preserving the decomposition $V := \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} V_l$. We consider the space $M_{\underline{d}}^{-1} \subset M_{\underline{d}}$ of representations of the *quadratic chainsaw quiver* formed by all the selfadjoint collections $A_l^* = A_l$, $B_l^* = B_{-l-1}$, $p_l^* = q_{-l}$. We denote by $M_{\underline{d}}^{-1} \subset M_{\underline{d}}^{-1}$ the scheme-theoretic intersection $M_{\underline{d}} \cap M_{\underline{d}}^{-1}$.

We prove that $M_{\underline{d}}^{-1} \cap M_{\underline{d}}^s = M_{\underline{d}}^{-1} \cap M_{\underline{d}}^c \subset M_{\underline{d}}^{-1}$ is dense, while $M_{\underline{d}}^{-1} \subset M_{\underline{d}}^{-1}$ is a complete intersection. We deduce that the categorical quotient $M_{\underline{d}}^{-1}//O(V_{\bullet})$ is reduced, irreducible and normal. Furthermore, we prove that the action of $O(V_{\bullet})$ on $M_{\underline{d}}^{-1} \cap M_{\underline{d}}^s = M_{\underline{d}}^{-1} \cap M_{\underline{d}}^c$ is free, and the quotient $(M_{\underline{d}}^{-1} \cap M_{\underline{d}}^s)/O(V_{\bullet})$ is naturally isomorphic to the moduli space $\overset{\circ}{Z}^{\bar{d}}(\widehat{\mathfrak{sp}}_N)$ of based maps of degree \bar{d} from the projective line to the Kashiwara flag scheme of the affine Lie algebra $\widehat{\mathfrak{sp}}_N$. Moreover, the categorical quotient $M_{\underline{d}}^{-1}//G(V_{\bullet})$ is naturally isomorphic to the Drinfeld zastava closure $Z^{\bar{d}}(\widehat{\mathfrak{sp}}_N) \supset \overset{\circ}{Z}^{\bar{d}}(\widehat{\mathfrak{sp}}_N)$.

1.3

Quite naturally, we would like to extend the above results to the case of the affine orthogonal Lie algebra $\widehat{\mathfrak{so}}_N$. To this end we change the parities of the bilinear forms in 1.2. That is, we equip W with a nondegenerate *symmetric* bilinear form, and we equip V with a *symplectic* form. The corresponding space of representations of the quadratic chainsaw quiver is denoted by $M_{\underline{d}}^1$, and the corresponding Levi subgroup of $Sp(V)$ is denoted by $Sp(V_{\bullet})$. It is still true that the action of $Sp(V_{\bullet})$ on $M_{\underline{d}}^1 \cap M_{\underline{d}}^s = M_{\underline{d}}^1 \cap M_{\underline{d}}^c$ is free, and the quotient $(M_{\underline{d}}^1 \cap M_{\underline{d}}^s)/Sp(V_{\bullet})$ is naturally isomorphic to $\overset{\circ}{Z}^{\bar{d}}(\widehat{\mathfrak{so}}_N)$.

However, we encounter the following mysterious obstacle: $M_{\underline{d}}^1$ is *not* irreducible in general, and $M_{\underline{d}}^1 \cap M_{\underline{d}}^s = M_{\underline{d}}^1 \cap M_{\underline{d}}^c$ is only dense in one of its irreducible components. For this reason the categorical quotient $M_{\underline{d}}^1//Sp(V_{\bullet})$ is *not* isomorphic to the zastava space $Z^{\bar{d}}(\widehat{\mathfrak{so}}_N)$ in general. The simplest example occurs when \bar{d} is the affine simple coroot of $\widehat{\mathfrak{so}}_N$, that is $\underline{d} = (\dots, 0, 1, 2, 1, 0, \dots)$.

1.4

Following [4], we describe the natural Poisson structure on $Z^{\bar{d}}(\widehat{\mathfrak{sp}}_N)$ in quiver terms. It is obtained by the Hamiltonian reduction of a Poisson subvariety of the dual vector space of a (nonsemisimple) Lie algebra $\mathfrak{a}_{\underline{d}}^{-1}$ with its Lie-Kirillov-Kostant bracket. Now the ring of functions $\mathbb{C}[Z^{\bar{d}}(\widehat{\mathfrak{sp}}_N)]$ admits a natural quantization $\mathcal{Y}_{\bar{d}}^{-1}$ as the quantum Hamiltonian reduction of a quotient algebra of the universal enveloping algebra $U(\mathfrak{a}_{\underline{d}}^{-1})$. The algebra $\mathcal{Y}_{\bar{d}}^{-1}$ admits a homomorphism from the Borel subalgebra \mathcal{Y}^{-1} of the Yangian of type C in the case of finite Zastava space. We prove that this homomorphism is surjective. In the affine situation, there is an affine analog $\widehat{\mathcal{Y}}^{-1}$ of \mathcal{Y}^{-1} (it is no longer a subalgebra in the Yangian of $\widehat{\mathfrak{sp}}_N$), and we define it explicitly by generators and relations. We prove that there is a surjective homomorphism $\widehat{\mathcal{Y}}^{-1} \rightarrow \mathcal{Y}_{\bar{d}}^{-1}$. Moreover, we write down certain elements in the kernel of this homomorphism and conjecture that they generate the kernel (as a two-sided ideal). These elements are similar to the generators of the kernel of the Kamnitzer-Webster-Weekes-Yacobi homomorphism from *shifted Yangian* to the quantization of the transversal slices in the affine Grassmannian. In fact, as explained in [10] $\mathcal{Y}_{\bar{d}}^{-1}$ as a filtered algebra is the limit of a sequence of quantum coordinate rings of transversal slices.

2. A quiver approach to Drinfeld Zastava for symplectic groups

2.1 Quadratic spaces

We will recall the convenient terminology introduced in [8]. Let U be an N -dimensional complex vector space equipped with a nondegenerate bilinear form (\cdot, \cdot) such that $(u, v) = \varepsilon(v, u)$. It will be called a *quadratic space of type ε* (shortly an *orthogonal space* in case $\varepsilon = 1$, a *symplectic space* in case $\varepsilon = -1$). We denote by $G_{\varepsilon}(U)$ the subgroup of $GL(U)$ leaving the form invariant. So we have $G_{\varepsilon}(U) = O(N)$ or $Sp(N)$ according to $\varepsilon = 1$ or $\varepsilon = -1$.

Let $A \mapsto A^*$, $\text{End}(U) \rightarrow \text{End}(U)$ be the canonical involution associated to the form, i.e. $(Au, v) = (u, A^*v)$ for any $u, v \in U$. More generally, for a linear operator $B \in \text{Hom}(U, 'U)$ we denote by B^* the adjoint (or transposed) operator $B^* \in \text{Hom}('U^*, U^*)$.

We choose a basis w_0, \dots, w_{N-1} in a quadratic space W of type $\varepsilon = -1$ such that for $0 \leq l < N/2$ we have $(w_l, w_m) = \delta_{m, N-1-l}$ (note that N is necessarily even). The linear span of w_l will be denoted by $W_l \cong \mathbb{C}$. We will often parametrize the base vectors by the elements of $\mathbb{Z}/N\mathbb{Z}$. We define $I := \{0, 1, \dots, N/2\} \subset \{0, \dots, N-1\} = \mathbb{Z}/N\mathbb{Z}$. We set $I = I_0 \sqcup I_1$ where

$$I_0 = \{1, \dots, \frac{N}{2} - 1\}, \quad I_1 = \{0, \frac{N}{2}\}.$$

We choose another quadratic space V of type $-\varepsilon = 1$ decomposed into direct sum $V = \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} V_l$ such that V_l is orthogonal to V_m unless $l + m = 0 \in \mathbb{Z}/N\mathbb{Z}$. Let d_l denote the dimension of V_l . We set $\underline{d} := (d_l)_{l \in I}$.

We denote by $G_{-\varepsilon}(V_\bullet)$ the Levi subgroup of $G_{-\varepsilon}(V)$ preserving the decomposition $V = \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} V_l$. It is isomorphic to $O(V_0) \times O(V_{N/2}) \times \prod_{0 < l < N/2} GL(V_l)$.

2.2 Quadratic Chainsaw Quivers

Following [4, 2.3] we consider the affine space $M_{\underline{d}}^\varepsilon$ of collections $(A_l, B_l, p_l, q_l)_{l \in \mathbb{Z}/N\mathbb{Z}}$ where $A_l \in \text{End}(V_l)$, $B_l \in \text{Hom}(V_l, V_{l+1})$, $p_l \in \text{Hom}(W_{l-1}, V_l)$, $q_l \in \text{Hom}(V_l, W_l)$ satisfy the following selfadjointness conditions: $A_l^* = A_{-l}$, $B_l^* = B_{-l-1}$, $p_l^* = q_{-l}$. Here we view p_l (resp. q_l) as a vector (resp. covector) of V_l using the identification of all W_m with \mathbb{C} .

Following *loc. cit.* we consider the subscheme $M_{\underline{d}}^\varepsilon \subset M_{\underline{d}}^\varepsilon$ parametrizing the \underline{d} -dimensional representations of the Chainsaw Quiver with bilinear form (or the *Quadratic Chainsaw Quiver* for short), cut out by the equations $A_{l+1}B_l - B_lA_l + p_{l+1}q_l = 0 \forall l$.

Clearly, $M_{\underline{d}}^\varepsilon$ is acted upon by the Levi subgroup $G_{-\varepsilon}(V_\bullet)$, and we denote by $\mathfrak{Z}_{\underline{d}}^\varepsilon$ the categorical quotient $M_{\underline{d}}^\varepsilon // G_{-\varepsilon}(V_\bullet)$.

Assumption. From now on we set $\varepsilon = -1$.

2.3 Examples

We consider three basic examples in types C_1, C_2, \tilde{C}_1 .

2.3.1 C_1 We take $N \geq 2$, $d_0 = \dots = d_{N/2-1} = 0$, $d_{N/2} = d$. We have $V_{N/2} = V = \mathbb{C}^d$, $A_{N/2} = A = A^* \in \text{End}(V)$, $B_1 = 0$, $p_1 = p \in V$, $q_1 = q \in V^*$, $q(v) = (p, v)$. Thus $M_{\underline{d}}^\varepsilon = \text{End}^+(V) \oplus V$, and $\mathfrak{Z}_{\underline{d}}^\varepsilon = (\text{End}^+(V) \oplus V) // O(V)$ where $\text{End}^+(V) \subset \text{End}(V)$ stands for the linear subspace of selfadjoint operators (symmetric matrices). By the classical Invariant Theory, the ring of $O(V)$ -invariant functions on $\text{End}^+(V) \oplus V$ is freely generated by the functions $a_1, \dots, a_d, b_0, \dots, b_{d-1}$ where $a_m := \text{Tr}(A^m)$, and $b_m = (p, A^m p)$. Hence $\mathfrak{Z}_{\underline{d}}^\varepsilon \simeq \mathbb{A}^{2d}$.

2.3.2 C_2 We take $N = 4$, $d_0 = 0$, $d_1 = d_2 = d_3 = 1$. We have $V_1 = V_2 = V_3 = \mathbb{C}$, and hence all our linear operators act between one-dimensional vector spaces, and can be written just as numbers. Hence $M_{\underline{d}}^\varepsilon$ has coordinates $A_1 = A_3, A_2, B_1 = B_2, q_2 = p_3, q_3 = p_1$, and $M_{\underline{d}}^\varepsilon$ is cut out by a single equation $B_1(A_1 - A_2) = p_2 p_3$. The group $G_{-\varepsilon}(V_\bullet)$ is the product $GL(V_1) \otimes O(V_2) \simeq \mathbb{C}^* \times \{\pm 1\}$ with coordinates $c \in \mathbb{C}^*$, $s = \pm 1$. It acts on $M_{\underline{d}}^\varepsilon$ as follows: $(c, s) \cdot (A_1, A_2, B_1, p_1, p_2, p_3) = (A_1, A_2, csB_1, c^{-1}p_1, sp_2, cp_3)$. The ring of $\mathbb{C}^* \times \{\pm 1\}$ -invariant functions on $M_{\underline{d}}^\varepsilon$ is generated by the functions $A_1, A_2, b_{12} := p_2^2, b_{01} := p_1 p_3, b_{02} := p_2 B_1 p_1, b_{03} := B_1^2 p_1^2$ with three quadratic relations: $b_{02}(A_1 - A_2) = b_{01} b_{12}$, $b_{03}(A_1 - A_2) = b_{01} b_{02}$, $b_{02}^2 = b_{12} b_{03}$. Thus $\mathfrak{Z}_{\underline{d}}^\varepsilon$ is a 4-dimensional (noncomplete) intersection of 3 quadrics in \mathbb{A}^6 . According to [12], $\mathfrak{Z}_{\underline{d}}^\varepsilon$ is reduced, not \mathbb{Q} -Gorenstein, but Cohen-Macaulay, normal, and has rational singularities.

2.3.3 \tilde{C}_1 We take $N = 2$, $d_0 = d_1 = 1$. We have $V_1 = \mathbb{C} = V_2$, and hence all our linear operators act between one-dimensional vector spaces, and can be written just as numbers. Hence $M_{\underline{d}}^\varepsilon$ has coordinates $A_1, A_2, B_0 = B_1, q_0 = p_0, q_1 = -p_1$, and $M_{\underline{d}}^\varepsilon$ is cut out by a single equation $B_0(A_1 - A_0) + p_1 q_0 = 0$. The group $G_{-\varepsilon}(V_\bullet)$ is the product $O(V_0) \times O(V_1) \simeq \{\pm 1\} \times \{\pm 1\}$ with coordinates (s_1, s_2) . The ring of $\{\pm 1\} \times \{\pm 1\}$ -invariant functions on $M_{\underline{d}}^\varepsilon$ is generated by the

functions $A_0, A_1, b_1 := p_1^2, b_0 := p_0^2, s := B_0^2$ with a single relation $b_1 b_0 - s(A_0 - A_1)^2 = 0$. Note the coincidence with the output of [4, Example 2.8.3].

2.4 Dimension of $M_{\underline{d}}^\varepsilon$

We define the *factorization morphism* $\Upsilon : M_{\underline{d}}^\varepsilon \rightarrow \mathbb{A}^{\underline{d}} = \prod_{l \in I} (\mathbb{A}^{(1)})^{(d_l)}$ so that the component Υ_l is just $\text{Spec } A_l$.

PROPOSITION 2.5. *Every fiber of Υ has dimension $\dim G_{-\varepsilon}(V_\bullet) + \sum_{l \in I} d_l$.*

Proof. The same argument as in the proof of [4, Proposition 2.11]. We just list the minor changes necessary in the quadratic case. The dimension estimate for a general fiber is reduced to the the dimension estimate for the zero fiber where all A_l are nilpotent. By the adjointness condition, $(A_\bullet, B_\bullet, p_\bullet, q_\bullet)$ is determined by its components $A_l, 0 \leq l \leq N/2; B_l, 0 \leq l < N/2; p_l, 0 < l \leq N/2; q_l, 0 \leq l < N/2$, and $A_0 \in \text{End}^+(V_0), A_{N/2} \in \text{End}^+(V_{N/2})$. Note that $\dim \text{End}^+(V_l) = \frac{d_l(d_l+1)}{2}, l = 0, N/2$, while $\dim O(V_l) = \frac{d_l(d_l-1)}{2}, l = 0, N/2$. The dimension of the space of nilpotent selfadjoint operators in $\text{End}^+(V_l)$ equals $\frac{d_l(d_l-1)}{2}, l = 0, N/2$. More generally, the space $\mathbb{O}_\lambda^+ \subset \text{End}^+(V_l)$ of nilpotent selfadjoint operators of Jordan type λ (a partition of d_l) is a finite union of $O(V_l)$ -orbits all of the same dimension $\dim \mathbb{O}_\lambda^+ = \frac{1}{2} \dim \mathbb{O}_\lambda$ (where \mathbb{O}_λ is the nilpotent $GL(V_l)$ -orbit consisting of nilpotent matrices of Jordan type λ), see [9, Proposition 5]. The argument of the proof of [4, Proposition 2.11] implies that, say, the dimension of the space of $(A_0, A_1, B_0, p_1, q_0)$ subject to $A_1 B_0 - B_0 A_0 + p_1 q_0 = 0$ is at most $\frac{d_0(d_0-1)}{2} + d_1^2 - d_1 + \min(d_0, d_1) + \max(d_0, d_1)$. Summing up the similar estimates over $0 \leq l < N/2$ we obtain the desired inequality $\dim \Upsilon^{-1}(0, \dots, 0) \leq \dim G_{-\varepsilon}(V_\bullet) + \sum_{l \in I} d_l$. The opposite inequality follows from the computation of the generic fiber of Υ , and the proposition follows. \square

The following corollary is proved the same way as [4, Corollary 2.12].

COROLLARY 2.6. $M_{\underline{d}}^\varepsilon$ is an irreducible reduced complete intersection in $M_{\underline{d}}^\varepsilon$. \square

THEOREM 2.7. $\mathfrak{Z}_{\underline{d}}^\varepsilon$ is a reduced irreducible normal scheme.

Proof. The same argument as in the proof of [4, Theorem 2.7.a)]. We just list the minor changes necessary in the quadratic case. As in *loc. cit.* we have to check the normality of the open subscheme $U \subset \mathfrak{Z}_{\underline{d}}^\varepsilon$ defined as the preimage under the factorization morphism $\Phi : \mathfrak{Z}_{\underline{d}}^\varepsilon \rightarrow \mathbb{A}^{\underline{d}}$ of an open subset $\hat{U} \subset \mathbb{A}^{\underline{d}}$ formed by all the colored configurations where at most 2 points collide. As in *loc. cit.* this reduces to a few basic checks we already performed in the Examples: 2.3.1 when two points of the same color 0 or $N/2$ (*outmost color* for short) collide; Example [4, 2.8.1] when two points of the same color $l, 0 < l < N/2$ (*innermost color* for short) collide; Example 2.3.2 when a point of an outmost color collides with a point of an inmost color; Example 2.3.3 when two points of different outmost colors collide; Example [4, 2.8.2] when two points of different innermost colors collide. This completes the proof of the theorem. \square

2.8 Symplectic zastava

For $\bar{d} = (d_0, \dots, d_{N/2}) \in \mathbb{N}^I$ we consider the affine zastava space for symplectic Lie group $G = Sp_N$ introduced in [2] and denoted $\mathfrak{U}_{G,B}^{\bar{d}}$. In the present paper we will denote it $Z_\varepsilon^{\bar{d}}$. It is a reduced irreducible affine scheme containing as an open subscheme the (smooth) moduli space $\overset{\circ}{Z}_\varepsilon^{\bar{d}}$ of degree \bar{d} based maps from \mathbb{P}^1 to the affine flag scheme of $G = Sp_N$.

Recall that Sp_N is the fixed point subgroup of the involutive pinning-preserving outer automorphism $\sigma : SL_N \rightarrow SL_N$. This automorphism acts on the affine flag scheme of SL_N , and on the zastava spaces $Z^{\underline{d}}$ for SL_N . More precisely, we define $\underline{d} = (\tilde{d}_0, \dots, \tilde{d}_{N-1}) \in \mathbb{N}^N$ as follows: for $0 \leq l \leq N/2$ we have $\tilde{d}_l = d_l$, and for $0 < l < N/2$ we set $\tilde{d}_{N-l} = d_l$. Then σ acts on $Z^{\underline{d}}$, and the fixed point subscheme (with the reduced closed subscheme structure) is isomorphic to $Z_{\varepsilon}^{\bar{\underline{d}}}$. In other words, $Z_{\varepsilon}^{\bar{\underline{d}}}$ is the closure in $Z^{\underline{d}}$ of $\overset{\circ}{Z}_{\varepsilon}^{\bar{\underline{d}}} \cong (\overset{\circ}{Z}^{\underline{d}})^{\sigma}$.

Recall the chainsaw quiver variety $\mathfrak{Z}_{\underline{d}}$ introduced in [4]. Theorem [4, 2.7.b)] constructs a morphism $\eta : \mathfrak{Z}_{\underline{d}} \rightarrow Z^{\underline{d}}$, and [3, Theorem 3.5] proves that η is an isomorphism. We will identify $\mathfrak{Z}_{\underline{d}}$ and $Z^{\underline{d}}$ via η . Under this identification the open subscheme $\overset{\circ}{Z}^{\underline{d}}$ corresponds to the open subscheme $\overset{\circ}{\mathfrak{Z}}_{\underline{d}}$ formed by the closed orbits of stable and costable quadruples $(A_{\bullet}, B_{\bullet}, p_{\bullet}, q_{\bullet})$, i.e. such that V_{\bullet} is generated by the action of A_{\bullet}, B_{\bullet} from the image of p_{\bullet} , and also V_{\bullet} contains no nonzero subspaces in $\text{Ker } q_{\bullet}$ closed with respect to A_{\bullet}, B_{\bullet} . The fixed point subscheme $(\overset{\circ}{\mathfrak{Z}}_{\underline{d}})^{\sigma}$ coincides with $\overset{\circ}{\mathfrak{Z}}_{\underline{d}}^{\varepsilon}$: the open subscheme of $\mathfrak{Z}_{\underline{d}}^{\varepsilon}$ formed by the closed orbits of stable (equivalently, costable) quadruples $(A_{\bullet}, B_{\bullet}, p_{\bullet}, q_{\bullet}) \in M_{\underline{d}}^{\varepsilon}$, cf. [1, Table 1 and Proposition 3.3].

LEMMA 2.9. *The closed embedding $\overset{\circ}{\mathfrak{Z}}_{\underline{d}}^{\varepsilon} \cong (\overset{\circ}{\mathfrak{Z}}_{\underline{d}})^{\sigma} \hookrightarrow \overset{\circ}{\mathfrak{Z}}_{\underline{d}}$ extends to the closed embedding $\mathfrak{Z}_{\underline{d}}^{\varepsilon} \hookrightarrow \mathfrak{Z}_{\underline{d}}$.*

Proof. It suffices to check that any $G_{-\varepsilon}(V_{\bullet})$ -invariant function on $M_{\underline{d}}^{\varepsilon}$ extends to a $\prod_{l=0}^{N-1} GL(V_l)$ -invariant function on $M_{\underline{d}}$. This is immediately seen on the generators provided by the classical Invariant Theory. \square

Now since $\overset{\circ}{\mathfrak{Z}}_{\underline{d}}^{\varepsilon}$ is dense in $\mathfrak{Z}_{\underline{d}}^{\varepsilon} = (\mathfrak{Z}_{\underline{d}})^{\sigma}$ we conclude that $\mathfrak{Z}_{\underline{d}}^{\varepsilon} \subset \mathfrak{Z}_{\underline{d}} = Z^{\underline{d}}$ coincides with the closure of $(\overset{\circ}{\mathfrak{Z}}_{\underline{d}})^{\sigma}$ in $\mathfrak{Z}_{\underline{d}} = Z^{\underline{d}}$. Since the symplectic zastava scheme $Z_{\varepsilon}^{\bar{\underline{d}}}$ also coincides with this closure, we arrive at the following

THEOREM 2.10. *There is a canonical isomorphism $\eta : \mathfrak{Z}_{\underline{d}}^{\varepsilon} \xrightarrow{\sim} Z_{\varepsilon}^{\bar{\underline{d}}}$ making the following diagram commutative:*

$$\begin{array}{ccc} \mathfrak{Z}_{\underline{d}}^{\varepsilon} & \longrightarrow & \mathfrak{Z}_{\underline{d}} \\ \eta \downarrow & & \downarrow \eta \\ Z_{\varepsilon}^{\bar{\underline{d}}} & \longrightarrow & Z_{\underline{d}} \end{array}$$

where the horizontal morphisms are the closed embeddings of the σ -fixed points subschemes. \square

COROLLARY 2.11. *The symplectic zastava scheme $Z_{\varepsilon}^{\bar{\underline{d}}}$ is normal.* \square

3. Hamiltonian reduction


3.1 Poisson structures

According to [5] (cf. [4, 3.1,3.3]), the smooth scheme $\overset{\circ}{Z}^{\underline{d}}$ carries a canonical symplectic structure which extends as a Poisson structure to $\mathfrak{Z}_{\underline{d}} = Z^{\underline{d}}$. This Poisson structure was constructed in [4] via Hamiltonian reduction. The restriction of the symplectic form on $\overset{\circ}{Z}^{\underline{d}}$ to $\overset{\circ}{Z}_{\varepsilon}^{\bar{\underline{d}}} \cong (\overset{\circ}{Z}^{\underline{d}})^{\sigma}$ coincides with the canonical symplectic form [5] on $\overset{\circ}{Z}_{\varepsilon}^{\bar{\underline{d}}}$. We conclude that the canonical symplectic structure on $\overset{\circ}{Z}_{\varepsilon}^{\bar{\underline{d}}}$ extends as a Poisson structure to $Z_{\varepsilon}^{\bar{\underline{d}}} \cong \mathfrak{Z}_{\underline{d}}^{\varepsilon}$, and the σ -fixed point embedding $\mathfrak{Z}_{\underline{d}}^{\varepsilon} \hookrightarrow \mathfrak{Z}_{\underline{d}}$ is

Poisson. In the next subsection we will construct this Poisson structure on $\mathfrak{Z}_d^\varepsilon$ via Hamiltonian reduction.

3.2 Classical

Recall the Hamiltonian reduction definition of the Poisson bracket on Zastava spaces in type A (see [4]). We “triangulate” the chainsaw quiver in the following way:



For a pair of vector spaces U, V define the following 2-step nilpotent Lie algebra:

$$\mathfrak{n}(U, V) := U \oplus V^* \oplus (U \otimes V^*),$$

where the space $U \otimes V^*$ is central, $[U, U] = [V^*, V^*] = 0$, and for $u \in U$, $v^\vee \in V^*$ one has $[u, v^\vee] = u \otimes v^\vee$.

To define the Poisson structure, we attach to each triangle of our graph the following Lie algebra

$$\mathfrak{a}_l := (\mathfrak{gl}(V_l) \oplus \mathfrak{gl}(V_{l+1})) \rtimes \mathfrak{n}(V_l, V_{l+1})$$

(the semidirect sum is with respect to the tautological action of $\mathfrak{gl}(V_l)$ on V_l and $\mathfrak{gl}(V_{l+1})$ on V_{l+1}^*).

Consider the Lie algebra

$$\mathfrak{a}_d := \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} \mathfrak{a}_l = \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} (\mathfrak{gl}(V_l) \oplus \mathfrak{gl}(V_{l+1})) \rtimes \mathfrak{n}(V_l, V_{l+1})$$

The coadjoint representation of \mathfrak{a}_d is the space $\mathfrak{a}_d^* = \{(A_l, A'_l, B_l, p_l, q_l)_{l \in \mathbb{Z}/N\mathbb{Z}}\}$, where

$$A_l \in \text{End}(V_l), \quad A'_l \in \text{End}(V_l), \quad B_l \in \text{Hom}(V_l, V_{l+1}), \quad p_l \in V_l, \quad q_l \in V_l^*.$$

Consider the subvariety $S_d \subset \mathfrak{a}_d^*$ defined by the following equations:

$$B_l A_l + A'_{l+1} B_l + p_{l+1} q_l = 0, \quad l \in \mathbb{Z}/N\mathbb{Z}. \quad (1)$$

Let $\mathfrak{gl}(V_l)_{\text{diag}}$ be the diagonal $\mathfrak{gl}(V_l)$ inside $\mathfrak{gl}(V_l) \oplus \mathfrak{gl}(V_l) \subset \mathfrak{a}_d$ and $\pi : \mathfrak{a}_d^* \rightarrow \mathfrak{gl}(V_l)_{\text{diag}}^*$ be the projection. Then the Drinfeld Zastava space \mathfrak{Z}_d is identified with the Hamiltonian reduction $S_d // \bigoplus_{l \in \mathbb{Z}/N\mathbb{Z}} \mathfrak{gl}(V_l)_{\text{diag}} = \pi^{-1}(0) \cap S_d // \prod_{l \in \mathbb{Z}/N\mathbb{Z}} GL(V_l)_{\text{diag}}$. This provides a natural Poisson bracket on \mathfrak{Z}_d .

The involution σ acts on the space \mathfrak{a}_d^* as follows:

$$A_l \mapsto -A_{N-l}^*, \quad A'_l \mapsto -A_{N-l}^*, \quad B_l \mapsto B_{N-l}^*, \quad p_l \mapsto q_{N-l}^*, \quad q_l \mapsto p_{N-l}^*.$$

REMARK 3.3. Strictly speaking, σ is not an involution on $\mathfrak{a}_{\underline{d}}^*$ since $p^{**} = -p$ and $q^{**} = -q$, but becomes an involution after the Hamiltonian reduction.

To describe the fixed point set, we consider the half of the chainsaw quiver formed by the vertices $l \in I$. Define the Lie algebra

$$\mathfrak{a}_{\underline{d}}^\varepsilon := \bigoplus_{l=0}^{\frac{N}{2}-1} \mathfrak{a}_l = \bigoplus_{l=1}^{\frac{N}{2}-1} (\mathfrak{gl}(V_l) \oplus \mathfrak{gl}(V_{l+1})) \rtimes \mathfrak{n}(V_l, V_{l+1}).$$

The coadjoint representation of $\mathfrak{a}_{\underline{d}}^\varepsilon$ is the space $\mathfrak{a}_{\underline{d}}^{\varepsilon*} = \{(A_l, A'_l, B_l, p_l, q_l)_{l \in I_0}, A_0, q_0, B_0, A'_{\frac{N}{2}}, p_{\frac{N}{2}}\}$, where

$$A_l \in \text{End}(V_l), \quad A'_l \in \text{End}(V_l), \quad B_l \in \text{Hom}(V_l, V_{l+1}), \quad p_l \in V_l, \quad q_l \in V_l^*.$$

The invariant subvariety $S_{\underline{d}}^\varepsilon \subset \mathfrak{a}_{\underline{d}}^{\varepsilon*}$ is again defined by the equations (1):

$$B_l A_l + A'_{l+1} B_l + p_{l+1} q_l = 0, \quad l = 0, \dots, \frac{N}{2} - 1.$$

Let $\mathfrak{o}(V_l) \subset \mathfrak{gl}(V_l) \subset \mathfrak{a}_{\underline{d}}^\varepsilon$ for $l \in I_1$ be the orthogonal Lie subalgebra and let $\pi : \mathfrak{a}_{\underline{d}}^{\varepsilon*} \rightarrow \bigoplus_{l \in I_0} \mathfrak{gl}(V_l)_{\text{diag}}^* \oplus \bigoplus_{l \in I_1} \mathfrak{o}(V_l)^*$ be the projection. Then the symplectic Drinfeld Zastava space $\mathfrak{Z}_{\underline{d}}^\varepsilon$ is identified (as a Poisson variety) with the Hamiltonian reduction:

$$S_{\underline{d}}^\varepsilon // \bigoplus_{l \in I_0} \mathfrak{gl}(V_l)_{\text{diag}} \oplus \bigoplus_{l \in I_1} \mathfrak{o}(V_l) = (\pi^{-1}(0) \cap S_{\underline{d}}^\varepsilon) // \prod_{l \in I_0} GL(V_l)_{\text{diag}} \times \prod_{l \in I_1} O(V_l).$$

We denote the group $\prod_{l \in I_0} GL(V_l)_{\text{diag}} \times \prod_{l \in I_1} O(V_l)$ simply by $G_{\underline{d}}$, and the corresponding Lie algebra $\bigoplus_{l \in I_0} \mathfrak{gl}(V_l)_{\text{diag}} \oplus \bigoplus_{l \in I_1} \mathfrak{o}(V_l)$ by $\mathfrak{g}_{\underline{d}}$.

3.4 Quantum

The natural quantization of the coordinate ring $\mathbb{C}[\mathfrak{a}_{\underline{d}}^{\varepsilon*}]$ is the enveloping algebra $U(\mathfrak{a}_{\underline{d}}^\varepsilon)$. It will be convenient to gather the generators of $U(\mathfrak{a}_{\underline{d}}^\varepsilon)$ (i.e. the basis elements of the Lie algebra $\mathfrak{a}_{\underline{d}}^\varepsilon$) into the following $U(\mathfrak{a}_{\underline{d}}^\varepsilon)$ -valued matrices:

$$A_k, B_k, q_k, A'_l, p_l, \quad 0 \leq k < \frac{N}{2}, \quad 0 < l \leq \frac{N}{2}.$$

According to [4] the coefficients of the following matrices form a subspace $R \subset U(\mathfrak{a}_{\underline{d}}^\varepsilon)$ invariant with respect to the adjoint action:

$$B_l A_l + A'_{l+1} B_l + p_{l+1} q_l, \quad l = 0, \dots, \frac{N}{2} - 1, \quad i = 1, \dots, d_{l+1}, \quad j = 1, \dots, d_l. \quad (2)$$

Equivalently, $U(\mathfrak{a}_{\underline{d}}^\varepsilon)R$ is a two-sided ideal in $U(\mathfrak{a}_{\underline{d}}^\varepsilon)$.

The natural quantization of the coordinate ring of the space $\mathfrak{Z}_{\underline{d}}^\varepsilon$ is the *quantum Hamiltonian reduction* $\mathcal{Y}_{\underline{d}}^\varepsilon := \left(U(\mathfrak{a}_{\underline{d}}^\varepsilon) / U(\mathfrak{a}_{\underline{d}}^\varepsilon)(R + \mathfrak{g}_{\underline{d}}) \right)^{G_{\underline{d}}}$. The ring $\mathcal{Y}_{\underline{d}}^\varepsilon$ has a natural filtration coming from the PBW filtration on $U(\mathfrak{a}_{\underline{d}}^\varepsilon)$.

PROPOSITION 3.5 PBW property. We have $\text{gr } \mathcal{Y}_{\underline{d}}^\varepsilon = \mathbb{C}[\mathfrak{Z}_{\underline{d}}^\varepsilon]$.

Proof. The proof is a word-to-word repetition of that of Proposition 3.28 from [4]. \square

We consider the following elements of $\mathcal{Y}_{\underline{d}}^\varepsilon$:

$$a_{l,r} := \text{Tr } A_l^r, \quad r = 1, 2, \dots, \quad l \in I;$$

$$b_{l,s} := q_l A_l^s p_l, \quad s = 0, 1, \dots, \quad l \in I.$$

We also introduce the following elements:

$$b_{k,l;s_k,\dots,s_l} := q_l A_l^{s_l} B_{l-1} A_{l-1}^{s_{l-1}} B_{l-2} \dots B_k A_k^{s_k} p_k, \quad k \leq l \in \mathbb{Z}, \quad s_i \in \mathbb{Z}_{\geq 0}. \quad (3)$$

$$c_{k,l;s_k,\dots,s_l} := B_l A_l^{s_l} B_{l-1} A_{l-1}^{s_{l-1}} B_{l-2} \dots B_k A_k^{s_k}, \quad l = k + mN, \quad s_i \in \mathbb{Z}_{\geq 0}. \quad (4)$$

From the definitions we get the following relations:

LEMMA 3.6. *Let $k < l + 1$. Then*

$$[b_{k,l;0,\dots,0}, b_{l+1,0}] = \begin{cases} b_{k,l+1;0,\dots,0} & l \pm k \neq 0, 2 \pmod{N}, \quad 2l + 2 \neq 0 \pmod{N} \\ 2b_{k,l+1;0,\dots,0} & 2l + 2 = 0 \pmod{N} \\ b_{k,l+1;0,\dots,0} - b_{k-1,l;0,\dots,0} & l + k = 0 \pmod{N} \\ 2(b_{k,l+1;0,\dots,0} - b_{k-1,l;0,\dots,0}) & l + k = 0 \pmod{N} \text{ and } 2l + 2 = 0 \pmod{N} \\ b_{k,l+1;0,\dots,0} - b_{k-1,l;0,\dots,0} & l - k = 2 \pmod{N} \end{cases}$$

LEMMA 3.7. *For $k \leq m \leq l$, we have $[a_{m,r}, b_{k,l;s_k,\dots,s_l}] = \lambda b_{k,l;s_k,\dots,s_{m+r-1},\dots,s_l} + L$, where $\lambda \in \mathbb{C} \setminus \{0\}$, $L \in \mathcal{Y}_{\underline{d}}^\varepsilon$ is expressed in $b_{k',l';s_{k'},\dots,s_{l'}}$ with $l' - k' < l - k$, and $\deg L \leq \deg b_{k,l;s_k,\dots,s_{m+r-1},\dots,s_l}$ with respect to the PBW filtration.*

Proof. Straightforward. □

LEMMA 3.8. (C_1 case) *Let p_0, p_1, A_0, A_1, B be the (matrices of) generators of the algebra $\mathfrak{a}_{\underline{d}}$ for $N = 2$. Then the algebra $\mathcal{Y}_{\underline{d}}^\varepsilon$ is generated by $a_{l,r} := \text{Tr } A_l^r$, $b_{l,0} := p_l^* p_l$ with $l = 0, 1, r = 1, \dots, d_l$.*

Proof. According to Lemma 3.7, it suffices to check that the invariants of the form $p_0^*(B^*B)^m p_0$, $p_1^*B(B^*B)^m p_0$, $p_0^*(B^*B)^m B^* p_1$, $p_1^*(BB^*)^m p_0$ and $\text{Tr}(BB^*)^m$ can be expressed in $a_{l,r}, b_{l,0}$. This is easily checked by induction on m . □

PROPOSITION 3.9. *The algebra $\mathcal{Y}_{\underline{d}}^\varepsilon$ is generated by $a_{l,r}, b_{l,s}$ with $l \in I$, $r = 1, \dots, d_l$, $s = 0, \dots, d_l - 1$.*

Proof. Arguing in the same way as in Proposition 3.35 of [4] we reduce the problem to expressing $b_{k,l;0,0,\dots,0}$ and $c_{0,mN;0,\dots,0}$ via $a_{l,r}, b_{l,s}$.

By Lemma 3.6 for $l - k < N - 1$ we have $b_{k,l;0,0,\dots,0} = \lambda[\dots [b_{k,0} b_{k+1,0}] \dots, b_{l-1,0}], b_{l,0}]$, where λ is a nonzero number. Thus $b_{k,l;0,0,\dots,0}$ with $l - k < n$ are expressed via $a_{l,r}, b_{l,s}$. Suppose that $b_{k,k+mN-1;0,0,\dots,0}$ for some $m \in \mathbb{Z}_+$ is expressed via $a_{l,r}, b_{l,s}$, then for $l - k < N$ we have $b_{k,l+mN;0,0,\dots,0} = \lambda[\dots [b_{k,k+mN-1;0,0,\dots,0} b_{k+N+1,0}] \dots, b_{l+N-1,0}], b_{l+N,0}]$. Thus $b_{k,l+mN;0,0,\dots,0}$ with $l - k < N$ are expressed via $a_{l,r}, b_{l,s}$ as well. So, the problem reduces to expressing $b_{k,k+mN-1;0,\dots,0,0}$ and $c_{0,mN;0,\dots,0}$ via $a_{l,r}, b_{l,s}$.

Let $\underline{D} = (d_0, d_{\frac{N}{2}})$. Define the homomorphism $\Phi : U(\mathfrak{a}_{\underline{D}}^\varepsilon) \rightarrow U(\mathfrak{a}_{\underline{d}}^\varepsilon)$ as

$$\Phi(A_0) = A_0, \quad \Phi(A_1) = A_{\frac{N}{2}}, \quad \Phi(B) = B_{\frac{N}{2}-1} \cdot B_{\frac{N}{2}-2} \dots B_0, \quad \Phi(p_0) = p_0, \quad \Phi(p_1) = B_{\frac{N}{2}-1} \dots B_1 p_1.$$

Note that $\Phi(\mathcal{Y}_{\underline{D}}^\varepsilon) \subset \mathcal{Y}_{\underline{d}}^\varepsilon$. By Lemma 3.8, $\mathcal{Y}_{\underline{D}}^\varepsilon$ is generated by the elements $a_{l,r} := \text{Tr } A_l^r$, $b_{l,0} := p_l^* p_l$ with $l = 0, 1, r = 1, \dots, d_l$. We have $\Phi(a_{0,r}) = a_{0,r}$, $\Phi(a_{1,r}) = a_{\frac{N}{2},r}$, $\Phi(b_{0,0}) = b_{0,0}$,

$\Phi(b_{1,0}) = b_{1,N-1;0,\dots,0}$. Thus everything from $\Phi(\mathcal{Y}_D^\varepsilon)$ is expressed via $a_{l,r}, b_{l,s}$. On the other hand, $\Phi(b_{0,2m-1;0,\dots,0}) = b_{0,mN-1;0,\dots,0}$ and $\Phi(c_{0,2m;0,\dots,0}) = c_{0,mN;0,\dots,0}$. \square

4. Yangians

4.1 Yangian of \mathfrak{sp}_N

Let $(c_{kl})_{k,l=1,2,\dots,\frac{N}{2}}$ stand for the symmetrized Cartan matrix of \mathfrak{sp}_N . That is $c_{kk} = 4$ for $k = \frac{N}{2}$; $c_{kk} = 2$ for $0 < k < \frac{N}{2}$; $c_{kl} = 0$ for $|k-l| > 1$; $c_{kl} = -1$ for $0 < k, l < \frac{N}{2}$ and $l = k \pm 1$; $c_{kl} = -2$ otherwise.

The Yangian $Y(\mathfrak{sp}_N)$ is generated by $\mathbf{x}_{k,r}^\pm, \mathbf{h}_{k,r}$, $k = 1, 2, \dots, \frac{N}{2}$, $r \in \mathbb{N}$, with the following relations:

$$[\mathbf{h}_{k,r}, \mathbf{h}_{l,s}] = 0, \quad [\mathbf{h}_{k,0}, \mathbf{x}_{l,s}^\pm] = \pm c_{kl} \mathbf{x}_{l,s}^\pm, \quad (5)$$

$$2[\mathbf{h}_{k,r+1}, \mathbf{x}_{l,s}^\pm] - 2[\mathbf{h}_{k,r}, \mathbf{x}_{l,s+1}^\pm] = \pm c_{kl} (\mathbf{h}_{k,r} \mathbf{x}_{l,s}^\pm + \mathbf{x}_{l,s}^\pm \mathbf{h}_{k,r}), \quad (6)$$

$$[\mathbf{x}_{k,r}^+, \mathbf{x}_{l,s}^-] = \delta_{kl} \mathbf{h}_{k,r+s}, \quad (7)$$

$$2[\mathbf{x}_{k,r+1}^\pm, \mathbf{x}_{l,s}^\pm] - 2[\mathbf{x}_{k,r}^\pm, \mathbf{x}_{l,s+1}^\pm] = \pm c_{kl} (\mathbf{x}_{k,r}^\pm \mathbf{x}_{l,s}^\pm + \mathbf{x}_{l,s}^\pm \mathbf{x}_{k,r}^\pm), \quad (8)$$

$$[\mathbf{x}_{k,r}^\pm, [\mathbf{x}_{k,p}^\pm, \mathbf{x}_{l,s}^\pm]] + [\mathbf{x}_{k,p}^\pm, [\mathbf{x}_{k,r}^\pm, \mathbf{x}_{l,s}^\pm]] = 0, \quad k = l \pm 1, \quad k \in I, l \in I_0, \quad \forall p, r, s \in \mathbb{N}. \quad (9)$$

$$\sum_{\sigma \in S_3} [\mathbf{x}_{k,r_{\sigma(3)}}^\pm, [\mathbf{x}_{k,r_{\sigma(2)}}^\pm, [\mathbf{x}_{k,r_{\sigma(1)}}^\pm, \mathbf{x}_{l,s}^\pm]]] = 0, \quad k = l \pm 1, \quad k \in I, l \in I_1, \quad \forall r_1, r_2, r_3, s \in \mathbb{N}. \quad (10)$$

We will consider the ‘‘Borel subalgebra’’ \mathcal{Y}^ε of the Yangian, generated by $\mathbf{x}_{k,r}^+$ and $\mathbf{h}_{k,r}$. For a formal variable u we introduce the generating series $\mathbf{h}_k(u) := 1 + \sum_{r=0}^\infty \mathbf{h}_{k,r} u^{-r-1}$; $\mathbf{x}_k^+(u) := \sum_{r=0}^\infty \mathbf{x}_{k,r}^+ u^{-r-1}$.

We also consider a bigger algebra $\mathcal{D}\mathcal{Y}^\varepsilon$, the ‘‘Borel subalgebra of the Yangian double’’, generated by all Fourier components of the series $\mathbf{h}_k(u) := 1 + \sum_{r=0}^\infty \mathbf{h}_{k,r} u^{-r-1}$; $\mathbf{x}_k^+(u) := \sum_{r=-\infty}^\infty \mathbf{x}_{k,r}^+ u^{-r-1}$ (i.e. the generating series $\mathbf{x}_k^+(u)$ are infinite in both positive and negative directions) with the defining relations (5,6,8,9,10). The algebra \mathcal{Y}^ε is then the subalgebra generated by negative Fourier components of $\mathbf{x}_k^+(u)$ and $\mathbf{h}_k(u)$ due to PBW property of the Yangians. We can then rewrite the equations (6,8) in the following form

$$\mathbf{h}_k(u) \mathbf{x}_l^+(v) \frac{2u - 2v - c_{kl}}{2u - 2v + c_{kl}} = \mathbf{x}_l^+(v) \mathbf{h}_k(u). \quad (11)$$

$$\mathbf{x}_k^+(u) \mathbf{x}_l^+(v) (2u - 2v - c_{kl}) = (2u - 2v + c_{kl}) \mathbf{x}_l^+(v) \mathbf{x}_k^+(u). \quad (12)$$

The function $\frac{2u-2v-c_{kl}}{2u-2v+c_{kl}}$ here is understood as a formal power series in u^{-1} , v^{-1} , $u^{-1}v$, hence the equation (11) is well-defined.

Following [4], we will use a little bit different generators of the Cartan subalgebra of the Yangian,

$$\mathbf{A}_k(u) := u^{d_k} + A_{k,0}u^{d_k-1} + \dots + A_{k,r}u^{d_k-r-1} + \dots, \quad (13)$$

obtained as the (unique) solution of the system of functional equations:

$$\mathbf{h}_k(u) = \mathbf{A}_k(u + \frac{1}{2})^{-1} \mathbf{A}_k(u - \frac{1}{2})^{-1} \mathbf{A}_{k-1}(u) \mathbf{A}_{k+1}(u) (u + \frac{1}{2})^{d_k} (u - \frac{1}{2})^{d_k} u^{-d_{k-1}} u^{-d_{k+1}}, \quad (14)$$

for $k = 1, 2, \dots, \frac{N}{2} - 1$, and

$$\mathbf{h}_{\frac{N}{2}}(u) = \mathbf{A}_{\frac{N}{2}}(u+1)^{-1} \mathbf{A}_{\frac{N}{2}}(u-1)^{-1} \mathbf{A}_{\frac{N}{2}-1}(u) \mathbf{A}_{\frac{N}{2}-1}(u+\frac{1}{2})(u+1)^{\frac{d_N}{2}} (u-1)^{\frac{d_N}{2}} u^{-d_{\frac{N}{2}-1}} (u+\frac{1}{2})^{-d_{\frac{N}{2}-1}}. \quad (15)$$

Here we take $\mathbf{A}_0(u) = 1$

LEMMA 4.2. *The generators $\mathbf{A}_k(u)$ of \mathcal{DY}^ε satisfy the relations*

$$\mathbf{A}_k(u) \mathbf{x}_l^+(v) \frac{2u - 2v + \frac{c_{kk}\delta_{kl}}{2}}{2u - 2v - \frac{c_{kk}\delta_{kl}}{2}} = \mathbf{x}_l^+(v) \mathbf{A}_k(u). \quad (16)$$

LEMMA 4.3. *Let $\mathbf{A}_k(u)$ and $\mathbf{x}_l^+(u)$ be the generating series of \mathcal{DY}^ε . Then the series*

$$\mathbf{a}_k(u) = \frac{\mathbf{A}_k(u - \frac{c_{kk}}{4})}{\mathbf{A}_k(u + \frac{c_{kk}}{4})} = 1 - d_k u^{-1} - \sum_{r=1}^{\infty} \mathbf{a}_{k,r} u^{-r-1}, \quad \mathbf{x}_l^+(u)$$

satisfies the following commutator relations

$$[\mathbf{a}_k(u), \mathbf{x}_l^+(v)](u-v) = -\frac{c_{kk}^2 \delta_{kl}}{u-v} \mathbf{x}_l^+(v) \mathbf{a}_k(u), \quad [\mathbf{a}_k(u), \mathbf{a}_l(v)] = 0. \quad (17)$$

The series $\mathbf{a}_k(u)$, $\mathbf{x}_l^+(u)$ generate \mathcal{DY}^ε with the defining relations (17), (8) and (9), and their negative Fourier components generate \mathcal{Y}^ε .

Proof. For $k \neq l$ the relation is obvious, for $k = l$ we have

$$\mathbf{a}_k(u) \mathbf{x}_k^+(v) \frac{u - \frac{c_{kk}}{4} - v + \frac{c_{kk}}{4}}{u - \frac{c_{kk}}{4} - v - \frac{c_{kk}}{4}} \cdot \frac{u + \frac{c_{kk}}{4} - v - \frac{c_{kk}}{4}}{u + \frac{c_{kk}}{4} - v + \frac{c_{kk}}{4}} = \mathbf{x}_k^+(v) \mathbf{a}_k(u).$$

therefore

$$\mathbf{a}_k(u) \mathbf{x}_k^+(v) \frac{(u-v)^2}{(u-v)^2 - \frac{c_{kk}^2}{4}} = \mathbf{x}_k^+(v) \mathbf{a}_k(u).$$

One can inductively express $\mathbf{A}_{k,r}$ via $\mathbf{a}_{k,s}$ with $s \leq r+1$, hence \mathcal{DY}^ε is generated by $\mathbf{a}_k(u)$ and $\mathbf{x}_l^+(u)$. On the other hand, the quotient of $\mathbb{C}[\mathbf{a}_{k,r}]_{r=1}^{\infty} \cdot \mathcal{DY}^+$ by the relation (17) is $\mathbb{C}[\mathbf{a}_{k,r}]_{r=1}^{\infty} \otimes \mathcal{DY}^+$ as a filtered vector space. The same argumentation for \mathcal{Y}^ε . Hence the assertion. \square

4.4 Yangian of $\widehat{\mathfrak{sp}}_N$

Let $(c_{kl})_{k,l \in I}$ stand for the symmetrized Cartan matrix of $\widehat{\mathfrak{sp}}_N$. That is $c_{kk} = 4$ for $k = 0$ or $k = \frac{N}{2}$; $c_{kk} = 2$ for $0 < k < \frac{N}{2}$; $c_{kl} = 0$ for $|k-l| > 1$; $c_{kl} = -1$ for $0 < k, l < \frac{N}{2}$ and $l = k \pm 1$; $c_{kl} = -2$ otherwise.

As for the finite case, we will consider the ‘‘affine Borel Yangian’’. This is an associative algebra $\widehat{\mathcal{Y}}^\varepsilon$ generated by the series

$$\mathbf{x}_k^+(u) := 1 + \sum_{r=0}^{\infty} \mathbf{x}_{k,r} u^{-r-1}, \quad (18)$$

$$\mathbf{A}_k(u) := u^{d_k} + \sum_{r=0}^{\infty} \mathbf{A}_{k,r} u^{d_k-r-1}, \quad (19)$$

with $k \in \mathbb{Z}$ subject to the relations

$$\mathbf{A}_k(u)\mathbf{A}_l(v) = \mathbf{A}_l(v)\mathbf{A}_k(u), \quad (20)$$

$$\mathbf{x}_k^\pm(u)\mathbf{x}_l^\pm(v)(2u - 2v \mp c_{kl}) = \mathbf{x}_l^\pm(v)\mathbf{x}_k^\pm(u)(2u - 2v \pm c_{kl}), \quad (21)$$

where (c_{kl}) stands for the symmetrized Cartan matrix of \widetilde{C}_n ;

$$\mathbf{A}_k(u)\mathbf{x}_l^+(v) \frac{2u - 2v + \frac{c_{kl}\delta_{kl}}{2}}{2u - 2v - \frac{c_{kl}\delta_{kl}}{2}} = \mathbf{x}_l^+(v)\mathbf{A}_k(u), \quad (22)$$

in the sense that negative Fourier components of LHS and RHS are equal, and the Serre relations (9) and (10).

4.5 Symplectic Yangian and symplectic Zastava spaces

THEOREM 4.6. *The algebra $\mathcal{Y}_{\underline{d}}^\varepsilon$ is a quotient of the Borel Yangian $\widehat{\mathcal{Y}}^\varepsilon$ of $\widehat{\mathfrak{sp}}_N$ by some ideal containing $\mathbf{A}_{k,r} = 0$ for $r > d_k$.*

Proof. For $k \in I$, $l \in I_0$, introduce the following generating series in $\mathcal{Y}_{\underline{d}}^\varepsilon$:

$$a_k(u) := 1 - d_k u^{-1} - \sum_{r=1}^{\infty} a_{k,r} u^{-r-1}, \quad b_l(u) := \sum_{s=0}^{\infty} b_{l,s} u^{-s-1} \quad (23)$$

For $l \in I_1, i = 1, 2, \dots, d_l$, introduce the following generating series in $U(\mathfrak{a}_{\underline{d}}^\varepsilon)$ (warning: not in $\mathcal{Y}_{\underline{d}}^\varepsilon$):

$$b_l^{(i)}(u) := \sum_{s=0}^{\infty} b_{l,s}^{(i)} u^{-s-1}, \quad b_{l,s}^{(i)} := (A_l^s p_l)^{(i)}, \quad (24)$$

the i -th coordinate of the vector $A_l^s p_l$ in the orthonormal basis of V_l . The following relations hold (see [4]):

LEMMA 4.7. *The following relations hold:*

$$(u - v)[b_k(u), b_k(v)] = (b_k(u)b_k(v) + b_k(v)b_k(u)) \text{ for } k \in I_0, \quad (25)$$

$$2(u - v)[b_k(u), b_l(v + d_l)] = -(b_k(u)b_l(v + d_l) + b_l(v + d_l)b_k(u)) \text{ for } k, l \in I_0, l = k + 1, \quad (26)$$

$$(u - v)[a_k(u), b_l(v)] = -\frac{\delta_{kl}}{u - v} b_l(v) a_k(u) \text{ for } k \in I, l \in I_0. \quad (27)$$

$$2(u - v)[b_k^{(i)}(u), b_k^{(j)}(v)] = c_{kl}(b_k^{(i)}(u)b_k^{(j)}(v) + b_k^{(j)}(v)b_k^{(i)}(u)) \text{ for } k \in I_1, i, j = 1, \dots, d_l, \quad (28)$$

$$2(u - v)[b_k^{(i)}(u), b_l(v + d_l)] = c_{kl}(b_k^{(i)}(u)b_l(v + d_l) + b_l(v + d_l)b_k^{(i)}(u)) \text{ for } k \in I_1, l = k + 1, i = 1, \dots, d_l, \quad (29)$$

$$2(u-v)[b_k(u), b_l^{(i)}(v+d_l)] = c_{kl}(b_k(u)b_l^{(i)}(v+d_l)+b_l^{(i)}(v+d_l)b_k(u)) \text{ for } l \in I_1, l = k+1, i = 1, \dots, d_l, \quad (30)$$

$$(u-v)[a_k(u), b_l^{(i)}(v)] = -\frac{\delta_{kl}}{u-v}b_l^{(i)}(v)a_k(u) \text{ for } k \in I, l \in I_1, i = 1, \dots, d_l. \quad (31)$$

Proof. This follows from Propositions 3.24 and 3.32 of [4]. \square

LEMMA 4.8. *We have*

$$[b_{k,r_2}, [b_{k,r_1}, b_{l,s}]] + [b_{k,r_1}, [b_{k,r_2}, b_{l,s}]] = 0 \text{ for } k, l \in I_0, |k-l| = 1. \quad (32)$$

$$[b_{k,r_2}, [b_{k,r_1}, b_{l,s}^{(i)}]] + [b_{k,r_1}, [b_{k,r_2}, b_{l,s}^{(i)}]] = 0 \text{ for } k \in I, l \in I_1, |k-l| = 1, i = 1, \dots, d_l, \quad (33)$$

$$[b_{k,r_2}^{(i)}, [b_{k,r_1}^{(j)}, b_{l,s}]] + [b_{k,r_1}^{(j)}, [b_{k,r_2}^{(i)}, b_{l,s}]] = 0 \text{ for } k \in I_1, l \in I, |k-l| = 1, i, j = 1, \dots, d_k. \quad (34)$$

Proof. This follows from Proposition 3.32 of [4]. \square

For $l \in I$, let $D_l(u)$ be the (unique) solution of the functional equation

$$a_l(u) := D_l(u - \frac{1}{2})D_l(u + \frac{1}{2})^{-1}. \quad (35)$$

We have

$$D_l(u)b_k(v)\frac{2u-2v+\delta_{kl}}{2u-2v-\delta_{kl}} = b_k(v)D_l(u) \text{ for } k \in I_0,$$

and

$$D_l(u)b_k^{(i)}(v)\frac{2u-2v+\delta_{kl}}{2u-2v-\delta_{kl}} = b_k^{(i)}(v)D_k(u) \text{ for } k \in I_1.$$

Set $\widetilde{b}_k^{(i)}(u) := D_k(u - \frac{1}{2})^{-1}b_k^{(i)}(u)$. From Lemma 4.7, we have

$$\widetilde{b}_k^{(i)}(u)\widetilde{b}_k^{(j)}(v) = \widetilde{b}_k^{(j)}(v)\widetilde{b}_k^{(i)}(u).$$

Note that the rest of the relations for $b_k^{(i)}$ from lemma 4.7 remain the same for $\widetilde{b}_k^{(i)}$.

For $l \in I_1$, $i, j = 1, \dots, d_l$, set

$$\widetilde{b}_l^{(ij)}(u) := D(u - \frac{1}{2})\widetilde{b}_l^{(i)}(u)\widetilde{b}_l^{(j)}(u+1). \quad (36)$$

Note that

$$\widetilde{b}_l^{(ii)}(u)\widetilde{b}_l^{(jj)}(v)\frac{u-v+2}{u-v-2} = \widetilde{b}_l^{(jj)}(v)\widetilde{b}_l^{(ii)}(u) \quad (37)$$

and

$$\widetilde{b}_l^{(ii)}(u)b_k(v)\frac{u-v+1}{u-v-1} = b_k(v)\widetilde{b}_l^{(ii)}(u) \quad (38)$$

for $|k-l| = 1$.

For $l \in I_1$ set $\widetilde{b}_l(u) = \sum_{i=1}^{d_l} \widetilde{b}_l^{(ii)}(u)$. From Proposition 3.9, we see that the algebra $\mathcal{Y}_{\underline{d}}^{\varepsilon}$ is generated by (Fourier coefficients of) $D_l(u)$, $b_k(u)$ for $k \in I_0$ and $\widetilde{b}_k(u)$ for $k \in I_1$. Now the Theorem reduces to the following

LEMMA 4.9. *There is a homomorphism $\varphi_{\underline{d}} : \widehat{\mathcal{Y}}^\varepsilon \rightarrow \mathcal{Y}_{\underline{d}}^\varepsilon$ sending $\mathbf{A}_k(u)$ to $D_k(u + \sum_{m=1}^k d_m)$ and $\mathbf{x}_l^+(u)$ to $b_l(u + \sum_{m=1}^l d_m)$ for $l \in I_0$ and to $\widetilde{b}_l(u + \sum_{m=1}^l d_m)$ for $l \in I_1$.*

Proof. We need to prove the quadratic and the Serre relations for the elements $D_l(u)$, $b_k(u)$. The quadratic relations follow from the relations (37) and (38). The proof of the Serre relations is entirely similar to that of Proposition 3.32 from [4]. \square

According to the Newton identity (see Theorem 7.1.3 of [11]), we have

$$a_l(u) = \frac{C_l(-u + d_l)}{C_l(-u + d_l - 1)}, \quad (39)$$

where $C_l(u)$ is the Capelli determinant. This means that $D(u) = C(-u + d_l - \frac{1}{2})$. In particular, $D_{l,r} = 0$ for $r > d_l$. \square

CONJECTURE 4.10. $\mathcal{Y}_{\underline{d}}^\varepsilon = \widehat{\mathcal{Y}}^\varepsilon / \{\mathbf{A}_{k,r} \mid r > d_k\}$.

ACKNOWLEDGEMENTS

We are grateful to D. Panyushev and D. Timashev for the help with references.

REFERENCES

- 1 J. Bryan, M. Sanders, *Instantons on S^4 and \overline{CP}^2 , rank stabilization, and Bott periodicity*, Topology **39** (2000), no. 2, 331–352.
- 2 A. Braverman, M. Finkelberg, D. Gaitsgory, *Uhlenbeck spaces via affine Lie algebras*, Progress in Math. **244** (2006), 17–135.
- 3 A. Braverman, M. Finkelberg, *Semi-infinite Schubert varieties and quantum K -theory of flag manifolds*, arXiv:1111.2266.
- 4 M. Finkelberg, L. Rybnikov, *Quantization of Drinfeld Zastava in type A* , arXiv:1009.0676, to appear in Journal of Eur. Math. Soc.
- 5 M. Finkelberg, A. Kuznetsov, N. Markarian, I. Mirković, *A note on a symplectic structure on the space of G -monopoles*, Commun. Math. Phys. **201** (1999), 411–421.
- 6 A. Gerasimov, S. Kharchev, D. Lebedev, S. Oblezin, *On a class of representations of the Yangian and moduli space of monopoles*, Comm. Math. Phys. **260** (2005), 511–525.
- 7 N. Guay, *Affine Yangians and deformed double current algebras in type A* , Advances in Math. **211** (2007), 436–484.
- 8 H. Kraft, C. Procesi, *On the geometry of conjugacy classes in classical groups*, Comment. Math. Helvetici **57** (1982), 539–602.
- 9 B. Kostant, S. Rallis, *Orbits and representations associated with symmetric spaces*, Amer. J. Math. **93** (1971), 753–809.
- 10 J. Kamnitzer, B. Webster, A. Weekes, O. Yacobi, *Yangians and quantizations of slices in the affine Grassmannian*, arXiv:1209.0349.
- 11 A. I. Molev, *Yangians and classical Lie algebras*, Math. Surveys and Monographs **143** AMS, Providence, RI (2007).
- 12 S. Kovács, *private communication*; G. Leuschke, A. Verbitsky, *computations with the package Macaulay2* (2013).

QUANTIZATION OF DRINFELD ZASTAVA IN TYPE C

Michael Finkelberg fnklberg@gmail.com

IMU, IITP, and National Research University Higher School of Economics, Department of Mathematics, 20 Myasnitskaya st, Moscow 101000, Russia

Leonid Rybnikov leo.rybnikov@gmail.com

IITP, and National Research University Higher School of Economics, Department of Mathematics, 20 Myasnitskaya st, Moscow 101000, Russia