

On large deviations of coupled diffusions with time scale separation

Anatolii A. Puhalskii
University of Colorado Denver and
Institute for Problems in Information Transmission

June 28, 2018

Abstract

We consider two Itô equations that evolve on different time scales. The equations are fully coupled in the sense that all of the coefficients may depend on both the “slow” and the “fast” motion and the diffusion terms may be correlated. The diffusion term in the slow process is small. A large deviation principle is obtained for the joint distribution of the slow process and of the empirical process of the fast variable. By projecting on the slow and fast variables, we arrive at new results on large deviations in the averaging framework and on large deviations of the empirical measures of ergodic diffusions, respectively. The proof of the main result relies on the property that exponential tightness implies large deviation relative compactness. The identification of the large deviation rate function is accomplished by analysing the large deviation limit of an exponential martingale.

1 Introduction

Consider the coupled diffusions specified by the stochastic differential equations

$$\begin{aligned} dX_t^\epsilon &= A(X_t^\epsilon, x_t^\epsilon) dt + \sqrt{\epsilon} B(X_t^\epsilon, x_t^\epsilon) dW_t^\epsilon, \\ dx_t^\epsilon &= \frac{1}{\epsilon} a(X_t^\epsilon, x_t^\epsilon) dt + \frac{1}{\sqrt{\epsilon}} b(X_t^\epsilon, x_t^\epsilon) dW_t^\epsilon, \end{aligned} \tag{1.1}$$

where $\epsilon > 0$ is a small parameter. Here, $A(u, x)$, where $u \in \mathbb{R}^n$ and $x \in \mathbb{R}^l$, is an n -vector, $B(u, x)$ is an $n \times k$ -matrix, $a(u, x)$ is an l -vector, $b(u, x)$ is an $l \times k$ -matrix, and $W^\epsilon = (W_t^\epsilon, t \in \mathbb{R}_+)$ is an \mathbb{R}^k -valued standard Wiener process. Accordingly, the stochastic process $X^\epsilon = (X_t^\epsilon, t \in \mathbb{R}_+)$ takes values in \mathbb{R}^n and the stochastic process $x^\epsilon = (x_t^\epsilon, t \in \mathbb{R}_+)$ takes values in \mathbb{R}^l . The processes X^ϵ and x^ϵ are seen to evolve on different time scales in that time for x^ϵ is accelerated by a factor of $1/\epsilon$. In a number of application areas, one is concerned with finding the logarithmic asymptotics of large deviations for the “slow” process X^ϵ as $\epsilon \rightarrow 0$. (As a matter of fact, our interest in this setup has been aroused by an application to optimal portfolio selection.) When no diffusion term is present in the equation for the slow process, this sort of result is usually referred to as “the averaging principle”. For contributions, see Freidlin [18], Veretennikov [50, 52, 57, 58], Feng and Kurtz [17, Section 11], and references therein. A different perspective has been offered by Liptser [28] whose insight was to consider the joint distribution of the slow process and of the empirical process associated with the fast variable. For the case where the processes X^ϵ and x^ϵ are one-dimensional, the coefficients $a(u, x)$ and $b(u, x)$ do not depend on the first variable and the Wiener

processes driving the diffusions can be taken independent, they derived a large deviation principle (LDP) for the pair $(X^\epsilon, \mu^\epsilon)$ and identified the associated large deviation rate function, where μ^ϵ represents the empirical process associated with x_t^ϵ . The large deviation principle for the slow process then follows by projection.

In this paper, we extend the joint LDP in Liptser [28] to the multidimensional case. It is assumed that the process dimensions are arbitrary and that all coefficients may depend on both variables in a continuous fashion, on the time variable in a measurable fashion, and on ϵ . The diffusions driving the slow and the fast processes do not have to be uncorrelated. We prove the large deviation principle for the distribution of $(X^\epsilon, \mu^\epsilon)$ and produce the large deviation rate function. Projections on the first and second coordinates yield LDPs for X^ϵ and μ^ϵ , respectively.

The results in the literature that obtain an LDP and identify the large deviation rate function for X^ϵ , with a nondegenerate diffusion term being present in the first of equations (1.1), concern the time homogeneous case where the diffusion coefficient in the equation for the fast process does not depend on the slow process, Veretennikov [55, 54, 58], Liptser [28], Feng and Kurtz [17, Section 11]. The latter restriction can be removed in the setting of the averaging principle provided the state space of the fast process is compact, Veretennikov [50, 56, 57]. This paper fills in the gaps by tackling a case of fully coupled diffusions in a noncompact state space. In addition, the coefficients may depend on the time variable explicitly. The results cover both the setup with a nondegenerate diffusion term and the setup with no diffusion term in the equation for the slow process. The form of the large deviation rate function for the slow process is new. For the time-homogeneous case, the continuity and nondegeneracy conditions on the coefficients are similar to those in the literature, except that additional smoothness properties are assumed of $b(u, x)$ as a function of x , as it is done in Liptser [28]. In return, we obtain that the probability measures for which the large deviation rate functions are finite must have weakly differentiable densities whose square roots belong to the Sobolev space $\mathbb{W}^{1,2}(\mathbb{R}^l)$. In particular, additional insight is gained into the LDP for the empirical measures of ergodic diffusion processes. On the other hand, the ergodicity requirements on the fast process in the nongradient case are more restrictive than those in some of the literature.

As in Liptser [28], an important part in our approach is played by the property that exponential tightness implies large deviation relative compactness so that once exponential tightness has been shown, establishing that a large deviation limit point is unique concludes an LDP proof. Liptser [28] identifies the large deviation rate function by evaluating limits of the probabilities that the process in question resides in small balls. We use a different device. The general idea is to consider a characterisation of stochastic processes that admits taking the large deviation limit. Such a characterisation may be the property that a certain process be a martingale, it may also arise out of the description of the process dynamic. The large deviation rate function is identified by the limiting relation, cf. Puhalskii [40, 41, 42], Puhalskii and Vladimirov [43]. In this paper, similarly to Puhalskii [40, 41], the large deviation limit is taken in an exponential martingale problem that has the distribution of $(X^\epsilon, \mu^\epsilon)$ as a solution. We then undertake a study of the limit equation. On the one hand, regularity properties of solutions are investigated. That analysis has much in common with and uses the results and methods of the regularity theory of elliptic partial differential equations. On the other hand, the domain of the validity of the equation is expanded. Put together, those tools enable us to show that the equation has a unique solution and to identify that solution.

The rest of the paper is organised as follows. In Section 2, the main results are stated, their implications are discussed, and earlier contributions are given a more detailed consideration. Section 3 outlines the proof strategy. It is implemented in Sections 4–8. The proof is completed in Section 9.

We conclude the introduction by giving a list of notation and conventions adopted in the paper. The blackboard bold font is reserved for topological spaces, the boldface font is used for entities associated with probability. Vectors are treated as column vectors. The Euclidean length of a vector

$x = (x_1, \dots, x_d)$ from \mathbb{R}^d , where $d \in \mathbb{N}$, is denoted by $|x|^T$ stands for the transpose of a matrix or a vector. For a matrix A , $\|A\|$ denotes the operator norm and A^\oplus denotes the Moore-Penrose pseudoinverse, if A is square then $\text{tr}(A)$ represents the trace of A . Given a positive semidefinite symmetric matrix Δ and a matrix z of a suitable dimension, which may be a vector, we define $\|z\|_\Delta^2 = z^T \Delta z$. Derivatives are understood as weak, or Sobolev, derivatives. For the definitions and basic properties, the reader is referred either to Adams and Fournier [1] or to Gilbarg and Trudinger [22]. For an \mathbb{R} -valued function f on \mathbb{R}^d , Df denotes the gradient and D^2f denotes the Hessian matrix of f . If f assumes its values in \mathbb{R}^{d_1} , then Df is the $d \times d_1$ -matrix with entries $\partial f_i / \partial x_j$ and $\text{div} f$ represents the divergence of f , where $d_1 \in \mathbb{N}$. Subscripts may be added to indicate that differentiation is carried out with respect to a specific variable. For instance, for an \mathbb{R} -valued function $f(t, u, x)$, where $u = (u_1, \dots, u_d)$ and $x = (x_1, \dots, x_{d_1})$, $D_x f$ and $D_u f$ refer to gradients in the third and the second variables, respectively, $D_{uu}^2 f$ is the matrix with entries $\partial^2 f / \partial u_i \partial u_j$, $D_{xx}^2 f$ is the matrix with entries $\partial^2 f / \partial x_i \partial x_j$, and $D_{ux}^2 f$ is the matrix with entries $\partial^2 f / \partial u_i \partial x_j$. The divergence of a matrix is computed rowwise. If $q > 1$, we will denote by q' the conjugate: $q' = q/(q-1)$. We use standard notation for spaces of differentiable functions, e.g., $\mathbb{C}^{1,2}(\Upsilon)$ denotes the space of \mathbb{R} -valued functions that are continuously differentiable once in the first variable and twice in the second variable over a domain Υ in \mathbb{R}^d , $\mathbb{C}_0^{1,2}(\Upsilon)$ is the subspace of $\mathbb{C}^{1,2}(\Upsilon)$ of functions of compact support, $\mathbb{C}_0^1(\Upsilon)$ is the space of continuously differentiable functions of compact support, and $\mathbb{C}_0^\infty(\Upsilon)$ is the space of infinitely differentiable functions of compact support. Given a measurable function $c(x)$ on Υ with values in the set of positive definite symmetric $d \times d$ -matrices and an \mathbb{R}_+ -valued measurable function $m(x)$ on Υ , we will denote by $\mathbb{L}^2(\Upsilon, \mathbb{R}^d, c(x), m(x) dx)$ the Hilbert space of \mathbb{R}^d -valued measurable functions on Υ with the norm $\|f\|_{c(\cdot), m(\cdot)} = (\int_\Upsilon \|f(x)\|_{c(x)}^2 m(x) dx)^{1/2}$. If $c(x)$ is the identity matrix, the notation will be shortened to $\mathbb{L}^2(\Upsilon, \mathbb{R}^d, m(x) dx)$ and to $\mathbb{L}^2(\Upsilon, \mathbb{R}^d)$ if, in addition, $m(x) = 1$. Spaces $\mathbb{L}^2(\Upsilon, m(x) dx)$ and $\mathbb{L}^2(\Upsilon)$ are defined similarly and consist of \mathbb{R} -valued functions. Space $\mathbb{L}^2(\Upsilon, \mathbb{R}^d, \mu(dx))$ is defined via integration with respect to measure μ . Also, standard notation for Sobolev spaces is adhered to, e.g., $\mathbb{W}^{1,2}(\Upsilon)$ is the Hilbert space of \mathbb{R} -valued functions f that possess the first Sobolev derivatives with the norm $\|f\|_{\mathbb{W}^{1,2}(\Upsilon)} = \|f\|_{\mathbb{L}^2(\Upsilon)} + \|Df\|_{\mathbb{L}^2(\Upsilon, \mathbb{R}^d)}$. The local version of a function space, e.g., $\mathbb{W}_{\text{loc}}^{1,2}(\Upsilon)$, consists of functions whose products with arbitrary \mathbb{C}_0^∞ -functions belong to that space, i.e., $\mathbb{W}^{1,2}(\Upsilon)$ in this case, and is endowed with the weakest topology under which the mappings that associate with functions such products are continuous. We let $\mathbb{W}^{1,2}(\Upsilon, m(x) dx)$ denote the set of functions $f \in \mathbb{W}_{\text{loc}}^{1,1}(\Upsilon)$ such that $f \in \mathbb{L}^2(\Upsilon, m(x) dx)$ and $Df \in \mathbb{L}^2(\Upsilon, \mathbb{R}^d, m(x) dx)$ equipped with the norm $\|f\|_{\mathbb{W}^{1,2}(\Upsilon, m(x) dx)} = \|f\|_{\mathbb{L}^2(\Upsilon, m(x) dx)} + \|Df\|_{\mathbb{L}^2(\Upsilon, \mathbb{R}^d, m(x) dx)}$ and let $\mathbb{H}^{1,2}(\Upsilon, m(x) dx)$ denote the closure of the set $\mathbb{C}^\infty(\Upsilon) \cap \mathbb{W}^{1,2}(\Upsilon, m(x) dx)$ in $\mathbb{W}^{1,2}(\Upsilon, m(x) dx)$. Spaces $\mathbb{W}^{1,2}(\Upsilon, c(x), m(x) dx)$ and $\mathbb{H}^{1,2}(\Upsilon, c(x), m(x) dx)$ are defined similarly. We let $\mathbb{L}_0^{1,2}(\Upsilon, \mathbb{R}^d, c(x), m(x) dx)$ represent the closure of the set of the gradients of functions from $\mathbb{C}_0^\infty(\Upsilon)$ in $\mathbb{L}^2(\Upsilon, \mathbb{R}^d, c(x), m(x) dx)$. The space of continuous functions on \mathbb{R}_+ with values in metric space \mathbb{S} is denoted by $\mathbb{C}(\mathbb{R}_+, \mathbb{S})$. It is endowed with the compact-open topology. If function $X = (X_s, s \in \mathbb{R}_+)$ from $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^d)$ is absolutely continuous w.r.t. Lebesgue measure, \dot{X}_s denotes its derivative at s . We let $\mathbb{M}(\mathbb{R}^d)$ (respectively, $\mathbb{M}_1(\mathbb{R}^d)$) represent the set of finite (respectively, probability) measures on \mathbb{R}^d endowed with the weak topology, see, e.g., Topsøe[49]; $\mathbb{P}(\mathbb{R}^d)$ denotes the set of probability densities $m(x)$ on \mathbb{R}^d such that $m \in \mathbb{W}_{\text{loc}}^{1,1}(\mathbb{R}^d)$ and $\sqrt{m} \in \mathbb{W}^{1,2}(\mathbb{R}^d)$. Topological spaces are equipped with Borel σ -algebras, except for \mathbb{R}_+ which is equipped with the Lebesgue σ -algebra, products of topological spaces are equipped with product topologies, and products of measurable spaces are equipped with product σ -algebras. The ‘‘overbar’’ notation is reserved for the closures of sets, $\mathbf{1}_\Gamma$ denotes the indicator function of a set Γ , $[a]$ stands for the integer part of real number a , $a \wedge b = \min(a, b)$, $a \vee b = \max(a, b)$, and $a^+ = a \vee 0$. Notation $U \subset\subset V$, where U and V are open subsets of \mathbb{R}^d , is to signify that the

closure of U is a compact subset of V . Throughout, the conventions that $\inf_{\emptyset} = \infty$ and $0/0 = 0$ are adopted. The terms “absolutely continuous”, “a.e.”, “almost all” refer to Lebesgue measure unless specified otherwise. All suprema in the time variable are understood as essential suprema with respect to Lebesgue measure.

We say that a net of probability measures \mathbf{P}^ϵ , where $\epsilon > 0$, defined on metric space \mathbb{S} obeys the large deviation principle (LDP) with a (tight) large deviation (rate) function \mathbf{I} for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ if \mathbf{I} is a function from \mathbb{S} to $[0, \infty]$ such that the sets $\{z \in \mathbb{S} : \mathbf{I}(z) \leq \delta\}$ are compact for all $\delta \in \mathbb{R}_+$, $\liminf_{\epsilon \rightarrow 0} \epsilon \ln \mathbf{P}^\epsilon(G) \geq -\inf_{z \in G} \mathbf{I}(z)$ for all open sets $G \subset \mathbb{S}$, and $\limsup_{\epsilon \rightarrow 0} \epsilon \ln \mathbf{P}^\epsilon(F) \leq -\inf_{z \in F} \mathbf{I}(z)$ for all closed sets $F \subset \mathbb{S}$. We say that the net \mathbf{P}^ϵ is exponentially tight for rate $1/\epsilon$ if $\inf_K \limsup_{\epsilon \rightarrow 0} (\mathbf{P}^\epsilon(\mathbb{S} \setminus K))^\epsilon = 0$ where K ranges over the collection of compact subsets of \mathbb{S} .

2 Main results

We will consider a time nonhomogeneous version of (1.1) in which the coefficients may depend on ϵ as well:

$$dX_t^\epsilon = A_t^\epsilon(X_t^\epsilon, x_t^\epsilon) dt + \sqrt{\epsilon} B_t^\epsilon(X_t^\epsilon, x_t^\epsilon) dW_t^\epsilon, \quad (2.1a)$$

$$dx_t^\epsilon = \frac{1}{\epsilon} a_t^\epsilon(X_t^\epsilon, x_t^\epsilon) dt + \frac{1}{\sqrt{\epsilon}} b_t^\epsilon(X_t^\epsilon, x_t^\epsilon) dW_t^\epsilon. \quad (2.1b)$$

As above, $A_t^\epsilon(u, x)$ is an n -vector, $B_t^\epsilon(u, x)$ is an $n \times k$ -matrix, $a_t^\epsilon(u, x)$ is an l -vector, $b_t^\epsilon(u, x)$ is an $l \times k$ -matrix, and $W^\epsilon = (W_t^\epsilon, t \in \mathbb{R}_+)$ is an \mathbb{R}^k -valued standard Wiener process. The stochastic process $X^\epsilon = (X_t^\epsilon, t \in \mathbb{R}_+)$ takes values in \mathbb{R}^n and the stochastic process $x^\epsilon = (x_t^\epsilon, t \in \mathbb{R}_+)$ takes values in \mathbb{R}^l . We assume that the functions $A_t^\epsilon(u, x)$, $a_t^\epsilon(u, x)$, $B_t^\epsilon(u, x)$, and $b_t^\epsilon(u, x)$ are measurable and locally bounded in (t, u, x) and are such that the equations (2.1a) and (2.1b) admit a weak solution (X^ϵ, x^ϵ) with trajectories in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n \times \mathbb{R}^l)$ for every initial condition $(X_0^\epsilon, x_0^\epsilon)$. More specifically, we assume that there exists a complete probability space $(\Omega^\epsilon, \mathcal{F}^\epsilon, \mathbf{P}^\epsilon)$ with filtration $\mathbf{F}^\epsilon = (\mathcal{F}_t^\epsilon, t \in \mathbb{R}_+)$ such that $(W_t^\epsilon, t \in \mathbb{R}_+)$ is a Wiener process relative to \mathbf{F}^ϵ , the processes $X^\epsilon = (X_t^\epsilon, t \in \mathbb{R}_+)$ and $x^\epsilon = (x_t^\epsilon, t \in \mathbb{R}_+)$ are \mathbf{F}^ϵ -adapted, have continuous trajectories, and the relations (2.1a) and (2.1b) hold for all $t \in \mathbb{R}_+$ \mathbf{P}^ϵ -a.s. (To ensure uniqueness which we do not assume apriori, one may require, in addition to the above hypotheses, that the coefficients be Lipschitz continuous.) For background information, see Ethier and Kurtz [15], Ikeda and Watanabe [24], Stroock and Varadhan [48]. We note that since the dimensions n , k , and l are arbitrary, the assumption that both X^ϵ and x^ϵ are driven by the same Wiener process does not constitute a loss of generality.

Let us denote $C_t^\epsilon(u, x) = B_t^\epsilon(u, x)B_t^\epsilon(u, x)^T$ and $c_t^\epsilon(u, x) = b_t^\epsilon(u, x)b_t^\epsilon(u, x)^T$. We introduce the boundedness and growth conditions that for all $N > 0$ and $t > 0$

$$\limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq N} \|c_s^\epsilon(u, x)\| < \infty, \quad (2.2a)$$

$$\limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq N} |A_s^\epsilon(u, x)| < \infty, \quad (2.2b)$$

$$\limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n} \frac{u^T A_s^\epsilon(u, x)}{1 + |u|^2} < \infty, \quad (2.2c)$$

and

$$\limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n} \frac{\|C_s^\epsilon(u, x)\|}{1 + |u|^2} < \infty. \quad (2.2d)$$

We also assume as given "limit coefficients": $A_t(u, x)$ is an n -vector, $B_t(u, x)$ is an $n \times k$ -matrix, $a_t(u, x)$ is an l -vector, and $b_t(u, x)$ is an $l \times k$ -matrix. Let $C_t(u, x) = B_t(u, x)B_t(u, x)^T$ and $c_t(u, x) = b_t(u, x)b_t(u, x)^T$. The following regularity properties will be needed.

Condition 2.1. *The functions $A_t(u, x)$, $B_t(u, x)$, and $b_t(u, x)$ are measurable and are bounded locally in (t, u) and globally in x and are continuous in (u, x) , the function $a_t(u, x)$ is measurable and locally bounded in (t, u, x) and is Lipschitz continuous in x locally uniformly in (t, u) , the functions $a_t(u, x)$ and $c_t(u, x)$ are continuous in u locally uniformly in t and uniformly in x , $c_t(u, x)$ is of class \mathbb{C}^1 in x , with the first partial derivatives being bounded and Lipschitz continuous in x locally uniformly in (t, u) , and $\text{div}_x c_t(u, x)$ is continuous in (u, x) .*

Another set of regularity requirements is furnished by the next condition. We introduce

$$G_t(u, x) = B_t(u, x)b_t(u, x)^T. \quad (2.3)$$

Condition 2.2. *The matrix $c_t(u, x)$ is positive definite uniformly in x and locally uniformly in (t, u) . Either $C_t(u, x) = 0$ for all (t, u, x) and $A_t(u, x)$ is locally Lipschitz continuous in u locally uniformly in t and uniformly in x , or the matrix $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) .*

Finally, certain stability properties will be required: for all $N > 0$ and $t > 0$,

$$\lim_{M \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l: |x| \geq M} \sup_{u \in \mathbb{R}^n: |u| \leq N} a_s^\epsilon(u, x)^T \frac{x}{|x|} = -\infty \quad (2.4a)$$

and

$$\lim_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq N} a_s(u, x)^T \frac{x}{|x|} = -\infty. \quad (2.4b)$$

Let $\mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ represent the subset of $\mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ of functions $\mu = (\mu_t, t \in \mathbb{R}_+)$ such that $\mu_t - \mu_s$ is an element of $\mathbb{M}(\mathbb{R}^l)$ for $t \geq s$ and $\mu_t(\mathbb{R}^l) = t$. It is endowed with the subspace topology and is a complete separable metric space, being closed in $\mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$. The stochastic process $\mu^\epsilon = (\mu_t^\epsilon, t \in \mathbb{R}_+)$, where

$$\mu_t^\epsilon(\Theta) = \int_0^t \mathbf{1}_\Theta(x_s^\epsilon) ds,$$

for $\Theta \in \mathcal{B}(\mathbb{R}^l)$, is a random element of $\mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$. We will regard $(X^\epsilon, \mu^\epsilon)$ as a random element of $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$. It is worth noting that the elements of $\mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ can be also regarded as σ -finite measures on $\mathbb{R}_+ \times \mathbb{R}^l$. We will then use notation $\mu(dt, dx)$ for μ .

Let Γ represent the set of (X, μ) such that the function $X = (X_s, s \in \mathbb{R}_+)$ from $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ is absolutely continuous w.r.t. Lebesgue measure on \mathbb{R}_+ and function $\mu = (\mu_s, s \in \mathbb{R}_+)$ from $\mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, when considered as a measure on $\mathbb{R}_+ \times \mathbb{R}^l$, is absolutely continuous w.r.t. Lebesgue measure on $\mathbb{R}_+ \times \mathbb{R}^l$, i.e., $\mu(ds, dx) = m_s(x) dx ds$, where $m_s(x)$, as a function of x , belongs to $\mathbb{P}(\mathbb{R}^l)$ for almost all s . Given $(X, \mu) \in \Gamma$, we define

$$\begin{aligned} \mathbf{I}'(X, \mu) = & \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) - \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) m_s(x) dx}^2 \right. \\ & + \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \text{div}_x (c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) \right. \\ & \left. \left. - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \right) dx \right) ds. \quad (2.5) \end{aligned}$$

We let $\mathbf{I}'(X, \mu) = \infty$ if $(X, \mu) \notin \Gamma$. It follows, on letting $\lambda = 0$, that if $\mathbf{I}'(X, \mu) < \infty$ then, for $t \in \mathbb{R}_+$,

$$\int_0^t \sup_{h \in \mathcal{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}_x (c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx ds < \infty$$

so that, thanks to $G_s(X_s, x)$ being bounded, for all $\lambda \in \mathbb{R}^n$,

$$\int_0^t \sup_{h \in \mathcal{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}_x (c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx ds < \infty. \quad (2.6)$$

We introduce the following convergence condition.

Condition 2.3. *If $\mathbf{I}'(X, \mu) < \infty$, then there exists a nonincreasing $[0, 1]$ -valued $\mathcal{C}_0^1(\mathbb{R}_+)$ -function $\eta(y)$ that equals 1 for $y \in [0, 1]$ and equals 0 for $y \geq 2$, such that*

$$\int_1^2 \frac{|D\eta(y)|^2}{1 - \eta(y)} dy < \infty, \quad (2.7)$$

and, for arbitrary $t \in \mathbb{R}_+$ and $\lambda \in \mathbb{R}^n$,

$$\begin{aligned} & \lim_{r \rightarrow \infty} \int_0^t \sup_{h \in \mathcal{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}_x (c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) \\ & \quad - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \eta\left(\frac{|x|}{r}\right)^2 dx ds \\ & = \int_0^t \sup_{h \in \mathcal{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}_x (c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) \\ & \quad - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx ds, \quad (2.8) \end{aligned}$$

where $m_s(x) = \mu(ds, dx)/(ds dx)$.

We note that (2.7) is satisfied if $\eta(y) = 1 - e^{-1/(y-1)}$ in a right neighbourhood of 1. The next lemma, whose proof is relegated to the appendix, furnishes a way of verifying Condition 2.3.

Lemma 2.1. *Suppose that Conditions 2.1 and 2.2 hold and that $\mathbf{I}'(X, \mu) < \infty$. If*

$$\int_0^t \int_{\mathbb{R}^l} |a_s(X_s, x)|^2 m_s(x) dx ds < \infty, \quad (2.9)$$

then Condition 2.3 holds.

If either

$$\limsup_{|x| \rightarrow \infty} \sup_{s \in [0, t]} a_s(X_s, x)^T \frac{x}{|x|^2} < 0 \quad (2.10)$$

or there exists real-valued function $\hat{a}_s(x)$ which belongs to $\mathbb{W}_{loc}^{1, q}(\mathbb{R}^l)$ in x , where $q > 2$ and $q \geq l$, such that

$$c_s(X_s, x)^{-1} \left(a_s(X_s, x) - \frac{1}{2} \operatorname{div}_x c_s(X_s, x) \right) = D_x \hat{a}_s(x), \quad (2.11)$$

then (2.9) holds.

We state the main result.

Theorem 2.1. *Let (2.2a)–(2.2d), (2.4a), (2.4b), and Conditions 2.1, 2.2, and 2.3 hold. If the net X_0^ϵ obeys the LDP in \mathbb{R}^n with large deviation function \mathbf{I}_0 for rate $1/\epsilon$ as $\epsilon \rightarrow 0$, the net X_0^ϵ is exponentially tight in \mathbb{R}^l for rate $1/\epsilon$ as $\epsilon \rightarrow 0$, and, for all $t > 0$ and $N > 0$, the convergences*

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l: |x| \leq N} \sup_{u \in \mathbb{R}^n: |u| \leq N} & \left(|A_s^\epsilon(u, x) - A_s(u, x)| + |a_s^\epsilon(u, x) - a_s(u, x)| \right. \\ & \left. + \|B_s^\epsilon(u, x) - B_s(u, x)\| + \|b_s^\epsilon(u, x) - b_s(u, x)\| \right) = 0 \end{aligned} \quad (2.12)$$

hold, then the net $(X^\epsilon, \mu^\epsilon)$ obeys the LDP in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ with large deviation function \mathbf{I} defined as follows:

$$\mathbf{I}(X, \mu) = \begin{cases} \mathbf{I}_0(X_0) + \mathbf{I}'(X, \mu), & \text{if } (X, \mu) \in \Gamma, \\ \infty, & \text{otherwise.} \end{cases}$$

Remark 2.1. Condition 2.3 may be superfluous as far as the validity of Theorem 2.1 is concerned. It is used at the final stage of the proof only, see Theorem 8.1.

Remark 2.2. By Lemma 6.7, $\mathbf{I}(X, \mu) = 0$ provided that a.e.

$$\dot{X}_s = \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx$$

and

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr} (c_s(X_s, x) D^2 p(x)) + Dp(x)^T a_s(X_s, x) \right) m_s(x) dx = 0,$$

where X_0 satisfies the equality $\mathbf{I}_0(X_0) = 0$ and the latter equation holds for all $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$. Consequently, $m_s(\cdot)$ is the invariant density of the diffusion process with the infinitesimal drift $a_s(X_s, \cdot)$ and diffusion matrix $c_s(X_s, \cdot)$.

Remark 2.3. Conditions 2.1 and (2.12) imply that

$$\limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l: |x| \leq N} \sup_{u \in \mathbb{R}^n: |u| \leq N} |a_s^\epsilon(u, x)| < \infty. \quad (2.13)$$

Conditions (2.2a)–(2.2d) and (2.12) imply that

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq N} \|c_s(u, x)\| < \infty, \quad (2.14a)$$

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq N} |A_s(u, x)| < \infty, \quad (2.14b)$$

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n} \frac{u^T A_s(u, x)}{1 + |u|^2} < \infty, \quad (2.14c)$$

and

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n} \frac{\|C_s(u, x)\|}{1 + |u|^2} < \infty. \quad (2.14d)$$

In particular, some of the boundedness requirements in Condition 2.1 are consequences of the other hypotheses of Theorem 2.1.

Remark 2.4. If $c_t(u, x)$ and $C_t(u, x)$ are positive definite uniformly in x and locally uniformly in (t, u) , then, since $b_t(u, x)^T c_t(u, x)^{-1} b_t(u, x)$ is the orthogonal projection operator onto the range of $b_t(u, x)^T$, the condition that $C_t(u, x) - G_t(u, x) c_t(u, x)^{-1} G_t(u, x)^T$ be positive definite uniformly in x and locally uniformly in (t, u) is implied by the following ‘‘angle condition’’: for any bounded region of (t, u) , there exists $\ell \in (0, 1)$ such that $|y_1^T y_2| \leq \ell |y_1| |y_2|$ for all y_1 and y_2 from the ranges of $B_t(u, x)^T$ and $b_t(u, x)^T$, respectively, where x is arbitrary and (t, u) belongs to the region. To put it another way, the condition requires that the angles between the elements of the range of $B_t(u, x)^T$, on the one hand, and the elements of the range of $b_t(u, x)^T$, on the other hand, be bounded away from zero uniformly in x and locally uniformly in (t, u) . It ensures that the processes X^ϵ and x^ϵ are ‘‘sufficiently random’’ in relation to each other. Under that condition, the ranges of $B_t(u, x)^T$ and $b_t(u, x)^T$ do not have common nontrivial subspaces and $k \geq n + l$. On the other hand, if $\|C_t(u, x)\|$ is bounded uniformly in x and locally uniformly in (t, u) , as is the case under the hypotheses of Theorem 2.1 according to (2.14d), the converse is also true: if $C_t(u, x) - G_t(u, x) c_t(u, x)^{-1} G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) , then the angle condition holds.

The solution of the variational problem in (2.5) plays an important part in the proof below, so we proceed with describing it. Let $c(x)$ represent a measurable function defined for $x \in \mathbb{R}^d$ and taking values in the space of positive definite symmetric $d \times d$ -matrices, let $m(x)$ represent a probability density on \mathbb{R}^d , and let S_i represent an open ball of radius i centred at the origin in \mathbb{R}^d , where $d \in \mathbb{N}$ and $i \in \mathbb{N}$. For function $\psi_j \in \mathbb{L}_0^{1,2}(S_j, \mathbb{R}^d, c(x), m(x) dx)$ and $j \geq i$, where $j \in \mathbb{N}$, we let $\pi_{ji} \psi_j$ denote the orthogonal projection of the restriction of ψ_j to S_i onto $\mathbb{L}_0^{1,2}(S_i, \mathbb{R}^d, c(x), m(x) dx)$ in $\mathbb{L}^2(S_i, \mathbb{R}^d, c(x), m(x) dx)$. Thus, the function $\pi_{ji} \psi_j$ is the element of $\mathbb{L}_0^{1,2}(S_i, \mathbb{R}^d, c(x), m(x) dx)$ such that $\int_{S_i} Dp(x)^T c(x) \pi_{ji} \psi_j(x) m(x) dx = \int_{S_i} Dp(x)^T c(x) \psi_j(x) m(x) dx$ for all $p \in \mathbb{C}_0^\infty(S_i)$. We note that if the density $m(x)$ is locally bounded away from zero, then $\pi_{ji} \psi_j$ is a certain gradient: $\pi_{ji} \psi_j = D\chi_{ji}$, where χ_{ji} is the weak solution of the Dirichlet problem $\operatorname{div}(c(x)m(x) D\chi_{ji}(x)) = \operatorname{div}(c(x)m(x)\psi_j(x))$ for $x \in S_i$ with a zero boundary condition (cf. the proof of Lemma 7.3). Since, for $i \leq j \leq k$, $\pi_{ji} \circ \pi_{kj} = \pi_{ki}$, the family $(\mathbb{L}_0^{1,2}(S_j, \mathbb{R}^d, c(x), m(x) dx), \pi_{ji})$ is a projective (or inverse) system in the category of sets. Given a function $\phi \in \mathbb{L}_{\text{loc}}^2(\mathbb{R}^d, \mathbb{R}^d, c(x), m(x) dx)$, the orthogonal projections ϕ_i of the restrictions of ϕ to S_i onto $\mathbb{L}_0^{1,2}(S_i, \mathbb{R}^d, c(x), m(x) dx)$ in $\mathbb{L}^2(S_i, \mathbb{R}^d, c(x), m(x) dx)$ are such that $\pi_{ji} \phi_j = \phi_i$, provided $i \leq j$, so they specify an element of the projective (or inverse) limit of $(\mathbb{L}_0^{1,2}(S_j, \mathbb{R}^d, c(x), m(x) dx), \pi_{ji})$, which we denote by $\Pi_{c(\cdot), m(\cdot)} \phi$. On extending the ϕ_i by zero outside of S_i , one has that, for $i \leq j$, $\|\phi_j\|_{c(\cdot), m(\cdot)}^2 - \|\phi_i\|_{c(\cdot), m(\cdot)}^2 = \|\phi_j - \phi_i\|_{c(\cdot), m(\cdot)}^2$, where the norms are taken in \mathbb{R}^d . Hence, if $\lim_{i \rightarrow \infty} \|\phi_i\|_{c(\cdot), m(\cdot)}^2 < \infty$, then the sequence ϕ_i converges in $\mathbb{L}^2(\mathbb{R}^d, \mathbb{R}^d, c(x), m(x) dx)$ as $i \rightarrow \infty$ and one can identify $\Pi_{c(\cdot), m(\cdot)} \phi$ with the limit, so $\Pi_{c(\cdot), m(\cdot)} \phi \in \mathbb{L}^2(\mathbb{R}^d, \mathbb{R}^d, c(x), m(x) dx)$. It is uniquely specified by the requirements that $\Pi_{c(\cdot), m(\cdot)} \phi \in \mathbb{L}_0^{1,2}(\mathbb{R}^d, \mathbb{R}^d, c(x), m(x) dx)$ and that, for all $p \in \mathbb{C}_0^\infty(\mathbb{R}^d)$,

$$\int_{\mathbb{R}^d} Dp(x)^T c(x) \Pi_{c(\cdot), m(\cdot)} \phi(x) m(x) dx = \int_{\mathbb{R}^d} Dp(x)^T c(x) \phi(x) m(x) dx. \quad (2.15)$$

In particular, if ϕ is an element of $\mathbb{L}^2(\mathbb{R}^d, \mathbb{R}^d, c(x), m(x) dx)$, then $\Pi_{c(\cdot), m(\cdot)} \phi$ is the orthogonal

projection of ϕ onto $\mathbb{L}_0^{1,2}(\mathbb{R}^d, \mathbb{R}^d, c(x), m(x) dx)$. For results on the existence and uniqueness for equation (2.15) when $\Pi_{c(\cdot), m(\cdot)}\phi$ is a gradient, see Pardoux and Veretennikov [35].

In the setting of Theorem 2.1, $d = l$. Since, under the hypotheses of Theorem 2.1, the matrix functions $c_t(u, \cdot)^{-1}G_t(u, \cdot)^T$ are bounded, so the matrix function $\Pi_{c_t(u, \cdot), m(\cdot)}(c_t(u, \cdot)^{-1}G_t(u, \cdot)^T)$, whose columns are the projections of the n columns of $c_t(u, \cdot)^{-1}G_t(u, \cdot)^T$ onto the space $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_t(x), m(x) dx)$, is a well defined element of the space $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^{l \times n}, c_t(u, x), m(x) dx)$ and we denote it by $\Psi_{t, m(\cdot), u}$. We also define

$$Q_{t, m(\cdot)}(u, x) = C_t(u, x) - \|\Psi_{t, m(\cdot), u}(x)\|_{c_t(u, x)}^2. \quad (2.16)$$

The function $Q_{t, m(\cdot)}(u, x)$ assumes values in the space of positive semi-definite $n \times n$ -matrices. If the matrix $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) , then the matrix $\int_{\mathbb{R}^l} Q_{t, m(\cdot)}(u, x)m(x) dx$ is positive definite locally uniformly in (t, u) . We also introduce $\Phi_{t, m(\cdot), u} = \Pi_{c_t(u, \cdot), m(\cdot)}(c_t(u, \cdot)^{-1}(a_t(u, \cdot) - \operatorname{div}_x c_t(u, \cdot)/2))$. Since $a_t(u, \cdot)$ is not necessarily square integrable with respect to $m(x) dx$, the function $\Pi_{c_t(u, \cdot), m(\cdot)}(c_t(u, \cdot)^{-1}(a_t(u, \cdot) - \operatorname{div}_x c_t(u, \cdot)/2))$, as a function of $x \in \mathbb{R}^l$, might not be an element of $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_t(u, x), m(x) dx)$.

For future reference, we note that, according to (2.15), a.e.,

$$\int_{\mathbb{R}^l} Dp(x)^T c_s(u, x) \Phi_{s, m(\cdot), u}(x) m(x) dx = \int_{\mathbb{R}^l} Dp(x)^T (a_s(u, x) - \frac{1}{2} \operatorname{div}_x c_s(u, x)) m(x) dx \quad (2.17a)$$

and

$$\int_{\mathbb{R}^l} Dp(x)^T c_s(u, x) \Psi_{s, m(\cdot), u}(x) m(x) dx = \int_{\mathbb{R}^l} Dp(x)^T G_s(u, x)^T m(x) dx, \quad (2.17b)$$

for all $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$. In addition, (2.17b) extends to Dp representing an arbitrary element of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(u, x), m(x) dx)$. A similar extension property holds for (2.17a), provided $a_s(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(u, x), m(x) dx)$.

Proposition 2.1. *If, under the hypotheses of Theorem 2.1, $\mathbf{I}'(X, \mu) < \infty$, then $\Phi_{s, m_s(\cdot), X_s}$ belongs to the space $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ for almost all s and*

$$\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{D_x m_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx$$

belongs to the range of $\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx$ for almost all s . Furthermore, $\Phi_{s, m_s(\cdot), X_s}(x)$ and $\Psi_{s, m_s(\cdot), X_s}(x)$ are measurable in (s, x) so that in the statement of Theorem 2.1,

$$\begin{aligned} \mathbf{I}(X, \mu) = & \mathbf{I}_0(X_0) + \frac{1}{2} \int_0^\infty \left(\int_{\mathbb{R}^l} \left\| \frac{D_x m_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right\|_{c_s(X_s, x)}^2 m_s(x) dx \right. \\ & \left. + \|\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx \right. \\ & \left. - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{D_x m_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx \right\|_{\left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^\oplus}^2 ds. \end{aligned} \quad (2.18)$$

Remark 2.5. If $m(x)$ is an element of $\mathbb{W}_{\text{loc}}^{1,1}(\mathbb{R}^l)$, then $Dm(x) = 0$ for almost all x on the set where $m(x) = 0$, so we will assume throughout that $Dm(x)/m(x) = 0$ a.e. on that set.

Remark 2.6. The expression on the righthand side of (2.18) serves both the case where $C_t(u, x) = 0$ for all (t, u, x) and $A_t(u, x)$ is locally Lipschitz continuous in u locally uniformly in t and uniformly in x , and the case where $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) . In each of the two cases, however, it simplifies as follows. If $C_t(u, x) = 0$ for all (t, u, x) and $A_t(u, x)$ is locally Lipschitz continuous in u locally uniformly in t and uniformly in x , then $Q_{s,m_s(\cdot)}(u, x) = 0$ for all (s, u, x) , so, in order for $\mathbf{I}(X, \mu)$ to be finite, it is necessary that, a.e.,

$$\dot{X}_s = \int_{\mathbb{R}^l} A_s(X_s, x)m_s(x) dx$$

so that

$$\mathbf{I}(X, \mu) = \mathbf{I}_0(X_0) + \frac{1}{2} \int_0^\infty \int_{\mathbb{R}^l} \left\| \frac{D_x m_s(x)}{2m_s(x)} - \Phi_{s,m_s(\cdot),X_s}(x) \right\|_{c_s(X_s,x)}^2 m_s(x) dx ds. \quad (2.19)$$

If $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) , then the matrix $\int_{\mathbb{R}^l} Q_{s,m_s(\cdot)}(X_s, x)m_s(x) dx$ is invertible, so its pseudo-inverse is the same as the inverse and the range condition in the statement of Proposition 2.1 is superfluous.

Remark 2.7. By Theorem 6.1, in order for $\mathbf{I}(X, \mu)$ to be finite it is necessary that $\int_0^t \int_{\mathbb{R}^l} (|D_x m_s(x)|^2/m_s(x) dx ds + |\Phi_{s,m_s(\cdot),X_s}(x)|^2) dx ds < \infty$ for all $t \in \mathbb{R}_+$.

Remark 2.8. The large deviation function in (2.18) can also be written as

$$\mathbf{I}(X, \mu) = \mathbf{I}_0(X_0) + \frac{1}{2} \int_0^\infty \int_{\mathbb{R}^l} |B_s(X_s, x)^T \hat{\lambda}_s + b_s(X_s, x)^T \hat{g}_s(x)|^2 m_s(x) dx ds,$$

where the pair $(\hat{\lambda}_s, \hat{g}_s(x))$ attains the supremum in (2.5), with \hat{g}_s assuming the role of Dh :

$$\begin{aligned} \hat{\lambda}_s &= \left(\int_{\mathbb{R}^l} Q_{s,m_s(\cdot)}(X_s, x)m_s(x) dx \right)^\oplus \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x)m_s(x) dx \right. \\ &\quad \left. - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{D_x m_s(x)}{2m_s(x)} - \Phi_{s,m_s(\cdot),X_s}(x) \right) m_s(x) dx \right) \end{aligned}$$

and

$$\hat{g}_s(x) = \frac{D_x m_s(x)}{2m_s(x)} - \Phi_{s,m_s(\cdot),X_s}(x) - \Psi_{s,m_s(\cdot),X_s}(x) \hat{\lambda}_s.$$

In the symmetric case where $c_t(u, x)^{-1}(2a_t(u, x) - \text{div}_x c_t(u, x)) = D_x \hat{m}_t(u, x)/\hat{m}_t(u, x)$, for some positive probability density $\hat{m}_t(u, \cdot)$ from $\mathbb{W}_{\text{loc}}^{1,1}(\mathbb{R}^l)$, one can identify $\Phi_{t,m_t(\cdot),u}$ with $D_x \hat{m}_t(u, \cdot)/(2\hat{m}_t(u, \cdot))$. (We note that the diffusion process with the infinitesimal drift coefficient $a_t(u, \cdot)$ and diffusion matrix $c_t(u, \cdot)$ has $\hat{m}_t(u, \cdot)$ as an invariant density.) One can then write the large deviation function in (2.19) by using a Dirichlet form:

$$\mathbf{I}(X, \mu) = \mathbf{I}_0(X_0) + \frac{1}{2} \int_0^\infty \int_{\mathbb{R}^l} \left\| D_x \sqrt{\frac{m_s(x)}{\hat{m}_s(X_s, x)}} \right\|_{c_s(X_s,x)}^2 \hat{m}_s(X_s, x) dx ds,$$

provided $D_x \hat{m}_t(u, \cdot) / \hat{m}_t(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_t(x), m_t(x) dx)$.

Let us look at a one-dimensional example:

$$\begin{aligned} dX_t^\epsilon &= A_t(X_t^\epsilon, x_t^\epsilon) dt + \sqrt{\epsilon} B_t(X_t^\epsilon, x_t^\epsilon) dW_{1,t}^\epsilon, \\ dx_t^\epsilon &= \frac{1}{\epsilon} a_t(X_t^\epsilon, x_t^\epsilon) dt + \frac{1}{\sqrt{\epsilon}} b_t(X_t^\epsilon, x_t^\epsilon) dW_{2,t}^\epsilon, \end{aligned}$$

where all coefficients are scalars and $W_{1,t}^\epsilon$ and $W_{2,t}^\epsilon$ are one-dimensional standard Wiener processes. Assuming that $\mathbf{E}^\epsilon(W_{1,t}^\epsilon W_{2,t}^\epsilon) = \rho t$, where $|\rho| < 1$, this setup can be cast as (2.1a) and (2.1b) with $W_t^\epsilon = (W_{1,t}^\epsilon, W_{3,t}^\epsilon)^T$, $B_t^\epsilon(u, x) = (B_t(u, x), 0)$, and $b_t^\epsilon(u, x) = (\rho b_t(u, x), \sqrt{1 - \rho^2} b_t(u, x))$, where $W_{3,t}^\epsilon$ represents a standard one-dimensional Wiener process that is independent of $W_{1,t}^\epsilon$. If $B_t(u, x)$ is bounded away from zero, the large deviation function in (2.18) takes the form

$$\begin{aligned} \mathbf{I}(X, \mu) &= \mathbf{I}_0(X_0) + \int_0^\infty \left(\frac{1}{8} \int_{\mathbb{R}} \left| \frac{D_x m_s(x)}{m_s(x)} - \frac{D_x \hat{m}_s(X_s, x)}{\hat{m}_s(X_s, x)} \right|^2 b_s(X_s, x)^2 m_s(x) dx \right. \\ &\quad + \frac{1}{2(1 - \rho^2)} \frac{1}{\int_{\mathbb{R}} B_s(X_s, x)^2 m_s(x) dx} \left| \dot{X}_s - \int_{\mathbb{R}} A_s(X_s, x) m_s(x) dx \right. \\ &\quad \left. \left. - \frac{\rho}{2} \int_{\mathbb{R}} B_s(X_s, x) b_s(X_s, x) \left(\frac{D_x m_s(x)}{m_s(x)} - \frac{D_x \hat{m}_s(X_s, x)}{\hat{m}_s(X_s, x)} \right) m_s(x) dx \right|^2 ds \right). \end{aligned}$$

If $B_t(u, x) = 0$, then according to (2.19),

$$\mathbf{I}(X, \mu) = \mathbf{I}_0(X_0) + \frac{1}{8} \int_0^\infty \int_{\mathbb{R}} \left| \frac{D_x m_s(x)}{m_s(x)} - \frac{D_x \hat{m}_s(X_s, x)}{\hat{m}_s(X_s, x)} \right|^2 b_s(X_s, x)^2 m_s(x) dx ds,$$

provided $\dot{X}_s = \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx$ a.e. For the special case that $A_s(u, x)$ and $B_s(u, x)$ do not depend on s , $a_s(u, x)$ and $b_s(u, x)$ do not depend on either s or u , and $\rho = 0$, this large deviation function appears in Liptser [28].

We now project to obtain an LDP for X^ϵ . The device of Lemma 6.5 and the minimax theorem, see, e.g., Theorem 7 on p.319 in Aubin and Ekeland [3], yield the following expression for $\inf_\mu \mathbf{I}(X, \mu)$.

Corollary 2.1. *Under the hypotheses of Theorem 2.1, the net X^ϵ obeys the LDP in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ with large deviation function \mathbf{I}^X defined as follows. If function $X = (X_s, s \in \mathbb{R}_+)$ from $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ is absolutely continuous w.r.t. Lebesgue measure on \mathbb{R}_+ , then*

$$\begin{aligned} \mathbf{I}^X(X) &= \mathbf{I}_0(X_0) + \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T \dot{X}_s - \sup_{m \in \mathbb{P}(\mathbb{R}^l)} \left(\lambda^T \int_{\mathbb{R}^l} A_s(X_s, x) m(x) dx + \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) m(x) dx}^2 \right. \right. \\ &\quad \left. \left. - \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div}_x (c_s(X_s, x) m(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m(x) \right. \right. \right. \right. \\ &\quad \left. \left. \left. - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m(x) \right) dx \right) \right) ds. \end{aligned}$$

Otherwise, $\mathbf{I}^X(X) = \infty$.

If X^ϵ is decoupled from x^ϵ , i.e., $A_t(u, x)$ and $B_t(u, x)$ do not depend on x , then Corollary 2.1 yields the LDP for Itô processes with small diffusions, cf. Freidlin and Wentzell [19]: with $A_s(u, x) = A_s(u)$, $B_s(u, x) = B_s(u)$, and $C_s(u) = B_s(u)B_s(u)^T$,

$$\mathbf{I}^X(X) = \mathbf{I}_0(X_0) + \int_0^\infty \frac{1}{2} \|\dot{X}_s - A_s(X_s)\|_{C_s(X_s)^\oplus}^2 ds,$$

provided $\dot{X}_s - A_s(X_s)$ belongs to the range of $C_s(X_s)$ a.e. and $\mathbf{I}^X(X) = \infty$, otherwise.

If one projects the LDP of Theorem 2.1 on the second variable, then an LDP for μ^ϵ is obtained. In particular, if x^ϵ is decoupled from X^ϵ so that $a_t(u, x)$ and $b_t(u, x)$ do not depend on u , we have the following results on the large deviations of the empirical processes and empirical measures of diffusion processes.

Corollary 2.2. *Suppose that*

$$d\tilde{x}_t^\epsilon = \frac{1}{\epsilon} \tilde{a}_t^\epsilon(\tilde{x}_t^\epsilon) dt + \frac{1}{\sqrt{\epsilon}} \tilde{b}_t^\epsilon(\tilde{x}_t^\epsilon) d\tilde{W}_t^\epsilon,$$

where $\tilde{x}_t^\epsilon \in \mathbb{R}^l$, $\tilde{a}_t^\epsilon(x) \in \mathbb{R}^l$, $\tilde{b}_t^\epsilon(x) \in \mathbb{R}^{l \times k}$, and $\tilde{W}_t^\epsilon \in \mathbb{R}^k$, with the coefficients being locally bounded. Assume that, for all $t \in \mathbb{R}_+$,

$$\begin{aligned} \limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \|\tilde{b}_s^\epsilon(x) \tilde{b}_s^\epsilon(x)^T\| &< \infty, \\ \lim_{M \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l: |x| \geq M} \tilde{a}_s^\epsilon(x)^T \frac{x}{|x|} &= -\infty. \end{aligned}$$

If, for all $t \in \mathbb{R}_+$ and all $N \in \mathbb{R}_+$,

$$\lim_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l: |x| \leq N} (|\tilde{a}_s^\epsilon(x) - \tilde{a}_s(x)| + \|\tilde{b}_s^\epsilon(x) - \tilde{b}_s(x)\|) = 0,$$

the matrix $\tilde{c}_t(x) = \tilde{b}_t(x) \tilde{b}_t(x)^T$ is positive definite uniformly in x and locally uniformly in t , is of class \mathbb{C}^1 in x , with the first partial derivatives being Lipschitz continuous and bounded in x locally uniformly in t , $\tilde{a}_t(x)$ is Lipschitz continuous in x locally uniformly in t , $\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \|\tilde{c}_s(x)\| < \infty$, $\limsup_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \tilde{a}_s(x)^T x / |x|^2 < 0$ for all $t \in \mathbb{R}_+$, and the net x_0^ϵ is exponentially tight in \mathbb{R}^l for rate $1/\epsilon$ as $\epsilon \rightarrow 0$, then the net $\tilde{\mu}^\epsilon$, where $\tilde{\mu}_t^\epsilon(dx) = \int_0^t \mathbf{1}_{dx}(\tilde{x}_s^\epsilon) ds$, obeys the LDP in $\mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ with large deviation function \mathbf{J} defined as follows.

If function $\mu = (\mu_s, s \in \mathbb{R}_+)$ from $\mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, when considered as a measure on $\mathbb{R}_+ \times \mathbb{R}^l$, is absolutely continuous w.r.t. Lebesgue measure on $\mathbb{R}_+ \times \mathbb{R}^l$, i.e., $\mu(ds, dx) = m_s(x) dx ds$, $m_s(x)$, as a function of x , belongs to $\mathbb{P}(\mathbb{R}^l)$ for almost all s , and $\tilde{\Phi}_{s, m_s(\cdot)}$, which represents $\Pi_{\tilde{c}_s(\cdot), m_s(\cdot)}(\tilde{c}_s(\cdot)^{-1}(\tilde{a}_s(\cdot) - \text{div}_x \tilde{c}_s(\cdot)/2))$, is an element of $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, \tilde{c}_s(x), m_s(x) dx)$ for almost all s , then

$$\begin{aligned} \mathbf{J}(\mu) &= \int_0^\infty \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \text{div}_x (\tilde{c}_s(x) m_s(x)) - \tilde{a}_s(x) m_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{\tilde{c}_s(x)}^2 m_s(x) \right) dx ds \\ &= \frac{1}{2} \int_0^\infty \int_{\mathbb{R}^l} \left\| \frac{D_x m_s(x)}{2m_s(x)} - \tilde{\Phi}_{s, m_s(\cdot)}(x) \right\|_{\tilde{c}_s(x)}^2 m_s(x) dx ds. \end{aligned}$$

Otherwise, $\mathbf{J}(\mu) = \infty$.

Corollary 2.3. *Suppose that*

$$dY_t = \check{a}(Y_t) dt + \check{b}(Y_t) d\check{W}_t, \quad Y_0 = 0,$$

where $Y_t \in \mathbb{R}^l$, $\check{a}(x) \in \mathbb{R}^l$, $\check{b}(x) \in \mathbb{R}^{l \times k}$, and $\check{W}_t \in \mathbb{R}^k$, with the coefficients being locally bounded.

If the matrix $\check{c}(x) = \check{b}(x)\check{b}(x)^T$ is uniformly positive definite, $\|\check{c}(x)\|$ is bounded, $\check{c}(\cdot) \in \mathbb{C}^1(\mathbb{R}^l, \mathbb{R}^{l \times l})$, with Lipschitz continuous bounded first partial derivatives, $\check{a}(\cdot)$ is Lipschitz continuous, and $\limsup_{|x| \rightarrow \infty} \check{a}(x)^T x / |x|^2 < 0$, then the empirical measures $(1/t) \int_0^t \mathbf{1}_{dx}(Y_s) ds$ obey the LDP in $\mathbb{M}_1(\mathbb{R}^l)$ for rate t as $t \rightarrow \infty$ with the large deviation function

$$\begin{aligned} \check{\mathbf{J}}(\mu) &= \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div}(\check{c}(x)m(x)) - \check{a}(x)m(x) \right) - \frac{1}{2} \|Dh(x)\|_{\check{c}(x)}^2 m(x) \right) dx \\ &= \frac{1}{2} \int_{\mathbb{R}^l} \left\| \frac{Dm(x)}{2m(x)} - \check{\Phi}_{m(\cdot)}(x) \right\|_{\check{c}(x)}^2 m(x) dx \end{aligned}$$

provided probability measure μ on \mathbb{R}^l has density m , which is an element of $\mathbb{P}(\mathbb{R}^l)$, and $\check{\Phi}_{m(\cdot)} = \Pi_{\check{c}(\cdot), m(\cdot)}(\check{c}(\cdot)^{-1}(\check{a}(\cdot) - \operatorname{div}\check{c}(\cdot)/2))$ is an element of $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, \check{c}(x), m(x) dx)$. Otherwise, $\check{\mathbf{J}}(\mu) = \infty$.

In order to derive Corollary 2.3 from Corollary 2.2, one takes $\epsilon = 1/t$ and defines $\tilde{x}_s^\epsilon = Y_{st}$.

One can thus write the large deviation function of Theorem 2.1 as

$$\begin{aligned} \mathbf{I}(X, \mu) &= \mathbf{I}_0(X_0) + \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) \nu_s(dx)) - \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) \nu_s(dx)}^2 \right. \\ &\quad \left. + \mathbf{J}^{s, X_s, \lambda}(\nu_s) \right) ds, \quad (2.20) \end{aligned}$$

with $\nu_s(dx) = m_s(x) dx$, and the large deviation function of Corollary 2.1 as

$$\begin{aligned} \mathbf{I}^X(X) &= \mathbf{I}_0(X_0) + \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T \dot{X}_s - \sup_{\nu \in \mathbb{M}_1(\mathbb{R}^l)} \left(\lambda^T \int_{\mathbb{R}^l} A_s(X_s, x) \nu(dx) + \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) \nu(dx)}^2 \right. \right. \\ &\quad \left. \left. - \mathbf{J}^{s, X_s, \lambda}(\nu) \right) \right) ds, \quad (2.21) \end{aligned}$$

where $\mathbf{J}^{s, u, \lambda}$ represents the large deviation function for the empirical measures $\nu_t^{s, u, \lambda}(dx) = (1/t) \int_0^t \mathbf{1}_{dx}(y_r^{s, u, \lambda}) dr$ for rate t as $t \rightarrow \infty$ and

$$dy_t^{s, u, \lambda} = (a_s(u, y_t^{s, u, \lambda}) + G_s(u, y_t^{s, u, \lambda})^T \lambda) dt + b_s(u, y_t^{s, u, \lambda}) dw_t, \quad y_0^{s, u, \lambda} = 0,$$

(w_t) being a k -dimensional standard Wiener process. In particular, if $G_t(u, x) = 0$ so that the diffusions driving the slow and the fast processes are virtually uncorrelated, then $\mathbf{J}^{s, u, \lambda}$ does not depend on λ and by Corollary 2.2, Corollary 2.3, and (2.20) the large deviation function $\mathbf{I}(X, \mu)$ is the sum of the large deviation function of the slow process, with the coefficients being averaged over the "current" empirical measure of the fast variable, and of the large deviation function of the empirical process of the fast variable, with the coefficients "frozen" at the current value of the slow variable.

The first results on large deviation asymptotics for the system (1.1) in the setup of the averaging principle available in the literature appear in Freidlin [18], see also the exposition in Freidlin and Wentzell [19, Section 9 of Chapter 7]. Freidlin [18] considers the equations

$$\begin{aligned}\dot{x}_t^\epsilon &= b(x_t^\epsilon, y_t^\epsilon), \\ \dot{y}_t^\epsilon &= \frac{1}{\epsilon} [B(x_t^\epsilon, y_t^\epsilon) + g(y_t^\epsilon)] + \frac{1}{\sqrt{\epsilon}} c(y_t^\epsilon) \dot{w}_t.\end{aligned}$$

It is assumed that the state space is a compact manifold. A noncompact setting is considered by Veretennikov [52]. Veretennikov [50, 56, 57] allows the diffusion coefficient in the fast process to depend on both variables:

$$\begin{aligned}dX_t^\epsilon &= f(X_t^\epsilon, Y_t^\epsilon) dt, \\ dY_t^\epsilon &= \epsilon^{-2} B(X_t^\epsilon, Y_t^\epsilon) dt + \epsilon^{-1} C(X_t^\epsilon, Y_t^\epsilon) dW_t.\end{aligned}$$

The state space of the fast process is a compact manifold.

Veretennikov [54, 55, 58] tackles the case where the slow process has a small diffusion term and the state space of the fast process may be noncompact but the diffusion coefficient in the equation for the fast process does not depend on the slow process so that

$$dX_t^\epsilon = f(X_t^\epsilon, Y_t^\epsilon) dt + \epsilon(\sigma_1(X_t^\epsilon, Y_t^\epsilon) dW_t^1 + \sigma_3(X_t^\epsilon, Y_t^\epsilon) dW_t^3), \quad (2.22)$$

$$dY_t^\epsilon = \epsilon^{-2} B(X_t^\epsilon, Y_t^\epsilon) dt + \epsilon^{-1}(C_1(Y_t^\epsilon) dW_t^1 + C_2(Y_t^\epsilon) dW_t^2),$$

where the Wiener processes are independent. The stability condition on the slow process is similar to (2.4a) and (2.4b).

In those papers, results on the LDP for the slow processes are obtained in the space of continuous functions on the $[0, L]$ interval endowed with uniform norm, where $L > 0$. The large deviation rate functions are of the form

$$\mathbf{I}(X) = \int_0^L \sup_{\lambda} (\lambda^T \dot{X}_t - H(X_t, \lambda)) dt,$$

provided X_t , $t \in [0, L]$, is an absolutely continuous function with a suitable initial condition. Otherwise, $\mathbf{I}(X) = \infty$. Here, with the notation of (2.22),

$$H(u, \lambda) = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \mathbf{E} \exp\left(\int_0^t (\lambda^T f(u, y_s^{u, \lambda}) + \frac{1}{2} \lambda^T (\sigma_1 \sigma_1^T(u, y_s^{u, \lambda}) + \sigma_3 \sigma_3^T(u, y_s^{u, \lambda})) \lambda) ds\right), \quad (2.23)$$

where

$$dy_t^{u, \lambda} = (B(u, y_t^{u, \lambda}) + C_1(y_t^{u, \lambda}) \sigma_1(u, y_t^{u, \lambda})^T \lambda) dt + (C_1(Y_t^{u, \lambda}) dW_t^1 + C_2(Y_t^{u, \lambda}) dW_t^2), \quad y_0^{u, \lambda} = 0.$$

The existence of the limit in (2.23) is proved by invoking the Frobenius theorem for compact positive operators.

Let us note that if one assumes the LDP at rate t as $t \rightarrow \infty$ of the empirical measures $\nu_t^{u, \lambda}(dx) = (1/t) \int_0^t \mathbf{1}_{dx}(y_s^{u, \lambda}) ds$ with large deviation rate function $\mathbf{J}^{u, \lambda}$, then, in view of Varadhan's lemma and (2.23), under suitable assumptions,

$$H(u, \lambda) = \sup_{\nu \in \mathbb{M}_1(\mathbb{R}^l)} \left(\int_{\mathbb{R}^l} (\lambda^T f(u, x) + \frac{1}{2} \lambda^T (\sigma_1 \sigma_1^T(u, x) + \sigma_3 \sigma_3^T(u, x)) \lambda) \nu(dx) - \mathbf{J}^{u, \lambda}(\nu) \right),$$

which is consistent with (2.21).

Section 11.6 of Feng and Kurtz [17] is concerned with the process X^ϵ satisfying equations (1.1). Conditions for the LDP to hold are obtained. They require the existence of functions with certain properties and are not easily translated into conditions on the coefficients. When the authors give explicit conditions on the coefficients, they need, in particular, $b(u, x)$ not to depend on u (see Lemma 11.60 on p.278). The large deviation rate function is identified as having the form (2.21) corresponding to the time-homogeneous setting, provided $B(u, x)b(u, x)^T = 0$ and certain additional hypotheses hold (see Theorem 11.6.5 on p.282). The authors choose not to pursue the setup of the averaging principle.

The LDP for the empirical measures of continuous-time Markov processes, such as in Corollary 2.3, is a well explored subject, see Donsker and Varadhan [11, 12], Deuschel and Stroock [10]. The canonical form of the large deviation rate function is $\sup_f \int_{\mathbb{R}^l} -\mathcal{L}f/f d\mu$, where \mathcal{L} represents the infinitesimal generator of the Markov process, see, e.g., Theorem 4.2.43 in Deuschel and Stroock [10]. The form in Corollary 2.3 follows by taking $f(x) = e^{-h(x)}$. Gärtner [20] and Veretennikov [51] characterise the large deviation functions via limits similar to that in (2.23), the latter author allowing discontinuous coefficients. Theorem 12.7 on p.291 of Feng and Kurtz [17] tackles associated empirical processes, cf. Corollary 2.2.

3 Some generalities

This section contains general results on the LDP that underlie the proof of Theorem 2.1, cf. Puhalskii [41]. Let Σ represent a directed set, let \mathbf{P}_σ , where $\sigma \in \Sigma$, represent a net of probability measures on a metric space \mathbb{S} indexed with the elements of Σ and let r_σ represent an \mathbb{R}_+ -valued function which tends to infinity as $\sigma \in \Sigma$. A $[0, \infty]$ -valued function \mathbf{I} on \mathbb{S} is referred to as a large deviation function if the sets $K_\delta = \{z \in \mathbb{S} : \mathbf{I}(z) \leq \delta\}$ are compact for all $\delta \in \mathbb{R}_+$. We say that the net \mathbf{P}_σ obeys the LDP with a large deviation function \mathbf{I} for rate r_σ as $\sigma \in \Sigma$ if $\liminf_{\sigma \in \Sigma} r_\sigma^{-1} \ln \mathbf{P}_\sigma(G) \geq -\inf_{z \in G} \mathbf{I}(z)$ for all open sets $G \subset \mathbb{S}$ and $\limsup_{\sigma \in \Sigma} r_\sigma^{-1} \ln \mathbf{P}_\sigma(F) \leq -\inf_{z \in F} \mathbf{I}(z)$ for all closed sets $F \subset \mathbb{S}$. We say that \mathbf{I} is a large deviation (LD) limit point of \mathbf{P}_σ for rate r_σ if there exists a subsequence σ_i , where $i \in \mathbb{N}$, such that \mathbf{P}_{σ_i} satisfies the LDP with \mathbf{I} for rate r_{σ_i} as $i \rightarrow \infty$. We say that the net \mathbf{P}_σ is sequentially large deviation (LD) relatively compact for rate r_σ as $\sigma \in \Sigma$ if any subsequence \mathbf{P}_{σ_i} of \mathbf{P}_σ contains a further subsequence $\mathbf{P}_{\sigma_{i_j}}$ which satisfies the LDP for rate $r_{\sigma_{i_j}}$ with some large deviation function as $j \rightarrow \infty$. We say that the net \mathbf{P}_σ is exponentially (or large deviation) tight for rate r_σ as $\sigma \in \Sigma$ if for arbitrary $\kappa > 0$ there exists compact $K \subset \mathbb{S}$ such that $\limsup_{\sigma \in \Sigma} \mathbf{P}_\sigma(\mathbb{S} \setminus K)^{1/r_\sigma} < \kappa$. We say that the net \mathbf{P}_σ is sequentially exponentially tight for rate r_σ as $\sigma \in \Sigma$ if any subsequence \mathbf{P}_{σ_i} is exponentially tight for rate r_{σ_i} as $i \rightarrow \infty$. We say that a net Y_σ of random elements of \mathbb{S} obeys the LDP, respectively, is sequentially LD relatively compact, respectively, is exponentially tight, respectively, is sequentially exponentially tight if the net of their laws has the indicated property.

The cornerstone of our approach is the next result (Puhalskii [36, 37, 38, 41], see also Feng and Kurtz [17] and references therein).

Theorem 3.1. *If the net \mathbf{P}_σ is sequentially exponentially tight for rate r_σ as $\sigma \in \Sigma$, then the net \mathbf{P}_σ is sequentially LD relatively compact for rate r_σ as $\sigma \in \Sigma$.*

The proof of the following theorem is standard.

Theorem 3.2. *If the net \mathbf{P}_σ is sequentially LD relatively compact for rate r_σ as $\sigma \in \Sigma$ and \mathbf{I} is a unique LD limit point of the \mathbf{P}_σ , then the net \mathbf{P}_σ satisfies the LDP with \mathbf{I} for rate r_σ as $\sigma \in \Sigma$.*

The next theorem is essentially Varadhan's lemma, see, e.g., Deuschel and Stroock [10]. It will be used to obtain equations for LD limit points.

Theorem 3.3. *Suppose the net \mathbf{P}_σ is sequentially exponentially tight for rate r_σ as $\sigma \in \Sigma$ and let \mathbf{I} represent an LD limit point of \mathbf{P}_σ . Let U_σ be a net of uniformly bounded real valued functions on \mathbb{S} such that $\int_{\mathbb{S}} \exp(r_\sigma U_\sigma(z)) \mathbf{P}_\sigma(dz) = 1$. If $U_\sigma \rightarrow U$ uniformly on compact sets as $\sigma \in \Sigma$, where the function U is continuous, then $\sup_{z \in \mathbb{S}} (U(z) - \mathbf{I}(z)) = 0$.*

Identification of LD limit points will be carried out with the aid of the next result.

Theorem 3.4. *Suppose \mathbf{I} is a large deviation function on \mathbb{S} and \mathcal{U} is a collection of functions on \mathbb{S} such that $\sup_{z \in \mathbb{S}} (U(z) - \mathbf{I}(z)) = 0$ for all $U \in \mathcal{U}$. Let $\mathbf{I}^{**}(z) = \sup_{U \in \mathcal{U}} U(z)$ and $K_\delta = \{z \in \mathbb{S} : \mathbf{I}(z) \leq \delta\}$, where $\delta \in \mathbb{R}_+$.*

1. *Let $\tilde{\mathcal{U}}$ represent a set of functions U such that $\sup_{z \in K_\delta} (U(z) - \mathbf{I}(z)) = 0$ for suitable $\delta \in \mathbb{R}_+$. Suppose $\hat{z} \in \mathbb{S}$ is such that $\mathbf{I}^{**}(\hat{z}) = \hat{U}(\hat{z})$ for some function $\hat{U} \in \tilde{\mathcal{U}}$. Suppose there exists sequence $U_i \in \tilde{\mathcal{U}}$ with the following properties: $\sup_{z \in K_\delta} (U_i(z) - \mathbf{I}(z)) = 0$ for some common δ , the functions U_i are continuous when restricted to K_δ and if z_i is a convergent sequence of elements of K_δ such that $U_i(z_i) = \mathbf{I}(z_i)$, then $U_i(z_i) \rightarrow \hat{U}(\hat{z})$ and $z_i \rightarrow \hat{z}$ as $i \rightarrow \infty$. Then $\mathbf{I}(\hat{z}) = \mathbf{I}^{**}(\hat{z})$.*
2. *If for every $z \in \mathbb{S}$ such that $\mathbf{I}^{**}(z) < \infty$ there exists a sequence of points z_i such that $\mathbf{I}(z_i) = \mathbf{I}^{**}(z_i)$, $z_i \rightarrow z$, and $\mathbf{I}^{**}(z_i) \rightarrow \mathbf{I}^{**}(z)$ as $i \rightarrow \infty$, then $\mathbf{I}(z) = \mathbf{I}^{**}(z)$ for all $z \in \mathbb{S}$.*

Proof. Let us first note that $\mathbf{I}(z) \geq \mathbf{I}^{**}(z)$ for all z , so, one needs to prove that $\mathbf{I}(z) \leq \mathbf{I}^{**}(z)$ if $\mathbf{I}^{**}(z) < \infty$. We prove part 1. Since $\sup_{z \in K_\delta} (U_i(z) - \mathbf{I}(z)) = 0$, K_δ is compact, and $U_i(z) - \mathbf{I}(z)$ is upper semicontinuous when restricted to K_δ , there exist $z_i \in K_\delta$ such that $U_i(z_i) = \mathbf{I}(z_i)$. One may assume that the sequence converges. Since $U_i(z_i) \rightarrow \hat{U}(\hat{z})$, $z_i \rightarrow \hat{z}$ and \mathbf{I} is lower semicontinuous, $\hat{U}(\hat{z}) \geq \mathbf{I}(\hat{z})$, so $\mathbf{I}^{**}(\hat{z}) \geq \mathbf{I}(\hat{z})$. The proof of part 2 is similar. \square

In the rest of the paper, the above framework is used to prove Theorem 2.1. In Section 4, LD relative compactness is established, see Theorem 4.1. In Section 5, equations along the lines of Theorem 3.3 are derived, see Theorem 5.1. Section 6 is concerned with regularity properties of (X, μ) for which the function \mathbf{I}^{**} as defined in Theorem 3.4 assumes finite values. It is also shown to be of the form given in Proposition 2.1, see Theorem 6.1. In Theorem 7.1 of Section 7, the large deviation function is identified for a large class of (X, μ) , which implements the recipe of part 1 of Theorem 3.4. In Theorem 8.1 of Section 8, it is proved that that class is dense in the sense of part 2 of Theorem 3.4. In Section 9, the proof of Theorem 2.1 is completed.

4 LD relative compactness

The main result of this section is the following theorem.

Theorem 4.1. *Suppose that conditions (2.2a) – (2.2d) and (2.4a) hold and that the net $(X_0^\epsilon, x_0^\epsilon)$ is exponentially tight for rate $1/\epsilon$ as $\epsilon \rightarrow 0$. Then the net $(X^\epsilon, \mu^\epsilon)$ is sequentially LD relatively compact in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ for rate $1/\epsilon$ as $\epsilon \rightarrow 0$.*

We precede the proof with a criterion of sequential LD relative compactness in $\mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$. Let $d(\cdot, \cdot)$ represent the Lipschitz metric on $\mathbb{M}(\mathbb{R}^l)$: $d(\tilde{\mu}, \hat{\mu}) = \sup\{|\int_{\mathbb{R}^l} f(x) \tilde{\mu}(dx) - \int_{\mathbb{R}^l} f(x) \hat{\mu}(dx)|\}$, with the supremum being taken over functions $f : \mathbb{R}^l \rightarrow \mathbb{R}$ such that $\sup_{x \in \mathbb{R}^l} |f(x)| \leq 1$ and $\sup_{x, y \in \mathbb{R}^l, x \neq y} |f(x) - f(y)|/|x - y| \leq 1$, see, e.g., p.395 in Dudley [13]. The proof of the next lemma is relegated to the appendix.

Lemma 4.1. 1. A net $\{\nu_\epsilon, \epsilon > 0\}$, where $\nu_\epsilon = (\nu_{\epsilon,t}, t \in \mathbb{R}_+)$, of random elements of $\mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ defined on respective probability spaces $(\Omega_\epsilon, \mathcal{F}_\epsilon, \mathbf{P}_\epsilon)$ is sequentially exponentially tight for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ if and only if for all $t \in \mathbb{R}_+$ and all $\eta > 0$,

$$\lim_{N \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \mathbf{P}_\epsilon(\nu_{\epsilon,t}(x \in \mathbb{R}^l : |x| > N) > \eta)^\epsilon = 0$$

and

$$\lim_{\delta \rightarrow 0} \limsup_{\epsilon \rightarrow 0} \sup_{s_1 \in [0,t]} \mathbf{P}_\epsilon\left(\sup_{s_2 \in [s_1, s_1 + \delta]} d(\nu_{\epsilon,s_1}, \nu_{\epsilon,s_2}) > \eta\right)^\epsilon = 0.$$

2. A net $\{Y_\epsilon, \epsilon > 0\}$, where $Y_\epsilon = (Y_{\epsilon,t}, t \in \mathbb{R}_+)$, of random elements of $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ defined on respective probability spaces $(\Omega_\epsilon, \mathcal{F}_\epsilon, \mathbf{P}_\epsilon)$ is sequentially exponentially tight for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ if and only if

$$\lim_{N \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \mathbf{P}_\epsilon(|Y_{\epsilon,0}| > N)^\epsilon = 0$$

and, for all $t \in \mathbb{R}_+$ and all $\eta > 0$,

$$\lim_{\delta \rightarrow 0} \limsup_{\epsilon \rightarrow 0} \sup_{s_1 \in [0,t]} \mathbf{P}_\epsilon\left(\sup_{s_2 \in [s_1, s_1 + \delta]} |Y_{\epsilon,s_2} - Y_{\epsilon,s_1}| > \eta\right)^\epsilon = 0.$$

Remark 4.1. The form of the conditions is due to Feng and Kurtz [17].

Proof of Theorem 4.1. Since $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ is a closed subset of $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ and $\mathbf{P}^\epsilon((X^\epsilon, \mu^\epsilon) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))) = 1$, it is sufficient to prove that the net $((X^\epsilon, \mu^\epsilon), \epsilon > 0)$ is sequentially LD relatively compact in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$. By Theorem 3.1, the latter property holds if $(X^\epsilon, \mu^\epsilon)$ is sequentially exponentially tight, which is the case if the nets X^ϵ and μ^ϵ are each sequentially exponentially tight.

We show that the net X^ϵ is sequentially exponentially tight first. By (2.1a) and Itô's lemma, on denoting $g_1(x) = D^2 \ln(1 + |x|^2)$,

$$\begin{aligned} \ln(1 + |X_t^\epsilon|^2) &= \ln(1 + |X_0^\epsilon|^2) + \int_0^t \frac{2(X_s^\epsilon)^T A_s^\epsilon(X_s^\epsilon, x_s^\epsilon)}{1 + |X_s^\epsilon|^2} ds + \frac{\epsilon}{2} \int_0^t \text{tr}(C_s^\epsilon(X_s^\epsilon, x_s^\epsilon) g_1(X_s^\epsilon)) ds \\ &\quad + \sqrt{\epsilon} \int_0^t \frac{2(X_s^\epsilon)^T}{1 + |X_s^\epsilon|^2} B_s^\epsilon(X_s^\epsilon, x_s^\epsilon) dW_s^\epsilon. \end{aligned}$$

Given $N > 0$, let $\tau_N^\epsilon = \inf\{s \in \mathbb{R}_+ : |X_s^\epsilon| \geq N\}$. Since τ_N^ϵ is an \mathbf{F}^ϵ -stopping time and

$$\exp\left(\frac{1}{\sqrt{\epsilon}} \int_0^t \frac{2(X_s^\epsilon)^T}{1 + |X_s^\epsilon|^2} B_s^\epsilon(X_s^\epsilon, x_s^\epsilon) dW_s^\epsilon - \frac{1}{2\epsilon} \int_0^t \left\| \frac{2X_s^\epsilon}{1 + |X_s^\epsilon|^2} \right\|_{C_s^\epsilon(X_s^\epsilon, x_s^\epsilon)}^2 ds\right), t \in \mathbb{R}_+,$$

is an \mathbf{F}^ϵ -local martingale,

$$\begin{aligned} \mathbf{E}^\epsilon \exp\left(\frac{1}{\epsilon} \ln(1 + |X_{t \wedge \tau_N^\epsilon}^\epsilon|^2) - \frac{1}{\epsilon} \ln(1 + |X_0^\epsilon|^2) - \frac{1}{\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \frac{2(X_s^\epsilon)^T A_s^\epsilon(X_s^\epsilon, x_s^\epsilon)}{1 + |X_s^\epsilon|^2} ds \right. \\ \left. - \frac{1}{2} \int_0^{t \wedge \tau_N^\epsilon} \text{tr}(C_s^\epsilon(X_s^\epsilon, x_s^\epsilon) g_1(X_s^\epsilon)) ds - \frac{1}{2\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \left\| \frac{2X_s^\epsilon}{1 + |X_s^\epsilon|^2} \right\|_{C_s^\epsilon(X_s^\epsilon, x_s^\epsilon)}^2 ds\right) \leq 1. \quad (4.1) \end{aligned}$$

Since

$$\operatorname{tr} (C_s^\epsilon(X_s^\epsilon, x)g_1(X_s^\epsilon)) \leq \sqrt{\operatorname{tr} g_1(X_s^\epsilon)^2} \sqrt{\operatorname{tr} C_s^\epsilon(X_s^\epsilon, x)^2} \leq 2n\sqrt{n} \frac{\|C_s^\epsilon(X_s^\epsilon, x_s^\epsilon)\|}{1 + |X_s^\epsilon|^2}, \quad (4.2)$$

on recalling (2.2c) and (2.2d), we have that there exists $L > 0$, which does not depend either on t or on N , such that for all $\epsilon > 0$ small enough,

$$\mathbf{E}^\epsilon \exp\left(\frac{1}{\epsilon} \ln(1 + |X_{t \wedge \tau_N^\epsilon}^\epsilon|^2) - \frac{1}{\epsilon} \ln(1 + |X_0^\epsilon|^2) - \frac{Lt}{\epsilon}\right) \leq 1.$$

For $\tilde{N} > 0$,

$$\begin{aligned} \mathbf{P}^\epsilon(\sup_{s \in [0, t]} |X_s^\epsilon| \geq N) &= \mathbf{P}^\epsilon(|X_{t \wedge \tau_N^\epsilon}^\epsilon| \geq N) \leq \mathbf{P}^\epsilon(|X_0^\epsilon| > \tilde{N}) \\ &+ \mathbf{E}^\epsilon \exp\left(\frac{1}{\epsilon} \ln(1 + |X_{t \wedge \tau_N^\epsilon}^\epsilon|^2) - \frac{1}{\epsilon} \ln(1 + N^2)\right) \mathbf{1}_{\{|X_0^\epsilon| \leq \tilde{N}\}} \\ &\leq \mathbf{P}^\epsilon(|X_0^\epsilon| > \tilde{N}) + \exp\left(\frac{1}{\epsilon} \ln(1 + \tilde{N}^2) + \frac{Lt}{\epsilon} - \frac{1}{\epsilon} \ln(1 + N^2)\right), \end{aligned}$$

so

$$\limsup_{N \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \mathbf{P}^\epsilon(\sup_{s \in [0, t]} |X_s^\epsilon| > N)^\epsilon \leq \limsup_{\epsilon \rightarrow 0} \mathbf{P}^\epsilon(|X_0^\epsilon| > \tilde{N})^\epsilon.$$

Since X_0^ϵ is exponentially tight and \tilde{N} is arbitrary, we conclude that

$$\lim_{N \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \mathbf{P}^\epsilon(\sup_{s \in [0, t]} |X_s^\epsilon| > N)^\epsilon = 0. \quad (4.3)$$

By (2.1a), for $s \in [0, t]$, $\delta > 0$, and $\eta > 0$,

$$\begin{aligned} \mathbf{P}^\epsilon(\sup_{\tilde{s} \in [s, s+\delta]} |X_{\tilde{s}}^\epsilon - X_s^\epsilon| > \eta) &\leq \mathbf{P}^\epsilon(\tau_N^\epsilon \leq t) + \mathbf{P}^\epsilon(\sup_{|u| \leq N} \sup_{x \in \mathbb{R}^l} |A_s^\epsilon(u, x)|\delta \\ &+ \sqrt{\epsilon} \sup_{\tilde{s} \in [s, s+\delta]} \left| \int_{s \wedge \tau_N^\epsilon}^{\tilde{s} \wedge \tau_N^\epsilon} B_r^\epsilon(X_r^\epsilon, x_r^\epsilon) dW_r^\epsilon \right| > \eta). \end{aligned}$$

Let e_i , for $i = 1, 2, \dots, n$, denote the i th unit vector of \mathbb{R}^n . Thanks to (2.2b) and (2.2d), for small enough δ and arbitrary $\alpha > 0$, provided $\epsilon > 0$ is small enough, on using Doob's inequality,

$$\begin{aligned} \mathbf{P}^\epsilon(\sup_{\tilde{s} \in [s, s+\delta]} |X_{\tilde{s}}^\epsilon - X_s^\epsilon| > \eta) &\leq \mathbf{P}^\epsilon(\tau_N^\epsilon \leq t) + \mathbf{P}^\epsilon\left(\sqrt{\epsilon} \sup_{\tilde{s} \in [s, s+\delta]} \left| \int_{s \wedge \tau_N^\epsilon}^{\tilde{s} \wedge \tau_N^\epsilon} B_r^\epsilon(X_r^\epsilon, x_r^\epsilon) dW_r^\epsilon \right| > \frac{\eta}{2}\right) \\ &\leq \mathbf{P}^\epsilon(\tau_N^\epsilon \leq t) + \sum_{i=1}^n \mathbf{P}^\epsilon\left(\sqrt{\epsilon} \sup_{\tilde{s} \in [s, s+\delta]} \left(e_i^T \int_{s \wedge \tau_N^\epsilon}^{\tilde{s} \wedge \tau_N^\epsilon} B_r^\epsilon(X_r^\epsilon, x_r^\epsilon) dW_r^\epsilon \right) > \frac{\eta}{2n}\right) \leq \mathbf{P}^\epsilon(\tau_N^\epsilon \leq t) \\ &+ \sum_{i=1}^n \mathbf{P}^\epsilon\left(\sup_{\tilde{s} \in [s, s+\delta]} \exp\left(\frac{\alpha}{\sqrt{\epsilon}} e_i^T \int_{s \wedge \tau_N^\epsilon}^{\tilde{s} \wedge \tau_N^\epsilon} B_r^\epsilon(X_r^\epsilon, x_r^\epsilon) dW_r^\epsilon - \frac{\alpha^2}{2\epsilon} \int_{s \wedge \tau_N^\epsilon}^{\tilde{s} \wedge \tau_N^\epsilon} e_i^T C_r^\epsilon(X_r^\epsilon, x_r^\epsilon) e_i dr\right) \right. \\ &> e^{\alpha\eta/(2n\epsilon)} \exp\left(-\frac{\alpha^2\delta}{2\epsilon} \sup_{r \in [0, t]} \sup_{|u| \leq N} \sup_{x \in \mathbb{R}^l} \|C_r^\epsilon(u, x)\|\right) \leq \mathbf{P}^\epsilon(\sup_{\tilde{s} \in [0, t]} |X_{\tilde{s}}^\epsilon| \geq N) \\ &\quad + n e^{-\alpha\eta/(2n\epsilon)} \exp\left(\frac{\alpha^2\delta}{2\epsilon} \sup_{r \in [0, t]} \sup_{|u| \leq N} \sup_{x \in \mathbb{R}^l} \|C_r^\epsilon(u, x)\|\right). \end{aligned}$$

By (2.2d), (4.3) and the fact that α can be chosen arbitrarily great,

$$\limsup_{\delta \rightarrow 0} \limsup_{\epsilon \rightarrow 0} \sup_{s \in [0, t]} \mathbf{P}^\epsilon \left(\sup_{\tilde{s} \in [s, s+\delta]} |X_s^\epsilon - X_{\tilde{s}}^\epsilon| > \eta \right)^\epsilon = 0.$$

The sequential exponential tightness of X^ϵ follows from part 2) of Lemma 4.1.

We prove now that μ^ϵ is sequentially exponentially tight. Let f represent an \mathbb{R} -valued twice continuously differentiable function on \mathbb{R}^l . By (2.1b) and Itô's lemma,

$$\begin{aligned} f(x_t^\epsilon) &= f(x_0^\epsilon) + \frac{1}{\epsilon} \int_0^t Df(x_s^\epsilon)^T a_s^\epsilon(X_s^\epsilon, x_s^\epsilon) ds + \frac{1}{2\epsilon} \int_0^t \text{tr} (c_s^\epsilon(X_s^\epsilon, x_s^\epsilon) D^2 f(x_s^\epsilon)) ds \\ &\quad + \frac{1}{\sqrt{\epsilon}} \int_0^t Df(x_s^\epsilon)^T b_s^\epsilon(X_s^\epsilon, x_s^\epsilon) dW_s^\epsilon. \end{aligned}$$

Therefore, on identifying μ^ϵ with measure $\mu^\epsilon(dt, dx)$, we have that, in analogy with (4.1),

$$\begin{aligned} \mathbf{E}^\epsilon \exp \left(f(x_{t \wedge \tau_N^\epsilon}^\epsilon) - f(x_0^\epsilon) - \frac{1}{\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \int_{\mathbb{R}^l} Df(x)^T a_s^\epsilon(X_s^\epsilon, x) \mu^\epsilon(ds, dx) \right. \\ \left. - \frac{1}{2\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \int_{\mathbb{R}^l} \text{tr} (c_s^\epsilon(X_s^\epsilon, x) D^2 f(x)) \mu^\epsilon(ds, dx) - \frac{1}{2\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \int_{\mathbb{R}^l} \|Df(x)\|_{c_s^\epsilon(X_s^\epsilon, x)}^2 \mu^\epsilon(ds, dx) \right) \leq 1. \end{aligned}$$

Let $g_2(u)$, where $u \in \mathbb{R}_+$, be an \mathbb{R}_+ -valued nondecreasing \mathbb{C}^2 -function with a bounded second derivative such that $Dg_2(0) = D^2g_2(0) = 0$ and $g_2(u) = u$ for $u \geq 1$. For given $\check{N} > 0$, we let $f(x) = g_2(|x| - \check{N}^+)$, where $x \in \mathbb{R}^l$. By (2.4a), if \check{N} is great enough, then for all ϵ small enough, $(x/|x|)^T a_{s \wedge \tau_N^\epsilon}^\epsilon(X_{s \wedge \tau_N^\epsilon}^\epsilon, x) \leq 0$ provided $|x| \geq \check{N}$. Since g_2 is a nondecreasing function,

$$Df(x)^T a_{s \wedge \tau_N^\epsilon}^\epsilon(X_{s \wedge \tau_N^\epsilon}^\epsilon, x) = Dg_2(|x| - \check{N}^+)(x/|x|)^T a_{s \wedge \tau_N^\epsilon}^\epsilon(X_{s \wedge \tau_N^\epsilon}^\epsilon, x) \leq 0.$$

In addition, like in (4.2), $\text{tr} (c_s^\epsilon(X_s^\epsilon, x) D^2 |x|) \leq l\sqrt{l-1} \|c_s^\epsilon(X_s^\epsilon, x)\|/|x|$. We obtain that

$$\begin{aligned} \mathbf{E}^\epsilon \exp \left(-f(x_0^\epsilon) - \frac{1}{\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \int_{|x| > \check{N}+1} \frac{x^T}{|x|} a_s^\epsilon(X_s^\epsilon, x) \mu^\epsilon(ds, dx) \right. \\ \left. - \frac{1}{2\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \int_{\check{N} \leq |x| \leq \check{N}+1} (\text{tr} (c_s^\epsilon(X_s^\epsilon, x) D^2 f(x)) + \|Df(x)\|_{c_s^\epsilon(X_s^\epsilon, x)}^2) \mu^\epsilon(ds, dx) \right. \\ \left. - \frac{1}{2\epsilon} \int_0^{t \wedge \tau_N^\epsilon} \int_{|x| > \check{N}+1} \left(\frac{\sqrt{l-1}}{|x|} l \|c_s^\epsilon(X_s^\epsilon, x)\| + \left\| \frac{x}{|x|} \right\|_{c_s^\epsilon(X_s^\epsilon, x)}^2 \right) \mu^\epsilon(ds, dx) \right) \leq 1. \quad (4.4) \end{aligned}$$

Since $\|c_s^\epsilon(u, x)\|$ is asymptotically bounded locally in (s, u) and globally in x , see (2.2a), there exists $\tilde{L} > 0$ such that $|\text{tr} (c_{s \wedge \tau_N^\epsilon}^\epsilon(X_{s \wedge \tau_N^\epsilon}^\epsilon, x) D^2 f(x)) + \|Df(x)\|_{c_{s \wedge \tau_N^\epsilon}^\epsilon(X_{s \wedge \tau_N^\epsilon}^\epsilon, x)}^2| \leq \tilde{L}$ for all $s \leq t$, all \check{N} , and

all x such that $|x| \in [\check{N}, \check{N} + 1]$, provided $\epsilon > 0$ is small enough. We can also assume that \tilde{L} is an upper bound on $\|c_{s \wedge \tau_N^\epsilon}^\epsilon(X_{s \wedge \tau_N^\epsilon}^\epsilon, x)\|$. We thus obtain from (4.4), on recalling that $\mu^\epsilon([0, t], \mathbb{R}^l) = t$, that provided ϵ is small enough and \check{N} is great enough,

$$\mathbf{E}^\epsilon \exp\left(-f(x_0^\epsilon) + \frac{1}{\epsilon} M^\epsilon \mu^\epsilon([0, t \wedge \tau_N^\epsilon], \{x : |x| > \check{N} + 1\}) - \frac{3\tilde{L}t}{2\epsilon}\right) \leq 1,$$

where

$$M^\epsilon = - \sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n : |u| \leq N} \sup_{x \in \mathbb{R}^l : |x| > \check{N} + 1} \frac{x^T}{|x|} a_s^\epsilon(u, x) > 0.$$

It follows that for arbitrary $\delta > 0$, all ϵ small enough, and all \check{N} great enough,

$$\begin{aligned} \mathbf{P}^\epsilon(\mu^\epsilon([0, t \wedge \tau_N^\epsilon], \{x \in \mathbb{R}^l : |x| > \check{N} + 1\}) > \delta) &\leq \mathbf{P}^\epsilon(|x_0^\epsilon| > \check{N}) \\ &+ \mathbf{E}^\epsilon \exp\left(\frac{M^\epsilon}{\epsilon} \mu^\epsilon([0, t \wedge \tau_N^\epsilon], \{x : |x| > \check{N} + 1\})\right) \mathbf{1}_{\{|x_0^\epsilon| \leq \check{N}\}} \exp\left(-\frac{M^\epsilon}{\epsilon} \delta\right) \\ &\leq \mathbf{P}^\epsilon(|x_0^\epsilon| > \check{N}) + \exp\left(\frac{3\tilde{L}t}{2\epsilon} - \frac{M^\epsilon \delta}{\epsilon} + g_2(0)\right), \end{aligned}$$

so by the facts that $\liminf_{\epsilon \rightarrow 0} M^\epsilon \rightarrow \infty$ and $\limsup_{\epsilon \rightarrow 0} \mathbf{P}^\epsilon(|x_0^\epsilon| \geq \check{N})^\epsilon \rightarrow 0$ as $\check{N} \rightarrow \infty$, and that (4.3) holds, we obtain that

$$\lim_{\check{N} \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \mathbf{P}^\epsilon(\mu^\epsilon([0, t], \{x \in \mathbb{R}^l : |x| > \check{N} + 1\}) > \delta)^\epsilon = 0.$$

Since $|\mu_t^\epsilon(\Theta) - \mu_s^\epsilon(\Theta)| \leq |t - s|$, for $\Theta \in \mathcal{B}(\mathbb{R}^l)$, the sequential exponential tightness of μ^ϵ follows from part 1 of Lemma 4.1. \square

Remark 4.2. Since $(X^\epsilon, \mu^\epsilon)$ is continuous in ϵ in distribution, one can prove that $(X^\epsilon, \mu^\epsilon)$ is exponentially tight.

5 The equation for the large deviation function

In this section, we derive an equation for large deviation limit points of $(X^\epsilon, \mu^\epsilon)$ that is to be used for identifying the large deviation function. For $0 = t_0 < t_1 < \dots < t_i$, let

$$\lambda(t, X) = \sum_{j=1}^i \lambda_j(X_{t_{j-1}}) \mathbf{1}_{[t_{j-1}, t_j)}(t), \quad (5.1)$$

where $X = (X_s, s \in \mathbb{R}_+) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ and the functions $\lambda_j(u)$, for $u \in \mathbb{R}^n$, are \mathbb{R}^n -valued and continuous. We define

$$\int_0^t \lambda(s, X) dX_s = \sum_{j=1}^i \lambda_i(X_{t_{j-1} \wedge t})^T (X_{t \wedge t_j} - X_{t \wedge t_{j-1}}). \quad (5.2)$$

Let $f(t, u, x)$ represent a $\mathcal{C}^{1,2,2}(\mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^l)$ -function with compact support in x locally uniformly in (t, u) and let, with $(X, \mu) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$,

$$\begin{aligned}
U_t^{\lambda(\cdot), f}(X, \mu) &= \int_0^t \lambda(s, X) dX_s - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T A_s(X_s, x) \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} D_x f(s, X_s, x)^T a_s(X_s, x) \mu(ds, dx) - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr}(c_s(X_s, x) D_{xx}^2 f(s, X_s, x)) \mu(ds, dx) \\
&\quad - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|\lambda(s, X)\|_{C_s(X_s, x)}^2 \mu(ds, dx) - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|D_x f(s, X_s, x)\|_{c_s(X_s, x)}^2 \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T G_s(X_s, x) D_x f(s, X_s, x) \mu(ds, dx). \quad (5.3)
\end{aligned}$$

Under condition 2.1, $U_t^{\lambda(\cdot), f}(X, \mu)$ is a continuous function of (X, μ) .

Let $\tau(X, \mu)$ represent a continuous function of $(X, \mu) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ that is also a stopping time relative to the flow $\mathbf{G} = (\mathcal{G}_t, t \in \mathbb{R}_+)$ on $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, where the σ -algebra \mathcal{G}_t is generated by the mappings $X \rightarrow X_s$ and $\mu \rightarrow \mu_s$ for $s \leq t$. (We note that the flow \mathbf{G} is not right continuous, so τ is a strict stopping time, see Jacod and Shiryaev [25].) Suppose also that $X_{t \wedge \tau(X, \mu)}$ is a bounded function of (X, μ) .

Theorem 5.1. *Suppose that conditions 2.1, (2.2a), (2.2b), (2.2d), and (2.12) hold. If $\tilde{\mathbf{I}}$ is a large deviation limit point of $(X^\epsilon, \mu^\epsilon)$ for rate $1/\epsilon$ as $\epsilon \rightarrow 0$, then*

$$\sup_{(X, \mu) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))} (U_{t \wedge \tau(X, \mu)}^{\lambda(\cdot), f}(X, \mu) - \tilde{\mathbf{I}}(X, \mu)) = 0. \quad (5.4)$$

Proof. The process $(\lambda(t, X^\epsilon), t \in \mathbb{R}_+)$ is \mathbf{F}^ϵ -adapted so that by (2.1a) and (5.2),

$$\int_0^t \lambda(s, X^\epsilon) dX_s^\epsilon = \int_0^t \lambda(s, X^\epsilon)^T A_s^\epsilon(X_s^\epsilon, x_s^\epsilon) ds + \sqrt{\epsilon} \int_0^t \lambda(s, X^\epsilon)^T B_s^\epsilon(X_s^\epsilon, x_s^\epsilon) dW_s^\epsilon. \quad (5.5)$$

By (2.1a), (2.1b), and Itô's lemma,

$$\begin{aligned}
f(t, X_t^\epsilon, x_t^\epsilon) &= f(0, X_0^\epsilon, x_0^\epsilon) + \int_0^t \frac{\partial f(s, X_s^\epsilon, x_s^\epsilon)}{\partial s} ds + \int_0^t D_u f(s, X_s^\epsilon, x_s^\epsilon)^T A_s^\epsilon(X_s^\epsilon, x_s^\epsilon) ds \\
&\quad + \sqrt{\epsilon} \int_0^t D_u f(s, X_s^\epsilon, x_s^\epsilon)^T B_s^\epsilon(X_s^\epsilon, x_s^\epsilon) dW_s^\epsilon + \frac{1}{\epsilon} \int_0^t D_x f(s, X_s^\epsilon, x_s^\epsilon)^T a_s^\epsilon(X_s^\epsilon, x_s^\epsilon) ds \\
&\quad + \frac{1}{\sqrt{\epsilon}} \int_0^t D_x f(s, X_s^\epsilon, x_s^\epsilon)^T b_s^\epsilon(X_s^\epsilon, x_s^\epsilon) dW_s^\epsilon + \frac{\epsilon}{2} \int_0^t \text{tr}(C_s^\epsilon(X_s^\epsilon, x_s^\epsilon) D_{uu}^2 f(s, X_s^\epsilon, x_s^\epsilon)) ds \\
&\quad + \frac{1}{2\epsilon} \int_0^t \text{tr}(c_s^\epsilon(X_s^\epsilon, x_s^\epsilon) D_{xx}^2 f(s, X_s^\epsilon, x_s^\epsilon)) ds + \int_0^t \text{tr}(G_s^\epsilon(X_s^\epsilon, x_s^\epsilon) D_{ux}^2 f(s, X_s^\epsilon, x_s^\epsilon)) ds, \quad (5.6)
\end{aligned}$$

where $G_s^\epsilon(u, x) = B_s^\epsilon(u, x)b_s^\epsilon(u, x)^T$. We denote

$$\begin{aligned}
U_t^\epsilon(X, \mu) &= \int_0^t \lambda(s, X) dX_s - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T A_s^\epsilon(X_s, x) \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} D_x f(s, X_s, x)^T a_s^\epsilon(X_s, x) \mu(ds, dx) - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr}(c_s^\epsilon(X_s, x) D_{xx}^2 f(s, X_s, x)) \mu(ds, dx) \\
&\quad - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|\lambda(s, X)\|_{C_s^\epsilon(X_s, x)}^2 \mu(ds, dx) - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|D_x f(s, X_s, x)\|_{c_s^\epsilon(X_s, x)}^2 \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T G_s^\epsilon(X_s, x) D_x f(s, X_s, x) \mu(ds, dx)
\end{aligned}$$

and

$$\begin{aligned}
V_t^\epsilon(X, \mu) &= f(t, X_t, x_t) - f(0, X_0, x_0) - \int_0^t \int_{\mathbb{R}^l} \frac{\partial f(s, X_s, x)}{\partial s} \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} D_u f(s, X_s, x)^T A_s^\epsilon(X_s, x) \mu(ds, dx) - \frac{\epsilon}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr}(C_s^\epsilon(X_s, x) D_{uu}^2 f(s, X_s, x)) \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} \text{tr}(G_s^\epsilon(X_s, x) D_{ux}^2 f(s, X_s, x)) \mu(ds, dx) - \frac{\epsilon}{2} \int_0^t \int_{\mathbb{R}^l} \|D_u f(s, X_s, x)\|_{C_s^\epsilon(X_s, x)}^2 \mu(ds, dx) \\
&\quad - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T C_s^\epsilon(X_s, x) D_u f(s, X_s, x) \mu(ds, dx) - \int_0^t \int_{\mathbb{R}^l} D_u f(s, X_s, x)^T G_s^\epsilon(X_s, x) D_x f(s, X_s, x) \mu(ds, dx).
\end{aligned}$$

Since the function $\lambda(s, u)$ is locally bounded, the function $f(s, u, x)$ and its derivatives are locally bounded and are of compact support in x , conditions (2.2a), (2.2b), (2.2d), and (2.13) hold, and $X_{t \wedge \tau(X, \mu)}$ is bounded, we have that there exists number $R(t) > 0$ such that for all ϵ small enough uniformly over (X, μ) ,

$$|U_{t \wedge \tau(X, \mu)}^\epsilon(X, \mu)| + |V_{t \wedge \tau(X, \mu)}^\epsilon(X, \mu)| \leq R(t). \quad (5.7)$$

Since X_s^ϵ and μ_s^ϵ are \mathcal{F}_s^ϵ -measurable, $\tau(X^\epsilon, \mu^\epsilon)$ is a stopping time relative to \mathbf{F}^ϵ . By (5.5), (5.6), and (5.7), the process $(\exp((1/\epsilon)U_{t \wedge \tau(X^\epsilon, \mu^\epsilon)}^\epsilon(X^\epsilon, \mu^\epsilon) + V_{t \wedge \tau(X^\epsilon, \mu^\epsilon)}^\epsilon(X^\epsilon, \mu^\epsilon)), t \in \mathbb{R}_+)$ is a bounded \mathbf{F}^ϵ -martingale, so

$$\mathbf{E}^\epsilon \exp\left(\frac{1}{\epsilon} U_{t \wedge \tau(X^\epsilon, \mu^\epsilon)}^\epsilon(X^\epsilon, \mu^\epsilon) + V_{t \wedge \tau(X^\epsilon, \mu^\epsilon)}^\epsilon(X^\epsilon, \mu^\epsilon)\right) = 1.$$

Since the function $f(s, u, x)$ is of compact support in x , the convergence hypotheses in (2.12) and the bound in (5.7) imply that $U_{t \wedge \tau(X, \mu)}^\epsilon(X, \mu) \rightarrow U_{t \wedge \tau(X, \mu)}^{\lambda(\cdot), f}(X, \mu)$ as $\epsilon \rightarrow 0$ uniformly over compact sets. By Theorem 3.3, $\sup_{(X, \mu) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))} (U_{t \wedge \tau(X, \mu)}^{\lambda(\cdot), f}(X, \mu) - \tilde{\mathbf{I}}(X, \mu)) = 0$. \square

Remark 5.1. One can see that there exists compact $K \subset \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ such that $\sup_{(X, \mu) \in K} (U_{t \wedge \tau(X, \mu)}^{\lambda(\cdot), f}(X, \mu) - \tilde{\mathbf{I}}(X, \mu)) = 0$.

6 Regularity properties

Let $\tilde{\mathbf{I}}$ represent a large deviation limit point of $(X^\epsilon, \mu^\epsilon)$ for rate $1/\epsilon$ as $\epsilon \rightarrow 0$ such that $\tilde{\mathbf{I}}(X, \mu) = \infty$ unless $X_0 = \hat{u}$, where \hat{u} is a preselected element of \mathbb{R}^n . Let, as in Theorem 3.4, for $(X, \mu) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$,

$$\mathbf{I}^{**}(X, \mu) = \sup_{\lambda(\cdot), f, t, \tau} U_{t \wedge \tau(X, \mu)}^{\lambda(\cdot), f}(X, \mu), \quad (6.1)$$

with the supremum being taken over $\lambda(t, X)$, $f(t, u, x)$, and $\tau(X, \mu)$ satisfying the requirements of Theorem 5.1 and over $t \geq 0$. We note that, under Condition 2.1, $\mathbf{I}^{**}(X, \mu)$ is a lower semicontinuous function of (X, μ) and that by Theorem 5.1,

$$\mathbf{I}^{**}(X, \mu) \leq \tilde{\mathbf{I}}(X, \mu). \quad (6.2)$$

The rest of the paper is concerned mostly with proving that equality prevails in (6.2), provided $X_0 = \hat{u}$. Since the case where $\mathbf{I}^{**}(X, \mu) < \infty$ needs to be considered only, in this section we undertake a study of the properties of (X, μ) such that $\mathbf{I}^{**}(X, \mu) < \infty$. We then prove that if $\mathbf{I}^{**}(X, \mu) < \infty$ and $X_0 = \hat{u}$, then $\mathbf{I}^{**}(X, \mu) = \mathbf{I}(X, \mu)$, where $\mathbf{I}(X, \mu)$ is given in the statements of Theorem 2.1 and Proposition 2.1 with $\mathbf{I}_0(\hat{u}) = 0$, see Theorem 6.1. We assume throughout conditions 2.1, 2.2, (2.4b), (2.14c) and (2.14d) to hold.

Lemma 6.1. *If $\mu \in \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, then μ is of the form $\mu(ds, dx) = \nu_s(dx) ds$, where $\nu_s(dx)$ is a transition probability kernel from \mathbb{R}_+ to \mathbb{R}^l . If $(X, \mu) \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ is such that $\mathbf{I}^{**}(X, \mu) < \infty$, then X is absolutely continuous with respect to Lebesgue measure.*

Proof. As $\mu \in \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, we have that $\mu(ds, dx) = \nu_s(dx)\mu(ds, \mathbb{R}^l)$, where $\nu_s(dx)$ is a transition kernel from \mathbb{R}_+ to \mathbb{R}^l , see, e.g., Theorem 8.1 on p.502 of Ethier and Kurtz [15].

Since $\mu(ds, \mathbb{R}^l)$ is Lebesgue measure on \mathbb{R}_+ , $\mu(ds, dx) = \nu_s(dx) ds$.

On taking $f = 0$ in (5.3) and assuming $\lambda(s, X)$ not to depend on X , so the piece of notation $\lambda(s)$ can be used instead, we have by (5.3), (6.1), and the part of the lemma just proved that if $\mathbf{I}^{**}(X, \mu) < \infty$, then

$$\int_0^t \lambda(s) dX_s \leq \int_0^t \int_{\mathbb{R}^l} \lambda(s)^T A_s(X_s, x) \nu_s(dx) ds + \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|\lambda(s)\|_{C_s(X_s, x)}^2 \nu_s(dx) ds + \mathbf{I}^{**}(X, \mu).$$

Replacing $\lambda(s)$ with $\delta\lambda(s)$, where $\delta > 0$, dividing through by δ , and minimising the righthand side over δ obtains that

$$\int_0^t \lambda(s) dX_s \leq \int_0^t \int_{\mathbb{R}^l} \lambda(s)^T A_s(X_s, x) \nu_s(dx) ds + \sqrt{2} \sqrt{\mathbf{I}^{**}(X, \mu)} \sqrt{\int_0^t \int_{\mathbb{R}^l} \|\lambda(s)\|_{C_s(X_s, x)}^2 \nu_s(dx) ds}.$$

It follows that X is absolutely continuous with respect to ds . □

By Lemma 6.1, if $\mathbf{I}^{**}(X, \mu) < \infty$, then (5.3) takes the form

$$\begin{aligned}
U_t^{\lambda(\cdot), f}(X, \mu) &= \int_0^t \lambda(s, X)^T \dot{X}_s ds - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T A_s(X_s, x) \nu_s(dx) ds \\
&\quad - \int_0^t \int_{\mathbb{R}^l} D_x f(s, X_s, x)^T a_s(X_s, x) \nu_s(dx) ds - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr}(c_s(X_s, x) D_{xx}^2 f(s, X_s, x)) \nu_s(dx) ds \\
&\quad - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|\lambda(s, X)\|_{C_s(X_s, x)}^2 \nu_s(dx) ds - \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|D_x f(s, X_s, x)\|_{c_s(X_s, x)}^2 \nu_s(dx) ds \\
&\quad - \int_0^t \int_{\mathbb{R}^l} \lambda(s, X)^T G_s(X_s, x) D_x f(s, X_s, x) \nu_s(dx) ds. \quad (6.3)
\end{aligned}$$

The next step is to show that $\nu_s(dx)$ has to be absolutely continuous with respect to dx and establish its integrability properties. We need, however, to lay the groundwork. The proofs of the following two lemmas are relegated to the appendix. The first one is essentially due to Röckner and Zhang [44, pp.204,205], [45], see also Bogachev, Krylov, and Röckner [6]. The second one addresses regularity of the invariant measures of diffusions and may be of interest in its own right.

Lemma 6.2. *Let $d \in \mathbb{N}$ and let O represent either \mathbb{R}^d or an open ball in \mathbb{R}^d . If $m(x)$ is an \mathbb{R}_+ -valued measurable function on \mathbb{R}^d such that $m \in \mathbb{W}_{loc}^{1,1}(\mathbb{R}^d)$ and $\sqrt{m} \in \mathbb{W}^{1,2}(O)$, then $\mathbb{H}^{1,2}(O, m(x) dx) = \mathbb{W}^{1,2}(O, m(x) dx)$.*

Lemma 6.3. *For $d \in \mathbb{N}$ and $x \in \mathbb{R}^d$, let $c(x)$ represent a locally Lipschitz continuous function with values in the set of symmetric positive definite $d \times d$ -matrices and let $b(x)$ represent an \mathbb{R}^d -valued measurable function. Suppose $m(x)$ is a probability density on \mathbb{R}^d such that $m(\ln m)^2 \in \mathbb{L}_{loc}^1(\mathbb{R}^d)$, $b \in \mathbb{L}_{loc}^2(\mathbb{R}^d, \mathbb{R}^d, m(x) dx)$, and*

$$\int_{\mathbb{R}^d} \text{tr}(c(x) D^2 p(x)) m(x) dx + \int_{\mathbb{R}^d} Dp(x)^T b(x) m(x) dx = 0$$

for all $p \in \mathbb{C}_0^\infty(\mathbb{R}^d)$, where we assume that $0(\ln 0)^2 = 0$.

Then $m \in \mathbb{W}_{loc}^{1,1}(\mathbb{R}^d)$ and $\sqrt{m} \in \mathbb{W}_{loc}^{1,2}(\mathbb{R}^d)$. Furthermore, given open ball S from \mathbb{R}^d , there exists constant M which depends on S , on the Lipschitz constant of $c(x)$ on S , and on $\inf_{x \in S} x^T c(x) x / |x|^2$ only, such that

$$\int_S \frac{|Dm(x)|^2}{m(x)} dx \leq M \left(1 + \int_S (\ln m(x))^2 m(x) dx + \int_S |b(x)|^2 m(x) dx \right). \quad (6.4)$$

The latter lemma is a local version of the result by Bogachev, Krylov, and Röckner [6] that if $b \in \mathbb{L}^2(\mathbb{R}^d, m(x) dx)$ then $\sqrt{m} \in \mathbb{W}^{1,2}(\mathbb{R}^d)$, see also Metafune, Pallara, and Rhandi [31].

For the next lemma, we recall that, according to our conventions, $q' = q/(q-1)$, provided $q > 1$.

Lemma 6.4. *Suppose that $\mathbf{I}^{**}(X, \mu) < \infty$, where $\mu(ds, dx) = \nu_s(dx) ds$. Then, for almost all s , the transition kernel $\nu_s(dx)$ is absolutely continuous with respect to Lebesgue measure, the density*

$m_s(x) = \nu_s(dx)/dx$ is an element of $\mathbb{L}_{loc}^\beta(\mathbb{R}^l)$ for all $\beta \in [1, l/(l-1))$ and is an element of $\mathbb{W}_{loc}^{1,\alpha}(\mathbb{R}^l)$ for all $\alpha \in [1, 2l/(2l-1))$, and $\sqrt{m_s(\cdot)} \in \mathbb{W}_{loc}^{1,2}(\mathbb{R}^l)$. Furthermore, for arbitrary $t > 0$ and open ball $S \subset \mathbb{R}^l$,

$$\int_0^t \int_S \frac{|Dm_s(x)|^2}{m_s(x)} dx ds < \infty. \quad (6.5)$$

If, in addition, $\sqrt{m_s(\cdot)} \in \mathbb{W}_0^{1,2}(\mathbb{R}^l)$, then $Dm_s(\cdot)/m_s(\cdot) \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$.

If $\kappa > 0$, $q \geq 2$, and $q > l$, then

$$\sup_{(X,\mu): \mathbf{I}^{**}(X,\mu) \leq \kappa} \int_0^t \int_S m_s(x)^{q'} dx ds < \infty. \quad (6.6)$$

Proof. By (6.1) and (6.3) with $\lambda(s, X) = 0$ and $f(s, u, x) = \phi(s, x)$, where $\phi \in \mathbb{C}^{1,2}(\mathbb{R}_+ \times \mathbb{R}^l)$ and the support of ϕ in x is bounded locally uniformly in s ,

$$\begin{aligned} & -\frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr}(c_s(X_s, x) D_{xx}^2 \phi(s, x)) \nu_s(dx) ds - \int_0^t \int_{\mathbb{R}^l} D_x \phi(s, x)^T a_s(X_s, x) \nu_s(dx) ds \\ & \leq \mathbf{I}^{**}(X, \mu) + \frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|D_x \phi(s, x)\|_{c_s(X_s, x)}^2 \nu_s(dx) ds. \end{aligned}$$

Replacing $\phi(s, x)$ with $\delta \phi(s, x)$, where $\delta > 0$, dividing through by δ , and minimising the righthand side over δ yields

$$\begin{aligned} & -\frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr}(c_s(X_s, x) D_{xx}^2 \phi(s, x)) \nu_s(dx) ds - \int_0^t \int_{\mathbb{R}^l} D_x \phi(s, x)^T a_s(X_s, x) \nu_s(dx) ds \\ & \leq \sqrt{2} \mathbf{I}^{**}(X, \mu)^{1/2} \left(\int_0^t \int_{\mathbb{R}^l} \|D_x \phi(s, x)\|_{c_s(X_s, x)}^2 \nu_s(dx) ds \right)^{1/2}. \quad (6.7) \end{aligned}$$

Let $\mathbb{L}_0^{1,2}([0, t] \times \mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx) ds)$ denote the closure in $\mathbb{L}^2([0, t] \times \mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx) ds)$ of the space of functions $D_x \phi$. By (6.7), the lefthand side extends to a continuous functional $T_t(g)$ on $\mathbb{L}_0^{1,2}([0, t] \times \mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx) ds)$. By the Riesz representation theorem, there exists a unique $\psi \in \mathbb{L}_0^{1,2}([0, t] \times \mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx) ds)$ such that

$$T_t(g) = \int_0^t \int_{\mathbb{R}^l} g(s, x)^T c_s(X_s, x) \psi(s, x) \nu_s(dx) ds,$$

for all $g \in \mathbb{L}_0^{1,2}([0, t] \times \mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx) ds)$, and

$$\left(\int_0^t \int_{\mathbb{R}^l} \|\psi(s, x)\|_{c_s(X_s, x)}^2 \nu_s(dx) ds \right)^{1/2} \leq \sqrt{2} \mathbf{I}^{**}(X, \mu)^{1/2}. \quad (6.8)$$

By uniqueness, ψ can be extended to a function on $\mathbb{R}_+ \times \mathbb{R}^l$ so that for all $t > 0$,

$$\begin{aligned} -\frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \text{tr} (c_s(X_s, x) D_{xx}^2 \phi(s, x)) \nu_s(dx) ds - \int_0^t \int_{\mathbb{R}^l} D_x \phi(s, x)^T a_s(X_s, x) \nu_s(dx) ds \\ = \int_0^t \int_{\mathbb{R}^l} D_x \phi(s, x)^T c_s(X_s, x) \psi(s, x) \nu_s(dx) ds. \end{aligned} \quad (6.9)$$

It follows that for almost all s and for all $h \in \mathbb{C}_0^2(\mathbb{R}^l)$,

$$\begin{aligned} -\frac{1}{2} \int_{\mathbb{R}^l} \text{tr} (c_s(X_s, x) D^2 h(x)) \nu_s(dx) = \int_{\mathbb{R}^l} Dh(x)^T a_s(X_s, x) \nu_s(dx) \\ + \int_{\mathbb{R}^l} Dh(x)^T c_s(X_s, x) \psi(s, x) \nu_s(dx). \end{aligned} \quad (6.10)$$

Since $\psi \in \mathbb{L}_0^{1,2}([0, t] \times \mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx) ds)$, we have that, for almost all s , $\psi(s, \cdot)$ belongs to the closure of the set of the $D_x h$ in $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), \nu_s(dx))$. In particular, $\int_{\mathbb{R}^l} |\psi(s, x)|^2 \nu_s(dx) < \infty$. Since $a_s(X_s, \cdot)$ and $\psi(s, \cdot)$ are locally integrable with respect to $\nu_s(dx)$ and $c_s(X_s, \cdot)$ is uniformly positive definite and is of class \mathbb{C}^1 , (6.10) and Theorem 2.1 in Bogachev, Krylov, and Röckner [7] imply that the measure $\nu_s(dx)$ has density $m_s(x)$ with respect to Lebesgue measure which belongs to $L_{\text{loc}}^\beta(\mathbb{R}^l)$ for all $\beta < l'$. It follows, since $a_s(X_s, \cdot)$ and $c_s(X_s, \cdot)$ are locally bounded and $\int_{\mathbb{R}^l} |\psi(s, x)|^2 \nu_s(dx) < \infty$, that, for arbitrary open ball S in \mathbb{R}^l , there exists $M > 0$ such that for all $h \in \mathbb{C}_0^2(S)$

$$\left| \int_S \text{tr} (c_s(X_s, x) D^2 h(x)) m_s(x) dx \right| \leq M \|Dh\|_{\mathbb{L}^{2\beta'}(S, \mathbb{R}^l)}.$$

Since $c_s(u, \cdot)$ is uniformly positive definite and is of class \mathbb{C}^1 , by Theorem 6.1 in Agmon [2], the density $m_s(\cdot)$ belongs to $\mathbb{W}_{\text{loc}}^{1,\alpha}(S)$ for all $\alpha < 2l/(2l-1)$. The inclusion $\sqrt{m_s(\cdot)} \in \mathbb{W}_{\text{loc}}^{1,2}(\mathbb{R}^l)$ follows from Lemma 6.3 and (6.10). For the inequality (6.5), we also recall (6.4) and (6.8).

Let $\mathbb{L}^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ represent the closure in $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ of the set of gradients of $\mathbb{C}^1(\mathbb{R}^l)$ -functions such that those gradients belong to $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$. Let $\varphi_M(y)$ be a smooth function such that $\varphi_M(y) = y$ when $|y| \leq M$ and $\varphi_M(y) = (y/|y|)(M+1)$ when $|y| \geq M+1$. Let $\eta(x)$ represent a $[0, 1]$ -valued continuously differentiable nonincreasing function defined for $x \geq 0$ such that $\eta(x) = 1$ for $x \in [0, 1]$ and $\eta(x) = 0$ for $x \geq 2$. Let $\eta_r(x) = \eta(|x|/r)$ where $x \in \mathbb{R}^l$ and $r > 0$. By associating with $h \in \mathbb{L}^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ the function $\varphi_M(h(x))\eta_r(x)$ and taking limits as $r \rightarrow \infty$ and $M \rightarrow \infty$, one can show that $\mathbb{L}^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx) = \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$. The property that $Dm_s(\cdot)/m_s(\cdot) \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$ when $\sqrt{m_s(\cdot)} \in \mathbb{W}^{1,2}(\mathbb{R}^l)$ now follows from Lemma 6.2.

We now adapt the proof of Theorem 2.1 in Bogachev, Krylov, and Röckner [7] in order to obtain the bound in (6.6). Let S_1 represent an open ball which contains S . By (6.8), (6.9), and local boundedness of $a_s(u, x)$ and $c_s(u, x)$, assuming that $\phi(s, x)$ in (6.9) is supported by S_1 in x for all $s \in [0, t]$, there exists $L_1 > 0$ such that for all (X, μ) with $\mathbf{I}^{**}(X, \mu) \leq \delta$,

$$\left| \int_0^t \int_{S_1} \text{tr} (c_s(X_s, x) D_{xx}^2 \phi(s, x)) m_s(x) dx ds \right| \leq L_1 \left(\int_0^t \sup_{x \in S_1} |D_x \phi(s, x)|^2 ds \right)^{1/2}. \quad (6.11)$$

An approximation argument shows that one may assume that $\phi(s, x)$ is measurable in (s, x) and is of class \mathbb{C}^2 in x . Let $\zeta(x)$ represent a \mathbb{C}_0^∞ -function on \mathbb{R}^l with support in S_1 that equals 1 on S and let $\varphi(s, x)$ be a measurable function that is of class \mathbb{C}^∞ in x . On letting $\phi(s, x) = \zeta(x)\varphi(s, x)$ in (6.11), we have that there exists $L_2 > 0$ such that for all $\varphi(s, x)$ and all (X, μ) that satisfy the inequality $\mathbf{I}^{**}(X, \mu) \leq \delta$,

$$\left| \int_0^t \int_{S_1} \text{tr} (c_s(X_s, x) D_{xx}^2 \varphi(s, x)) \zeta(x) m_s(x) dx ds \right| \leq L_2 \left(\int_0^t \left(\sup_{x \in S_1} |\varphi(s, x)|^2 + \sup_{x \in S_1} |D_x \varphi(s, x)|^2 \right) ds \right)^{1/2}.$$

By Sobolev's imbedding, $\mathbb{W}^{2,q}(S_1)$ is continuously imbedded into $\mathbb{W}^{1,\infty}(S_1)$ provided $q > l$ (see, e.g., Theorem 4.12 on p.85 in Adams and Fournier [1]), hence,

$$\left| \int_0^t \int_{S_1} \text{tr} (c_s(X_s, x) D_{xx}^2 \varphi(s, x)) \zeta(x) m_s(x) dx ds \right| \leq L_3 \left(\int_0^t \|\varphi(s, \cdot)\|_{\mathbb{W}^{2,q}(S_1)}^2 ds \right)^{1/2}, \quad (6.12)$$

where $L_3 > 0$. The latter inequality extends to $\varphi(s, \cdot) \in \mathbb{C}^2(\overline{S_1})$. Given a bounded continuous function $f(s, x)$ such that $f(s, \cdot) \in \mathbb{C}_0^\infty(S_1)$, let $\varphi(s, \cdot) \in \mathbb{C}^2(\overline{S_1})$ be such that $\text{tr} (c_s(X_s, x) D_{xx}^2 \varphi(s, x)) = f(s, x)$ and $\varphi(s, x) = 0$ on the boundary of S_1 , see Theorem 6.14 on p.107 of Gilbarg and Trudinger [22]. By Theorem 9.13 on p.239 in Gilbarg and Trudinger [22], where we take $\Omega' = \Omega = S_1$, and on recalling that the norms $\|c_s(u, \cdot)\|_{\mathbb{W}^{2,q}(S_1)}$ are bounded locally in (s, u) , we have that

$$\|\varphi(s, \cdot)\|_{\mathbb{W}^{2,q}(S_1)} \leq L_4 (\|\varphi(s, \cdot)\|_{\mathbb{L}^q(S_1)} + \|f(s, \cdot)\|_{\mathbb{L}^q(S_1)})$$

locally uniformly in s . By Theorem 9.1 on p.220 in Gilbarg and Trudinger [22], $\sup_{x \in S_1} |\varphi(s, x)| \leq L_5 \|f(s, \cdot)\|_{\mathbb{L}^q(S_1)}$ locally uniformly in s . We obtain that there exists $L_6 > 0$ such that $\|\varphi(s, \cdot)\|_{\mathbb{W}^{2,q}(S_1)} \leq L_6 \|f(s, \cdot)\|_{\mathbb{L}^q(S_1)}$. By (6.12), if $q \geq 2$, then, for some $L_7 > 0$,

$$\left| \int_0^t \int_{S_1} f(s, x) \zeta(x) m_s(x) dx ds \right| \leq L_7 \left(\int_0^t \int_{S_1} |f(s, x)|^q dx ds \right)^{1/q}.$$

Since the functions $f(s, x)$ are dense in $\mathbb{L}^q([0, t] \times S_1)$,

$$\left(\int_0^t \int_{S_1} |\zeta(x) m_s(x)|^{q'} dx ds \right)^{1/q'} \leq L_7,$$

which yields the required bound (6.6) if one recalls that $\zeta(x) = 1$ on S . □

Remark 6.1. As a byproduct of the proof, the function $\psi(s, \cdot)$ is an element of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ for almost all s .

We now work toward proving that \mathbf{I}^{**} is the same as \mathbf{I} in Theorem 2.1 and Proposition 2.1. The following lemma will be useful for calculating \mathbf{I}^{**} , cf. Lemma A.2 on p.460 in Puhalskii [41].

Lemma 6.5. *Let V represent a complete separable metric space, let U represent a dense subspace, and let \mathbb{R} -valued function $f(s, y)$ be defined on $\mathbb{R}_+ \times V$, be measurable in s and continuous in y .*

Suppose also that $f(s, \lambda(s))$ is locally integrable with respect to Lebesgue measure for all measurable functions $\lambda(s)$ that assume values in U . Then, for all $t \in \mathbb{R}_+$,

$$\sup_{\lambda(\cdot) \in \Lambda} \int_0^t f(s, \lambda(s)) ds = \int_0^t \sup_{y \in U} f(s, y) ds,$$

where Λ represents the set of measurable functions assuming values in U .

In the rest of the paper we denote D_x by D , divergencies are understood with respect to x . The next lemma is the key to proving that $\sqrt{m_s(\cdot)} \in \mathbb{W}^{1,2}(\mathbb{R}^l)$ in the statement of Theorem 2.1.

Lemma 6.6. *Let $m_s(x)$, where $x \in \mathbb{R}^l$ and $s \in \mathbb{R}_+$, represent an \mathbb{R}_+ -valued measurable function which is a probability density on \mathbb{R}^l and an element of $\mathbb{W}_{loc}^{1,1}(\mathbb{R}^l)$ for almost all s . If, for some $t > 0$ and $L_1 > 0$, we have that $\int_0^t \int_S |Dm_s(x)|^2 / m_s(x) dx ds < \infty$, for all open balls S , and*

$$\int_0^t \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x)m_s(x)) - a_s(X_s, x)m_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \right) dx ds \leq L_1, \quad (6.13)$$

then there exists $L_2 > 0$, which depends on L_1 and t only, such that

$$\int_0^t \int_{\mathbb{R}^l} \frac{|Dm_s(x)|^2}{m_s(x)} dx ds \leq L_2.$$

Proof. Let $\eta(x)$ represent a $[0, 1]$ -valued twice continuously differentiable nonincreasing function defined for $x \geq 0$ such that $\eta(x) = 1$ for $x \in [0, 1]$ and $\eta(x) = 0$ for $x \geq 2$. Let $\eta_r(x) = \eta(|x|/r)$ where $x \in \mathbb{R}^l$ and $r > 0$. We note that the bound in (6.13) extends to functions $h(x)$ from the closure $\mathbb{H}_0^{1,2}(S_{2r+1}, m_s(x) dx) \in \mathbb{C}_0^\infty(S_{2r+1})$ in $\mathbb{W}^{1,2}(S_{2r+1}, m_s(x) dx)$, where S_{2r+1} represents the open ball of radius $2r+1$ centred at the origin in \mathbb{R}^l . Let $\delta > 1$. Since (the restriction of) $\ln(m_s(\cdot) \wedge \delta \vee \delta^{-1})$ to S_{2r+1} is an element of $\mathbb{W}^{1,2}(S_{2r+1}, m_s(x) dx)$ a.e. and since by Lemma 6.2, $\mathbb{W}^{1,2}(S_{2r+1}, m_s(x) dx) = \mathbb{H}^{1,2}(S_{2r+1}, m_s(x) dx)$, we have that $\ln(m_s(\cdot) \wedge \delta \vee \delta^{-1}) \in \mathbb{H}^{1,2}(S_{2r+1}, m_s(x) dx)$, so, $\ln(m_s(\cdot) \wedge \delta \vee \delta^{-1}) \eta_r(\cdot)^2$ is an element of $\mathbb{H}_0^{1,2}(S_{2r+1}, m_s(x) dx)$. Hence, one can take $h(x) = (1/4) \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x)^2$ in (6.13) to obtain

$$\begin{aligned} & \int_0^t \int_{\mathbb{R}^l} \left(\left(\frac{1}{4} \frac{Dm_s(x)}{m_s(x)} \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 + \frac{1}{2} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x) \right)^T \right. \\ & \quad \left(\frac{1}{2} c_s(X_s, x) \frac{Dm_s(x)}{m_s(x)} + \frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) m_s(x) \\ & \quad - \frac{1}{16} \left\| \frac{Dm_s(x)}{m_s(x)} \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 \right\|_{c_s(X_s, x)}^2 m_s(x) \\ & \quad \left. - \frac{1}{4} \left\| \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x) \right\|_{c_s(X_s, x)}^2 m_s(x) \right) dx ds \leq L_1. \end{aligned}$$

Therefore,

$$\begin{aligned}
& \int_0^t \int_{\mathbb{R}^l} \left(\frac{1}{8} \frac{\|Dm_s(x)\|_{c_s(X_s, x)}^2}{m_s(x)} \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 \right. \\
& \quad \left. - \frac{1}{16} \frac{\|Dm_s(x)\|_{c_s(X_s, x)}^2}{m_s(x)} \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^4 \right) dx ds \\
& - \int_0^t \int_{\mathbb{R}^l} \left(\frac{1}{4} Dm_s(x)^T \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) \right. \\
& \quad \left. + \frac{1}{4} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x)^T c_s(X_s, x) Dm_s(x) \right. \\
& \quad \left. + \frac{1}{2} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x)^T \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) m_s(x) \right. \\
& \quad \left. - \frac{1}{4} (\ln(m_s(x) \wedge \delta \vee \delta^{-1}))^2 \eta_r(x)^2 \|D\eta_r(x)\|_{c_s(X_s, x)}^2 m_s(x) \right) dx ds \leq L_1. \quad (6.14)
\end{aligned}$$

We bound the terms on the righthand side. Integration by parts yields

$$\begin{aligned}
& \int_{\mathbb{R}^l} Dm_s(x)^T \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) dx \\
& = - \int_{\mathbb{R}^l} (m_s(x) \wedge \delta \vee \delta^{-1}) \operatorname{div} (\eta_r(x)^2 \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right)) dx \\
& = - \int_{\mathbb{R}^l} (m_s(x) \wedge \delta \vee \delta^{-1}) (\eta_r(x)^2 \operatorname{div} \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) \\
& \quad + 2\eta_r(x) D\eta_r(x)^T \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right)) dx.
\end{aligned}$$

By Condition 2.1, there exists $M_1 > 0$ such that $|\operatorname{div}((1/2) \operatorname{div} c_s(X_s, x) - a_s(X_s, x))| \leq M_1$ and $|(1/2) \operatorname{div} c_s(X_s, x) - a_s(X_s, x)| \leq M_1(1 + |x|)$, for all $0 \leq s \leq t$ and $x \in \mathbb{R}^l$, so

$$\left| \int_{\mathbb{R}^l} (m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x)^2 \operatorname{div} \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) dx \right| \leq M_1$$

and, letting M_2 represent an upper bound on the absolute values of the first derivative of $\eta(x)$,

$$\begin{aligned}
& \left| \int_{\mathbb{R}^l} (m_s(x) \wedge \delta \vee \delta^{-1}) 2\eta_r(x) D\eta_r(x)^T \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) dx \right| \\
& = \left| \int_{\mathbb{R}^l} (m_s(x) \wedge \delta \vee \delta^{-1}) 2\eta_r(x) \frac{1}{r} \frac{x^T}{|x|} D\eta \left(\frac{|x|}{r} \right) \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) dx \right| \\
& \leq 2M_1 M_2 \frac{1}{r} \int_{x \in \mathbb{R}^l: r \leq |x| \leq 2r} m_s(x) (1 + |x|) dx \leq 2M_1 M_2 \frac{2r + 1}{r}
\end{aligned}$$

Therefore,

$$\left| \int_{\mathbb{R}^l} Dm_s(x)^T \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) dx \right| \leq M_1 + 2M_1 M_2 \frac{2r+1}{r}. \quad (6.15)$$

Similarly,

$$\begin{aligned} & \int_{\mathbb{R}^l} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x)^T c_s(X_s, x) Dm_s(x) dx \\ &= - \int_{\mathbb{R}^l} \operatorname{div}(\ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) c_s(X_s, x) D\eta_r(x)) m_s(x) dx \\ &= - \int_{\mathbb{R}^l} \eta_r(x) D\eta_r(x)^T c_s(X_s, x) Dm_s(x) \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) dx \\ &\quad - \int_{\mathbb{R}^l} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \operatorname{div}(\eta_r(x) c_s(X_s, x) D\eta_r(x)) m_s(x) dx. \end{aligned}$$

For the terms on the righthand side, we have that, for suitable $M_3 > 0$,

$$\left| \int_{\mathbb{R}^l} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \operatorname{div}(\eta_r(x) c_s(X_s, x) D\eta_r(x)) m_s(x) dx \right| \leq M_3 \frac{|\ln \delta|}{r}$$

and, for arbitrary $\kappa > 0$,

$$\begin{aligned} & \left| \int_{\mathbb{R}^l} \eta_r(x) D\eta_r(x)^T c_s(X_s, x) Dm_s(x) \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) dx \right| \\ & \leq \frac{1}{2\kappa r^2} \int_{\mathbb{R}^l} \|D\eta\left(\frac{|x|}{r}\right)\|_{c_s(X_s, x)}^2 m_s(x) dx + \frac{\kappa}{2} \int_{\mathbb{R}^l} \frac{\|Dm_s(x)\|_{c_s(X_s, x)}^2}{m_s(x)} \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) \eta_r(x)^2 dx. \end{aligned}$$

Hence,

$$\begin{aligned} & \left| \int_{\mathbb{R}^l} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x)^T c_s(X_s, x) Dm_s(x) dx \right| \\ & \leq M_3 \frac{|\ln \delta|}{r} + \frac{M_2^2}{2\kappa r^2} \int_{\mathbb{R}^l} \|c_s(X_s, x)\| m_s(x) dx + \frac{\kappa}{2} \int_{\mathbb{R}^l} \frac{\|Dm_s(x)\|_{c_s(X_s, x)}^2}{m_s(x)} \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) dx. \end{aligned} \quad (6.16)$$

The remaining two terms on the righthand side of (6.14) are bounded as follows:

$$\begin{aligned} & \left| \int_{\mathbb{R}^l} \ln(m_s(x) \wedge \delta \vee \delta^{-1}) \eta_r(x) D\eta_r(x)^T \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) m_s(x) dx \right| \\ & \leq M_1 M_2 |\ln \delta| \frac{2r+1}{r} \int_{x \in \mathbb{R}^l: |x| \geq r} m_s(x) dx \end{aligned} \quad (6.17)$$

and

$$\int_{\mathbb{R}^l} (\ln(m_s(x) \wedge \delta \vee \delta^{-1}))^2 \eta_r(x)^2 \|D\eta_r(x)\|_{c_s(X_s, x)}^2 m_s(x) dx \leq \frac{M_2^2 (\ln \delta)^2}{r^2} \int_{\mathbb{R}^l} \|c_s(X_s, x)\| m_s(x) dx. \quad (6.18)$$

We obtain by (6.14) – (6.18) that there exists $M_4 > 0$ such that, given arbitrary $\delta > 1$, for all r great enough (depending on δ),

$$\frac{1}{16} \int_0^t \int_{\mathbb{R}^l} \frac{\|Dm_s(x)\|_{c_s(X_s, x)}^2}{m_s(x)} \eta_r(x)^2 \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) (2 - \eta_r(x)^2 - 2\kappa) dx ds \leq L_1 + M_4 t$$

so that assuming $\kappa < 1/2$,

$$\int_0^t \int_{\mathbb{R}^l} \frac{\|Dm_s(x)\|_{c_s(X_s, x)}^2}{m_s(x)} \eta_r(x)^2 \mathbf{1}_{\{\delta^{-1} \leq m_s(x) \leq \delta\}}(x) dx ds \leq \frac{16}{1 - 2\kappa} (L_1 + M_4 t),$$

which implies the assertion of the lemma by letting $r \rightarrow \infty$ and $\delta \rightarrow \infty$. \square

The next theorem establishes the equality $\mathbf{I}^{**}(X, \mu) = \mathbf{I}(X, \mu)$ provided $\mathbf{I}^{**}(X, \mu) < \infty$, $X_0 = \hat{u}$, and $\mathbf{I}_0(\hat{u}) = 0$.

Theorem 6.1. *Suppose that conditions 2.1, 2.2, (2.4b), (2.14c) and (2.14d) hold and that $\mathbf{I}^{**}(X, \mu) < \infty$. Then $\mu(ds, dx) = m_s(x) dx ds$, where $m_s(\cdot) \in \mathbb{P}(\mathbb{R}^l)$ a.e. and $\int_0^t \int_{\mathbb{R}^l} |Dm_s(x)|^2 / m_s(x) dx ds < \infty$ for all $t \in \mathbb{R}_+$. The projection $\Phi_{s, m_s(\cdot), X_s}(x)$ belongs to $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ as a function of x for almost every s , $\Phi_{s, m_s(\cdot), X_s}(x)$ and $\Psi_{s, m_s(\cdot), X_s}(x)$ are measurable in (s, x) , and $\int_0^t \int_{\mathbb{R}^l} \|\Phi_{s, m_s(\cdot), X_s}(x)\|^2 m_s(x) dx ds < \infty$ for all $t \in \mathbb{R}_+$. We also have that*

$$\begin{aligned} \mathbf{I}^{**}(X, \mu) &= \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) \right. \\ &\quad - \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) m_s(x) dx}^2 + \sup_{h \in C_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) \right. \right. \\ &\quad \left. \left. - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \right) dx \right) ds \\ &= \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) - \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) m_s(x) dx}^2 \right. \\ &\quad \left. + \sup_{g \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)} \int_{\mathbb{R}^l} \left(g(x)^T c_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} \right. \right. \right. \\ &\quad \left. \left. \left. - \Phi_{s, m_s(\cdot), X_s}(x) - \Psi_{s, m_s(\cdot), X_s}(x) \lambda \right) - \frac{1}{2} \|g(x)\|_{c_s(X_s, x)}^2 \right) m_s(x) dx \right) ds. \end{aligned} \quad (6.20)$$

The vector $\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) (Dm_s(x)/(2m_s(x)) - \Phi_{s, m_s(\cdot), X_s}(x)) m_s(x) dx$ is in the range of $\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx$ a.e. and the supremum in (6.20) is attained at

$$\hat{\lambda}_s = \left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^\oplus \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx \right) \quad (6.21)$$

and

$$\hat{g}_s(x) = \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) - \Psi_{s, m_s(\cdot), X_s}(x) \hat{\lambda}_s \quad (6.22)$$

so that

$$\mathbf{I}^{**}(X, \mu) = \int_0^\infty \left(\frac{1}{2} \int_{\mathbb{R}^l} \left\| \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right\|_{c_s(X_s, x)}^2 m_s(x) dx + \frac{1}{2} \left\| \dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx \right\|_{\left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^\oplus}^2 \right) ds. \quad (6.23)$$

Proof. We recall the expression (6.1) for $\mathbf{I}^{**}(X, \mu)$, where the supremum is taken over $t \in \mathbb{R}_+$, functions $\lambda(s, X)$ given by (5.1), and $\mathcal{C}^{1,2,2}(\mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^l)$ -functions $f(s, u, x)$ that are compactly supported in x locally uniformly in (t, u) . According to Lemma 6.4, if $\mathbf{I}^{**}(X, \mu) < \infty$, then $\nu_s(dx) = m_s(x) dx$, where $m_s(\cdot) \in \mathbb{W}_{\text{loc}}^{1,1}(\mathbb{R}^l)$, so one can integrate by parts in (6.3) to obtain

$$\begin{aligned} U_t^{\lambda(\cdot), f}(X, \mu) &= \int_0^t \left(\lambda(s, X)^T \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx \right) - \frac{1}{2} \int_{\mathbb{R}^l} \|\lambda(s, X)\|_{C_s(X_s, x)}^2 m_s(x) dx \right. \\ &\quad \left. + \int_{\mathbb{R}^l} Df(s, X_s, x)^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) dx \right. \\ &\quad \left. - \frac{1}{2} \int_{\mathbb{R}^l} \|Df(s, X_s, x)\|_{c_s(X_s, x)}^2 m_s(x) dx - \int_{\mathbb{R}^l} \lambda(s, X)^T G_s(X_s, x) Df(s, X_s, x) m_s(x) dx \right) ds. \quad (6.24) \end{aligned}$$

An approximation argument using mollifiers implies that the supremum will not change if $\lambda(s, X)$ is assumed bounded and measurable in s and if $f(s, u, x)$ is assumed measurable, continuously differentiable in x with bounded first partial derivatives and compactly supported in x locally

uniformly in (s, u) . Therefore, on noting that X is kept fixed,

$$\begin{aligned} \mathbf{I}^{**}(X, \mu) = & \sup \int_0^t \left(\lambda(s)^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) - \frac{1}{2} \|\lambda(s)\|_{\int_{\mathbb{R}^l} c_s(X_s, x) m_s(x) dx}^2 \right. \\ & + \int_{\mathbb{R}^l} D\phi(s, x)^T \left(\frac{1}{2} \operatorname{div}_x (c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) dx - \frac{1}{2} \int_{\mathbb{R}^l} \|D\phi(s, x)\|_{c_s(X_s, x)}^2 m_s(x) dx \\ & \left. - \int_{\mathbb{R}^l} \lambda(s)^T G_s(X_s, x) D\phi(s, x) m_s(x) dx \right) ds, \end{aligned}$$

where the supremum is taken over $t \in \mathbb{R}_+$, bounded measurable functions $\lambda(s)$, and measurable functions $\phi(s, x)$ that are continuously differentiable in x with bounded first partial derivatives and are compactly supported in x locally uniformly in s . By Lemma 6.5, one can optimise with respect to $\lambda(s)$ and $D\phi(s, x)$ inside the ds -integral which yields (6.19). In some more detail, we apply Lemma 6.5 with U being the Cartesian product of the closed ball of radius i in \mathbb{R}^n and of the set $U_i = \{Dh : h \in \mathbb{C}_0^1(\mathbb{R}^l), \sup_{x \in \mathbb{R}^l} |Dh(x)| \leq i \text{ and } h(x) = 0 \text{ if } |x| \geq i\}$ and with V being the Cartesian product of the closed ball of radius i and of the closure of U_i in the space of continuous functions with support in the open ball of radius i centred at the origin in \mathbb{R}^l that are bounded above by i in absolute value, the latter space being endowed with the sup-norm topology, where $i \in \mathbb{N}$, and let $i \rightarrow \infty$.

Integration by parts in (6.10), with $\nu_s(dx) = m_s(x) dx$, yields

$$\int_{\mathbb{R}^l} Dh(x)^T \left(\frac{1}{2} \operatorname{div} (c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) dx = \int_{\mathbb{R}^l} Dh(x)^T c_s(X_s, x) \psi(s, x) m_s(x) dx.$$

On recalling that $\psi(s, \cdot) \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ for almost all s by Remark 6.1, we have that the function $-\psi(s, x)$ represents the orthogonal projection of $c_s(X_s, x)^{-1} (a_s(X_s, x) - (1/2) \operatorname{div} (c_s(X_s, x) m_s(x)) / m_s(x))$ onto $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$. Since by (6.19), Lemma 6.4, and Lemma 6.6, $Dm_s(x) / m_s(x)$ is a member of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ for almost all s , we have that the function $-\psi(s, x) + (1/2) Dm_s(x) / m_s(x)$ belongs to $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$ for almost all s , so, by (2.17a), it equals $\Phi_{s, m_s(\cdot), X_s}(x)$.

We show that $\Phi_{s, m_s(\cdot), X_s}(x)$ and $\Psi_{s, m_s(\cdot), X_s}(x)$ are properly measurable. Let \mathcal{U}_s represent the closure of the set $\{c_s(X_s, \cdot)^{1/2} \sqrt{m_s(\cdot)} Dp(\cdot) : p \in \mathbb{C}_0^\infty(\mathbb{R}^l, \mathbb{R}^n)\}$ in $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^{l \times n})$. Introducing $\varphi_s(x) = c_s(X_s, x)^{-1/2} G_s(X_s, x)^T \sqrt{m_s(x)}$ and $\hat{\varphi}_s(x) = c_s(X_s, x)^{1/2} \Psi_{s, m_s(\cdot), X_s}(x) \sqrt{m_s(x)}$, we have that $\hat{\varphi}_s$ is the orthogonal projection of φ_s onto \mathcal{U}_s (see (2.17b) and (2.3)). By Corollary 8.2.13 on p.317 in Aubin and Frankowska [4], $\hat{\varphi}_s$ is a measurable function from \mathbb{R}_+ to $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^{l \times n})$. (We note that $s \rightarrow \mathcal{U}_s$ is a measurable set-valued map by part vi) of Theorem 8.1.4 on p.310 in Aubin and Frankowska [4].) This implies that the mapping $(s, x) \rightarrow \Psi_{s, m_s(\cdot), X_s}(x)$ is measurable. The reasoning for $\Phi_{s, m_s(\cdot), X_s}$ is similar.

The representation in (6.20) follows from (2.17b), (2.17a), (2.3), and (6.19). Since the function

$$\tilde{g}_s(x) = \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) - \Psi_{s, m_s(\cdot), X_s}(x) \lambda$$

is a member of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$, it attains the supremum in (6.20), which yields

$$\begin{aligned} \mathbf{I}^{**}(X, \mu) &= \int_0^\infty \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) - \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) m_s(x) dx}^2 \right. \\ &\quad \left. + \frac{1}{2} \int_{\mathbb{R}^l} \|\Phi_{s, m_s(\cdot), X_s}(x) - \frac{Dm_s(x)}{2m_s(x)} - \Psi_{s, m_s(\cdot), X_s}(x)\lambda\|_{c_s(X_s, x)}^2 m_s(x) dx \right) ds. \end{aligned} \quad (6.25)$$

Since the matrix $Q_{s, m_s(\cdot)}(u, x) = C_s(u, x) - \|\Psi_{s, m_s(\cdot), u}(x)\|_{c_s(u, x)}^2$ (see (2.16)) is positive semidefinite, the supremum over λ in (6.25) is attained at

$$\begin{aligned} \tilde{\lambda} &= \left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^\oplus \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx \right. \\ &\quad \left. - \int_{\mathbb{R}^l} \Psi_{s, m_s(\cdot), X_s}(x)^T c_s(X_s, x) \left(\Phi_{s, m_s(\cdot), X_s}(x) - \frac{Dm_s(x)}{2m_s(x)} \right) m_s(x) dx \right) \end{aligned}$$

and equals

$$\frac{1}{2} \int_{\mathbb{R}^l} \left\| \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s} \right\|_{c_s(X_s, x)}^2 m_s(x) dx + \frac{1}{2} \|\tilde{\lambda}\|_{\left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^\oplus},$$

provided

$$\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} \Psi_{s, m_s(\cdot), X_s}(x)^T c_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx$$

is in the range of $\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx$ a.e. Otherwise, the supremum equals infinity. The fact that $\tilde{\lambda} = \hat{\lambda}_s$ and the expression in (6.23) follow from (2.17b) and (6.25). The properties that $\int_0^t \int_{\mathbb{R}^l} |Dm_s(x)|^2 / m_s(x) dx ds$ and $\int_0^t \int_{\mathbb{R}^l} \|\Phi_{s, m_s(\cdot), X_s}(x)\|^2 dx ds$ are finite follow from Lemma 6.6, (6.19), and (6.23). \square

Motivated by (6.23) in Theorem 6.1, let us introduce, provided $\mathbf{I}^{**}(X, \mu) < \infty$ so that $\mu(ds, dx) = m_s(x) dx ds$ and $\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) (Dm_s(x)/(2m_s(x)) - \Phi_{s, m_s(\cdot), X_s}(x)) m_s(x) dx$ is in the range of $\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx$ a.e.,

$$\begin{aligned} \mathbf{I}_t^{**}(X, \mu) &= \int_0^t \left(\frac{1}{2} \int_{\mathbb{R}^l} \left\| \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right\|_{c_s(X_s, x)}^2 m_s(x) dx \right. \\ &\quad \left. + \frac{1}{2} \left\| \dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} \right. \right. \right. \\ &\quad \left. \left. \left. - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx \right\|_{\left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^\oplus}^2 \right) ds. \end{aligned} \quad (6.26)$$

Similarly to the proof of Theorem 6.1, we also have that

$$\begin{aligned} \mathbf{I}_t^{**}(X, \mu) &= \int_0^t \sup_{\lambda \in \mathbb{R}^n} \left(\lambda^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) - \frac{1}{2} \|\lambda\|_{\int_{\mathbb{R}^l} C_s(X_s, x) m_s(x) dx}^2 \right. \\ &\quad + \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) \right. \\ &\quad \left. \left. - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x) m_s(x)}^2 \right) dx \right) ds. \end{aligned} \quad (6.27)$$

For the proof of Theorem 8.1, it will be needed to extend (X, μ) defined on $[0, t]$ past t in such a way that $\mathbf{I}_t^{**}(X, \mu) = \mathbf{I}^{**}(X, \mu)$. That is done in the following lemma which also concerns the zeros of $\mathbf{I}^{**}(X, \mu)$.

Lemma 6.7. *For $t \in \mathbb{R}_+$ and $z \in \mathbb{R}^n$, the system of equations*

$$\dot{X}_s = \int_{\mathbb{R}^l} A_{s+t}(X_s, x) m_s(x) dx, \quad X_0 = z, \quad (6.28)$$

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr}(c_{s+t}(X_s, x) D^2 p(x)) + a_{s+t}(X_s, x)^T Dp(x) \right) m_s(x) dx = 0 \quad (6.29)$$

where $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$ is otherwise arbitrary, has a solution $(X^\dagger, (m_s^\dagger(x)))$ such that X^\dagger is locally Lipschitz continuous, $m_s^\dagger(x)$ is measurable, and $m_s^\dagger(\cdot) \in \mathbb{P}(\mathbb{R}^l)$. If, given (X, μ) such that $\mathbf{I}^{**}(X, \mu) < \infty$, one defines $(\hat{X}, \hat{\mu})$ by the relations $\hat{X}_s = X_s$ and $\hat{\mu}_s = \mu_s$ for $s \leq t$, and $\hat{X}_s = X_{s-t}^\dagger$ and $\hat{\mu}_s(dx) = \mu_t(dx) + \int_0^{s-t} m_r^\dagger(dx) dr$ for $s > t$, where $z = X_t$, then $\mathbf{I}^{**}(\hat{X}, \hat{\mu}) = \mathbf{I}_t^{**}(X, \mu)$. In particular, if $t = 0$, then $\mathbf{I}^{**}(X^\dagger, \mu^\dagger) = 0$.

Proof. Since $a_s(u, x)$ is locally bounded, since $c_s(u, x)$ is bounded, is positive definite, and is of class \mathbb{C}^1 in x , and since $a_s(u, x)^T x/|x| \rightarrow -\infty$ as $|x| \rightarrow \infty$ by (2.4b), applications of Theorem 1.4.1 in Bogachev, Krylov, and Röckner [8] (with $V(x) = \sqrt{1 + |x|^2}$) and of Theorem 2.2 and Proposition 2.4 in Metafuno, Pallara, and Rhandi [31], show that for every $s, t \in \mathbb{R}_+$ and $u \in \mathbb{R}^n$ there exists a unique probability density $m_s(x)$ satisfying the equation

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr}(c_{s+t}(u, x) D^2 p(x)) + Dp(x)^T a_{s+t}(u, x) \right) m_s(x) dx = 0. \quad (6.30)$$

We apply the method of successive approximations: let $X_s^0 = z$ and, for $i \in \mathbb{N}$,

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr}(c_{s+t}(X_s^i, x) D^2 p(x)) + Dp(x)^T a_{s+t}(X_s^i, x) \right) m_s^i(x) dx = 0, \quad (6.31)$$

$$\dot{X}_s^{i+1} = \int_{\mathbb{R}^l} A_{s+t}(X_s^{i+1}, x) m_s^i(x) dx, \quad X_0^{i+1} = z. \quad (6.32)$$

We note that $m_s^i(x)$ is a measurable function of (s, x) (one can use, e.g., Theorem 8.2.9 on p.315 in Aubin and Frankowska [4]). By (2.14c), we have that given $L > 0$, there exists $M > 0$ such that a.e. in $s \in [0, L]$, $d|X_s^{i+1}|^2/ds \leq M(1 + |X_s^{i+1}|^2)$. Gronwall's inequality implies that

$\sup_{i \in \mathbb{N}} \sup_{s \in [0, L]} |X_s^i| < \infty$. By (6.32) and (2.14b), the derivatives \dot{X}_s^{i+1} are bounded uniformly in $i \in \mathbb{N}$ and $s \in [0, L]$, so the sequence $(X_s^i, s \in [0, L])$ is relatively compact for the uniform norm on $[0, L]$. Let X_s^\dagger represent a limit point. It is a locally Lipschitz continuous function.

As in Metafune, Pallara, and Rhandi [31, Proposition 2.4], we have that, for arbitrary $\delta > 0$ and $L > 0$,

$$\sup_{s \in [0, L]} \sup_{i \in \mathbb{N}} \int_{\mathbb{R}^l} e^{\delta|x|} m_s^i(x) dx < \infty. \quad (6.33)$$

In some more detail, let for a function p which is twice differentiable at x ,

$$\mathcal{L}_s^i p(x) = \frac{1}{2} \text{tr} (c_{s+t}(X_s^i, x) D^2 p(x)) + Dp(x)^T a_{s+t}(X_s^i, x).$$

Since, for $|x| > 0$,

$$\mathcal{L}_s^i e^{\delta|x|} = \left(\frac{1}{2} \text{tr} (c_{s+t}(X_s^i, x) \left(\frac{\delta}{|x|} \left(I - \frac{xx^T}{|x|^2} \right) + \delta^2 \frac{xx^T}{|x|^2} \right)) + \delta a_{s+t}^i(X_s^i, x)^T \frac{x}{|x|} \right) e^{\delta|x|},$$

where I represents the $l \times l$ identity matrix, and $\sup_{i \in \mathbb{N}} a_{s+t}(X_s^i, x)^T x / |x| \rightarrow -\infty$ as $|x| \rightarrow \infty$, there exists $R > 1$ such that $\mathcal{L}_s^i e^{\delta|x|} \leq 0$ and $e^{\delta|x|} \leq |\mathcal{L}_s^i e^{\delta|x|}|$ for all $s \in [0, t]$ and all $i \in \mathbb{N}$ provided $|x| > R$. Let F be a $C^\infty(\mathbb{R}^l)$ -function such that $F(x) = e^{\delta|x|}$ if $|x| \geq 1$. Arguing as in the proof of Proposition 2.3 in Metafune, Pallara, and Rhandi [31], one can see that

$$\int_{x \in \mathbb{R}^l: |x| > R} |\mathcal{L}_s^i e^{\delta|x|}| m_s^i(x) dx \leq \int_{x \in \mathbb{R}^l: |x| \leq R} \mathcal{L}_s^i F(x) m_s^i(x) dx$$

so that

$$\int_{x \in \mathbb{R}^l: |x| > R} e^{\delta|x|} m_s^i(x) dx \leq \int_{x \in \mathbb{R}^l: |x| \leq R} \mathcal{L}_s^i F(x) m_s^i(x) dx, \quad (6.34)$$

which implies (6.33).

Hence, given $s \in [0, L]$, the sequence of probability measures $m_s^i(x) dx$ is tight. Proposition 2.16 in Bogachev, Krylov, and Röckner [7] implies that the $m_s^i(x)$ converge in the variation norm along a subsequence to density $m_s^\dagger(x)$. Since the local \mathbb{L}^q -norms of the $m_s^i(x)$ are uniformly bounded for all $q > 1$ (see (2.26) in Bogachev, Krylov, and Röckner [7]), $\sup_{i \in \mathbb{N}} |a_{s+t}(X_s^i, x)|$ grows at most linearly with $|x|$ by Lipschitz continuity and the fact that $\sup_{x \in \mathbb{R}^l} \sup_{i \in \mathbb{N}} \|c_{s+t}^i(X_s^i, x)\| < \infty$ (see Condition 2.1), and $\sup_{i \in \mathbb{N}} \int_{\mathbb{R}^l} e^{\delta|x|} m_s^i(x) dx < \infty$, on taking a limit in (6.31), we have by dominated convergence that (6.29) holds. Since density $m_s^\dagger(x)$ is specified uniquely by (6.29) $m_s^i(x) \rightarrow m_s^\dagger(x)$ as $i \rightarrow \infty$ along a subsequence such that the X^i converge to X^\dagger . Since $\sup_{i \in \mathbb{N}} \sup_{x \in \mathbb{R}^l} |A_{s+t}(X_s^i, x)| < \infty$ by (2.14b), a similar reasoning shows that taking the above subsequential limit in (6.32) obtains (6.28). Since (6.33) implies that $\int_{\mathbb{R}^l} |a_s(X_s, x)|^2 m_s^\dagger(x) dx < \infty$, by Theorem 1.1 in Bogachev, Krylov, and Röckner [6], $\sqrt{m_s^\dagger(\cdot)} \in \mathbb{W}^{1,2}(\mathbb{R}^l)$.

On noting that (6.30) can be written as

$$\int_{\mathbb{R}^l} Dp(x)^T (a_{s+t}(u, x) - \frac{1}{2} \text{div} c_{s+t}(u, x)) m_s(x) dx = \frac{1}{2} \int_{\mathbb{R}^l} Dp(x)^T c_{s+t}(u, x) Dm_s(x) dx,$$

we have that $\Phi_{s+t, m_s^\dagger(\cdot), X_s^\dagger}(x) = Dm_s^\dagger(x) / (2m_s^\dagger(x))$ which implies, by (6.23) and (6.26), that $\mathbf{I}^{**}(\hat{X}, \hat{\mu}) = \mathbf{I}_t^{**}(X, \mu)$. \square

Remark 6.2. By Proposition 2.4 and Theorem 6.1 (with $\beta = 1$) in Metafunne, Pallari, and Randi [31], $m_s^\dagger(\cdot)$ decays exponentially at infinity. It is also positive and Hölder continuous, see Bogachev, Krylov, and Röckner [7, Theorem 2.8, Corollary 2.10, Corollary 2.11] and Bogachev, Krylov, and Röckner [8].

7 Identifying the large deviation function

The purpose of this section is to show that $\tilde{\mathbf{I}} = \mathbf{I}^{**}$ for sufficiently regular functions (X, μ) . More specifically, we will prove the following theorem.

Theorem 7.1. *Suppose that conditions 2.1, 2.2, (2.4b), and (2.14d) hold. Suppose that $\tilde{\mathbf{I}}$ is a large deviation function that satisfies the assertion of Theorem 5.1 and is such that $\tilde{\mathbf{I}}(X, \mu) = \infty$ unless $X_0 = \hat{u}$. Suppose that $(\hat{X}, \hat{\mu})$ is such that $\hat{X}_0 = \hat{u}$, $\mathbf{I}^{**}(\hat{X}, \hat{\mu}) < \infty$, \hat{X} is locally Lipschitz continuous and that $\hat{m}_s(x) = \hat{\mu}(ds, dx)/(ds dx)$ is of the form*

$$\hat{m}_s(x) = M_s(\tilde{m}_s(x)\hat{\eta}^2\left(\frac{|x|}{r}\right) + e^{-\alpha|x|}(1 - \hat{\eta}^2\left(\frac{|x|}{r}\right)))$$

where $\tilde{m}_s(x)$ is a probability density in x which is locally bounded away from zero and belongs to $\mathbb{C}^1(\mathbb{R}^l)$ as a function of x , with $|Dm_s(x)|$ being locally bounded in (s, x) , $\hat{\eta}(y)$ is a nonincreasing $[0, 1]$ -valued $\mathbb{C}_0^1(\mathbb{R}_+)$ -function, where $y \in \mathbb{R}_+$, that equals 1 for $y \in [0, 1]$ and equals 0 for $y \geq 2$, $r > 0$, $\alpha > 0$, and M_s is the normalising constant. Then, for given $\tilde{m}_s(x)$, $\hat{\eta}(y)$, and r , there exists $\alpha_0 > 0$ such that $\tilde{\mathbf{I}}(\hat{X}, \hat{\mu}) = \mathbf{I}^{**}(\hat{X}, \hat{\mu})$ for all $\alpha > \alpha_0$.

We assume throughout the section the hypotheses of Theorem 7.1 to hold. We start by extending the assertion of Theorem 5.1 to a larger set of functions $(\lambda(\cdot), f)$. For economy of notation, we denote $\gamma = (X, \mu)$ and recall that Γ represents the set of γ such that X is absolutely continuous and μ admits density $m_s(x)$ that is an element of $\mathbb{P}(\mathbb{R}^l)$ in x , for almost all s . Let $\lambda(s, X)$, where $s \in \mathbb{R}_+$ and $X \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$, represent an \mathbb{R}^n -valued measurable function and let $h_s(u, x)$, where $s \in \mathbb{R}_+$, $u \in \mathbb{R}^n$ and $x \in \mathbb{R}^l$, represent an \mathbb{R} -valued measurable function, which is an element of $\mathbb{W}_{\text{loc}}^{1,1}(\mathbb{R}^l)$ in x and is of bounded support in x locally uniformly over (s, u) . If, for all $t \in \mathbb{R}_+$ and all $\gamma \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, $\int_0^t \int_{\mathbb{R}^l} (|\lambda(s, X)|^2 + |Dh_s(X_s, x)|^2) \mu(dx, ds) < \infty$, we define, given $N \in \mathbb{N}$,

$$\tau^N(\gamma) = \inf\{t \in \mathbb{R}_+ : \int_0^t \int_{\mathbb{R}^l} \left(\|\lambda(s, X)\|_{C_s(X_s, x)}^2 + \|Dh_s(X_s, x)\|_{C_s(X_s, x)}^2 \right) \mu(dx, ds) + X_t^* + t \geq N\} \quad (7.1a)$$

and, provided $\gamma \in \Gamma$,

$$\begin{aligned} \theta^N(\gamma) &= \int_0^{\tau^N(\gamma)} \left(\lambda(s, X)^T \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx \right) - \frac{1}{2} \|\lambda(s, X)\|_{C_s(X_s, x)}^2 \int_{\mathbb{R}^l} c_s(X_s, x) m_s(x) dx \right. \\ &\quad \left. + \int_{\mathbb{R}^l} (Dh_s(X_s, x))^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) \right. \\ &\quad \left. - \frac{1}{2} \|Dh_s(X_s, x)\|_{C_s(X_s, x)}^2 \int_{\mathbb{R}^l} m_s(x) dx - \int_{\mathbb{R}^l} \lambda(s, X)^T G_s(X_s, x) Dh_s(X_s, x) m_s(x) dx \right) ds. \end{aligned} \quad (7.1b)$$

For the latter definition, we assume that, in addition,

$$\int_0^t (|\dot{X}_s|^2 + \int_{\mathbb{R}^l} \frac{|Dm_s(x)|^2}{m_s(x)} dx) ds < \infty, \quad (7.2)$$

for all $t \in \mathbb{R}_+$, and use the piece of notation $X_t^* = \sup_{s \in [0, t]} |X_s|$. (The definition of $\theta^N(\gamma)$ is modelled on the expression for $U_t^{\lambda(\cdot), f}(X, \mu)$ in (6.24).) We note that $\tau^N(\gamma) \leq N$. Furthermore, we have the following lemma, for which we reuse the piece of notation of Theorem 3.4 that, for $\delta \in \mathbb{R}_+$,

$$K_\delta = \{\gamma : \tilde{\mathbf{I}}(\gamma) \leq \delta\}$$

and recall that K_δ is a compact in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ and that $K_\delta \subset \Gamma$. Theorem 6.1 implies that (7.2) holds on K_δ . For the definition of the essential supremum of a family of measurable functions used in the next lemma, see, e.g., Proposition II.4.1 on p.44 of Neveu [34].

Lemma 7.1. *Let $\lambda^i(s, X)$ and $h_s^i(u, x)$ be sequences of functions satisfying the same hypotheses as $\lambda(s, X)$ and $h_s(u, x)$, respectively, and let $\tau^{N, i}(\gamma)$ and $\theta^{N, i}(\gamma)$ be defined by the respective equations (7.1a) and (7.1b), with $\lambda^i(s, X)$ and $h_s^i(u, x)$ being substituted for $\lambda(s, X)$ and $h_s(u, x)$, respectively. If, in addition, the functions $h_s^i(u, x)$ are of bounded support in x uniformly over i and locally uniformly over (s, u) , then*

$$\int_0^N \operatorname{ess\,sup}_{\gamma \in K_\delta} |\lambda(s, X)|^2 ds + \int_0^N \operatorname{ess\,sup}_{\gamma \in K_\delta} \int_{\mathbb{R}^l} |Dh_s(X_s, x)|^2 m_s(x) dx ds < \infty, \quad (7.3)$$

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} \int_0^N |\lambda(s, X) - \lambda^i(s, X)|^2 ds = 0, \quad (7.4a)$$

and

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} \int_0^N \int_{\mathbb{R}^l} |Dh_s(X_s, x) - Dh_s^i(X_s, x)|^2 m_s(x) dx ds = 0, \quad (7.4b)$$

then

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} |\tau^N(\gamma) - \tau^{N, i}(\gamma)| = 0 \quad (7.5a)$$

and

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} |\theta^N(\gamma) - \theta^{N, i}(\gamma)| = 0. \quad (7.5b)$$

Proof. Let us note that under the hypotheses,

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} \int_0^N \int_{\mathbb{R}^l} \left| \|\lambda^i(s, X)\|_{C_s(X_s, x)}^2 - \|\lambda(s, X)\|_{C_s(X_s, x)}^2 \right| m_s(x) dx ds = 0, \quad (7.6a)$$

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} \int_0^N \int_{\mathbb{R}^l} \left| \|Dh_s^i(X_s, x)\|_{c_s(X_s, x)}^2 - \|Dh_s(X_s, x)\|_{c_s(X_s, x)}^2 \right| m_s(x) dx ds = 0, \quad (7.6b)$$

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} \int_0^N \left| (\lambda^i(s, X) - \lambda(s, X))^T \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx \right) \right| ds = 0, \quad (7.6c)$$

and

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} \int_0^N \left| \int_{\mathbb{R}^l} (Dh_s^i(X_s, x) - Dh_s(X_s, x))^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) dx \right| ds = 0. \quad (7.6d)$$

The first two convergences are implied by (7.4a), (2.14d), and (7.4b), (2.14a), respectively, and (7.3). The convergence in (7.6c) follows via Cauchy's inequality from (7.4a) and the fact that, according to (6.23) in Theorem 6.1,

$$\sup_{(X, \mu): \mathbf{I}^{**}(X, \mu) \leq \delta} \int_0^N \left| \dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx \right|^2 ds < \infty. \quad (7.7)$$

Similarly, (7.6d) is a consequence of (7.4b), if one recalls that the functions involved are of uniformly bounded support in x and takes into account part (6.5) of Lemma 6.4.

The convergence in (7.5a) follows from (7.6a), (7.6b) and the observation that by (7.1a)

$$\begin{aligned} |\tau^N(\gamma) - \tau^{N,i}(\gamma)| &\leq \int_0^N \int_{\mathbb{R}^l} \left(\|\lambda(s, X)\|_{C_s(X_s, x)}^2 - \|\lambda^i(s, X)\|_{C_s(X_s, x)}^2 \right. \\ &\quad \left. + \|Dh_s(X_s, x)\|_{c_s(X_s, x)}^2 - \|Dh_s^i(X_s, x)\|_{c_s(X_s, x)}^2 \right) m_s(x) dx ds. \end{aligned}$$

The convergence in (7.5b) follows by (7.1b), (7.5a), (7.6a)–(7.6d), and (7.7), if one notes that, thanks to (7.3),

$$\sup_{\gamma \in K_\delta} \int_0^t |\lambda(s, X)|^2 ds, \quad \sup_{\gamma \in K_\delta} \int_0^t \int_{\mathbb{R}^l} |Dh_s(X_s, x)|^2 m_s(x) dx ds,$$

and

$$\sup_{\gamma \in K_\delta} \int_0^t \int_{\mathbb{R}^l} Dh_s(X_s, x)^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) dx ds$$

are continuous functions of $t \in [0, N]$. □

Lemma 7.2. *Let $\lambda_s(u)$ represent an \mathbb{R}^n -valued function of $(s, u) \in \mathbb{R}_+ \times \mathbb{R}^n$, which is measurable in s , is continuous in u for almost all s and is such that $\int_0^N \sup_{|u| \leq L} |\lambda_s(u)|^2 ds < \infty$ for all $L > 0$. Suppose that the function $h_s(u, x)$, in addition to being measurable and being of class $\mathbb{W}_{loc}^{1,1}$ in x , vanishes when x is outside of some open ball in \mathbb{R}^l locally uniformly in (s, u) , that the function $Dh_s(u, x)$ is continuous in (u, x) for almost all $s \in \mathbb{R}_+$, and that $\int_0^N \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_{\mathbb{R}^l} |Dh_s(u, x)|^q dx ds < \infty$ for all $q > 1$ and $L > 0$. Then, under the hypotheses of Theorem 7.1, the function $\theta^N(\gamma)$, where $\lambda(s, X) = \lambda_s(X_s)$, is continuous in γ when restricted to K_δ ,*

$$\sup_{\gamma \in \Gamma} (\theta^N(\gamma) - \tilde{\mathbf{I}}(\gamma)) = 0$$

and the latter supremum is attained. Furthermore,

$$\sup_{\gamma \in K_{2N+2}} (\theta^N(\gamma) - \tilde{\mathbf{I}}(\gamma)) = 0.$$

Proof. The functions $|\lambda_s(u)| \mathbf{1}_{\{|\lambda_s(u)| \geq r\}}(s, u)$ are upper semicontinuous in u and monotonically decreasing in r , so by Dini's theorem $|\lambda_s(u)|^2 \mathbf{1}_{\{|\lambda_s(u)| \geq r\}}(s, u) \rightarrow 0$ as $r \rightarrow \infty$ uniformly on $\{u \in \mathbb{R}^n : |u| \leq L\}$. Let r_i be such that $\int_0^N \sup_{u \in \mathbb{R}^n: |u| \leq L} |\lambda_s(u)|^2 \mathbf{1}_{\{|\lambda_s(u)| \geq r_i\}}(s, u) ds < 1/i$, where $L = \sup_{\gamma \in K_\delta} \sup_{s \in [0, t]} |X_s|$ and $i \in \mathbb{N}$. Since $\lambda_s(u)$ is a Carathéodory function, as a consequence of the Scorza-Dragnoni theorem, see, e.g., p.235 in Ekeland and Temam [14], there exists a measurable function $\check{\lambda}_s^i(u)$ that is continuous in (s, u) , is bounded above in absolute value by r_i , and is such that $\int_0^N \mathbf{1}_{\{\lambda_s(\cdot) \neq \check{\lambda}_s^i(\cdot)\}}(s) ds < 2/(ir_i^2)$. Letting $\lambda^i(s, X) = \check{\lambda}_{[j(i)s]/j(i)}^i(X_{[j(i)s]/j(i)})$, where $j(i)$ is great enough and $j(i) \rightarrow \infty$ as $i \rightarrow \infty$, we have that (7.4a) holds.

Similarly, let

$$h_s^i(u, x) = \int_{\mathbb{R} \times \mathbb{R}^l} \rho_{1/i}(\tilde{s}, y) h_{s-\tilde{s}}(u, x - y) d\tilde{s} dy,$$

where $\rho_\kappa(\tilde{s}, y) = (\tilde{\rho}_1(\tilde{s}/\kappa)/\kappa)(\tilde{\rho}_2(y/\kappa)/\kappa^l)$, $\tilde{\rho}_1(\tilde{s})$ is a mollifier on \mathbb{R} such that $\tilde{\rho}_1(\tilde{s}) = 0$ if $|\tilde{s}| > 1$, $\tilde{\rho}_2(y)$ is a mollifier on \mathbb{R}^l such that $\tilde{\rho}_2(y) = 0$ if $|y| > 1$, and $h_s(u, x) = 0$ if $s \leq 0$. The function $h_s^i(u, x)$ is an element of $\mathbb{C}^\infty(\mathbb{R}_+ \times \mathbb{R}^l)$ in (s, x) for all u and $Dh_s^i(u, x) = \int_{\mathbb{R} \times \mathbb{R}^l} \rho_{1/i}(\tilde{s}, y) Dh_{s-\tilde{s}}(u, x - y) d\tilde{s} dy$, cf. Theorem 2.29 on p.36 in Adams and Fournier [1]. In addition, $Dh_s^i(u, x)$ is a continuous function for every i . We also have that, for all open balls S , all $L > 0$ and all $q > 1$,

$$\lim_{i \rightarrow \infty} \int_0^N \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_S |Dh_s(u, x) - Dh_s^i(u, x)|^q dx ds = 0, \quad (7.8)$$

which can be shown as follows. If, in addition, $Dh_s(u, x)$ is continuous in all variables, then $Dh_s^i(u, x)$ converges to $Dh_s(u, x)$ locally uniformly in (s, u, x) , cf. Theorem 2.29 on p.36 in Adams and Fournier [1], so, (7.8) holds. In the general case, in analogy with the above reasoning, there exist r_j such that $\int_0^{N+1} \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_{\tilde{S}} |Dh_s(u, x)|^q \mathbf{1}_{\{|Dh_s(u, x)| \geq r_j\}}(s, u, x) dx ds < 1/j$ where \tilde{S} represents the open ball in \mathbb{R}^l centred at the origin of radius one greater than that of S , and there exists a continuous function $\check{h}_s^j(u, x)$, which is bounded above in absolute value by r_j , such that $\int_0^{N+1} \mathbf{1}_{\{Dh_s(\cdot, \cdot) \neq \check{h}_s^j(\cdot, \cdot)\}}(s) ds < 2/(jr_j^q)$. Calculations show that

$$\int_0^N \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_S |Dh_s(u, x) - \check{h}_s^j(u, x)|^q dx ds \leq \frac{2^{q-1}}{j} + \frac{2^{q+1}}{j}$$

and

$$\int_0^N \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_S |Dh_s^i(u, x) - \int_{\mathbb{R} \times \mathbb{R}^l} \rho_{1/i}(\tilde{s}, y) \check{h}_{s-\tilde{s}}^j(u, x-y) d\tilde{s} dy|^q dx ds \leq \frac{2^{q-1}}{j} + \frac{2^{q+1}V(S)}{j},$$

where $V(S)$ represents the volume of the ball S . Hence, (7.8) holds.

By an application of Hölder's inequality, it follows from (7.8), (6.6) in Lemma 6.4 and $h_s(u, x)$ having compact support in x locally uniformly over (s, u) that (7.4b) holds. Also, (7.3) holds.

Let $\tau^{N,i}$ and $\theta^{N,i}$ be defined as in Lemma 7.1. The functions $h_s^i(u, x)$, $\lambda^i(s, X)$, and $\tau^{N,i}(X, \mu)$ satisfy the requirements imposed on the respective functions $f(s, u, x)$, $\lambda(s, X)$, and $\tau(X, \mu)$ when deriving (5.4). Furthermore, integration by parts on the righthand side of (5.3) with $\mu(ds, dx) = m_s(x) dx ds$, implies that $\theta^{N,i}(\gamma) = U_{N \wedge \tau^{N,i}(\gamma)}^{\lambda^i(\cdot), h^i}(\gamma)$ provided $\gamma \in \Gamma$. In addition, by (7.1a), $|X_{\tau^{N,i}(\gamma)}| \leq N$ and $\tau^{N,i}(\gamma)$ is a continuous function of $\gamma \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$, cf. Theorem 2 on p.510 and Theorem 3 on p.511 in Liptser and Shiryaev [29]. We obtain by equation (5.4) of Theorem 5.1 and the fact that $\tilde{\mathbf{I}}(\gamma) = \infty$ unless $\gamma \in \Gamma$ (see Theorem 6.1) that

$$\sup_{\gamma \in \Gamma} (\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) = 0. \quad (7.9)$$

Let us show that, for all $\delta > 2N + 1$,

$$\sup_{\gamma \in K_\delta} (\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) = 0. \quad (7.10)$$

Let, for $\gamma \in \Gamma$,

$$\begin{aligned} \tilde{\theta}^{N,i}(\gamma) &= \int_0^{\tau^{N,i}(\gamma)} \left(2\lambda^{i,j}(s, X)^T (\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx) - \frac{1}{2} \int_{\mathbb{R}^l} \|2\lambda^{i,j}(s, X)\|_{C_s(X_s, x)}^2 m_s(x) dx \right. \\ &\quad \left. + \int_{\mathbb{R}^l} (2Dh_s^i(X_s, x))^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) - a_s(X_s, x) m_s(x) \right) \right. \\ &\quad \left. - \frac{1}{2} \|2Dh_s^i(X_s, x)\|_{C_s(X_s, x)}^2 m_s(x) dx - \int_{\mathbb{R}^l} 4\lambda^{i,j}(s, X)^T G_s(X_s, x) Dh_s^i(X_s, x) m_s(x) dx \right) ds. \end{aligned}$$

By (5.3), $\tilde{\theta}^{N,i}(\gamma) = U_{N \wedge \tau^{N,i}(\gamma)}^{2\lambda^i(\cdot), 2h^i}(\gamma)$, provided $\gamma \in \Gamma$, so in analogy with (7.9),

$$\sup_{\gamma \in \Gamma} (\tilde{\theta}^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) = 0.$$

On noting that $\tilde{\theta}^{N,i}(\gamma) \geq 2\theta^{N,i}(\gamma) - 2N$, we have that, for $M > 0$,

$$\begin{aligned} \sup_{\gamma: \theta^{N,i}(\gamma) \geq M} (\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) &\leq \sup_{\gamma: \theta^{N,i}(\gamma) \geq M} (2\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) - M \\ &\leq \sup_{\gamma: \theta^{N,i}(\gamma) \geq M} (\tilde{\theta}^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) + 2N - M \leq 2N - M. \end{aligned}$$

Since, by (7.9),

$$0 = \sup_{\gamma \in \Gamma} (\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) \leq \sup_{\gamma \in K_\delta} (\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) \vee \sup_{\gamma: \theta^{N,i}(\gamma) \geq M} (\theta^{N,i}(\gamma) - \tilde{\mathbf{I}}(\gamma)) \vee (M - \delta),$$

we conclude, on choosing $M = 2N + 1$, that (7.10) holds for $\delta > 2N + 1$.

Since by Lemma 7.1, for arbitrary $\delta \in \mathbb{R}_+$,

$$\lim_{i \rightarrow \infty} \sup_{\gamma \in K_\delta} |\theta^N(\gamma) - \theta^{N,i}(\gamma)| = 0, \quad (7.11)$$

we obtain by (7.10) that

$$\sup_{\gamma \in K_\delta} (\theta^N(\gamma) - \tilde{\mathbf{I}}(\gamma)) = 0. \quad (7.12)$$

Since $\theta^{N,i}(\gamma) = U_{\tau^{N,i}(\gamma)}^{\lambda^i(\cdot), h^i}(\gamma)$, the latter function is continuous in γ , and K_δ is compact, (7.11) implies that $\theta^N(\gamma)$ is continuous on K_δ . Since $\tilde{\mathbf{I}}(\gamma)$ is a lower semicontinuous function of γ , the supremum in (7.12) is attained. On the other hand, if $\tilde{\mathbf{I}}(\gamma) < \infty$, then by (7.9) and (7.11), $\sup_{\gamma \in \Gamma} (\theta^N(\gamma) - \tilde{\mathbf{I}}(\gamma)) \leq 0$. □

In order to prove that $\tilde{\mathbf{I}} = \mathbf{I}^{**}$, we will use $\hat{\lambda}$ and \hat{g} defined in (6.21) and (6.22), respectively, as $\lambda_s(u)$ and $Dh_s(u, x)$ in the preceding lemma. We therefore need $\Phi_{s, m_s(\cdot), u}$ and $\Psi_{s, m_s(\cdot), u}$ to be sufficiently regular. The next lemma addresses both regularity and growth-rate properties.

Lemma 7.3. *Suppose that conditions 2.1, 2.2, (2.4b), and (2.14d) hold. Let $m_s(x)$ represent an \mathbb{R}_+ -valued measurable function that is a probability density in x for almost every s . Suppose $m_s(x)$ is bounded away from zero on bounded sets of (s, x) , $m_s(\cdot) \in \mathcal{C}^1(\mathbb{R}^l)$, with $|Dm_s(x)|$ being locally bounded in (s, x) , and $m_s(x) = M_s e^{-\alpha|x|}$ for all $|x|$ great enough locally uniformly in s , where $\alpha > 0$. Then there exist \mathbb{R} -valued measurable function $w_s(u, x)$ and \mathbb{R}^n -valued measurable function $v_s(u, x)$ such that $w_s(u, \cdot) \in \mathbb{W}_{loc}^{2,q}(\mathbb{R}^l)$ and $v_s(u, \cdot) \in \mathbb{W}_{loc}^{2,q}(\mathbb{R}^l, \mathbb{R}^n)$, where $q > 1$ is otherwise arbitrary, $\Phi_{s, m_s(\cdot), u}(\cdot) = Dw_s(u, \cdot)$ and $\Psi_{s, m_s(\cdot), u}(\cdot) = Dv_s(u, \cdot)$ for almost all $s \in \mathbb{R}_+$ and all $u \in \mathbb{R}^n$, i.e.,*

$$\int_{\mathbb{R}^l} Dp(x)^T (a_s(u, x) - \frac{1}{2} \operatorname{div} c_s(u, x)) m_s(x) dx = \int_{\mathbb{R}^l} Dp(x)^T c_s(u, x) Dw_s(u, x) m_s(x) dx \quad (7.13a)$$

and

$$\int_{\mathbb{R}^l} Dp(x)^T G_s(u, x)^T m_s(x) dx = \int_{\mathbb{R}^l} Dp(x)^T c_s(u, x) Dv_s(u, x)^T m_s(x) dx \quad (7.13b)$$

for all $p \in \mathcal{C}_0^\infty(\mathbb{R}^l)$. Furthermore, $w_s(u, x)$, $Dw_s(u, x)$, $v_s(u, x)$, and $Dv_s(u, x)$ are continuous in (u, x) for almost all $s \in \mathbb{R}_+$, and, for all open balls $S \subset \mathbb{R}^l$, all $L > 0$ and all $t > 0$,

$$\begin{aligned} \sup_{s \in [0, t]} \sup_{u: |u| \leq L} (\|w_s(u, \cdot)\|_{\mathbb{W}^{2,q}(S)} + \|v_s(u, \cdot)\|_{\mathbb{W}^{2,q}(S, \mathbb{R}^n)} + \|Dw_s(u, \cdot)\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)} \\ + \|Dv_s(u, \cdot)\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l \times \mathbb{R}^n, m_s(x) dx)}) < \infty. \end{aligned}$$

Also, there exists α_0 which depends on the functions $a_t(u, x)$ and $c_t(u, x)$ only such that, if $\alpha > \alpha_0$, then for all $L > 0$ and all $t > 0$,

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq L} w_s(u, x) < \infty \quad (7.14a)$$

and

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq L} \left(\frac{|w_s(u, x)| + |Dw_s(u, x)|}{1 + |x|^2} + \frac{|v_s(u, x)| + \|Dv_s(u, x)\|}{1 + |x|} \right) < \infty, \quad (7.14b)$$

and, for all $|x|$ great enough locally uniformly in s ,

$$x^T \left(a_s(u, x) - \frac{1}{2} \operatorname{div} c_s(u, x) - c_s(u, x) D w_s(u, x) \right) = 0 \quad (7.15a)$$

and

$$(G_s(u, x) - D v_s(u, x) c_s(u, x)) x = 0. \quad (7.15b)$$

Proof. Since $a_s(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$ by the fact that $a_s(u, x)$ grows at most linearly in x and $m_s(x)$ decays exponentially, and $\operatorname{div} c_s(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$ for a similar reason, $\Phi_{s, m_s(\cdot), u}$ as defined by (2.17a), is an element of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(u, x), m_s(x) dx)$, being a projection in the Hilbert space $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(X_s, x), m_s(x) dx)$. In addition,

$$\int_{\mathbb{R}^l} \|\Phi_{s, m_s(\cdot), u}(x)\|_{c_s(u, x)}^2 m_s(x) dx \leq \int_{\mathbb{R}^l} \|c_s(u, x)^{-1} \left(a_s(u, x) - \frac{1}{2} \operatorname{div} c_s(u, x) \right)\|_{c_s(u, x)}^2 m_s(x) dx. \quad (7.16)$$

By Conditions 2.1 and 2.2 and $m_s(\cdot)$ decaying exponentially,

$$\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_{\mathbb{R}^l} |\Phi_{s, m_s(\cdot), u}(x)|^2 m_s(x) dx < \infty. \quad (7.17)$$

We prove that $\Phi_{s, m_s(\cdot), u}(\cdot)$ is a gradient. Let $D w_i \rightarrow \Phi_{s, m_s(\cdot), u}$ in $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(u, x), m_s(x) dx)$ as $i \rightarrow \infty$, where $w_i \in \mathbb{C}_0^\infty(\mathbb{R}^l)$. Then for every $f \in \mathbb{C}_0^\infty(\mathbb{R}^l, \mathbb{R}^l)$ such that $\operatorname{div} f(x) = 0$, we have that $\int_{\mathbb{R}^l} D w_i(x)^T f(x) dx = 0$. Since $m_s(\cdot)$ is bounded away from zero locally and $c_s(u, x)$ is positive definite, convergence in $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(u, x), m_s(x) dx)$ implies convergence in $\mathbb{L}_{\text{loc}}^2(\mathbb{R}^l, \mathbb{R}^l)$, so $D w_i \rightarrow \Phi_{s, m_s(\cdot), u}$ in $\mathbb{L}_{\text{loc}}^2(\mathbb{R}^l, \mathbb{R}^l)$. Therefore, $\int_{\mathbb{R}^l} \Phi_{s, m_s(\cdot), u}(x)^T f(x) dx = 0$. It follows that $\Phi_{s, m_s(\cdot), u}(x) = D \tilde{w}_s(u, x)$ in the sense of distributions, where $\tilde{w}_s(u, \cdot) \in \mathbb{W}_{\text{loc}}^{1,2}(\mathbb{R}^l)$, see, e.g., Lemma 2.2.1 on p.73 in Sohr [47]. (One could also invoke the Helmholtz decomposition, see, e.g., Farwig, Kozono, and Sohr [16].) Consequently, $\int_{\mathbb{R}^l} \chi(x)^T \Phi_{s, m_s(\cdot), u}(x) dx = - \int_{\mathbb{R}^l} \operatorname{div} \chi(x) \tilde{w}_s(u, x) dx$, for all $\chi \in \mathbb{C}_0^1(\mathbb{R}^l, \mathbb{R}^l)$.

By (2.17a) and condition 2.1, for $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$,

$$- \int_{\mathbb{R}^l} \operatorname{div} (c_s(u, x) m_s(x) D p(x)^T) \tilde{w}_s(u, x) dx = \int_{\mathbb{R}^l} D p(x)^T \left(a_s(u, x) - \frac{1}{2} \operatorname{div} c_s(u, x) \right) m_s(x) dx.$$

Thus, $\tilde{w}_s(u, \cdot)$ is a weak solution to the equation

$$\operatorname{div} (c_s(u, x) D \tilde{w}_s(u, x) m_s(x)) = \operatorname{div} \left(\left(a_s(u, x) - \frac{1}{2} \operatorname{div} c_s(u, x) \right) m_s(x) \right) \quad (7.18)$$

in that

$$\int_{\mathbb{R}^l} D p(x)^T c_s(u, x) D \tilde{w}_s(u, x) m_s(x) dx = \int_{\mathbb{R}^l} D p(x)^T \left(a_s(u, x) - \frac{1}{2} \operatorname{div} c_s(u, x) \right) m_s(x) dx. \quad (7.19)$$

We note that (7.19) uniquely specifies $D \tilde{w}_s(u, \cdot)$ as an element of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, c_s(u, x), m_s(x) dx)$.

Let S and \tilde{S} represent open balls in \mathbb{R}^l such that $S \subset\subset \tilde{S}$, let $\zeta(x)$ represent a \mathbb{C}_0^∞ -function with support in \tilde{S} such that $\zeta(x) = 1$ for $x \in S$, and let $\varphi(x)$ represent a $\mathbb{C}_0^\infty(\tilde{S})$ function. On letting $p(x) = \varphi(x)\zeta(x)$ in (7.19) and integrating by parts, we obtain that $\zeta(x)\tilde{w}_s(u, x)$ is a weak solution f to the Dirichlet problem

$$\begin{aligned} \operatorname{div}(c_s(u, x)m_s(x)Df(x)) &= \operatorname{div}(c_s(u, x)D\zeta(x)\tilde{w}_s(u, x)m_s(x)) + D\zeta(x)^T c_s(u, x)D\tilde{w}_s(u, x)m_s(x) \\ &+ \operatorname{div}\left(\left(a_s(u, x) - \frac{1}{2}\operatorname{div}c_s(u, x)\right)\zeta(x)m_s(x)\right) - D\zeta(x)^T\left(a_s(u, x) - \frac{1}{2}\operatorname{div}c_s(u, x)\right)m_s(x) \end{aligned} \quad (7.20)$$

on \tilde{S} with a zero boundary condition. By Theorem 8.3 on p.181 and Theorem 8.8 on p.183 in Gilbarg and Trudinger [22], $\zeta(x)\tilde{w}_s(u, x)$ is an element of $\mathbb{W}^{2,2}(S)$ and is a strong solution of (7.20). Therefore, $\tilde{w}_s(u, \cdot) \in \mathbb{W}_{\text{loc}}^{2,2}(\mathbb{R}^l)$ and (7.18) holds a.e. in x .

Differentiation in (7.18) and division by $m_s(x)$ yield

$$\begin{aligned} \operatorname{tr}(c_s(u, x)D^2\tilde{w}_s(u, x)) + (c_s(u, x)\frac{Dm_s(x)}{m_s(x)} + \operatorname{div}c_s(u, x))^T D\tilde{w}_s(u, x) \\ = \operatorname{div}(a_s(u, x) - \frac{1}{2}\operatorname{div}c_s(u, x)) + (a_s(u, x) - \frac{1}{2}\operatorname{div}c_s(u, x))\frac{Dm_s(x)}{m_s(x)}. \end{aligned} \quad (7.21)$$

On writing the lefthand side as $\mathcal{L}_{s,u}(x)\tilde{w}_s(u, x)$ and letting $f_s(u, x)$ represent the righthand side, we have that $\mathcal{L}_{s,u}(x)\tilde{w}_s(u, x) = f_s(u, x)$. Let $Y_{s,u}^y(t)$ represent the diffusion process in t with the infinitesimal generator $\mathcal{L}_{s,u}(\cdot)$ and initial condition y , defined on a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ with expectation denoted by \mathbf{E} . It is a strong Markov process by Conditions 2.1 and 2.2 and the hypotheses of the lemma. One can also choose $Y_{s,u}^y(t)$ to be measurable in all variables. (A possible line of reasoning invokes continuous dependence of solutions of stochastic differential equations on parameters, see, e.g., Gikhman and Skorokhod [21], or Krylov [26], and the Scorza-Dragoni theorem.) If $|x|$ is great enough so that $m_s(x) = M_s e^{-\alpha|x|}$, then

$$\frac{Dm_s(x)}{m_s(x)} = -\alpha \frac{x}{|x|}. \quad (7.22)$$

Hence, on recalling Condition 2.1, in particular that $|\operatorname{div}c_s(u, x)|$ is bounded in x locally uniformly in (s, u) , and (2.14a), we have that there exists α_0 which depends on $a_t(u, x)$ and $c_t(u, x)$ only such that if $\alpha > \alpha_0$, then $\limsup_{|x| \rightarrow \infty} (x/|x|)^T (c_s(u, x)Dm_s(x)/m_s(x) + \operatorname{div}c_s(u, x)) < 0$, so $Y_{s,u}^y(t)$ is an ergodic process, see, e.g., Has'minskii [23], Veretennikov [53], and Malyshkin [30]. Since, by the divergence theorem,

$$\int_{\mathbb{R}^l} \mathcal{L}_{s,u}(x)p(x)m_s(x)dx = \int_{\mathbb{R}^l} \operatorname{div}(c_s(u, x)Dp(x)m_s(x))dx = 0$$

for all $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$, $m_s(x)dx$ is the unique invariant measure. Similarly,

$$\int_{\mathbb{R}^l} f_s(u, x)m_s(x)dx = \int_{\mathbb{R}^l} \operatorname{div}\left(\left(a_s(u, x) - \frac{1}{2}\operatorname{div}c_s(u, x)\right)m_s(x)\right)dx = 0,$$

the latter equality being a consequence of $m_s(x)$ decaying exponentially as $|x| \rightarrow \infty$. By (7.21), (7.22), Lipschitz continuity of $a_s(u, \cdot)$ and of $\operatorname{div}c_s(u, \cdot)$, the boundedness property of $\operatorname{div}c_s(u, \cdot)$, and by (2.4b), we may assume that α_0 is such that if $\alpha > \alpha_0$, then $f_s(u, x) > 0$ for all $|x|$ great enough locally uniformly in (s, u) . Also,

$$\sup_{s \in [0, t]} \sup_{x \in \mathbb{R}^l} \sup_{u \in \mathbb{R}^n: |u| \leq L} \frac{|f_s(u, x)|}{1 + |x|} < \infty.$$

By Theorem 1 in Pardoux and Veretennikov [35], the function

$$\check{w}_s(u, x) = - \int_0^\infty \mathbf{E} f_s(u, Y_{s,u}^x(t)) dt \quad (7.23)$$

is well defined, belongs to $\mathbb{W}_{\text{loc}}^{2,q}(\mathbb{R}^l)$, for all $q > 1$, as a function of x , $D\check{w}_s(u, x)$ is of polynomial growth in x , in particular, $D\check{w}_s(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$, and $\mathcal{L}_{s,u}(x)\check{w}_s(u, x) = f_s(u, x)$. Since $D\check{w}_s(u, x)$ also satisfies (7.19), we have that $D\check{w}_s(u, x) = D\tilde{w}_s(u, x)$. In addition, $\check{w}_s(u, x)$ is measurable in (s, u, x) .

As in Pardoux and Veretennikov [35], by (7.23) and the strong Markov property, for $R > 0$,

$$\check{w}_s(u, x) = \mathbf{E}\check{w}_s(u, Y_{s,u}^x(\tau^R)) - \mathbf{E} \int_0^{\tau^R} f_s(u, Y_{s,u}^x(t)) dt, \quad (7.24)$$

where $\tau^R = \inf\{t \in \mathbb{R}_+ : |Y_{s,u}^x(t)| \leq R\} < \infty$. Since $|Y_{s,u}^x(\tau^R)| = R$ if $|x| > R$, by $f_s(u, x)$ being positive for all $|x|$ great enough, we have that if R is great enough then $\check{w}_s(u, x) \leq \check{w}_s(u, R)$, provided $|x| > R$. One can see that the bounds in the calculation of part (a) of the proof of Theorem 1 in Pardoux and Veretennikov [35] hold uniformly over $u \in [0, L]$ and $s \in [0, t]$, which shows that $\sup_{x: |x| \leq R} \sup_{\substack{s \in [0, t], \\ u \in \mathbb{R}^n: |u| \leq L}} |\check{w}_s(u, x)| < \infty$. Since the righthand side of (7.21) grows at

most linearly in $|x|$ locally uniformly in (s, u) , the arguments of part (b) of the proof of Theorem 2 (with $\beta = 2$ and $\alpha = 0$) and of part (e) of the proof of Theorem 1 in Pardoux and Veretennikov [35], along with (7.24), show that the functions $|\check{w}_s(u, x)|$ and $|D\check{w}_s(u, x)|$ grow at most quadratically in $|x|$ locally uniformly in (s, u) .

We define $w_s(u, x) = \check{w}_s(u, x) - V_1^{-1} \int_{S_1} \check{w}_s(u, y) dy$, where S_1 represents the unit open ball centred at the origin in \mathbb{R}^l and V_1 represents the volume of that ball. Obviously, the bounds on $\check{w}_s(u, x)$ we have found are also valid for $w_s(u, x)$. It also satisfies (7.13a). We prove that, for all $q > 1$,

$$\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \|w_s(u, \cdot)\|_{\mathbb{W}^{2,q}(S)} < \infty. \quad (7.25)$$

Since $Dw_s(u, x) = \Phi_{s, m_s(\cdot), u}$ and (7.17) holds, $\int_{S_1} w_s(u, x) dx = 0$, and $m_s(x)$ is locally bounded away from zero, an application of Poincaré's inequality yields $\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \|w_s(u, \cdot)\|_{\mathbb{L}^2(S_1)} < \infty$. If S_2 is a ball containing S_1 , then, for some $L_{S_1, S_2} > 0$,

$$\|w_s(u, \cdot)\|_{\mathbb{W}^{1,2}(S_2)}^2 \leq L_{S_1, S_2} (\|Dw_s(u, \cdot)\|_{\mathbb{L}^2(S_2)}^2 + \|w_s(u, \cdot)\|_{\mathbb{L}^2(S_1)}^2),$$

see p.299 in Kufner, John, and Fučík [27], also Theorem 7.4 on p.109 in Nečas [33]. Thus, on recalling that $S \subset\subset \tilde{S}$ and letting \check{S} represent an open ball in \mathbb{R}^l such that $\tilde{S} \subset\subset \check{S}$, we have that

$$\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \|w_s(u, \cdot)\|_{\mathbb{W}^{1,2}(\check{S})} < \infty. \quad (7.26)$$

By (7.13a), Theorem 5.5.5'(a) on p.156 in Morrey [32], the discussion on p.12 of Bogachev, Krylov, and Röckner [8], Shaposhnikov [46], and the fact that $\sup_{x \in \check{S}} |a_s(u, x)|$ and $\|c_s(u, \cdot)\|_{\mathbb{W}^{1,\infty}(\check{S}, \mathbb{R}^{l \times l})}$ are bounded locally uniformly in (s, u) , we have that $\|w_s(u, \cdot)\|_{\mathbb{W}^{1,q}(\check{S})} \leq M_{\check{S}, \check{S}, q} (1 + \|w_s(u, \cdot)\|_{\mathbb{L}^1(\check{S})})$ locally uniformly in (s, u) . By (7.26), $\sup_{s \in [0, t]} \sup_{|u| \leq L} \|w_s(u, \cdot)\|_{\mathbb{W}^{1,q}(\check{S})} < \infty$. By (7.13a), via a similar argument to the one used for $\zeta(\cdot)\tilde{w}_s(u, \cdot)$ above, $\zeta(\cdot)w_s(u, \cdot)$ is a strong solution to (7.20).

By Theorem 9.15 on p.241 in Gilbarg and Trudinger [22], $\zeta(\cdot)w_s(u, \cdot) \in \mathbb{W}^{2,q}(\tilde{S})$. By Theorem 9.11 on p.235 in Gilbarg and Trudinger [22], locally uniformly in (s, u) , for some $\tilde{M}_{S, \tilde{S}, q} > 0$,

$$\|w_s(u, \cdot)\|_{\mathbb{W}^{2,q}(S)} \leq \tilde{M}_{S, \tilde{S}, q}(1 + \|w_s(u, \cdot)\|_{\mathbb{L}^q(\tilde{S})})$$

which implies (7.25).

We now address the continuity of $w_s(u, x)$. Let $u_i \rightarrow u$. By (7.25) and Sobolev's imbedding, the sequences $w_s(u_i, \cdot)$ and $Dw_s(u_i, \cdot)$ are equicontinuous in $x \in S$, so they are relatively compact in $\mathbb{C}(S, \mathbb{R}^l)$. A similar property holds for $(a_s(u_i, \cdot) - (1/2)\text{div } c_s(u_i, \cdot))m_s(\cdot)$. Taking a subsequential limit in (7.13a) implies that $Dw_s(u_i, \cdot) \rightarrow Dw_s(u, \cdot)$ in $\mathbb{C}(S, \mathbb{R}^l)$. By Poincaré's inequality for $S = S_1$ and the fact that $\int_{S_1} w_s(u, x) dx = 0$, $w_s(u_i, \cdot) \rightarrow w_s(u, \cdot)$ in $\mathbb{L}^2(S_1)$. The bound

$$\|w_s(u_i, \cdot) - w_s(u, \cdot)\|_{\mathbb{W}^{1,2}(S_2)}^2 \leq L_{S_1, S_2}(\|Dw_s(u_i, \cdot) - Dw_s(u, \cdot)\|_{\mathbb{L}^2(S_2)}^2 + \|w_s(u_i, \cdot) - w_s(u, \cdot)\|_{\mathbb{L}^2(S_1)}^2)$$

shows that $w_s(u_i, \cdot) \rightarrow w_s(u, \cdot)$ in $\mathbb{L}^2(S_2)$. Since S_2 is an arbitrary ball that contains S_1 , $w_s(u_i, \cdot) \rightarrow w_s(u, \cdot)$ in $\mathbb{C}(S, \mathbb{R}^l)$. Hence, $w_s(u, x)$ and $Dw_s(u, x)$ are continuous in (u, x) for almost all s .

We prove (7.15a). Since $a_s(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$ and $Dw_s(u, \cdot) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$, (7.13a) extends to $\mathbb{C}^1(\mathbb{R}^l)$ -functions $p(x)$ such that $\int_{\mathbb{R}^l} (p(x)^2 + |Dp(x)|^2) m_s(x) dx < \infty$. For given $\kappa > 0$, $\delta > 0$, and $x_0 \in \mathbb{R}^l$, we let $p(x) = |x|^2 e^{-\delta[(|x-x_0|^2/\kappa-1)^+]^2}$. By dominated convergence,

$$\lim_{\delta \rightarrow \infty} \int_{\mathbb{R}^l} Dp(x)^T c_s(u, x) Dw_s(u, x) m_s(x) dx = 2 \int_{x \in \mathbb{R}^l: |x-x_0| < \kappa} x^T c_s(u, x) Dw_s(u, x) m_s(x) dx$$

and

$$\lim_{\delta \rightarrow \infty} \int_{\mathbb{R}^l} Dp(x)^T (a_s(u, x) - \frac{1}{2} \text{div } c_s(u, x)) m_s(x) dx = 2 \int_{x \in \mathbb{R}^l: |x-x_0| < \kappa} x^T (a_s(u, x) - \frac{1}{2} \text{div } c_s(u, x)) m_s(x) dx.$$

Dividing the righthand sides by the volume of the ball of radius κ centred at x_0 , letting $\kappa \rightarrow 0$ and accounting for (7.13a) and for $Dw_s(u, x)$, $m_s(x)$, $c_s(u, x)$ and $a_s(u, x)$ being continuous in x , yields (7.15a).

The part that concerns $v_s(u, x)$ is dealt with similarly, except that one uses Theorem 2 of Pardoux and Veretennikov [35] with $\beta = 1$ to bound the growth rate of the second term of the sum in (7.14b). \square

We now take on the proof of Theorem 7.1. Let $\hat{w}_s(u, x)$ and $\hat{v}_s(u, x)$ represent $w_s(u, x)$ and $v_s(u, x)$, respectively, in the statement of Lemma 7.3 for $m_s(x) = \hat{m}_s(x)$. We define, guided by (6.21) and (6.22), on recalling (2.17b) and (2.3),

$$\begin{aligned} \hat{\lambda}_s(u) = & \left(\int_{\mathbb{R}^l} Q_{s, \hat{m}_s(\cdot)}(u, x) \hat{m}_s(x) dx \right)^{-1} \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(u, x) \hat{m}_s(x) dx \right. \\ & \left. - \int_{\mathbb{R}^l} G_s(u, x) \left(\frac{D\hat{m}_s(x)}{2\hat{m}_s(x)} - D\hat{w}_s(u, x) \right) \hat{m}_s(x) dx \right) \end{aligned} \quad (7.27)$$

if $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) and $\hat{\lambda}_s(u) = 0$ if $C_t(u, x) = 0$ for all (t, u, x) , and

$$\hat{h}_s(u, x) = \frac{1}{2} \ln \hat{m}_s(x) - \hat{w}_s(u, x) - \hat{v}_s(u, x)^T \hat{\lambda}_s(u) \quad (7.28)$$

so that

$$D\hat{h}_s(u, x) = \frac{D\hat{m}_s(x)}{2\hat{m}_s(x)} - D\hat{w}_s(u, x) - D\hat{v}_s(u, x)^T \hat{\lambda}_s(u). \quad (7.29)$$

We note that by (2.16),

$$Q_{s, \hat{m}_s(\cdot)}(u, x) = C_s(u, x) - \|D\hat{v}_s(u, x)\|_{C_s(u, x)}^2. \quad (7.30)$$

The continuity properties of $w_s(u, x)$ and $v_s(u, x)$ established in Lemma 7.3 imply that $\hat{\lambda}_s(u)$ is continuous in u and that $\hat{h}_s(u, x)$ and $D\hat{h}_s(u, x)$ are continuous in (u, x) , for almost all $s \in \mathbb{R}_+$.

If $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) , then, the analogue of (7.16) for $D\hat{w}_s(u, x)$, (7.27) and Condition 2.1 imply that, for some $\vartheta_1 > 0$,

$$|\hat{\lambda}_s(u)| \leq \vartheta_1 (|\dot{X}_s| + \sup_{x \in \mathbb{R}^l} |A_s(u, x)| + \left(\int_{\mathbb{R}^l} |a_s(u, x)|^2 \hat{m}_s(x) dx \right)^{1/2} + \sup_{x \in \mathbb{R}^l} \frac{|D\hat{m}_s(x)|}{\hat{m}_s(x)}). \quad (7.31)$$

Since $\hat{m}_s(x) = M_s e^{-\alpha|x|}$ for $|x| > 2r$ and $|a_s(u, x)|$ grows at most linearly in x , we conclude that $|\hat{\lambda}_s(u)|$ is locally bounded in (s, u) .

Therefore, by Lemma 7.3, for all $L > 0$, all open balls S in \mathbb{R}^l , and all $q > 1$,

$$\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} |\hat{\lambda}_s(u)| + \sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_S |D\hat{h}_s(u, x)|^q dx < \infty. \quad (7.32)$$

By Theorem 6.1, the supremum in (6.20) is attained at $\lambda = \hat{\lambda}_s(u)$ and $g = D\hat{h}_s(u, x)$, however, the function $\hat{h}_s(u, x)$ might not be of compact support in x so in order to use it in Lemma 7.2, we need to restrict it to a compact set. Let $\eta(y)$ represent an \mathbb{R}_+ -valued nonincreasing $C_0^\infty(\mathbb{R}_+)$ -function such that $\eta(y) = 1$ for $0 \leq y \leq 1$ and $\eta(y) = 0$ for $y \geq 2$. Let $\hat{w}_s^i(u, x) = \hat{w}_s(u, x)\eta(|x|/i)$ and $\hat{v}_s^i(u, x) = \hat{v}_s(u, x)\eta(|x|/i)$. We note that

$$D\hat{w}_s^i(u, x) = \eta\left(\frac{|x|}{i}\right)D\hat{w}_s(u, x) + \frac{x}{i|x|} D\eta\left(\frac{|x|}{i}\right) \hat{w}_s(u, x) \quad (7.33a)$$

and

$$D\hat{v}_s^i(u, x) = D\hat{v}_s(u, x)\eta\left(\frac{|x|}{i}\right) + \frac{x}{i|x|} \hat{v}_s(u, x) D\eta\left(\frac{|x|}{i}\right)^T. \quad (7.33b)$$

We define, in analogy with (7.27),

$$\begin{aligned} \hat{\lambda}_s^i(u) = & \left(\int_{\mathbb{R}^l} Q_{s, \hat{m}_s(\cdot)}(u, x) \hat{m}_s(x) dx \right)^{-1} \left(\dot{X}_s - \int_{\mathbb{R}^l} A_s(u, x) \hat{m}_s(x) dx \right. \\ & \left. - \int_{\mathbb{R}^l} G_s(u, x) \left(\frac{1}{2} D\left(\eta\left(\frac{|x|}{i}\right) \ln \hat{m}_s(x)\right) - D\hat{w}_s^i(u, x) \right) \hat{m}_s(x) dx \right) \end{aligned} \quad (7.34)$$

if $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) , and $\hat{\lambda}_s^i(u) = 0$ if $C_t(u, x) = 0$. We let, in analogy with (7.28),

$$\hat{h}_s^i(u, x) = \frac{1}{2} \eta\left(\frac{|x|}{i}\right) \ln \hat{m}_s(x) - \hat{w}_s^i(u, x) - \hat{v}_s^i(u, x)^T \hat{\lambda}_s^i(u). \quad (7.35)$$

In analogy with (7.31) and in view of (7.33a) and (7.14b) in Lemma 7.3, one can see that the $|\hat{\lambda}_s^i(u)|$ are bounded uniformly in i and locally uniformly in (s, u) , where the bound may depend on α . Also, $\hat{\lambda}_s^i(u)$ is continuous in u , so it satisfies the hypotheses of Lemma 7.2.

If $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) , then by (7.33a), Lemma 7.3, (7.27), and (7.34),

$$\lim_{i \rightarrow \infty} \sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} |\hat{\lambda}_s^i(u) - \hat{\lambda}_s(u)| = 0. \quad (7.36)$$

The latter convergence also holds if $C_t(u, x) = 0$ in that $\hat{\lambda}_s^i(u) = \hat{\lambda}_s(u) = 0$.

Similarly, since by (7.33a), (7.33b), and (7.35),

$$D\hat{h}_s^i(u, x) = \eta\left(\frac{|x|}{i}\right)D\hat{h}_s(u, x) + \frac{1}{i} \frac{x}{|x|} D\eta\left(\frac{|x|}{i}\right) \left(\frac{1}{2} \ln \hat{m}_s(x) - \hat{w}_s(u, x) - \hat{v}_s(u, x)^T \hat{\lambda}_s^i(u)\right), \quad (7.37)$$

we have that

$$\lim_{i \rightarrow \infty} \sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \int_S |D\hat{h}_s^i(u, x) - D\hat{h}_s(u, x)|^q dx = 0, \quad (7.38)$$

for all $L > 0$, all open balls S in \mathbb{R}^l , and all $q > 1$. The functions $\hat{h}_s^i(u, x)$ also satisfy the hypotheses of Lemma 7.2.

Another auxiliary lemma is in order.

Lemma 7.4. *Suppose, for $i \in \mathbb{N}$, $X^i \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$, $X \in \mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$, and $m_s^i(x)$ and $m_s(x)$ are measurable functions which are probability densities in x on \mathbb{R}^l for almost all s such that*

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr}(c_s(X_s^i, x) D^2 p(x)) + Dp(x)^T (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i)) \right. \\ \left. - \frac{1}{2} \operatorname{div} c_s(X_s^i, x) + c_s(X_s^i, x) D\hat{h}_s^i(X_s^i, x) \right) m_s^i(x) dx = 0$$

and

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr}(c_s(X_s, x) D^2 p(x)) + Dp(x)^T (a_s(X_s, x) + G_s(X_s, x)^T \hat{\lambda}_s(X_s)) \right. \\ \left. - \frac{1}{2} \operatorname{div} c_s(X_s, x) + c_s(X_s, x) D\hat{h}_s(X_s, x) \right) m_s(x) dx = 0, \quad (7.39)$$

for all $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$. If $X^i \rightarrow X$ as $i \rightarrow \infty$, then, for all α great enough and for all $t > 0$,

$$\lim_{i \rightarrow \infty} \int_0^t \int_{\mathbb{R}^l} |m_s^i(x) - m_s(x)| dx ds = 0, \quad (7.40a)$$

$$\lim_{i \rightarrow \infty} \int_0^t \int_{\mathbb{R}^l} \|D\hat{h}_s^i(X_s^i, x)\|_{c_s(X_s^i, x)}^2 m_s^i(x) dx ds = \int_0^t \int_{\mathbb{R}^l} \|D\hat{h}_s(X_s, x)\|_{c_s(X_s, x)}^2 m_s(x) dx ds, \quad (7.40b)$$

and

$$\lim_{i \rightarrow \infty} \int_0^t \|\hat{\lambda}_s^i(X_s^i)\|_{\int_{\mathbb{R}^l} c_s(X_s^i, x) m_s^i(x) dx}^2 ds = \int_0^t \|\hat{\lambda}_s(X_s)\|_{\int_{\mathbb{R}^l} c_s(X_s, x) m_s(x) dx}^2 ds. \quad (7.40c)$$

Proof. Let us first address existence and uniqueness of $m_s^i(x)$ and $m_s(x)$. Since $\sup_{|u| \leq L} (x/|x|)^T a_s(u, x) \rightarrow -\infty$ as $|x| \rightarrow \infty$, the function $G_s(u, x)$ is bounded, the function $\hat{\lambda}_s^i(u)$ is bounded locally in (s, u) and the function $\hat{h}_s^i(u, x)$ is of compact support in x locally uniformly in (s, u) , we have that

$$\lim_{|x| \rightarrow \infty} \frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x) + c_s(X_s^i, x) D \hat{h}_s^i(X_s^i, x)) = -\infty, \quad (7.41)$$

which implies that $m_s^i(x)$ is well defined and is specified uniquely, see, e.g., Metafuno, Pallara, and Rhandi [31, Theorem 2.2, Proposition 2.4].

By (7.29), relations (7.15a) and (7.15b) of Lemma 7.3 imply that

$$\begin{aligned} x^T (a_s(X_s, x) + G_s(X_s, x)^T \hat{\lambda}_s(X_s) - \frac{1}{2} \operatorname{div} c_s(X_s, x) + c_s(X_s, x) D \hat{h}_s(X_s, x)) \\ = \frac{x^T}{2} c_s(X_s, x) \frac{D \hat{m}_s(x)}{\hat{m}_s(x)}. \end{aligned}$$

If $|x| > 2r$, then $D \hat{m}_s(x) / \hat{m}_s(x) = -\alpha x / |x|$, so, locally uniformly in s ,

$$\limsup_{|x| \rightarrow \infty} \frac{x^T}{|x|} (a_s(X_s, x) + G_s(X_s, x)^T \hat{\lambda}_s(X_s) - \frac{1}{2} \operatorname{div} c_s(X_s, x) + c_s(X_s, x) D \hat{h}_s(X_s, x)) < 0,$$

which ensures the existence and uniqueness of $m_s(x)$.

As in the proof of Lemma 6.7, we will show that, for arbitrary $\delta > 0$, there exists $\alpha > 0$ such that for all $t > 0$

$$\sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \int_{\mathbb{R}^t} e^{\delta |x|} m_s^i(x) dx < \infty. \quad (7.42)$$

We begin by establishing a uniform version of (7.41):

$$\begin{aligned} \lim_{\alpha \rightarrow \infty} \limsup_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x) \\ + c_s(X_s^i, x) D \hat{h}_s^i(X_s^i, x)) = -\infty. \end{aligned} \quad (7.43)$$

By (7.37), (7.29), (7.15a), and (7.14a), for $|x| > 2r$, on recalling that $\hat{m}_s(x) = M_s e^{-\alpha |x|}$,

$$\begin{aligned} \frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x) + c_s(X_s^i, x) D \hat{h}_s^i(X_s^i, x)) \\ = \frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) (1 - \eta(\frac{|x|}{i})) \\ - \frac{\alpha}{2} \left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 (\eta(\frac{|x|}{i}) + \frac{|x|}{i} D \eta(\frac{|x|}{i})) \\ - \frac{1}{i} \left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 D \eta(\frac{|x|}{i}) \left(\frac{1}{2} \ln M_s - \hat{w}_s(X_s^i, x) - \hat{v}_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) \right). \end{aligned} \quad (7.44)$$

Let $\kappa \in (0, 1)$ be such that $\eta(y) + y D \eta(y) > 0$ if $\eta(y) \in [1 - \kappa, 1]$. If $\eta(|x|/i) \geq 1 - \kappa$, then $\eta(|x|/i) \in [1 - \kappa, 1]$, so substituting $1 - \kappa$ and 1 for $\eta(|x|/i)$, we have that

$$\begin{aligned} \frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) (1 - \eta(\frac{|x|}{i})) \\ \leq \left(\kappa \frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) \right)^+. \end{aligned} \quad (7.45)$$

In addition, for some $\vartheta > 0$,

$$\left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \left(\eta\left(\frac{|x|}{i}\right) + \frac{|x|}{i} D\eta\left(\frac{|x|}{i}\right) \right) \geq \vartheta \inf_{y: \eta(y) \geq 1-\kappa} (\eta(y) + y D\eta(y)) > 0.$$

The righthand side of (7.45) being equal to 0 if $|x|$ is great enough, uniformly in i and locally uniformly in s , it follows that there exists $\hat{M}_1 > 0$ such that for all $|x|$ great enough, depending on $\alpha > 0$ and $t > 0$, uniformly in $i \in \mathbb{N}$ and $s \in [0, t]$,

$$\begin{aligned} & \left(\frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) (1 - \eta\left(\frac{|x|}{i}\right)) \right. \\ & \quad \left. - \frac{\alpha}{2} \left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \left(\eta\left(\frac{|x|}{i}\right) + \frac{|x|}{i} D\eta\left(\frac{|x|}{i}\right) \right) \right) \mathbf{1}_{\{\eta(|x|/i) \geq 1-\kappa\}}(x) \leq -\hat{M}_1 \alpha \mathbf{1}_{\{\eta(|x|/i) \geq 1-\kappa\}}(x). \end{aligned}$$

If $\eta(|x|/i) < 1 - \kappa$, then, analogously, given arbitrary $\hat{M}_2 > 0$, we have that, for all $|x|$ great enough, depending on α and $t > 0$, all $i \in \mathbb{N}$, and all $s \in [0, t]$,

$$\frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) (1 - \eta\left(\frac{|x|}{i}\right)) \leq -\hat{M}_2.$$

Also,

$$\left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \left(\eta\left(\frac{|x|}{i}\right) + \frac{|x|}{i} D\eta\left(\frac{|x|}{i}\right) \right) \geq \vartheta \inf_{y: \eta(y) \leq 1-\kappa} (\eta(y) + y D\eta(y)).$$

Hence, given arbitrary $\hat{M}_3 > 0$, for all $|x|$ great enough, depending on α and t , uniformly in $i \in \mathbb{N}$ and $s \in [0, t]$,

$$\begin{aligned} & \left(\frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) (1 - \eta\left(\frac{|x|}{i}\right)) \right. \\ & \quad \left. - \frac{\alpha}{2} \left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \left(\eta\left(\frac{|x|}{i}\right) + \frac{|x|}{i} D\eta\left(\frac{|x|}{i}\right) \right) \right) \mathbf{1}_{\{\eta(|x|/i) \leq 1-\kappa\}}(x) \leq -\hat{M}_3 \mathbf{1}_{\{\eta(|x|/i) \leq 1-\kappa\}}(x). \end{aligned}$$

We conclude that

$$\begin{aligned} & \lim_{\alpha \rightarrow \infty} \limsup_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \left(\frac{x^T}{|x|} (a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) \right. \\ & \quad \left. - \frac{1}{2} \operatorname{div} c_s(X_s^i, x)) (1 - \eta\left(\frac{|x|}{i}\right)) - \frac{\alpha}{2} \left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \left(\eta\left(\frac{|x|}{i}\right) + \frac{|x|}{i} D\eta\left(\frac{|x|}{i}\right) \right) \right) = -\infty. \end{aligned} \quad (7.46)$$

We now work with line 3 of (7.44). Since $D\eta(y) \leq 0$, and since, by (7.14a), the $\hat{w}_s(X_s^i, x)$ are bounded from above in x and i locally uniformly in s ,

$$\limsup_{\alpha \rightarrow \infty} \limsup_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \frac{1}{i} \left(-\left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \right) D\eta\left(\frac{|x|}{i}\right) \hat{w}_s(X_s^i, x) < \infty. \quad (7.47)$$

Since $D\eta(y) = 0$ unless $y \in [1, 2]$, we have that $(1/i)|D\eta(|x|/i)\hat{v}_s(X_s^i, x)| \leq 4|D\eta(|x|/i)|\hat{v}_s(X_s^i, x)/(1 + |x|)$, which is bounded in i and x locally uniformly in s for α great enough by (7.14b) of Lemma 7.3. It follows that

$$\limsup_{\alpha \rightarrow \infty} \limsup_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \frac{1}{i} \left(-\left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 \right) D\eta\left(\frac{|x|}{i}\right) \hat{v}_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) < \infty. \quad (7.48)$$

In addition,

$$\lim_{|x| \rightarrow \infty} \sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \frac{1}{i} \left\| \frac{x}{|x|} \right\|_{c_s(X_s^i, x)}^2 D\eta\left(\frac{|x|}{i}\right) \frac{1}{2} \ln M_s = 0. \quad (7.49)$$

Combining (7.44), (7.46), (7.47), (7.48), and (7.49) proves (7.43).

The rest of the proof is carried out similarly to the proof of Lemma 6.7, with (7.43) assuming the role of the condition that $\sup_{i \in \mathbb{N}} a_{s+t}(X_s^i, x)^T x/|x| \rightarrow -\infty$. Firstly, letting $f_s^i(x) = a_s(X_s^i, x) + G_s(X_s^i, x)^T \hat{\lambda}_s^i(X_s^i) - \operatorname{div} c_s(X_s^i, x)/2 + c_s(X_s^i, x) D\hat{h}_s^i(X_s^i, x)$ and $\mathcal{L}_s^i p(x) = (1/2) \operatorname{tr}(c_s(X_s^i, x) D^2 p(x)) + f_s^i(x)^T Dp(x)$ one derives the inequality in (6.34), where F represents a $\mathbb{C}^\infty(\mathbb{R}^l)$ -function such that $F(x) = e^{\delta|x|}$ if $|x| \geq 1$. The inequality in (7.42) follows if one recalls (7.37), that the $\hat{\lambda}_s^i(u)$ are bounded uniformly in i and locally uniformly in (s, u) , and that according to Lemma 7.3,

$$\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \|\hat{w}_s(u, \cdot)\|_{\mathbb{W}^{2, q}(S)} + \sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \|\hat{v}_s(u, \cdot)\|_{\mathbb{W}^{2, q}(S, \mathbb{R}^n)} < \infty$$

for all $q > 1$, all $L > 0$, and all open balls $S \subset \mathbb{R}^l$ so that by Sobolev's imbedding

$$\sup_{s \in [0, t]} \sup_{u \in \mathbb{R}^n: |u| \leq L} \sup_{x \in \mathbb{R}^l: |x| \leq R} (|\hat{w}_s(u, x)| + |D\hat{w}_s(u, x)| + |\hat{v}_s(u, \cdot)| + \|D\hat{v}_s(u, x)\|) < \infty.$$

Since the $\lambda_s^i(u)$ are bounded uniformly in i and locally uniformly in (s, u) and (7.32) and (7.42) hold, by Proposition 2.16 in Bogachev, Krylov, and Röckner [7], for almost all s the functions $m_s^i(\cdot)$ converge in the variation norm along a subsequence to probability density $\tilde{m}_s(\cdot)$. By (7.29), (7.37), the bounds (7.14b), and by (7.42), we have that

$$\sup_{s \in [0, t]} \sup_{i \in \mathbb{N}} \int_{\mathbb{R}^l} \|D\hat{h}_s^i(X_s^i, x)\|_{c_s(X_s^i, x)}^3 m_s^i(x) dx < \infty. \quad (7.50)$$

Since $\sup_{i \in \mathbb{N}} |\hat{\lambda}_s^i| < \infty$, the convergences in (7.36) and (7.38) imply that $\tilde{m}_s(x)$ must satisfy (7.39), so $\tilde{m}_s(x) = m_s(x)$ and $m_s^i(\cdot) \rightarrow m_s(\cdot)$ in the variation norm. The limit in (7.40a) follows by dominated convergence. The convergence in (7.40c) follows from (7.36), (7.32), and (2.14d). For (7.40b), we also take into account (7.50). \square

We finish the proof of Theorem 7.1. Let, given $N \in \mathbb{N}$, $\hat{\gamma}^{N, i}$ and $\hat{\theta}^{N, i}$ be defined by the respective equations (7.1a) and (7.1b) with $\hat{\lambda}_s^i(u)$ and $\hat{h}_s^i(u, x)$ as $\lambda_s(u)$ and $\hat{h}_s(u, x)$, respectively. Since the functions $\hat{\lambda}_s^i(u)$ and $\hat{h}_s^i(u, x)$ satisfy the hypotheses of Lemma 7.2, there exist $\gamma^{N, i} = (X^{N, i}, \mu^{N, i}) \in \Gamma$ such that $\hat{\theta}^{N, i}(\gamma^{N, i}) = \tilde{\mathbf{I}}(\gamma^{N, i})$ and $\gamma^{N, i} \in K_{2N+2}$ for all i . In particular, $X_0^{N, i} = \hat{u}$, $\mu^{N, i}(ds, dx) = m_s^{N, i}(x) dx ds$, where $m_s^{N, i}(\cdot) \in \mathbb{P}(\mathbb{R}^l)$ (see Theorem 6.1), and the set $\{\gamma^{N, i}, i = 1, 2, \dots\}$ is relatively compact. Since $\tilde{\mathbf{I}}(\gamma^{N, i}) \geq \mathbf{I}^{**}(\gamma^{N, i})$, on the one hand, and $\hat{\theta}^{N, i}(\gamma^{N, i}) \leq \mathbf{I}^{**}(\gamma^{N, i})$ by (6.19) and (7.1b), on the other hand, we have that

$$\hat{\theta}^{N, i}(\gamma^{N, i}) = \mathbf{I}^{**}(\gamma^{N, i}) = \tilde{\mathbf{I}}(\gamma^{N, i}). \quad (7.51)$$

Let $\mu^{N, i} \rightarrow \mu^N$ in $\mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ and $X^{N, i} \rightarrow X^N$ in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ along a subsequence of i , which we still denote by i .

By (7.1b) and (7.51), the suprema in (6.19) for $(X, \mu) = (X^{N, i}, \mu^{N, i})$ are attained at $(\hat{\lambda}_s^i(X_s^{N, i}), \hat{h}_s^i(X_s^{N, i}, x))$ when $s \leq \hat{\tau}^{N, i}(\gamma^{N, i})$. In particular, since the supremum over h for $\lambda = \hat{\lambda}_s^i(X_s^{N, i})$ is attained at $h(x) = \hat{h}_s^i(X_s^{N, i}, x)$, we have that

$$\begin{aligned} D\hat{h}_s^i(X_s^{N, i}, x) &= \Pi_{c_s(X_s^{N, i}, \cdot), m_s^{N, i}(\cdot)} \left(\frac{Dm_s^{N, i}(x)}{2m_s^{N, i}(x)} \right. \\ &\quad \left. + c_s(X_s^{N, i}, \cdot)^{-1} \left(\frac{1}{2} \operatorname{div} c_s(X_s^{N, i}, \cdot) - a_s(X_s^{N, i}, \cdot) - G_s(X_s^{N, i}, \cdot)^T \hat{\lambda}_s^i(X_s^{N, i}) \right) \right)(x). \end{aligned} \quad (7.52)$$

Hence, for $p \in \mathbb{C}_0^\infty(\mathbb{R}^l)$,

$$\int_{\mathbb{R}^l} Dp(x)^T (c_s(X_s^{N,i}, x) \frac{Dm_s^{N,i}(x)}{2m_s^{N,i}(x)} + \frac{1}{2} \operatorname{div} c_s(X_s^{N,i}, x) - a_s(X_s^{N,i}, x) \\ - G_s(X_s^{N,i}, x)^T \hat{\lambda}_s^i(X_s^{N,i}) - c_s(X_s^{N,i}, x) D\hat{h}_s^i(X_s^{N,i}, x)) m_s^{N,i}(x) dx = 0.$$

Integration by parts yields

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr} (c_s(X_s^{N,i}, x) D^2 p(x)) + Dp(x)^T (a_s(X_s^{N,i}, x) - \frac{1}{2} \operatorname{div} c_s(X_s^{N,i}, x) \\ + G_s(X_s^{N,i}, x)^T \hat{\lambda}_s^i(X_s^{N,i}) + c_s(X_s^{N,i}, x) D\hat{h}_s^i(X_s^{N,i}, x)) \right) m_s^{N,i}(x) dx = 0. \quad (7.53)$$

Thus, $m_s^{N,i}(x) dx$ is an invariant probability for a diffusion. By (7.13a), (7.13b) (for $\hat{w}_s(u, x)$ and $\hat{v}_s(u, x)$), and (7.29), via a similar manipulation,

$$\int_{\mathbb{R}^l} \left(\frac{1}{2} \operatorname{tr} (c_s(X_s^N, x) D^2 p(x)) + Dp(x)^T (a_s(X_s^N, x) - \frac{1}{2} \operatorname{div} c_s(X_s^N, x) \\ + G_s(X_s^N, x)^T \hat{\lambda}_s(X_s^N) + c_s(X_s^N, x) D\hat{h}_s(X_s^N, x)) \right) \hat{m}_s(x) dx = 0.$$

Let $\tilde{m}_s^{N,i}(x)$ represent a probability density that solves (7.53) for all $s \in \mathbb{R}_+$ rather than for $s \leq \hat{\tau}^{N,i}(\gamma^{N,i})$. The existence of $\tilde{m}_s^{N,i}(x)$ is established as in the proof of Lemma 7.4, more specifically, see (7.41). Lemma 7.4 implies that $\tilde{m}_s^{N,i}(x) \rightarrow \hat{m}_s(x)$ in $\mathbb{L}^1([0, t] \times \mathbb{R}^l)$ as $i \rightarrow \infty$, that

$$\lim_{i \rightarrow \infty} \int_0^t \int_{\mathbb{R}^l} \|D\hat{h}_s^i(X_s^{N,i}, x)\|_{c_s(X_s^{N,i}, x)}^2 \tilde{m}_s^{N,i}(x) dx ds = \int_0^t \int_{\mathbb{R}^l} \|D\hat{h}_s(X_s^N, x)\|_{c_s(X_s^N, x)}^2 \hat{m}_s(x) dx ds \quad (7.54a)$$

and that

$$\lim_{i \rightarrow \infty} \int_0^t \|\hat{\lambda}_s^i(X_s^{N,i})\|_{\int_{\mathbb{R}^l} C_s(X_s^{N,i}, x) \tilde{m}_s^{N,i}(x) dx}^2 ds = \int_0^t \|\hat{\lambda}_s(X_s^N)\|_{\int_{\mathbb{R}^l} C_s(X_s^N, x) \hat{m}_s(x) dx}^2 ds. \quad (7.54b)$$

By Lemma 7.1, $\hat{\tau}^{N,i}(\tilde{\gamma}^{N,i}) \rightarrow \hat{\tau}^N(\gamma^N)$ as $i \rightarrow \infty$, where $\tilde{\gamma}^{N,i} = (X_s^{N,i}, \tilde{\mu}^{N,i})$ and $\tilde{\mu}^{N,i}(dx, ds) = \tilde{m}_s^{N,i}(x) dx ds$. Since $\hat{\tau}^{N,i}(\tilde{\gamma}^{N,i}) = \hat{\tau}^{N,i}(\gamma^{N,i})$, we obtain that $\hat{\tau}^{N,i}(\gamma^{N,i}) \rightarrow \tau^N(\gamma^N)$ and that $\tilde{m}_s^{N,i}(x) \rightarrow \hat{m}_s(x)$ in $\mathbb{L}^1([0, \tau^N(\gamma^N)] \times \mathbb{R}^l)$, so $\mu_s^N(dx) = \hat{m}_s(x) dx$ for almost all $s \leq \tau^N(\gamma^N)$.

We now use the fact that the supremum in (6.19) over λ for $h(x) = \hat{h}_s^i(X_s^{N,i}, x)$ is attained at $\lambda = \hat{\lambda}_s^i(X_s^{N,i})$. If $C_t(u, x) = 0$ and $A_t(u, x)$ is locally Lipschitz continuous in u locally uniformly in t and uniformly in x , then $\hat{\lambda}_s^i(X_s^{N,i}) = 0$, so $\dot{X}_s^{N,i} = \int_{\mathbb{R}^l} A_s(X_s^{N,i}, x) m_s^{N,i}(x) dx$, which, as in the proof of Lemma 6.7, implies since $X_s^{N,i} \rightarrow X_s^N$ in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n)$ and $(m_s^{N,i}(x)) \rightarrow (\hat{m}_s(x))$ in $\mathbb{L}^1([0, \tau^N(\gamma^N)] \times \mathbb{R}^l)$ as $i \rightarrow \infty$ that $\dot{X}_s^N = \int_{\mathbb{R}^l} A_s(X_s^N, x) \hat{m}_s(x) dx$ a.e. for $s \leq \tau^N(\gamma^N)$. By uniqueness, $X_s^N = \hat{X}_s$ for $s \leq \tau^N(\gamma^N)$. As a byproduct, $\dot{X}_s^{N,i} \rightarrow \dot{\hat{X}}_s$ as $i \rightarrow \infty$ a.e. on $[0, \tau^N(\gamma^N)]$.

Suppose that $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite locally uniformly in (t, u) and uniformly in x . Then the maximisation condition is

$$\begin{aligned} \dot{X}_s^{N,i} = & \int_{\mathbb{R}^l} A_s(X_s^{N,i}, x)m_s^{N,i}(x) dx + \int_{\mathbb{R}^l} G_s(X_s^{N,i}, x)D\hat{h}_s^i(X_s^{N,i}, x) m_s^{N,i}(x) dx \\ & + \int_{\mathbb{R}^l} C_s(X_s^{N,i}, x)\hat{\lambda}_s^i(X_s^{N,i})m_s^{N,i}(x) dx. \end{aligned}$$

On integrating both sides from 0 to t and letting $i \rightarrow \infty$, we have by the facts that $\gamma^{N,i} \rightarrow \gamma^N$, that $X^{N,i} \rightarrow X^N$, that $m_s^{N,i}(x) \rightarrow \hat{m}_s(x)$ in $\mathbb{L}^1([0, \tau^N(\gamma^N)] \times \mathbb{R}^l)$, and that $\hat{\lambda}_s^i(u) \rightarrow \hat{\lambda}_s(u)$ locally uniformly in (s, u) as $i \rightarrow \infty$ (see (7.36)), by (7.38), by (7.50), and by (7.29) that, for almost all $s \leq \tau^N(\gamma^N)$,

$$\begin{aligned} \dot{X}_s^N = & \int_{\mathbb{R}^l} A_s(X_s^N, x)\hat{m}_s(x) dx + \int_{\mathbb{R}^l} G_s(X_s^N, x) \left(\frac{D\hat{m}_s(x)}{2\hat{m}_s(x)} - D\hat{w}_s(X_s^N, x) \right) \hat{m}_s(x) dx \\ & + \int_{\mathbb{R}^l} (C_s(X_s^N, x) - G_s(X_s^N, x) D\hat{v}_s(X_s^N, x)) \hat{m}_s(x) dx \hat{\lambda}_s(X_s^N). \quad (7.55) \end{aligned}$$

Since $D\hat{v}_s(u, \cdot) \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^{n \times l}, c_s(x), \hat{m}_s(x) dx)$ and $G_s(u, \cdot)$ is bounded, (7.13b) extends to Dp representing an arbitrary element of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_t(x), \hat{m}_s(x) dx)$, so by (7.30),

$$\int_{\mathbb{R}^l} Q_{s, \hat{m}_s(\cdot)}(u, x) \hat{m}_s(x) dx = \int_{\mathbb{R}^l} (C_s(u, x) - G_s(u, x) D\hat{v}_s(u, x)) \hat{m}_s(x) dx.$$

Substitution of the latter expression in (7.27) and of (7.27) into (7.55) obtains that $\dot{X}_s^N = \dot{\hat{X}}_s$ a.e. on $[0, \tau^N(\gamma^N)]$, so on recalling that $X_0^N = \hat{X}_0 = \hat{u}$ we conclude that $X_s^N = \hat{X}_s$ for $s \leq \tau^N(\gamma^N)$. In addition, $\dot{X}_s^{N,i} \rightarrow \dot{\hat{X}}_s$ as $i \rightarrow \infty$ a.e. on $[0, \tau^N(\gamma^N)]$.

Hence, in either case, $\tau^N(\gamma^N) = \tau^N(\hat{\gamma})$ and $\gamma_s^N = \hat{\gamma}_s$ for $s \leq \tau^N(\hat{\gamma})$ so that $\theta^N(\gamma^N) = \theta^N(\hat{\gamma})$, where $\hat{\gamma} = (\hat{X}, \hat{\mu})$. We show that

$$\theta^N(\gamma^N) = \lim_{i \rightarrow \infty} \hat{\theta}^{N,i}(\gamma^{N,i}). \quad (7.56)$$

By (7.1b) and (7.52),

$$\begin{aligned} \hat{\theta}^{N,i}(\gamma^{N,i}) = & \int_0^{\hat{\tau}^{N,i}(\gamma^{N,i})} \left(\hat{\lambda}_s^i(X_s^{N,i})^T (\dot{X}_s^{N,i} - \int_{\mathbb{R}^l} A_s(X_s^{N,i}, x)m_s^{N,i}(x) dx) \right. \\ & \left. - \frac{1}{2} \|\hat{\lambda}_s^i(X_s^{N,i})\|_{\int_{\mathbb{R}^l} C_s(X_s^{N,i}, x)m_s^{N,i}(x) dx}^2 + \frac{1}{2} \int_{\mathbb{R}^l} \|D\hat{h}_s^i(X_s^{N,i}, x)\|_{C_s(X_s^{N,i}, x)}^2 m_s^{N,i}(x) dx \right) ds. \end{aligned}$$

Similarly,

$$\begin{aligned} \theta^N(\gamma^N) = & \int_0^{\tau^N(\gamma^N)} \left(\hat{\lambda}_s(X_s^N)^T (\dot{X}_s^N - \int_{\mathbb{R}^l} A_s(X_s^N, x)\hat{m}_s(x) dx) \right. \\ & \left. - \frac{1}{2} \|\hat{\lambda}_s(X_s^N)\|_{\int_{\mathbb{R}^l} C_s(X_s^N, x)\hat{m}_s(x) dx}^2 + \frac{1}{2} \int_{\mathbb{R}^l} \|D\hat{h}_s(X_s^N, x)\|_{C_s(X_s^N, x)}^2 \hat{m}_s(x) dx \right) ds. \end{aligned}$$

On recalling convergences (7.54a) and (7.54b) which are locally uniform in t , the fact that $\tilde{m}_s^{N,i}(x) = m_s^{N,i}(x)$ for $s \leq \tau^{N,i}(\gamma^{N,i})$, and the convergences $\hat{\tau}^{N,i}(\gamma^{N,i}) \rightarrow \tau^N(\gamma^N)$, $\gamma^{N,i} \rightarrow \gamma^N$, for (7.56), it remains to check that

$$\begin{aligned} & \lim_{i \rightarrow \infty} \int_0^{\hat{\tau}^{N,i}(\gamma^{N,i})} \hat{\lambda}_s^i(X_s^{N,i})^T (\dot{X}_s^{N,i} - \int_{\mathbb{R}^l} A_s(X_s^{N,i}, x) m_s^{N,i}(x) dx) ds \\ &= \int_0^{\tau^N(\gamma^N)} \hat{\lambda}_s(X_s^N)^T (\dot{X}_s^N - \int_{\mathbb{R}^l} A_s(X_s^N, x) \hat{m}_s(x) dx) ds. \end{aligned} \quad (7.57)$$

The convergences $\hat{\tau}^{N,i}(\gamma^{N,i}) \rightarrow \tau^N(\gamma^N)$, $\gamma^{N,i} \rightarrow \gamma^N$, and $\dot{X}_s^{N,i} \rightarrow \dot{X}_s^N$ for almost all $s < \tau^N(\gamma^N)$, imply that the $\mathbf{1}_{\{[0, \hat{\tau}^{N,i}(\gamma^{N,i})]\}}(s) \hat{\lambda}_s^i(X_s^{N,i})^T (\dot{X}_s^{N,i} - \int_{\mathbb{R}^l} A_s(X_s^{N,i}, x) m_s^{N,i}(x) dx)$ converge to $\mathbf{1}_{\{[0, \tau^N(\gamma^N)]\}}(s) \hat{\lambda}_s(X_s^N)^T (\dot{X}_s^N - \int_{\mathbb{R}^l} A_s(X_s^N, x) \hat{m}_s(x) dx)$ as $i \rightarrow \infty$ for almost all s . Since the $\hat{\lambda}_s^i(u)$ are bounded uniformly in i and locally uniformly in (s, u) , the uniform integrability needed to derive (7.57) follows by the bound $\sup_{\gamma \in K_\delta} \int_0^N |\dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx|^2 ds < \infty$, which is a consequence of (6.23).

By (7.51), (7.56), and part 1 of Theorem 3.4, $\mathbf{I}^{**}(\gamma^N) = \theta^N(\gamma^N) = \tilde{\mathbf{I}}(\gamma^N)$. (Alternatively, one can follow the proof of part 1 of Theorem 3.4 by letting $i \rightarrow \infty$ in (7.51) to obtain that $\mathbf{I}^{**}(\gamma^N) \geq \theta^N(\gamma^N) \geq \tilde{\mathbf{I}}(\gamma^N)$.) Therefore, $\mathbf{I}^{**}(\hat{\gamma}) \geq \theta^N(\hat{\gamma}) = \theta^N(\gamma^N) = \tilde{\mathbf{I}}(\gamma^N)$. Let $\pi_t(\gamma)$, where $\gamma = (X, \mu)$, denote the projection $((X_{s \wedge t}, \mu_{s \wedge t}(\cdot)), s \in \mathbb{R}_+)$. We have that

$$\tilde{\mathbf{I}}(\gamma^N) \geq \inf_{\gamma: \pi_{\tau^N(\gamma^N)}(\gamma) = \pi_{\tau^N(\gamma^N)}(\gamma^N)} \tilde{\mathbf{I}}(\gamma) = \inf_{\gamma: \pi_{\tau^N(\hat{\gamma})}(\gamma) = \pi_{\tau^N(\hat{\gamma})}(\hat{\gamma})} \tilde{\mathbf{I}}(\gamma).$$

The sets $\pi_{\tau^N(\hat{\gamma})}^{-1}(\pi_{\tau^N(\hat{\gamma})}(\hat{\gamma}))$ are closed and decrease to $\hat{\gamma}$ as $N \rightarrow \infty$, so the rightmost side converges to $\tilde{\mathbf{I}}(\hat{\gamma})$, by $\tilde{\mathbf{I}}$ being lower compact. We conclude that $\mathbf{I}^{**}(\hat{\gamma}) \geq \tilde{\mathbf{I}}(\hat{\gamma})$, so $\mathbf{I}^{**}(\hat{\gamma}) = \tilde{\mathbf{I}}(\hat{\gamma})$.

8 Approximating the large deviation function

By Theorem 3.4, in order to complete the proof of Theorem 2.1, it remains to establish an approximation theorem for \mathbf{I}^{**} along the lines of part 2 of Theorem 3.4. We state it next.

Theorem 8.1. *Suppose that conditions 2.1 – 2.3, (2.4b), and (2.14d) hold. If $\mathbf{I}^{**}(X, \mu) < \infty$, then there exists sequence $(X^{(j)}, \mu^{(j)})$ whose members satisfy the requirements on $(\hat{X}, \hat{\mu})$ in the statement of Theorem 7.1 such that $(X^{(j)}, \mu^{(j)}) \rightarrow (X, \mu)$ and $\mathbf{I}^{**}(X^{(j)}, \mu^{(j)}) \rightarrow \mathbf{I}^{**}(X, \mu)$ as $j \rightarrow \infty$.*

Proof. Let $\mu(ds, dx) = m_s(x) dx ds$ and

$$k_s(x) = \frac{1}{2m_s(x)} \operatorname{div} (c_s(X_s, x) m_s(x)) - a_s(X_s, x). \quad (8.1)$$

Since, by Theorem 6.1, $\int_0^t \int_{\mathbb{R}^l} |Dm_s(x)|^2 / m_s(x) dx ds < \infty$, for all $t \in \mathbb{R}_+$, we have that $k_s(\cdot) \in \mathbb{L}_{\text{loc}}^2(\mathbb{R}^l, \mathbb{R}^l, m_s(x) dx)$ a.e.

Let function η be as in Condition 2.3. We introduce $\eta_r(x) = \eta(|x|/r)$ and

$$k_s^r(x) = \frac{1}{2\eta_r^2(x)m_s(x)} \operatorname{div} (c_s(X_s, x) \eta_r^2(x) m_s(x)) - a_s(X_s, x), \quad (8.2)$$

where $x \in \mathbb{R}^l$ and $r > 0$. We also let S_r represent the open ball in \mathbb{R}^l of radius r centred at the origin.

We first prove that one can choose $(X^{(j)}, \mu^{(j)})$ of the required form that converge to (X, μ) as $j \rightarrow \infty$ and are such that $\mathbf{I}_t^{**}(X^{(j)}, \mu^{(j)}) \rightarrow \mathbf{I}_t^{**}(X, \mu)$ for all t , where \mathbf{I}_t^{**} is defined by (6.26).

Let us begin with the case where $C_t(u, x) = 0$ for all (t, u, x) and $A_t(u, x)$ is Lipschitz continuous in u locally uniformly in t and uniformly in x . By Theorem 6.1, $\dot{X}_s = \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx$ a.e., the latter equation having a unique solution. Let $\rho_\kappa(x) = (1/\kappa^l)\rho(x/\kappa)$ for $\kappa > 0$, where $\rho(x)$ is a mollifier on \mathbb{R}^l . We define, for $i, j, j' \in \mathbb{N}$ and $\alpha > 0$,

$$m_s^{i,j,j'}(x) = M_s^{i,j,j'}(\hat{m}_s^{i,j'}(x)\eta_j^2(x) + e^{-\alpha|x|}(1 - \eta_j^2(x))) \quad (8.3a)$$

and

$$M_s^{i,j,j'} = \left(\int_{\mathbb{R}^l} (\hat{m}_s^{i,j'}(x)\eta_j^2(x) + e^{-\alpha|x|}(1 - \eta_j^2(x))) dx \right)^{-1}, \quad (8.3b)$$

where

$$\hat{m}_s^{i,j'}(x) = \int_{\mathbb{R}^l} \rho_{1/i}(\tilde{x}) \hat{m}_s^{j'}(x - \tilde{x}) d\tilde{x}, \quad \hat{m}_s^{j'}(x) = m_s(x) \wedge j' \vee \frac{1}{j'}. \quad (8.4)$$

We note that, thanks to Theorem 6.1, $\hat{m}_s^{j'} \in \mathbb{W}_{\text{loc}}^{1,2}(\mathbb{R}^l)$.

We use Lemma 6.7 to define $X^{i,j,j'}$ as the solution of the equation

$$\dot{X}_s^{i,j,j'} = \int_{\mathbb{R}^l} A_s(X_s^{i,j,j'}, x) m_s^{i,j,j'}(x) dx,$$

with $X_0^{i,j,j'} = X_0$. The densities $m_s^{i,j,j'}(x)$ are of class \mathbb{C}^1 in x , with bounded derivatives, and are locally bounded away from zero, and the $X^{i,j,j'}$ are locally Lipschitz continuous by Lemma 6.7.

We introduce further

$$M_s^{j,j'} = \left(\int_{\mathbb{R}^l} (\hat{m}_s^{j'}(x)\eta_j^2(x) + e^{-\alpha|x|}(1 - \eta_j^2(x))) dx \right)^{-1}, \quad (8.5a)$$

$$m_s^{j,j'}(x) = M_s^{j,j'}(\hat{m}_s^{j'}(x)\eta_j^2(x) + e^{-\alpha|x|}(1 - \eta_j^2(x))), \quad (8.5b)$$

and

$$\dot{X}_s^{j,j'} = \int_{\mathbb{R}^l} A_s(X_s^{j,j'}, x) m_s^{j,j'}(x) dx, \quad X_0^{j,j'} = X_0. \quad (8.5c)$$

Let also

$$M_s^j = \left(\int_{\mathbb{R}^l} (m_s(x)\eta_j^2(x) + e^{-\alpha|x|}(1 - \eta_j^2(x))) dx \right)^{-1}, \quad (8.6a)$$

$$m_s^j(x) = M_s^j(m_s(x)\eta_j^2(x) + e^{-\alpha|x|}(1 - \eta_j^2(x))), \quad (8.6b)$$

and

$$\dot{X}_s^j = \int_{\mathbb{R}^l} A_s(X_s^j, x) m_s^j(x) dx, \quad X_0^j = X_0.$$

We have that

$$\lim_{i \rightarrow \infty} M_s^{i,j,j'} = M_s^{j,j'}, \quad \lim_{i \rightarrow \infty} \int_{\mathbb{R}^l} |m_s^{i,j,j'}(x) - m_s^{j,j'}(x)| dx = 0, \quad \lim_{i \rightarrow \infty} X_s^{i,j,j'} = X_s^{j,j'}, \quad (8.7a)$$

$$\lim_{j' \rightarrow \infty} M_s^{j,j'} = M_s^j, \quad \lim_{j' \rightarrow \infty} \int_{\mathbb{R}^l} |m_s^{j,j'}(x) - m_s^j(x)| dx = 0, \quad \lim_{j' \rightarrow \infty} X_s^{j,j'} = X_s^j, \quad (8.7b)$$

and

$$\lim_{j \rightarrow \infty} M_s^j = 1, \quad \lim_{j \rightarrow \infty} \int_{\mathbb{R}^l} |m_s^j(x) - m_s(x)| dx = 0, \quad \lim_{j \rightarrow \infty} X_s^j = X_s. \quad (8.7c)$$

The third convergence on each line is proved by a similar compactness argument to the one used in the proof of Lemma 6.7.

By (6.27),

$$\begin{aligned} \mathbf{I}_t^{**}(X, \mu) &= \int_0^t \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \left(\int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s, x) m_s(x)) \right. \right. \\ &\quad \left. \left. - a_s(X_s, x) m_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx \right) ds \end{aligned} \quad (8.8)$$

and

$$\begin{aligned} \mathbf{I}_t^{**}(X^{i,j,j'}, \mu^{i,j,j'}) &= \int_0^t \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \left(\int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}(c_s(X_s^{i,j,j'}, x) m_s^{i,j,j'}(x)) \right. \right. \\ &\quad \left. \left. - a_s(X_s^{i,j,j'}, x) m_s^{i,j,j'}(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s^{i,j,j'}, x)}^2 m_s^{i,j,j'}(x) dx \right) ds. \end{aligned}$$

By (8.3a),

$$\mathbf{I}_t^{**}(X^{i,j,j'}, \mu^{i,j,j'}) \leq M_s^{i,j,j'} \left(\int_0^t I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j,j'}, s) ds + \int_0^t I_2^j(X_s^{i,j,j'}, s) ds \right), \quad (8.9)$$

where, for generic \tilde{X}_s and \tilde{m}_s ,

$$\begin{aligned} I_1^j(\tilde{X}_s, \tilde{m}_s, s) &= \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}(c_s(\tilde{X}_s, x) \eta_j^2(x) \tilde{m}_s(x)) \right. \\ &\quad \left. - a_s(\tilde{X}_s, x) \eta_j^2(x) \tilde{m}_s(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(\tilde{X}_s, x)}^2 \eta_j^2(x) \tilde{m}_s(x) dx \end{aligned} \quad (8.10a)$$

and

$$I_2^j(\tilde{X}_s, s) = \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x)^T \left(\frac{1}{2} \operatorname{div}(c_s(\tilde{X}_s, x) e^{-\alpha|x|} (1 - \eta_j^2(x)) \right. \\ \left. - a_s(\tilde{X}_s, x) e^{-\alpha|x|} (1 - \eta_j^2(x)) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(\tilde{X}_s, x)}^2 e^{-\alpha|x|} (1 - \eta_j^2(x))) dx. \quad (8.10b)$$

We prove that

$$\lim_{i \rightarrow \infty} I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) = I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s). \quad (8.11)$$

Let, in analogy with (8.2),

$$k_s^{i,j,j'}(x) = \frac{1}{2\eta_j^2(x)\hat{m}_s^{i,j'}(x)} \operatorname{div}(c_s(X_s^{i,j,j'}, x)\eta_j^2(x)\hat{m}_s^{i,j'}(x)) - a_s(X_s^{i,j,j'}, x). \quad (8.12)$$

This function is an element of $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, \eta_j^2(x)\hat{m}_s^{i,j'}(x) dx)$.

The supremum in $I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s)$ is attained at a unique element $g_s^{i,j,j'}$ of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, \eta_j^2(x)\hat{m}_s^{i,j'}(x) dx)$ such that

$$\int_{\mathbb{R}^l} Dp(x)^T k_s^{i,j,j'}(x)\eta_j^2(x)\hat{m}_s^{i,j'}(x) dx = \int_{\mathbb{R}^l} Dp(x)^T c_s(X_s^{i,j,j'}, x)g_s^{i,j,j'}(x)\eta_j^2(x)\hat{m}_s^{i,j'}(x) dx \quad (8.13)$$

for all $p \in \mathbb{C}_0^1(\mathbb{R}^l)$ and

$$I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) = \int_{\mathbb{R}^l} \frac{1}{2} \|g_s^{i,j,j'}(x)\|_{c_s(X_s^{i,j,j'}, x)}^2 \hat{m}_s^{i,j'}(x)\eta_j^2(x) dx. \quad (8.14)$$

Similarly, the supremum in $I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s)$ is attained at a unique element $g_s^{j,j'}$ of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, \eta_j^2(x)\hat{m}_s^{j'}(x) dx)$ such that

$$\int_{\mathbb{R}^l} Dp(x)^T k_s^{j,j'}(x)\hat{m}_s^{j'}(x)\eta_j^2(x) dx = \int_{\mathbb{R}^l} Dp(x)^T c_s(X_s^{j,j'}, x)g_s^{j,j'}(x)\hat{m}_s^{j'}(x)\eta_j^2(x) dx \quad (8.15)$$

and

$$I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s) = \int_{\mathbb{R}^l} \frac{1}{2} \|g_s^{j,j'}(x)\|_{c_s(X_s^{j,j'}, x)}^2 \hat{m}_s^{j'}(x)\eta_j^2(x) dx, \quad (8.16)$$

where

$$k_s^{j,j'}(x) = \frac{1}{2\eta_j^2(x)\hat{m}_s^{j'}(x)} \operatorname{div}(c_s(X_s^{j,j'}, x)\eta_j^2(x)\hat{m}_s^{j'}(x)) - a_s(X_s^{j,j'}, x). \quad (8.17)$$

Let

$$Q_1 = \int_{\mathbb{R}^l} \|Dp(x)\|_{c_s(X_s^{i,j,j'}, x)}^2 \hat{m}_s^{i,j'}(x)\eta_j^2(x) dx, \\ Q_2 = \int_{\mathbb{R}^l} \|k_s^{i,j,j'}(x)\hat{m}_s^{i,j'}(x) - k_s^{j,j'}(x)\hat{m}_s^{j'}(x)\|_{c_s(X_s^{i,j,j'}, x)^{-1} \frac{\eta_j^2(x)}{\hat{m}_s^{i,j'}(x)}}^2 dx,$$

and

$$Q_3 = \int_{\mathbb{R}^l} \|c_s(X_s^{j,j'}, x) g_s^{j,j'}(x)\|_{c_s(X_s^{i,j,j'}, x)^{-1}}^2 \frac{\hat{m}_s^{j'}(x)^2 \eta_j^2(x)}{\hat{m}_s^{i,j'}(x)} dx.$$

By (8.13) and (8.15), we have that

$$\begin{aligned} & \int_{\mathbb{R}^l} Dp(x)^T c_s(X_s^{i,j,j'}, x) g_s^{i,j,j'}(x) \hat{m}_s^{i,j'}(x) \eta_j^2(x) dx \\ &= \int_{\mathbb{R}^l} Dp(x)^T (k_s^{i,j,j'}(x) \hat{m}_s^{i,j'}(x) - k_s^{j,j'}(x) \hat{m}_s^{j'}(x)) \eta_j^2(x) dx \\ &+ \int_{\mathbb{R}^l} Dp(x)^T c_s(X_s^{j,j'}, x) g_s^{j,j'}(x) \hat{m}_s^{j'}(x) \eta_j^2(x) dx \leq \sqrt{Q_1} \sqrt{Q_2} + \sqrt{Q_1} \sqrt{Q_3}. \end{aligned}$$

Hence,

$$\begin{aligned} & \sqrt{\int_{\mathbb{R}^l} \|g_s^{i,j,j'}(x)\|_{c_s(X_s^{i,j,j'}, x)}^2 \hat{m}_s^{i,j'}(x) \eta_j^2(x) dx} \\ &= \sup_{p \in \mathbb{C}_0^1(\mathbb{R}^l): Q_1 \leq 1} \int_{\mathbb{R}^l} Dp(x) c_s(X_s^{i,j,j'}, x) g_s^{i,j,j'}(x) \hat{m}_s^{i,j'}(x) \eta_j^2(x) dx \leq \sqrt{Q_2} + \sqrt{Q_3}. \end{aligned}$$

By (8.14), for arbitrary $\kappa > 0$,

$$I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) \leq \frac{1}{2} \left(1 + \frac{1}{\kappa}\right) Q_2 + \frac{1}{2} (1 + \kappa) Q_3.$$

By (8.4), $\|\hat{m}_s^{i,j'} - \hat{m}_s^{j'}\|_{\mathbb{W}^{1,2}(S_{2j})} \rightarrow 0$ as $i \rightarrow \infty$, see, e.g., Lemma 3.16 on p.66 in Adams and Fournier [1], so, on recalling (8.12), (8.17), and Condition 2.2, we have that $Q_2 \rightarrow 0$ as $i \rightarrow \infty$. The integrand in Q_3 tends to $\|g_s^{j,j'}(x)\|_{c_s(X_s^{j,j'}, x)}^2 \hat{m}_s^{j'}(x) \eta_j^2(x)$ in Lebesgue measure, see (8.7a). Since the function $\hat{m}_s^{j'}(x)/\hat{m}_s^{i,j'}(x)$ is bounded in x and i , by dominated convergence, Q_3 converges to $\int_{\mathbb{R}^l} \|g_s^{j,j'}(x)\|_{c_s(X_s^{j,j'}, x)}^2 \hat{m}_s^{j'}(x) \eta_j^2(x) dx$ as $i \rightarrow \infty$ so that, on recalling (8.16),

$$\limsup_{i \rightarrow \infty} I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) \leq I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s).$$

On the other hand, by (8.10a) and integration by parts,

$$\begin{aligned} I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) &= \sup_{h \in \mathbb{C}_0^2(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(-\frac{1}{2} \operatorname{tr}(c_s(X_s^{i,j,j'}, x) D^2 h(x)) - Dh(x)^T a_s(X_s^{i,j,j'}, x) \right. \\ &\quad \left. - \frac{1}{2} \|Dh(x)\|_{c_s(X_s^{i,j,j'}, x)}^2 \right) \hat{m}_s^{i,j'}(x) \eta_j^2(x) dx \end{aligned}$$

and a similar representation holds for $I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s)$, which facts imply, in view of (8.7a) and the continuity properties in Condition 2.1, that

$$\liminf_{i \rightarrow \infty} I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) \geq I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s). \quad (8.18)$$

We have proved (8.11). We now show that integrals with respect to s converge too. Let us note that, by (8.13) and (8.14),

$$I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) \leq \int_{\mathbb{R}^l} \frac{1}{2} \|k_s^{i,j,j'}(x)\|_{c_s(X_s^{i,j,j'}, x)^{-1}}^2 \hat{m}_s^{i,j'}(x) \eta_j^2(x) dx,$$

so, by (8.12), and conditions 2.1 and 2.2 there exists $M > 0$ such that

$$I_1^{i,j,j'}(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) \leq M \left(1 + \int_{\mathbb{R}^l} \frac{|D\hat{m}_s^{i,j'}(x)|^2}{\hat{m}_s^{i,j'}(x)} \eta_j^2(x) dx\right).$$

Accounting for (8.4) we have that

$$\begin{aligned} \frac{1}{2} \frac{|D\hat{m}_s^{i,j'}(x)|^2}{\hat{m}_s^{i,j'}(x)} &= \sup_{y \in \mathbb{R}^l} (y^T D\hat{m}_s^{i,j'}(x) - \frac{1}{2} |y|^2 \hat{m}_s^{i,j'}(x)) \\ &\leq \int_{\mathbb{R}^l} \rho_{1/i}(\tilde{x}) \sup_{y \in \mathbb{R}^l} (y^T D\hat{m}_s^{j'}(x - \tilde{x}) - \frac{1}{2} |y|^2 \hat{m}_s^{j'}(x - \tilde{x})) d\tilde{x} = \frac{1}{2} \int_{\mathbb{R}^l} \rho_{1/i}(\tilde{x}) \frac{|D\hat{m}_s^{j'}(x - \tilde{x})|^2}{\hat{m}_s^{j'}(x - \tilde{x})} d\tilde{x}. \end{aligned}$$

Therefore, recalling the definition of $\hat{m}_s^{j'}(x)$ in (8.4),

$$\int_{\mathbb{R}^l} \frac{|D\hat{m}_s^{i,j'}(x)|^2}{\hat{m}_s^{i,j'}(x)} \eta_j^2(x) dx \leq \int_{\mathbb{R}^l} \frac{|D\hat{m}_s^{j'}(x)|^2}{\hat{m}_s^{j'}(x)} dx \leq \int_{\mathbb{R}^l} \frac{|Dm_s(x)|^2}{m_s(x)} dx.$$

Since $\int_0^t \int_{\mathbb{R}^l} |Dm_s(x)|^2 / m_s(x) dx ds < \infty$ by Theorem 6.1, (8.11) and Fatou's lemma yield the convergence

$$\lim_{i \rightarrow \infty} \int_0^t I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j'}, s) ds = \int_0^t I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s) ds. \quad (8.19)$$

Let us show that

$$\lim_{j' \rightarrow \infty} \int_0^t I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s) ds = \int_0^t I_1^j(X_s^j, m_s, s) ds. \quad (8.20)$$

We have that

$$\begin{aligned} &|I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s) - I_1^j(X_s^{j,j'}, m_s, s)| \\ &\leq \frac{1}{2} \int_{\mathbb{R}^l} \left(\left\| \frac{1}{2\eta_j^2(x)m_s(x)} \operatorname{div}(c_s(X_s^{j,j'}, x)\eta_j^2(x)m_s(x)) - a_s(X_s^{j,j'}, x) \right\|_{c_s(X_s^{j,j'}, x)^{-1}}^2 \right. \\ &+ \left. \left\| \frac{1}{2\eta_j^2(x)} \operatorname{div}(c_s(X_s^{j,j'}, x)\eta_j^2(x)) - a_s(X_s^{j,j'}, x) \right\|_{c_s(X_s^{j,j'}, x)^{-1}}^2 \eta_j^2(x)m_s(x)(1 - \mathbf{1}_{[1/j', j']}(m_s(x))) \right) dx, \end{aligned}$$

so, by dominated convergence,

$$\lim_{j' \rightarrow \infty} \int_0^t |I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s) - I_1^j(X_s^{j,j'}, m_s, s)| ds = 0. \quad (8.21)$$

Let $\vartheta > 0$ be such that $\|y\|_{c_s(u,x)}^2 \geq \vartheta|y|^2$, for all $s \in [0, t]$, for all u from a large enough ball, all x , and all y . By the convergence of $X^{j,j'}$ to X^j as $j' \rightarrow \infty$, the continuity of $c_s(u, x)$ in u locally uniformly in s and uniformly in x , and by $c_s(u, x)$ being positive definite uniformly in x and locally uniformly in (s, u) , given arbitrary $\delta \in (0, 1)$ and $\kappa \in (0, 1)$, for all j' great enough, locally uniformly in s ,

$$\begin{aligned}
I_1^j(X_s^{j,j'}, m_s, s) &\leq \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T k_s^j(x) - \frac{1}{2} (1 - \delta)(1 - \kappa) \|Dh(x)\|_{c_s(X_s^j, x)}^2 m_s(x) \eta_j^2(x) dx \\
&\quad + \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \left(\int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div}((c_s(X_s^{j,j'}, x) - c_s(X_s^j, x)) m_s(x) \eta_j^2(x)) \right. \right. \\
&\quad \left. \left. - (a_s(X_s^{j,j'}, x) - a_s(X_s^j, x)) m_s(x) \eta_j^2(x) - \frac{1}{2} \delta(1 - \kappa) \|Dh(x)\|_{c_s(X_s^j, x)}^2 m_s(x) \eta_j^2(x) \right) dx \right) \\
&\leq (1 - \delta)^{-1} (1 - \kappa)^{-1} \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T k_s^j(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s^j, x)}^2 m_s(x) \eta_j^2(x) dx \\
&\quad + \delta^{-1} (1 - \kappa)^{-1} \frac{\vartheta^{-1}}{2} \int_{\mathbb{R}^l} \frac{1}{2} \frac{\operatorname{div}((c_s(X_s^{j,j'}, x) - c_s(X_s^j, x)) m_s(x) \eta_j^2(x))}{m_s(x) \eta_j^2(x)} \\
&\quad \quad \quad - (a_s(X_s^{j,j'}, x) - a_s(X_s^j, x))^2 m_s(x) \eta_j^2(x) dx. \quad (8.22)
\end{aligned}$$

By the convergence of $X_s^{j,j'}$ to X_s^j as $j' \rightarrow \infty$, Condition 2.1 and the convergence of $\int_0^t \int_{\mathbb{R}^l} |Dm_s(x)|^2 / m_s(x) dx ds$, the integral from 0 to t of the second integral on the rightmost side of (8.22) tends to zero as $j' \rightarrow \infty$. Therefore, by (8.21), (8.2), (8.8), and (8.10a),

$$\limsup_{j' \rightarrow \infty} \int_0^t I_1^j(X_s^{j,j'}, \hat{m}_s^{j'}, s) ds \leq \int_0^t I_1^j(X_s^j, m_s, s) ds$$

and by an analogue of (8.18), we obtain (8.20).

We now take a limit as $j \rightarrow \infty$. By a similar reasoning to the one used in (8.22), given arbitrary $\delta \in (0, 1)$ and $\kappa \in (0, 1)$, for all j great enough, locally uniformly in s ,

$$\begin{aligned}
I_1^j(X_s^j, m_s, s) &\leq (1 - \delta)^{-2} (1 - \kappa)^{-1} \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T k_s(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \eta_j^2(x) dx \\
&\quad + \delta^{-1} (1 - \delta)^{-1} (1 - \kappa)^{-1} \frac{1}{2} \int_{\mathbb{R}^l} \|D\eta_j(x)\|_{c_s(X_s, x)}^2 m_s(x) dx \\
&\quad + \delta^{-1} (1 - \kappa)^{-1} \frac{\vartheta^{-1}}{2} \int_{\mathbb{R}^l} \frac{1}{2} \frac{\operatorname{div}((c_s(X_s^j, x) - c_s(X_s, x)) m_s(x) \eta_j^2(x))}{m_s(x) \eta_j^2(x)} \\
&\quad \quad \quad - (a_s(X_s^j, x) - a_s(X_s, x))^2 m_s(x) \eta_j^2(x) dx \quad (8.23)
\end{aligned}$$

so that, by Condition 2.3 (with $\lambda = 0$) and Condition 2.1, we have, on recalling (8.8), that

$$\lim_{j \rightarrow \infty} \int_0^t I_1^j(X_s^j, m_s, s) ds = \mathbf{I}_t^{**}(X, \mu). \quad (8.24)$$

Putting together (8.19), (8.20), and (8.24) yields the convergence

$$\lim_{j \rightarrow \infty} \lim_{j' \rightarrow \infty} \lim_{i \rightarrow \infty} \int_0^t I_1^j(X_s^{i,j,j'}, \hat{m}_s^{i,j,j'}, s) ds = \mathbf{I}_t^{**}(X, \mu). \quad (8.25)$$

We now show that the term I_2^j is inconsequential. On recalling that $|a_s(u, x)|$ grows at most linearly in $|x|$ and $|\operatorname{div} c_s(u, x)|$ and $\|c_s(u, x)\|$ are bounded in x locally uniformly in (s, u) , we have that, for some $L > 0$, all (i, j) , and all $s \leq t$,

$$\begin{aligned} I_2^j(X_s^{i,j,j'}, s) &\leq \int_{x \in \mathbb{R}^l: |x| \geq j} \sup_{y \in \mathbb{R}^l} \left(y^T \left(\frac{1}{2} \frac{\operatorname{div}(c_s(X_s^{i,j,j'}, x)(1 - \eta_j^2(x)))}{1 - \eta_j^2(x)} \right. \right. \\ &\quad \left. \left. - \alpha c_s(X_s^{i,j,j'}, x) \frac{x}{2|x|} - a_s(X_s^{i,j,j'}, x) \right) - \frac{1}{2} \|y\|_{c_s(X_s^{i,j,j'}, x)}^2 \right) (1 - \eta_j^2(x)) e^{-\alpha|x|} dx \\ &\leq \int_{x \in \mathbb{R}^l: |x| \geq j} L(1 + \alpha^2 + |x|^2 + \frac{1}{j^2} \frac{|D\eta(|x|/j)|^2}{1 - \eta^2(|x|/j)}) e^{-\alpha|x|} dx. \end{aligned}$$

Since $\eta(y) = 0$ for $y \geq 2$ and (2.7) holds, the latter integral tends to 0 as $j \rightarrow \infty$, so,

$$\lim_{j \rightarrow \infty} \limsup_{j' \rightarrow \infty} \limsup_{i \rightarrow \infty} \int_0^t I_2^j(X_s^{i,j,j'}, s) ds = 0. \quad (8.26)$$

By (8.7a), (8.7b), (8.7c), (8.9), (8.25), and (8.26),

$$\limsup_{j \rightarrow \infty} \limsup_{j' \rightarrow \infty} \limsup_{i \rightarrow \infty} \mathbf{I}_t^{**}(X^{i,j,j'}, \mu^{i,j,j'}) \leq \mathbf{I}_t^{**}(X, \mu).$$

Thus, there exist sequences $j'(j) \rightarrow \infty$ and $i(j) \rightarrow \infty$ as $j \rightarrow \infty$ such that $(X^{i(j),j,j'(j)}, \mu^{i(j),j,j'(j)}) \rightarrow (X, \mu)$ and

$$\limsup_{j \rightarrow \infty} \mathbf{I}_t^{**}(X^{i(j),j,j'(j)}, \mu^{i(j),j,j'(j)}) \leq \mathbf{I}_t^{**}(X, \mu).$$

The reverse inequality follows from the lower semicontinuity of $\mathbf{I}_t^{**}(X, \mu)$ (see (6.27), where we let $\mathbf{I}_t^{**}(X, \mu) = \infty$ if $\mathbf{I}^{**}(X, \mu) = \infty$), so

$$\lim_{j \rightarrow \infty} \mathbf{I}_t^{**}(X^{i(j),j,j'(j)}, \mu^{i(j),j,j'(j)}) = \mathbf{I}_t^{**}(X, \mu), \quad (8.27)$$

and one can take $(X^{(j)}, \mu^{(j)}) = (X^{i(j),j,j'(j)}, \mu^{i(j),j,j'(j)})$.

Suppose now that $C_t(u, x) - G_t(u, x)c_t(u, x)^{-1}G_t(u, x)^T$ is positive definite uniformly in x and locally uniformly in (t, u) . We proceed similarly to the case where $C_t(u, x) = 0$ and define $m_s^{i,j,j'}(x)$, $M_s^{i,j,j'}$, M_s^j , and $m_s^j(x)$ by the respective relations (8.3a), (8.3b), (8.4), (8.5a)–(8.5c), (8.6a), and (8.6b). We let

$$\dot{X}_s^{i,j,j'} = \dot{X}_s^{j,j'} = \dot{X}_s^j = \dot{X}_s \mathbf{1}_{\{|\dot{X}_s| \leq j\}}(s), \quad X_0^{i,j,j'} = X_0^{j,j'} = X_0^j = X_0. \quad (8.28)$$

The convergences in (8.7a), (8.7b), and (8.7c) still hold.

Replacing $a_s(X_s, x)$ with $a_s(X_s, x) + G_s(X_s, x)^T \lambda$ in the proof above, we conclude that in analogy with (8.27), there exist sequences $i(j) \rightarrow \infty$ and $j'(j) \rightarrow \infty$ as $j \rightarrow \infty$ such that, for all $\lambda \in \mathbb{R}^n$ with rational components,

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \int_0^t \sup_{h \in C_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div} (c_s(X_s^j, x) m_s^{i(j), j, j'(j)}(x)) \right. \right. \\
& \quad \left. \left. - (a_s(X_s^j, x) + G_s(X_s^j, x)^T \lambda) m_s^{i(j), j, j'(j)}(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s^j, x)}^2 m_s^{i(j), j, j'(j)}(x) \right) dx ds \\
& = \int_0^t \sup_{h \in C_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div} (c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) \right. \\
& \quad \left. - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \right) dx ds,
\end{aligned}$$

which, in particular, implies that

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \sup_{h \in C_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div} (c_s(X_s^j, x) m_s^{i(j), j, j'(j)}(x)) \right. \right. \\
& \quad \left. \left. - (a_s(X_s^j, x) + G_s(X_s^j, x)^T \lambda) m_s^{i(j), j, j'(j)}(x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s^j, x)}^2 m_s^{i(j), j, j'(j)}(x) \right) dx \\
& = \sup_{h \in C_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} \left(Dh(x)^T \left(\frac{1}{2} \operatorname{div} (c_s(X_s, x) m_s(x)) - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) m_s(x) \right) \right. \\
& \quad \left. - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) \right) dx,
\end{aligned}$$

for almost all $s \in [0, t]$.

We write these convergences, relabelling $m^{i(j), j, j'(j)}$ as $m^{(j)}$ and X_s^j as $X_s^{(j)}$, as

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \int_0^t \int_{\mathbb{R}^l} \left\| \frac{Dm_s^{(j)}(x)}{2m_s^{(j)}(x)} - \Phi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) - \Psi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \lambda \right\|_{c_s(X_s^{(j)}, x)}^2 m_s^{(j)}(x) dx ds \\
& = \int_0^t \int_{\mathbb{R}^l} \left\| \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) - \Psi_{s, m_s(\cdot), X_s}(x) \lambda \right\|_{c_s(X_s, x)}^2 m_s(x) dx ds \quad (8.29a)
\end{aligned}$$

and

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \int_{\mathbb{R}^l} \left\| \frac{Dm_s^{(j)}(x)}{2m_s^{(j)}(x)} - \Phi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) - \Psi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \lambda \right\|_{c_s(X_s^{(j)}, x)}^2 m_s^{(j)}(x) dx \\
& = \int_{\mathbb{R}^l} \left\| \frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) - \Psi_{s, m_s(\cdot), X_s}(x) \lambda \right\|_{c_s(X_s, x)}^2 m_s(x) dx, \quad (8.29b)
\end{aligned}$$

respectively.

By (6.26) and (8.29a) with $\lambda = 0$, it remains to prove that

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \int_0^t \left\| \dot{X}_s^{(j)} - \int_{\mathbb{R}^l} A_s(X_s^{(j)}, x) m_s^{(j)}(x) dx - \int_{\mathbb{R}^l} G_s(X_s^{(j)}, x) \left(\frac{Dm_s^{(j)}(x)}{2m_s^{(j)}(x)} \right. \right. \\
& \quad \left. \left. - \Phi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \right) m_s^{(j)}(x) dx \right\|_{\left(\int_{\mathbb{R}^l} Q_{s, m_s^{(j)}(\cdot)}(X_s^{(j)}, x) m_s^{(j)}(x) dx \right)^{-1}} ds \\
& = \int_0^t \left\| \dot{X}_s - \int_{\mathbb{R}^l} A_s(X_s, x) m_s(x) dx - \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} \right. \right. \\
& \quad \left. \left. - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx \right\|_{\left(\int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx \right)^{-1}} ds. \quad (8.30)
\end{aligned}$$

On subtracting two versions of (8.29b) for λ differing by a sign,

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \int_{\mathbb{R}^l} \Psi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x)^T c_s(X_s^{(j)}, x) \left(\frac{Dm_s^{(j)}(x)}{2m_s^{(j)}(x)} - \Phi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \right) m_s^{(j)}(x) dx \\
& = \int_{\mathbb{R}^l} \Psi_{s, m_s(\cdot), X_s}(x)^T c_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx, \quad (8.31)
\end{aligned}$$

so, on using (2.17b),

$$\begin{aligned}
& \lim_{j \rightarrow \infty} \int_{\mathbb{R}^l} G_s(X_s^{(j)}, x) \left(\frac{Dm_s^{(j)}(x)}{2m_s^{(j)}(x)} - \Phi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \right) m_s^{(j)}(x) dx \\
& = \int_{\mathbb{R}^l} G_s(X_s, x) \left(\frac{Dm_s(x)}{2m_s(x)} - \Phi_{s, m_s(\cdot), X_s}(x) \right) m_s(x) dx. \quad (8.32)
\end{aligned}$$

By (8.29b) and (8.31),

$$\lim_{j \rightarrow \infty} \int_{\mathbb{R}^l} \left\| \Psi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \right\|_{c_s(X_s^{(j)}, x)}^2 m_s^{(j)}(x) dx = \int_{\mathbb{R}^l} \left\| \Psi_{s, m(\cdot), X_s}(x) \right\|_{c_s(X_s, x)}^2 m_s(x) dx,$$

so, by (2.16),

$$\lim_{j \rightarrow \infty} \int_{\mathbb{R}^l} Q_{s, m_s^{(j)}(\cdot)}(X_s^{(j)}, x) m_s^{(j)}(x) dx = \int_{\mathbb{R}^l} Q_{s, m_s(\cdot)}(X_s, x) m_s(x) dx. \quad (8.33)$$

By (8.28), (8.7a), (8.7b), (8.7c), (A.2), and (8.33), one has pointwise convergence of the integrands with respect to ds on the lefthand side of (8.30) to the integrand on the righthand side.

Since the matrices $\int_{\mathbb{R}^l} Q_{s, m_s^{(j)}(\cdot)}(X_s^{(j)}, x)$

$m_s^{(j)}(x) dx$ are uniformly positive definite, the j th integrand is bounded above by

$$M \left(|\dot{X}_s|^2 + 1 + \int_{\mathbb{R}^l} \left| \frac{Dm_s^{(j)}(x)}{2m_s^{(j)}(x)} - \Phi_{s, m_s^{(j)}(\cdot), X_s^{(j)}}(x) \right|^2 m_s^{(j)}(x) dx \right),$$

for some $M > 0$. By (8.29a) and (8.29b), by the fact that the right hand side of (6.23) is finite, so $\int_0^t |\dot{X}_s|^2 ds < \infty$, and by dominated convergence, we conclude that (8.30) holds.

We have thus proved that in both cases there exist $(X^{(j)}, \mu^{(j)})$ with needed regularity properties that converge to (X, μ) and are such that $\mathbf{I}_t^{**}(X^{(j)}, \mu^{(j)}) \rightarrow \mathbf{I}_t^{**}(X, \mu)$ for all $t \in \mathbb{R}_+$. Picking a suitable subsequence, we can assume that $\mathbf{I}_j^{**}(X^{(j)}, \mu^{(j)}) \leq \mathbf{I}_j^{**}(X, \mu) + 1/j$. We redefine the subsequence $(X_t^{(j)}, \mu_t^{(j)})$ for $t \geq j$ such that $\mathbf{I}_j^{**}(X^{(j)}, \mu^{(j)}) = \mathbf{I}^{**}(X^{(j)}, \mu^{(j)})$, thanks to Lemma 6.7. The obtained sequence will still converge to (X, μ) . In addition, $\limsup_{j \rightarrow \infty} \mathbf{I}^{**}(X^{(j)}, \mu^{(j)}) \leq \mathbf{I}^{**}(X, \mu)$, which yields the assertion of Theorem 8.1 by the lower semicontinuity of \mathbf{I}^{**} . \square

9 Proof of Theorem 2.1

Suppose that $\lim_{\epsilon \rightarrow 0} \mathbf{P}^\epsilon(|X_0^\epsilon - \hat{u}| > \kappa)^\epsilon = 0$, for arbitrary $\kappa > 0$. Then any large deviation limit point $\tilde{\mathbf{I}}$ of \mathbf{P}^ϵ is such that $\tilde{\mathbf{I}}(X, \mu) = \infty$ unless $X_0 = \hat{u}$. If (X, μ) is such that $X_0 = \hat{u}$ and $\mathbf{I}^{**}(X, \mu) < \infty$, by Theorems 5.1, 7.1, and 8.1, there exist (X^i, μ^i) , which satisfy the hypotheses on $(\tilde{X}, \tilde{\mu})$ in Theorem 7.1, such that $\mathbf{I}^{**}(X^i, \mu^i) = \tilde{\mathbf{I}}(X^i, \mu^i)$, $(X^i, \mu^i) \rightarrow (X, \mu)$ as $i \rightarrow \infty$, and $\mathbf{I}^{**}(X^i, \mu^i) \rightarrow \mathbf{I}^{**}(X, \mu)$ as $i \rightarrow \infty$. By Theorem 3.4 (with the role of \mathcal{U} being played by the set of functions $U_{t \wedge \tau}^{\lambda(\cdot), f}$ in Theorem 5.1 and with the role of $\tilde{\mathcal{U}}$ being played by the set of functions θ^N in Lemma 7.2) and Theorem 6.1, $\tilde{\mathbf{I}}(X, \mu) = \mathbf{I}^{**}(X, \mu) = \mathbf{I}(X, \mu)$ for all (X, μ) .

In the general setting of Theorem 2.1, let $\mathcal{L}_{\hat{u}}^\epsilon$ denote the regular conditional distribution of $(X^\epsilon, \mu^\epsilon)$ given that $X_0^\epsilon = \hat{u}$, where $\hat{u} \in \mathbb{R}^n$ and is otherwise arbitrary. By what has been proved, if $u^\epsilon \rightarrow \hat{u}$ as $\epsilon \rightarrow 0$, then the $\mathcal{L}_{u^\epsilon}^\epsilon$ obey the LDP in $\mathbb{C}(\mathbb{R}_+, \mathbb{R}^n) \times \mathbb{C}_\uparrow(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ with the large deviation function $\tilde{\mathbf{I}}_{\hat{u}}$ as defined in the statement of Theorem 2.1, where $\mathbf{I}_0(\hat{u}) = 0$ and $\mathbf{I}_0(u) = \infty$ if $u \neq \hat{u}$. Since by the hypotheses of Theorem 2.1 the distributions of X_0^ϵ obey the LDP with a large deviation function \mathbf{I}_0 , it follows that the distributions of $(X^\epsilon, \mu^\epsilon)$ obey the LDP with $\mathbf{I}(X, \mu) = \mathbf{I}_0(X_0) + \tilde{\mathbf{I}}_{X_0}(X, \mu)$, see, e.g., Chaganty [9], Puhalskii [39]. Theorem 2.1 has been proved.

A Appendix

Proof of Lemma 2.1. Let $\eta_r(x) = \eta(|x|/r)$ and

$$k_s(x, \lambda) = c_s(X_s, x)^{-1} \left(\frac{\operatorname{div}(c_s(X_s, x)m_s(x))}{2m_s(x)} - (a_s(X_s, x) + G_s(X_s, x)^T \lambda) \right). \quad (\text{A.1})$$

By (2.9) and Theorem 6.1, $k_s(\cdot, \lambda) \in \mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)$, for almost all s . We prove that, for those s ,

$$\begin{aligned} \lim_{r \rightarrow \infty} \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ = \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx. \quad (\text{A.2}) \end{aligned}$$

Since

$$\begin{aligned} \lim_{r \rightarrow \infty} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ = \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx, \end{aligned}$$

we have that

$$\begin{aligned} \liminf_{r \rightarrow \infty} \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ \geq \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx, \end{aligned}$$

so, we need to prove that

$$\begin{aligned} \limsup_{r \rightarrow \infty} \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ \leq \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx. \quad (\text{A.3}) \end{aligned}$$

On denoting by H_r the Hilbert space $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), \eta_r^2(x) m_s(x) dx)$, one can write (A.3) as

$$\begin{aligned} \limsup_{r \rightarrow \infty} \sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ \leq \sup_{Dh \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx. \quad (\text{A.4}) \end{aligned}$$

Given $R > 0$ and $\kappa \in (0, 1)$, for all r great enough,

$$\begin{aligned} \sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ \leq \sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1-\kappa}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ + \int_{\mathbb{R}^l} \sup_{y \in \mathbb{R}^l} (y^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| > R\}}(x) - \frac{\kappa}{2} \|y\|_{c_s(X_s, x)}^2) \eta_r^2(x) m_s(x) dx \\ = \frac{1}{1-\kappa} \sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x))^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\ + \frac{1}{2\kappa} \int_{\mathbb{R}^l} \|k_s(x, \lambda)\|_{c_s(X_s, x)}^2 \mathbf{1}_{\{|x| > R\}}(x) \eta_r^2(x) m_s(x) dx \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{1-\kappa} \sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x)) m_s(x) dx \\
&\quad + \frac{1}{2\kappa} \int_{\mathbb{R}^l} \|k_s(x, \lambda)\|_{c_s(X_s, x)}^2 \mathbf{1}_{\{|x| > R\}}(x) m_s(x) dx. \quad (\text{A.5})
\end{aligned}$$

If $\|Dh(\cdot)\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), \eta_r^2(x) m_s(x) dx)} > 2\|k_s(x, \lambda)\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)}$, then, by the Cauchy–Schwarz inequality, the first integral on the rightmost side of (A.5) is negative. Therefore, on denoting by K_r the subset of H_r of (the equivalence classes of) functions Dh such that $\|Dh\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), \eta_r^2(x) m_s(x) dx)} \leq 2\|k_s(x, \lambda)\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)}$, we have that

$$\begin{aligned}
&\sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x)) m_s(x) dx \\
&= \sup_{Dh \in K_r} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x)) m_s(x) dx. \quad (\text{A.6})
\end{aligned}$$

Being closed subspaces of the reflexive Banach spaces $\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), \eta_r^2(x) m_s(x) dx)$, the spaces H_r are reflexive Banach spaces in their own right. The sets K_r are weakly compact subsets of the H_r , being convex, bounded, and closed. The K_r make up an inverse system of compact Hausdorff spaces with identifications of elements of K_r as elements of $K_{r'}$, for $r > r'$, as bonding maps $\pi_{rr'}$. The inverse limit of the K_r , as $r \rightarrow \infty$, which we denote by K is a compact Hausdorff space and can be identified with the subspace of $\mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), (x) m_s(x) dx)$ of the elements Dh such that $\|Dh\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)} \leq 2\|k_s(x, \lambda)\|_{\mathbb{L}^2(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)}$. The projections from K to K_r are identifications of the elements of K as elements of K_r . For each r , the integral in (A.6) is a concave and continuous function of Dh for the norm topology on H_r , so, it is weakly upper semicontinuous on H_r , and, hence, on K_r . Those integrals decrease as r increases and converge to $\int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \|Dh(x)\|_{c_s(X_s, x)}^2 / 2) m_s(x) dx$, as $r \rightarrow \infty$. By the version of Dini's lemma in Remark A.1,

$$\begin{aligned}
&\lim_{r \rightarrow \infty} \sup_{Dh \in K_r} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x)) m_s(x) dx = \\
&\quad \sup_{Dh \in K} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2) m_s(x) dx. \quad (\text{A.7})
\end{aligned}$$

By (A.5), (A.6), and (A.7),

$$\begin{aligned}
&\limsup_{r \rightarrow \infty} \sup_{Dh \in H_r} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2) \eta_r^2(x) m_s(x) dx \\
&\leq \frac{1}{1-\kappa} \sup_{Dh \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x) dx)} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) \\
&\quad - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2) m_s(x) dx + \frac{1}{2\kappa} \int_{\mathbb{R}^l} \|k_s(x, \lambda)\|_{c_s(X_s, x)}^2 \mathbf{1}_{\{|x| > R\}}(x) m_s(x) dx. \quad (\text{A.8})
\end{aligned}$$

Since, similarly to (A.7),

$$\begin{aligned}
& \sup_{Dh \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x))} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2) m_s(x) dx \\
& \leq \frac{1}{1 - \kappa} \sup_{Dh \in \mathbb{L}_0^{1,2}(\mathbb{R}^l, \mathbb{R}^l, c_s(x, X_s), m_s(x))} \int_{\mathbb{R}^l} (Dh(x)^T c_s(X_s, x) k_s(x, \lambda) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2) m_s(x) dx \\
& \quad + \frac{1}{2\kappa} \int_{\mathbb{R}^l} \|k_s(x, \lambda)\|_{c_s(X_s, x)}^2 \mathbf{1}_{\{|x| > R\}}(x) m_s(x) dx,
\end{aligned}$$

one obtains (A.4) by letting $R \rightarrow \infty$ and $\kappa \rightarrow 0$ on the righthand side of (A.8) and accounting for $k_s(\cdot, \lambda)$ being square integrable. The convergence in (2.8) is obtained by an application of Lebesgue's dominated convergence theorem.

By Theorem 6.1, if $\mathbf{I}'(X, \mu) < \infty$, then

$$\int_0^t \sup_{h \in \mathbb{C}_0^1(\mathbb{R}^l)} \int_{\mathbb{R}^l} (Dh(x)^T (\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x)) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2) m_s(x) dx ds < \infty. \quad (\text{A.9})$$

Suppose (2.10) holds and let L denote an upper bound for the lefthand side of (A.9). By (6.6) in the statement of Lemma 6.4 and Condition 2.1, L is also an upper bound on the integrals on the left of (A.9) for $h \in \mathbb{W}_0^{1,q}(S)$, where S is an open ball in \mathbb{R}^l , $q \geq 2$, and $q > l$. On taking $h(x) = \kappa(|x|^2 \vee r_1^2 \wedge r_2^2 - r_2^2)$, where $\kappa > 0$ and $0 < r_1 < r_2$, we have that

$$\int_0^t \int_{x \in \mathbb{R}^l: r_1 \leq |x| \leq r_2} (\kappa x^T (\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x)) - \kappa^2 \|c_s(X_s, x)\| |x|^2) m_s(x) dx ds \leq L.$$

If r_1 is great enough, there exists $\delta > 0$ such that $x^T a_s(X_s, x) \leq -\delta|x|^2$ if $|x| \geq r_1$. Therefore, for small enough $\kappa > 0$, great enough r_1 , and all $r_2 > r_1$,

$$\frac{\kappa\delta}{2} \int_0^t \int_{x \in \mathbb{R}^l: r_1 \leq |x| \leq r_2} |x|^2 m_s(x) dx ds \leq L.$$

The square integrability of $a_s(X_s, x)$ now follows by Condition 2.1.

Suppose now (2.11) holds. We take, for given s ,

$$h(x) = -(\hat{a}_s(X_s, x) \vee (-\delta) \wedge \delta) \eta_r(x).$$

Then

$$Dh(x) = -D\hat{a}_s(X_s, x) \mathbf{1}_{\{|\hat{a}_s(X_s, x)| \leq \delta\}}(x) \eta_r(x) - (\hat{a}_s(X_s, x) \vee (-\delta) \wedge \delta) \frac{1}{r} \frac{x}{|x|} D\eta\left(\frac{|x|}{r}\right),$$

$h \in \mathbb{W}_0^{1,q}(S)$, for large enough ball S , and, for $\kappa \in (0, 1)$,

$$\begin{aligned}
& \int_{\mathbb{R}^l} (Dh(x))^T \left(\frac{1}{2} \operatorname{div} c_s(X_s, x) - a_s(X_s, x) \right) - \frac{1}{2} \|Dh(x)\|_{c_s(X_s, x)}^2 m_s(x) dx \\
& \geq \frac{1-\kappa}{2} \int_{\mathbb{R}^l} \|D_x \hat{a}_s(X_s, x)\|_{c_s(X_s, x)}^2 \mathbf{1}_{\{|\hat{a}_s(X_s, x)| \leq \delta\}}(x) \eta_r(x) m_s(x) dx \\
& - \int_{\mathbb{R}^l} (\hat{a}_s(X_s, x) \vee (-\delta) \wedge \delta) \frac{1}{r} D\eta\left(\frac{|x|}{r}\right) \frac{x^T}{|x|} c_s(X_s, x) D_x \hat{a}_s(X_s, x) m_s(x) dx \\
& - \frac{1}{2r^2} \left(1 + \frac{1}{\kappa}\right) \int_{\mathbb{R}^l} (\hat{a}_s(X_s, x) \vee (-\delta) \wedge \delta)^2 \|D\eta\left(\frac{|x|}{r}\right)\|_{c_s(X_s, x)}^2 m_s(x) dx.
\end{aligned}$$

As $r \rightarrow \infty$, the integrals from 0 to t of the latter two integrals converge to zero (we recall that by Theorem 6.1, $\int_0^t \int_{\mathbb{R}^l} |x^T a_s(X_s, x)|/|x| m_s(x) dx ds < \infty$, so $\int_0^t \int_{\mathbb{R}^l} |x^T c_s(X_s, x) D_x \hat{a}_s(X_s, x)|/|x| m_s(x) dx ds < \infty$). Therefore,

$$\frac{1}{2} \int_0^t \int_{\mathbb{R}^l} \|a_s(X_s, x) - \frac{1}{2} \operatorname{div} c_s(X_s, x)\|_{c_s(X_s, x)^{-1}}^2 m_s(x) dx ds \leq L,$$

which implies the square integrability of $a_s(X_s, \cdot)$ thanks to Conditions 2.1 and 2.2. \square

Remark A.1. The version of Dini's lemma invoked in the proof is as follows. Let $(K_\sigma, \pi_{\sigma\tau}, \Sigma)$ be an inverse system of compact Hausdorff topological spaces over directed set Σ . In particular, for $\tau \leq \sigma$, $\pi_{\sigma\tau}$ is a continuous map from K_σ to K_τ . Let K represent the inverse limit of the K_σ and let $\pi_\sigma : K \rightarrow K_\sigma$ be the canonical projections. We note that K is a compact Hausdorff topological space. Let real-valued functions f_σ be defined and upper semicontinuous on the K_σ . Suppose that the sequence f_σ is monotonically decreasing in the sense that if $x_\sigma \in K_\sigma$, then $f_\sigma(x_\sigma) \leq f_\tau(\pi_{\sigma\tau} x_\sigma)$, for all $\tau \leq \sigma$. Let $x \in K$. Since, for $\tau \leq \sigma$, $f_\tau(\pi_\tau x) = f_\tau(\pi_{\sigma\tau} \circ \pi_\sigma x) \geq f_\sigma(\pi_\sigma x)$, the net $f_\sigma(\pi_\sigma x)$ is monotonically decreasing. Let $f(x)$ represent the limit of $f_\sigma(\pi_\sigma x)$, as $\sigma \in \Sigma$. Then $\sup_{x_\sigma \in K_\sigma} f_\sigma(x_\sigma) \rightarrow \sup_{x \in K} f(x)$. For a proof, one notes that the sets $\pi_\sigma^{-1}(\{x_\sigma \in K_\sigma : f_\sigma(x_\sigma) \geq \sup_{x \in K} f(x) + \epsilon\})$ form a nested collection of closed subsets of K with an empty intersection. (There is also a sequential version of this result.)

Remark A.2. A "down-to-earth" version of the proof of (A.7) goes as follows. Let the supremum on the lefthand side be attained at Dh_r . All of the Dh_r belong to K_{r_0} , provided $r \geq r_0$. By the diagonal process, we may assume that the Dh_r converge weakly to some $D\hat{h}$ that belongs to all K_{r_0} , so it belongs to K . We also have that

$$\begin{aligned}
& \int_{\mathbb{R}^l} (Dh_r(x))^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh_r(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x) m_s(x) dx \\
& \leq \int_{\mathbb{R}^l} (Dh_r(x))^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh_r(x)\|_{c_s(X_s, x)}^2 \eta_{r_0}^2(x) m_s(x) dx.
\end{aligned}$$

By the upper semicontinuity in K_{r_0} ,

$$\begin{aligned} \limsup_{r \rightarrow \infty} \int_{\mathbb{R}^l} (Dh_r(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh_r(x)\|_{c_s(X_s, x)}^2 \eta_{r_0}^2(x)) m_s(x) dx \\ \leq \int_{\mathbb{R}^l} (D\hat{h}(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|D\hat{h}(x)\|_{c_s(X_s, x)}^2 \eta_{r_0}^2(x)) m_s(x) dx. \end{aligned}$$

Letting $r_0 \rightarrow \infty$ yields

$$\begin{aligned} \limsup_{r \rightarrow \infty} \int_{\mathbb{R}^l} (Dh_r(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|Dh_r(x)\|_{c_s(X_s, x)}^2 \eta_r^2(x)) m_s(x) dx \\ \leq \int_{\mathbb{R}^l} (D\hat{h}(x)^T c_s(X_s, x) k_s(x, \lambda) \mathbf{1}_{\{|x| \leq R\}}(x) - \frac{1}{2} \|D\hat{h}(x)\|_{c_s(X_s, x)}^2) m_s(x) dx. \end{aligned}$$

Proof of Lemma 4.1. By Ascoli's theorem, set $\Gamma \subset \mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$ is relatively compact if and only if for each $t \in \mathbb{R}_+$ there exists compact $K_t \subset \mathbb{M}(\mathbb{R}^l)$ such that $\tilde{\nu}_t \in K_t$ for all $\tilde{\nu} = (\tilde{\nu}_s, s \in \mathbb{R}_+) \in \Gamma$ and

$$\limsup_{\delta \rightarrow 0} \sup_{\tilde{\nu} \in \Gamma} \sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| \leq \delta} d(\tilde{\nu}_{s_1}, \tilde{\nu}_{s_2}) = 0. \quad (\text{A.10})$$

Suppose, the net ν_ϵ is sequentially exponentially tight for rate $1/\epsilon$. Given arbitrary $t > 0$ and $\eta > 0$, let $\{\epsilon_i, i \in \mathbb{N}\}$ be a sequence converging to zero as $i \rightarrow \infty$ such that

$$\lim_{i \rightarrow \infty} \mathbf{P}_{\epsilon_i}(\nu_{\epsilon_i, t}(x \in \mathbb{R}^l : |x| > i) > \eta)^{\epsilon_i} = \lim_{N \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \mathbf{P}_\epsilon(\nu_{\epsilon, t}(x \in \mathbb{R}^l : |x| > N) > \eta)^\epsilon$$

and

$$\lim_{i \rightarrow \infty} \sup_{s_1 \in [0, t]} \mathbf{P}_{\epsilon_i} \left(\sup_{s_2 \in [s_1, s_1 + 1/i]} d(\nu_{\epsilon_i, s_1}, \nu_{\epsilon_i, s_2}) > \eta \right)^{\epsilon_i} = \lim_{\delta \rightarrow 0} \limsup_{\epsilon \rightarrow 0} \sup_{s_1 \in [0, t]} \mathbf{P}_\epsilon \left(\sup_{s_2 \in [s_1, s_1 + \delta]} d(\nu_{\epsilon, s_1}, \nu_{\epsilon, s_2}) > \eta \right)^\epsilon.$$

Since the sequence $\{\nu_{\epsilon_i}, i \in \mathbb{N}\}$ is exponentially tight for rate $1/\epsilon_i$, given $\kappa > 0$, there exists compact Γ such that $\limsup_{i \rightarrow \infty} \mathbf{P}_{\epsilon_i}(\nu_{\epsilon_i} \notin \Gamma)^{\epsilon_i} < \kappa$. Hence, for the associated K_t , $\limsup_{i \rightarrow \infty} \mathbf{P}_{\epsilon_i}(\nu_{\epsilon_i, t} \notin K_t)^{\epsilon_i} < \kappa$. By compactness of K_t , for arbitrary $\eta > 0$ there exists i such that $\sup_{\tilde{\nu} \in K_t} \tilde{\nu}(x : |x| > i) \leq \eta$, so $\limsup_{i \rightarrow \infty} \mathbf{P}_{\epsilon_i}(\nu_{\epsilon_i, t}(x : |x| > i) > \eta)^{\epsilon_i} < \kappa$. Similarly, choosing i in such a way that $\sup_{\nu \in \Gamma} \sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| < 1/i} d(\nu_{s_1}, \nu_{s_2}) \leq \eta$ we obtain that $\limsup_{i \rightarrow \infty} \mathbf{P}_{\epsilon_i}(\sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| < 1/i} d(\nu_{\epsilon_i, s_1}, \nu_{\epsilon_i, s_2}) > \eta)^{\epsilon_i} < \kappa$, which implies that $\limsup_{i \rightarrow \infty} \sup_{s_1 \in [0, t]} \mathbf{P}_{\epsilon_i}(\sup_{s_2 \in [s_1, s_1 + 1/i]} d(\nu_{\epsilon_i, s_1}, \nu_{\epsilon_i, s_2}) > \eta)^{\epsilon_i} < \kappa$.

Suppose now that the convergences in the hypotheses hold. We first show that

$$\lim_{\delta \rightarrow 0} \limsup_{\epsilon \rightarrow 0} \mathbf{P}_\epsilon \left(\sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| \leq \delta} d(\nu_{\epsilon, s_1}, \nu_{\epsilon, s_2}) > \eta \right)^\epsilon = 0.$$

It follows by the bound (cf. Billingsley [5, Chapter 2])

$$\begin{aligned} \mathbf{P}_\epsilon \left(\sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| \leq \delta} d(\nu_{\epsilon, s_1}, \nu_{\epsilon, s_2}) > \eta \right) &\leq \mathbf{P}_\epsilon \left(\bigcup_{i=0}^{\lfloor t/\delta \rfloor} \{3 \sup_{s \in [i\delta, (i+1)\delta]} d(\nu_{\epsilon, i\delta}, \nu_{\epsilon, s}) > \eta\} \right) \\ &\leq \sum_{i=0}^{\lfloor t/\delta \rfloor} \mathbf{P}_\epsilon \left(3 \sup_{s \in [i\delta, (i+1)\delta]} d(\nu_{\epsilon, i\delta}, \nu_{\epsilon, s}) > \eta \right) \leq \left(\frac{t}{\delta} + 1 \right) \sup_{s_1 \in [0, t]} \mathbf{P}_\epsilon \left(\sup_{s_2 \in [s_1, s_1 + \delta]} d(\nu_{\epsilon, s_1}, \nu_{\epsilon, s_2}) > \frac{\eta}{3} \right). \end{aligned}$$

Let $\epsilon_i \rightarrow 0$ as $i \rightarrow \infty$. Given $\kappa > 0$, $t \in \mathbb{R}_+$, and $j \in \mathbb{N}$, one can choose $\delta_{j,t} > 0$ such that $\delta_{j,t} \rightarrow 0$ as $j \rightarrow \infty$ and

$$\limsup_{i \rightarrow \infty} \mathbf{P}_{\epsilon_i} \left(\sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| \leq \delta_{j,t}} d(\nu_{\epsilon_i, s_1}, \nu_{\epsilon_i, s_2}) > \frac{1}{j} \right)^{\epsilon_i} < \frac{\kappa}{2^j 2^t},$$

for all $i \in \mathbb{N}$ and all $j \in \mathbb{N}$. Since the space $\mathbb{M}(\mathbb{R}^l)$ is complete and separable, each of the measures ν_ϵ has a tight distribution, so we can choose $\tilde{\delta}_{j,t} > 0$ such that $\tilde{\delta}_{j,t} \rightarrow 0$ as $j \rightarrow \infty$ and the inequality

$$\mathbf{P}_{\epsilon_i} \left(\sup_{s_1, s_2 \in [0, t]: |s_1 - s_2| \leq \tilde{\delta}_{j,t}} d(\nu_{\epsilon_i, s_1}, \nu_{\epsilon_i, s_2}) > \frac{1}{j} \right)^{\epsilon_i} < \frac{\kappa}{2^j 2^t} \quad (\text{A.11})$$

holds for all i . Similarly, one can choose $N_{j,t} \rightarrow \infty$ as $j \rightarrow \infty$ such that the inequality

$$\mathbf{P}_{\epsilon_i} \left(\max_{L=1, 2, \dots, \lfloor t/\tilde{\delta}_{j,t} \rfloor} \nu_{\epsilon_i, L\tilde{\delta}_{j,t}}(x \in \mathbb{R}^l : |x| > N_{j,t}) > \frac{1}{j} \right)^{\epsilon_i} < \frac{\kappa}{2^j 2^t} \quad (\text{A.12})$$

holds for all $i \in \mathbb{N}$.

Let

$$\Gamma = \bigcap_{M \in \mathbb{N}} \bigcap_{j \in \mathbb{N}} \left(\{ \tilde{\nu} \in \mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l)) : \sup_{s_1, s_2 \in [0, M]: |s_1 - s_2| \leq \tilde{\delta}_{j,M}} d(\tilde{\nu}_{s_1}, \tilde{\nu}_{s_2}) \leq \frac{1}{j} \} \right. \\ \left. \bigcap_{L=1}^{\lfloor M/\tilde{\delta}_{j,M} \rfloor} \{ \tilde{\nu} \in \mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l)) : \tilde{\nu}_{L\tilde{\delta}_{j,M}}(|x| > N_{j,M}) \leq \frac{1}{j} \} \right).$$

According to the definition, (A.10) holds for all $t \in \mathbb{R}_+$. In addition, if $M \geq t$, then, for $\tilde{\nu} \in \Gamma$, using the definition of $d(\cdot, \cdot)$,

$$\tilde{\nu}_t(|x| > N_{j,M} + 1) \leq \max_{L=1, 2, \dots, \lfloor M/\tilde{\delta}_{j,M} \rfloor} \tilde{\nu}_{L\tilde{\delta}_{j,M}}(|x| > N_{j,M}) + \frac{1}{j} \leq \frac{2}{j}.$$

Hence,

$$\lim_{j \rightarrow \infty} \sup_{\tilde{\nu} \in \Gamma} \tilde{\nu}_t(|x| > N_{j,M} + 1) = 0,$$

so by Prohorov's theorem, the $\tilde{\nu}_t$ belong to a compact set. It follows that Γ is relatively compact in $\mathbb{C}(\mathbb{R}_+, \mathbb{M}(\mathbb{R}^l))$.

In addition, by (A.11) and (A.12), assuming $\epsilon_i < 1$,

$$\mathbf{P}_{\epsilon_i}(\nu_{\epsilon_i} \notin \Gamma)^{\epsilon_i} \leq \sum_{M=1}^{\infty} \sum_{j=1}^{\infty} \left(\mathbf{P}_{\epsilon_i} \left(\sup_{s_1, s_2 \in [0, M]: |s_1 - s_2| \leq \tilde{\delta}_{j,M}} d(\nu_{\epsilon_i, s_1}, \nu_{\epsilon_i, s_2}) > \frac{1}{j} \right)^{\epsilon_i} \right. \\ \left. + \mathbf{P}_{\epsilon_i} \left(\max_{L=1, 2, \dots, \lfloor M/\tilde{\delta}_{j,M} \rfloor} \nu_{\epsilon_i, L\tilde{\delta}_{j,M}}(x \in \mathbb{R}^l : |x| > N_{j,M}) > \frac{1}{j} \right)^{\epsilon_i} \right) \leq 2\kappa.$$

Thus, the sequence $\{\nu_{\epsilon_i}, i \in \mathbb{N}\}$ is exponentially tight for rate $1/\epsilon_i$ as $i \rightarrow \infty$. This completes the proof of the first part of the lemma. The second part is proved similarly. \square

Proof of Lemma 6.2. We follow the approach of Röckner and Zhang [44, pp.204,205], [45], see also Bogachev, Krylov, and Röckner [6]. Let ψ be a bounded function from $\mathbb{W}^{1,2}(O, m(x) dx)$. Let, for $j \in \mathbb{N}$, b_j represent an \mathbb{R}_+ -valued \mathbb{C}_0^∞ -function on \mathbb{R} such that, for $y \in \mathbb{R}$, $\mathbf{1}_{[-j, j]}(y) \leq b_j(y) \leq$

$\mathbf{1}_{(-j-1, j+1)}(y)$ and $|Db_j(y)| \leq 2$, and let $\phi_j(x) = b_j(\ln m(x))$ if $m(x) > 0$ and $\phi_j(x) = 0$ if $m(x) = 0$, where $x \in \mathbb{R}^d$. It is noteworthy that $\phi_j \in \mathbb{W}^{1,2}(O)$. We have that

$$\int_O |\psi(x) - \phi_j(x)\psi(x)|^2 m(x) dx \leq \int_O |\psi(x)|^2 \mathbf{1}_{\{|\ln m(x)| \geq j\}}(x) m(x) dx \rightarrow 0 \text{ as } j \rightarrow \infty.$$

In addition,

$$\begin{aligned} \int_O |D\psi(x) - D(\phi_j(x)\psi(x))|^2 m(x) dx &\leq 2 \int_O |D\psi(x)|^2 \mathbf{1}_{\{|\ln m(x)| \geq j\}}(x) m(x) dx \\ &+ 8 \int_O |\psi(x)|^2 \mathbf{1}_{\{|\ln m(x)| \geq j\}}(x) \frac{|Dm(x)|^2}{m(x)} dx \rightarrow 0 \text{ as } j \rightarrow \infty. \end{aligned}$$

Thus, $\phi_j\psi \rightarrow \psi$ in $\mathbb{W}^{1,2}(O, m(x) dx)$ as $j \rightarrow \infty$.

We have that

$$\int_O |\phi_j(x)\psi(x)|^2 dx = \int_O |\phi_j(x)\psi(x)|^2 \mathbf{1}_{\{m(x) > e^{-j-1}\}} dx \leq e^{j+1} \int_O |\psi(x)|^2 m(x) dx.$$

Similarly,

$$\int_O |D(\phi_j(x)\psi(x))|^2 dx \leq 8e^{j+1} \int_O \frac{|Dm(x)|^2}{m(x)} |\psi(x)|^2 dx + 2e^{j+1} \int_O |D\psi(x)|^2 m(x) dx.$$

Thus, $\phi_j\psi \in \mathbb{W}^{1,2}(O)$, so there exists sequence ψ_i of $\mathbb{C}^\infty(O)$ -functions with bounded $|\psi_i(x)|$ and $|D\psi_i(x)|$ that converges to $\phi_j\psi$ in $\mathbb{W}^{1,2}(O)$ as $i \rightarrow \infty$. The following inequalities show that $\phi_{j+1}\psi_i \rightarrow \phi_j\psi$ in $\mathbb{W}^{1,2}(O, m(x) dx)$ as $i \rightarrow \infty$: since $\phi_j = \phi_j\phi_{j+1}$,

$$\begin{aligned} &\int_O (|\phi_j(x)\psi(x) - \phi_{j+1}(x)\psi_i(x)|^2 + |D(\phi_j(x)\psi(x)) - D(\phi_{j+1}(x)\psi_i(x))|^2) m(x) dx \\ &\leq \int_O (9|\phi_j(x)\psi(x) - \psi_i(x)|^2 + 2|D(\phi_j(x)\psi(x)) - D\psi_i(x)|^2) \mathbf{1}_{\{|\ln m(x)| \leq j+2\}}(x) m(x) dx \\ &\leq e^{j+2} \int_O (9|\phi_j(x)\psi(x) - \psi_i(x)|^2 + 2|D(\phi_j(x)\psi(x)) - D\psi_i(x)|^2) dx \rightarrow 0 \text{ as } i \rightarrow \infty. \end{aligned}$$

It remains to check that $\phi_{j+1}\psi_i \in \mathbb{H}^{1,2}(O, m(x) dx)$, which follows provided $\phi_{j+1} \in \mathbb{H}^{1,2}(O, m(x) dx)$. If O is an open ball in \mathbb{R}^d , we let $\eta(x)$ represent a $\mathbb{C}_0^\infty(\mathbb{R}^d)$ -function such that $\eta(x) \in [0, 1]$ and $\eta(x) = 1$ on O and let $\tilde{m}(x) = m(x)\eta(x) + (1 - \eta(x))/(1 + |x|^{d+1})$ and $\tilde{\phi}_{j+1}(x) = b_j(\ln \tilde{m}(x))$. Then $\tilde{m}(x) = m(x)$ and $\tilde{\phi}_{j+1}(x) = \phi_{j+1}(x)$ for $x \in O$. If $O = \mathbb{R}^d$, we let $\tilde{m}(x) = m(x)$ and $\tilde{\phi}_{j+1}(x) = \phi_{j+1}(x)$ for all $x \in \mathbb{R}^d$. It suffices to prove that $\tilde{\phi}_{j+1} \in \mathbb{H}^{1,2}(\mathbb{R}^d, \tilde{m}(x) dx)$. We have that $\sqrt{\tilde{m}} \in \mathbb{W}^{1,2}(\mathbb{R}^d)$ for both cases where O is an open ball and $O = \mathbb{R}^d$. By R\"ockner and Zhang [45] (see pp.461–463), $\tilde{\phi}_{j+1}$ belongs to the domain of the closure of the Dirichlet form \mathcal{E}^0 on $\mathbb{L}^2(\mathbb{R}^d, \tilde{m}(x) dx)$ defined by $\mathcal{E}^0(f, g) = \int_{\mathbb{R}^d} Df(x)^T Dg(x) \tilde{m}(x) dx$, where $f, g \in \mathbb{C}_0^\infty(\mathbb{R}^d)$. Thus, there exist $q_i \in \mathbb{C}_0^\infty(\mathbb{R}^d)$ such that $\int_{\mathbb{R}^d} |Dq_i(x) - D\tilde{\phi}_{j+1}(x)|^2 \tilde{m}(x) dx + \int_{\mathbb{R}^d} |q_i(x) - \tilde{\phi}_{j+1}(x)|^2 \tilde{m}(x) dx \rightarrow 0$ as $i \rightarrow \infty$ which implies the needed property.

If $\psi \in \mathbb{H}^{1,2}(\mathbb{R}^d)$ is not bounded, then it is the limit of the functions $\psi \wedge i \vee (-i)$ in $\mathbb{W}^{1,2}(\mathbb{R}^d)$ as $i \rightarrow \infty$. □

Proof of Lemma 6.3. We follow the approach of Bogachev, Krylov, and Röckner [6], see also Metafunne, Pallara, and Rhandi [31]. Let

$$\rho(x) = \begin{cases} \left(\int_{y \in \mathbb{R}^d: |y| < 1} e^{1/(|y|^2-1)} dy \right)^{-1} e^{1/(|x|^2-1)}, & \text{if } |x| < 1 \\ 0, & \text{if } |x| \geq 1, \end{cases}$$

let $\rho_\epsilon(x) = (1/\epsilon^d) \rho(x/\epsilon)$, and let $p_\epsilon(x) = \int_{\mathbb{R}^d} \rho_\epsilon(x-y) p(y) dy$, where $\epsilon > 0$ and $p \in \mathbb{C}_0^\infty(\mathbb{R}^d)$. We note that $\int_{\mathbb{R}^d} \rho_\epsilon(x) dx = 1$, $p_\epsilon \in \mathbb{C}_0^\infty(\mathbb{R}^d)$, and $\int_{\mathbb{R}^d} c(x) \rho_\epsilon(x-\cdot) m(x) dx \in \mathbb{C}_0^\infty(\mathbb{R}^d, \mathbb{R}^{d \times d})$. One can see that, similarly to the calculation on p.227 of Bogachev, Krylov, and Röckner [6],

$$\begin{aligned} \int_{\mathbb{R}^d} \text{tr}(c(x) D^2 p_\epsilon(x)) m(x) dx + \int_{\mathbb{R}^d} b(x)^T D p_\epsilon(x) m(x) dx &= - \int_{\mathbb{R}^d} D p(x)^T c(x) \int_{\mathbb{R}^d} D \rho_\epsilon(x-y) m(y) dy dx \\ &- \int_{\mathbb{R}^d} D p(x)^T \int_{\mathbb{R}^d} (c(y) - c(x)) D \rho_\epsilon(x-y) m(y) dy dx + \int_{\mathbb{R}^d} D p(x)^T \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x-y) m(y) dy dx. \end{aligned}$$

Since the lefthand side equals zero, introducing

$$m_\epsilon(x) = \int_{\mathbb{R}^d} \rho_\epsilon(x-y) m(y) dy, \quad (\text{A.13})$$

we have that

$$\begin{aligned} \int_{\mathbb{R}^d} D p(x)^T c(x) D m_\epsilon(x) dx &= - \int_{\mathbb{R}^d} D p(x)^T \int_{\mathbb{R}^d} (c(y) - c(x)) D \rho_\epsilon(x-y) m(y) dy dx \\ &+ \int_{\mathbb{R}^d} D p(x)^T \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x-y) m(y) dy dx. \quad (\text{A.14}) \end{aligned}$$

Let $\eta \in \mathbb{C}_0^\infty(\mathbb{R}^d)$ be such that $|\eta(x)| \leq 1$, let $\kappa > 0$, and let $p(x) = \eta^2(x) \ln(m_\epsilon(x) + \kappa)$. By (A.14),

$$\begin{aligned} \int_{\mathbb{R}^d} \eta^2(x) \frac{D m_\epsilon(x)^T c(x) D m_\epsilon(x)}{m_\epsilon(x) + \kappa} dx &= - \int_{\mathbb{R}^d} \ln(m_\epsilon(x) + \kappa) 2\eta(x) D\eta(x)^T c(x) D m_\epsilon(x) dx \\ &- \int_{\mathbb{R}^d} \eta^2(x) \frac{D m_\epsilon(x)^T}{m_\epsilon(x) + \kappa} \int_{\mathbb{R}^d} (c(y) - c(x)) D \rho_\epsilon(x-y) m(y) dy dx \\ &- \int_{\mathbb{R}^d} \ln(m_\epsilon(x) + \kappa) 2\eta(x) D\eta(x)^T \int_{\mathbb{R}^d} (c(y) - c(x)) D \rho_\epsilon(x-y) m(y) dy dx \\ &+ \int_{\mathbb{R}^d} \eta^2(x) \frac{D m_\epsilon(x)^T}{m_\epsilon(x) + \kappa} \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x-y) m(y) dy dx \\ &+ \int_{\mathbb{R}^d} \ln(m_\epsilon(x) + \kappa) 2\eta(x) D\eta(x)^T \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x-y) m(y) dy dx. \end{aligned}$$

The next step is to provide bounds on the terms on the righthand side. By the Cauchy inequality, for arbitrary $\delta > 0$,

$$\begin{aligned}
& \left| \int_{\mathbb{R}^d} \ln(m_\epsilon(x) + \kappa) 2\eta(x) D\eta(x)^T c(x) Dm_\epsilon(x) dx \right| \leq \delta \int_{\mathbb{R}^d} \eta^2(x) \frac{Dm_\epsilon(x)^T c(x) Dm_\epsilon(x)}{m_\epsilon(x) + \kappa} dx \\
& + \frac{1}{\delta} \int_{\mathbb{R}^d} (\ln(m_\epsilon(x) + \kappa))^2 (m_\epsilon(x) + \kappa) D\eta(x)^T c(x) D\eta(x) dx, \\
& \left| \int_{\mathbb{R}^d} \eta^2(x) \frac{Dm_\epsilon(x)^T}{m_\epsilon(x) + \kappa} \int_{\mathbb{R}^d} (c(y) - c(x)) D\rho_\epsilon(x - y) m(y) dy dx \right| \\
& \leq \delta \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} |Dm_\epsilon(x)|^2 dx + \frac{1}{4\delta} \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} \left| \int_{\mathbb{R}^d} (c(y) - c(x)) D\rho_\epsilon(x - y) m(y) dy \right|^2 dx, \\
& \left| \int_{\mathbb{R}^d} \ln(m_\epsilon(x) + \kappa) 2\eta(x) D\eta(x)^T \int_{\mathbb{R}^d} (c(y) - c(x)) D\rho_\epsilon(x - y) m(y) dy dx \right| \\
& \leq \int_{\mathbb{R}^d} (\ln(m_\epsilon(x) + \kappa))^2 (m_\epsilon(x) + \kappa) |D\eta(x)|^2 dx + \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} \left| \int_{\mathbb{R}^d} (c(y) - c(x)) D\rho_\epsilon(x - y) m(y) dy \right|^2 dx, \\
& \left| \int_{\mathbb{R}^d} \eta^2(x) \frac{Dm_\epsilon(x)^T}{m_\epsilon(x) + \kappa} \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x - y) m(y) dy dx \right| \\
& \leq \delta \int_{\mathbb{R}^d} \eta^2(x) \frac{|Dm_\epsilon(x)|^2}{m_\epsilon(x) + \kappa} dx + \frac{1}{4\delta} \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} \left| \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x - y) m(y) dy \right|^2 dx, \\
& \int_{\mathbb{R}^d} \ln(m_\epsilon(x) + \kappa) 2\eta(x) D\eta(x)^T \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x - y) m(y) dy dx \\
& \leq \int_{\mathbb{R}^d} (\ln(m_\epsilon(x) + \kappa))^2 (m_\epsilon(x) + \kappa) |D\eta(x)|^2 dx + \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} \left| \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x - y) m(y) dy \right|^2 dx.
\end{aligned}$$

Therefore, assuming $\vartheta > 0$ is such that $\vartheta \leq (y/|y|)^T c(x) (y/|y|) \leq \vartheta^{-1}$ for all $x, y \in \mathbb{R}^d$,

$$\begin{aligned}
& (\vartheta - \delta\vartheta^{-1} - 2\delta) \int_{\mathbb{R}^d} \eta^2(x) \frac{|Dm_\epsilon(x)|^2}{m_\epsilon(x) + \kappa} dx \leq \left(\frac{\vartheta^{-1}}{\delta} + 2 \right) \int_{\mathbb{R}^d} (\ln(m_\epsilon(x) + \kappa))^2 (m_\epsilon(x) + \kappa) |D\eta(x)|^2 dx \\
& + \left(\frac{1}{4\delta} + 1 \right) \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} \left| \int_{\mathbb{R}^d} (c(y) - c(x)) D\rho_\epsilon(x - y) m(y) dy \right|^2 dx \\
& + \left(\frac{1}{4\delta} + 1 \right) \int_{\mathbb{R}^d} \frac{\eta^2(x)}{m_\epsilon(x) + \kappa} \left| \int_{\mathbb{R}^d} b(y) \rho_\epsilon(x - y) m(y) dy \right|^2 dx.
\end{aligned}$$

By the Cauchy-Schwarz inequality and (A.13),

$$(\vartheta - \delta\vartheta^{-1} - 2\delta) \int_{\mathbb{R}^d} \eta^2(x) \frac{|Dm_\epsilon(x)|^2}{m_\epsilon(x) + \kappa} dx \leq \left(\frac{\vartheta^{-1}}{\delta} + 2 \right) \int_{\mathbb{R}^d} (\ln(m_\epsilon(x) + \kappa))^2 (m_\epsilon(x) + \kappa) |D\eta(x)|^2 dx$$

$$\begin{aligned}
& + \left(\frac{1}{4\delta} + 1\right) \int_{\mathbb{R}^d} \eta^2(x) \int_{|x-y|<\epsilon} \frac{|(c(y) - c(x))D\rho_\epsilon(x-y)|^2}{\rho_\epsilon(x-y)} m(y) dy dx \\
& \quad + \left(\frac{1}{4\delta} + 1\right) \int_{\mathbb{R}^d} \eta^2(x) \int_{\mathbb{R}^d} |b(y)|^2 \rho_\epsilon(x-y) m(y) dy dx. \quad (\text{A.15})
\end{aligned}$$

Let $N > 0$ be such that $\eta(x) = 0$ if $|x| \geq N$. By Jensen's inequality,

$$\begin{aligned}
& \int_{\mathbb{R}^d} (\ln(m_\epsilon(x) + \kappa))^2 (m_\epsilon(x) + \kappa) |D\eta(x)|^2 dx \\
& \leq \sup_{|x| \leq N} |D\eta(x)|^2 \int_{|x| \leq N} \int_{\mathbb{R}^d} (\ln(m(y) + \kappa))^2 (m(y) + \kappa) \rho_\epsilon(x-y) dy dx \\
& \leq \sup_{|x| \leq N} |D\eta(x)|^2 \int_{|y| \leq N+\epsilon} (\ln(m(y) + \kappa))^2 (m(y) + \kappa) dy. \quad (\text{A.16})
\end{aligned}$$

Since $c(x)$ is locally Lipschitz continuous, there exists $L > 0$ which only depends on N such that $\|c(y) - c(x)\| \leq L|x-y|$ for all $\epsilon > 0$ small enough provided $|x-y| \leq \epsilon$ and $|x| \leq N$. It follows that

$$\begin{aligned}
& \int_{\mathbb{R}^d} \eta^2(x) \int_{|x-y|<\epsilon} \frac{|(c(y) - c(x))D\rho_\epsilon(x-y)|^2}{\rho_\epsilon(x-y)} m(y) dy dx \\
& \leq \int_{\mathbb{R}^d} \int_{|x-y|<\epsilon} \eta^2(x) \frac{\|c(y) - c(x)\|^2}{\epsilon^2} \frac{4|(y-x)/\epsilon|^2}{(|(y-x)/\epsilon|^2 - 1)^4} \rho_\epsilon(y-x) dx m(y) dy \\
& \leq 4L^2 \int_{|x|<1} \frac{|x|^4}{(|x|^2 - 1)^4} \rho(x) dx. \quad (\text{A.17})
\end{aligned}$$

Also,

$$\begin{aligned}
& \int_{\mathbb{R}^d} \eta^2(x) \int_{\mathbb{R}^d} |b(y)|^2 \rho_\epsilon(x-y) m(y) dy dx \leq \int_{|x| \leq N} \int_{|y| \leq N+\epsilon} |b(y)|^2 \rho_\epsilon(x-y) m(y) dy dx \\
& \leq \int_{|y| \leq N+\epsilon} |b(y)|^2 m(y) dy. \quad (\text{A.18})
\end{aligned}$$

Combining (A.15), (A.16), (A.17), and (A.18) and assuming that $\delta > 0$ is small enough yields

$$\begin{aligned}
& \limsup_{\epsilon \rightarrow 0} \int_{\mathbb{R}^d} \eta^2(x) \frac{|Dm_\epsilon(x)|^2}{m_\epsilon(x)} dx \leq \frac{1}{\vartheta - \delta\vartheta^{-1} - 2\delta} \left(\left(\frac{\vartheta^{-1}}{\delta} + 2\right) \sup_{|x| \leq N} |D\eta(x)|^2 \int_{|x| \leq N} (\ln m(x))^2 m(x) dx \right. \\
& \quad \left. + \left(\frac{1}{4\delta} + 1\right) 4L^2 \int_{|x|<1} \frac{|x|^4}{(|x|^2 - 1)^4} \rho(x) dx + \left(\frac{1}{4\delta} + 1\right) \int_{|x| \leq N} |b(x)|^2 m(x) dx \right). \quad (\text{A.19})
\end{aligned}$$

Thus, the net $\sqrt{m_\epsilon}$ is weakly relatively compact in $\mathbb{W}^{1,2}(S)$, so it is strongly relatively compact in $\mathbb{L}^2(S)$ for any open ball $S \subset \mathbb{R}^d$, see the Rellich-Kondrashov theorem on p.168 of Adams and

Fournier [1]. Hence, the net m_ϵ is strongly relatively compact in $\mathbb{L}^1(S)$. Furthermore, any limit point of $\sqrt{m_\epsilon}$ in the weak topology of $\mathbb{W}^{1,2}(S)$ is a limit point of m_ϵ in $\mathbb{L}^1(S)$. On the other hand, by (A.13), the net m_ϵ converges to m in $\mathbb{L}^1(\mathbb{R}^d)$. It follows that \sqrt{m} is the unique weak limit of $\sqrt{m_\epsilon}$ in $\mathbb{W}^{1,2}(S)$. Thus, $\sqrt{m} \in \mathbb{W}_{\text{loc}}^{1,2}(\mathbb{R}^d)$. Letting $\epsilon \rightarrow 0$ in (A.19) yields (6.4). □

References

- [1] R.A. Adams and J.J.F. Fournier. *Sobolev spaces*. Academic Press, 2nd edition, 2003. Pure and Applied Mathematics, Vol. 140.
- [2] Sh. Agmon. The L_p approach to the Dirichlet problem. I. Regularity theorems. *Ann. Scuola Norm. Sup. Pisa (3)*, 13:405–448, 1959.
- [3] J.-P. Aubin and I. Ekeland. *Applied Nonlinear Analysis*. Wiley, 1984.
- [4] J.-P. Aubin and H. Frankowska. *Set-valued analysis*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2009. Reprint of the 1990 edition.
- [5] P. Billingsley. *Convergence of Probability Measures*. Wiley, 1968.
- [6] V. I. Bogachev, N. Krylov, and M. Röckner. Regularity of invariant measures: the case of non-constant diffusion part. *J. Funct. Anal.*, 138(1):223–242, 1996.
- [7] V. I. Bogachev, N. V. Krylov, and M. Röckner. On regularity of transition probabilities and invariant measures of singular diffusions under minimal conditions. *Comm. Partial Differential Equations*, 26(11-12):2037–2080, 2001.
- [8] V. I. Bogachev, N. V. Krylov, and M. Röckner. Elliptic and parabolic equations for measures. *Uspekhi Mat. Nauk*, 64(6(390)):5–116, 2009.
- [9] N. R. Chaganty. Large deviations for joint distributions and statistical applications. *Sankhyā Ser. A*, 59(2):147–166, 1997.
- [10] J. D. Deuschel and D. W. Stroock. *Large Deviations*. American Mathematical Society, second edition, 2001.
- [11] M.D. Donsker and S.R.S. Varadhan. Asymptotic evaluation of certain Markov process expectations for large time I. *Comm. Pure Appl. Math.*, 28:1–47, 1975.
- [12] M.D. Donsker and S.R.S. Varadhan. Asymptotic evaluation of certain Markov process expectations for large time III. *Comm. Pure Appl. Math.*, 29:389–461, 1976.
- [13] R. M. Dudley. *Real analysis and probability*, volume 74 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. Revised reprint of the 1989 original.
- [14] I. Ekeland and R. Temam. *Convex Analysis and Variational Problems*. North Holland, 1976.
- [15] S. N. Ethier and T. G. Kurtz. *Markov Processes. Characterization and Convergence*. Wiley, 1986.

- [16] R. Farwig, H. Kozono, and H. Sohr. On the Helmholtz decomposition in general unbounded domains. *Arch. Math.*, 88(3):239–248, 2007.
- [17] J. Feng and T. G. Kurtz. *Large Deviations for Stochastic Processes*, volume 131 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2006.
- [18] M. Freidlin. The averaging principle and theorems on large deviations. *Uspekhi Mat. Nauk*, 33(5(203)):107–160, 238, 1978.
- [19] M.I. Freidlin and A.D. Wentzell. *Random Perturbations of Dynamical Systems*. Springer, 2nd edition, 1998.
- [20] J. Gärtner. On large deviations from an invariant measure. *Teor. Veroyatnost. i Primenen.*, 22(1):27–42, 1977.
- [21] I. I. Gikhman and A. V. Skorokhod. *Stokhasticheskie differentsialnye uravneniya i ikh prilozheniya*. “Naukova Dumka”, Kiev, 1982.
- [22] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order*, volume 224 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1983.
- [23] R.Z. Has'minskii. *Stochastic Stability of Differential Equations*. Sijthoff & Noordhoff, 1980. (Original title: Ustoicivost' sistem differencial'nyh uravnenii pri sluchainyh vozmuscheniyah ih parametrov, Nauka, Moscow, 1969).
- [24] N. Ikeda and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes*. North Holland, 2nd edition, 1989.
- [25] J. Jacod and A.N. Shiryaev. *Limit Theorems for Stochastic Processes*. Springer, 1987.
- [26] N. V. Krylov. *Controlled diffusion processes*, volume 14 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2009. Translated from the 1977 Russian original by A. B. Aries, Reprint of the 1980 edition.
- [27] A. Kufner, O. John, and S. Fučík. *Function spaces*. Noordhoff International Publishing, Leyden, 1977. Monographs and Textbooks on Mechanics of Solids and Fluids; Mechanics: Analysis.
- [28] R. Liptser. Large deviations for two scaled diffusions. *Probab. Theory Related Fields*, 106(1):71–104, 1996.
- [29] R.Sh. Liptser and A.N. Shiryaev. *Theory of Martingales*. Kluwer, 1989.
- [30] M.N. Malyskin. Subexponential estimates of the rate of convergence to the invariant measure for stochastic differential equations. *Theory Prob. Appl.*, 45(3):466–479, 2000.
- [31] G. Metafune, D. Pallara, and A. Rhandi. Global properties of invariant measures. *J. Funct. Anal.*, 223(2):396–424, 2005.
- [32] Ch.B. Morrey, Jr. *Multiple integrals in the calculus of variations*. Classics in Mathematics. Springer-Verlag, Berlin, 2008. Reprint of the 1966 edition [MR0202511].
- [33] J. Nečas. *Direct methods in the theory of elliptic equations*. Springer-Verlag, 2012.

- [34] J. Neveu. *Mathematical foundations of the calculus of probability*. Translated by Amiel Feinstein. Holden-Day, Inc., San Francisco, Calif.-London-Amsterdam, 1965.
- [35] E. Pardoux and A. Yu. Veretennikov. On the Poisson equation and diffusion approximation. I. *Ann. Probab.*, 29(3):1061–1085, 2001.
- [36] A. Puhalskii. On functional principle of large deviations. In V.Sazonov and T.Sherwashidze, editors, *New Trends in Probability and Statistics*, volume 1, pages 198–218. VSP/Moks’las, 1991.
- [37] A. Puhalskii. Weak convergence theory approach to large deviations. In *Large deviations and applications*, Oberwolfach, 1992.
- [38] A. Puhalskii. On the theory of large deviations. *Theory Probab. Appl.*, 38:490–497, 1993.
- [39] A. Puhalskii. Large deviation analysis of the single server queue. *Queueing Systems Theory Appl.*, 21(1-2):5–66, 1995.
- [40] A. Puhalskii. Large deviations of semimartingales: a maxingale problem approach. I. Limits as solutions to a maxingale problem. *Stoch. Stoch. Rep.*, 61:141–243, 1997.
- [41] A. Puhalskii. *Large Deviations and Idempotent Probability*. Chapman & Hall/CRC, 2001.
- [42] A.A. Puhalskii. Stochastic processes in random graphs. *Ann. Probab.*, 33(1):337–412, 2005.
- [43] A.A. Puhalskii and A.A. Vladimirov. A large deviation principle for join the shortest queue. *Math. Oper. Res.*, 32(3):700–710, 2007.
- [44] M. Röckner and T.S. Zhang. Uniqueness of generalized Schrödinger operators and applications. *J. Funct. Anal.*, 105(1):187–231, 1992.
- [45] M. Röckner and T.S. Zhang. Uniqueness of generalized Schrödinger operators. II. *J. Funct. Anal.*, 119(2):455–467, 1994.
- [46] S. V. Shaposhnikov. On interior estimates for the Sobolev norms of solutions of elliptic equations. *Mat. Zametki*, 83(2):316–320, 2008.
- [47] H. Sohr. *The Navier-Stokes Equations*. Birkhäuser, 2001.
- [48] D. W. Stroock and S. R. S. Varadhan. *Multidimensional Diffusion Processes*. Springer, 1979.
- [49] F. Topsøe. *Topology and Measure*, volume 133 of *Lecture Notes in Mathematics*. Springer, 1970.
- [50] A. Veretennikov. On large deviations in the averaging principle for SDE’s with a “full dependence”, revisited. *Discrete Contin. Dyn. Syst. Ser. B*, 18(2):523–549, 2013.
- [51] A. Yu. Veretennikov. On large deviations for ergodic process empirical measures. In *Topics in nonparametric estimation*, volume 12 of *Adv. Soviet Math.*, pages 125–133. Amer. Math. Soc., Providence, RI, 1992.
- [52] A. Yu. Veretennikov. Large deviations in averaging principle for stochastic differential equation systems (noncompact case). *Stochastics Stochastics Rep.*, 48(1-2):83–96, 1994.

- [53] A. Yu. Veretennikov. On polynomial mixing bounds for stochastic differential equations. *Stoch. Proces. Appl.*, 70:115–127, 1997.
- [54] A. Yu. Veretennikov. Letter to the editors: “On large deviations for stochastic differential equations with small diffusion and averaging”. *Teor. Veroyatnost. i Primenen.*, 43(4):819, 1998.
- [55] A. Yu. Veretennikov. On large deviations for stochastic differential equations with small diffusion and averaging. *Teor. Veroyatnost. i Primenen.*, 43(2):349–351, 1998.
- [56] A. Yu. Veretennikov. On large deviations in the averaging principle for stochastic differential equations with “complete dependence”. *Teor. Veroyatnost. i Primenen.*, 43(4):765–767, 1998.
- [57] A. Yu. Veretennikov. On large deviations in the averaging principle for SDEs with a “full dependence”. *Ann. Probab.*, 27(1):284–296, 1999.
- [58] A. Yu. Veretennikov. On large deviations for SDEs with small diffusion and averaging. *Stochastic Process. Appl.*, 89(1):69–79, 2000.