

Notes on the Level Curves of a Meromorphic Function

Trevor Richards*

Department of Mathematics, University of Florida, Gainesville, United States

November 27, 2024

Abstract

The subject of this paper is the bounded level curves of a meromorphic function f with domain G such that each component of ∂G consists of a level curve of f . (A primary example of such a function being a ratio of finite Blaschke products of different degrees, with domain \mathbb{D} .) We will first prove several facts about a single bounded level curve of a f in isolation from the other level curves of f . We will then study how the level curves of f lie with respect to each other. It is natural to expect that the sets $\{z : |f(z)| = \epsilon\}$ vary continuously as ϵ varies. We will make this notion explicit, and use this continuity to prove several results about the bounded level curves of f . It is well known that if z_0 is a zero or a pole of f , then f is conformally equivalent to the function $z \mapsto z^k$ (for some $k \in \mathbb{Z}$) in a neighborhood of z_0 . We generalize this fact by finding a natural decomposition of G into finitely many sub-regions (also bounded by level curves of f), on each of which f is conformally equivalent to $z \mapsto z^k$ (for some $k \in \mathbb{Z}$). Also included is a new proof, using level curves, of the Gauss–Lucas theorem that the critical points of a polynomial are contained in the convex hull of the polynomial’s zeros.

Keywords: complex analysis; meromorphic functions; level curves; Gauss–Lucas theorem

1 Setting, Background, and Major Results

The study of the level curves of a meromorphic function f (sets of the form $\{z : |f(z)| = \epsilon\}$) or tracts of the function (sets of the form $\{z : |f(z)| < \epsilon\}$) has seemed to cluster around two major classes of questions. We will briefly mention a couple of results from each class to give a taste of the existing research on this subject. The first class of questions has to do with properties of a level curve (or tract) of a single function such as arc length of the level curve, area of the tract, and convexity or star-shapeness of the tract. In 1981, W. K. Hayman and J. M. Wu [7] showed that if $f : \mathbb{D} \rightarrow \mathbb{C}$ is an injective analytic function, then any level set of f has length less than $2 \cdot 10^{35}$. On the other hand, in 1980, P. W. Jones [8] constructed an analytic bounded function $f : \mathbb{D} \rightarrow \mathbb{C}$ each of whose level sets is either empty or of infinite length. Questions about convexity and star-shapeness have mostly had to do with polynomial tracts, and several counter-examples have been given by G. Pirinian and others [5, 6, 9].

The other class of questions has to do with meromorphic functions which share a level curve or a tract. A number of people have given results showing the relationship between two meromorphic functions which share a level curve, starting with G. Valiron [12] in 1937, whose work implied the following result.

Theorem 1.1. *Let f_1 and f_2 be entire functions such that $|f_1| \equiv 1 \equiv |f_2|$ on a simple closed curve Γ . Then there is an entire function h with $|h| \equiv 1$ on Γ , and $f_1 \equiv b_1 \circ h$ and $f_2 \equiv b_2 \circ h$ for some finite Blaschke products b_1 and b_2 .*

A nice summation of this result and following results of the same flavor may be found in a 1983 paper by K. Stephenson and C. Sundberg [11] in which they give the following stronger result for inner functions.

*Email: trevor1r@ufl.edu

Theorem 1.2. *Let ϕ_1 and ϕ_2 be inner functions such that the sets $\{z : |\phi_1(z)| = c\}$ and $\{z : |\phi_2(z)| = c\}$ have a sub-arc in common for some $c \in (0, 1)$. Then $\phi_1 = \lambda\phi_2$ for some unimodular constant λ .*

Since then K. Stephenson [10] has given a more general result which implies much of the earlier work.

In future work we intend to prove a result somewhat in the vein of the results from this second class of questions. Namely we will show that if the configuration of the critical level curves of two finite Blaschke products f and g (that is, the level curves of f and g which contain critical points of the respective functions) are the same (modulo homeomorphism), then f and g are conformally equivalent (that is, there is some conformal $\phi : \mathbb{D} \rightarrow \mathbb{D}$ such that $g \equiv f \circ \phi$ on \mathbb{D}). Some of the motivation behind this paper is to develop the preliminaries necessary for this future work, however the results found here are of independent interest as well. The proofs found in this paper are elementary, using well known results about meromorphic functions such as the Maximum Modulus Theorem, the Open Mapping Theorem, and the Argument Principle. These and the other facts needed may be found in many texts on complex analysis including [2] and [3].

In order to introduce the work in this paper, we make the following standard definition.

Definition: For $D \subset \mathbb{C}$ an open set, and $f : D \rightarrow \bar{\mathbb{C}}$ a meromorphic function, and $\epsilon \in [0, \infty]$, the set $E_{f,\epsilon} := \{z \in D : |f(z)| = \epsilon\}$ is called a level set of f , and a component of $E_{f,\epsilon}$ is called a level curve of f .

We now establish the setting used in Section 2 in order to describe the results found therein. Let $G \subset \mathbb{C}$ be an open set. Let f be a function meromorphic on an open set containing $cl(G)$. Let $\epsilon \in (0, \infty)$ be given, and let $\Lambda \subset G$ be a fixed bounded component of $E_{f,\epsilon}$ which is contained in G . In Section 2, we examine the single level curve Λ . We observe that Λ forms a planar graph in \mathbb{C} , whose vertices are critical points of f , and we determine exactly which planar graphs we might see as bounded level curves of f . As an application of this, we give a new proof using level curves of the Gauss-Lucas theorem that the critical points of a polynomial are contained in the convex hull of the polynomial's zeros. We then give a formula for counting the faces of Λ in terms of the multiplicities of the critical points of f in Λ .

In Section 3 we impose the following further restrictions of G and f .

- $|f| \equiv 1$ on any unbounded component of ∂G . For each bounded component K of ∂G , there is some $\rho \in (0, \infty)$ such that $|f| \equiv \rho$ on K . (That is, each component of ∂G is contained in some level curve of f in G' .)
- If G is unbounded, then $\lim_{z \in G, z \rightarrow \infty} |f(z)| = 1$.
- G does not contain any unbounded level curve of f .
- G is connected and G^c has finitely many bounded components.
- No level curve of f in G intersects ∂G .

It seems reasonable to expect that as the "height" ϵ of f varies continuously, the corresponding level curves of f will in some sense vary continuously as well. Proposition 3.7 makes this notion of continuous variance explicit. Theorem 3.8 shows that if there are any two level curves L_1 and L_2 of f in G , each of which is exterior to the other, then there is a critical point of f in G which is exterior to both L_1 and L_2 . This implies that if we remove from G each level curve of f which contains a critical point, and each zero and pole of f in G , then each component of the remaining set is conformally equivalent to an annulus. In order to explain the final major result of Section 3, we first recall a property of meromorphic functions.

It is well known that if g is a function meromorphic on some region $D \subset \mathbb{C}$, and $z \in D$ is a zero or pole of g , then there is some region $D_z \subset D$ containing z , and ϕ a conformal mapping which maps D_z to a disk centered at 0 such that $f \equiv \phi^n$ on D_z for some $n \in \mathbb{Z} \setminus \{0\}$. Our final major result in Section 3 extends this fact by in a sense decomposing G into finitely many regions on each of which f is conformally equivalent to a pure power of z . That is, if A is one of the regions remaining after removing the level curves of f in G which contain critical points of f , along with the zeros and poles of f in G , as noted above A is conformally equivalent to an annulus. In Theorem 3.11 we find explicitly which annulus A is conformally equivalent to, and construct a conformal mapping ϕ from A to this annulus, such that for some $n \in \mathbb{Z} \setminus \{0\}$, $f \equiv \phi^n$ on A .

2 Level Curves as Planar Graphs

Definition: By a planar graph we mean a finite graph embedded in a plane or sphere, whose edges do not cross. For the purposes of this paper we will include simple closed paths as planar graphs.

In the existing literature on the subject of level curves of a meromorphic function, it is generally assumed without proof that Λ is a planar graph whose vertices are the critical points of f in Λ , and whose edges are paths in Λ with endpoints at the critical points of f . However because part of the subject of this paper is precisely the geometry of Λ , we make that fact explicit in Proposition 2.1, and give an indication as to the nature of the proof. As much of the proof amounts to an exercise in analytic continuation and compactness, we leave out most of the details.

Proposition 2.1. *Λ is a planar graph.*

Proof. Since Λ is bounded and f is non-constant, there are only finitely many critical points of f in Λ . It will turn out that the critical points of f are exactly the intersections of Λ . Choose some $z_0 \in \Lambda$ which is not a critical point of f . We wish to show that either z_0 lies in a path in Λ whose endpoints are critical points of f , or that Λ contains no critical points of f , in which case Λ consists of a simple closed path.

Since $f'(z_0) \neq 0$, there is some open simply connected set D which contains z_0 such that an inverse g of f may be defined on $f(D)$. $f(z_0)$ lies in a circle of radius ϵ . Our strategy is to continue g analytically around that circle as far as possible, and consider the path that this traces out back in the domain of f .

Case 2.1.1. *g may be continued indefinitely around the ϵ -circle in both directions.*

By the compactness of Λ , there are only finitely many points in Λ at which f takes the value $f(z_0)$. Therefore if g is continued analytically along the ϵ -circle in one direction enough times, the path traced out in the domain intersects itself. Let σ denote this path. Because f is invertible at each point in σ we may conclude that σ is simple (as f would not be injective at a crossing point of such a path) and does not contain any critical point of f . Furthermore, we can use the invertibility of f at each point in σ , and the connectedness of Λ , to show that the path σ is all of Λ . Thus Λ is a simple closed path which does not contain any critical points of f .

Case 2.1.2. *g may not be continued indefinitely around the ϵ -circle in at least one direction.*

Again let σ denote the path obtained by continuing g around the ϵ -circle as far as may be in the direction in which it cannot be continued indefinitely. Clearly σ is contained in Λ and does not contain any critical points of f (because f is invertible at each point in σ). One can use the compactness of Λ to show that σ approaches a unique point in Λ , and one can show that this point is a critical point of f (otherwise one could continue g further along the ϵ -circle). g may not be continued indefinitely around the ϵ circle in the opposite direction because this would imply that Λ does not contain a critical point of f (as seen in Case 2.1.1 above). Therefore if we allow σ now to denote the path obtained by continuing g along the ϵ -circle in the opposite direction, we obtain another path from z_0 to a critical point of f . We conclude that z_0 lies in a path in Λ whose end points are critical points of f . Furthermore this path is simple because f is invertible at each point in this path other than the end points.

Finally we may again use the compactness of Λ to show that Λ contains only finitely many edges and vertices, which gives us the desired result. \square

By inspecting the above proof we may make the following observation.

Corollary 2.2. *The critical points of f contained in Λ are the vertices of Λ . If Λ does not contain any critical point of f , then Λ is a simple closed path.*

Due to the nice behavior of rational functions near ∞ , one can easily extend the result of the above proposition to all level curves of a rational function. Since the proof merely requires one to pre-compose the rational function with a Möbius function, we omit it here.

Corollary 2.3. *If $g : \bar{\mathbb{C}} \rightarrow \bar{\mathbb{C}}$ is a rational function, every level curve of g is a planar graph.*

We will now uncover some restrictions on which planar graphs we may see as level curves of our function f . First in Proposition 2.5 we see that Λ must have bounded faces, and that each edge of Λ is adjacent to at least one of its bounded faces. In Proposition 2.6, we see that each vertex of Λ must have evenly many edges of Λ incident to it. First some topology.

Lemma 2.4. *For any disjoint closed sets $X, Y \subset \bar{\mathbb{C}}$, and $x, y \in \bar{\mathbb{C}} \setminus (X \cup Y)$, if x and y are in the same component of X^c and the same component of Y^c , then x and y are in the same component of $(X \cup Y)^c$.*

Proof. Given X, Y, x, y as in the statement of the lemma, suppose by way of contradiction that x and y are in different components of $(X \cup Y)^c$. Assume without loss of generality that $y = \infty$. Let A_1 denote the component of $(X \cup Y)^c$ which contains x , and let B_1 denote the component of $(X \cup Y)^c$ which contains y . Let Z denote the union of all bounded components of A_1^c . Define $A_2 := A_1 \cup Z$. Since B_1 is open and contains ∞ , and A_2 does not intersect B_1 , we may conclude that A_2 is bounded. Therefore A_2 has only a single unbounded component, so A_2 is simply connected. And the boundary of a simply connected set is connected, so $\partial A_2 \subset X \cup Y$ is connected. Because X and Y are disjoint and compact, this implies that ∂A_2 is either contained in X or contained in Y . However this is a contradiction because x and y are in the same component of X^c and in the same component of Y^c . \square

It may easily be seen that this lemma implies that if A is an open connected set, and $X \subset A$ is compact, such that X^c has a single component, then $A \setminus X$ is connected. This fact will be used several times in this paper, and we will cite the above lemma when it is needed. We now proceed to the proposition.

Proposition 2.5. *Λ has one or more bounded faces and a single unbounded face, and each edge of Λ is adjacent to two distinct faces of Λ .*

Proof. Since Λ is bounded, there is a single unbounded face of Λ , so to show that Λ has bounded faces, it suffices to show the second result of the proposition, that each edge of Λ is adjacent to two distinct faces of Λ . Define $E := \{z \in E_{f,\epsilon} \cap G : z \notin \Lambda\}$. Then $cl(E)$ is a closed set contained in $cl(G)$, and $cl(E)$ does not intersect Λ .

Choose some $z_0 \in \Lambda$ which is not a zero of f' . That is, z_0 is a point in one of the edges of Λ . Choose some $\delta > 0$ small enough so that $B_\delta(z_0) \subset G \setminus E$. (So for all $z \in B_\delta(z_0)$, if $|f(z)| = \epsilon$, then $z \in \Lambda$.) By the Open Mapping Theorem, there are points $x, y \in B_\delta(z_0)$ such that $|f(x)| < \epsilon$ and $|f(y)| > \epsilon$. Suppose by way of contradiction that x and y are in the same face of Λ . Lemma 2.4 (with $\Lambda = X$ and $E \cup G^c = Y$) gives that x and y are in the same component of $(\Lambda \cup E \cup G^c)^c$ which is equal to $G \setminus E_{f,\epsilon}$. Since open connected sets in \mathbb{C} are path connected, there is a path $\gamma : [0, 1] \rightarrow G \setminus E_{f,\epsilon}$ from x to y . But f is continuous, and $|f(x)| < \epsilon < |f(y)|$, so there is some $s \in (0, 1)$ such that $|f(\gamma(s))| = \epsilon$, and thus $\gamma(s) \in E_{f,\epsilon}$, and we have our contradiction. So we conclude that x and y are in different faces of Λ , and thus Λ has bounded faces.

Furthermore, since δ was arbitrarily small, and the points x and y we found in $B_\delta(z_0)$ were shown to be in different components of Λ^c , we have that z_0 is in the boundary of more than one component of Λ^c . Since the edge of Λ containing z_0 is locally conformally equivalent to the ϵ -circle, we have that the edge of Λ containing z_0 is in the boundary of at most two faces of Λ , and we are done. \square

One more restriction may be made on which planar graphs we may see as level curves of f . As the method of proof is largely the same as that used in the proof of Proposition 2.1, we will leave out most of the details. First a definition.

Definition: If $z \in G$ is a zero of f' , then let $mult(z)$ denote the multiplicity of z as a zero of f' .

Proposition 2.6. *If z is a vertex of Λ , then the number of edges of Λ which are incident to z is exactly $2(mult(z) + 1)$.*

Proof. As already noted, if $z \in \Lambda$ is not a zero of f' , then z is not a vertex of Λ . Suppose $z \in \Lambda$ is a zero of f' . Then f is precisely $(\text{mult}(z) + 1)$ -to-1 in a neighborhood of z . One can show (using the same tricks found in the proof of Proposition 2.1 of analytic continuation of branches of the inverse of f) that there are precisely $2(\text{mult}(z) + 1)$ edges of Λ incident to z . \square

In future work we will show that for any planar graph which satisfies the restrictions described in the above propositions (namely a planar graph Λ such that each edge is in the boundary of two distinct faces of Λ^c and each vertex of Λ has an evenly many, and four or more, edges incident to it), there is some region G and function f as above such that some level curve of f in G is homeomorphic to the graph Λ . In Figure 1 we have several examples of planar graphs that satisfy the above restrictions, and in Figure 2 we have several examples of planar graphs which do not satisfy these restrictions.




Figure 1: Admissible Graphs




Figure 2: Non Admissible Graphs

Definition: For $K \subset G$, define $B_K := \{z \in K : f'(z) = 0, f(z) \neq 0\}$.

We are now able to count precisely the number of faces of Λ in terms of the critical points of f contained in Λ as follows.

Proposition 2.7. Λ has exactly $\left(\sum_{a \in B_\Lambda} \text{mult}(a)\right) + 1$ bounded faces and one unbounded face.

Proof. Note first that if Λ does not contain any zeros of f' (so B_Λ is empty), then from Corollary 2.2, Λ is a single simple closed path, and thus Λ has two faces, one bounded and one unbounded, and this agrees with the formula in the statement above.

Now suppose that B_Λ is non-empty. Let \mathcal{V} denote the number of points in B_Λ (therefore \mathcal{V} is the number of vertices of Λ). For $a \in B_\Lambda$, f' has a zero with multiplicity $\text{mult}(a)$ at a , so by Proposition 2.6, exactly $2(\text{mult}(a) + 1)$ edges of Λ meet at a . Each line in Λ has two endpoints, so this implies that there are

$$\frac{1}{2} \sum_{a \in B_\Lambda} 2(\text{mult}(a) + 1) = \left(\sum_{a \in B_\Lambda} \text{mult}(a)\right) + \mathcal{V} \text{ edges in } \Lambda.$$

Euler's characteristic formula (see for example [1]) for a connected planar graph states that for any planar graph,

$$\text{number of faces} = \text{number of edges} - \text{number of vertices} + 2.$$

Using this, we obtain that the number of faces of Λ is exactly

$$\left(\left(\sum_{a \in B_\Lambda} \text{mult}(a) \right) + \mathcal{V} \right) - \mathcal{V} + 2 = \left(\sum_{a \in B_\Lambda} \text{mult}(a) \right) + 2.$$

Of these, one is unbounded and the rest are bounded. □

Here are several examples.

Example: Let $f(z) = z^n$ for some $n \in \{1, 2, 3, \dots\}$, and $\epsilon \in (0, \infty)$. Then $E_{f,\epsilon}$ is the circle $\{z \in \mathbb{C} : |z| = \epsilon^{\frac{1}{n}}\}$.

Example: Let $f(z) = z^n - 1$ for some $n \in \{2, 3, \dots\}$. If $\epsilon \in (0, 1)$, then $E_{f,\epsilon}$ has n components, each a simple closed path which contains a single zero of f in its bounded face. If $\epsilon \in (1, \infty)$, then $E_{f,\epsilon}$ consists of a single simple closed path which contains all n zeros of f in its bounded face. Finally, $E_{f,1}$ consists of a single component with a single vertex (at 0), n edges, and n bounded faces, each of which contains a single zero of f . In Figure 3 below we see the example $f(z) = z^5 - 1$. The level sets shown are the sets $E_{f,.5}$, $E_{f,1}$, and $E_{f,1.5}$.




Figure 3: $f(z) = z^5 - 1$

As an application of the above results, we have a new proof of the following theorem of Gauss. First a definition.

Definition: For $z \in G$, define Λ_z to be the component of $E_{f,|f(z)|}$ which contains z . If $z \in \partial G$, let Λ_z denote the component of ∂G which contains z .

Theorem 2.8. *Let p be a polynomial. The critical points of p are contained in the convex hull of the zeros of p .*

Proof. Let $w_1, w_2, \dots, w_n \in \mathbb{C}$ be the zeros of p repeated according to multiplicity. Assume by way of contradiction that there is some critical point z_0 of p which is not in the convex hull of $\{w_i : 1 \leq i \leq n\}$. By pre-composing with a linear map, we may assume that all zeros of p are contained in the disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$, and that $z_0 \in (1, \infty)$. Let G denote one of the bounded faces of Λ_{z_0} which is incident to z_0 , and let $\gamma : [0, 1] \rightarrow \mathbb{C}$ be a parameterization of the boundary of G . Thus, γ is a simple closed path with $\gamma(0) = \gamma(1) = z_0$. The Maximum Modulus Theorem implies that G contains a zero of p , and therefore ∂G intersects the line $L := \{z \in \mathbb{C} : \text{Re}(z) = 1\}$. Define $r_1, r_2 \in (0, 1)$ by $r_1 = \min\{r \in [0, 1] : \gamma(r) \in L\}$ and $r_2 = \max\{r \in [0, 1] : \gamma(r) \in L\}$.

Then $\gamma(0, r_1)$ and $\gamma(r_2, 1)$ are paths from z_0 to L which do not intersect except at z_0 (and possibly in L). Therefore since there are at least two bounded faces of Λ_{z_0} which are incident to z_0 , there are at least four paths in Λ_{z_0} from z_0 to L which do not intersect except at z_0 and in L . It is not hard to see then that there is some $s \in (-1, 1) \setminus \{0\}$ such that two of the paths intersect the set $\{z \in \mathbb{C} : \operatorname{Re}(z) > 1, \operatorname{Im}(z) = s\}$.




Figure 4: Gauss' Theorem

Let z_1, z_2 be distinct points in $\{z \in \mathbb{C} : \operatorname{Re}(z) > 1, \operatorname{Im}(z) = s\}$ which are contained in Λ_{z_0} , with $\operatorname{Re}(z_1) < \operatorname{Re}(z_2)$, as in Figure 4. Then for each zero w_i of p , $|z_1 - w_i| < |z_2 - w_i|$. Therefore

$$|p(z_1)| = \prod_{i=1}^n |z_1 - w_i| < \prod_{i=1}^n |z_2 - w_i| = |p(z_2)|,$$

which is a contradiction because z_1 and z_2 are in the same level curve of p . Thus we conclude that there is no critical point z_0 of p outside of the convex hull of the zeros of p . \square

3 Assorted Properties of Level Curves

3.1 Setting

We now impose the following additional restrictions on the set G and the function f .

- $|f| \equiv 1$ on any unbounded component of ∂G . For each bounded component K of ∂G , there is some $\rho \in (0, \infty)$ such that $|f| \equiv \rho$ on K . (That is, each component of ∂G is contained in some level curve of f in G' .)

Note: If one of the components of G^c is a single point $\{z\}$ for some $z \in \mathbb{C}$ (and thus z is a removable discontinuity of f restricted to G), then we replace G with $G \cup \{z\}$.

- If G is unbounded, then $\lim_{z \in G, z \rightarrow \infty} |f(z)| = 1$.
- G does not contain any unbounded level curve of f .
- G is connected and G^c has finitely many bounded components.
- No level curve of f in G intersects ∂G . That is, for any $z \in G$, $\operatorname{cl}(\lambda_z) \subset G$.

Note: Of particular interest to us for future work is the case $f = \frac{B_1}{B_2}$ where B_1 and B_2 are finite Blaschke products with $\deg(B_1) \neq \deg(B_2)$, with $G = \mathbb{D}$. Note that when B_1 and B_2 are finite degree Blaschke products of the same degree, $f = \frac{B_1}{B_2}$ and $G = \mathbb{D}$ violates the final restriction above. For example, if we set $B_1(z) = \frac{z - \frac{1}{2}}{1 - \frac{1}{2}z}$ and $B_2(z) = \frac{z + \frac{1}{2}}{1 + \frac{1}{2}z}$, and $f = \frac{B_1}{B_2}$ and $G = \mathbb{D}$, this choice of f and G satisfy the first four restrictions above, but the level curve of f which contains 0 consists of the imaginary axis along with the unit circle, and thus the final restriction is violated.

3.2 Properties

In the previous section we examined one level curve of f in isolation from the others. In this section we catalogue some facts about the level curves of f globally in G .

The restriction $\lim_{z \in G, z \rightarrow \infty} |f(z)| = 1$ immediately implies that there are only finitely many zeros and poles of f contained in G . Our first result in this section extends this to the level curves of f associated to values between 0 and ∞ . We also show that there are only finitely many level curves of f in G which contain critical points of f . We will first introduce some vocabulary and define a partial ordering on the set of all level curves of f in G which we will use in the proof.

Definition: A level curve of f which is not a zero or pole of f , and which contains a critical point of f , we will call a critical level curve of f . A level curve of f which is not a critical level curve we call a non-critical level curve of f .

Definition: If Λ_1 and Λ_2 are level curves of f in G then we say $\Lambda_1 \prec \Lambda_2$ if Λ_1 lies in one of the bounded faces of Λ_2 . Let D be an open sub-set of G . If Λ_1 is a critical level curve of f contained in D , then we say that Λ_1 is \prec -maximal with respect to G if there is no other critical level curve Λ_2 of f contained in D such that $\Lambda_1 \prec \Lambda_2$.

Proposition 3.1. G contains only finitely many components of $E_{f,\epsilon}$, and only finitely many critical level curves.

Proof. Suppose by way of contradiction that $\{L_i\}_{i=1}^{\infty}$ is a sequence of distinct components of $E_{f,\epsilon}$ contained in G . Then there is some subsequence $\{L_{i_n}\}_{n=1}^{\infty}$ such that one of the following occurs:

- $L_{i_1} \prec L_{i_2} \prec L_{i_3} \prec \dots$
- $L_{i_1} \succ L_{i_2} \succ L_{i_3} \succ \dots$
- For each $p, q \in \{1, 2, \dots\}$ with $p \neq q$, L_{i_p} is contained in the unbounded face of L_{i_q} .

Case 3.1.1. $L_{i_1} \prec L_{i_2} \prec L_{i_3} \prec \dots$

For each $j \in \{1, 2, \dots\}$, let D_j denote the face of $L_{i_{j+1}}$ which contains L_{i_j} . Let \widetilde{D}_j denote the subset of D_j which is exterior to L_{i_j} . With this definition, we have a sequence of bounded disjoint open connected sets $\{\widetilde{D}_j\}_{j=1}^{\infty}$, such that $\partial\widetilde{D}_j \subset G$ for each j . (One can use Lemma 2.4 to show that each \widetilde{D}_j is connected.)

Moreover, since there are only finitely many bounded components of G^c , all but finitely many of the $\{\widetilde{D}_j\}_{j=1}^{\infty}$ are contained in G .

Thus (by dropping finitely many of the original sequence) we may assume that each of $\{\widetilde{D}_j\}_{j=1}^{\infty}$ is contained entirely in G . However since f is meromorphic on G , and $|f| \equiv \epsilon$ on $\partial\widetilde{D}_j$ for each $j \in \{1, 2, \dots\}$, the Maximum Modulus Theorem implies that f has a zero or pole in each \widetilde{D}_j . Thus we have a contradiction because as noted earlier, f has only finitely many zeros and poles in G .

Case 3.1.2. $L_{i_1} \succ L_{i_2} \succ L_{i_3} \dots$

The exact same argument works as in the previous case with the appropriate changes.

Case 3.1.3. For each $p, q \in \{1, 2, \dots\}$ with $p \neq q$, L_{i_p} is contained in the unbounded face of L_{i_q} .

For each $j \in \{1, 2, \dots\}$, let D_j denote one of the bounded components of $L_{i_j}^c$. Again, by dropping finitely many of the original $\{D_j\}_{j=1}^\infty$, we may assume that each of $\{D_j\}_{j=1}^\infty$ is contained in G , and thus must contain a zero or pole of f , which gives us the desired contradiction.

Thus by cases we have a contradiction of the original assumption, so we conclude that G contains only finitely many level curves of $E_{f,\epsilon}$.

The fact that G contains only finitely many critical level curves follows from an almost identical argument, letting $\{L_i\}_{i=1}^\infty$ be a sequence of critical level curves, and again obtaining a contradiction by using the fact that there are only finitely many zeros and poles of f in G and only finitely many components of G^c . \square

In the next result we show that a closed set contained in the complement of a level curve of f may be separated from that level curve by a different non-critical level curve of f . This begins to show that the level sets of f vary continuously. First a topological fact that we will use several times throughout this paper, the proof of which follows directly from the compactness of $\bar{\mathbb{C}}$.

Fact 3.2. Let $D \subset \bar{\mathbb{C}}$ be connected and open (either in \mathbb{C} or in $\bar{\mathbb{C}}$). Let $K \subset D$ be closed (in \mathbb{C} or in $\bar{\mathbb{C}}$). Then there is a closed connected set $\tilde{K} \subset D$ which contains K . Furthermore if K is bounded, then \tilde{K} may be chosen to be bounded.

Definition: For $D \subset \mathbb{C}$, let $sc(D)$ denote the union of D with each bounded component of D^c . (The "sc" stands for "simply connected.")

Proposition 3.3. Let L denote some level curve of f in G , or some component of ∂G , and let $\eta \in [0, \infty]$ be the value that $|f|$ takes on L . Let $K \subset sc(G)$ denote some closed set contained in a single bounded component of L^c , such that at least one of L or K is bounded. If $z \in G$ is sufficiently close to L and in the same component of L^c as K , the following holds.

- $|f(z)| \neq \eta$.
- Λ_z is a non-critical level curve of f in G .
- K and L are in different faces of Λ_z .
- If L is bounded and K is in the unbounded face of L , then K is in the unbounded face of Λ_z . Otherwise K is in the bounded face of Λ_z .

Proof. Let D denote the intersection of the component of L^c which contains K with $sc(G)$.

Case 3.3.1. L is unbounded.

Since L is unbounded, K must be bounded. Let $K_1 \subset D$ be the set of all points $z \in D$ such that z satisfies one of the following.

- $z \in K$.
- $z \in G$ and $f(z) = 0$ or $f(z) = \infty$.
- $z \in G$ and $|f(z)| = \eta$.
- $z \in G$ and $f'(z) = 0$.
- $z \in G^c$.

K_1 is closed (as it is a finite union of closed sets), and by Proposition 3.1, K_1 is bounded and thus compact. Therefore by Fact 3.2, we may find some compact connected set $\widetilde{K}_1 \subset D$ which contains K_1 . We can then find a compact connected set $K_2 \subset D$ such that \widetilde{K}_1 is contained in the interior of K_2 (for example a finite union of closed balls contained in D). Since \widetilde{K}_1 does not intersect ∂K_2 , $|f|$ does not take the value η on ∂K_2 . Thus since ∂K_2 is compact, if we set $\iota := \inf_{z \in \partial K_2} (||f(z)| - \eta|)$, then $\iota > 0$. We will now use this ι to determine how close the "sufficiently close" from the statement of the proposition is.

Claim 3.3.1. *Some $\delta > 0$ may be found small enough so that if $z \in G$ is within δ of L , then $z \notin K_2$, and $||f(z)| - \eta| \in (0, \iota)$.*

Since each level curve of f in G is bounded, L must be an unbounded component of ∂G , and thus $\eta = 1$. However we will keep writing "η", as we will be using essentially the same argument for the next case, when η may not equal 1. Since K_2 is compact, and L is closed, $\delta > 0$ may be chosen so that if $z \in G$ is within δ of L , then $z \notin K_2$. Since $\lim_{z \in G, z \rightarrow \infty} |f(z)| = \eta$, there is some $R > 0$ such that if $z \in G$ with $|z| > R$, then $||f(z)| - \eta| < \iota$. Furthermore, Proposition 3.1 implies that the set $\{z \in G : |f(z)| = \eta\}$ is bounded, so by choosing R larger if necessary, we may conclude that for $z \in G$ with $|z| > R$, $||f(z)| - \eta| \in (0, \iota)$. On the other hand, by the compactness of $\{z \in L : |z| \leq R\}$, we can find some $\delta > 0$ smaller than before such that if $z \in G$ is within δ of L , and $|z| \leq R$, then $||f(z) - \eta| \in (0, \iota)$, which gives us the result of the claim.

Claim 3.3.2. *With $\delta > 0$ chosen as in the previous claim, if $z \in D$ is less than δ away from L , then Λ_z is a non-critical level curve of f contained in G , such that L is in the unbounded face of Λ_z , and K_2 is in the bounded face of Λ_z , and $|f(z)| \neq \eta$.*

Let $z \in D$ be less than δ away from L . By the previous claim, $||f(z)| - \eta| \in (0, \iota)$ (and thus $|f(z)| \neq \eta$), so by definition of ι , Λ_z does not intersect ∂K_2 . Since Λ_z does intersect K_2^c (namely at z), and Λ_z is connected, we may conclude that Λ_z does not intersect K_2 . And K_2 is connected, so K_2 is entirely contained in one of the faces of Λ_z . Since K_2 contains all critical points of f in D , Λ_z is a non-critical level curve of f , and therefore has only one bounded face. By the Maximum Modulus Theorem this bounded face must contain either a zero or pole of f , or a point in G^c . But each zero and pole of f and point in G^c which is in D is contained in K_2 , so we may conclude that K_2 is contained in the bounded face of Λ_z . Since L is unbounded, L is contained in the unbounded face of Λ_z .

Case 3.3.2. *L is bounded.*

If K is contained in a bounded component of L^c , then the same argument as above works with minor changes, so let us assume that K is contained in the unbounded component of L^c . We use a similar argument, but we want to construct K_2 in a way that will guarantee that when we have found our δ , if $z \in D$ is within δ of L , then K_2 will be contained in the unbounded face of Λ_z . Therefore we define K_1 identically as before except in addition we include ∞ in K_1 . Constructing K_2 as before, we obtain a closed connected set contained in D which contains K_1 in its interior. Let ι and δ be defined as above. The same argument as above with minor changes allows us to conclude if $z \in D$ is within δ of L , then Λ_z is a non-critical level curve of f in G with $|f| \neq \eta$ on Λ_z , and L is in the bounded face of Λ_z , and K_2 (and therefore K) is in the unbounded face of Λ_z . \square

This last proposition hints that in some sense the components of $E_{f,\eta}$ vary continuously as η varies. And indeed this proposition does most of the work of showing this continuous variance, as it is used extensively in the proof of Proposition 3.7, which makes very explicit the continuous variance. First two corollaries.

Definition: For $K \subset G$, define $\Lambda_K := \bigcup_{z \in K} \Lambda_z$.

Corollary 3.4. *Let $K \subset G$ be compact. Then Λ_K is compact.*

Proof. We will first show that Λ_K is closed in \mathbb{C} . Let $z_0 \in \Lambda_K^c$ be given. We will show that there is some $\delta > 0$ such that $B_\delta(z_0) \subset \Lambda_K^c$.

Case 3.4.1. $z_0 \in cl(G)$.

Since $z_0 \notin \Lambda_K$, $\Lambda_{z_0} \cap K = \emptyset$. Thus by Proposition 3.3, there is some non-critical level curve L of f in G such that Λ_{z_0} is contained in one face of L and K is contained in the other face of Λ_{z_0} . Let D_1 denote the face of L which contains z_0 and let D_2 denote the face of L which contains K . For each $z \in K$, $z \in D_2$ and Λ_z is connected and does not intersect L , so Λ_z is contained in D_2 . Therefore $\Lambda_K \subset D_2$. If we choose $\delta > 0$ small enough that $B_\delta(z_0) \subset D_1$, then $B_\delta(z_0) \subset \Lambda_K^c$.

Case 3.4.2. $z_0 \notin cl(G)$.

Since $z_0 \notin cl(G)$, there is some $\delta > 0$ such that $B_\delta(z_0) \subset G^c$, and thus $B_\delta(z_0) \subset \Lambda_K^c$.

We conclude that Λ_K^c is open in \mathbb{C} , and thus Λ_K is closed in \mathbb{C} .

Now we wish to show that Λ_K is bounded. Of course if G is bounded then we are done, so suppose not. Let K_1 denote the union of K with all components of $E_{f,1}$ contained in G . By Proposition 3.1, there are only finitely many components of $E_{f,1}$ contained in G , each of which is compact, so K_1 is compact. Since K_1 is compact, there is a sequence of balls B_1, B_2, \dots, B_N each contained in G such that $K_1 \subset \bigcup_{i=1}^N B_i$. Define

$K_2 := cl\left(\bigcup_{i=1}^N B_i\right)$. K_2 is compact, so ∂K_2 is compact. Therefore we may define $\rho := \min_{z \in \partial K_2} (|f(z)| - 1)$.

Since $\{z \in G : |f(z)| = 1\}$ is contained in the interior of K_2 , there is no point on the boundary of K_2 at which $|f|$ takes the value 1. Therefore we conclude that $\rho > 0$. We will show that Λ_{K_2} is bounded, and thus Λ_K is bounded as well.

We now break Λ_{K_2} into two different sets, each of which we will show is bounded. Define $F_1 := \{z \in K_2 : |f(z)| \in (1 - \rho, 1 + \rho)\}$ and $F_2 := \{z \in K_2 : |f(z)| \notin (1 - \rho, 1 + \rho)\}$. $K_2 = F_1 \cup F_2$, so $\Lambda_{K_2} = \Lambda_{F_1} \cup \Lambda_{F_2}$. For any $z \in F_1$, z is contained in the interior of K_2 , and Λ_z does not intersect ∂K_2 , so $\Lambda_z \subset K_2$. Therefore $\Lambda_{F_1} \subset K_2$, so Λ_{F_1} is bounded. On the other hand, for each $z \in \Lambda_{F_2}$, $|f(z)| \notin (1 - \rho, 1 + \rho)$, and $\lim_{w \in G, w \rightarrow \infty} |f(w)| = 1$, so Λ_{F_2} is bounded.

Therefore we conclude that Λ_{K_2} is bounded, and $K \subset K_2$, so Λ_K is bounded. Finally we conclude that Λ_K is compact. \square

Definition: If D is a face of Λ , say that f is increasing into D if there is some $\iota > 0$ such that $|f| > \epsilon$ on $\{z \in D : d(\{z\}, \Lambda) < \iota\}$, and say that f is decreasing into D if there is some $\iota > 0$ such that $|f| < \epsilon$ on $\{z \in D : d(\{z\}, \Lambda) < \iota\}$.

Corollary 3.5. *Let D be a face of Λ . Then either f is increasing into D or f is decreasing into D .*

Proof. We begin by assuming that D is one of the bounded faces of Λ .

Define A to be the union of the set $\{z \in D \cap G : |f(z)| = \epsilon\}$ with any components of G^c contained in D . If A defined as such is empty, then let A be just some single point in D . A is closed, so by Proposition 3.3, there is some non-critical level curve of f contained in $D \cap G$ which separates Λ from A . Call this level curve L . Let $\eta > 0$ be such that $|f| \equiv \eta$ on L . Since L separates A from Λ , L does not intersect A , so $\eta \neq \epsilon$.

Case 3.5.1. $\eta > \epsilon$.

Let D_1 denote the face of L which contains Λ , and let D_2 denote the face of L which contains A . Define $D' := D_1 \cap D$, the portion of D which is "between" L and Λ . $D \setminus D_1 = cl(D_2)$, so $D \setminus D_1$ is compact and simply connected. We may use Lemma 2.4 to show that $D \setminus cl(D_2) = D \cap D_1$ is connected. Therefore since $|f| \neq \epsilon$ on $D \cap D_1$, and $|f|$ is continuous, either $|f| > \epsilon$ on $D \cap D_1$, or $|f| < \epsilon$ on $D \cap D_1$. But there are points in $\partial(D \cap D_1)$ at which $|f|$ takes values greater than ϵ , namely any point in L , so there must be points in $D \cap D_1$ at which $|f|$ takes values greater than ϵ by the continuity of $|f|$. Thus $|f| > \epsilon$ on $D \cap D_1$. Define $\iota := d(L, \Lambda)$. If $w \in \{z \in D : d(\{z\}, \Lambda) < \iota\}$, then $w \in D \cap D_1$, so $|f(w)| > \epsilon$. Thus f is increasing into D .

Case 3.5.2. $\eta < \epsilon$.

The same argument as above works, with the conclusion that f is decreasing into D .
If D is the unbounded face of Λ , the same argument works with the appropriate minor changes. \square

Now we finally make explicit the notion that the components of $E_{f,\eta}$ "vary continuously in η ". Recall that the Hausdorff distance \check{d} (defined below) may be characterized by saying that for two sets $X, Y \subset \mathbb{C}$ to be close with respect to \check{d} , each $x \in X$ must be close to Y , and each $y \in Y$ must be close to X .

Definition: For $X, Y \subset \mathbb{C}$, define $d_1(X, Y) := \sup_{x \in X} (d(\{x\}, Y))$, and define $d_2(X, Y) := \sup_{y \in Y} (d(X, \{y\}))$. Then the distance function we are interested in is $\check{d}(X, Y) := \max(d_1(X, Y), d_2(X, Y))$. If either X or Y are empty, we define $\check{d}(X, Y) := \infty$.

The following proposition states that if $\eta > 0$ is sufficiently close to ϵ , then some collection of level curves from $E_{f,\eta}$ is close to Λ with respect to \check{d} . First we will list a fact to be used in the proof.

Fact 3.6. *Let $D \subset \mathbb{C}$ be a non-empty set such that ∂D is bounded. Then if $\delta > 0$ is given, there is some finite subset $A \subset D$ such that $\check{d}(A, \partial D) < \delta$.*

This follows from the compactness of ∂D .

Proposition 3.7. *Let $\delta > 0$ be given. Then there is some $\eta \in (0, \epsilon)$ such that for each $\zeta \in (\epsilon - \eta, \epsilon + \eta)$, there is some collection L_1, \dots, L_N of level curves of f contained in G such that $|f| \equiv \zeta$ on each L_i , and $\check{d}\left(\left(\bigcup_{i=1}^N L_i\right), \Lambda\right) < \delta$.*

Proof. Let $\delta > 0$ be given. Let E_1, E_2, \dots, E_N be an enumeration of the edges of Λ , and for each $i \in \{1, 2, \dots, N\}$, let z_i be some fixed interior point of E_i (that is, a point in E_i which is not an endpoint of E_i). Our first goal is to find an $\iota > 0$ small enough so that each of the following hold:

1. For each $i \in \{1, 2, \dots, N\}$, the only faces of Λ that intersect $B_\iota(z_i)$ are the two faces adjacent to E_i .
2. If D is one of the faces of Λ , and $z \in D$ is less than ι away from Λ , then $\check{d}(\Lambda_z, \partial D) < \frac{\delta}{2}$.
3. If D is one of the faces of Λ , and we define $A := \{z \in D : d(\{z\}, \Lambda) < \iota\}$, then either $|f| < \epsilon$ on A or $|f| > \epsilon$ on A .

Suppose an ι may be found which satisfies each of these three items. Our next goal is to find some $\eta \in (0, \epsilon)$ small enough so that if $\zeta \in (\epsilon - \eta, \epsilon + \eta)$, then for each $i \in \{1, 2, \dots, N\}$, there is some point in $B_\iota(z_i)$ at which $|f|$ takes the value ζ . Suppose such an $\eta > 0$ may be found. We now show that the statement of the proposition holds for this δ and η .

Let $\zeta \in (\epsilon - \eta, \epsilon + \eta)$ be given. Of course if $\zeta = \epsilon$, then by putting $N = 1$ and $L_1 = \Lambda$, the statement of the proposition obviously holds, so let us assume that $\zeta \neq \epsilon$. For each $i \in \{1, 2, \dots, N\}$, let w_i be a fixed point in $B_\iota(z_i)$ at which $|f|$ takes the value ζ , which may be found by Item 3 above. Define $\mathcal{L} := \bigcup_{i=1}^N \Lambda_{w_i}$.

Claim 3.7.1. $\check{d}(\mathcal{L}, \Lambda) < \delta$.

Let $x \in \mathcal{L}$ be given. Let $i \in \{1, 2, \dots, N\}$ be such that $x \in \Lambda_{w_i}$. Since $d(\{w_i\}, \Lambda) < \iota$, we have that $\check{d}(\Lambda_{w_i}, \Lambda) < \frac{\delta}{2}$ by Item 2, and thus $d(\{x\}, \Lambda) < \frac{\delta}{2}$. Therefore $d_1(\mathcal{L}, \Lambda) < \delta$.

Let $x \in \Lambda$ be given. Let $i \in \{1, 2, \dots, N\}$ be such that $x \in E_i$. Let D denote the face of Λ which contains w_i . $d(\{z_i\}, \{w_i\}) < \iota$, so by Item 1 above, $E_i \subset \partial D$. And $d(\{w_i\}, \Lambda) < \iota$, so by Item 2 above, $\check{d}(\Lambda_{w_i}, \partial D) < \frac{\delta}{2}$. Since $x \in E_i \subset \partial D$, we conclude that $d(\Lambda_{w_i}, \{x\}) < \frac{\delta}{2}$. Since $x \in \Lambda$ was chosen arbitrarily, we conclude that $d_2(\mathcal{L}, \Lambda) < \delta$.

Finally we conclude that $\check{d}(\mathcal{L}, \Lambda) < \delta$, and we are done, subject to finding the specified ι and η .

Of course if we show that a positive constant may be chosen to satisfy each of the three above items individually, then the minimum of the three constants will have the properties desired in a choice of ι .

We first show that ι may be chosen to satisfy Item 1. Let $i \in \{1, 2, \dots, N\}$ be given. Define E_i' to be the points in E_i which are not endpoints of E_i . Then $\Lambda \setminus E_i'$ is closed, so $r_i := d(\{z_i\}, \Lambda \setminus E_i') > 0$. Now if D is a face of Λ such that $z_i \notin \partial D$ (and thus $\partial D \subset \Lambda \setminus E_i'$), and $w \in D$, then any path from w to z_i intersects ∂D , in particular the straight line path. Therefore

$$d(\{z_i\}, \{w\}) \geq d(\{z_i\}, \partial D) \geq d(\{z_i\}, \Lambda \setminus E_i') = r_i,$$

and therefore we conclude that D does not intersect $B_{r_i}(z_i)$. If we choose $\iota > 0$ smaller than each r_i , this ι satisfies Item 1.

We now show that ι may be chosen to satisfy Item 2. Let D be one of the faces of Λ . By Fact 3.6, there is a finite sequence of points $x_1, x_2, \dots, x_k \in D$ such that $\check{d}\left(\bigcup_{i=1}^k \{x_i\}, \partial D\right) < \frac{\delta}{4}$. Define

$$K := \{x_i\}_{i=1}^k \cup \left\{z \in D : d(\{z\}, \partial D) \geq \frac{\delta}{4}\right\}.$$

By Proposition 3.3, there is some $\iota_D > 0$ such that if $z \in D$ is less than ι_D away from Λ , then Λ_z contains Λ in one face and K in the other. Fix some $w \in D$ less than ι_D away from Λ .

Claim 3.7.2. $\check{d}(\Lambda_w, \partial D) < \frac{\delta}{2}$.

Let $w_0 \in \Lambda_w$ be given. Since Λ_w is contained in $D \setminus K$, we have $\Lambda_w \subset \{z \in D : d(\{z\}, \partial D) < \frac{\delta}{4}\}$. Thus $d(\{w_0\}, \partial D) < \frac{\delta}{4}$, and thus $d_1(\Lambda_w, \partial D) < \frac{\delta}{2}$.

Let $z_0 \in \partial D$ be given. Then for some $i \in \{1, 2, \dots, k\}$, $d(\{x_i\}, \{z_0\}) < \frac{\delta}{4}$. But $x_i \in K$, so x_i and z_0 are in different faces of Λ_w . Since x_i and z_0 are in different faces of Λ_w , the straight line path from z_0 to x_i intersects Λ_w , and thus $d(\Lambda_w, \{z_0\}) \leq d(\{x_i\}, \{z_0\}) < \frac{\delta}{4}$. Thus $d_2(\Lambda_w, \partial D) < \frac{\delta}{2}$.

Therefore we conclude that $\check{d}(\Lambda_w, \partial D) < \frac{\delta}{2}$. Define $\iota := \min(\iota_D : D \text{ is a face of } \Lambda)$. Then this ι satisfies Item 2.

Finally the fact that ι may be chosen to satisfy Item 3 follows directly from Corollary 3.5.

The fact that an $\eta > 0$ may be found with the desired property follows directly from the Open Mapping Theorem. Since the desired $\iota > 0$ and $\eta > 0$ may be found, we are done. \square

The next result is one of the most important results of the paper. This theorem states that if any two level curves of f in G are exterior to each other, then there is a critical level curve of f in G which contains the two level curves in different bounded faces. We will conclude from this that if we remove all critical level curves from G , each component of the remaining set will be conformally equivalent to a disk or an annulus. If, in addition, we remove the zeros and poles of f from G , then each component of the remaining set will be conformally equivalent to an annulus.

Theorem 3.8. *Let each of L_1 and L_2 be a level curve of f contained in G or a bounded components of G^c . Let F_1 denote the unbounded face of L_1 and F_2 the unbounded face of L_2 . If $L_1 \subset F_2$, and $L_2 \subset F_1$, then there is some $w \in G$ which lies in $F_1 \cap F_2$, such that $f'(w) = 0$ and L_1 and L_2 are contained in different bounded faces of Λ_w .*

Proof. Define $\epsilon_1 := |f(L_1)|$, and $\epsilon_2 := |f(L_2)|$.

Case 3.8.1. *Both L_1 and L_2 are level curves of f in G .*

$G \cap F_1 \cap F_2$ is open, and by Lemma 2.4, it is connected. From this it is easy to show that we can find a path $\gamma : [0, 1] \rightarrow G$ such that $\gamma(0) \in L_1$ and $\gamma(1) \in L_2$, and for all $r \in (0, 1)$, $\gamma(r) \in G \cap F_1 \cap F_2$. For $r \in (0, 1)$, define $K_r := \Lambda_{\gamma(r)}$, and define $A \subset (0, 1)$ be the set such that $r \in A$ if and only if K_r contains L_1 in one of its bounded faces L_2 in its unbounded face. Clearly if L is any level curve of f in G such that L_1 and L_2 are in different faces of L , then L intersects the path γ . Thus Proposition 3.3 guarantees that A is non-empty. So if we define $r_1 := \sup\{r \in (0, 1) : r \in A\}$, we have $r_1 \in (0, 1]$.

Claim 3.8.1. $r_1 < 1$.

Proposition 3.3 also implies that we may find some $s \in (0, 1)$ such that K_s contains L_2 in one of its bounded faces and L_1 in its unbounded face. Let D denote the face of K_s which contains L_2 . Since K_s and L_2 are compact, the distance between K_s and L_2 is greater than zero, so because γ is continuous, there is some $\eta > 0$ such that if $r \in (1 - \eta, 1]$, $\gamma(r)$ is contained in D . Therefore for all $r \in (1 - \eta, 1)$, K_r is contained in D , so the bounded faces of K_r are contained in D . Thus for all $r \in (1 - \eta, 1)$, r is not in A . From this we may conclude that $r_1 < 1$.

We will now show that K_{r_1} contains a critical point of f , and that L_1 and L_2 are contained in different bounded faces of K_{r_1} .

Claim 3.8.2. $L_1 \prec K_{r_1}$.

Since $\gamma(r_1) \in F_1 \cap F_2$, $K_{r_1} \subset F_1 \cap F_2$. Suppose by way of contradiction that L_1 is contained in the unbounded face of K_{r_1} . Then by Proposition 3.3, there is some non-critical level curve Λ_1 of f contained in G such that K_{r_1} is contained in the bounded face of Λ_1 and L_1 is contained in the unbounded face of Λ_1 . Then one may easily use the continuity of γ to show there is some $s \in (0, r_1)$ such that for each $r \in (s, r_1]$, K_r does not contain L_1 in any of its bounded faces (since $\gamma(r)$ is in the bounded face of L_1). Therefore for each $r \in (s, r_1]$, $r \notin A$, which is a contradiction of the definition of r_1 . Thus we conclude that $L_1 \prec K_{r_1}$.

Claim 3.8.3. $L_2 \prec K_{r_1}$.

Suppose by way of contradiction that L_2 is in the unbounded face of K_{r_1} . Then Proposition 3.3 gives that there is some non-critical level curve Λ_2 of f in G such that K_{r_1} is in the bounded face of Λ_2 , and L_2 is in the unbounded face of Λ_2 . Thus $\gamma(r_1)$ is in the bounded face of Λ_2 , and $\gamma(1)$ is in the unbounded face of Λ_2 , so by the continuity of γ , there is some $r \in (r_1, 1)$ such that $\gamma(r) \in \Lambda_2$ (and thus $K_r = \Lambda_2$). But L_1 is in one of the bounded faces of K_{r_1} , so L_1 is in the bounded face of Λ_2 as well, and thus $r \in A$. However this is a contradiction of the definition of r_1 . We conclude that L_2 is contained in one of the bounded faces of K_{r_1} . That is, $L_2 \prec K_{r_1}$.

Thus L_1 and L_2 are each contained in a bounded face of K_{r_1} . We now wish to show that L_1 and L_2 are contained in different bounded faces of K_{r_1} .

Claim 3.8.4. L_1 and L_2 are contained in different bounded faces of K_{r_1} .

We use an almost identical argument to that found in Claim 3.8.2. Suppose by way of contradiction that L_1 and L_2 are contained in the same bounded face of K_{r_1} . Let D denote the face of K_{r_1} containing L_1 and L_2 . Again by Proposition 3.3, there is some non-critical level curve Λ_3 of f contained in D such that L_1 and L_2 are in the bounded face of Λ_3 , and K_{r_1} is in the unbounded face of Λ_3 . Therefore $\gamma(0)$ is in the bounded face of Λ_3 and $\gamma(r_1)$ is in the unbounded face of Λ_3 . Thus by the continuity of γ there is some $s \in (0, r_1)$ such that for each $r \in (s, r_1]$, $\gamma(r)$ is in the unbounded face of Λ_3 . Fix for the moment some $r \in (s, r_1]$. K_r is contained in the unbounded face of Λ_3 , and L_1 and L_2 are both contained in the bounded face of Λ_3 . Therefore since K_r does not intersect Λ_3 , L_1 is in the bounded face of K_r if and only if L_2 is as well. Thus $r \notin A$ for each $r \in (s, r_1]$, which contradicts the definition of r_1 .

We conclude that L_1 and L_2 are contained in different bounded faces K_{r_1} , which implies that K_{r_1} contains a zero of f' by Corollary 2.2, and we are done.

Case 3.8.2. *At least one of L_1 and L_2 are bounded components of ∂G .*

Suppose that L_1 is a bounded component of G^c . L_2 is compact, so by Proposition 3.3, we may find a non-critical level curve L_1' of f in G such that L_1 is contained in the bounded face of L_1' and L_2 is contained in the unbounded face of L_1' . If L_2 is a bounded component of G^c , the same use of Proposition 3.3 gives us a level curve L_2' of f in G such that L_2 is contained in the bounded face of L_2' and L_1' (and thus L_1) is contained in the unbounded face of L_2' . Then we can use the preceding case with L_1' and L_2' , which then gives us the desired result for L_1 and L_2 . \square

Let us now extend the \prec -ordering to the bounded components of ∂G as follows.

Definition: If $K \subset G$ is a bounded component of ∂G , and L is a level curve of f in G such that K is contained in one of the bounded faces of L , then we write $K \prec L$.

Next we have two corollaries to the previous theorem guaranteeing the existence of a unique \prec -maximal critical level curve of f in certain regions of G , but first a definition.

Definition: Define

$$\mathcal{B} := \{z \in G : f'(z) = 0 \text{ or } f(z) = 0 \text{ or } f(z) = \infty\},$$

and define

$$\mathcal{C} := \left(\bigcup_{z \in \mathcal{B}} \Lambda_z \right) \cup \{z \in \partial G : z \text{ is in a bounded component of } \partial G\}.$$

Recall that since $\lim_{z \in G, z \rightarrow \infty} f(z) = 1$, f has only finitely many zeros and poles in G , and G^c was stipulated to have only finitely many bounded components, so Proposition 3.1 immediately implies that \mathcal{B} is finite, and thus that \mathcal{C} has finitely many components.

Corollary 3.9. *There is a unique \prec -maximal component of \mathcal{C} .*

Proof. By the Maximum Modulus Theorem, \mathcal{C} is non-empty. Since \mathcal{C} has finitely many components, it must have at least one \prec -maximal component. Suppose by way of contradiction that there are two distinct components L_1 and L_2 of \mathcal{C} which are both \prec -maximal. Then each is in the unbounded component of the other, so Theorem 3.8 guarantees that there is some other critical level curve $\Lambda' \subset \mathcal{C}$ such that $L_i \prec \Lambda'$ for $i = 1, 2$. But this is a contradiction of the \prec -maximality of L_1 and L_2 . \square

More generally we have the following corollary.

Corollary 3.10. *If D is a bounded face of Λ then there is a unique component of \mathcal{C} which is \prec -maximal with respect to D .*

Proof. Just replace G with D , and then the desired result is just the result from Corollary 3.9 in the new setting. \square

In light of our following final result, the name "critical level curve" has new meaning. The following theorem says that on any component of $G \setminus \mathcal{C}$, f is very simple, being conformally equivalent to the function $z \mapsto z^n$ for some $n \in \{1, 2, \dots\}$. First a bit of notation.

Definition: For any path γ , let $-\gamma$ denote the same curve γ equipped with the opposite orientation.

Theorem 3.11. *Let D be a component of $G \setminus \mathcal{C}$. Then the following hold.*

1. D is conformally equivalent to some annulus A .
2. Let E_1 denote the inner boundary of D , and let E_2 denote the outer boundary of D . Then there is some $\epsilon_1, \epsilon_2 \in [0, \infty]$ such that $\epsilon_1 \neq \epsilon_2$, and $|f| \equiv \epsilon_1$ on E_1 , and $|f| \equiv \epsilon_2$ on E_2 .
3. Let $i_1, i_2 \in \{1, 2\}$ be chosen so that $\epsilon_{i_1} < \epsilon_{i_2}$. Then there is some $N \in \{1, 2, \dots\}$ such that $A = \text{ann}(0; \epsilon_{i_1}^{\frac{1}{N}}, \epsilon_{i_2}^{\frac{1}{N}})$, and some conformal mapping $\phi : D \rightarrow \text{ann}(0; \epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$ such that $f \equiv \phi^M$ on D , where $M = \pm N$.
4. The conformal map ϕ described in Item 3 extends continuously to E_2 and to all points in E_1 which are not zeros of f' . If $z \in E_1$ is a zero of f' , and $\gamma : [0, 1] \rightarrow G$ is a path such that $\gamma([0, 1)) \subset D$, and $\gamma(1) = z$, then $\lim_{r \rightarrow 1} \gamma(r)$ exists.

Proof. Let D be some component of $G \setminus \mathcal{C}$. Since D does not contain a zero or pole of f , the Maximum Modulus Theorem implies that D^c must have at least one bounded component. Suppose by way of contradiction that D^c has two distinct bounded components. Replacing G with D in the statement of Theorem 3.8, we may conclude that D contains a zero of f' , which is a contradiction because all zeros of f' in G are contained in \mathcal{C} . We conclude that D^c has exactly one bounded component. Thus D is conformally equivalent to an annulus (see for example [4]). Let E_1 denote the interior boundary of D and E_2 denote the exterior boundary of D . Each component of the boundary of D is contained in a level curve of f or a component of ∂G . Therefore we may define $\epsilon_1 \in [0, \infty]$ to be the value of $|f|$ on E_1 and $\epsilon_2 \in [0, \infty]$ to be the value of $|f|$ on E_2 . By the Maximum Modulus Theorem since D does not contain a zero or pole of f and $D \subset G$, we may conclude that $\epsilon_1 \neq \epsilon_2$. Assume throughout that $\epsilon_1 < \epsilon_2$, otherwise make the appropriate minor changes.

Claim 3.11.1. *There is some $N \in \{\pm 1, \pm 2, \dots\}$ such that for any $z \in D$, the change in $\arg(f)$ as Λ_z is traversed in the positive direction is exactly $2\pi N$.*

Let $z_1, z_2 \in D$ be given. Since D contains no critical points of f , Λ_{z_1} and Λ_{z_2} are non-critical level curves of f , and Theorem 3.8 implies that either $\Lambda_{z_1} \prec \Lambda_{z_2}$ or $\Lambda_{z_2} \prec \Lambda_{z_1}$. Rename z_1 and z_2 if necessary so that $\Lambda_{z_1} \prec \Lambda_{z_2}$. By the Maximum Modulus Theorem, the bounded face of Λ_{z_1} contains some point in \mathcal{C} . However D contains no points of \mathcal{C} , so E_1 is contained in the bounded face of Λ_{z_1} . Let Figure 5 represent this choice of D , z_1 , and z_2 . Let σ denote the path in Figure 6.




Figure 5: D , z_1 , and z_2 .




Figure 6: Definition of σ .

Since σ may be retracted to a point in D , the total change in $\arg(f)$ along σ is zero. Therefore the sum of the changes in $\arg(f)$ along Λ_{z_2} and $-\Lambda_{z_1}$ equals zero. Therefore the change in $\arg(f)$ along Λ_{z_1} is the same as the change in $\arg(f)$ along Λ_{z_2} . We conclude that the change in $\arg(f)$ along Λ_z is independent of the choice of $z \in D$. Let β denote this common number. It is well known (see for example [3]) that if γ is any closed path in \mathbb{C} and g is a function analytic on γ with no zeros on γ , the change in $\arg(g)$ along γ is well defined and there is some $n \in \mathbb{Z}$ such that the change in $\arg(g)$ along γ is $2\pi n$. Thus we may choose $N \in \mathbb{Z}$ such that $\beta = 2\pi N$. By the same argument as above, the changes in $\arg(f)$ along E_1 and E_2 are both $2\pi N$ as well.

Furthermore since D does not contain any zero of f' , f is injective at each point in D , so if L is some level curve of f in D , $\arg(f)$ is either strictly increasing or strictly decreasing as L is traversed in the positive direction. Therefore $N \neq 0$, so $N \in \{\pm 1, \pm 2, \dots\}$.

Let us now adopt the convention that unless otherwise specified, any time we are referring to a level curve of f in D as a path, we are traversing that level curve in the direction in which $\arg(f)$ is increasing. By a similar argument to the one in the claim above one may show that this is either the positive orientation for all level curves of f in D or the negative orientation for all level curves of f in D . Thus with this assumed orientation we may say that the change in $\arg(f)$ along any level curve L in D is $2\pi N$ for some $N \in \{1, 2, \dots\}$ which is independent of L .

Claim 3.11.2. *Let γ be a closed path in D . Then the change in $\arg(f)$ along γ is an integer multiple of $2\pi N$.*

Let k denote the number of times γ winds around E_1 . Then since D contains no zeros or poles of f , the Argument Principle implies that the change in $\arg(f)$ along γ is k times the change in $\arg(f)$ along E_1 . Thus the change in $\arg(f)$ along γ is $k2\pi N$.

We now wish to define the conformal mapping described in the statement of the theorem. Fix some z_0 in D at which f takes a positive real value. (To see that such a point z_0 exists, observe that for any $z \in D$, the change in $\arg(f)$ along Λ_z is $2\pi N$, so there are N different points in Λ_z at which f takes positive real values.) We wish to define a map $\phi : D \rightarrow \mathbb{C}$ which we will show is a conformal map with $\phi(D) = \text{ann}(0; \epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$. For $w \in D$, let $\gamma : [0, 1] \rightarrow D$ be a path such that $\gamma(0) = z_0$ and $\gamma(1) = w$. Define α to be the change in $\arg(f)$ along γ . Then define $\phi(w) := |f(w)|^{\frac{1}{N}} e^{i\frac{\alpha}{N}}$. This may be shown to be the value at w of analytic continuation of $f^{\frac{1}{N}}$ along the path γ , and thus (if well defined) is analytic in a neighborhood of w . Therefore if ϕ defined as such is well defined, then ϕ is analytic on D . Let $w \in D$ be given, and let γ_1 and γ_2 be paths in D from z_0 to w . Let $\phi_1(w)$ denote the value of ϕ at w obtained using $\gamma = \gamma_1$ and let $\phi_2(w)$ denote the value of ϕ at w obtained by using $\gamma = \gamma_2$. Let α_1 denote the change in $\arg(f)$ along γ_1 and let α_2 denote the change in $\arg(f)$ along γ_2 . Let γ_3 denote the path in D obtained by traversing γ_1 and then traversing $-\gamma_2$. γ_3 is a closed path in D , so by the claim above, the change in $\arg(f)$ along γ_3 (which is the change in $\arg(f)$ along γ_1 minus the change in $\arg(f)$ along γ_2) is an integer multiple of $2\pi N$. Thus $\alpha_1 = \alpha_2 + k2\pi N$ for some $k \in \mathbb{Z}$. Thus

$$\phi_1(w) = |f(w)|^{\frac{1}{N}} e^{i\frac{\alpha_1}{N}} = |f(w)|^{\frac{1}{N}} e^{i\frac{\alpha_2 + k2\pi N}{N}} = |f(w)|^{\frac{1}{N}} e^{i\frac{\alpha_2}{N}} e^{i\frac{k2\pi N}{N}} = \phi_2(w).$$

Therefore whether we define $\phi(w)$ using γ_1 or using γ_2 we obtain the same value. (Note that we are essentially just showing that we may take an N^{th} root of f on D .)

Claim 3.11.3. $\phi(D) = \text{ann}(0; \epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$.

Note that by the Maximum Modulus Theorem, for each $z \in D$, since $D \subset G$ and D does not contain any zero or pole of f , $|f(z)| \in (\epsilon_1, \epsilon_2)$, and thus $|\phi(z)| \in (\epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$. Therefore $\phi(D) \subset \text{ann}(0; \epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$.

Let $\xi \in \text{ann}(0; \epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$ be given. Choose some $x \in D$ such that $|f(x)|^{\frac{1}{N}} = |\xi|$. Such an x exists because $|f| \equiv \epsilon_1$ on E_1 and $|f| \equiv \epsilon_2$ on E_2 , and $|f|$ is continuous. Let $\rho \in [0, 2\pi)$ be such that $\arg(\xi) = \arg(\phi(x)) + \rho$. (And thus $\phi(x)e^{i\rho} = \xi$.) Since the change in $\arg(f)$ along Λ_x is $2\pi N$ with $N \neq 0$, there is some point x' in Λ_x such that if Λ_x is traversed from x to x' , then the change in $\arg(f)$ along this portion of Λ_x is exactly ρN . One can easily then show that $\phi(x') = \xi$.

Therefore we conclude that $\phi(D) = \text{ann}(0; \epsilon_1^{\frac{1}{N}}, \epsilon_2^{\frac{1}{N}})$.

Claim 3.11.4. ϕ is injective.

Let $w_1, w_2 \in D$ be given such that $\phi(w_1) = \phi(w_2)$. From the definition of ϕ , $|f(w_1)| = |f(w_2)|$. Therefore w_1 and w_2 are in the same level curve of f . The Maximum Modulus Theorem and Theorem 3.8 together imply that there is only one level curve of f in D on which $|f|$ takes the value $|f(w_1)|$. Thus we conclude that there is some level curve L of f in D which contains both w_1 and w_2 . Let γ_1 be a path in D from z_0 to w_1 . Let γ_2 be a path obtained by traversing L in the direction of increase of $\arg(f)$ from w_1 to w_2 . (If $w_1 = w_2$ let γ_2 be constant.) Let γ_3 denote the path from z_0 to w_2 obtained by first traversing γ_1 and then traversing γ_2 . Let α_i denote the change in $\arg(f)$ along γ_i for $i \in \{1, 2\}$. Since $\arg(f)$ is strictly increasing as γ_2 is traversed, and the total change in $\arg(f)$ as all of L is traversed is $2\pi N$, we conclude that $\alpha_2 \in [0, 2\pi N)$. Since $\phi(w_1) = \phi(w_2)$, we have the following equation, calculating $\phi(w_1)$ using γ_1 , and calculating $\phi(w_2)$ using γ_3 .

$$|f(w_1)|^{\frac{1}{N}} e^{i\frac{\alpha_1}{N}} = \phi(w_1) = \phi(w_2) = |f(w_2)|^{\frac{1}{N}} e^{i\frac{\alpha_1 + \alpha_2}{N}} = |f(w_1)|^{\frac{1}{N}} e^{i\frac{\alpha_1}{N}} e^{i\frac{\alpha_2}{N}}.$$

Dividing both sides by $|f(w_1)|^{\frac{1}{N}} e^{i\frac{\alpha_1}{N}}$, we obtain $1 = e^{i\frac{\alpha_2}{N}}$. Thus α_2 is an integer multiple of $2\pi N$. Since $\alpha_2 \in [0, 2\pi N)$, we conclude that $\alpha_2 = 0$, and thus $w_1 = w_2$.

It remains to show that ϕ extends continuously all points in the boundary of D except possibly the zeros of f' in E_1 . To see this we just observe that the definition of ϕ may be defined in the same way as above for all points in ∂D . This extension is well defined by the same argument used above except at the zeros of f' in E_1 . If z is one of the zeros of f' in E_1 , then for any path $\gamma : [0, 1] \rightarrow G$ such that $\gamma([0, 1)) \subset D$, and $\gamma(1) = z$, it may be shown using the continuity of f that $\lim_{r \rightarrow 1^-} \phi(\gamma(r))$ exists. \square

References

- [1] P. S. Alexandroff. *Combinatorial topology*. Dover, New York, 1998.
- [2] L. Alfors. *Complex analysis: an introduction to the theory of analytic functions of one complex variable*. McGraw-Hill, New York, 1966.
- [3] J. Conway. *Functions of one complex variable I*. Springer, New York, 2 edition, 1978.
- [4] J. Conway. *Functions of one complex variable II*. Springer, New York, 1995.
- [5] P. Erdős, F. Herzog, and G. Piranian. Metric properties of polynomials. *J. Analyse Math.*, 6:125–148, 1958.
- [6] A. W. Goodman. On the convexity of the level curves of a polynomial. *Proc. Amer. Math. Soc.*, 17:358–361, 1966.
- [7] W. K. Hayman and J. M. G. Wu. Level sets of univalent functions. *Comment. Math. Helv.*, 3:366–403, 1981.
- [8] P. W. Jones. Bounded holomorphic functions with all level sets of infinite length. *Mich. Math. J.*, pages 75–80, 1980.
- [9] G. Piranian. The shape of level curves. *Proc. Amer. Math. Soc.*, 17:1276–1279, 1966.
- [10] K. Stephenson. Analytic functions sharing level curves and tracts. *Ann. of Math. (2)*, 123:107–144, 1986.
- [11] K. Stephenson and C. Sundberg. Level curves of inner functions. *Proc. London Math. Soc. (3)*, 51:77–94, 1985.
- [12] M. G. Valiron. Sur les courbes de module constant des fonctions entieres. *C. R. Acad. Sci. Paris*, 204:402–404, 1937.