

Countable Partially Exchangeable Mixtures

Cecilia Prosdocimi ^{*} Lorenzo Finesso [†]

October 15, 2018

Abstract

Partially exchangeable sequences representable as mixtures of Markov chains are completely specified by de Finetti's mixing measure. The paper characterizes, in terms of a subclass of hidden Markov models, the partially exchangeable sequences with mixing measure concentrated on a countable set, for sequences of random variables both on a discrete and a Polish space.

Keywords— exchangeability, partial exchangeability, Markov exchangeability, countable mixtures of Markov chains, hidden Markov model, mixing measure

1 Introduction

In the Hewitt-Savage generalization of de Finetti's theorem, the distributions of exchangeable sequences of Polish valued random variables are shown to be in one to one correspondence with distributions of mixtures of i.i.d. sequences, and ultimately with the mixing measure defining the mixture. The mixing measure thus acts as a model of the exchangeable sequence and its properties shed light on the random mechanism generating the sequence. In this regard [5], using the connection with the Markov moment problem,

^{*}Dipartimento di Economia e Finanza, Università LUISS, viale Romania 32, 00197 Roma, Italy, prosdocimi.cecilia@gmail.com

[†]Istituto di Elettronica e di Ingegneria dell'Informazione e delle Telecomunicazioni, Consiglio Nazionale delle Ricerche, via Gradenigo 6/a, Padova, Italy, lorenzo.finesso@ieiit.cnr.it

characterizes the subclass of exchangeable sequences of discrete valued random variables whose mixing measures are absolutely continuous, and have densities in L^p . It is also of interest to characterize the subclass for which the mixing measure is discrete (*i.e.* concentrated on a countable set). A contribution in this direction has been given by [3], where it is proved that an exchangeable sequence of discrete valued random variables has de Finetti mixing measure concentrated on a countable set if and only if it is a hidden Markov model (HMM), see Definition 2.8 for the precise notion.

The more general class of partially exchangeable sequences is in one to one correspondence with mixtures of Markov chains. As noted in [5], the results on the regularity of the mixing measure carry to partially exchangeable sequences. On the other hand, to the best of our knowledge, no results have been reported concerning partially exchangeable sequences with discrete mixing measures.

The goal of the present paper is to characterize, in the spirit of [3], countable mixtures of Markov chains. Our results hold for sequences of both discrete and Polish space valued random variables. This has required the development of a few special results for HMMs, previously not available in the literature. Of independent interest are Propositions 3.1 and 4.1 on the rows of the array of the successors of an HMM, and most of the Appendix, on properties of sequences of stopping times with respect to the filtration generated by an HMM and its underlying Markov chain. For sequences of Polish valued random variables it has also been necessary to first extend [3] to show the equivalence between exchangeable HMMs and countable mixtures of i.i.d. sequences.

In Section 2 we review the basic definitions in the setup most convenient for our purpose. The reader should be aware of the fact that slightly different notions of partial exchangeability coexist in the literature for sequences of discrete valued random variables. We recall the original definition, introduced in [2] and elaborated in [8]. The latter paper clarifies the relationship with the alternative definition given in [4]. Section 3 deals with sequences of discrete valued random variables, Section 4 with sequences of Polish valued random variables. In Section 3.1 we constructively prove Proposition 3.1, which is instrumental in the balance of the paper. In Section 3.2 we characterize countable mixtures of Markov chains taking values on a discrete space. In Section 4 we extend the result of [3] to exchangeable sequences of Polish valued random variables. This allows us to prove Theorem 4.3, the main

result of the paper. Section 5 contains final remarks and hints at possible extensions. In the Appendix are collected results on sequences of stopping times for HMMs, unavailable elsewhere in the literature, and a technical result for a special class of HMMs representable as mixtures of i.i.d. sequences.

2 Preliminaries

$Y = (Y_n)_{n \geq 0}$ denotes a sequence of random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, taking values in a Polish space S endowed with the Borel σ -field \mathcal{S} . The generic element of S is denoted y , and $y_0^N = y_0 y_1 \dots y_N$ is an element (a string) of the $N + 1$ -th fold Cartesian product S^{N+1} , likewise $(Y_0^N = y_0^N)$ is the event $(Y_0 = y_0, Y_1 = y_1, \dots, Y_N = y_N)$.

Exchangeable and partially exchangeable sequences. The notions of exchangeability and partial exchangeability both originate in the work of de Finetti. The former notion is well established in the literature, the latter has been presented in various disguises. For ease of reference we give below the definitions used in the paper.

Definition 2.1. *[[9], page 24] A sequence of random variables (Y_n) on (S, \mathcal{S}) is exchangeable if, for any $N \in \mathbb{N}$, and distinct natural numbers m_1, \dots, m_N ,*

$$(Y_{m_1}, \dots, Y_{m_N}) \stackrel{d}{=} (Y_1, \dots, Y_N),$$

where $\stackrel{d}{=}$ denotes equality in distribution.

We adopt the definition of partial exchangeability given in [8], close to de Finetti's original [2]. The relation with the definition of partial exchangeability given in [4] is clarified in [8]. The definition requires the introduction of the *successors array* V of a given random sequence (Y_n) on S , and the extension of S to $S^* = S \cup \{\partial\}$, where $\partial \notin S$ is a fictitious state.

If S is discrete (as in Section 3 of the paper) $V = (V_{y,n})_{y \in S, n \geq 1}$ is defined setting $V_{y,n}$ equal to the value of Y immediately following the n -th visit to y of Y . If Y visits y only $m < \infty$ time, to avoid rows of V of finite length, one assigns $V_{y,n} = \partial$ for all $n > m$.

If S is uncountable (as in Section 4 of the paper) let $\mathcal{E} = \{E_j\}_{j \geq 0}$ be a fixed countable partition of S^* , with $E_0 = \{\partial\}$. The matrix $V = (V_{j,n})_{j,n \geq 1}$ is defined setting $V_{j,n}$ equal to the value of Y immediately following the n -th visit to E_j . As in the discrete case, if Y visits E_j only $m < \infty$ times, set

$V_{j,n} = \partial$ for all $n > m$. In the uncountable case V depends on the partition \mathcal{E} .

Definition 2.2. *[[8]] A sequence of random variables (Y_n) on (S, \mathcal{S}) is partially exchangeable if its successors array V is distributionally invariant under finite, not necessarily identical, permutations within each of its rows.*

Note that, if (Y_n) is partially exchangeable, the rows of its successors array V are exchangeable.

Mixtures of i.i.d. sequences and of Markov chains. The set of all probability measures on (S, \mathcal{S}) is denoted $\mathcal{M}_{\mathcal{S}}$ and it is equipped with the σ -field generated by the maps $p \mapsto p(A)$, varying p in $\mathcal{M}_{\mathcal{S}}$ and A in \mathcal{S} .

Definition 2.3. *(Y_n) is a mixture of i.i.d. sequences if there exists a random probability measure \tilde{p} on $\mathcal{M}_{\mathcal{S}}$ such that, for any $N \in \mathbb{N}$ and any $A_0, \dots, A_N \in \mathcal{S}$,*

$$\mathbb{P}(Y_0 \in A_0, \dots, Y_N \in A_N \mid \tilde{p}) = \tilde{p}(A_0) \dots \tilde{p}(A_N), \quad \mathbb{P} - a.s. \quad (1)$$

Definition 2.4. *A mixture of i.i.d. sequences is countable (finite) if the random probability measure \tilde{p} is concentrated on a countable (finite) subset of $\mathcal{M}_{\mathcal{S}}$.*

If (Y_n) is a countable (finite) mixture of i.i.d. sequences let $(p_h(\cdot))_{h \in H}$, where H is countable (finite), be the set of measures on which \tilde{p} is concentrated then, integrating Equation (1) over Ω , one has

$$\mathbb{P}(Y_0 \in A_0, \dots, Y_N \in A_N) = \sum_{h \in H} \mu_h p_h(A_0) \dots p_h(A_N), \quad (2)$$

where $\mu_h := \mathbb{P}(\tilde{p} = p_h) > 0$ and $\sum_{h \in H} \mu_h = 1$.

A time homogenous Markov chain with values in S^* is characterized by a transition kernel $k : S^* \times \mathcal{S}^* \rightarrow [0, 1]$. For the purposes of the paper it is enough to consider Markov chains whose transition kernels are constant, with respect to the first variable, on the elements of a fixed partition $\mathcal{E} = (E_j)_{j \geq 0}$ of S . Kernels k in this subclass can be represented in terms of the simpler kernels in the class T^* made up of all kernels $t : \mathbb{N}_0 \times \mathcal{S}^* \rightarrow [0, 1]$, where $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For a given $t \in T^*$ one defines $k = k_t$ as follows

$$k_t(y, A) := \sum_{j \geq 0} \mathbb{I}_{E_j}(y) t(j, A), \quad (3)$$

where $\mathbb{I}_{E_j}(\cdot)$ is the indicator function of E_j . The reader is referred to [8] for a more detailed discussion. We are ready to give the following

Definition 2.5. *Let (Y_n) be a sequence of random variables with $\mathbb{P}(Y_0 = y_0) = 1$, for some $y_0 \in E_1$, then (Y_n) is a mixture of homogeneous Markov chains if there exists a random kernel \tilde{t} on T^* such that, for any $N \in \mathbb{N}$, and $A_1, \dots, A_N \in \mathcal{S}^*$,*

$$\mathbb{P}(Y_1 \in A_1, \dots, Y_N \in A_N \mid \tilde{t}) = \int_{A_1} \cdots \int_{A_N} k_{\tilde{t}}(y_0, dy_1) \cdots k_{\tilde{t}}(y_{N-1}, dy_N) \quad \mathbb{P}\text{-a.s.} \quad (4)$$

By the definition of k_t in (3), Equation (4) gives

$$\mathbb{P}(Y_1 \in A_1, \dots, Y_N \in A_N \mid \tilde{t}) = \sum_{j_1^N \in (\mathbb{N}_0)^N} \int_{A_1 \cap E_{j_1}} \cdots \int_{A_N \cap E_{j_N}} \tilde{t}(1, dy_1) \cdots \tilde{t}(j_{N-1}, dy_N) \quad \mathbb{P}\text{-a.s.} \quad (5)$$

Definition 2.6. *A mixture of Markov chains is countable (finite) if the random kernel \tilde{t} is countably (finitely) valued.*

For a countable (finite) mixture of Markov chains, Equation (4) reads, after integration over Ω ,

$$\begin{aligned} & \mathbb{P}(Y_1 \in A_1, \dots, Y_N \in A_N) \\ &= \sum_{h \in H} \mu_h \int_{A_1} \cdots \int_{A_N} k_{t_h}(y_0, dy_1) \cdots k_{t_h}(y_{N-1}, dy_N) \end{aligned} \quad (6)$$

where $(t_h)_{h \in H}$ are the kernels on which \tilde{t} is concentrated, $\mu_h := \mathbb{P}(\tilde{t} = t_h)$, and H is a countable (finite) set. Finally we can write the finite distributions of a countable mixtures of Markov chains as

$$\begin{aligned} & \mathbb{P}(Y_1 \in A_1, \dots, Y_N \in A_N) \\ &= \sum_{h \in H} \mu_h \sum_{j_1^N \in (\mathbb{N}_0)^N} \int_{A_1 \cap E_{j_1}} \cdots \int_{A_N \cap E_{j_N}} t_h(1, dy_1) \cdots t_h(j_{N-1}, dy_N). \end{aligned} \quad (7)$$

Remark 2.7. *To define mixtures of Markov chains when S^* is discrete, less technicalities are needed since transition kernels reduce to transition matrices. When S^* is discrete Equation (4) reduces to*

$$\mathbb{P}(Y_1^N = y_1^N \mid \tilde{P}) = \tilde{P}_{y_0 y_1} \tilde{P}_{y_1 y_2} \cdots \tilde{P}_{y_{N-1} y_N} \quad \mathbb{P}\text{-a.s.}, \quad (8)$$

where $\tilde{P} = (\tilde{P}_{i,j})_{i,j \in S^*}$ is a random matrix varying on the set \mathcal{P}^* of transition matrices, and $P(Y_0 = y_0) = 1$ for some $y_0 \in S$. The analog of Equation (6) is

$$\mathbb{P}(Y_1^N = y_1^N) = \sum_{h \in H} \mu_h P_{y_0 y_1}^h \cdots P_{y_{N-1} y_N}^h, \quad (9)$$

where $(P^h)_{h \in H}$ are the possible values taken by \tilde{P} .

Representation theorems. The classic de Finetti's representation theorem characterizes exchangeable sequences as mixtures of i.i.d. sequences. The statement, the proof, and an extensive discussion of the ramifications of the theorem can be found in [1] and [9]. The representation theorem characterizing partially exchangeable sequences as mixtures of Markov chains was first proved in [4], for S discrete and using a slightly different notion of partial exchangeability, and later extended to S a general Polish space in [8].

Hidden Markov models. Let S be a Polish space endowed with the Borel σ -field \mathcal{S} .

Definition 2.8. *A random sequence (Y_n) is an hidden Markov model (HMM) if there exists a pair (X_n, \tilde{Y}_n) taking values on $\mathcal{X} \times S$ for a discrete space \mathcal{X} , such that*

1. (X_n) is a time homogeneous Markov chain on \mathcal{X} ,
2. (conditional independence property) for any $N \in \mathbb{N}$, and for any $S_0, \dots, S_N \in \mathcal{S}$ it holds

$$\mathbb{P}(\tilde{Y}_0 \in S_0, \dots, \tilde{Y}_N \in S_N \mid X_0^N = x_0^N) = \prod_{n=0}^N \mathbb{P}(\tilde{Y}_n \in S_n \mid X_n = x_n),$$

3. (Y_n) and (\tilde{Y}_n) have the same distributions, i.e. for any $N \in \mathbb{N}$, and for any $S_0, \dots, S_N \in \mathcal{S}$ it holds

$$P(Y_0 \in S_0, \dots, Y_N \in S_N) = P(\tilde{Y}_0 \in S_0, \dots, \tilde{Y}_N \in S_N).$$

It is often possible to verify the second property directly for the sequence (Y_n) .

A HMM is characterized by the initial distribution π on \mathcal{X} , by the transition matrix $P = (P_{ij})_{i,j \in \mathcal{X}}$ of the Markov chain (X_n) , and by the *read-out* distributions $f_x(\bar{S})$, with $\bar{S} \in \mathcal{S}$, where

$$f_x(\bar{S}) := \mathbb{P}(\tilde{Y}_n \in \bar{S} \mid X_n = x). \quad (10)$$

We refer to the sequence (X_n) as the "underlying Markov chain" of the HMM.

For discrete S , there are many equivalent definitions of HMMs (see [11]), not all making sense for S Polish.

Remark 2.9. *A countable mixture of i.i.d. sequences as in Equation (2) is an HMM: take as (X_n) the Markov chain with values in H , with identity transition matrix, initial distribution $(\mu_1, \dots, \mu_h, \dots)_{h \in H}$, and read-out distributions $\mathbb{P}(Y_n \in \bar{S} \mid X_n = h) = p_h(\bar{S})$.*

3 Countable Markov mixtures with discrete state space

In this section S is a discrete set.

3.1 The successors array of hidden Markov models

The following result will be instrumental later, and it is also of independent interest. It is based on some useful properties of HMMs, that can be found in the Appendix. HMMs and the successors array are defined in Section 2.

Proposition 3.1. *Let (Y_n) be a HMM on a discrete space S with recurrent¹ underlying Markov chain, then each row of the successors array $(V_{y,n})$ is a HMM with recurrent underlying Markov chain.*

Proof. Denote with \mathcal{X} the discrete state space of the Markov chain (X_n) , underlying the process (Y_n) and let, for any $x \in \mathcal{X}$ and $y \in S$,

$$f_x(y) := \mathbb{P}(Y_n = y \mid X_n = x)$$

¹A time homogeneous Markov chain (X_n) is recurrent if $\mathbb{P}(X_n = x \text{ i.o. } n \mid X_1 = x) = 1$, for all $x \in \mathcal{X}$ such that $\mathbb{P}(X_1 = x) > 0$. Such Markov chains have no transient states but possibly more than one recurrence class.

be the read-out distributions. Fix $y \in S$. To prove the theorem we construct a recurrent Markov chain $(W_n^y)_{n \geq 1}$ such that the pair $(W_n^y, V_{y,n})_{n \geq 1}$ satisfies the conditions of Definition 2.8 of HMM (note that for convenience we let time start at $n = 1$). The proof is divided in three main steps.

Step 1 *Construction of the Markov chain (W_n^y) .* To construct the Markov chain (W_n^y) , define inductively the random times of the n -th visit of (Y_n) to state y :

$$\begin{aligned}\tau_1^y &:= \inf\{t \geq 0 \mid Y_t = y\}, \\ \tau_n^y &:= \inf\{t > \tau_{n-1}^y \mid Y_t = y\},\end{aligned}$$

with the usual convention $\inf \emptyset = +\infty$. The random times (τ_n^y) are stopping times with respect to the filtration spanned by (Y_n) , and so are the times $(\tau_n^y + 1)$. The random times $(\tau_n^y + 1)$ are actually hitting-times of $\mathcal{X} \times y$, according to the Definition 6.4 in the Appendix. Define the sequence

$$W_n^y := \begin{cases} \varepsilon & \text{for } \tau_n^y = +\infty, \\ X_{\tau_n^y + 1} & \text{for } \tau_n^y < +\infty, \end{cases} \quad (1)$$

where $\varepsilon \notin \mathcal{X}$ is a fictitious state. The sequence W_n^y is either identically equal to ε , or it never hits it since the times τ_n^y are either all finite or all infinite².

Let us check that (W_n^y) is a Markov chain. If the case $W_n^y \equiv \varepsilon$ obtains, (W_n^y) is a (recurrent) Markov chain. Otherwise a direct computation gives, for any $N \in \mathbb{N}$, and any $x_1, \dots, x_N \in \mathcal{X}$,

$$\begin{aligned}\mathbb{P}(W_N^y = x_N \mid W_{N-1}^y = x_{N-1}, \dots, W_1^y = x_1) \\ &= \mathbb{P}(X_{\tau_N^y + 1} = x_N \mid X_{\tau_{N-1}^y + 1} = x_{N-1}, \dots, X_{\tau_1^y + 1} = x_1) \\ &= \mathbb{P}(X_{\tau_N^y + 1} = x_N \mid X_{\tau_{N-1}^y + 1} = x_{N-1}) = \mathbb{P}(W_N^y = x_N \mid W_{N-1}^y = x_{N-1}),\end{aligned}$$

where Remark 6.7 in the Appendix applies. Thus (W_n^y) is Markov.

Step 2 *Check of the recurrence of (W_n^y) .* Since

$$\mathbb{P}(W_n^y = x \text{ i.o. } n \mid W_1^y = x) = \mathbb{P}(X_{\tau_n^y + 1} = x \text{ i.o. } n \mid X_{\tau_1^y + 1} = x),$$

to check the recurrence of (W_n^y) we have to verify that, for all $x \in \mathcal{X}$,

$$\mathbb{P}(X_{\tau_n^y + 1} = x \text{ i.o. } n \mid X_{\tau_1^y + 1} = x) = 1. \quad (2)$$

²If $Y_n = y$, for some finite n , then $X_n = x$, for some x such that $f_x(y) > 0$. Since X is recurrent it hits x infinitely many times, thus Y hits y infinitely many times.

Fix $x \in \mathcal{X}$ such that $\mathbb{P}(X_{\tau_1^y+1} = x) > 0$, and choose $\bar{x} \in \mathcal{X}$ such that $f_{\bar{x}}(y) > 0$ and $\mathbb{P}(X_{n+1} = x \mid X_n = \bar{x}) > 0$, (there exists at least one such \bar{x}). Define the auxiliary sequence of hitting times:

$$\begin{aligned}\sigma_1^{\bar{x},y} &:= \inf\{t \geq 0 \mid X_t = \bar{x}, Y_t = y\}, \\ \sigma_n^{\bar{x},y} &:= \inf\{t > \sigma_{n-1}^{\bar{x},y} \mid X_t = \bar{x}, Y_t = y\}.\end{aligned}$$

The hitting times $(\sigma_n^{\bar{x},y})$ are finite whenever (τ_n^y) are finite, and the sequence $(\sigma_n^{\bar{x},y})$ is a subsequence of (τ_n^y) , thus

$$(X_{\sigma_n^{\bar{x},y}+1} = x) \subseteq \bigcup_{m \geq n} (X_{\tau_m^y+1} = x), \quad (3)$$

and trivially

$$(X_{\sigma_n^{\bar{x},y}+1} = x \text{ i.o. } n) \subseteq (X_{\tau_n^y+1} = x \text{ i.o. } n). \quad (4)$$

The events $(X_{\sigma_n^{\bar{x},y}+1} = x)_n$ are independent under the law $\mathbb{P}(\cdot \mid X_{\tau_1^y+1} = x)$, since $\{(X_{\sigma_n^{\bar{x},y}+1} = x)_n, (X_{\tau_1^y+1} = x)\}$ is a \mathbb{P} -independent set. In fact, for any $N \in \mathbb{N}$, and for any choice of $m_1 < \dots < m_N \in \mathbb{N}$,

$$\begin{aligned}\mathbb{P}(X_{\sigma_{m_N}^{\bar{x},y}+1} = x, X_{\sigma_{m_{N-1}}^{\bar{x},y}+1} = x, \dots, X_{\sigma_{m_1}^{\bar{x},y}+1} = x, X_{\tau_1^y+1} = x) \\ &= \mathbb{P}(X_{\sigma_{m_N}^{\bar{x},y}+1} = x, X_{\sigma_{m_N}^{\bar{x},y}} = \bar{x}, \dots, X_{\sigma_{m_1}^{\bar{x},y}+1} = x, X_{\sigma_{m_1}^{\bar{x},y}} = \bar{x}, X_{\tau_1^y+1} = x) \\ &= \mathbb{P}(X_{\sigma_{m_N}^{\bar{x},y}+1} = x \mid X_{\sigma_{m_N}^{\bar{x},y}} = \bar{x}) \mathbb{P}(X_{\sigma_{m_N}^{\bar{x},y}} = \bar{x} \mid X_{\sigma_{m_{N-1}}^{\bar{x},y}+1} = x, \dots, X_{\sigma_{m_1}^{\bar{x},y}+1} = x, X_{\sigma_{m_1}^{\bar{x},y}} = \bar{x}, X_{\tau_1^y+1} = x) \\ &\quad \times \dots \times \mathbb{P}(X_{\sigma_{m_1}^{\bar{x},y}+1} = x \mid X_{\sigma_{m_1}^{\bar{x},y}} = \bar{x}) \times \mathbb{P}(X_{\sigma_{m_1}^{\bar{x},y}} = \bar{x} \mid X_{\tau_1^y+1} = x) \mathbb{P}(X_{\tau_1^y+1} = x) \\ &= \mathbb{P}(X_{\sigma_{m_N}^{\bar{x},y}+1} = x) \dots \mathbb{P}(X_{\sigma_{m_1}^{\bar{x},y}+1} = x) \mathbb{P}(X_{\tau_1^y+1} = x),\end{aligned}$$

where the second equality follows by Remark 6.3 in the Appendix, and the first and last equality follow noting that $X_{\sigma_{m_N}^{\bar{x},y}} = \bar{x}$ for any $\omega \in \Omega$ by definition of $\sigma^{\bar{x},y}$. Note that by definition $\sigma_n^{\bar{x},y} > \tau_1^y$ for any $n > 1$, but it could happen $\sigma_1^{\bar{x},y} = \tau_1^y$, so the computation above needs some care for $m_1 = 1$, but can be easily recovered also in this case.

The events $(X_{\sigma_n^{\bar{x},y}+1} = x)$ are equiprobable, with strictly positive probability. By the Borel-Cantelli lemma

$$\mathbb{P}(X_{\sigma_n^{\bar{x},y}+1} = x \text{ i.o. } n \mid X_{\tau_1^y+1} = x) = 1. \quad (5)$$

Equations (4) and (5) taken together give

$$\mathbb{P}(X_{\tau_n^y+1} = x \text{ i.o. } n \mid X_{\tau_1^y+1} = x) \geq \mathbb{P}(X_{\sigma_n^{\bar{x},y}+1} = x \text{ i.o. } n \mid X_{\tau_1^y+1} = x) = 1.$$

Condition (2) is satisfied, thus the recurrence of (W_n^y) is proved.

Step 3 *Verification that the pair $(W_n^y, V_{y,n})$ is a HMM.* Let us check that the pair $(W_n^y, V_{y,n})$ is as in Definition 2.8. Set

$$\mathbb{P}(V_{y,n} = \delta \mid W_n^y = \varepsilon) = 1.$$

For $\varepsilon \neq x \in \mathcal{X}$ and $\delta \neq \bar{y} \in S$, the pair $(W_n^y, V_{y,n})$ inherits the read-out distributions of (X_n, Y_n) :

$$\mathbb{P}(V_{y,n} = \bar{y} \mid W_n^y = x) = \mathbb{P}(Y_{\tau_n^y+1} = \bar{y} \mid X_{\tau_n^y+1} = x) = f_x(\bar{y}), \quad (6)$$

see Lemma 6.8 in the Appendix. Let us verify the conditional independence property, i.e. that for any $N \in \mathbb{N}$ and any $y_1^N \in S^N$ and any $x_1^N \in \mathcal{X}^N$

$$\begin{aligned} & \mathbb{P}(V_{y,1} = y_1, \dots, V_{y,N} = y_N \mid W_1^y = x_1, \dots, W_N^y = x_N) \\ &= \prod_{n=1}^N \mathbb{P}(V_{y,n} = y_n \mid W_n^y = x_n). \end{aligned}$$

It follows from the direct computation,

$$\begin{aligned} & \mathbb{P}(V_{y,1} = y_1, \dots, V_{y,N} = y_N \mid W_1^y = x_1, \dots, W_N^y = x_N) \\ &= \mathbb{P}(Y_{\tau_1^y+1} = y_1, \dots, Y_{\tau_N^y+1} = y_N \mid X_{\tau_1^y+1} = x_1, \dots, X_{\tau_N^y+1} = x_N) \\ &= \prod_{n=1}^N \mathbb{P}(Y_{\tau_n^y+1} = y_n \mid X_{\tau_n^y+1} = x_n) = \prod_{n=1}^N \mathbb{P}(V_{y,n} = y_n \mid W_n^y = x_n), \end{aligned}$$

where the second equality is a direct consequence of Lemma 6.9 of the Appendix. The sequence $(V_{y,n})_n$ is therefore a HMM with recurrent underlying Markov chain, and this concludes the proof of the proposition. \square \square

3.2 Representation of countable mixtures

In [3] Dharmadhikari gives a characterization of countable mixtures of i.i.d. sequences, linking HMMs to the class of exchangeable sequences. The main result of [3] can be rephrased as follows (see Section 2 for the definitions of exchangeable sequences, mixture of i.i.d. sequences and HMM).

Theorem 3.2. (*Dharmadhikari*) *Let (Y_n) be an exchangeable sequence on a discrete state space S . The sequence (Y_n) is a countable mixture of i.i.d. sequences if and only if (Y_n) is a HMM with recurrent underlying Markov chain.*

In the original formulation of Theorem 3.2 the stationarity of the underlying Markov chain is one of the hypotheses, but close inspection of the proof in [3] reveals that only the absence of transient states is required.

The aim of this section is to extend the above theorem to partially exchangeable sequences, i.e. to characterize *countable* mixtures of Markov chains. The analog of Theorem 3.2 for mixtures of Markov chains is as follows (we refer to Section 2 for the definition of partially exchangeable sequences, mixture of Markov chains and HMMs).

Theorem 3.3. *Let (Y_n) be a partially exchangeable sequence on a discrete state space S , with $\mathbb{P}(Y_0 = y_0) = 1$ for some $y_0 \in S$. The sequence (Y_n) is a countable mixture of Markov chains if and only if (Y_n) is a HMM with recurrent underlying Markov chain.*

Proof. The standing hypothesis is that (Y_n) is a partially exchangeable sequence. We first prove that if (Y_n) is a HMM, then it is a countable mixture of Markov chains i.e., in the notations of Remark 2.7, \tilde{P} takes countably many values. By the partial exchangeability of (Y_n) , the row $(V_{y,n})$, for any $y \in S$, is exchangeable, and therefore a mixture of i.i.d. sequences. As proved e.g. in Lemma 2.15 of [1] or in Proposition 1.1.4 of [9],

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbb{I}_{V_{y,n}}(\cdot) = \tilde{p}_y(\cdot) \quad \mathbb{P} - a.s., \quad (7)$$

where the limit has to be interpreted in the topology of weak convergence, and where \tilde{p}_y is the random probability measure with values in \mathcal{M}_S corresponding to \tilde{p} in Definition 2.3. It follows from the proof of Theorem 1 in [8], that the random probability measure \tilde{p}_y in Equation (7) is the y -th row of the random matrix \tilde{P} . By Proposition 3.1 above, each row $(V_{y,n})$ is a HMM, and therefore, by Theorem 3.2 above, it is a countable mixture of i.i.d. sequences. The random probability measure \tilde{p}_y is thus concentrated on a countable set, and so is the y -th row of \tilde{P} . Since this holds for each y , the conclusion is that \tilde{P} takes countably many values.

To prove the converse one has to show that if (Y_n) is a given countable mixture of Markov chains, i.e. if Equation (9) holds for some countable family $(P^h)_{h \in H}$, then (Y_n) is a HMM. We construct a pair (X_n, \tilde{Y}_n) satisfying the conditions of Definition 2.8, with (X_n) recurrent, and such that (\tilde{Y}_n) and the given (Y_n) have the same distributions.

The Markov chain (X_n) is defined on the state space³ $S \times H$, with transition probability matrix \mathbf{P} , the direct sum of the transition matrices P^h ,

$$\mathbf{P} := \begin{pmatrix} P^1 & 0 & 0 & \dots \\ 0 & P^2 & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

and initial distribution π defined, for any $y \in S$ and $h \in H$, as

$$\pi(y, h) = \begin{cases} \mu_h & \text{for } y = y_0 \\ 0 & \text{for } y \neq y_0, \end{cases}$$

and $\mu_h := \mathbb{P}(\tilde{P} = P^h)$. To show that (X_n) is recurrent note that by Theorem 1 in [8], (Y_n) is conditionally recurrent, therefore the matrices $\{P^h\}$ in the mixture correspond to recurrent chains. Since \mathbf{P} is the direct sum of such matrices, (X_n) is recurrent.

Consider now a sequence (\tilde{Y}_n) , with fixed initial state $\tilde{Y}_0 = y_0$, conditionally independent given (X_n) , and with read-out distributions defined as follows

$$f_{(x,h)}(y) := \mathbb{P}(\tilde{Y}_n = y \mid X_n = (x, h)) = \delta_{x,y},$$

where $\delta_{\cdot, \cdot}$ is the Kronecker symbol. Let us compute the finite distributions of (\tilde{Y}_n) , for any $N \in \mathbb{N}$ and any $y_1^N \in (S^*)^N$,

³E.g. ordering the states in first lexical order as follows: $(1, 1), (2, 1), \dots, (1, 2), (2, 2), \dots$

$$\begin{aligned}
\mathbb{P}(\tilde{Y}_1^N = y_1^N) &= \sum_{(x_0^N, h_0^N) \in S^{N+1} \times H^{N+1}} \mathbb{P}(\tilde{Y}_0^N = y_0^N, X_0^N = (x_0^N, h_0^N)) \\
&= \sum_{(x_0^N, h_0^N) \in S^{N+1} \times H^{N+1}} \mathbb{P}(X_0 = (x_0, h_0)) \prod_{n=0}^N \mathbb{P}(\tilde{Y}_n = y_n \mid X_n = (x_n, h_n)) \\
&\quad \times \prod_{n=0}^{N-1} \mathbb{P}(X_{n+1} = (x_{n+1}, h_{n+1}) \mid X_n = (x_n, h_n)) \\
&= \sum_{(x_0^N, h_0^N) \in S^{N+1} \times H^{N+1}} \pi(x_0, h_0) \prod_{n=0}^N f_{(x_n, h_n)}(y_n) \prod_{n=0}^{N-1} \mathbf{P}_{(x_n, h_n)(x_{n+1}, h_{n+1})} \\
&= \sum_{h \in H} \mu_h P_{y_0 y_1}^h \cdots P_{y_{N-1} y_N}^h,
\end{aligned}$$

where the second equality follows from the conditional independence of (\tilde{Y}_n) given (X_n) , and the fourth from the definition of the read-out densities and by the block structure of \mathbf{P} . Comparing (9) with the last expression, we have that (Y_n) and (\tilde{Y}_n) have the same distributions, thus (Y_n) is a HMM with recurrent underlying Markov chain and the theorem is proved. \square

Note that, if S finite, in Theorems 3.2 and 3.3 the state space of the underlying Markov chain is *finite* if and only if the mixture is *finite*.

4 Countable Markov mixtures with Polish state space

In this section S is a Polish space.

4.1 The successors array

The proposition below is the analog of Proposition 3.1 for uncountable state space S (the definitions of HMM and of successors array are in Section 2).

Proposition 4.1. *Let (Y_n) be a HMM on a Polish space S with recurrent underlying Markov chain, then each row of the successors array $(V_{j,n})$ is a HMM with recurrent underlying Markov chain.*

Proof. Let (X_n) be the underlying Markov chain of (Y_n) . Consider the partition $\mathcal{E} = (E_j)_{j \geq 0}$ of S^* , and for any element E_j of the partition define

$$\tau_1^{E_j} := \inf\{t \geq 0 \mid Y_t \in E_j\}, \quad \tau_n^{E_j} := \inf\{t > \tau_{n-1}^{E_j} \mid Y_t \in E_j\}.$$

The proof can be carried out exactly as the proof of Proposition 3.1, substituting τ_n^y there with $\tau_n^{E_j}$, and $\sigma_n^{\bar{x},y}$ with $\sigma_n^{\bar{x},E_j}$, defined below

$$\begin{aligned} \sigma_1^{\bar{x},E_j} &:= \inf\{t \geq 0 \mid X_t = \bar{x}, Y_t \in E_j\}, \\ \sigma_n^{\bar{x},E_j} &:= \inf\{t > \sigma_{n-1}^{\bar{x},y} \mid X_t = \bar{x}, Y_t \in E_j\}, \end{aligned}$$

where $\bar{x} \in \mathcal{X}$ is such that $f_{\bar{x}}(E_j) > 0$ and $\mathbb{P}(X_{n+1} = x \mid X_n = \bar{x}) > 0$, (x has the same role as in Equation (2)). \square

4.2 Representation of countable mixtures

This subsection mirrors Subsection 3.2 for the case of Polish state space S . Theorem 4.2 below extends Theorem 3.2 to Polish state spaces. To the best of our knowledge the extension is not available in the literature. Based on Theorem 4.2 we prove Theorem 4.3 which is the counterpart of Theorem 3.3 and the main result of the subsection.

Note that Theorem 3.2, i.e. Dharmadhikari's original result [3], can not be directly generalized as it relies on a definition of HMMs unsuitable for general state spaces.

4.2.1 Representation of countable i.i.d. mixtures

Exchangeable sequences, mixture of i.i.d. sequences, and HMMs are defined in Section 2.

Theorem 4.2. *Let (Y_n) be an exchangeable sequence on a Polish space. The sequence (Y_n) is a countable mixture of i.i.d. sequences if and only if (Y_n) is a HMM with recurrent underlying Markov chain.*

Proof. If (Y_n) is a countable mixture of i.i.d., then it is a HMM by Remark 2.9. To prove the converse let (Y_n) be an exchangeable HMM, whose recurrent underlying Markov chain (X_n) has transition probability matrix P and initial distribution π . The Markov chain (X_n) has no transient states, but possibly more than one recurrence class. As noted in [3], by the exchangeability of

(Y_n) , one can substitute P with the Cesàro limit $P^* := \lim_{n \rightarrow \infty} 1/n \sum_{k=1}^n P^k$, where P^k is the k -power of P . By the ergodic theorem P^* has a block structure, being the direct sum of matrices P_h with identical rows, one block P_h for each recurrence class. By Lemma 6.10 of the Appendix (Y_n) is a countable mixture of i.i.d. sequences. \square

4.2.2 Representation of countable Markov mixtures

See Section 2 for the definitions of partially exchangeable sequence, mixture of Markov chains and HMM.

Theorem 4.3. *Let (Y_n) be a partially exchangeable sequence on a Polish space with $\mathbb{P}(Y_0 = y_0) = 1$, for some $y_0 \in E_1$. The sequence (Y_n) is a countable mixture of homogeneous Markov chains if and only if (Y_n) is a HMM with recurrent underlying Markov chain.*

Proof. The partially exchangeability of (Y_n) is a standing hypothesis. Let (Y_n) be a HMM with recurrent underlying Markov chain. To prove that (Y_n) is a countable mixture of Markov chains imitate the proof of Theorem 3.3. Note first that $\frac{1}{N} \sum_{n=1}^N \mathbb{I}_{V_{j,n}}(\cdot) \rightarrow \theta_j(\cdot)$, where θ_j is a probability measure on \mathcal{S} . As in the proof of Theorem 4 in [8], for any $j \in \mathbb{N}$ define $\tilde{t}(j, \cdot) := \theta_j(\cdot)$. To conclude use Proposition 4.1 and Theorem 4.2.

For the converse assume that (Y_n) is a countable mixture of Markov chains, with random kernel \tilde{t} taking values $(t_h)_{h \in H}$ and with $\mu_h = \mathbb{P}(\tilde{t} = t_h)$, and finite distributions as in Equation (7). To prove that (Y_n) is a HMM with recurrent underlying Markov chain we construct a recurrent Markov chain (X_n) and a sequence (\tilde{Y}_n) satisfying the first two conditions in Definition 2.8, then showing that (\tilde{Y}_n) has the same distributions of (Y_n) ⁴. Consider thus a Markov chain (X_n) taking values in $H \times \mathbb{N}_0 \times \mathbb{N}_0$, with components $X_n = (h_n, i_n, j_n)$ representing the index of the running chain in the mixture, the discretized value of Y_{n-1} (i.e. the elements of the partition to which Y_{n-1} belongs), and the discretized value of Y_n respectively. The initial distribution of (X_n) is taken to be

$$\mathbb{P}(X_0 = (h, i, j)) = \mu_h \frac{1}{2^{i+1}} \delta_{j,1},$$

⁴ The construction used for the proof of Theorem 3.3 can not be used here, in fact the Markov chain (X_n) there takes values in the product space $H \times S$, which can be now uncountable, while we need a discrete underlying Markov chain.

where $\delta_{\cdot, \cdot}$ is again the Kronecker symbol, and its transition probabilities

$$\mathbb{P}\left(X_n = (h_n, i_n, j_n) \mid X_{n-1} = (h_{n-1}, i_{n-1}, j_{n-1})\right) = \delta_{h_{n-1}, h_n} \delta_{i_n, j_{n-1}} t_{h_n}(i_n, E_{j_n}).$$

The Markov chain (X_n) is recurrent since the kernels t_h correspond to recurrent Markov chains by Theorem 4 in [8]. Consider now a sequence (\tilde{Y}_n) jointly distributed with (X_n) , with fixed initial value $\tilde{Y}_0 = y_0$, conditionally independent given (X_n) , and with read-out distributions defined as follows for any $A \in \mathcal{S}$

$$\mathbb{P}(\tilde{Y}_n \in A \mid X_n = (h, i, j)) = \begin{cases} 0 & \text{for } t_h(i, E_j) = 0 \\ \frac{1}{t_h(i, E_j)} \int_{A \cap E_j} t_h(i, dy) & \text{for } t_h(i, E_j) \neq 0. \end{cases}$$

For any $N \in \mathbb{N}$, and any $A_1, \dots, A_N \in \mathcal{S}^*$, the distributions of (\tilde{Y}_n) are

computed as follows

$$\begin{aligned}
& \mathbb{P}(\tilde{Y}_1 \in A_1, \dots, \tilde{Y}_N \in A_N) \\
&= \sum_{h_1^N \in H^N, i_1^N \in (\mathbb{N}_0)^N, j_1^N \in (\mathbb{N}_0)^N} \mathbb{P}(\tilde{Y}_1 \in A_1, \dots, \tilde{Y}_N \in A_N, X_1 = (h_1, i_1, j_1), \dots, X_N = (h_N, i_N, j_N)) \\
&= \sum_{h_1^N \in H^N, i_1^N \in (\mathbb{N}_0)^N, j_1^N \in (\mathbb{N}_0)^N} \mathbb{P}(\tilde{Y}_1 \in A_1, \dots, \tilde{Y}_N \in A_N \mid X_1 = (h_1, i_1, j_1), \dots, X_N = (h_N, i_N, j_N)) \\
&\quad \times \mathbb{P}(X_1 = (h_1, i_1, j_1), \dots, X_N = (h_N, i_N, j_N)) \\
&= \sum_{h_0^N \in H^N, i_0^N \in (\mathbb{N}_0)^{N+1}, j_0^N \in (\mathbb{N}_0)^{N+1}} \prod_{n=1}^N \mathbb{P}(\tilde{Y}_n \in A_n \mid X_n = (h_n, i_n, j_n)) \\
&\quad \times \prod_{n=1}^N \mathbb{P}(X_n = (h_n, i_n, j_n) \mid X_{n-1} = (h_{n-1}, i_{n-1}, j_{n-1})) \mathbb{P}(X_0 = (h_0, i_0, j_0)) \\
&= \sum_{h_0^N \in H^N, i_0^N \in (\mathbb{N}_0)^{N+1}, j_0^N \in (\mathbb{N}_0)^{N+1}} \prod_{n=1}^N \frac{1}{t_{h_n}(i_n, E_{j_n})} \int_{A_n \cap E_{j_n}} t_{h_n}(i_n, dy) \\
&\quad \times \prod_{n=1}^N \delta_{h_{n-1}, h_n} \delta_{i_n, j_{n-1}} t_{h_n}(i_n, E_{j_n}) \mu_{h_0} \delta_{j_0, 1} \frac{1}{2^{i_0+1}} \\
&= \sum_{h \in H, i_0 \in \mathbb{N}_0, j_0^N \in (\mathbb{N}_0)^{N+1}} \prod_{n=1}^N \int_{A_n \cap E_{j_n}} t_h(j_{n-1}, dy) \mu_h \delta_{j_0, 1} \frac{1}{2^{i_0+1}} \\
&= \sum_{h \in H, j_1^N \in (\mathbb{N}_0)^N} \mu_h \int_{A_1 \cap E_{j_1}} t_h(1, dy) \prod_{n=2}^N \int_{A_n \cap E_{j_n}} t_h(j_{n-1}, dy).
\end{aligned}$$

Comparing the expression above with Equation (7), one concludes that the distributions of (\tilde{Y}_n) coincide with those of (Y_n) , therefore proving that (Y_n) is a HMM. \square

5 Concluding remarks

Throughout the paper we referred to the notion of partial exchangeability originally given by de Finetti and to the corresponding representation theorem as given in [8]. For discrete state space partial exchangeability can be defined in a slightly different way, and a representation theorem in this

alternative framework is proved in [4]. According to [4], a sequence of random variables is partially exchangeable if the probability is invariant under all permutations of a string that preserves the first value and the transition counts between any couple of states. A characterization of countable mixtures of Markov chains can be given also in the setup of [4], using different mathematical tools. The result is in [7], but for a complete proof see [10]. By the same token the characterization of countable mixtures of Markov chains of order k holds true, for the proof see [10]. Unfortunately the approach of [7] and [10] does not readily generalize to Polish state space.

Based on the results in [8], a de Finetti's type representation theorem for mixtures of semi-Markov processes have been proved in [6]. The authors are confident that a characterization of countable mixtures of semi-Markov processes in terms of HMMs can be given properly adapting the proof of Proposition 4.1 and Theorem 4.3.

6 Appendix

6.1 Strong Markov and strong conditional independence for HMMs

This section contains some useful properties of HMMs.

Lemma 6.1. (*Splitting property*) *Let (Y_n) be a HMM with underlying Markov chain (X_n) . Then the pair (X_n, Y_n) is a Markov chain. Moreover for any $N \in \mathbb{N}$, for any $x_1, \dots, x_N \in \mathcal{X}$, and $S_1, \dots, S_N \in \mathcal{S}$ such that $\mathbb{P}(X_1^{N-1} = x_1^{N-1}, Y_1^{N-1} \in S_1^{N-1}) > 0$ we have*

$$\begin{aligned} & \mathbb{P}(X_N = x, Y_N \in S_N \mid X_1^{N-1} = x_1^{N-1}, Y_1^{N-1} \in S_1^{N-1}) \\ & = \mathbb{P}(X_N = x, Y_N \in S_N \mid X_{N-1} = x_{N-1}). \end{aligned}$$

Proof.

$$\begin{aligned}
& \mathbb{P}(X_N = x, Y_N \in S_N \mid X_1^{N-1} = x_1^{N-1}, Y_1^{N-1} \in S_1^{N-1}) \\
&= \frac{\mathbb{P}(X_1^N = x_1^N, Y_1^N \in S_1^N)}{\mathbb{P}(X_1^{N-1} = x_1^{N-1}, Y_1^{N-1} \in S_1^{N-1})} \\
&= \frac{\prod_{n=1}^N \mathbb{P}(Y_n \in S_n \mid X_n = x_n) \mathbb{P}(X_1^N = x_1^N)}{\prod_{n=1}^{N-1} \mathbb{P}(Y_n \in S_n \mid X_n = x_n) \mathbb{P}(X_1^{N-1} = x_1^{N-1})} \\
&= \mathbb{P}(Y_N \in S_N \mid X_N = x_N) \mathbb{P}(X_N = x_N \mid X_{N-1} = x_{N-1}) \\
&= \mathbb{P}(Y_N \in S_N \mid X_N = x_N, X_{N-1} = x_{N-1}) \mathbb{P}(X_N = x_N \mid X_{N-1} = x_{N-1}) \\
&= \mathbb{P}(Y_N \in S_N, X_N = x_N \mid X_{N-1} = x_{N-1}),
\end{aligned}$$

where the second and fourth equality follow by the conditional independence of the observations (Y_n) in the definition of HMM. \square

Lemma 6.2. (*Strong splitting property*) *Let (Y_n) be a HMM with underlying Markov chain (X_n) , and γ be a stopping time for (X_n, Y_n) , then, for any $x, \tilde{x}, \bar{x} \in \mathcal{X}$, and any $S_1, S_2, S_3 \in \mathcal{S}$ such that $\mathbb{P}(X_\gamma = \tilde{x}, Y_\gamma \in S_2, X_{\gamma \wedge n} = \bar{x}, Y_{\gamma \wedge n} \in S_1) > 0$ it holds that*

$$\begin{aligned}
& \mathbb{P}(X_{\gamma+k} = x, Y_{\gamma+k} \in S_3 \mid X_\gamma = \tilde{x}, Y_\gamma \in S_2, X_{\gamma \wedge n} = \bar{x}, Y_{\gamma \wedge n} \in S_1) \\
&= \mathbb{P}(X_{\gamma+k} = x, Y_{\gamma+k} \in S_3 \mid X_\gamma = \tilde{x}). \tag{1}
\end{aligned}$$

Proof. We manipulate separately the left-hand side (LHS) and the right-hand side (RHS) of Equation (1). For readability denote $C_r := (\gamma = r, X_r = \tilde{x}, Y_r \in S_2, X_{r \wedge n} = \bar{x}, Y_{r \wedge n} \in S_1)$. Applying Lemma 6.1, the numerator of the conditional probability on the LHS of Equation (1) is

$$\begin{aligned}
& \mathbb{P}(X_{\gamma+k} = x, Y_{\gamma+k} \in S_3, X_\gamma = \tilde{x}, Y_\gamma \in S_2, X_{\gamma \wedge n} = \bar{x}, Y_{\gamma \wedge n} \in S_1) \\
&= \sum_{r \geq 1} \mathbb{P}(X_{r+k} = x, Y_{r+k} \in S_3 \mid C_r) \mathbb{P}(C_r) \\
&= \sum_{r \geq 1} \mathbb{P}(X_{r+k} = x, Y_{r+k} \in S_3 \mid X_r = \tilde{x}) \mathbb{P}(C_r) \\
&= \sum_{r \geq 1} \mathbb{P}(Y_{r+k} \in S_3 \mid X_{r+k} = x) \mathbb{P}(X_{r+k} = x \mid X_r = \tilde{x}) \mathbb{P}(C_r) \\
&= f_x(S_3) P_{\tilde{x}, x}^{(k)} \sum_{r \geq 1} \mathbb{P}(C_r) \\
&= f_x(S_3) P_{\tilde{x}, x}^{(k)} \mathbb{P}(X_\gamma = \tilde{x}, Y_\gamma \in S_2, X_{\gamma \wedge n} = \bar{x}, Y_{\gamma \wedge n} \in S_1),
\end{aligned}$$

where $P_{\tilde{x}, x}^{(k)}$ is the \tilde{x}, x -entry of the k -step transition matrix of the Markov chain (X_n) . The numerator of the conditional probability on the RHS of Equation (1), again applying Lemma 6.1, is

$$\begin{aligned}
& \mathbb{P}(X_{\gamma+k} = x, Y_{\gamma+k} \in S_3, X_\gamma = \tilde{x}) \\
&= \sum_{r \geq 1} \mathbb{P}(X_{r+k} = x, Y_{r+k} \in S_3 \mid \gamma = r, X_r = \tilde{x}) \mathbb{P}(\gamma = r, X_r = \tilde{x}) \\
&= \sum_{r \geq 1} \mathbb{P}(X_{r+k} = x, Y_{r+k} \in S_3 \mid X_r = \tilde{x}) \mathbb{P}(\gamma = r, X_r = \tilde{x}) \\
&= f_x(S_3) P_{\tilde{x}, x}^{(k)} \mathbb{P}(X_\gamma = \tilde{x}).
\end{aligned}$$

The lemma is proved comparing the expressions of the LHS and the RHS derived above. \square

Taking $S_1 = S_2 = S_3 = S$ we have

Remark 6.3. *Let $(X_n), \gamma$ be as in Lemma 6.2 then, for any $x, \tilde{x}, \bar{x} \in \mathcal{X}$, such that $\mathbb{P}(X_\gamma = \tilde{x}, X_{\gamma \wedge n} = \bar{x}) > 0$ it holds that*

$$\mathbb{P}(X_{\gamma+k} = x, \mid X_\gamma = \tilde{x}, X_{\gamma \wedge n} = \bar{x}) = \mathbb{P}(X_{\gamma+k} = x, \mid X_\gamma = \tilde{x}). \quad (2)$$

Definition 6.4. Let (Y_n) be a HMM with underlying Markov chain (X_n) , and let $A \subset \mathcal{X} \times \mathcal{S}$. We say that the sequence of random times $(\gamma_n)_{n \geq 1}$ is a sequence of hitting times of A if

$$\begin{aligned}\gamma_1 &:= \inf\{t \geq 0 \mid (X_t, Y_t) \in A\}, \\ \gamma_n &:= \inf\{t > \gamma_{n-1} \mid (X_t, Y_t) \in A\}.\end{aligned}$$

Lemma 6.5. (*Generalized strong splitting property*) Let (Y_n) be a HMM with underlying Markov chain (X_n) . Let (γ_n) be a sequence of hitting times of A for (X_n, Y_n) , where $A \subset \mathcal{X} \times \mathcal{S}$. Then for any N , and any $(x_1, S_1), \dots, (x_N, S_N) \in A$ such that $\mathbb{P}(X_{\gamma_1}^{\gamma_1} = x_1^{\gamma_1}, Y_{\gamma_1}^{\gamma_1} \in S_1^{\gamma_1}) > 0$ it holds

$$\begin{aligned}\mathbb{P}(X_{\gamma_N} = x_N, Y_{\gamma_N} \in S_N \mid X_{\gamma_1}^{\gamma_1} = x_1^{\gamma_1}, Y_{\gamma_1}^{\gamma_1} \in S_1^{\gamma_1}) \\ = \mathbb{P}(X_{\gamma_N} = x_N, Y_{\gamma_N} \in S_N \mid X_{\gamma_{N-1}} = x_{N-1}).\end{aligned}$$

Proof. Denote with A^c the complement of A in $\mathcal{X} \times \mathcal{S}$, and with $(A^c)^r$ the r -th fold Cartesian product of A^c . Let $B := (X_{\gamma_1}^{\gamma_1} = x_1^{\gamma_1}, Y_{\gamma_1}^{\gamma_1} \in S_1^{\gamma_1})$. Applying Lemma 6.2 in the third equality below, the numerator of the conditional probability on the LHS is

$$\begin{aligned}\mathbb{P}(X_{\gamma_1}^{\gamma_1} = x_1^{\gamma_1}, Y_{\gamma_1}^{\gamma_1} \in S_1^{\gamma_1}) \\ &= \sum_{r \geq 1} \mathbb{P}(\gamma_N = \gamma_{N-1} + r, X_{\gamma_{N-1}+r} = x_N, Y_{\gamma_{N-1}+r} \in S_N \mid B) \mathbb{P}(B) \\ &= \sum_{r \geq 1} \mathbb{P}(X_{\gamma_{N-1}+r} = x_N, Y_{\gamma_{N-1}+r} \in S_N, (X_{\gamma_{N-1}+1}^{\gamma_{N-1}+r-1}, Y_{\gamma_{N-1}+1}^{\gamma_{N-1}+r-1}) \in (A^c)^r \mid B) \mathbb{P}(B) \\ &= \sum_{r \geq 1} \mathbb{P}(X_{\gamma_{N-1}+r} = x_N, Y_{\gamma_{N-1}+r} \in S_N, (X_{\gamma_{N-1}+1}^{\gamma_{N-1}+r-1}, Y_{\gamma_{N-1}+1}^{\gamma_{N-1}+r-1}) \in (A^c)^r \mid X_{\gamma_{N-1}} = x_{N-1}) \mathbb{P}(B) \\ &= \sum_{r \geq 1} \mathbb{P}(\gamma_N = \gamma_{N-1} + r, X_{\gamma_{N-1}+r} = x_N, Y_{\gamma_{N-1}+r} \in S_N \mid X_{\gamma_{N-1}} = x_{N-1}) \mathbb{P}(B) \\ &= \mathbb{P}(X_{\gamma_N} = x_N, Y_{\gamma_N} \in S_N \mid X_{\gamma_{N-1}} = x_{N-1}) \mathbb{P}(B),\end{aligned}$$

and dividing by $\mathbb{P}(B)$ the lemma is proved. \square

Remark 6.6. By the same token, for any $(x_1, S_1), \dots, (x_N, S_N) \in \mathcal{X} \times \mathcal{S}$,

$$\begin{aligned}\mathbb{P}(X_{\gamma_{N+1}} = x_N, Y_{\gamma_{N+1}} \in S_N \mid X_{\gamma_1}^{\gamma_1+1} = x_1^{\gamma_1+1}, Y_{\gamma_1}^{\gamma_1+1} \in S_1^{\gamma_1+1}) \\ = \mathbb{P}(X_{\gamma_{N+1}} = x_N, Y_{\gamma_{N+1}} \in S_N \mid X_{\gamma_{N-1}+1} = x_{N-1}).\end{aligned}$$

Taking $S_1 = \dots = S_N = S$, Remark 6.6 gives

Remark 6.7. For any $x_1, \dots, x_N \in \mathcal{X}$,

$$\mathbb{P}(X_{\gamma_N+1} = x_N \mid X_{\gamma_1+1}^{\gamma_{N-1}+1} = x_1^{N-1}) = \mathbb{P}(X_{\gamma_N+1} = x_N \mid X_{\gamma_{N-1}+1} = x_{N-1}).$$

As a consequence of the conditional independence property of HMMs we have

Lemma 6.8. Let (Y_n) be a HMM with underlying Markov chain (X_n) . Then for any $N \in \mathbb{N}$, for any $x_1, \dots, x_N \in \mathcal{X}$, and $S_1, \dots, S_N \in \mathcal{S}$ such that $\mathbb{P}(X_1^N = x_1^N, Y_1^{N-1} \in S_1^{N-1}) > 0$ we have

$$\mathbb{P}(Y_N \in S_N \mid X_1^N = x_1^N, Y_1^{N-1} \in S_1^{N-1}) = \mathbb{P}(Y_N \in S_N \mid X_N = x_N). \quad (3)$$

Moreover let σ, τ two stopping times for (X_n, Y_n) such that $\sigma < \tau$. Then for any $\bar{S} \in \mathcal{S}^*$, and any $x_1, x_2 \in \mathcal{X}$ we have

$$\mathbb{P}(Y_{\tau+1} \in \bar{S} \mid X_{\tau+1} = x_2) = f_{x_2}(\bar{S}), \quad (4)$$

$$\mathbb{P}(Y_{\tau+1} \in \bar{S} \mid X_{\tau+1} = x_2, X_{\sigma+1} = x_1) = \mathbb{P}(Y_{\tau+1} \in \bar{S} \mid X_{\tau+1} = x_2). \quad (5)$$

Proof. Equation (3) can be easily proved using the conditional independence property. Equation (4) can be seen disintegrating the stopping time $\tau + 1$,

$$\begin{aligned} & \mathbb{P}(Y_{\tau+1} \in \bar{S}, X_{\tau+1} = x) \\ &= \sum_{m \in \mathbb{N}} \mathbb{P}(\tau + 1 = m, Y_m \in \bar{S}, X_m = x) \\ &= \sum_{m \in \mathbb{N}} \mathbb{P}(Y_m \in \bar{S} \mid \tau + 1 = m, X_m = x) \mathbb{P}(\tau + 1 = m, X_m = x) \\ &= \sum_{m \in \mathbb{N}} \mathbb{P}(Y_m \in \bar{S} \mid X_m = x) \mathbb{P}(\tau + 1 = m, X_m = x) \\ &= f_x(\bar{S}) \sum_{m \in \mathbb{N}} \mathbb{P}(\tau + 1 = m, X_m = x) = f_x(\bar{S}) \mathbb{P}(X_{\tau+1} = x), \end{aligned}$$

where the third equality follows by Equation (3), and the result then follows by definition of conditional probability.

To get Equation (5) statement write

$$\begin{aligned}
& \mathbb{P}(Y_{\tau+1} \in \bar{S}, X_{\tau+1} = x_2, X_{\sigma+1} = x_1) \\
&= \sum_{m>n} \sum_{n \in \mathbb{N}} \mathbb{P}(\tau = m, \sigma = n, Y_{m+1} \in \bar{S}, X_{m+1} = x_2, X_{n+1} = x_1) \\
&= \sum_{m>n} \sum_{n \in \mathbb{N}} \mathbb{P}(Y_{m+1} \in \bar{S} \mid \tau = m, \sigma = n, X_{m+1} = x_2, X_{n+1} = x_1) \mathbb{P}(\tau = m, \sigma = n, X_{m+1} = x_2, X_{n+1} = x_1) \\
&= \sum_{m>n} \sum_{n \in \mathbb{N}} \mathbb{P}(Y_{m+1} \in \bar{S} \mid X_{m+1} = x_2) \mathbb{P}(\tau = m, \sigma = n, X_{m+1} = x_2, X_{n+1} = x_1) \\
&= f_{x_2}(\bar{S}) \sum_{m>n} \sum_{n \in \mathbb{N}} \mathbb{P}(\tau = m, \sigma = n, X_{m+1} = x_2, X_{n+1} = x_1) \\
&= f_{x_2}(\bar{S}) \mathbb{P}(X_{\tau+1} = x_2, X_{\sigma+1} = x_1),
\end{aligned}$$

where the third equality follows by Equation (3), and the result follows by the first statement and again by definition of conditional probability. \square

Lemma 6.9. *Let (Y_n) be a HMM with underlying Markov chain (X_n) , and (γ_n) be a sequence of hitting times for A , where $A \subset \mathcal{S}$, then, for any $N \in \mathbb{N}$, and for any $(x_1, S_1) \dots, (x_N, S_N) \in \mathcal{X} \times \mathcal{S}$,*

$$\mathbb{P}(Y_{\gamma_1+1}^{\gamma_{N+1}} \in S_1^N \mid X_{\gamma_1+1}^{\gamma_{N+1}} = x_1^N) = \prod_{k=1}^N \mathbb{P}(Y_{\gamma_k+1} \in S_k \mid X_{\gamma_k+1} = x_k). \quad (6)$$

Proof. Let $C := (Y_{\gamma_1+1}^{\gamma_{N-1}+1} \in S_1^{N-1}, X_{\gamma_1+1}^{\gamma_{N-1}+1} = x_1^{N-1})$ for readability.

$$\begin{aligned}
\mathbb{P}(Y_{\gamma_1+1}^{\gamma_{N+1}} \in S_1^N, X_{\gamma_1+1}^{\gamma_{N+1}} = x_1^N) &= \mathbb{P}(Y_{\gamma_N+1} \in S_N, X_{\gamma_N+1} = x_N \mid C) \mathbb{P}(C) \\
&= \mathbb{P}(Y_{\gamma_N+1} \in S_N, X_{\gamma_N+1} = x_N \mid X_{\gamma_{N-1}+1} = x_{N-1}) \mathbb{P}(C) \\
&= \mathbb{P}(Y_{\gamma_N+1} \in S_N \mid X_{\gamma_N+1} = x_N) \mathbb{P}(X_{\gamma_N+1} = x_N \mid X_{\gamma_{N-1}+1} = x_{N-1}) \mathbb{P}(C) \\
&= \prod_{k=1}^N \mathbb{P}(Y_{\gamma_k+1} \in S_k \mid X_{\gamma_k+1} = x_k) \mathbb{P}(X_{\gamma_1}^{\gamma_{N-1}+1} = x_1^{N-1}),
\end{aligned}$$

where the second equality follows by Remark 6.6, the third by Lemma 6.8, and the last equality follows iterating the procedure and using Remark 6.7. \square

6.2 HMMs and countable mixtures of i.i.d. sequences

The following fact was used in the proof of Theorem 4.2. If the HMM (Y_n) has an underlying Markov chain with block structured transition probability matrix, with identical rows within blocks, then (Y_n) is a countable mixture of i.i.d. sequences.

Consider a Markov chain (X_n) with values in \mathcal{X} and transition matrix P as follows

$$P := \begin{pmatrix} P^1 & 0 & \dots & \dots & 0 & 0 \\ 0 & P^2 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \\ 0 & \dots & \dots & 0 & P^h & 0 \\ 0 & \dots & \dots & \dots & \dots & \ddots \end{pmatrix}, \quad P^h := \begin{pmatrix} p_{c_1^h}^h & p_{c_2^h}^h & \dots & p_{c_{l_h}^h}^h \\ p_{c_1^h}^h & p_{c_2^h}^h & \dots & p_{c_{l_h}^h}^h \\ \vdots & \vdots & \vdots & \vdots \\ p_{c_1^h}^h & p_{c_2^h}^h & \dots & p_{c_{l_h}^h}^h \end{pmatrix}, \quad (7)$$

with $h \in H$, a countable set. The block P^h has size l_h . Some of the p_c^h can be null. The Markov chain (X_n) has clearly H recurrence classes, one for each block, and no transient states. Let us indicate with C_h the h -th recurrence class, corresponding to the states of the h -th block, set $C_h = \{c_1^h, \dots, c_{l_h}^h\}$, where l_h can be infinite. Trivially $\mathcal{X} = \cup_{h \in H} C_h$. An invariant distribution associated with the h -th block is $\mathbf{p}^h := (p_{c_1^h}^h, \dots, p_{c_{l_h}^h}^h)$, and for any sequence $\mu_h > 0$ with $\sum_{h \in H} \mu_h = 1$, the vector

$$\pi = (\mu_1 \mathbf{p}^1, \dots, \mu_h \mathbf{p}^h, \dots) \quad (8)$$

is an invariant distribution for P .

Lemma 6.10. *Consider a HMM (Y_n) where the underlying Markov chain (X_n) has transition matrix P as in (7), invariant measure π as in (8), and assigned read-out distributions $f_x(\bar{S})$, then (Y_n) is a countable mixtures of i.i.d. sequences where \tilde{p} takes values in the set $\{F_h, h \in H\}$, with*

$$F_h(\bar{S}) := p_{c_1^h}^h f_{c_1^h}(\bar{S}) + \dots + p_{c_{l_h}^h}^h f_{c_{l_h}^h}(\bar{S}),$$

and $\mathbb{P}(\tilde{p} = F_h) = \mu_h$.

Proof. Let us compute the finite distributions of (Y_n) . For any $N \in \mathbb{N}$, let $S_0, \dots, S_N \in \mathcal{S}$:

$$\begin{aligned}
\mathbb{P}(Y_0^N \in S_0^N) &= \sum_{x_0^N \in \mathcal{X}} \mathbb{P}(Y_0^N \in S_0^N, X_0^N = x_0^N) \\
&= \sum_{x_0^N \in \mathcal{X}} P(X_0 = x_0) \prod_{n=0}^N \mathbb{P}(Y_n \in S_n \mid X_n = x_n) \prod_{n=1}^N \mathbb{P}(X_n = x_n \mid X_{n-1} = x_{n-1}) \\
&= \sum_{x_0^N \in \mathcal{X}} \pi_{x_0} \prod_{n=0}^N f_{x_n}(S_n) \prod_{n=1}^N P_{x_{n-1}, x_n} \\
&= \sum_{x_0^N \in \mathcal{X}} \pi_{x_0} f_{x_0}(S_0) P_{x_0, x_1} f_{x_1}(S_1) \dots P_{x_{N-1}, x_N} f_{x_N}(S_N) \\
&= \sum_{h \in H} \sum_{x_0^N \in C_h} \mu_h p_{x_0}^h f_{x_0}(S_0) p_{x_1}^h f_{x_1}(S_1) \dots p_{x_N}^h f_{x_N}(S_N) \\
&= \sum_{h \in H} \mu_h \sum_{x_0 \in C_h} p_{x_0}^h f_{x_0}(S_0) \left(\sum_{x_1 \in C_h} p_{x_1}^h f_{x_1}(S_1) \dots \left(\sum_{x_N \in C_h} p_{x_N}^h f_{x_N}(S_N) \right) \right) \\
&= \sum_{h \in H} \mu_h F_h(S_0) F_h(S_1) \dots F_h(S_N),
\end{aligned}$$

where the second equality follows by the HMM properties, the fifth equality follows noting that $P_{x_n, x_{n+1}}$ is null for x_n and x_{n+1} in different recurrence classes, and it is equal to $p_{x_{n+1}}^h$ for x_n and x_{n+1} in the same recurrence class C_h . The expression above coincides with the representation of countable mixtures of i.i.d. sequences given in (2), thus completing the proof. \square

References

- [1] Aldous, D.J. (1985) *Exchangeability and related topics in Ecole d'Été de Probabilités de Saint-Flour XIII - 1983*, Lecture Notes in Mathematics 1117, Springer, Berlin
- [2] de Finetti, B. (1938) *Sur la condition d'équivalence partielle* in *Actualité Scientifiques et Industrielles*, Hermann, Paris, **739**, 5-18
- [3] Dharmadhikari, S.W. (1964) *Exchangeable processes which are function of stationary Markov chains* in *The Annals of Mathematical Statistics*, **35**, 429-430

- [4] Diaconis, P. and Freedman, D. (1980) *de Finetti's theorem for Markov chains* in *The Annals of Probability*, **8**, 115-130
- [5] Diaconis, P. and Freedman, D. (2004) *The Markov Moment Problem and de Finetti's Theorem, Part I and Part II* in *Mathematische Zeitschrift*, **247**, 183-212
- [6] Epifani, I. and Fortini, S. and Ladelli, L. (2002) *A characterization for mixtures of semi-Markov processes*, in *Statistics and Probability Letters* **60**, 445-457
- [7] Finesso, L. and Prosdocimi, C. (2009) *Partially exchangeable hidden Markov models*, in *Proceeding of European Control Conference 2009*, 3910–3914
- [8] Fortini, S., Ladelli, L., Petris, G. and Regazzini, E. (2002) *On mixtures of distribution of Markov chains* in *Stochastic Processes and their Applications*, **100**, 147-165
- [9] Kallenberg, O. (2005) in *Probabilistic Symmetries and Invariance Principles* , Springer
- [10] Prosdocimi, C. (2010) *Partial exchangeability and change detection for hidden Markov models*, in *PhD Dissertation, cycle XXII, University of Padova*
- [11] Vidyasagar, M. (2011) *The complete realization problem for hidden Markov models: A survey and some new results* in *Mathematics of Control, Signals and Systems*, **23** (1), 1-65