Three proofs of the Casas-Alvero conjecture

Luis J. Fernández de las Heras^{a,*}, María J. Fernández de las Heras^b

 ^aDpto. de Matemática Aplicada, E.T.S. de Ingenieros Industriales, Universidad Politécnica de Madrid, José G. Abascal 2, 28006 Madrid, Spain.
 ^bCSIC, Centro de Acústica Aplicada y Evaluación no Destructiva, Serrano 144, 28006 Madrid. Spain.

Abstract

The Casas-Alvero conjecture claims that a complex univariate polynomial having roots in common with each of its derivatives must be a power of a linear polynomial. Up to now, only partial proofs and numerical evidences have been presented. In this paper we give three different proofs of the conjecture.

Keywords: Polynomial interpolation, univariate polynomials.

2010 MSC: 41A05, 30E05

1. Introduction

This paper is concerned with the following question posed by E. Casas-Alvero more than a decade ago.

Casas-Alvero Conjecture. Let f be a monic complex polynomial of degree n in a single variable z. Suppose that $gcd(f, f^{(k)}) \neq 1$ for $k = 1, \ldots, n-1$, where $f^{(k)}$ denotes the k-th derivative of f. Then, there exists a constant $a \in \mathbb{C}$ such that $f(z) = (z-a)^n$.

It may be proven that if the conjecture is true over $\mathbb C$ then it is true over all fields of characteristic 0. In contrast, the conjecture is not true in prime characteristic.

In [2] the Casas-Alvero conjecture was proven for polynomials of degree n less than or equal to 7. The conjecture has also been proven for infinitely many values of n, see [1].

^{*}Corresponding author. Fax: +34 91 336 3001. Phone: +34 91 336 3106 Email addresses: lfernandez@etsii.upm.es (Luis J. Fernández de las Heras), mjose@caend.upm-csic.es (María J. Fernández de las Heras)

We can rewrite the conjecture in terms of interpolation polynomials on the complex plane \mathbb{C} , see [3, 4].

Theorem 1.1. Let z_1, z_2, \ldots, z_n be n complex numbers and let p be a monic complex polynomial of degree n. Suppose that the polynomial p satisfies

$$p(z_k) = 0, k = 1, ..., n;$$

 $p^{(k)}(z_{k+1}) = 0, k = 1, ..., n - 1.$

Then

$$p(z) = (z - a)^n, \qquad a \in \mathbb{C}.$$

In the sequel any monic polynomial satisfying the conditions of Theorem 1.1 will be called a Casas-Alvero polynomial.

2. Birkhoff interpolation

First, we pose the following problems which may be considered as particular cases of Birkhoff or lacunary interpolation problems [5]. They are closely related to the Casas-Alvero conjecture.

Problem 2.1. Given n complex numbers $\alpha_1, \ldots \alpha_n$, find all monic polynomials p that satisfy

$$p^{(k)}(\alpha_{k+1}) = 0, \qquad k = 0, 1, \dots, n-1.$$

Solution. A monic polynomial p_n of degree n can be expressed in the form

$$p_n(z) = \sum_{j=0}^n a_j z^j, \tag{1}$$

where $a_n = 1$. For k = 1, ..., n, its k-th derivative is given by

$$p_n^{(k)}(z) = \sum_{j=k}^n a_j \frac{j!}{(j-k)!} z^{j-k}.$$

If we evaluate the above expressions at the nodes, we obtain

$$p_n^{(k)}(\alpha_{k+1}) = \sum_{j=k}^n a_j \frac{j!}{(j-k)!} \alpha_{k+1}^{j-k} = 0, \quad k = 0, 1, \dots, n-1.$$

We can write the above equations in matrix form as

$$A \cdot \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \\ a_n \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ n! \end{pmatrix},$$

where

$$A = \begin{pmatrix} 1 & \alpha_1 & \alpha_1^2 & \cdots & \alpha_1^{n-2} & \alpha_1^{n-1} & \alpha_1^n \\ 0 & 1 & 2\alpha_2 & \cdots & \frac{n-2}{1}\alpha_2^{n-3} & \frac{n-1}{1}\alpha_2^{n-2} & n\alpha_2^{n-1} \\ 0 & 0 & 2! & \cdots & \frac{(n-2)!}{(n-4)!}\alpha_3^{n-4} & \frac{(n-1)!}{(n-3)!}\alpha_3^{n-3} & \frac{n!}{(n-2)!}\alpha_3^{n-2} \\ \vdots & \vdots & \vdots & \cdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & (n-2)! & \frac{(n-1)!}{1}\alpha_{n-1} & \frac{n!}{2!}\alpha_{n-1}^2 \\ 0 & 0 & 0 & \cdots & 0 & (n-1)! & n!\alpha_n \\ 0 & 0 & 0 & \cdots & 0 & 0 & n! \end{pmatrix}.$$

Clearly the interpolation problem has a solution and it is unique, because the matrix is of full rank. The solution of the system is trivial but laborious. The incidence matrix will have as many rows as the number of different nodes. In the extreme case that there is only one node, the problem is of Hermite type. As in all cases the matrix is triangular, there is always a unique solution.

A straightforward calculation shows that the unique solution p_n of Problem 2.1 admits the following integral representation

$$p_n(z) = n! \int_{\alpha_1}^z dx_1 \int_{\alpha_2}^{x_1} dx_2 \dots \int_{\alpha_{n-1}}^{x_{n-2}} dx_{n-1} \int_{\alpha_n}^{x_{n-1}} dx_n.$$
 (2)

We can study now the inverse problem.

Problem 2.2. Given n complex numbers z_1, z_2, \ldots, z_n , let

$$\hat{z} = (z_1, \dots, z_n)$$
 and $p(z) = \prod_{k=1}^{n} (z - z_k)$.

Find all the vectors of complex numbers

$$\alpha = (\alpha_1, \alpha_2, ..., \alpha_n)$$

$$p^{(k)}(\alpha_{k+1}) = 0, \qquad k = 0, 1, ..., n-1.$$

Solution. The problem has a solution. We begin with the last equation of the system given by Problem 2.1 to obtain α_n . Then we solve the (n-1)-th equation to get the two values of α_{n-1} and we continue until considering the first equation of degree n which allows us to obtain the n values of α_1 .

The problem has a solution but the solution is not unique. The problem may have until n! different solutions. Of course, it is possible to approximate the solutions by numerical methods, but it should be noticed that this problem is invariant under permutation of the symmetric group of the n components of \hat{z} . If we change the order of the roots in \hat{z} , the polynomial p does not change and the solutions of the problem are the same ones as before. The number of solutions is just equal to the number of permutations of \hat{z} .

We can make Problem 2.2 more complicated considering additional conditions like requiring the nodes of interpolation α to be equal to the zeros of the polynomial p. Thus, the polynomial p and its derivatives will have common zeros.

3. The first proof

There are several different ways to tackle the proof. We have chosen to consider it as a problem of the same type as that of Problem 2.2. That is, assuming that the roots of the polynomial are known, find the nodes of interpolation. Subsequently, impose the condition that the roots coincide with the nodes. Then this problem is invariant under permutations of the zeros of the polynomial as in Problem 2.2. But if we permute the zeros of the polynomial, the interpolation nodes are automatically interchanged. Thus the polynomial and its derivatives up to order n-1 must take the value zero at all the zeros of the polynomial. Therefore, all the zeros are of multiplicity n. But this is possible only if there is a single zero of multiplicity n. Thus, the Casas-Alvero conjecture is proven.

4. A second proof by induction

Taking account of the representation (2), for n = 2, the Casas-Alvero polynomial is

$$p_2(z) = 2! \int_{\alpha_1}^z dx_1 \int_{\alpha_2}^{x_1} dx_2 = 2! \int_{\alpha_2}^z dx_1 \int_{\alpha_2}^{x_1} dx_2$$
$$= (z - \alpha_2)^2 - (\alpha_1 - \alpha_2)^2 = (z - \alpha_2)^2 = (z - \alpha_1)^2.$$

Now suppose that, for $k=1,\ldots,n-1,$ any Casas-Alvero polynomial is of the form

$$p_k(z) = (z - \alpha_1)^k.$$

Then

$$p_n(z) = n! \int_{\alpha_1}^z dx_1 \int_{\alpha_2}^{x_1} dx_2 \cdots \int_{\alpha_{n-1}}^{x_{n-2}} dx_{n-1} \int_{\alpha_n}^{x_{n-1}} dx_n.$$

Notice that $p_n(z)$ takes the value zero at $z = \alpha_k$ from k = 1 to n. So, for $k = 1, \ldots, n$, we have

$$p_n(z) = n! \int_{\alpha_k}^z dx_1 \int_{\alpha_2}^{x_1} dx_2 \cdots \int_{\alpha_{n-1}}^{x_{n-2}} dx_{n-1} \int_{\alpha_n}^{x_{n-1}} dx_n.$$

Therefore

$$p_{n}(z) = n! \int_{\alpha_{k}}^{z} dx_{1} \int_{\alpha_{2}}^{x_{1}} dx_{2} \cdots \int_{\alpha_{n-1}}^{x_{n-2}} (x_{n-1} - \alpha_{n}) dx_{n-1}$$
$$= n! \int_{\alpha_{k}}^{z} dx_{1} \int_{\alpha_{2}}^{x_{1}} dx_{2} \cdots \int_{\alpha_{n-1}}^{x_{n-2}} x_{n-1} dx_{n-1} - n\alpha_{n} p_{n-1}(z),$$

for k = 1, ..., n, where p_{n-1} is a Casas-Alvero polynomial of degree n-1. Then

$$p_{n-1}(z) = (z - \alpha_1)^{n-1}$$

and

$$\alpha_1 = \alpha_2 = \cdots = \alpha_{n-1}$$
.

Therefore

$$p_{n}(z) = n! \int_{\alpha_{1}}^{z} dx_{1} \int_{\alpha_{1}}^{x_{1}} dx_{2} \cdots \int_{\alpha_{1}}^{x_{n-3}} \frac{1}{2} (x_{n-2}^{2} - \alpha_{1}^{2}) dx_{n-2} - n\alpha_{n} p_{n-1}(z)$$

$$= n! \int_{\alpha_{1}}^{z} dx_{1} \int_{\alpha_{1}}^{x_{1}} dx_{2} \cdots \int_{\alpha_{1}}^{x_{n-3}} \frac{1}{2!} x_{n-2}^{2} dx_{n-2}$$

$$- \frac{n(n-1)}{2} \alpha_{1}^{2} p_{n-2}(z) - n\alpha_{n} p_{n-1}(z),$$

where $p_{n-2}(z) = (z - \alpha_1)^{n-2}$.

If we continue the process of calculating the iterated integral, we arrived at

$$p_{n}(z) = z^{n} - \alpha_{1}^{n} - \sum_{k=1}^{n-2} {n \choose k} \alpha_{1}^{n-k} (z - \alpha_{1})^{k} - n\alpha_{n} p_{n-1}(z)$$

$$= z^{n} - \sum_{k=0}^{n-1} {n \choose k} \alpha_{1}^{n-k} (z - \alpha_{1})^{k} + n(\alpha_{1} - \alpha_{n})(z - \alpha_{1})^{n-1}$$

$$= z^{n} + (z - \alpha_{1})^{n} - \sum_{k=0}^{n} {n \choose k} \alpha_{1}^{n-k} (z - \alpha_{1})^{k}$$

$$+ n(\alpha_{1} - \alpha_{n})(z - \alpha_{1})^{n-1}$$

$$= (z - \alpha_{1})^{n} + n(\alpha_{1} - \alpha_{n})(z - \alpha_{1})^{n-1},$$

since the number α_n is a root of the polynomial p_n . Actually

$$p_n(\alpha_n) = (\alpha_n - \alpha_1)^n + n(\alpha_1 - \alpha_n)(\alpha_n - \alpha_1)^{n-1}$$

= $-(n-1)(\alpha_n - \alpha_1)^n = 0.$

Then, $\alpha_n = \alpha_1$ and $p_n(z) = (z - \alpha_1)^n$, as we wanted to prove.

5. The third proof

Although we think that the previous two proofs are correct, we encourage the reader to continue reading the article.

Lemma 5.1. Every Casas-Alvero polynomial has a root equal to the geometric center of gravity of its roots.

Proof. Let p_n be a monic polynomial of degree n expressed in the form (1). Then

$$p_n^{(n-1)}(z) = n! z + (n-1)! a_{n-1}.$$

Evaluating at the zero z_n of the polynomial $p_n^{(n-1)}$, we obtain

$$p_n^{(n-1)}(z_n) = n!z_n + (n-1)!a_{n-1} = 0,$$

or, equivalently,

$$z_n = -\frac{a_{n-1}}{n} = \frac{1}{n} \sum_{j=1}^n z_j.$$

Now, let p_n be a monic polynomial of degree n. The polynomial p_n may be written in the form

$$p_n(z) = \sum_{k=0}^{n} \binom{n}{k} (-1)^k c_k z^{n-k},$$
 (3)

where

$$c_0 = 1,$$
 $c_1 = \binom{n}{1}^{-1} \sum_{j=1}^n z_j,$ $c_k = \binom{n}{k}^{-1} \sum_{j_1+j_2+\dots+j_n=k}^{j_i=0,1} \left(\prod_{i=1}^n z_i^{j_i}\right), \quad k = 2,\dots,n.$

Taking derivatives in (3), we obtain

$$p'_{n}(z) = \sum_{k=0}^{n-1} {n \choose k} (-1)^{k} c_{k} (n-k) z^{n-k-1}$$

$$= n \sum_{k=0}^{n-1} {n \choose k} (-1)^{k} c_{k} \frac{n-k}{n} z^{n-k-1}$$

$$= n \sum_{k=0}^{n-1} {n-1 \choose k} (-1)^{k} c_{k} z^{n-k-1}.$$

The above calculation indicates that the zeros of the derivative of a polynomial of degree n and the zeros of this polynomial have in common the following quantities: the average of their zeros, the mean double product of their zeros, and so forth until the average (n-1) product of their zeros.

If p_n is a Casas-Alvero polynomial, then, for k = 1, ..., n - 1, the polynomial $p_n^{(k)}$ has the same coefficients c_j , j = 1, ..., n - k, as p_n .

For k = n - 1, the zero shared by the polynomial and its derivative of order n - 1 is precisely c_1 and this determines the coefficient of degree n - 1 of the polynomial p_n .

For k = n - 2, which is the zero shared by the polynomial and the derivative of order n - 2? The two zeros determine the same value for c_2 .

Therefore, whatever zero p_n shares with its derivative we obtain same result for the polynomial. But this reasoning is valid for all the derivatives of p_n . Then all the zeros of the successive derivatives of p_n must be zeros of p_n , therefore all the zeros must be equal.

6. Concluding remarks

We conclude that the Casas-Alvero conjecture is true. In connection with the work carried out in this paper, we are currently studying an optimization problem in which symmetries play an important role. The results related to this problem will appear elsewhere.

In our opinion solving a math problem is not to close the door but rather open a window to new challenges. For this reason, we propose a degenerated Birkhoff interpolation problem in which the number of equations is greater than the number of unknowns. Among the different possibilities we have chosen the problem stated below. Find the conditions to be met by the interpolation nodes and their values for the following problem to be solvable.

Problem 6.1. Given 2n complex numbers $\alpha_1, \ldots, \alpha_n$ and c_0, \ldots, c_{n-1} , find all the polynomials p of degree n that satisfy

$$p(\alpha_{k+1}) = p^{(k)}(\alpha_{k+1}) = c_k, \qquad k = 0, 1, \dots, n-1.$$

References

- [1] H.-C. Graf von Bothmer, O. Labs, J. Schicho, C. van de Woestijne, The Casas-Alvero conjecture for infinitely many degrees, J. Algebra 316 (2007) 224–230.
- [2] G. Diaz-Toca, L. González-Vega, On a conjecture about univariate polynomials and their roots, In: Algorithmic Algebra and Logic 2005, Norderstedt, Germany, 2005, pp. 83–90.
- [3] J. Draisma, J.P. de Jong, On the Casas-Alvero conjecture, Eur. Math. Soc. Newsl. 80 (2011) 29–33.
- [4] E. Casas-Alvero, Singularities of plane curves, London Mathematical Society Lecture Note Series 276, Cambridge University Press, Cambridge, 2000.
- [5] G.G. Lorentz, K. Jetter, S.D. Riemenschneider, Birkhoff Interpolation, Addison-Wesley, Reading, Mass., 1982.