

Symmetry Classification of quasi-linear PDE's.

II: an exceptional case

Giampaolo Cicogna

*Dip. di Fisica "E.Fermi", Università di Pisa and INFN, Sez. di Pisa
Largo B. Pontecorvo 3, Ed. B-C, I-56127, Pisa, Italy
(fax: +39-050-2214887; e-mail: cicogna@df.unipi.it)*

Abstract

This short note completes the symmetry analysis of a class of quasi-linear partial differential equations considered in the previous paper (Nonlinear Dynamics **51**, 309–316 (2008)): it deals with the presence of an “exceptional” Lie point symmetry, not previously examined, which is admitted only if the involved parameters are fixed by precise relationships. The peculiarity of this symmetry is enhanced by the fact that it leads to a solution relevant in the theory of plasma physics, and also related to the presence of a nontrivial example of a conditional symmetry of weak type.

Key words: symmetry classification; quasi-linear PDE's; Grad-Schlüter-Shafranov equation; weak conditional symmetry

Running title: Symmetry Classification II: an exceptional case

This short note is an Addendum to my previous paper [1], and completes the “symmetry classification” of a class of quasi-linear partial differential equations. It provides the case of an “exceptional” symmetry, not previously considered, and a related solution which is relevant in plasma physics.

One of the examples examined in [1] (Sect. 5) concerned equations of the form (for the unknown function $u = u(x, y)$)

$$u_{xx} + u_{yy} + \frac{a}{x}u_x = \alpha(x, y) F_1(u) + F_2(u) \quad (a = \text{const} \neq 0) \quad (1)$$

where α, F_1, F_2 are given. Equations of this type include several interesting cases in mathematical physics, as reduced forms of wave equation, heat and

Schrödinger equation, and – more specifically, as we shall see – an important equation in plasma physics.

In [1] it was been shown that equation (1) may admit Lie point symmetries only if both functions F_1, F_2 have the form $F_i \propto u^{c_i}$ or $F_i \propto \exp(c_i u)$, where c_i are constants. Even in these cases, in general, only nearly trivial symmetries (scalings and translations) are admitted.

There is actually an “exception”, not examined in [1], which occurs when

$$\alpha(x, y) = x^r \quad , \quad F_1(u) = \gamma_1 u^{c_1} \quad , \quad F_2(u) = \gamma_2 u^{c_2} \quad (2)$$

where $r, c_1, c_2, \gamma_1, \gamma_2$ are constants, and if the two following special relations hold

$$c_1 = 1 + 2(r + 2)/a \quad , \quad c_2 = 1 + (4/a) \quad . \quad (3)$$

In this case indeed the new nontrivial symmetry generated by the vector field

$$X = 2xy \frac{\partial}{\partial x} + (y^2 - x^2) \frac{\partial}{\partial y} - ayu \frac{\partial}{\partial u} \quad (4)$$

is admitted, as it can be easily checked.

The presence of this symmetry implies that if $u = u(x, y)$ is any solution of Eq. (1) with the choice (2) and (3), then also $\tilde{u} = \tilde{u}(x, y)$ defined by

$$\tilde{u}(x, y) = [C(x, y, \lambda)]^{-a/2} u(\tilde{x}(x, y, \lambda), \tilde{y}(x, y, \lambda)) \quad (5)$$

where

$$C(x, y, \lambda) = 1 + \lambda^2(x^2 + y^2) + 2\lambda y$$

$$\tilde{x} = x [C(x, y, \lambda)]^{-1} \quad \text{and} \quad \tilde{y} = [y + \lambda(x^2 + y^2)] [C(x, y, \lambda)]^{-1} \quad (5')$$

solves the equation for any $\lambda \in \mathbf{R}$.

More specifically, let me now fix $r = 2$ and then consider the equation

$$u_{xx} + u_{yy} + \frac{a}{x} u_x = \gamma_1 x^2 u^{1+(8/a)} + \gamma_2 u^{1+(4/a)} \quad (6)$$

according to (3). The choice $r = 2$ is relevant not only because the resulting equation has the same form as the Grad-Schlüter-Shafranov (also called Bragg–Hawthorne) equation, well known in magnetohydrodynamics and plasma physics (see e.g. [2] and also [3, 4], in this case one has $a = -1$), but also because this equation admits an interesting family of solutions, thanks to the exceptional symmetry given above, and to the following property.

It is easy to verify that the vector field

$$Y = y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} \quad (7)$$

generates neither a standard (exact) symmetry nor a (properly defined) conditional symmetry (see e.g. [5, 6, 7, 8]) for Eq. (6). Indeed, introducing as usual the additional Y -invariance condition expressed by the first-order equation

$$\Delta^{(1)} \equiv Y u = 0 \quad (8)$$

and writing Eq. (6) as $\Delta = 0$, one has not only $(\Pr^{(2)}Y)\Delta|_{\Delta=0} \neq 0$ but also

$$(\Pr^{(2)}Y)\Delta|_{\Delta=\Delta^{(1)}=0} \neq 0$$

where as usual $\Pr^{(2)}Y$ denotes the second prolongation of Y . However, putting $(\Pr^{(2)}Y)\Delta \equiv \tilde{\Delta}$, and imposing the new condition $\tilde{\Delta} = 0$, it can be verified that

$$(\Pr^{(2)}Y)\Delta|_{\Delta=\Delta^{(1)}=\tilde{\Delta}=0} = 0$$

is satisfied. In this sense (see [9, 10, 11]), I say that Y is a *weak* conditional symmetry for Eq. (6). Notice that a reduction of similar type was already considered [12, 13], with the name of “anti-reduction”. Accordingly, a reduction procedure for Eq. (6) is possible. To this purpose, one can directly and more simply rewrite this equation in terms of the Y -invariant variable $s = x^2 - y^2$ (which solves Eq. (8)). Writing $v = v(s) = u(x^2 - y^2)$, Eq. (6) becomes

$$-4sv_{ss} + 2av_s + 8x^2v_{ss} = \gamma_1 x^2 v^{1+(8/a)} + \gamma_2 v^{1+(4/a)} .$$

The presence of the term x^2 in this equation confirms that Y is not a proper conditional symmetry, but the two separated equations

$$-4sv_{ss} + 2av_s = \gamma_2 v^{1+(4/a)} \quad \text{and} \quad 8v_{ss} = \gamma_1 v^{1+(8/a)}$$

admit a common solution $v = s^{-a/4}$, i.e.

$$u(x, y) = (x^2 - y^2)^{-a/4} \quad (9)$$

with

$$\gamma_1 = (a/2)(a + 4) \quad \gamma_2 = -(a/4)(3a + 4) .$$

The most interesting situation occurs when $a \neq \pm 4, \pm 8, \dots$: in this case the solution (9) exists only in the region $|x| \geq |y|$.

It can be noticed that equation (6) admits also one of the trivial symmetries mentioned at the beginning of this paper (and discussed in [1]), precisely the scaling

$$X' = x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} - \frac{a}{2} u \frac{\partial}{\partial u}$$

and – in addition – that the above solution (9) turns out to be invariant under this symmetry (see also [4] for the reduction of Eq. (1) obtained thanks to the symmetry X'). In other words, one can say that solution (9) exhibits the peculiar property of being simultaneously invariant under the “standard” symmetry X' and the weak conditional one Y .

Taking now advantage from the “exceptional” symmetry X [Eq.(4)] which is present in this case, we can construct from the above solution (9) a continuous family of solutions to Eq. (6) using the rule (5-5'): we obtain

$$u(x, y) = [x^2 - (y + \lambda(x^2 + y^2))^2]^{-a/4} \quad (10)$$

which holds only in the interior of the two circles centered resp. in $x_0 = 1/(2\lambda)$, $y_0 = -1/(2\lambda)$ and $x_0 = -1/(2\lambda)$, $y_0 = -1/(2\lambda)$, both of radius $1/(\sqrt{2}\lambda)$, and excluding their intersection (it is not restrictive to assume $\lambda > 0$).

A solution of this form is relevant in the context of plasma physics because it describes a specially interesting “double-D” shaped toroidal plasma equilibrium which exhibits a rather flat plasma profile with strong pressure gradients and current gradients at the edge (see [4] for some comment on this point). This is another aspect that enhances the exceptional character of the symmetry X considered in the present note and the interest in it.

Acknowledgments. It is a pleasure to thank the referees for their valuable remarks, and in particular one of them for pointing out to me the two references [12-13].

References

- [1] Cicogna, G.: Symmetry classification of quasi-linear PDE's containing arbitrary functions, *Nonlinear Dynamics* **51**, 309–316 (2008)
- [2] Wesson, J.: *Tokamaks*, The Oxford Engineering Ser. 48, Oxford, Clarendon (1997)
- [3] Frewer, M., Oberlack, M., Guenther, S.: Symmetry investigations on the incompressible stationary axisymmetric Euler equations with swirl, *Fluid Dyn. Research* **39**, 647–664 (2007)
- [4] Cicogna, G., Pegoraro, F., Ceccherini, F.: Symmetries, weak symmetries and related solutions of the Grad-Shafranov equation, *Phys. of Plasmas* **17**, 102506-1/8 (2010)
- [5] Bluman, G.W. and Cole, J.D.: The general similarity solution of the heat equation, *J. Math. Mech.* **18**, 1025–1042 (1969)
- [6] Levi, D. and Winternitz, P.: Non-classical symmetry reduction: example of the Boussinesq equation, *J. Phys. A* **22**, 2915–2924 (1989)

- [7] Fushchych, W.I. (Ed.), *Symmetry analysis of equations of mathematical physics*, Inst. of Math. Acad. of Science of Ukraine, Kiev (1992), and *Conditional symmetries of the equations of mathematical physics*, in Modern group analysis: advanced analytical and computational methods in mathematical physics, (Ibragimov N.H., Torrisi M. and Valenti A., Eds.), Kluwer, Dordrecht (1993)
- [8] Kunzinger, M. and Popovych, R.O.: Is a nonclassical symmetry a symmetry?, Proc. 4th Workshop Group Analysis of Differential Equations & Integrable systems, pp. 107–120, Cyprus (2008); arXiv:0903.0821 (2009)
- [9] Olver, P.J. and Rosenau, Ph.: The construction of special solutions to partial differential equations, Phys. Lett. A **114**, 107–112 (1986), and Group-invariant solutions of differential equations, SIAM J. Appl. Math. **47**, 263–278 (1986)
- [10] Cicogna, G.: A discussion on the different notions of symmetry of differential equations, Proc. Inst. Math. N.A.S. Ukr. **50**, 77–84, 2004
- [11] Cicogna, G. and Laino, M.: On the notion of conditional symmetry of differential equations, Rev. Math. Phys. **18**, 1–18 (2006)
- [12] Fushchych, W.I.: Conditional symmetry of the equations of nonlinear mathematical physics, Ukrainian Math. J. **43**, No. 11, 1350–1364 (1991)
- [13] Fushchych, W.I. and Zhdanov, R.Z.: Anti-reduction of the nonlinear wave equation, Reports of N.A.S. of Ukraine **11**, 37–41 (1993), translated in: Fushchych, W.I., Scientific Works **5**, 116–119 (2003).