

Inequalities for the Jacobian Elliptic Functions with Complex Modulus*

Klaus Schiefermayr[†]

Abstract

Despite the fact that there is a huge amount on papers and books devoted to the theory of Jacobian elliptic functions, very little is known when the modulus k of these functions lies outside the unit interval $[0, 1]$. In this note, we prove some simple inequalities for the absolute value of Jacobian elliptic functions with complex modulus.

Mathematics Subject Classification (2000): 33E05

Keywords: Inequality, Jacobian elliptic function

1 Introduction and Main Result

Consider the Jacobian elliptic functions $\operatorname{sn}(z, k)$, $\operatorname{cn}(z, k)$, $\operatorname{dn}(z, k)$ with complex parameter (modulus) $k \in \mathbb{C}$. Starting with the works of Jacobi in the 1820's until now, there exists a huge amount on papers and books devoted to the theory of Jacobian elliptic functions, see, e.g., [3], [2], or [6]. Almost all of these contributions are restricted to the case when the modulus k is in the unit interval $[0, 1]$. Exceptions are, e.g., the articles of Walker [4], [5]. Since the functions $\operatorname{sn}(z, k)$, $\operatorname{cn}(z, k)$, and $\operatorname{dn}(z, k)$ depend on k^2 rather than k , we shall use $m = k^2$ as parameter but use the same notation $\operatorname{sn}(z, m)$, $\operatorname{cn}(z, m)$, and $\operatorname{dn}(z, m)$.

There seems to exist no estimates for the absolute value of $\operatorname{sn}(z, m)$, $\operatorname{cn}(z, m)$, and $\operatorname{dn}(z, m)$ in terms of elementary functions. In this paper, we give such estimates.

Theorem 1. *Let $m \in \mathbb{C}$, $|m| \leq 1$, and $z \in \mathbb{C}$, $|z| \leq R < \frac{\pi}{2}$. Then the inequalities*

$$|\operatorname{sn}(z, m)| \leq \frac{\operatorname{sn}(|z|, m_1)}{\operatorname{cn}(|z|, m_1)} \leq \tan |z|, \quad (1)$$

$$|\operatorname{cn}(z, m)| \leq \frac{1}{\operatorname{cn}(|z|, m_1)} \leq \frac{1}{\cos |z|}, \quad (2)$$

$$|\operatorname{dn}(z, m)| \leq \frac{\operatorname{dn}(|z|, m_1)}{\operatorname{cn}(|z|, m_1)} \leq \frac{1}{\cos |z|}, \quad (3)$$

hold, where $m_1 := 1 - |m| \in [0, 1]$. For all inequalities we have equality if $z = 0$ or if $z = iy$, $-R \leq y \leq R$, and $m = 1$. Further, in the first inequalities of (1)–(3), equality is attained if $z = iy$, $-R \leq y \leq R$, and $m = |m|$.

*published in: Journal of Mathematical Inequalities **6** (2012), 91–94.

[†]University of Applied Sciences Upper Austria, School of Engineering and Environmental Sciences, Stelzhamerstrasse 23, 4600 Wels, Austria, KLAUS.SCHIEFERMAYR@FH-WELS.AT

2 Proof

Proof of Theorem 1. Starting point for the proof are the Taylor expansions of $\operatorname{sn}(z, m)$, $\operatorname{cn}(z, m)$, and $\operatorname{dn}(z, m)$, respectively, given by [4, Eq. (3.6)]

$$\operatorname{sn}(z, m) = \sum_{n=0}^{\infty} (-1)^n s_n(m) \frac{z^{2n+1}}{(2n+1)!}, \quad (4)$$

$$\operatorname{cn}(z, m) = \sum_{n=0}^{\infty} (-1)^n c_n(m) \frac{z^{2n}}{(2n)!}, \quad (5)$$

$$\operatorname{dn}(z, m) = \sum_{n=0}^{\infty} (-1)^n d_n(m) \frac{z^{2n}}{(2n)!}, \quad (6)$$

where $s_n(m)$, $c_n(m)$, and $d_n(m)$ are polynomials with positive integer coefficients. Thus, we have the inequalities

$$|s_n(m)| \leq s_n(|m|) \leq s_n(1), \quad (7)$$

$$|c_n(m)| \leq c_n(|m|) \leq c_n(1), \quad (8)$$

$$|d_n(m)| \leq d_n(|m|) \leq d_n(1). \quad (9)$$

Note that each of the power series (7)–(9) are absolutely convergent for $|m| \leq 1$ and $|z| < \frac{\pi}{2}$, see [4, Thm. 3.2]. Especially,

$$\operatorname{sn}(iz, m) = i \sum_{n=0}^{\infty} s_n(m) \frac{z^{2n+1}}{(2n+1)!}, \quad (10)$$

$$\operatorname{cn}(iz, m) = \sum_{n=0}^{\infty} c_n(m) \frac{z^{2n}}{(2n)!}, \quad (11)$$

$$\operatorname{dn}(iz, m) = \sum_{n=0}^{\infty} d_n(m) \frac{z^{2n}}{(2n)!}. \quad (12)$$

If we put $m = 1$ in formulae (10)–(12), we get (note that $c_n(1) = d_n(1)$)

$$\tanh(iz) = i \sum_{n=0}^{\infty} s_n(1) \frac{z^{2n+1}}{(2n+1)!}, \quad (13)$$

$$\frac{1}{\cosh(iz)} = \sum_{n=0}^{\infty} c_n(1) \frac{z^{2n}}{(2n)!} = \sum_{n=0}^{\infty} d_n(1) \frac{z^{2n}}{(2n)!}. \quad (14)$$

Hence, the inequalities

$$\begin{aligned} |\operatorname{sn}(z, m)| &\leq \sum_{n=0}^{\infty} |s_n(m)| \frac{|z|^{2n+1}}{(2n+1)!} \quad \text{by (4)} \\ &\leq \sum_{n=0}^{\infty} s_n(|m|) \frac{|z|^{2n+1}}{(2n+1)!} =: (*_1) \quad \text{by (7)} \\ &= \frac{1}{i} \operatorname{sn}(i|z|, |m|) \quad \text{by (10)} \\ &= \frac{\operatorname{sn}(|z|, 1 - |m|)}{\operatorname{cn}(|z|, 1 - |m|)} \quad \text{by [2, Eq. (2.6.12)]} \end{aligned}$$

and

$$\begin{aligned}
 |\operatorname{cn}(z, m)| &\leq \sum_{n=0}^{\infty} |c_n(m)| \frac{|z|^{2n}}{(2n)!} && \text{by (5)} \\
 &\leq \sum_{n=0}^{\infty} c_n(|m|) \frac{|z|^{2n}}{(2n)!} =: (*_2) && \text{by (8)} \\
 &= \operatorname{cn}(i|z|, |m|) && \text{by (11)} \\
 &= \frac{1}{\operatorname{cn}(|z|, 1 - |m|)} && \text{by [2, Eq. (2.6.12)]}
 \end{aligned}$$

and

$$\begin{aligned}
 |\operatorname{dn}(z, m)| &\leq \sum_{n=0}^{\infty} |d_n(m)| \frac{|z|^{2n}}{(2n)!} && \text{by (6)} \\
 &\leq \sum_{n=0}^{\infty} d_n(|m|) \frac{|z|^{2n}}{(2n)!} =: (*_3) && \text{by (9)} \\
 &= \operatorname{dn}(i|z|, |m|) && \text{by (12)} \\
 &= \frac{\operatorname{dn}(|z|, 1 - |m|)}{\operatorname{cn}(|z|, 1 - |m|)} && \text{by [2, Eq. (2.6.12)]}
 \end{aligned}$$

hold and we have proved the first inequalities of (1)–(3), respectively. Analogously,

$$\begin{aligned}
 (*_1) &\leq \sum_{n=0}^{\infty} s_n(1) \frac{|z|^{2n+1}}{(2n+1)!} && \text{by (8)} \\
 &= \frac{1}{i} \tanh(i|z|) && \text{by (13)} \\
 &= \tan |z|
 \end{aligned}$$

and

$$\begin{aligned}
 (*_2), (*_3) &\leq \sum_{n=0}^{\infty} c_n(1) \frac{|z|^{2n}}{(2n)!} && \text{by (8)} \\
 &= \frac{1}{\cosh(i|z|)} && \text{by (14)} \\
 &= \frac{1}{\cos |z|}
 \end{aligned}$$

holds. □

Remark. Another (much more complicated) possibility for proving the second inequalities of (1)–(3), respectively, is the following: For fixed $u \in [0, \frac{\pi}{2}]$, consider the functions

$$f_1(m_1) := \frac{\operatorname{sn}(u, m_1)}{\operatorname{cn}(u, m_1)}, \quad f_2(m_1) := \frac{1}{\operatorname{cn}(u, m_1)}, \quad f_3(m_1) := \frac{\operatorname{dn}(u, m_1)}{\operatorname{cn}(u, m_1)}$$

for $0 \leq m_1 \leq 1$. Then one has to prove that these functions are strictly monotone decreasing in m_1 , i.e. $f'_j(m_1) < 0$, $j = 1, 2, 3$, using formulae (710.54), (710.57), and (710.60) of [1].

References

- [1] P.F. Byrd and M.D. Friedman, *Handbook of elliptic integrals for engineers and scientists*, Springer, 1971.
- [2] D.F. Lawden, *Elliptic functions and applications*, Springer, 1989.
- [3] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark (eds.), *NIST handbook of mathematical functions*, U.S. Department of Commerce National Institute of Standards and Technology, Washington, DC, 2010.
- [4] P. Walker, *The analyticity of Jacobian functions with respect to the parameter k* , R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **459** (2003), 2569–2574.
- [5] P.L. Walker, *The distribution of the zeros of Jacobian elliptic functions with respect to the parameter k* , Comput. Methods Funct. Theory **9** (2009), 579–591.
- [6] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, Cambridge University Press, 1962.