

WSLD operators II: the new fourth order difference approximations for space Riemann-Liouville derivative

MINGHUA CHEN AND WEIHUA DENG[†]

School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, P. R. China

[Received on 28 February 2018]

High order discretization schemes play more important role in fractional operators than classical ones. This is because usually for classical derivatives the stencil for high order discretization schemes is wider than low order ones; but for fractional operators the stencils for high order schemes and low order ones are the same. Then using high order schemes to solve fractional equations leads to almost the same computational cost with first order schemes but the accuracy is greatly improved. Using the fractional linear multistep methods, Lubich obtains the ν -th order ($\nu \leq 6$) approximations of the α -th derivative ($\alpha > 0$) or integral ($\alpha < 0$) [Lubich, SIAM J. Math. Anal., 17, 704-719, 1986], because of the stability issue the obtained scheme can not be directly applied to the space fractional operator with $\alpha \in (1, 2)$ for time dependent problem. By weighting and shifting Lubich's 2nd order discretization scheme, in [Chen & Deng, arXiv:1304.7425] we derive a series of effective high order discretizations for space fractional derivative, called WSLD operators there. As the sequel of the previous work, we further provide new high order schemes for space fractional derivatives by weighting and shifting Lubich's 3rd and 4th order discretizations. In particular, we prove that the obtained 4th order approximations are effective for space fractional derivatives. And the corresponding schemes are used to solve the space fractional diffusion equation with variable coefficients.

Keywords: Fractional derivatives; High order scheme; Weighted and shifted Lubich difference operators; Numerical stability

1. Introduction

Fractional calculus (i.e., integrals and derivatives of any arbitrary real or even complex order) has attracted considerable attention during the past several decades, due mainly to its demonstrated applications in seemingly diverse and widespread fields of science and engineering [Kilbas *et al.* (2006)]; and fractional derivatives provide an excellent tool for the description of memory and hereditary properties of various materials and processes [Podlubny (1999)]. With the ubiquitous applications of fractional calculus, fractional partial differential equations (PDEs) appear naturally. Effectively solving fractional PDEs becomes urgent, and intrigues mathematicians. It is still possible to analytically solve the linear fractional PDEs with constant coefficients by using Laplace or Fourier transform, but most of the time the solutions are represented by infinite series or transactional functions. Without doubt, the new challenges also exist in numerically solving fractional PDEs; but some basic ideas have been developed, for instance, finite difference method [Meerschaert & Tadjeran (2004); Sousa & Li (2011); Sun & Wu (2006); Tian *et al.* (2012); Zhuang *et al.* (2009)]; finite element method [Ervin & Roop (2006); Deng (2008)]; spectral method [Li & Xu (2009)].

In numerically solving fractional PDEs, besides a little bit complex numerical analysis, the big

[†]Corresponding author. Email: dengwh@lzu.edu.cn

challenge comes from the computational cost caused by the nonlocal properties of fractional operators. High order scheme is a natural idea to reduce the challenge of cost. Comparing with first order schemes, the high order schemes for fractional operators do not increase computational cost but greatly improve the accuracy. The reason is that both the derived matrixes corresponding to the higher order schemes and low order schemes are full and have the same structure [Chen *et al.* (2012)]. In fact, there are already some important progresses for the high order discretizations of fractional derivatives, including WSGD operator [Tian *et al.* (2012)], CWSGD operator [Zhou *et al.* (2013)], second order discretization [Sousa & Li (2011)], second order discretization for Riesz fractional derivative [Ortigueira (2006)], and WSLD operator [Chen & Deng (2013)]. This paper is the sequel of [Chen & Deng (2013)], i.e, based on Lubich's 3rd and 4th operators to provide new high order discretization schemes for space fractional derivatives.

Using the fractional linear multistep methods, Lubich (1986) obtains the v -th order ($v \leq 6$) approximations of the α -th derivative ($\alpha > 0$) or integral ($\alpha < 0$) by the corresponding coefficients of the generating functions $\delta^\alpha(\zeta)$, where

$$\delta^\alpha(\zeta) = \left(\sum_{i=1}^v \frac{1}{i} (1-\zeta)^i \right)^\alpha. \quad (1.1)$$

For $\alpha = 1$, the scheme reduces to the classical $(v+1)$ -point backward difference formula [Henrici (1962)]. For $v = 1$, $\alpha > 0$, the scheme (1.1) corresponds to the standard Grünwald discretization of α -th derivative with first order accuracy; unfortunately, for the time dependent equations all the difference discretizations are unstable. By weighting and shifting Lubich's 2nd order discretization, a class of effective high order schemes for space fractional derivatives are presented [Chen & Deng (2013)]. Is it possible to design the high order schemes for space fractional derivatives by using Lubich's 3rd, 4th, 5th, 6th order operators? This paper will answer that at least by applying Lubich's 3rd, 4th order operators, the new discretizations for space fractional derivatives can be constructed. The concrete discretizations will be presented, and the effectiveness of 4th order schemes for space fractional derivative will be proved. And we will also provide a simple application to solve the space fractional diffusion equation with variable coefficients.

The outline of this paper is as follows. In Section 2, we derive the new fourth order approximations for space fractional Riemann-Liouville derivatives, being effective in solving space fractional PDEs. A simple application of the new discretization schemes are presented in Section 3 to solve the space fractional diffusion equation with variable coefficients. And in Section 4, the numerical experiments are performed to show the effectiveness of the algorithm and verify the theoretical results. Finally, the paper is concluded with some remarks in the last section.

2. Derivation of new fourth order discretizations for space fractional operators

Based on Lubich's 3rd and 4th discretizations, we derive new fourth order approximations for Riemann-Liouville derivative, and prove that they are effective in solving space fractional PDEs, i.e., all the eigenvalues of the matrixes corresponding to the discretized operators have negative real parts.

DEFINITION 2.1 (Podlubny (1999)) The α -th ($n-1 < \alpha < n$) order left and right Riemann-Liouville fractional derivatives of the function $u(x)$ on $[x_L, x_R]$, $-\infty \leq x_L < x_R \leq \infty$ are, respectively, defined by

$${}_{x_L}D_x^\alpha u(x) = \frac{1}{\Gamma(n-\alpha)} \frac{\partial^n}{\partial x^n} \int_{x_L}^x (x-\xi)^{n-\alpha-1} u(\xi) d\xi,$$

and

$${}_x D_{x_R}^\alpha u(x) = \frac{1}{\Gamma(n-\alpha)} \frac{\partial^n}{\partial x^n} \int_x^{x_R} (\xi-x)^{n-\alpha-1} u(\xi) d\xi.$$

LEMMA 2.1 (Ervin & Roop (2006)) Let $\alpha > 0$, $u \in C_0^\infty(\Omega)$, $\Omega \subset \mathbb{R}$, then

$$\mathcal{F}({}_{-\infty} D_x^\alpha u(x)) = (-i\omega)^\alpha \widehat{u}(\omega) \quad \text{and} \quad \mathcal{F}({}_x D_\infty^\alpha u(x)) = (i\omega)^\alpha \widehat{u}(\omega),$$

where \mathcal{F} denotes Fourier transform operator and $\widehat{u}(\omega) = \mathcal{F}(u)$, i.e.,

$$\widehat{u}(\omega) = \int_{\mathbb{R}} e^{i\omega x} u(x) dx.$$

LEMMA 2.2 Let the function $f(z) = (1 + bz + cz^2 + dz^3 + ez^4)^\alpha$, then

$$f(0) = 1, \quad f'(0) = \alpha b, \quad f''(0) = \alpha(\alpha-1)b^2 + 2\alpha c,$$

$$f'''(0) = \alpha(\alpha-1)(\alpha-2)b^3 + 6\alpha(\alpha-1)bc + 6\alpha d,$$

$$f''''(0) = \alpha(\alpha-1)(\alpha-2)(\alpha-3)b^4 + 12\alpha(\alpha-1)(\alpha-2)b^2c + 12\alpha(\alpha-1)c^2 + 24\alpha(\alpha-1)bd + 24\alpha e,$$

and

$$f(z) = f(0+z) = f(0) + zf'(0) + \frac{z^2}{2!}f''(0) + \frac{z^3}{3!}f'''(0) + \frac{z^4}{4!}f''''(0) + \mathcal{O}(z^5).$$

Proof. It is easy to check that by the Taylor series expansion. \square

2.1 Derivation of the discretizations

In this subsection, based on Lubich's operator (1.1), we do the expansions to get the formulas of the coefficients when $\nu = 3, 4, 5$; by the technique of Fourier transform, prove that the operators have their respective desired convergent order; and by weighting and shifting Lubich's 3rd (and 4th) order operators, obtain new high order discretization schemes.

First, taking $\nu = 1$, for all $|\zeta| \leq 1$, Eq. (1.1) becomes the following equation [Podlubny (1999)],

$$(1-\zeta)^\alpha = \sum_{m=0}^{\infty} (-1)^m \binom{\alpha}{m} \zeta^m = \sum_{m=0}^{\infty} l_m^{1,\alpha} \zeta^m, \quad (2.1)$$

with the recursively formula

$$l_0^{1,\alpha} = 1, \quad l_m^{1,\alpha} = \left(1 - \frac{\alpha+1}{m}\right) l_{m-1}^{1,\alpha}, \quad m \geq 1. \quad (2.2)$$

Letting $\nu = 2$, for all $|\zeta| \leq 1$, Eq. (1.1) has the following form [Chen & Deng (2013)],

$$\left(\frac{3}{2} - 2\zeta + \frac{1}{2}\zeta^2\right)^\alpha = \left(\frac{3}{2}\right)^\alpha (1-\zeta)^\alpha \left(1 - \frac{1}{3}\zeta\right)^\alpha = \sum_{j=0}^{\infty} l_j^{2,\alpha} \zeta^j, \quad (2.3)$$

with

$$l_j^{2,\alpha} = \left(\frac{3}{2}\right)^\alpha \sum_{m=0}^j \mu_2^m l_m^{1,\alpha} l_{j-m}^{1,\alpha}, \quad (2.4)$$

where $\mu_2 = \frac{1}{3}$ and $l_m^{1,\alpha}$ is defined by (2.2).

Setting $\nu = 3$, for all $|\zeta| \leq 1$, Eq. (1.1) leads to the following form

$$\begin{aligned}
& \left(\frac{11}{6} - 3\zeta + \frac{3}{2}\zeta^2 - \frac{1}{3}\zeta^3 \right)^\alpha \\
&= (1 - \zeta)^\alpha \left(\frac{11}{6} - \frac{7}{6}\zeta + \frac{1}{3}\zeta^2 \right)^\alpha \\
&= \left(\frac{11}{6} \right)^\alpha (1 - \zeta)^\alpha (1 - \mu_3 \zeta)^\alpha (1 - \overline{\mu}_3 \zeta)^\alpha \\
&= \left(\frac{11}{6} \right)^\alpha \sum_{n=0}^{\infty} (-1)^n \binom{\alpha}{n} \zeta^n \sum_{m=0}^{\infty} (-\mu_3)^m \binom{\alpha}{m} \zeta^m \sum_{l=0}^{\infty} (-\overline{\mu}_3)^l \binom{\alpha}{l} \zeta^l \\
&= \left(\frac{11}{6} \right)^\alpha \sum_{n=0}^{\infty} (-1)^n \binom{\alpha}{n} \zeta^n \left\{ \sum_{j=0}^{\infty} \sum_{m=0}^j (-1)^j \mu_3^m \overline{\mu}_3^{j-m} \binom{\alpha}{m} \binom{\alpha}{j-m} \zeta^j \right\} \\
&= \sum_{k=0}^{\infty} l_k^{3,\alpha} \zeta^k,
\end{aligned} \tag{2.5}$$

with

$$l_k^{3,\alpha} = \left(\frac{11}{6} \right)^\alpha \sum_{j=0}^k \sum_{m=0}^j \mu_3^m \overline{\mu}_3^{j-m} l_m^{1,\alpha} l_{j-m}^{1,\alpha} l_{k-j}^{1,\alpha}, \tag{2.6}$$

where $\mu_3 = \frac{4}{7+\sqrt{39}i}$, $\overline{\mu}_3 = \frac{4}{7-\sqrt{39}i}$, $i = \sqrt{-1}$ and $k = j + n$, $j = l + m$, and $l_m^{1,\alpha}$ is defined by (2.2).

Taking $\nu = 4$, for all $|\zeta| \leq 1$, Eq. (1.1) reduces to the following form

$$\begin{aligned}
\left(\frac{25}{12} - 4\zeta + 3\zeta^2 - \frac{4}{3}\zeta^3 + \frac{1}{4}\zeta^4 \right)^\alpha &= (1 - \zeta)^\alpha \left(\frac{25}{12} - \frac{23}{12}\zeta + \frac{13}{12}\zeta^2 - \frac{1}{4}\zeta^3 \right)^\alpha \\
&= \left(\frac{25}{12} \right)^\alpha (1 - \zeta)^\alpha \left(-\frac{3}{25}\zeta^3 + \frac{13}{25}\zeta^2 - \frac{23}{25}\zeta + 1 \right)^\alpha \\
&= \left(\frac{25}{12} \right)^\alpha (1 - \zeta)^\alpha (1 - \nu_4 \zeta)^\alpha (1 - \mu_4 \zeta)^\alpha (1 - \overline{\mu}_4 \zeta)^\alpha;
\end{aligned}$$

then from Shengjin's Formulas [Fan (1989)], we obtain

$$\begin{aligned}
\nu_4 &= \frac{3a}{-b - (\sqrt[3]{Y_1} - \sqrt[3]{-Y_2})}; \\
\mu_4 &= \frac{3a}{-b + \frac{1}{2}(\sqrt[3]{Y_1} - \sqrt[3]{-Y_2}) + \frac{\sqrt{3}}{2}(\sqrt[3]{Y_1} + \sqrt[3]{-Y_2})i}; \\
\overline{\mu}_4 &= \frac{3a}{-b + \frac{1}{2}(\sqrt[3]{Y_1} - \sqrt[3]{-Y_2}) - \frac{\sqrt{3}}{2}(\sqrt[3]{Y_1} + \sqrt[3]{-Y_2})i},
\end{aligned}$$

where $a = -\frac{3}{25}$, $b = \frac{13}{25}$, $c = -\frac{23}{25}$, $d = 1$; $A = b^2 - 3ac$, $B = bc - 9ad$, $C = c^2 - 3bd$; $\Delta = B^2 - 4AC > 0$,

$Y_1 = Ab + \frac{3}{2}a(-B - \sqrt{\Delta}) > 0$, $Y_2 = Ab + \frac{3}{2}a(-B + \sqrt{\Delta}) < 0$, and $i = \sqrt{-1}$. Thus

$$\begin{aligned} & \left(\frac{25}{12} - 4\zeta + 3\zeta^2 - \frac{4}{3}\zeta^3 + \frac{1}{4}\zeta^4 \right)^\alpha \\ &= \left(\frac{25}{12} \right)^\alpha \sum_{n=0}^{\infty} (-1)^n \binom{\alpha}{n} \zeta^n \sum_{m=0}^{\infty} (-\mu_4)^m \binom{\alpha}{m} \zeta^m \sum_{l=0}^{\infty} (-\bar{\mu}_4)^l \binom{\alpha}{l} \zeta^l \sum_{p=0}^{\infty} (-\nu_4)^p \binom{\alpha}{p} \zeta^p \quad (2.7) \\ &= \sum_{n=0}^{\infty} l_n^{4,\alpha} \zeta^n, \end{aligned}$$

with

$$l_n^{4,\alpha} = \left(\frac{25}{12} \right)^\alpha \sum_{k=0}^n \sum_{j=0}^k \sum_{m=0}^j \nu_4^{n-k} \mu_4^m \bar{\mu}_4^{j-m} l_m^{1,\alpha} l_{j-m}^{1,\alpha} l_{k-j}^{1,\alpha} l_{n-k}^{1,\alpha}, \quad (2.8)$$

and $l_m^{1,\alpha}$ is defined by (2.2).

Taking $\nu = 5$, for all $|\zeta| \leq 1$, Eq. (1.1) has the following form

$$\begin{aligned} & \left(\frac{137}{60} - 5\zeta + 5\zeta^2 - \frac{10}{3}\zeta^3 + \frac{5}{4}\zeta^4 - \frac{1}{5}\zeta^5 \right)^\alpha \\ &= (1 - \zeta)^\alpha \left(\frac{137}{60} - \frac{163}{60}\zeta + \frac{137}{60}\zeta^2 - \frac{21}{20}\zeta^3 + \frac{1}{5}\zeta^4 \right)^\alpha \\ &= \left(\frac{137}{60} \right)^\alpha (1 - \zeta)^\alpha \left(1 - \frac{163}{137}\zeta + \zeta^2 - \frac{63}{137}\zeta^3 + \frac{12}{137}\zeta^4 \right)^\alpha \\ &= \left(\frac{1}{5} \right)^\alpha (1 - \zeta)^\alpha \left(\zeta^4 - \frac{21}{4}\zeta^3 + \frac{137}{12}\zeta^2 - \frac{163}{12}\zeta + \frac{137}{12} \right)^\alpha \\ &= \left(\frac{137}{60} \right)^\alpha (1 - \zeta)^\alpha (1 - \nu_5 \zeta)^\alpha (1 - \bar{\nu}_5 \zeta)^\alpha (1 - \mu_5 \zeta)^\alpha (1 - \bar{\mu}_5 \zeta)^\alpha. \end{aligned}$$

According to Ferrari's Formulas [Polyanin & Manzhirov (2007)] and Shengjin's Formulas [Fan (1989)], we obtain

$$\begin{aligned} \nu_5 &= \frac{4}{-(b+M) + \sqrt{(b+M)^2 - 16(y + \frac{N}{M})}}; \\ \bar{\nu}_5 &= \frac{4}{-(b-M) + \sqrt{(b-M)^2 - 16(y - \frac{N}{M})}}; \\ \mu_5 &= \frac{4}{-(b+M) - \sqrt{(b+M)^2 - 16(y + \frac{N}{M})}}; \\ \bar{\mu}_5 &= \frac{4}{-(b-M) - \sqrt{(b-M)^2 - 16(y - \frac{N}{M})}}, \end{aligned}$$

where $a = 1$, $b = -\frac{21}{4}$, $c = \frac{137}{12}$, $d = -\frac{163}{12}$, $e = \frac{137}{12}$; $\tilde{a} = 1$, $\tilde{b} = -\frac{137}{24}$, $\tilde{c} = \frac{1231}{192}$, $\tilde{d} = \frac{4259}{1536}$; $A = \tilde{b}^2 - 3\tilde{a}\tilde{c}$, $B = \tilde{b}\tilde{c} - 9\tilde{a}\tilde{d}$, $C = \tilde{c}^2 - 3\tilde{b}\tilde{d}$; $\Delta = B^2 - 4AC < 0$; $T = \frac{2A\tilde{b} - 3\tilde{a}B}{2A^{\frac{3}{2}}}$, $\theta = \arccos(T)$ $y = \frac{-\tilde{b} - 2A^{\frac{1}{2}} \cos \frac{\theta}{3}}{3a}$; and

6 of 22

$M = \sqrt{8y + b^2 - 4c}$, $N = \sqrt{by - d}$. Therefore, we have

$$\left(\frac{137}{60} - 5\zeta + 5\zeta^2 - \frac{10}{3}\zeta^3 + \frac{5}{4}\zeta^4 - \frac{1}{5}\zeta^5 \right)^\alpha = \sum_{q=0}^{\infty} l_q^{5,\alpha} \zeta^q, \quad (2.9)$$

where

$$l_q^{5,\alpha} = \left(\frac{137}{60} \right)^\alpha \sum_{n=0}^q \sum_{k=0}^n \sum_{j=0}^k \sum_{m=0}^j \frac{1}{v_5^{q-n} v_5^{n-k} \mu_5^m \bar{\mu}_5^{j-m}} l_m^{1,\alpha} l_{j-m}^{1,\alpha} l_{k-j}^{1,\alpha} l_{n-k}^{1,\alpha} l_{q-n}^{1,\alpha}, \quad (2.10)$$

and $l_m^{1,\alpha}$ is defined by (2.2).

In the following, by using the technique of Fourier transform, we again list and simply prove the convergent order of Lubich's operator.

LEMMA 2.3 (Case $v = 1$; Meerschaert & Tadjeran (2004)) Let u , ${}_{-\infty}D_x^{\alpha+1}u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$, and denote that

$${}_{1L}A_p^{1,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{m=0}^{\infty} l_m^{1,\alpha} u(x - (m-p)h),$$

where $l_m^{1,\alpha}$ is defined by (2.2) and p an integer. Then

$${}_{-\infty}D_x^\alpha u(x) = {}_{1L}A_p^{1,\alpha}u(x) + \mathcal{O}(h).$$

LEMMA 2.4 (Case $v = 2$; Chen & Deng (2013)) Let u , ${}_{-\infty}D_x^{\alpha+1}u(x)$ (or ${}_{-\infty}D_x^{\alpha+2}u(x)$) with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$ when $p \neq 0$ (or $p = 0$); and denote that

$${}_{1L}A_p^{2,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{j=0}^{\infty} l_j^{2,\alpha} u(x - (j-p)h),$$

where $l_j^{2,\alpha}$ is defined by (2.4) and p an integer. Then

$$\begin{aligned} {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{2,\alpha}u(x) + \mathcal{O}(h), & p \neq 0; \\ {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{2,\alpha}u(x) + \mathcal{O}(h^2), & p = 0. \end{aligned}$$

LEMMA 2.5 (Case $v = 3$) Let u , ${}_{-\infty}D_x^{\alpha+1}u(x)$ (or ${}_{-\infty}D_x^{\alpha+3}u(x)$) and their Fourier transforms belong to $L_1(\mathbb{R})$ when $p \neq 0$ (or $p = 0$); and denote that

$${}_{1L}A_p^{3,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{k=0}^{\infty} l_k^{3,\alpha} u(x - (k-p)h), \quad (2.11)$$

where $l_k^{3,\alpha}$ is defined by (2.6) and p an integer. Then

$$\begin{aligned} {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{3,\alpha}u(x) + \mathcal{O}(h), & p \neq 0; \\ {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{3,\alpha}u(x) + \mathcal{O}(h^3), & p = 0. \end{aligned}$$

Proof. According to (2.5), there exist

$$\begin{aligned}
\mathcal{F}({}_{1L}A_p^{3,\alpha}u)(\omega) &= h^{-\alpha} \sum_{k=0}^{\infty} l_k^{3,\alpha} \mathcal{F}(u(x - (k-p)h))(\omega) \\
&= h^{-\alpha} e^{-i\omega p h} \sum_{k=0}^{\infty} l_k^{3,\alpha} (e^{i\omega h})^k \widehat{u}(\omega) \\
&= h^{-\alpha} e^{-i\omega p h} (1 - e^{i\omega h})^\alpha \left(\frac{11}{6} - \frac{7}{6} e^{i\omega h} + \frac{1}{3} e^{2i\omega h} \right)^\alpha \widehat{u}(\omega) \\
&= (-i\omega)^\alpha e^{-i\omega p h} \left(\frac{1 - e^{i\omega h}}{-i\omega h} \right)^\alpha \left(\frac{11}{6} - \frac{7}{6} e^{i\omega h} + \frac{1}{3} e^{2i\omega h} \right)^\alpha \widehat{u}(\omega) \\
&= (-i\omega)^\alpha e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \left(\frac{11}{6} - \frac{7}{6} e^{-z} + \frac{1}{3} e^{-2z} \right)^\alpha \widehat{u}(\omega),
\end{aligned}$$

with $z = -i\omega h$. It is easy to check that

$$\begin{aligned}
&e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \\
&= 1 + \left(p - \frac{\alpha}{2} \right) z + \left(\frac{1}{2} p^2 - \frac{\alpha}{2} p + \frac{3\alpha^2 + \alpha}{24} \right) z^2 \\
&\quad + \left(\frac{1}{6} p^3 - \frac{\alpha}{4} p^2 + \frac{3\alpha^2 + \alpha}{24} p - \frac{\alpha^3 + \alpha^2}{48} \right) z^3 \\
&\quad + \left(\frac{1}{24} p^4 - \frac{\alpha}{12} p^3 + \frac{3\alpha^2 + \alpha}{48} p^2 - \frac{\alpha^3 + \alpha^2}{48} p + \frac{15\alpha^4 + 30\alpha^3 + 5\alpha^2 - 2\alpha}{5760} \right) z^4 + \mathcal{O}(z^5),
\end{aligned} \tag{2.12}$$

and from the Lemma 2.2, we have

$$\begin{aligned}
\left(\frac{11}{6} - \frac{7}{6} e^{-z} + \frac{1}{3} e^{-2z} \right)^\alpha &= \left(1 + \frac{1}{2} z + \frac{1}{12} z^2 - \frac{1}{4} z^3 + \mathcal{O}(z^4) \right)^\alpha \\
&= 1 + \frac{\alpha}{2} z + \frac{\alpha(3\alpha - 1)}{24} z^2 + \frac{\alpha(\alpha + 3)(\alpha - 4)}{48} z^3 + \mathcal{O}(z^4),
\end{aligned}$$

then from the above equation and (2.12) we obtain

$$e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \left(\frac{11}{6} - \frac{7}{6} e^{-z} + \frac{1}{3} e^{-2z} \right)^\alpha = 1 + pz + \frac{p^2}{2} z^2 + \frac{2p^3 - 3\alpha}{12} z^3 + \mathcal{O}(z^4). \tag{2.13}$$

Therefore, from Lemma 2.1, we obtain

$$\mathcal{F}({}_{1L}A_p^{3,\alpha}u)(\omega) = \mathcal{F}({}_{-\infty}D_x^\alpha u(x)) + \widehat{\phi}(\omega),$$

where $\widehat{\phi}(\omega) = (-i\omega)^\alpha \left(pz + \frac{p^2}{2} z^2 + \frac{2p^3 - 3\alpha}{12} z^3 + \mathcal{O}(z^4) \right) \widehat{u}(\omega)$, $z = -i\omega h$. Then there exists

$$\begin{aligned}
|\widehat{\phi}(\omega)| &\leq \widetilde{c} |i\omega|^{\alpha+1} |\widehat{u}(\omega)| \cdot h, & p \neq 0; \\
|\widehat{\phi}(\omega)| &\leq c |i\omega|^{\alpha+3} |\widehat{u}(\omega)| \cdot h^3, & p = 0.
\end{aligned}$$

Hence

$$|{}_{-\infty}D_x^\alpha u(x) - {}_{1L}A_p^{3,\alpha} u(x)| = |\phi(x)| \leq \frac{1}{2\pi} \int_{\mathbb{R}} |\widehat{\phi}(\omega)| dx = \begin{cases} \mathcal{O}(h), & p \neq 0; \\ \mathcal{O}(h^3), & p = 0. \end{cases}$$

□

LEMMA 2.6 (Case $\nu = 4$) Let u , ${}_{-\infty}D_x^{\alpha+1}u(x)$ (or ${}_{-\infty}D_x^{\alpha+4}u(x)$) with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$ when $p \neq 0$ (or $p = 0$); and denote that

$${}_{1L}A_p^{4,\alpha} u(x) = \frac{1}{h^\alpha} \sum_{n=0}^{\infty} l_n^{4,\alpha} u(x - (n-p)h), \quad (2.14)$$

where $l_n^{4,\alpha}$ is defined by (2.8) and p an integer. Then

$$\begin{aligned} {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{4,\alpha} u(x) + \mathcal{O}(h), & p \neq 0; \\ {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{4,\alpha} u(x) + \mathcal{O}(h^4), & p = 0. \end{aligned}$$

Proof. According to (2.7) we obtain

$$\begin{aligned} \mathcal{F}({}_{1L}A_p^{4,\alpha} u)(\omega) &= h^{-\alpha} \sum_{n=0}^{\infty} l_n^{4,\alpha} \mathcal{F}(u(x - (n-p)h))(\omega) \\ &= h^{-\alpha} e^{-i\omega p h} \sum_{n=0}^{\infty} l_n^{4,\alpha} \left(e^{i\omega h} \right)^n \widehat{u}(\omega) \\ &= h^{-\alpha} e^{-i\omega p h} \left(1 - e^{i\omega h} \right)^\alpha \left(\frac{25}{12} - \frac{23}{12} e^{i\omega h} + \frac{13}{12} e^{2i\omega h} - \frac{1}{4} e^{3i\omega h} \right)^\alpha \widehat{u}(\omega) \\ &= (-i\omega)^\alpha e^{-i\omega p h} \left(\frac{1 - e^{i\omega h}}{-i\omega h} \right)^\alpha \left(\frac{25}{12} - \frac{23}{12} e^{i\omega h} + \frac{13}{12} e^{2i\omega h} - \frac{1}{4} e^{3i\omega h} \right)^\alpha \widehat{u}(\omega) \\ &= (-i\omega)^\alpha e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \left(\frac{25}{12} - \frac{23}{12} e^{-z} + \frac{13}{12} e^{-2z} - \frac{1}{4} e^{-3z} \right)^\alpha \widehat{u}(\omega), \end{aligned}$$

where $z = -i\omega h$. From Lemma 2.2, there exist

$$\begin{aligned} \left(\frac{25}{12} - \frac{23}{12} e^{-z} + \frac{13}{12} e^{-2z} - \frac{1}{4} e^{-3z} \right)^\alpha &= \left(1 + \frac{1}{2}z + \frac{1}{12}z^2 - \frac{29}{144}z^4 + \mathcal{O}(z^5) \right)^\alpha \\ &= 1 + \frac{\alpha}{2}z + \frac{\alpha(3\alpha-1)}{24}z^2 + \frac{\alpha^2(\alpha-1)}{48}z^3 + \mathcal{O}(z^4), \end{aligned}$$

then we obtain

$$e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \left(\frac{25}{12} - \frac{23}{12} e^{-z} + \frac{13}{12} e^{-2z} - \frac{1}{4} e^{-3z} \right)^\alpha = 1 + pz + \frac{p^2}{2!}z^2 + \frac{p^3}{3!}z^3 + \mathcal{O}(z^4). \quad (2.15)$$

Therefore, from Lemma 2.1, we obtain

$$\mathcal{F}({}_{1L}A_p^{4,\alpha} u)(\omega) = \mathcal{F}({}_{-\infty}D_x^\alpha u(x)) + \widehat{\phi}(\omega),$$

where $\widehat{\phi}(\omega) = (-i\omega)^\alpha \left(pz + \frac{p^2}{2!}z^2 + \frac{p^3}{3!}z^3 + \mathcal{O}(z^4) \right) \widehat{u}(\omega)$. Then there exists

$$\begin{aligned} |\widehat{\phi}(\omega)| &\leq \widetilde{c}|i\omega|^{\alpha+1}|\widehat{u}(\omega)| \cdot h, & p \neq 0; \\ |\widehat{\phi}(\omega)| &\leq c|i\omega|^{\alpha+4}|\widehat{u}(\omega)| \cdot h^4, & p = 0. \end{aligned}$$

Hence

$$|_{-\infty}D_x^\alpha u(x) - {}_{1L}A_p^{4,\alpha}u(x)| = |\phi(x)| \leq \frac{1}{2\pi} \int_{\mathbb{R}} |\widehat{\phi}(\omega)| dx = \begin{cases} \mathcal{O}(h), & p \neq 0; \\ \mathcal{O}(h^4), & p = 0. \end{cases}$$

□

LEMMA 2.7 (Case $\nu = 5$) Let $u, {}_{-\infty}D_x^{\alpha+1}u(x)$ (or ${}_{-\infty}D_x^{\alpha+5}u(x)$) with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$ when $p \neq 0$ (or $p = 0$); and denote that

$${}_{1L}A_p^{5,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{q=0}^{\infty} l_q^{5,\alpha} u(x - (q-p)h), \quad (2.16)$$

where $l_q^{5,\alpha}$ is defined by (2.10) and p an integer. Then

$$\begin{aligned} {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{5,\alpha}u(x) + \mathcal{O}(h), & p \neq 0; \\ {}_{-\infty}D_x^\alpha u(x) &= {}_{1L}A_p^{5,\alpha}u(x) + \mathcal{O}(h^5), & p = 0. \end{aligned}$$

Proof. According to (2.9) we obtain

$$\begin{aligned} \mathcal{F}({}_{1L}A_p^{5,\alpha}u)(\omega) &= h^{-\alpha} \sum_{l=0}^{\infty} l_q^{5,\alpha} \mathcal{F}(u(x - (q-p)h))(\omega) \\ &= h^{-\alpha} e^{-i\omega p h} \sum_{l=0}^{\infty} l_q^{5,\alpha} \left(e^{i\omega h} \right)^q \widehat{u}(\omega) \\ &= h^{-\alpha} e^{-i\omega p h} \left(1 - e^{i\omega h} \right)^\alpha \left(\frac{137}{60} - \frac{163}{60} e^{i\omega h} + \frac{137}{60} e^{2i\omega h} - \frac{21}{20} e^{3i\omega h} + \frac{1}{5} e^{4i\omega h} \right)^\alpha \widehat{u}(\omega) \\ &= (-i\omega)^\alpha e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \left(\frac{137}{60} - \frac{163}{60} e^{-z} + \frac{137}{60} e^{-2z} - \frac{21}{20} e^{-3z} + \frac{1}{5} e^{-4z} \right)^\alpha \widehat{u}(\omega), \end{aligned}$$

where $z = -i\omega h$. Using Lemma 2.2 leads to

$$\begin{aligned} &\left(\frac{137}{60} - \frac{163}{60} e^{-z} + \frac{137}{60} e^{-2z} - \frac{21}{20} e^{-3z} + \frac{1}{5} e^{-4z} \right)^\alpha \\ &= \left(1 + \frac{1}{2}z + \frac{1}{12}z^2 - \frac{1}{720}z^4 + \mathcal{O}(z^5) \right)^\alpha \\ &= 1 + \frac{\alpha}{2}z + \frac{\alpha(3\alpha-1)}{24}z^2 + \frac{\alpha^2(\alpha-1)}{48}z^3 + \frac{15\alpha^4 - 30\alpha^3 + 5\alpha^2 + 2\alpha}{5760}z^4 + \mathcal{O}(z^5), \end{aligned}$$

then we obtain

$$\begin{aligned} &e^{pz} \left(\frac{1 - e^{-z}}{z} \right)^\alpha \left(\frac{137}{60} - \frac{163}{60} e^{-z} + \frac{137}{60} e^{-2z} - \frac{21}{20} e^{-3z} + \frac{1}{5} e^{-4z} \right)^\alpha \\ &= 1 + pz + \frac{p^2}{2!}z^2 + \frac{p^3}{3!}z^3 + \frac{p^4}{4!}z^4 + \mathcal{O}(z^5). \end{aligned} \quad (2.17)$$

Therefore, from Lemma 2.1, there exists

$$\mathcal{F}({}_{1L}A_p^{5,\alpha}u)(\omega) = \mathcal{F}(-{}_{\infty}D_x^\alpha u(x)) + \widehat{\phi}(\omega),$$

where $\widehat{\phi}(\omega) = (-i\omega)^\alpha \left(pz + \frac{p^2}{2!}z^2 + \frac{p^3}{3!}z^3 + \frac{p^4}{4!}z^4 + \mathcal{O}(z^5) \right) \widehat{u}(\omega)$. Then we get

$$\begin{aligned} |\widehat{\phi}(\omega)| &\leq \widetilde{c}|i\omega|^{\alpha+1}|\widehat{u}(\omega)| \cdot h, & p \neq 0; \\ |\widehat{\phi}(\omega)| &\leq c|i\omega|^{\alpha+5}|\widehat{u}(\omega)| \cdot h^5, & p = 0. \end{aligned}$$

Hence

$$|{}_{\infty}D_x^\alpha u(x) - {}_{1L}A_p^{5,\alpha}u(x)| = |\phi(x)| \leq \frac{1}{2\pi} \int_{\mathbb{R}} |\widehat{\phi}(\omega)| dx = \begin{cases} \mathcal{O}(h), & p \neq 0; \\ \mathcal{O}(h^5), & p = 0. \end{cases}$$

□

According to Lemmas 2.2-2.6, the fractional approximation operators have the same form

$${}_{1L}A_p^{v,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{k=0}^{\infty} l_k^{v,\alpha} u(x - (k-p)h), \quad v = 1, 2, 3, 4, 5,$$

where $l_k^{1,\alpha}$, $l_k^{2,\alpha}$, $l_k^{3,\alpha}$, $l_k^{4,\alpha}$ and $l_k^{5,\alpha}$ are, respectively, defined by (2.2), (2.4), (2.6), (2.8) and (2.10). By the same idea of the proof in [Chen & Deng (2013)], we can get the following Theorems 2.2-2.7; and for the simplicity, we omit the proofs here.

THEOREM 2.2 (Case $v = 3, 4$; Second order approximations for left Riemann-Liouville derivative) Let u , ${}_{\infty}D_x^{\alpha+2}u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$. Denote that

$$2LA_{p,q}^{v,\alpha}u(x) = w_p^v {}_{1L}A_p^{v,\alpha}u(x) + w_q^v {}_{1L}A_q^{v,\alpha}u(x), \quad v = 3, 4, \quad (2.18)$$

where ${}_{1L}A_p^{3,\alpha}$ and ${}_{1L}A_q^{3,\alpha}$ are defined in (2.11); ${}_{1L}A_p^{4,\alpha}$ and ${}_{1L}A_q^{4,\alpha}$ are defined in (2.14); $w_p^v = \frac{q}{q-p}$, $w_q^v = \frac{p}{p-q}$, $v = 3, 4$, and p, q are integers, $p \neq q$. Then

$${}_{\infty}D_x^\alpha u(x) = 2LA_{p,q}^{v,\alpha}u(x) + \mathcal{O}(h^2).$$

THEOREM 2.3 (Case $v = 3, 4$; Third order approximations for left Riemann-Liouville derivative) Let u , ${}_{\infty}D_x^{\alpha+3}u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$. Denote that

$$3LA_{p,q,r,s}^{v,\alpha}u(x) = w_{p,q}^v 2LA_{p,q}^{v,\alpha}u(x) + w_{r,s}^v 2LA_{r,s}^{v,\alpha}u(x), \quad v = 3, 4, \quad (2.19)$$

where $2LA_{p,q}^{v,\alpha}$ and $2LA_{r,s}^{v,\alpha}$ are defined in (2.18), $w_{p,q}^v = \frac{rs}{rs-pq}$, $w_{r,s}^v = \frac{pq}{pq-rs}$, $v = 3, 4$, and p, q, r, s are integers, $pq \neq rs$. Then

$${}_{\infty}D_x^\alpha u(x) = 3LA_{p,q,r,s}^{v,\alpha}u(x) + \mathcal{O}(h^3).$$

THEOREM 2.4 (Case $v = 3, 4$; Fourth order approximations for left Riemann-Liouville derivative) Let u , ${}_{\infty}D_x^{\alpha+4}u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$. Denote that

$$4LA_{p,q,r,s,\overline{p},\overline{q},\overline{r},\overline{s}}^{v,\alpha}u(x) = w_{p,q,r,s}^v 3LA_{p,q,r,s}^{v,\alpha}u(x) + w_{\overline{p},\overline{q},\overline{r},\overline{s}}^v 3LA_{\overline{p},\overline{q},\overline{r},\overline{s}}^{v,\alpha}u(x), \quad v = 3, 4, \quad (2.20)$$

where ${}_3LA_{p,q,r,s}^{v,\alpha}$ and ${}_3LA_{\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}$ are defined in (2.19), and

$$w_{p,q,r,s}^v = \frac{c_{\bar{p},\bar{q},\bar{r},\bar{s}}^v}{c_{\bar{p},\bar{q},\bar{r},\bar{s}}^v - c_{p,q,r,s}^v}; \quad w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v = \frac{c_{p,q,r,s}^v}{c_{p,q,r,s}^v - c_{\bar{p},\bar{q},\bar{r},\bar{s}}^v};$$

where

$$c_{p,q,r,s}^3 = \frac{2pqrs(r+s-p-q) + 3\alpha(pq-rs)}{12(rs-pq)}; \quad c_{\bar{p},\bar{q},\bar{r},\bar{s}}^3 = \frac{2\bar{p}\bar{q}\bar{r}\bar{s}(\bar{r}+\bar{s}-\bar{p}-\bar{q}) + 3\alpha(\bar{p}\bar{q}-\bar{r}\bar{s})}{12(\bar{r}\bar{s}-\bar{p}\bar{q})},$$

and

$$c_{p,q,r,s}^4 = \frac{pqrs(r+s-p-q)}{6(rs-pq)}; \quad c_{\bar{p},\bar{q},\bar{r},\bar{s}}^4 = \frac{\bar{p}\bar{q}\bar{r}\bar{s}(\bar{r}+\bar{s}-\bar{p}-\bar{q})}{6(\bar{r}\bar{s}-\bar{p}\bar{q})},$$

and $p, q, r, s, \bar{p}, \bar{q}, \bar{r}, \bar{s}$, are integers, $c_{p,q,r,s}^v \neq c_{\bar{p},\bar{q},\bar{r},\bar{s}}^v$. Then

$$-{}_{\infty}D_x^\alpha u(x) = {}_4LA_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha} u(x) + \mathcal{O}(h^4).$$

For the right Riemann-Liouville fractional derivative, denote that

$${}_1RA_p^{v,\alpha} u(x) = \frac{1}{h^\alpha} \sum_{k=0}^{\infty} l_k^{v,\alpha} u(x + (k-p)h), \quad v = 1, 2, 3, 4, 5, \quad (2.21)$$

where $l_k^{1,\alpha}, l_k^{2,\alpha}, l_k^{3,\alpha}, l_k^{4,\alpha}$ and $l_k^{5,\alpha}$ are, respectively, defined by (2.2), (2.4), (2.6), (2.8) and (2.10), and p is a integer. In particular, the coefficients in (2.22) are completely the same as the ones in (2.18); and the coefficients in (2.23) the same as the ones in (2.19); and the coefficients in (2.24) the same as the ones in (2.20).

THEOREM 2.5 (Case $v = 3, 4$; Second order approximations for right Riemann-Liouville derivative) Let $u, {}_x D_\infty^{\alpha+2} u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$, and denote that

$${}_2RA_{p,q}^{v,\alpha} u(x) = w_p^v {}_1RA_p^{v,\alpha} u(x) + w_q^v {}_1RA_q^{v,\alpha} u(x), \quad v = 3, 4, \quad (2.22)$$

then

$${}_x D_\infty^\alpha u(x) = {}_2RA_{p,q}^{v,\alpha} u(x) + \mathcal{O}(h^2).$$

THEOREM 2.6 (Case $v = 3, 4$; Third order approximations for right Riemann-Liouville derivative) Let $u, {}_x D_\infty^{\alpha+3} u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$, and denote that

$${}_3RA_{p,q,r,s}^{v,\alpha} u(x) = w_{p,q}^v {}_2RA_{p,q}^{v,\alpha} u(x) + w_{r,s}^v {}_2RA_{r,s}^{v,\alpha} u(x), \quad v = 3, 4, \quad (2.23)$$

then

$${}_x D_\infty^\alpha u(x) = {}_3RA_{p,q,r,s}^{v,\alpha} u(x) + \mathcal{O}(h^3).$$

THEOREM 2.7 (Case $v = 3, 4$; Fourth order approximations for right Riemann-Liouville derivative) Let $u, {}_x D_\infty^{\alpha+4} u(x)$ with $\alpha \in (1, 2)$ and their Fourier transforms belong to $L_1(\mathbb{R})$, and denote that

$${}_4RA_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha} u(x) = w_{p,q,r,s}^v {}_3RA_{p,q,r,s}^{v,\alpha} u(x) + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v {}_3RA_{\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha} u(x), \quad v = 3, 4, \quad (2.24)$$

then

$${}_x D_\infty^\alpha u(x) = {}_4RA_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha} u(x) + \mathcal{O}(h^4).$$

All the above schemes are applicable to finite domain, say, (x_L, x_R) , after performing zero extensions to the functions considered. Let $u(x)$ be the zero extended function from the finite domain (x_L, x_R) , and satisfy the requirements of the above corresponding theorems (Lemma 2.5 - Theorem 2.7). Taking

$${}_{1L}\tilde{A}_p^{v,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{k=0}^{\lfloor \frac{x-x_L}{h} \rfloor + p} l_k^{v,\alpha} u(x - (k-p)h), \quad v = 3, 4. \quad (2.25)$$

Then

$$\begin{aligned} {}_{x_L}D_x^\alpha u(x) &= {}_{1L}\tilde{A}_p^{v,\alpha}u(x) + \mathcal{O}(h), & p \neq 0; \\ {}_{x_L}D_x^\alpha u(x) &= {}_{1L}\tilde{A}_p^{v,\alpha}u(x) + \mathcal{O}(h^v), & p = 0; \\ {}_{x_L}D_x^\alpha u(x) &= {}_{2L}\tilde{A}_{p,q}^{v,\alpha}u(x) + \mathcal{O}(h^2), & {}_{2L}\tilde{A}_{p,q}^{v,\alpha}u(x) = w_p^v {}_{1L}\tilde{A}_p^{v,\alpha}u(x) + w_q^v {}_{1L}\tilde{A}_q^{v,\alpha}u(x); \\ {}_{x_L}D_x^\alpha u(x) &= {}_{3L}\tilde{A}_{p,q,r,s}^{v,\alpha}u(x) + \mathcal{O}(h^3), & {}_{3L}\tilde{A}_{p,q,r,s}^{v,\alpha}u(x) = w_{p,q}^v {}_{2L}\tilde{A}_{p,q}^{v,\alpha}u(x) + w_{r,s}^v {}_{2L}\tilde{A}_{r,s}^{v,\alpha}u(x); \\ {}_{x_L}D_x^\alpha u(x) &= {}_{4L}\tilde{A}_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}u(x) + \mathcal{O}(h^4); \end{aligned} \quad (2.26)$$

where

$${}_{4L}\tilde{A}_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}u(x) = w_{p,q,r,s}^v {}_{3L}\tilde{A}_{p,q,r,s}^{v,\alpha}u(x) + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v {}_{3L}\tilde{A}_{\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}u(x).$$

Denoting $x_i = x_L + ih$, $i = -m, \dots, 0, 1, \dots, N_x - 1, N_x, \dots, N_x + m$, and $h = (x_R - x_L)/N_x$ being the uniform spacestep, it can be note that

$$u(x_i) = 0, \quad \text{for } i = -m, -m+1, \dots, 0 \quad \text{and} \quad i = N_x, N_x+1, \dots, N_x+m,$$

where

$$m = \max(\text{abs}(p, q, r, s, \bar{p}, \bar{q}, \bar{r}, \bar{s})). \quad (2.27)$$

Then the approximation operator of (2.25) can be described as

$${}_{1L}\tilde{A}_p^{v,\alpha}u(x_i) = \frac{1}{h^\alpha} \sum_{k=0}^{i+p} l_k^{v,\alpha} u(x_{i-k+p}) = \frac{1}{h^\alpha} \sum_{k=m-p}^{i+m} l_{k+p-m}^{v,\alpha} u(x_{i-k+m}) = \frac{1}{h^\alpha} \sum_{k=0}^{i+m} l_{k+p-m}^{v,\alpha} u(x_{i-k+m}), \quad (2.28)$$

where $l_{k+p-m}^{v,\alpha} = 0$, when $k+p-m < 0$, and p is an integer. Then

$$\begin{aligned} {}_{x_L}D_x^\alpha u(x_i) &= {}_{1L}\tilde{A}_p^{v,\alpha}u(x_i) + \mathcal{O}(h) = \frac{1}{h^\alpha} \sum_{k=0}^{i+m} l_{k+p-m}^{v,\alpha} u(x_{i-k+m}) + \mathcal{O}(h), & p \neq 0; \\ {}_{x_L}D_x^\alpha u(x_i) &= {}_{1L}\tilde{A}_p^{v,\alpha}u(x_i) + \mathcal{O}(h^v) = \frac{1}{h^\alpha} \sum_{k=0}^{i+m} l_{k+p-m}^{v,\alpha} u(x_{i-k+m}) + \mathcal{O}(h^v), & p = 0; \\ {}_{x_L}D_x^\alpha u(x_i) &= {}_{2L}\tilde{A}_{p,q}^{v,\alpha}u(x_i) + \mathcal{O}(h^2) = \frac{1}{h^\alpha} \sum_{k=0}^{i+m} (w_p^v l_{k+p-m}^{v,\alpha} + w_q^v l_{k+q-m}^{v,\alpha}) u(x_{i-k+m}) + \mathcal{O}(h^2); \\ {}_{x_L}D_x^\alpha u(x_i) &= {}_{3L}\tilde{A}_{p,q,r,s}^{v,\alpha}u(x_i) + \mathcal{O}(h^3) = \frac{1}{h^\alpha} \sum_{k=0}^{i+m} (w_{p,q}^v w_p^v l_{k+p-m}^{v,\alpha} + w_{p,q}^v w_q^v l_{k+q-m}^{v,\alpha} \\ &\quad + w_{r,s}^v w_r^v l_{k+r-m}^{v,\alpha} + w_{r,s}^v w_s^v l_{k+s-m}^{v,\alpha}) u(x_{i-k+m}) + \mathcal{O}(h^3); \\ {}_{x_L}D_x^\alpha u(x_i) &= {}_{4L}\tilde{A}_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}u(x_i) + \mathcal{O}(h^4) = \frac{1}{h^\alpha} \sum_{k=0}^{i+m} \phi_k^{v,\alpha} u(x_{i-k+m}) + \mathcal{O}(h^4), \end{aligned} \quad (2.29)$$

where

$$\begin{aligned}\varphi_k^{v,\alpha} &= w_{p,q,r,s}^v w_{p,q}^v w_p^v l_{k+p-m}^{v,\alpha} + w_{p,q,r,s}^v w_{p,q}^v w_q^v l_{k+q-m}^{v,\alpha} + w_{p,q,r,s}^v w_{r,s}^v w_r^v l_{k+r-m}^{v,\alpha} \\ &+ w_{p,q,r,s}^v w_{r,s}^v w_s^v l_{k+s-m}^{v,\alpha} + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{p},\bar{q}}^v w_{\bar{p}}^v l_{k+\bar{p}-m}^{v,\alpha} + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{p},\bar{q}}^v w_{\bar{p}}^v l_{k+\bar{q}-m}^{v,\alpha} \\ &+ w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{r},\bar{s}}^v w_{\bar{r}}^v l_{k+\bar{r}-m}^{v,\alpha} + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{r},\bar{s}}^v w_{\bar{s}}^v l_{k+\bar{s}-m}^{v,\alpha}.\end{aligned}\quad (2.30)$$

Taking $U = [u(x_0), u(x_1), \dots, u(x_{N_x})]^T$, then (2.28) can be rewritten as matrix form

$${}_1L\tilde{A}_p^{v,\alpha}U = \frac{1}{h^\alpha}A_p^{v,\alpha}U, \quad (2.31)$$

where

$$A_p^{v,\alpha} = \begin{bmatrix} l_p^{v,\alpha} & l_{p-1}^{v,\alpha} & \dots & l_0^{v,\alpha} & & & & & \\ l_{p+1}^{v,\alpha} & l_p^{v,\alpha} & l_{p-1}^{v,\alpha} & \dots & l_0^{v,\alpha} & & & & \\ l_{p+2}^{v,\alpha} & l_{p+1}^{v,\alpha} & l_p^{v,\alpha} & l_{p-1}^{v,\alpha} & \dots & l_0^{v,\alpha} & & & \\ \vdots & \ddots & \ddots & \ddots & \ddots & \dots & \ddots & & \\ l_{n-2}^{v,\alpha} & \dots & \ddots & l_{p+1}^{v,\alpha} & l_p^{v,\alpha} & l_{p-1}^{v,\alpha} & \dots & l_0^{v,\alpha} & \\ \vdots & \ddots & \dots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ l_{p+n-3}^{v,\alpha} & \dots & \ddots & \dots & \ddots & l_{p+1}^{v,\alpha} & l_p^{v,\alpha} & l_{p-1}^{v,\alpha} & l_{p-2}^{v,\alpha} \\ l_{p+n-2}^{v,\alpha} & l_{p+n-3}^{v,\alpha} & \dots & l_{n-2}^{v,\alpha} & \dots & l_{p+2}^{v,\alpha} & l_{p+1}^{v,\alpha} & l_p^{v,\alpha} & l_{p-1}^{v,\alpha} \end{bmatrix}, \quad (2.32)$$

and $l_k^{v,\alpha} = 0$, when $k < 0$, and p is an integer. From (2.26) and (2.31) we obtain

$$\begin{aligned}2L\tilde{A}_{p,q}^{v,\alpha}U &= \frac{1}{h^\alpha}A_{p,q}^{v,\alpha}U, & A_{p,q}^{v,\alpha} &= w_p^v A_p^{v,\alpha} + w_q^v A_q^{v,\alpha}; \\ 3L\tilde{A}_{p,q,r,s}^{v,\alpha}U &= \frac{1}{h^\alpha}A_{p,q,r,s}^{v,\alpha}U, & A_{p,q,r,s}^{v,\alpha} &= w_{p,q}^v A_{p,q}^{v,\alpha} + w_{r,s}^v A_{r,s}^{v,\alpha}; \\ 4L\tilde{A}_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}U &= \frac{1}{h^\alpha}A_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}U, & A_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha} &= w_{p,q,r,s}^v A_{p,q,r,s}^{v,\alpha} + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v A_{\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}.\end{aligned}\quad (2.33)$$

Similarly, for the right Riemann-Liouville fractional derivative, assume that

$${}_1R\tilde{A}_p^{v,\alpha}u(x) = \frac{1}{h^\alpha} \sum_{k=0}^{[\frac{x_R-x}{h}]+p} l_k^{v,\alpha} u(x + (k-p)h),$$

then there exists

$${}_1R\tilde{A}_p^{v,\alpha}u(x_i) = \frac{1}{h^\alpha} \sum_{k=0}^{N_x-i+p} l_k^{v,\alpha} u(x_{i+k-p}) = \frac{1}{h^\alpha} \sum_{k=m-p}^{N_x-i+m} l_{k+p-m}^{v,\alpha} u(x_{i+k-m}) = \frac{1}{h^\alpha} \sum_{k=0}^{N_x-i+m} l_{k+p-m}^{v,\alpha} u(x_{i+k-m}),$$

where $l_{k+p-m}^{v,\alpha} = 0$, $k+p-m < 0$, and p is an integer. And the fourth order approximation is

$${}_x D_{x_R}^\alpha u(x_i) = 4R\tilde{A}_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}u(x_i) + \mathcal{O}(h^4) = \frac{1}{h^\alpha} \sum_{k=0}^{N_x-i+m} \varphi_k^{v,\alpha} u(x_{i+k-m}) + \mathcal{O}(h^4), \quad (2.34)$$

where $\varphi_k^{v,\alpha}$ is defined by (2.30), and the matrices forms are

$$\begin{aligned}
1R\tilde{A}_p^{v,\alpha}U &= \frac{1}{h^\alpha}B_p^{v,\alpha}U, & B_p^{v,\alpha} &= (A_p^{v,\alpha})^T \\
2R\tilde{A}_{p,q}^{v,\alpha}U &= \frac{1}{h^\alpha}B_{p,q}^{v,\alpha}U, & B_{p,q}^{v,\alpha} &= w_p^v B_p^{v,\alpha} + w_q^v B_q^{v,\alpha}; \\
3R\tilde{A}_{p,q,r,s}^{v,\alpha}U &= \frac{1}{h^\alpha}B_{p,q,r,s}^{v,\alpha}U, & B_{p,q,r,s}^{v,\alpha} &= w_{p,q}^v B_{p,q}^{v,\alpha} + w_{r,s}^v B_{r,s}^{v,\alpha}; \\
4R\tilde{A}_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}U &= \frac{1}{h^\alpha}B_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}U, & B_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha} &= w_{p,q,r,s}^v B_{p,q,r,s}^{v,\alpha} + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v B_{\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}.
\end{aligned} \tag{2.35}$$

REMARK 2.1 We want to emphasize that for $v = 3$ and 4 the coefficients in (2.18) are the same, but they correspond to different discretization schemes; the same thing also happens for (2.19), (2.22), and (2.23).

REMARK 2.2 When $p = 0$, $\alpha \in (1, 2)$ and $v = 3, 4$, then $A_p^{v,\alpha}$ in (2.32) reduces to the lower triangular matrix, and it can be easily checked that all the eigenvalues of $A_p^{v,\alpha}$ are greater than one. This is the reason that the scheme for time dependent problem is unstable when directly using the v -order ($v = 2, 3, 4, 5$) Lubich's operator with $\alpha \in (1, 2)$ to discretize space fractional derivative.

REMARK 2.3 If $\alpha < 0$, $\{I_k^{v,\alpha}\}_{k=0}^\infty$ ($v = 1, 2, 3, 4, 5$) correspond to the coefficients of the v -order approximation of fractional integral operators.

EXAMPLE 2.8 To numerically verify the truncation error given in Lemma 2.7 ($v = 5, p = 0$) in a bounded domain. We utilize the approximation (2.28) to simulate the following equation

$${}_0D_x^\alpha u(x) = \frac{\Gamma(9)}{\Gamma(9-\alpha)} x^{8-\alpha}, \quad x \in (0, 1),$$

when $\alpha < 0$, the fractional operator ${}_0D_x^\alpha$ is a fractional integral operator; if $\alpha \in (0, 1)$ we take $u(0) = 0$; and if $\alpha \in (1, 2)$ let $u(0) = 0$, $u(1) = 1$; the exact solution of the above equation is $u(x) = x^8$.

Table 1. The maximum errors and convergent orders for the second equation of (2.29), when $v = 5, p = 0$.

h	$\alpha = -0.5$	Rate	$\alpha = 0.5$	Rate	$\alpha = 1.8$	Rate
1/10	8.0041e-04		4.0005e-03		6.9882e-02	
1/20	4.9935e-05	4.0026	2.0652e-04	4.2758	2.8034e-03	4.6397
1/40	2.1214e-06	4.5570	7.8935e-06	4.7095	1.2005e-04	4.5454
1/60	3.0790e-07	4.7601	9.3316e-07	5.2661	1.3775e-05	5.3397

Table 1 numerically verifies Lemma 2.7, and shows that the truncation errors are $\mathcal{O}(h^5)$.

2.2 Effective fourth order discretization for space fractional derivatives

This subsection focuses on how to choose the parameters $p, q, r, s, \bar{p}, \bar{q}, \bar{r}, \bar{s}$ such that all the eigenvalues of the matrix $A_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}$ ($v = 3, 4$) have negative real parts; this means that the corresponding schemes work for space fractional derivatives. Since $B_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}$ is the transpose of $A_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}$, we don't need to discuss them separately.

LEMMA 2.8 (Quarteroni *et al.*, 2007, p. 28) A real matrix A of order n is positive definite if and only if its symmetric part $H = \frac{A+A^T}{2}$ is positive definite. Let $H \in \mathbb{R}^{n \times n}$ be symmetric, then H is positive definite if and only if the eigenvalues of H are positive.

LEMMA 2.9 (Quarteroni *et al.*, 2007, p. 184) If $A \in \mathbb{C}^{n \times n}$, let $H = \frac{A+A^H}{2}$ be the hermitian part of A , then for any eigenvalue λ of A , the real part $\Re(\lambda(A))$ satisfies

$$\lambda_{\min}(H) \leq \Re(\lambda(A)) \leq \lambda_{\max}(H),$$

where $\lambda_{\min}(H)$ and $\lambda_{\max}(H)$ are the minimum and maximum of the eigenvalues of H , respectively.

DEFINITION 2.9 (Chan & Jin, 2007, p. 13) Let $n \times n$ Toeplitz matrix T_n be of the following form:

$$T_n = \begin{bmatrix} t_0 & t_{-1} & \cdots & t_{2-n} & t_{1-n} \\ t_1 & t_0 & t_{-1} & \cdots & t_{2-n} \\ \vdots & t_1 & t_0 & \ddots & \vdots \\ t_{n-2} & \cdots & \ddots & \ddots & t_{-1} \\ t_{n-1} & t_{n-2} & \cdots & t_1 & t_0 \end{bmatrix};$$

i.e., $t_{i,j} = t_{i-j}$ and T_n is constant along its diagonals. Assume that the diagonals $\{t_k\}_{k=-n+1}^{n-1}$ are the Fourier coefficients of a function f , i.e.,

$$t_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx,$$

then the function f is called the generating function of T_n .

LEMMA 2.10 (Chan & Jin, 2007, p. 13-15) (Grenander-Szegö theorem) Let T_n be given by above matrix with a generating function f , where f is a 2π -periodic continuous real-valued functions defined on $[-\pi, \pi]$. Let $\lambda_{\min}(T_n)$ and $\lambda_{\max}(T_n)$ denote the smallest and largest eigenvalues of T_n , respectively. Then we have

$$f_{\min} \leq \lambda_{\min}(T_n) \leq \lambda_{\max}(T_n) \leq f_{\max},$$

where f_{\min} and f_{\max} is the minimum and maximum values of $f(x)$, respectively. Moreover, if $f_{\min} < f_{\max}$, then all eigenvalues of T_n satisfies

$$f_{\min} < \lambda(T_n) < f_{\max},$$

for all $n > 0$; In particular, if $f_{\min} > 0$, then T_n is positive definite.

LEMMA 2.11 Let $A_{p,q}^{v,\alpha}$ be given in (2.33), $(p,q) = (1,q)$, q is an integer ($q \neq 1$), $1 < \alpha < 2$ and denote that

$$H_{p,q}^v = \frac{A_{p,q}^{v,\alpha} + (A_{p,q}^{v,\alpha})^T}{2}, \quad v = 3, 4, \quad (2.36)$$

then the generating function of $H_{p,q}^v$ is

$$f_{p,q}^v(\alpha, x) = \left(2 \sin \frac{x}{2}\right)^\alpha (a_v^2 + b_v^2)^{\frac{\alpha}{2}} \cdot \left[w_p^v \cos\left(\alpha\left(\frac{x}{2} - \frac{\pi}{2} + \theta\right) - x\right) + w_q^v \cos\left(\alpha\left(\frac{x}{2} - \frac{\pi}{2} + \theta\right) - qx\right) \right], \quad (2.37)$$

16 of 22

where w_p^v and w_q^v are defined in (2.18), $\theta = -\arctan \frac{b_v}{a_v}$,

$$a_3 = \frac{1}{6}(11 - 7\cos(x) + 2\cos(2x)), \quad b_3 = \frac{1}{6}(7\sin(x) - 2\sin(2x));$$

and

$$a_4 = \frac{1}{12}(25 - 23\cos(x) + 13\cos(2x) - 3\cos(3x)), \quad b_4 = \frac{1}{12}(23\sin(x) - 13\sin(2x) + 3\sin(3x)).$$

Proof.

(1) For $(p, q) = (1, q)$, $q \leq 0$, we have $A_{p,q}^{v,\alpha} = w_p^v A_p^{v,\alpha} + w_q^v A_q^{v,\alpha}$ and

$$A_{p,q}^{v,\alpha} = \begin{bmatrix} \phi_1^{v,\alpha} & \phi_0^{v,\alpha} & & & \\ \phi_2^{v,\alpha} & \phi_1^{v,\alpha} & \phi_0^{v,\alpha} & & \\ \vdots & \ddots & \ddots & \ddots & \\ \phi_{N_x-2}^{v,\alpha} & \ddots & \ddots & \phi_1^{v,\alpha} & \phi_0^{v,\alpha} \\ \phi_{N_x-1}^{v,\alpha} & \phi_{N_x-2}^{v,\alpha} & \cdots & \phi_2^{v,\alpha} & \phi_1^{v,\alpha} \end{bmatrix},$$

with

$$\phi_k^{v,\alpha} = \begin{cases} w_p^v l_k^{v,\alpha}, & 0 \leq k \leq -q, \\ w_p^v l_k^{v,\alpha} + w_q^v l_{k+q-1}^{v,\alpha}, & k > -q. \end{cases}$$

The generating functions of $A_{p,q}^{v,\alpha}$ and $(A_{p,q}^{v,\alpha})^T$ are

$$f_{A_{p,q}^{v,\alpha}}(x) = \sum_{k=0}^{\infty} \phi_k^{v,\alpha} e^{i(k-1)x}, \quad \text{and} \quad f_{(A_{p,q}^{v,\alpha})^T}(x) = \sum_{k=0}^{\infty} \phi_k^{v,\alpha} e^{-i(k-1)x},$$

respectively. Then

$$f_{p,q}^v(\alpha, x) = \frac{f_{A_{p,q}^{v,\alpha}}(x) + f_{(A_{p,q}^{v,\alpha})^T}(x)}{2},$$

is the generating function of $H_{p,q}^v$. Since $f_{A_{p,q}^{v,\alpha}}(x)$ and $f_{(A_{p,q}^{v,\alpha})^T}(x)$ are mutually conjugated, then $f_{p,q}^v(\alpha, x)$ is a 2π -periodic continuous real-valued functions defined on $[-\pi, \pi]$. Moreover, $f_{p,q}^v(\alpha, x)$ is an even function, so we just need to consider its principal value on $[0, \pi]$. According to the following equations

$$(1 - e^{\pm ix})^\alpha = \left(2\sin\frac{x}{2}\right)^\alpha e^{\pm i\alpha(\frac{x}{2} - \frac{\pi}{2})}, \quad (a_v - b_v i)^\alpha = (a_v^2 + b_v^2)^{\frac{\alpha}{2}} e^{i\alpha\theta},$$

where $v = 3, 4$ and $\theta = -\arctan \frac{b_v}{a_v}$, then we have

$$\begin{aligned} f_{A_{p,q}^{v,\alpha}}(x) &= \sum_{k=0}^{\infty} \phi_k^{3,\alpha} e^{i(k-1)x} = w_p^3 e^{-ix} \sum_{k=0}^{\infty} l_k^{3,\alpha} e^{ikx} + w_q^3 e^{-iqx} \sum_{k=0}^{\infty} l_k^{3,\alpha} e^{ikx} \\ &= w_p^3 e^{-ix} (1 - e^{ix})^\alpha \left(\frac{11}{6} - \frac{7}{6}e^{ix} + \frac{1}{3}e^{2ix}\right)^\alpha + w_q^3 e^{-iqx} (1 - e^{ix})^\alpha \left(\frac{11}{6} - \frac{7}{6}e^{ix} + \frac{1}{3}e^{2ix}\right)^\alpha \\ &= w_p^3 e^{-ix} (1 - e^{ix})^\alpha (a_3 - b_3 i)^\alpha + w_q^3 e^{-iqx} (1 - e^{ix})^\alpha (a_3 - b_3 i)^\alpha \\ &= w_p^3 e^{-ix} \left(2\sin\frac{x}{2}\right)^\alpha e^{i\alpha(\frac{x}{2} - \frac{\pi}{2})} (a_3^2 + b_3^2)^{\frac{\alpha}{2}} e^{i\alpha\theta} + w_q^3 e^{-iqx} \left(2\sin\frac{x}{2}\right)^\alpha e^{i\alpha(\frac{x}{2} - \frac{\pi}{2})} (a_3^2 + b_3^2)^{\frac{\alpha}{2}} e^{i\alpha\theta} \\ &= \left(2\sin\frac{x}{2}\right)^\alpha (a_3^2 + b_3^2)^{\frac{\alpha}{2}} \left\{ w_p^3 e^{i[(\alpha(\frac{x}{2} - \frac{\pi}{2}) - \theta) - x]} + w_q^3 e^{i[(\alpha(\frac{x}{2} - \frac{\pi}{2}) + \theta) - qx]} \right\}; \end{aligned}$$

respectively. Then

$$f_{p,q}^v(\alpha, x) = \frac{f_{A_{p,q}^{v,\alpha}}(x) + f_{(A_{p,q}^{v,\alpha})^T}(x)}{2},$$

is the generating function of $H_{p,q}^v$.

By the similar way, for $q \geq 2$, their exists

$$f_{p,q}^v(\alpha, x) = \left(2 \sin \frac{x}{2}\right)^\alpha (a_v^2 + b_v^2)^{\frac{\alpha}{2}} \cdot \left[w_p^v \cos \left(\alpha \left(\frac{x}{2} - \frac{\pi}{2} + \theta \right) - x \right) + w_q^v \cos \left(\alpha \left(\frac{x}{2} - \frac{\pi}{2} + \theta \right) - qx \right) \right].$$

It can be noted that $f_{p,q}^v(\alpha, x)$ has the same form when $q \leq 0$ and $q \geq 2$, $p = 1$, Then $H_{p,q}^v$ has the generating function

$$f_{p,q}^v(\alpha, x) = \left(2 \sin \frac{x}{2}\right)^\alpha (a_v^2 + b_v^2)^{\frac{\alpha}{2}} \cdot \left[w_p^v \cos \left(\alpha \left(\frac{x}{2} - \frac{\pi}{2} + \theta \right) - x \right) + w_q^v \cos \left(\alpha \left(\frac{x}{2} - \frac{\pi}{2} + \theta \right) - qx \right) \right].$$

□

THEOREM 2.10 (Case $v = 3, 4$; Effective 4th order schemes) Let $A_{v,\alpha} =: A_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}$ with $1 < \alpha < 2$, be given in (2.33) and $(p, q, r, s, \bar{p}, \bar{q}, \bar{r}, \bar{s}) = (1, -1, 1, 2, 1, -1, 1, 3)$. Then any eigenvalue λ of $A_{v,\alpha}$ satisfies

$$\Re(\lambda(A_{v,\alpha})) < 0, \quad v = 3, 4,$$

and the matrices $A_{v,\alpha}$ and $A_{v,\alpha}^T$ are negative definite.

Proof. Take

$$H_\alpha^v = \frac{A_{v,\alpha} + A_{v,\alpha}^T}{2} = w_{p,q,r,s}^v w_{p,q}^v H_{p,q}^v + w_{p,q,r,s}^v w_{r,s}^v H_{r,s}^v + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{p},\bar{q}}^v H_{\bar{p},\bar{q}}^v + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{r},\bar{s}}^v H_{\bar{r},\bar{s}}^v,$$

where $H_{p,q}^v$, $H_{r,s}^v$, $H_{\bar{p},\bar{q}}^v$ and $H_{\bar{r},\bar{s}}^v$ are defined in (2.36). Then

$$f^v(\alpha, x) = w_{p,q,r,s}^v w_{p,q}^v f_{p,q}^v(\alpha, x) + w_{p,q,r,s}^v w_{r,s}^v f_{r,s}^v(\alpha, x) + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{p},\bar{q}}^v f_{\bar{p},\bar{q}}^v(\alpha, x) + w_{\bar{p},\bar{q},\bar{r},\bar{s}}^v w_{\bar{r},\bar{s}}^v f_{\bar{r},\bar{s}}^v(\alpha, x),$$

is the generating function of H_α^v , where $f_{p,q}^v(\alpha, x)$, $f_{r,s}^v(\alpha, x)$, $f_{\bar{p},\bar{q}}^v(\alpha, x)$ and $f_{\bar{r},\bar{s}}^v(\alpha, x)$ are given by (2.37). And there exists $f^v(\alpha, x) \leq 0$ (see Figs. 1-2), when $(p, q, r, s, \bar{p}, \bar{q}, \bar{r}, \bar{s}) = (1, -1, 1, 2, 1, -1, 1, 3)$.



Since $f^v(\alpha, x)$ is not identically zero for $\alpha \in (1, 2)$, from Lemma 2.10, it implies that $\lambda(H_\alpha^v) < 0$, and H_α^v is negative definite. Then we obtain $\Re(\lambda(A_{v,\alpha})) < 0$ by Lemma 2.9, and the matrices $A_{v,\alpha}$ and $A_{v,\alpha}^T$ are negative definite from Lemma 2.8. □

3. Simple application to space fractional diffusion equation

Similar to the discussions in [Chen & Deng (2013)], in this section, we apply the 4th order discretizations to solve the following fractional diffusion equation with variable coefficients

$$\frac{\partial u(x,t)}{\partial t} = d_+(x) {}_{x_L} D_x^\alpha u(x,t) + d_-(x) {}_x D_{x_R}^\alpha u(x,t) + f(x,t). \quad (3.1)$$

In the time direction, the Crank-Nicolson scheme is used. The 4th order left fractional approximation operator (2.29), and right fractional approximation operator (2.34) are respectively used to discretize the left Riemann-Liouville fractional derivative, and right Riemann-Liouville fractional derivative.

FIG. 1. $f^v(\alpha, x)$ for $v = 3$ FIG. 2. $f^v(\alpha, x)$ for $v = 4$

Let the mesh points $x_i = x_L + ih$, $i = -m, \dots, 0, 1, \dots, N_x - 1, N_x, \dots, N_x + m$, with m in (2.27) and $t_n = n\tau$, $0 \leq n \leq N_t$, where $h = (x_R - x_L)/N_x$, $\tau = T/N_t$, i.e., h is the uniform space step size and τ the time step size. Taking u_i^n as the approximated value of $u(x_i, t_n)$ and $d_{+,i} = d_+(x_i)$, $d_{-,i} = d_-(x_i)$, $f_i^{n+1/2} = f(x_i, t_{n+1/2})$, where $t_{n+1/2} = (t_n + t_{n+1})/2$.

The full discretization of (3.1) has the following form [Chen & Deng (2013)]

$$\left[I - \frac{\tau}{2h\alpha} (D_+ A_{v,\alpha} + D_- A_{v,\alpha}^T) \right] U^{n+1} = \left[I + \frac{\tau}{2h\alpha} (D_+ A_{v,\alpha} + D_- A_{v,\alpha}^T) \right] U^n + \tau F^{n+1/2}, \quad (3.2)$$

where $A_{v,\alpha} =: A_{p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}}^{v,\alpha}$ with $1 < \alpha < 2$, $v = 3, 4$, are given in (2.33), and

$$D_+ = \begin{bmatrix} d_{+,0} & & & & \\ & d_{+,1} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & d_{+,N_x} \end{bmatrix}, \quad D_- = \begin{bmatrix} d_{-,0} & & & & \\ & d_{-,1} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & d_{-,N_x} \end{bmatrix}, \quad (3.3)$$

and

$$U^n = [u_0^n, u_1^n, \dots, u_{N_x}^n]^T, \quad F^{n+1/2} = [f_0^{n+1/2}, f_1^{n+1/2}, \dots, f_{N_x}^{n+1/2}]^T.$$

By the same way given in [Chen & Deng (2013)], we can theoretically prove that the difference scheme is unconditionally stable and 4th order convergent in space directions and 2nd order convergent in time direction; the proofs are omitted here.

THEOREM 3.1 (Chen & Deng (2013)) Let $D_- = \kappa_\alpha D_+$, where κ_α is any given nonnegative constant, then the difference scheme (3.2) with $\alpha \in (1, 2)$ is unconditionally stable.

THEOREM 3.2 (Chen & Deng (2013)) Let $u(x_i, t_n)$ be the exact solution of (3.1) with $\alpha \in (1, 2)$, and u_i^n be the solution of the finite difference scheme (3.2). If $D_- = \kappa_\alpha D_+$, then there is a positive constant C such that

$$\|u(x_i, t_n) - u_i^n\| \leq C(\tau^2 + h^4), \quad i = 0, 2, \dots, N_x; n = 0, 1, \dots, N_t,$$

where κ_α is any given nonnegative constant.

4. Numerical Results

In this section, we numerically verify the above theoretical results including convergence rates and numerical stability. And the l_∞ norm is used to measure the numerical errors.

Consider the fractional diffusion equation (3.1) [Chen & Deng (2013)] in the domain $0 < x < 2$, $0 < t \leq 1$, with the variable coefficients $d_+(x) = x^\alpha$, $d_-(x) = 2x^\alpha$, the forcing function

$$f(x,t) = \cos(t+1)x^4(2-x)^4 - x^\alpha \sin(t+1) \left[\frac{\Gamma(9)}{\Gamma(9-\alpha)}(x^{8-\alpha} + 2(2-x)^{8-\alpha}) \right. \\ \left. - 8 \frac{\Gamma(8)}{\Gamma(8-\alpha)}(x^{7-\alpha} + 2(2-x)^{7-\alpha}) + 24 \frac{\Gamma(7)}{\Gamma(7-\alpha)}(x^{6-\alpha} + 2(2-x)^{6-\alpha}) \right. \\ \left. - 32 \frac{\Gamma(6)}{\Gamma(6-\alpha)}(x^{5-\alpha} + 2(2-x)^{5-\alpha}) + 16 \frac{\Gamma(5)}{\Gamma(5-\alpha)}(x^{4-\alpha} + 2(2-x)^{4-\alpha}) \right],$$

and the initial condition $u(x,0) = \sin(1)x^4(2-x)^4$, the boundary conditions $u(0,t) = u(1,t) = 0$, and the exact solution of the equation is $u(x,t) = \sin(t+1)x^4(2-x)^4$.

Table 2. The maximum errors and convergent orders for the scheme (3.2) of the fractional diffusion equation (3.1) at $t=1$ and $\tau = h^2$, where $(p,q,r,s,\bar{p},\bar{q},\bar{r},\bar{s}) = (1,-1,1,2,1,-1,1,3)$, $v = 3,4$.

$v = 3, h$	$\alpha = 1.1$	Rate	$\alpha = 1.5$	Rate	$\alpha = 1.8$	Rate
1/10	1.8349e-02		2.1073e-02		2.3337e-02	
1/20	1.4015e-03	3.7107	1.8381e-03	3.5191	2.3106e-03	3.3362
1/40	8.8517e-05	3.9849	1.2004e-04	3.9367	1.6131e-04	3.8404
1/80	5.2342e-06	4.0799	7.5382e-06	3.9931	1.0478e-05	3.9443
$v = 4, h$	$\alpha = 1.1$	Rate	$\alpha = 1.5$	Rate	$\alpha = 1.8$	Rate
1/10	1.7241e-02		9.6037e-03		5.8735e-03	
1/20	7.9269e-04	4.4430	5.2600e-04	4.1905	3.4793e-04	4.0774
1/40	3.4558e-05	4.5197	2.4926e-05	4.3994	2.1158e-05	4.0395
1/80	1.4824e-06	4.5430	1.1512e-06	4.4364	1.3045e-06	4.0196

Table 2 shows the maximum errors at time $t = 1$, and the time and space stepsizes are taken as $\tau = h^2$. The numerical results confirm the $\mathcal{O}(\tau^2 + h^4)$ order convergence.

5. Conclusions

For solving classical differential equations, usually people think that the high order schemes can reduce computational cost for getting some requested accuracy, i.e., they think that the improved accuracy can overtake the increased workload. For the issue of computational cost of fractional differential equations, the high order schemes play more fundamental role; since they can greatly increase the accuracy but without adding new cost. As the sequel of [Chen & Deng (2013)], based on Lubich's 3rd and 4th operators, this paper further provides new 4th order schemes for space fractional derivatives. The effectiveness of the new discretizations is verified theoretically and numerically.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under Grant No. 11271173, and the Program for New Century Excellent Talents in University under Grant No. NCET-09-0438.

References

- CHAN, R. H. & JIN, X. Q. (2007) *An Introduction to Iterative Toeplitz Solvers*. SIAM.
- CHEN, M. H. & DENG, W. H. (2013) WSLD operators: A class of fourth order difference approximations for space Riemann-Liouville derivative. arXiv: 1304.7425 [math.NA].
- CHEN, M. H., WANG, Y. T., CHENG, X. & DENG, W. H. (2012) Second-order LOD multigrid method for multidimensional Riesz fractional diffusion equation. arXiv:1301.2643v1 [math.NA].
- DENG, W. H. (2008) Finite element method for the space and time fractional Fokker-Planck equation. *SIAM J. Numer. Anal.*, **47**, 204-226.
- ERVIN, V. J. & ROOP, J. P. (2006) Variational formulation for the stationary fractional advection dispersion equation. *Numer. Methods Partial Differential Equations.*, **22**, 558-576.
- FAN, S. J. (1989) A new extracting formula and a new distinguishing means on the one variable cubic equation. *Natural science journal of Hainan teachers college, China.*, **2**, 2, 91-98.
- HENRICI, P. (1962) *Discrete Variable Methods in Ordinary Differential Equations*. New York: John Wiley.
- KILBAS, A., SRIVASTAVA, M. & TRUJILLO, J. (2006) *Theory and Applications of Fractional Differential Equations*. New York.
- LI, X. J. & XU, C. J. (2009) A space-time spectral method for the time fractional diffusion equation. *SIAM J. Numer. Anal.*, **47**, 2108-2131.
- LUBICH, CH. (1986) Discretized fractional calculus. *SIAM J. Math. Anal.*, **17**, 704-719.
- MEERSCHAERT, M. M. & TADJERAN, C. (2004) Finite difference approximations for fractional advection-dispersion flow equations. *J. Comput. Appl. Math.*, **172**, 65-77.
- ORTIGUEIRA, M. D. (2006) Riesz potential operators and inverses via fractional centred derivatives. *International J. Math. Math. Sci.*, **2006**, 1-12.
- PODLUBNY, I. (1999) *Fractional Differential Equations*. New York: Academic Press.
- POLYANIN, A. D. & MANZHIROV, A. V. (2007) *Handbook of Mathematics for Engineers and Scientists*. Chapman & Hall/CRC Press, New York.
- QUARTERONI, A., SACCO, R. & SALERI, F. (2007) *Numerical Mathematics, 2nd ed.* Springer.
- SOUSA, E. & LI, C. (2011) A weighted finite difference method for the fractional diffusion equation based on the Riemann-Liouville derivative. arXiv:1109.2345v1 [math.NA].
- SUN, Z. Z. & WU, X. N. (2006) A fully discrete difference scheme for a diffusion-wave system. *Appl. Numer. Math.*, **56**, 193-209.

- TIAN, W. Y., ZHOU, H. & DENG, W. H. (2012) A class of second order difference approximations for solving space fractional diffusion Equations. arXiv:1204.4870v1 math.NA.
- ZHOU, H., TIAN, W. Y. & DENG, W. H. (2013) Quasi-compact finite difference schemes for space fractional diffusion equations, *J. Sci. Comput.*, **56**(1), 45–66.
- ZHUANG, P., LIU, F., ANH, V. & TURNER, I. (2009) Numerical methods for the variable-order fractional advection-diffusion equation with a nonlinear source term. *SIAM J. Numer. Anal.*, **47**, 1760–1781.