

# An Upper Bound for the Logarithmic Capacity of Two Intervals\*

Klaus Schiefermayr<sup>†</sup>

## Abstract

The logarithmic capacity (also called Chebyshev constant or transfinite diameter) of two real intervals  $[-1, \alpha] \cup [\beta, 1]$  has been given explicitly with the help of Jacobi's elliptic and theta functions already by Achieser in 1930. By proving several inequalities for these elliptic and theta functions, an upper bound for the logarithmic capacity in terms of elementary functions of  $\alpha$  and  $\beta$  is derived.

*Mathematics Subject Classification (2000):* 33E05, 31A15

*Keywords:* Chebyshev constant, Logarithmic capacity, Jacobi's elliptic functions, Jacobi's theta functions, Transfinite diameter, Two intervals

## 1 Introduction

Let  $C$  be a compact set in the complex plane and let  $\mathbb{P}_n$  be the set of all polynomials of degree less or equal  $n$ . Then  $L_n(C)$ , defined by

$$L_n(C) := \inf_{p_{n-1} \in \mathbb{P}_{n-1}} \sup_{z \in C} |z^n - p_{n-1}(z)|, \quad (1)$$

is usually called the *minimum deviation* of degree  $n$  on  $C$ . The *logarithmic capacity* (or *Chebyshev constant*) of  $C$  is then defined by the limit

$$\text{cap}(C) := \lim_{n \rightarrow \infty} \sqrt[n]{L_n(C)}. \quad (2)$$

The logarithmic capacity of  $C$  can be given in a completely different way by

$$\text{cap}(C) = \lim_{n \rightarrow \infty} \sup_{z_i, z_j \in C} \left( \prod_{1 \leq i < j \leq n} |z_i - z_j| \right)^{\frac{2}{n(n-1)}} \quad (3)$$

and, in this connection, is also called the *transfinite diameter*. For further reading and weighted analogues concerning this constant, see [15].

In this paper, we consider the case of two intervals, i.e.,

$$C = C_{\alpha, \beta} := [-1, \alpha] \cup [\beta, 1], \quad -1 < \alpha < \beta < 1. \quad (4)$$

The logarithmic capacity of two intervals  $C_{\alpha, \beta}$  can only be given with the help of Jacobi's elliptic and theta functions (see Theorem 1). The purpose of this paper is to give an upper bound for  $\text{cap}(C_{\alpha, \beta})$  in terms of elementary functions of  $\alpha$  and  $\beta$ .

\*published in: Complex Variables and Elliptic Equations **53** (2008), 65–75.

<sup>†</sup>University of Applied Sciences Upper Austria, School of Engineering and Environmental Sciences, Stelzhamerstrasse 23, 4600 Wels, Austria, KLAUS.SCHIEFERMAYR@FH-WELS.AT

Let  $K \equiv K(k)$  and  $E \equiv E(k)$  be the complete elliptic integral of the first and second kind, respectively, where  $0 < k < 1$  is known as the modulus of  $K$  and  $E$ . Let  $k' := \sqrt{1 - k^2}$ ,  $K' \equiv K'(k) := K(k')$ , and let  $\operatorname{sn}(u) \equiv \operatorname{sn}(u, k)$ ,  $\operatorname{cn}(u) \equiv \operatorname{cn}(u, k)$ ,  $\operatorname{dn}(u) \equiv \operatorname{dn}(u, k)$  be Jacobi's elliptic functions with respect to the modulus  $k$ . Further, let  $\Theta(u) \equiv \Theta(u, k)$ ,  $H(u) \equiv H(u, k)$ ,  $H_1(u) \equiv H_1(u, k)$  and  $\Theta_1(u) \equiv \Theta_1(u, k)$  be the four theta functions of Jacobi (old notation).

Further, we need Jacobi's zeta function  $\operatorname{zn}(u) \equiv \operatorname{zn}(u, k)$  given by

$$\operatorname{zn}(u) = \int_0^u \operatorname{dn}^2(v) \, dv - \frac{uE}{K} = \frac{\Theta'(u)}{\Theta(u)}. \quad (5)$$

For the definitions and many important properties of these functions see, e.g., [9], [13], [14, chapter X] or [22].

A formula for the logarithmic capacity of two intervals in terms of Jacobi's elliptic and theta functions has been derived by N.I. Achieser [2]. For the background of [2] and some related results, see [23], [24], [1], [3], [16], and [17].

**Theorem 1** (Achieiser [2]). *Let  $C_{\alpha, \beta} := [-1, \alpha] \cup [\beta, 1]$ ,  $-1 < \alpha < \beta < 1$ , let the modulus  $k$ ,  $0 < k < 1$ , be given by*

$$k^2 := \frac{2(\beta - \alpha)}{(1 - \alpha)(1 + \beta)} \quad (6)$$

and let  $0 < \lambda < 1$  be such that

$$\operatorname{sn}^2(\lambda K) = \frac{1 - \alpha}{2}. \quad (7)$$

Then

$$\operatorname{cap}(C_{\alpha, \beta}) = \frac{\Theta^4(0)}{2 \operatorname{dn}^2(\lambda K) \Theta^4(\lambda K)} = \frac{2k'^2 K^2}{\pi^2 \operatorname{dn}^2(\lambda K) \Theta^4(\lambda K)}. \quad (8)$$

The problem in operating with formula (8) is the difficult connection of the modulus  $k$  and the parameter  $\lambda$  with the values  $\alpha$  and  $\beta$  via formulae (6) and (7), where the modulus  $k$  appear both in the term  $K \equiv K(k)$  and as the parameter of the elliptic function  $\operatorname{sn}(u) \equiv \operatorname{sn}(u, k)$ . With the help of some inequalities and monotonicity properties of Jacobi's elliptic and theta functions given and proved in Section 3, we are able to prove an upper bound for  $\operatorname{cap}(C_{\alpha, \beta})$  in terms of elementary functions of  $\alpha$  and  $\beta$  in Section 2. Concluding this section, we collect some useful properties of  $\alpha$  and  $\beta$  (as functions of  $k$  and  $\lambda$ ) and of the capacity  $\operatorname{cap}(C_{\alpha, \beta})$  in the following remark.

**Remark.**

(i) By (6) and (7),

$$\operatorname{cn}^2(\lambda K) = \frac{1 + \alpha}{2}, \quad \operatorname{dn}^2(\lambda K) = \frac{1 + \alpha}{1 + \beta} \quad (9)$$

and

$$k'^2 = \frac{(1 + \alpha)(1 - \beta)}{(1 - \alpha)(1 + \beta)}. \quad (10)$$

(ii) By (6) and (7), there is a one-to-one correspondence between  $\{\alpha, \beta\}$ ,  $-1 < \alpha < \beta < 1$ , and  $\{k, \lambda\}$ ,  $0 < k < 1$ ,  $0 < \lambda < 1$ . Moreover, for  $\alpha \equiv \alpha(k, \lambda)$  and  $\beta \equiv \beta(k, \lambda)$ , we get

$$\begin{aligned} \alpha(k, \lambda) &= 1 - 2 \operatorname{sn}^2(\lambda K), \\ \beta(k, \lambda) &= \frac{2 \operatorname{cn}^2(\lambda K)}{\operatorname{dn}^2(\lambda K)} - 1 = 2 \operatorname{sn}^2((1 + \lambda)K) - 1. \end{aligned} \quad (11)$$

- (iii) Let  $0 < k < 1$  be fixed. By (11), both,  $\alpha(k, \lambda)$  and  $\beta(k, \lambda)$ , are strictly monotone decreasing functions of  $\lambda$  with

$$\lim_{\lambda \rightarrow 0} \alpha(k, \lambda) = \lim_{\lambda \rightarrow 0} \beta(k, \lambda) = 1, \quad \lim_{\lambda \rightarrow 1} \alpha(k, \lambda) = \lim_{\lambda \rightarrow 1} \beta(k, \lambda) = -1.$$

- (iv) Let  $0 < \lambda < 1$  be fixed. By (11) and [11, Sect. 1 & 4],  $\alpha(k, \lambda)$  and  $\beta(k, \lambda)$  is a strictly monotone decreasing and increasing function of  $k$ , respectively. Moreover,

$$\lim_{k \rightarrow 0} \alpha(k, \lambda) = \lim_{k \rightarrow 0} \beta(k, \lambda) = \cos(\lambda\pi), \quad -\lim_{k \rightarrow 1} \alpha(k, \lambda) = \lim_{k \rightarrow 1} \beta(k, \lambda) = 1,$$

i.e., for the limiting case  $k \rightarrow 0$ , we get  $C_{\alpha, \beta} \rightarrow [-1, 1]$ , and for  $k \rightarrow 1$ ,  $C_{\alpha, \beta}$  vanishes.

- (v) By (11), for  $\lambda = \frac{1}{2}$ ,  $0 < \lambda < \frac{1}{2}$ ,  $\frac{1}{2} < \lambda < 1$ , there is  $\alpha + \beta = 0$ ,  $\alpha + \beta > 0$ ,  $\alpha + \beta < 0$ , respectively.

- (vi) If  $\lambda$  changes to  $1 - \lambda$ , then  $\{\alpha, \beta\}$  changes to  $\{-\beta, -\alpha\}$  and the capacity remains the same, i.e.,

$$\text{cap}(C_{\alpha, \beta}) = \frac{\Theta^4(0)}{2 \text{dn}^2((1 - \lambda)K) \Theta^4((1 - \lambda)K)} = \frac{2k'^2 K^2}{\pi^2 \text{dn}^2((1 - \lambda)K) \Theta^4((1 - \lambda)K)}. \quad (12)$$

- (vii) The capacity is monotone, i.e., for  $-1 < \alpha_1 \leq \alpha_2 < \beta_2 \leq \beta_1 < 1$ ,

$$\text{cap}(C_{\alpha_1, \beta_1}) \leq \text{cap}(C_{\alpha_2, \beta_2}).$$

- (viii) Let  $C$  be any compact set in the complex plane, then for  $a > 0$

$$\text{cap}(aC) = a \text{cap}(C). \quad (13)$$

- (ix) By (vii), for  $\alpha + \beta \geq 0$ , there is a trivial lower bound for  $\text{cap}(C_{\alpha, \beta})$  given by

$$\text{cap}(C_{\alpha, \beta}) \geq \text{cap}([-1, -\beta] \cup [\beta, 1]) = \frac{1}{2} \sqrt{1 - \beta^2}. \quad (14)$$

- (x) By (vii), for  $\alpha + \beta \geq 0$  and  $\alpha < 0$ , there is a trivial upper bound for  $\text{cap}(C_{\alpha, \beta})$  given by

$$\text{cap}(C_{\alpha, \beta}) \leq \text{cap}([-1, \alpha] \cup [-\alpha, 1]) = \frac{1}{2} \sqrt{1 - \alpha^2}. \quad (15)$$

Again by (vii), another trivial upper bound for all  $-1 < \alpha < \beta < 1$  is

$$\text{cap}(C_{\alpha, \beta}) \leq \text{cap}([-1, 1]) = \frac{1}{2}. \quad (16)$$

- (xi) Another lower bound can be extracted from [18]

$$\text{cap}(C_{\alpha, \beta}) \geq \text{cap}([-1, \alpha]) + \text{cap}([\beta, 1]) = \frac{1 + \alpha}{4} + \frac{1 - \beta}{4}. \quad (17)$$

- (xii) Finally, another upper bound can be derived from an inequality of Gillis [12], which, in our special case of two intervals, reads as follows:

$$\text{cap}(C_{\alpha, \beta}) \leq 2 \exp\left(\frac{\log \frac{1+\alpha}{8} \log \frac{1-\beta}{8}}{\log \frac{(1+\alpha)(1-\beta)}{64}}\right) \quad (18)$$

(xiii) Let  $\alpha \in (-1, 1)$  be fixed. Then, by (17) and (18),

$$\lim_{\beta \rightarrow 1} \text{cap}(C_{\alpha, \beta}) = \frac{1 + \alpha}{4} = \text{cap}([-1, \alpha]). \quad (19)$$

Note that this is not true for the minimum deviation  $L_n(C_{\alpha, \beta})$ .

(xiv) Let  $\varphi, \psi \in (0, \pi)$  such that  $\cos \varphi = \alpha$ ,  $\cos \psi = \beta$ , and let

$$A := \{z \in \mathbb{C} : |z| = 1, \arg z \in [-\psi, \psi] \cup [\varphi, 2\pi - \varphi]\}, \quad (20)$$

i.e.,  $C_{\alpha, \beta}$  is the projection of  $A$  onto the real axis. Then, by a result of Robinson [20],

$$\text{cap}(A) = \sqrt{2 \text{cap}(C_{\alpha, \beta})}. \quad (21)$$

## 2 Main Results

Let us first give a lower bound for the logarithmic capacity of two intervals  $C_{\alpha, \beta}$ , proved by Alexander Yu. Solynin [21, Sect. 2.2] (in fact he derived a lower bound for the logarithmic capacity of several intervals). In our notation, his result reads as follows:

**Theorem 2** (Solynin [21]). *Let  $-1 < \alpha < \beta < 1$  and define*

$$\varphi := \arccos(\alpha), \quad \psi := \arccos(\beta),$$

then

$$\text{cap}(C_{\alpha, \beta}) \geq \frac{1}{2} \max_{\delta \in [\psi, \varphi]} \left[ \left( \sin\left(\frac{\psi\pi}{2\delta}\right) \right)^{2\delta^2/\pi^2} \left( \sin\left(\frac{(\pi - \varphi)\pi}{2(\pi - \delta)}\right) \right)^{2(\pi - \delta)^2/\pi^2} \right]. \quad (22)$$

**Remark.** Numerical calculations suggest that the optimal  $\delta$  in (22), which cannot be calculated analytically, may be roughly approximated by  $\delta = (\psi + \varphi)/2$ . With the optimal  $\delta$ , the lower bound (22) is excellent, but also for  $\delta = (\psi + \varphi)/2$ , the lower bound (22) is very good.

In addition to (22), we give another lower bound for the capacity of  $C_{\alpha, \beta}$  in very simple functions of  $\alpha$  and  $\beta$ .

**Theorem 3.** *Let  $-1 < \alpha < \beta < 1$ , then*

$$\text{cap}(C_{\alpha, \beta}) \geq \frac{\sqrt{k'}}{1 + k'} = \frac{\sqrt[4]{(1 - \alpha^2)(1 - \beta^2)}}{\sqrt{(1 - \alpha)(1 + \beta)} + \sqrt{(1 + \alpha)(1 - \beta)}}, \quad (23)$$

where equality is attained for  $\alpha + \beta = 0$ .

*Proof.* Let  $0 < k < 1$  be fixed, then, by (8) and Lemma 3,  $\text{cap}(C_{\alpha, \beta})$  is strictly monotone decreasing in  $\lambda$ ,  $0 \leq \lambda \leq \frac{1}{2}$ . Thus, by Lemma 2 and  $\text{dn}(\frac{K}{2}) = \sqrt{k'}$ ,

$$\text{cap}(C_{\alpha, \beta}) \geq \frac{2k'^2 K^2}{\pi^2 \text{dn}^2(\frac{K}{2}) \Theta^4(\frac{K}{2})} = \frac{\sqrt{k'}}{1 + k'}.$$

For  $\frac{1}{2} < \lambda < 1$ , the inequality follows by (12).  $\square$

**Remark.** For any  $\alpha, \beta$ ,  $-1 < \alpha < \beta < 1$ ,  $\alpha + \beta \geq 0$ , the lower bound in (23) is greater (i.e. better) than the lower bound in (14).

The next theorem contains the main result of the paper, an upper bound for the logarithmic capacity of two intervals  $C_{\alpha, \beta}$ .

**Theorem 4.** Let  $-1 < \alpha < \beta < 1$  and let  $k \in (0, 1)$  and  $\lambda \in (0, 1)$  be given by (6) and (7), respectively, then

$$\text{cap}(C_{\alpha, \beta}) \leq \frac{1}{2} \text{dn}^2(\lambda K) \exp\left(2(E/K - k'^2) \log^2 \frac{1 + \text{sn}(\lambda K)}{\text{cn}(\lambda K)}\right), \quad (24)$$

where equality is attained for the limiting case  $\lambda \rightarrow 0$ .

*Proof.*

$$\begin{aligned} \text{cap}(C_{\alpha, \beta}) &= \frac{1}{2 \text{dn}^2((1 - \lambda)K)} \left( \frac{\Theta(0)}{\Theta((1 - \lambda)K)} \right)^4 \quad \text{by (12)} \\ &\leq \frac{k'^2}{2 \text{dn}^2((1 - \lambda)K)} \exp(2(E/K - k'^2)\lambda^2 K^2) \quad \text{by Lemma 7} \\ &\leq \frac{1}{2} \text{dn}^2(\lambda K) \exp(2(E/K - k'^2) \log^2 \frac{1 + \text{sn}(\lambda K)}{\text{cn}(\lambda K)}) \quad \text{by Lemma 5} \end{aligned}$$

□

**Remark.**

- (i) Upper bound (24) is only good in case of  $\alpha + \beta \geq 0$ , i.e.,  $0 < \lambda \leq \frac{1}{2}$ . For  $\alpha + \beta < 0$ , one has to replace  $\{\alpha, \beta\}$  by  $\{-\beta, -\alpha\}$ .
- (ii) Analogously to (24), we also derived a lower bound for  $\text{cap}(C_{\alpha, \beta})$ , which turned out to be less than the lower bound (22), thus we dismissed it.
- (iii) In Figure 1, we compare the capacity  $\text{cap}(C_{\alpha, \beta})$  with the upper bounds (24) and (18). The graphs of these three functions are plotted for  $\alpha \in \{\pm 0.7, \pm 0.4, \pm 0.1\}$  and  $|\alpha| \leq \beta \leq 1$ .
- (iv) By (6), (7), (9), and (10), inequality(24) of Theorem 4 may be written in the form

$$\text{cap}(C_{\alpha, \beta}) \leq \frac{1 + \alpha}{2(1 + \beta)} \exp\left[2\left(\frac{E}{K} - \frac{(1 + \alpha)(1 - \beta)}{(1 - \alpha)(1 + \beta)}\right) \log^2 \frac{\sqrt{2} + \sqrt{1 - \alpha}}{\sqrt{1 + \alpha}}\right] \quad (25)$$

In this bound, there appear  $K \equiv K(k)$  and  $E \equiv E(k)$ , which cannot be expressed by elementary functions of  $k$ . Nevertheless, there exists a couple of inequalities in the literature, see [4], [5], [6], [7], [8], [10], and [19], involving elementary functions of  $k$ , from which we collect the best ones in the following lemma. Hence, inequality (25), together with (26) and (29), using (6) and (10), gives an upper bound for  $\text{cap}(C_{\alpha, \beta})$  in terms of elementary functions of  $\alpha$  and  $\beta$ .

**Lemma 1.** Let  $0 < k < 1$ .




Figure 1: The capacity  $\text{cap}(C_{\alpha,\beta})$  (solid line) with the upper bounds (24) (dashed line) and (18) (dotted line) for  $\alpha \in \{-0.7, -0.4, -0.1, 0.1, 0.4, 0.7\}$  and  $|\alpha| \leq \beta \leq 1$

(i) For the complete elliptic integral of the first kind, we have the inequalities

$$\max\{K_1, K_2\} < K(k) < \min\{K_3, K_4, K_5\}, \quad (26)$$

where

$$\begin{aligned} K_1 &:= \frac{\pi}{2} \left( \frac{\tanh^{-1}(k)}{k} \right)^{3/4}, \\ K_2 &:= \left( 1 + \frac{k'^2}{4} \right) \log\left(\frac{4}{k'}\right) - \frac{k'^2}{4}, \end{aligned} \quad (27)$$

and

$$\begin{aligned} K_3 &:= \frac{\pi}{2} \left( \frac{\frac{3}{4} \log(k')}{k' - 1} + \frac{1}{2(1 + k')} \right), \\ K_4 &:= \log\left(\frac{4}{k'} + (e^{\pi/2} - 4)(k')^\gamma\right), \quad \gamma := \frac{4 - \frac{\pi}{4} e^{\pi/2}}{e^{\pi/2} - 4}, \\ K_5 &:= \left( 1 + \frac{k'^2}{4} \right) \log\left(\frac{4}{k'}\right). \end{aligned} \quad (28)$$

(ii) For the complete elliptic integral of the second kind, we have the inequalities

$$E_1 < E(k) < E_2, \quad (29)$$

where

$$E_1 := \frac{\pi}{2} \left( \frac{1 + k'^{3/2}}{2} \right)^{2/3}, \quad E_2 := \frac{\pi}{2} \left( \frac{1 + k'^{\delta}}{2} \right)^{1/\delta}, \quad \delta := \frac{\log(2)}{\log(\pi/2)}. \quad (30)$$

*Proof.*

(i)  $K_1 < K(k)$  is proved in [5, Theorem 18],  $K_2 < K(k)$  can be found in [10, equation (31)]. For  $K(k) < K_3$  and  $K(k) < K_4$  see Theorems 19 and 20 of [5],  $K(k) < K_5$  is proved in [4].

(ii) Both inequalities are proved by Alzer and Qiu in [5, Theorem 22].

□

**Remark.** Numerical computations show that

$$\max\{K_1, K_2\} = \begin{cases} K_1 & \text{for } 0 < k \leq 0.888\dots \\ K_2 & \text{for } 0.888\dots \leq k < 1 \end{cases} \quad (31)$$

and

$$\min\{K_3, K_4, K_5\} = \begin{cases} K_3 & \text{for } 0 < k \leq 0.971\dots \\ K_4 & \text{for } 0.971\dots \leq k \leq 0.990\dots \\ K_5 & \text{for } 0.990\dots \leq k < 1 \end{cases} \quad (32)$$

### 3 Auxiliary Results for Jacobi's Elliptic and Theta Functions

**Lemma 2.** *Let  $0 < k < 1$ . Then*

$$\begin{aligned} \Theta\left(\frac{K}{2}\right) &= \Theta_1\left(\frac{K}{2}\right) = \sqrt[4]{\frac{2}{\pi^2}(1+k')K^2\sqrt{k'}}, \\ H\left(\frac{K}{2}\right) &= H_1\left(\frac{K}{2}\right) = \sqrt[4]{\frac{2}{\pi^2}(1-k')K^2\sqrt{k'}}. \end{aligned} \quad (33)$$

*Proof.* For the four theta-functions, the following duplication formulas hold (see [13, p. 21] or formula (1051.25) of [9])

$$\begin{aligned} H(2u)\Theta(0)H_1(0)\Theta_1(0) &= 2H(u)\Theta(u)H_1(u)\Theta_1(u) \\ H_1(2u)H_1(0)\Theta^2(0) &= \Theta^2(u)H_1^2(u) - H^2(u)\Theta_1^2(u) \\ \Theta_1(2u)\Theta_1(0)\Theta^2(0) &= \Theta^2(u)\Theta_1^2(u) - H^2(u)H_1^2(u) \\ \Theta(2u)\Theta^3(0) &= \Theta_1^4(u) - H_1^4(u) = \Theta^4(u) - H^4(u) \end{aligned} \quad (34)$$

Further, by  $\Theta(-u) = \Theta(u)$  and  $\Theta_1(u+K) = \Theta(u)$ ,

$$\Theta\left(\frac{K}{2}\right) = \Theta\left(-\frac{K}{2}\right) = \Theta_1\left(-\frac{K}{2} + K\right) = \Theta_1\left(\frac{K}{2}\right),$$

and, by  $H(u) = -H(-u)$  and  $H_1(u+K) = -H(u)$ ,

$$H\left(\frac{K}{2}\right) = -H\left(-\frac{K}{2}\right) = H_1\left(-\frac{K}{2} + K\right) = H_1\left(\frac{K}{2}\right).$$

Hence, by (34), setting  $u = \frac{K}{2}$ ,

$$\begin{aligned}\Theta^4\left(\frac{K}{2}\right) - H^4\left(\frac{K}{2}\right) &= \Theta(K) \Theta^3(0) = \frac{4k'^{3/2} K^2}{\pi^2}, \\ 2\Theta^2\left(\frac{K}{2}\right) H^2\left(\frac{K}{2}\right) &= H(K) \Theta(0) H_1(0) \Theta_1(0) = \frac{4kK^2\sqrt{k'}}{\pi^2}.\end{aligned}$$

From these two equalities, the asserted formulas follow after some computation.  $\square$

**Lemma 3.** *Let  $0 < k < 1$ . Then the function  $\operatorname{dn}^2(u) \Theta^4(u)$  is strictly monotone increasing in  $u$ ,  $0 \leq u \leq \frac{K}{2}$ .*

*Proof.* Differentiating the above function gives

$$\frac{\partial}{\partial u} \{ \operatorname{dn}^2(u) \Theta^4(u) \} = 2 \operatorname{dn}^2(u) \Theta^4(u) \underbrace{\left( 2 \operatorname{zn}(u) - \frac{k^2 \operatorname{sn}(u) \operatorname{cn}(u)}{\operatorname{dn}(u)} \right)}_{=:g(u)}.$$

Note that  $g(0) = g\left(\frac{K}{2}\right) = 0$ . We prove that  $g(u) > 0$  for  $u \in (0, \frac{K}{2})$  by showing that  $g''(u) < 0$  for  $u \in (0, \frac{K}{2})$ . Indeed,

$$g''(u) = -\frac{2k^4 \operatorname{sn}(u) \operatorname{cn}(u)}{\operatorname{dn}^3(u)} (\operatorname{cn}(u) \operatorname{dn}(u) + k' \operatorname{sn}(u)) \underbrace{(\operatorname{cn}(u) \operatorname{dn}(u) - k' \operatorname{sn}(u))}_{=:h(u)}$$

and  $h(u)$  is strictly monotone decreasing for  $u \in (0, \frac{K}{2})$  and  $h\left(\frac{K}{2}\right) = 0$ , thus  $h(u) > 0$ , hence  $g''(u) < 0$  for  $u \in (0, \frac{K}{2})$ . This completes the proof.  $\square$

**Lemma 4.** *Let  $0 < k < 1$ , then, for  $0 < u < K$ ,  $\frac{\partial}{\partial k} \{ \operatorname{cn}(u, k) \} > 0$  and  $\frac{\partial}{\partial k} \{ \operatorname{dn}(u, k) \} < 0$ .*

*Proof.* By (710.52) and (710.53) of [9],

$$\begin{aligned}\frac{\partial}{\partial k} \{ \operatorname{cn}(u) \} &= \frac{\operatorname{sn}(u) \operatorname{dn}(u)}{kk'^2} \underbrace{\left( E(u) - k'^2 u - \frac{k^2 \operatorname{sn}(u) \operatorname{cn}(u)}{\operatorname{dn}(u)} \right)}_{=:f(u)} \\ \frac{\partial}{\partial k} \{ \operatorname{dn}(u) \} &= \frac{k \operatorname{sn}(u) \operatorname{cn}(u)}{k'^2} \underbrace{\left( E(u) - k'^2 u - \frac{\operatorname{sn}(u) \operatorname{dn}(u)}{\operatorname{cn}(u)} \right)}_{=:g(u)}\end{aligned}$$

Hence, it suffices to show that  $f(u) > 0$  and  $g(u) < 0$  for  $0 < u < K$ . Obviously,  $f(0) = g(0) = 0$ , and, using the formulas (121.00) of [9],

$$f'(u) = \frac{k^2 k'^2 \operatorname{sn}^2(u)}{\operatorname{dn}^2(u)}, \quad g'(u) = -\frac{k'^2}{\operatorname{cn}^2(u)},$$

thus  $f'(u) > 0$  and  $g'(u) < 0$  for  $0 < u < K$ , which gives  $f(u) > 0$  and  $g(u) < 0$ ,  $0 < u < K$ .  $\square$

**Lemma 5.** *Let  $0 < k < 1$ , then, for  $0 \leq u < K$ ,*

$$\frac{1}{\operatorname{dn}(u, k)} \leq \cosh(u) \leq \frac{1}{\operatorname{cn}(u, k)}, \quad (35)$$

and

$$\log \frac{1 + k \operatorname{sn}(u)}{\operatorname{dn}(u)} \leq u \leq \log \frac{1 + \operatorname{sn}(u)}{\operatorname{cn}(u)}. \quad (36)$$



*Proof.* By Lemma 4 and the fact that

$$\lim_{k \rightarrow 1} \operatorname{cn}(u, k) = \lim_{k \rightarrow 1} \operatorname{dn}(u, k) = \frac{1}{\cosh(u)}$$

we get  $\operatorname{cn}(u, k) \leq 1/\cosh(u)$  and  $\operatorname{dn}(u, k) \geq 1/\cosh(u)$  for every  $u \in (0, K)$ . The second inequality follows immediately by the well known formula  $\cosh^{-1}(u) = \log(u + \sqrt{u^2 - 1})$ .  $\square$

**Lemma 6.** *Let  $0 < k < 1$ , then, for  $0 \leq u \leq K$ ,*

$$\operatorname{zn}(u) \leq (E - k'^2 K)(1 - u/K). \quad (37)$$

*Proof.* Define

$$f(u) := (E - k'^2 K)(1 - u/K) - \operatorname{zn}(u),$$

then  $f(K) = 0$  and  $f'(u) = k'^2 - \operatorname{dn}^2(u) < 0$  for  $0 \leq u \leq K$ , thus  $f(u) \geq 0$  for  $0 \leq u \leq K$ .  $\square$

**Lemma 7.** *Let  $0 < k < 1$ , then, for  $0 \leq \lambda \leq 1$ ,*

$$\frac{\Theta(0)}{\Theta(\lambda K)} \leq \sqrt{k'} \exp\left(\frac{1}{2}(E/K - k'^2)(1 - \lambda)^2 K^2\right). \quad (38)$$

*Proof.* For  $0 \leq u \leq K$ , we have

$$\log \frac{\Theta(K)}{\Theta(u)} = -\log \frac{\Theta(u)}{\Theta(K)} = -\int_K^u \operatorname{zn}(v) \, dv = \int_u^K \operatorname{zn}(v) \, dv.$$

Thus, by Lemma 6,

$$\log \frac{\Theta(K)}{\Theta(u)} \leq \int_u^K (E - k'^2 K)(1 - v/K) \, dv = (E - k'^2 K)(K/2 - u + u^2/(2K))$$

Taking exp on both sides and using the relation  $\Theta(0) = \sqrt{k'} \Theta(K)$  gives (38).  $\square$

## References

- [1] N.I. Achieser, *Über einige Funktionen, die in gegebenen Intervallen am wenigsten von Null abweichen*, Bull. Phys. Math. **3** (1929), 1–69 (in German).
- [2] ———, *Sur les polynomes de Tschebyscheff pour deux segments*, C. R. Math. Acad. Sci. Paris **191** (1930), 754–756 (in French).
- [3] ———, *Über einige Funktionen, welche in zwei gegebenen Intervallen am wenigsten von Null abweichen*, Bull. Acad. Sci. URSS **7** (1932), 1163–1202 (in German).
- [4] H. Alzer, *Sharp inequalities for the complete elliptic integral of the first kind*, Math. Proc. Cambridge Philos. Soc. **124** (1998), 309–314.
- [5] H. Alzer and S.-L. Qiu, *Monotonicity theorems and inequalities for the complete elliptic integrals*, J. Comput. Appl. Math. **172** (2004), 289–312.

- [6] G.D. Anderson, M.K. Vamanamurthy, and M. Vuorinen, *Functional inequalities for complete elliptic integrals and their ratios*, SIAM J. Math. Anal. **21** (1990), 536–549.
- [7] ———, *Functional inequalities for hypergeometric functions and complete elliptic integrals*, SIAM J. Math. Anal. **23** (1992), 512–524.
- [8] ———, *Conformal invariants, inequalities, and quasiconformal maps*, Wiley, 1997.
- [9] P.F. Byrd and M.D. Friedman, *Handbook of elliptic integrals for engineers and scientists*, Springer, 1971.
- [10] B.C. Carlson and J.L. Gustafson, *Asymptotic approximations for symmetric elliptic integrals*, SIAM J. Math. Anal. **25** (1994), 288–303.
- [11] B.C. Carlson and J. Todd, *The degenerating behavior of elliptic functions*, SIAM J. Numer. Anal. **20** (1983), 1120–1129.
- [12] J. Gillis, *Tchebycheff polynomials and the transfinite diameter*, Amer. J. Math. **63** (1941), 283–290.
- [13] D.F. Lawden, *Elliptic functions and applications*, Springer, 1989.
- [14] W. Magnus, F. Oberhettinger, and R.P. Soni, *Formulas and theorems for the special functions of mathematical physics*, Springer, 1966.
- [15] H.N. Mhaskar and E.B. Saff, *Weighted analogues of capacity, transfinite diameter, and Chebyshev constant*, Constr. Approx. **8** (1992), 105–124.
- [16] F. Peherstorfer, *Elliptic orthogonal and extremal polynomials*, Proc. London Math. Soc. (3) **70** (1995), 605–624.
- [17] F. Peherstorfer and K. Schiefermayr, *Description of inverse polynomial images which consist of two Jordan arcs with the help of Jacobi's elliptic functions*, Comput. Methods Funct. Theory **4** (2004), 355–390.
- [18] C. Pommerenke, *Über die Kapazität der Summe von Kontinuen*, Math. Ann. **139** (1959), 127–132.
- [19] S.-L. Qiu and M.K. Vamanamurthy, *Sharp estimates for complete elliptic integrals*, SIAM J. Math. Anal. **27** (1996), 823–834.
- [20] R.M. Robinson, *On the transfinite diameters of some related sets*, Math. Z. **108** (1969), 377–380.
- [21] A.Yu. Solynin, *Extremal configurations in some problems on capacity and harmonic measure*, J. Math. Sci. **89** (1998), 1031–1049.
- [22] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, Cambridge University Press, 1962.
- [23] E.I. Zolotarev, *Sur la méthode d'intégration de M. Tchebichef*, J. Math. Pures Appl. **19** (1874), 161–188.

- [24] ———, *Applications of elliptic functions to problems of functions deviating least and most from zero (in Russian)*, Oeuvres de E.I. Zolotarev, Vol. 2, Izdat. Akad. Nauk SSSR, Leningrad (1932), 1–59.  
Available on <http://www.math.technion.ac.il/hat/papers.html>.