

Some New Properties of Jacobi's Theta Functions*

Klaus Schiefermayr[†]

Abstract

In this paper, a monotonicity property for the quotient of two Jacobi's theta functions with respect to the modulus k is proved.

Mathematics Subject Classification (2000): 33E05

Keywords: Jacobi's elliptic functions, Jacobi's theta functions

1 Introduction and Main Result

Let $k \in (0, 1)$ denote the modulus of Jacobi's elliptic functions $\operatorname{sn}(u)$, $\operatorname{cn}(u)$, and $\operatorname{dn}(u)$, of Jacobi's theta functions $\Theta(u)$, $H(u)$, $H_1(u)$, and $\Theta_1(u)$, and, finally, of Jacobi's zeta function, $\operatorname{zn}(u)$. Here we follow the notation of Carlson and Todd [3], in other references, like [8], Jacobi's zeta function is denoted by $Z(u)$. Let $k' := \sqrt{1 - k^2} \in (0, 1)$ be the complementary modulus, let $K \equiv K(k)$ and $E \equiv E(k)$ be the complete elliptic integral of the first and second kind, respectively, and let $K' \equiv K'(k) := K(k')$ and $E' \equiv E'(k) := E(k')$.

If we want to point out the dependence of these functions on the modulus k , we will write $K(k)$, $K'(k)$, $\operatorname{sn}(u, k)$, $\Theta(u, k)$, etc. In this paper, we follow the old notation of the theta functions, which goes back to Jacobi. There is another notation of the four theta functions given by $\Theta(u, k) = \vartheta_0(v, \tau) = \vartheta_4(v, \tau)$, $H(u, k) = \vartheta_1(v, \tau)$, $H_1(u, k) = \vartheta_2(v, \tau)$ and $\Theta_1(u, k) = \vartheta_3(v, \tau)$, where $\tau = iK'/K$ and $v = u/(2K)$ (note that in some references, like [2] and [4], $v = u\pi/(2K)$).

For the definitions and many important properties of these functions, see, e.g., [2], [4], [5], [7], and [1].

In [3], Carlson and Todd proved that, for each $\lambda \in (0, 1)$, the functions $\operatorname{sn}(\lambda K, k)$ and $\operatorname{zn}(\lambda K, k)$ are strictly monotone increasing with respect to the modulus k , $0 < k < 1$. In addition, they investigated the degenerating behaviour of these functions as $k \rightarrow 0$ and especially as $k \rightarrow 1$. Hence the question arises, whether analogous monotonicity properties hold for the theta functions. Unfortunately, $\Theta(\lambda K, k)$ does not have the same monotonicity behaviour with respect to the modulus k for every $\lambda \in (0, 1)$. Numerical examples show that for small λ ($\lambda \leq 0.5$), $\Theta(\lambda K)$ is strictly monotone decreasing, for large λ ($\lambda \geq 0.6$), $\Theta(\lambda K)$ is strictly monotone increasing in k and for some $\lambda \in (0.5, 0.6)$, $\Theta(\lambda K)$ is not monotone at all in the whole interval $(0, 1)$. However, we are able to prove that for each $\lambda, \mu \in \mathbb{R}$, the quotient $\Theta(\lambda K)/\Theta(\mu K)$ of two theta functions is strictly monotone. In addition, the degenerative behaviour of $\Theta(\lambda K)/\Theta(\mu K)$ as $k \rightarrow 0$ and $k \rightarrow 1$ is given.

*published in: Journal of Computational and Applied Mathematics **178** (2005), 419–424.

[†]University of Applied Sciences Upper Austria, School of Engineering and Environmental Sciences, Stelzhamerstrasse 23, 4600 Wels, Austria, KLAUS.SCHIEFERMAYR@FH-WELS.AT

Theorem 1.

- (i) Let $\lambda, \mu \in \mathbb{R}$. If $\cos(\lambda\pi) > \cos(\mu\pi)$ [$\cos(\lambda\pi) < \cos(\mu\pi)$], then $\Theta(\lambda K)/\Theta(\mu K)$ is a positive, strictly monotone decreasing [increasing] function of the modulus $k \in (0, 1)$. If $\cos(\lambda\pi) = \cos(\mu\pi)$, i.e., $\lambda = \mu + 2\nu$, $\nu \in \mathbb{Z}$, then $\Theta(\lambda K)/\Theta(\mu K) = 1$.
- (ii) Let $\lambda, \mu \in (0, 1)$, $k \in (0, 1)$, then $\Theta(\lambda K)/\Theta(\mu K) \rightarrow 1$ as $k \rightarrow 0$ and

$$\frac{\Theta(\lambda K)}{\Theta(\mu K)} \sim \left(\frac{k'}{4}\right)^{(\mu-\lambda)(1-(\lambda+\mu)/2)} \quad \text{as } k \rightarrow 1.$$

- (iii) Let $\mu \in (0, 1)$ and $k \in (0, 1)$. Then $f(\lambda) := \Theta((\mu - \lambda)K)/\Theta((\mu + \lambda)K)$ is a convex function of $\lambda \in (0, 1)$ with $f(0) = f(1) = 1$.

Remark. By the relation $\Theta_1(u) = \Theta(u + K)$, one gets analogous monotonicity properties for the quotients $\Theta_1(\lambda K)/\Theta_1(\mu K)$, $\Theta(\lambda K)/\Theta_1(\mu K)$, and $\Theta_1(\lambda K)/\Theta(\mu K)$.

In [6], we considered polynomials, whose $[-1, 1]$ inverse image consists of two Jordan arcs, i.e., we characterized polynomials P_n , for which $P_n^{-1}([-1, 1])$ consists of two Jordan arcs (in general, $P_n^{-1}([-1, 1])$ consists of n Jordan arcs). Since these polynomials P_n can be given with the help of an elliptic integral, Jacobi's elliptic and theta functions appear in a natural way. When describing the shape of the two Jordan arcs, we need the above theorem, see Theorem 22, Lemma 33 and Lemma 34 of [6].

2 Proof of the Main Result

First, we collect some derivation formulas, which are an immediate consequence of formula (710.00) of [2].

Lemma 2. Let $k \in (0, 1)$. Then

$$\begin{aligned} \frac{dK}{dk} &= \frac{E - k'^2 K}{kk'^2}, & \frac{dK'}{dk} &= \frac{k^2 K' - E'}{kk'^2}, \\ \frac{d}{dk} \left\{ \frac{1}{K} \right\} &= \frac{k'^2 K - E}{kk'^2 K^2}, & \frac{d}{dk} \left\{ \frac{K'}{K} \right\} &= \frac{-\pi}{2kk'^2 K^2} \end{aligned}$$

Concerning the limiting behaviour of $\text{sn}(\lambda K)$, etc., as $k \rightarrow 1$, Carlson and Todd [3] have proved the following.

Lemma 3. Let $0 < \lambda < 1$, then, as $k \rightarrow 1$, we have

$$\begin{aligned} K &\sim \log\left(\frac{4}{k'}\right), & \text{sn}(\lambda K) &\sim 1 - 2\left(\frac{k'}{4}\right)^{2\lambda}, & \text{cn}(\lambda K) &\sim \text{dn}(\lambda K) \sim 2\left(\frac{k'}{4}\right)^\lambda, \\ & & \text{zn}(\lambda K) &\sim 1 - \lambda - 2\left(\frac{k'}{4}\right)^{2\lambda}. \end{aligned}$$

Moreover, we will need the following formulas for the derivatives of the theta functions, which are a direct consequence of (1053.01), (1052.02) and (731.01)–(731.03) of [2].

Lemma 4. *The following relations hold:*

$$\begin{aligned}\frac{\partial}{\partial u}\{\Theta(u)\} &= \Theta(u)\text{zn}(u), \\ \frac{\partial}{\partial u}\{H(u)\} &= \sqrt{k}\Theta(u)(\text{cn}(u)\text{dn}(u) + \text{sn}(u)\text{zn}(u)), \\ \frac{\partial}{\partial u}\{H_1(u)\} &= \frac{\sqrt{k}}{\sqrt{k'}}\Theta(u)(-\text{sn}(u)\text{dn}(u) + \text{cn}(u)\text{zn}(u)), \\ \frac{\partial}{\partial u}\{\Theta_1(u)\} &= \frac{1}{\sqrt{k'}}\Theta(u)(-k^2\text{sn}(u)\text{cn}(u) + \text{dn}(u)\text{zn}(u)),\end{aligned}$$

and

$$\begin{aligned}\frac{\partial^2}{\partial u^2}\{\Theta(u)\} &= \Theta(u)(\text{dn}^2(u) + \text{zn}^2(u) - E/K), \\ \frac{\partial^2}{\partial u^2}\{H(u)\} &= \sqrt{k}\Theta(u)\left(-k^2\text{sn}(u)\text{cn}^2(u) + 2\text{cn}(u)\text{dn}(u)\text{zn}(u)\right. \\ &\quad \left.+ \text{sn}(u)(\text{zn}^2(u) - E/K)\right), \\ \frac{\partial^2}{\partial u^2}\{H_1(u)\} &= \frac{\sqrt{k}}{\sqrt{k'}}\Theta(u)\left(-k^2\text{sn}^2(u)\text{cn}(u) - 2\text{sn}(u)\text{dn}(u)\text{zn}(u)\right. \\ &\quad \left.+ \text{cn}(u)(\text{zn}^2(u) - E/K)\right), \\ \frac{\partial^2}{\partial u^2}\{\Theta_1(u)\} &= \frac{1}{\sqrt{k'}}\Theta(u)\left(\text{dn}(u)(1 - k^2\text{cn}^2(u)) - 2k^2\text{sn}(u)\text{cn}(u)\text{zn}(u)\right. \\ &\quad \left.+ \text{dn}(u)(\text{zn}^2(u) - E/K)\right).\end{aligned}$$

Remark. By (123.01), (123.03) of [2] and Lemma 4, the relation

$$\frac{\partial}{\partial u}\left\{\log\left(\frac{H(v-u)}{H(v+u)}\right)\right\} = \frac{2\text{sn}(v)\text{cn}(v)\text{dn}(v)}{\text{sn}^2(u) - \text{sn}^2(v)} - 2\text{zn}(v)$$

holds which implies

$$\log\left(\frac{H(v-u)}{H(v+u)}\right) = \int_0^u \frac{2\text{sn}(v)\text{cn}(v)\text{dn}(v)}{\text{sn}^2(u) - \text{sn}^2(v)} du - 2u\text{zn}(v). \quad (1)$$

Analogous formulas can be obtained for other ratios of theta functions.

The next lemma gives the derivatives with respect to the modulus k for the four theta functions $\Theta(u)$, $H(u)$, $H_1(u)$, $\Theta_1(u)$, where $u = \lambda K$. Note that λK as well as the theta functions themselves depend on the modulus k .

Lemma 5. *For $\lambda \in \mathbb{R}$, the derivatives with respect to k of the four theta functions $\Theta(\lambda K)$, $H(\lambda K)$, $H_1(\lambda K)$, $\Theta_1(\lambda K)$ are given by*

$$\begin{aligned}\frac{d}{dk}\{\Theta(\lambda K)\} &= -\frac{1}{2kk'^2}\frac{\partial^2}{\partial u^2}\{\Theta(u)\}\Big|_{u=\lambda K} \\ &= -\frac{1}{2kk'^2}\Theta(\lambda K)(\text{dn}^2(\lambda K) + \text{zn}^2(\lambda K) - E/K), \\ \frac{d}{dk}\{H(\lambda K)\} &= -\frac{1}{2kk'^2}\frac{\partial^2}{\partial u^2}\{H(u)\}\Big|_{u=\lambda K} \\ \frac{d}{dk}\{H_1(\lambda K)\} &= -\frac{1}{2kk'^2}\frac{\partial^2}{\partial u^2}\{H_1(u)\}\Big|_{u=\lambda K} \\ \frac{d}{dk}\{\Theta_1(\lambda K)\} &= -\frac{1}{2kk'^2}\frac{\partial^2}{\partial u^2}\{\Theta_1(u)\}\Big|_{u=\lambda K}\end{aligned}$$

Proof. The four theta functions $\vartheta_j(v, \tau)$, $j = 1, 2, 3, 4$, satisfy a differential equation of the form

$$\frac{\partial^2}{\partial v^2} \{\vartheta_j(v, \tau)\} = 4i\pi \frac{\partial}{\partial \tau} \{\vartheta_j(v, \tau)\},$$

see [5, p. 375]. By Lemma 2 and since $\Theta(u, k) = \vartheta_4(\frac{u}{2K}, \tau)$, where $\tau = iK'/K$,

$$\begin{aligned} \frac{\partial}{\partial k} \{\Theta(u, k)\} &= \frac{\partial}{\partial v} \{\vartheta_4(\frac{u}{2K}, \tau)\} \frac{d}{dk} \{\frac{u}{2K}\} + \frac{\partial}{\partial \tau} \{\vartheta_4(\frac{u}{2K}, \tau)\} \frac{d\tau}{dk} \\ &= \frac{u(k'^2K - E)}{kk'^2K} \frac{\partial}{\partial u} \{\Theta(u, k)\} - \frac{1}{2kk'^2} \frac{\partial^2}{\partial u^2} \{\Theta(u, k)\}. \end{aligned}$$

Thus, for the derivative with respect to k , by Lemma 2, we get

$$\begin{aligned} \frac{d}{dk} \{\Theta(\lambda K, k)\} &= \frac{d}{dk} \{\lambda K\} \frac{\partial}{\partial u} \{\Theta(u, k)\} \Big|_{u=\lambda K} + \frac{\partial}{\partial k} \{\Theta(u, k)\} \Big|_{u=\lambda K} \\ &= \frac{\lambda(E - k'^2K)}{kk'^2} \frac{\partial}{\partial u} \{\Theta(u, k)\} \Big|_{u=\lambda K} + \frac{\lambda K(k'^2K - E)}{kk'^2K} \frac{\partial}{\partial u} \{\Theta(u, k)\} \Big|_{u=\lambda K} \\ &\quad - \frac{1}{2kk'^2} \frac{\partial^2}{\partial u^2} \{\Theta(u, k)\} \Big|_{u=\lambda K} \\ &= -\frac{1}{2kk'^2} \frac{\partial^2}{\partial u^2} \{\Theta(u, k)\} \Big|_{u=\lambda K}. \end{aligned}$$

Since the last identity holds for the other three theta functions as well, this gives the assertion. \square

Proof of Theorem 1.

(i) By Lemma 5,

$$\begin{aligned} \frac{d}{dk} \left\{ \frac{\Theta(\lambda K)}{\Theta(\mu K)} \right\} &= -\frac{1}{2kk'^2} \frac{\Theta(\lambda K)}{\Theta(\mu K)} \\ &\quad \times (\operatorname{dn}^2(\lambda K) + \operatorname{zn}^2(\lambda K) - \operatorname{dn}^2(\mu K) - \operatorname{zn}^2(\mu K)). \end{aligned}$$

Thus, it remains to be shown that $g(u) := \operatorname{dn}^2(u) + \operatorname{zn}^2(u)$ satisfies the inequality $g(\lambda K) - g(\mu K) > 0$ [< 0] for all $\lambda, \mu \in \mathbb{R}$ with $\cos(\lambda\pi) > [<] \cos(\mu\pi)$. This property holds since $g(-u) = g(u)$, $g(u + 2K) = g(u)$ and $g(u)$ is a positive, strictly monotone decreasing function in $(0, K)$. Concerning the monotonicity of $g(u)$, note that

$$g'(u) = -2\operatorname{dn}^2(u) \underbrace{\left(\frac{k^2 \operatorname{sn}(u) \operatorname{cn}(u)}{\operatorname{dn}(u)} - \operatorname{zn}(u) \right)}_{=: h(u)} - \frac{2E \operatorname{zn}(u)}{K},$$

where $h(0) = h(K) = 0$ and

$$h''(u) = -\frac{2k^2k'^2 \operatorname{sn}(u) \operatorname{cn}(u)}{\operatorname{dn}^3(u)} < 0 \quad \text{for } u \in (0, K).$$

Thus $h(u) > 0$ and therefore $g'(u) < 0$ for $u \in (0, K)$.

- (ii) Assume that $\lambda < \mu$ and let $\alpha := (\lambda + \mu)/2$, $\beta := (\mu - \lambda)/2$, i.e. $\lambda = \alpha - \beta$, $\mu = \alpha + \beta$, $\alpha, \beta \in (0, 1)$. By (1052.02) of [2] and (1), we get

$$\begin{aligned} \log\left(\frac{\Theta(\lambda K)}{\Theta(\mu K)}\right) &= \log\left(\frac{\operatorname{sn}(\mu K)H(\lambda K)}{\operatorname{sn}(\lambda K)H(\mu K)}\right) = \log(\operatorname{sn}(\mu K)) - \log(\operatorname{sn}(\lambda K)) \\ &+ 2K\operatorname{sn}(\alpha K)\operatorname{cn}(\alpha K)\operatorname{dn}(\alpha K) \int_0^\beta \frac{d\nu}{\operatorname{sn}^2(\nu K) - \operatorname{sn}^2(\alpha K)} - 2\beta K\operatorname{zn}(\alpha K). \end{aligned}$$

Thus, as $k \rightarrow 0$,

$$\begin{aligned} \log\left(\frac{\Theta(\lambda K)}{\Theta(\mu K)}\right) &\sim \log(\sin(\frac{\mu\pi}{2})) - \log(\sin(\frac{\lambda\pi}{2})) \\ &+ \pi \sin(\frac{\alpha\pi}{2}) \cos(\frac{\alpha\pi}{2}) \int_0^\beta \frac{d\nu}{\sin^2(\frac{\nu\pi}{2}) - \sin^2(\frac{\alpha\pi}{2})} = 0, \end{aligned}$$

and, as $k \rightarrow 1$, by Lemma 3, $a := k'/4$,

$$\begin{aligned} \log\left(\frac{\Theta(\lambda K)}{\Theta(\mu K)}\right) &\sim \log(1 - 2a^{2\mu}) - \log(1 - 2a^{2\lambda}) \\ &- 2\log(a)(1 - 2a^{2\alpha})4a^{2\alpha} \int_0^\beta \frac{d\nu}{4(a^{2\alpha} - a^{2\nu})} + 2\beta(1 - \alpha - 2a^{2\alpha})\log(a) \\ &\sim 2\log(a) \int_0^\beta \frac{d\nu}{a^{2(\nu-\alpha)} - 1} + 2\beta(1 - \alpha)\log(a) \\ &= -2\beta\log(a) + \log(a^{2(\beta-\alpha)} - 1) - \log(a^{-2\alpha} - 1) + 2\beta(1 - \alpha)\log(a) \\ &\sim -2\beta\log(a) + 2(\beta - \alpha)\log(a) + 2\alpha\log(a) + 2\beta(1 - \alpha)\log(a) \\ &= 2\beta(1 - \alpha)\log(a). \end{aligned}$$

If $\lambda > \mu$, note that by the above proved result

$$\begin{aligned} \log\frac{\Theta(\lambda K)}{\Theta(\mu K)} &= -\log\frac{\Theta(\mu K)}{\Theta(\lambda K)} \sim -(\lambda - \mu)(1 - (\lambda + \mu)/2)\log a \\ &= (\mu - \lambda)(1 - (\lambda + \mu)/2)\log a. \end{aligned}$$

- (iii) Obviously, $f(0) = 1$ and, by (1051.02) and (1051.03) of [2],

$$f(1) = \frac{\Theta(\mu K - K)}{\Theta(\mu K + K)} = \frac{\Theta_1(-\mu K)}{\Theta_1(\mu K)} = 1.$$

Further, by Lemma 4,

$$\begin{aligned} f''(\lambda) &= K^2 f(\lambda) \left((\operatorname{zn}((\mu - \lambda)K) + \operatorname{zn}((\mu + \lambda)K))^2 \right. \\ &\quad \left. + \operatorname{dn}^2((\mu - \lambda)K) - \operatorname{dn}^2((\mu + \lambda)K) + \frac{2E}{K^2} \right) > 0 \end{aligned}$$

for every $\lambda \in (0, 1)$, which gives the assertion. \square

References

- [1] N.I. Achieser, *Elements of the theory of elliptic functions*, Translations of Mathematical Monographs, vol. 79, American Mathematical Society, 1990.
- [2] P.F. Byrd and M.D. Friedman, *Handbook of elliptic integrals for engineers and scientists*, Springer, 1971.
- [3] B.C. Carlson and J. Todd, *The degenerating behavior of elliptic functions*, SIAM J. Numer. Anal. **20** (1983), 1120–1129.
- [4] D.F. Lawden, *Elliptic functions and applications*, Springer, 1989.
- [5] W. Magnus, F. Oberhettinger, and R.P. Soni, *Formulas and theorems for the special functions of mathematical physics*, Springer, 1966.
- [6] F. Peherstorfer and K. Schiefermayr, *Description of inverse polynomial images which consist of two Jordan arcs with the help of Jacobi's elliptic functions*, Comput. Methods Funct. Theory **4** (2004), 355–390.
- [7] J. Tannery and J. Molk, *Éléments de la théorie des fonctions elliptiques. Tomes I, II. Calcul différentiel*, Chelsea Publishing, 1972.
- [8] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, Cambridge University Press, 1962.