

ON INVOLUTIVE CLUSTER AUTOMORPHISMS

NDOUNÉ NDOUNÉ

ABSTRACT. We construct a special embedding of the translation quiver $\mathbb{Z}Q$ in the three-dimensional affine space \mathbb{R}^3 where Q is a finite connected acyclic quiver and $\mathbb{Z}Q$ contains a local slice whose quiver is isomorphic to the opposite quiver Q^{op} of Q . Via this embedding, we show that there exists an involutive anti-automorphism of the translation quiver $\mathbb{Z}Q$. As an immediate consequence, we characterize explicitly the group of cluster automorphisms of the cluster algebras of seed (X, Q) , where Q and Q^{op} are mutation equivalent.

1. INTRODUCTION

Cluster algebras were invented by Fomin and Zelevinsky [11] in order to understand the canonical bases of quantized enveloping algebras associated to semi-simple Lie algebras [18]. A cluster algebra is a commutative ring with a distinguished set of generators, called *cluster variables*. The set of all cluster variables is constructed recursively from an initial set using a procedure called *mutation*.

Today the theory of cluster algebras relates with numerous fields of mathematics, for example, representation theory of algebras, Lie theory, Poisson geometry, Teichmüller theory and integrable systems. Following [3, 2], we are interested in studying the involutive cluster automorphisms. The objective of this paper is to solve an open problem posed in [3]; it was conjectured that if Q is a finite connected acyclic quiver which is mutation equivalent to Q^{op} , then the group of cluster automorphisms of the cluster algebra of a seed (X, Q) is a semi-direct product of the subgroup of the direct cluster automorphisms with \mathbb{Z}_2 . A partial answer was given in [3], in the case when the underlying graph of Q is a tree. We prove it here in general.

In this paper, we construct an embedding of a repetitive translation quiver $\mathbb{Z}Q$ in 3-space \mathbb{R}^3 , when Q is a finite connected acyclic quiver and the translation quiver contains a local slice whose quiver is isomorphic to Q^{op} . In particular, we construct an embedding of the transjective component of the Auslander Reiten quiver of the cluster category \mathcal{C}_Q . We then use this construction to prove the existence of an involutive inverse cluster automorphism of a cluster algebra with a seed (X, Q) , where Q is a finite connected acyclic quiver which is mutation equivalent to Q^{op} .

Our paper is organized as follows. After an introductory section 2, we construct in section 3, the required embedding η of $\mathbb{Z}Q$ in \mathbb{R}^3 , such that if Σ and $\tilde{\Sigma}$ are two local slices whose quivers are respectively isomorphic to Q and Q^{op} , then $\eta(\Sigma)$ and $\eta(\tilde{\Sigma})$ are symmetric with respect to a particular plane.

In section 4, we prove that, if Q is a finite acyclic quiver which is mutation equivalent to Q^{op} , then the group of cluster automorphisms of a cluster algebra of a seed (X, Q) is a semi-direct product of the subgroup of the direct cluster automorphisms with \mathbb{Z}_2 .

ACKNOWLEDGMENTS

This article is part of my PhD thesis, under the supervision of Ibrahim Assem and Vasilisa Shramchenko. I would like to thank them deeply for their patience and availability.

2. CLUSTER AUTOMORPHISMS

The notion of cluster automorphisms was introduced in [3] and it is connected with quiver automorphisms. We recall that a *quiver* is a quadruple $Q = (Q_0, Q_1, s, t)$ consisting of two sets, Q_0 (whose elements are called *vertices*) and Q_1 (whose elements are called *arrows*), and of two maps $s, t : Q_1 \rightarrow Q_0$ associating to each arrow $\alpha \in Q_1$ its *source* $s(\alpha)$ and its *target* $t(\alpha)$. If $i = s(\alpha)$ and $j = t(\alpha)$, we denote this situation by $i \xrightarrow{\alpha} j$. Given a vertex i , we set $i^+ = \{\alpha \in Q_1 | s(\alpha) = i\}$ and $i^- = \{\alpha \in Q_1 | t(\alpha) = i\}$.

Let Q be a finite connected quiver without oriented cycles of length one or two. Let $n = |Q_0|$ denote the number of vertices in Q , $X = \{x_1, x_2, \dots, x_n\}$ be a set of n variables and denote the vertices by $Q_0 = \{1, 2, \dots, n\}$, where we agree that the point i corresponds to the variable x_i . We consider the field $\mathcal{F} = \mathbb{Q}(x_1, x_2, \dots, x_n)$ of rational functions in x_1, x_2, \dots, x_n (with rational coefficients). The *cluster algebra* $\mathcal{A} = \mathcal{A}(X, Q)$ is the \mathbb{Z} -subalgebra of \mathcal{F} defined by a set of generators obtained recursively from X in the following manner. Let k be such that $1 \leq k \leq n$. We define the mutation $\mu_{x_k, X}$ (or μ_{x_k} or μ_k for brevity if there is no ambiguity) of the pair (X, Q) to be a new pair (X', Q') , that is $\mu_k(X, Q) = (X', Q')$, where Q' is the quiver obtained from Q by performing the following operations:

- (a) for any path $i \rightarrow k \rightarrow j$ of length two having k as mid-vertex, we insert a new arrow $i \rightarrow j$.
- (b) all arrows incident to the vertex k are reversed,
- (c) all cycles of length two are deleted.

On the other hand, X' is a set of variables defined as follows. $X' = (X \setminus \{x_k\}) \cup \{x'_k\}$ where $x'_k \in \mathcal{F}$ is obtained from X by the

so-called *exchange relation*:

$$x_k x'_k = \prod_{\alpha \in i^+} x_{t(\alpha)} + \prod_{\alpha \in i^-} x_{s(\alpha)}.$$

Let \mathcal{X} be the union of all possible sets of variables obtained from X by successive mutations. Then the *cluster algebra* $\mathcal{A} = \mathcal{A}(X, Q)$ is the \mathbb{Z} -subalgebra of \mathcal{F} generated by \mathcal{X} .

Each pair (\tilde{X}, \tilde{Q}) obtained from (X, Q) by successive mutations is called a *seed*, and the set \tilde{X} is called a *cluster*. The elements $\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n$ of a cluster \tilde{X} are called *cluster variables*. Each cluster is a transcendence basis for the field \mathcal{F} . The pair (X, Q) is called the *initial seed*, and X is called the *initial cluster*.

Gekhtman, Shapiro and Vainshtein showed in [13] that for every seed (\tilde{X}, \tilde{Q}) the quiver \tilde{Q} is uniquely defined by the cluster \tilde{X} , and we use the notation $Q(\tilde{X})$ for the quiver of a given cluster \tilde{X} . We write p_x for the point of $Q(\tilde{X})$ corresponding to the cluster variable $x \in \tilde{X}$. Following [11] we recall that the Laurent phenomenon asserts that each cluster variable in \mathcal{A} can be expressed as a Laurent polynomial in x_i , with $1 \leq i \leq n$, that is, every cluster variable is of the form

$$\frac{p(x_1, x_2, \dots, x_n)}{\prod_{l=1}^n x_l^{d_l}},$$

where $p \in \mathbb{Z}[x_1, x_2, \dots, x_n]$, and d_l is a positive integer (for $1 \leq l \leq n$).

As a consequence, $\mathcal{A} \subseteq \mathbb{Z}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$, see [11] for more details. The following definition is due to [3].

Definition 2.1. *Let \mathcal{A} be a cluster algebra, and let $f : \mathcal{A} \rightarrow \mathcal{A}$ be an automorphism of \mathbb{Z} -algebras. Then f is called *cluster automorphism* if there exists a seed (X, Q) of \mathcal{A} such that the following conditions are satisfied:*

- (CA1) $f(X)$ is a cluster;
- (CA2) $f(X)$ is compatible with mutations, that is, for every $x \in X$, we have $f(\mu_{x, X}(x)) = \mu_{f(x), f(X)}(f(x))$.

Now we state an equivalent characterization of cluster automorphisms see [3, Lemma 2.3].

Lemma 2.1. *Let f be a \mathbb{Z} -algebra automorphism of \mathcal{A} . Then f is a cluster automorphism if and only if there exists a seed (X, Q) such that $f(X)$ is a cluster and one of the following two conditions is satisfied:*

- (a) *there exists an isomorphism of quivers $\varphi : Q \rightarrow Q(f(X))$ such that $\varphi(p_x) = p_{f(x)}$ for all $p_x \in Q_0$, or*
- (b) *there exists an isomorphism of quivers $\varphi : Q^{op} \rightarrow Q(f(X))$ such that $\varphi(p_x) = p_{f(x)}$ for all $p_x \in Q_0$.*

A cluster automorphism f is called *direct* if it satisfies condition (a) of the above lemma, and f is called *inverse* if it satisfies condition (b).

The set of all cluster automorphisms forms a group under composition denoted by $\text{Aut}(\mathcal{A})$ and called the group of *cluster automorphisms*. The set of all direct automorphisms is denoted by $\text{Aut}^+(\mathcal{A})$; it is a normal subgroup of $\text{Aut}(\mathcal{A})$ of index at most two. For more results and properties of cluster automorphisms, we refer the reader to [3].

3. INVOLUTIVE AUTOMORPHISMS OF QUIVERS

In this section, we assume that Q is a finite connected acyclic quiver. Let K be an algebraically closed field, KQ the path algebra of Q and $\text{mod}KQ$ the category of finitely generated right KQ -modules. For the properties of $\text{mod}KQ$ and its Auslander-Reiten quiver $\Gamma(\text{mod}KQ)$ we refer the reader to [5, 4]. We denote by $\mathcal{D}^d(\text{mod}KQ)$ the bounded derived category over $\text{mod}KQ$. This is a triangulated Krull-Schmidt category having a Serre duality and hence almost split triangles. Since KQ is hereditary, the Auslander Reiten quiver $\Gamma(\mathcal{D}^d(\text{mod}KQ))$ of $\mathcal{D}^d(\text{mod}KQ)$ is well-understood, see [14].

Following [4] we give the definitions and some basic properties of a well-known translation quiver. If Q is a connected acyclic quiver, the *repetitive quiver* $\mathbb{Z}Q$ is the quiver whose set of vertices is $(\mathbb{Z}Q)_0 = \{(m, \nu) | m \in \mathbb{Z}, \nu \in Q_0\}$ and the set of arrows is $(\mathbb{Z}Q)_1 = \{(m, \alpha) : (m, \nu) \rightarrow (m, \nu') | m \in \mathbb{Z}, \alpha : \nu \rightarrow \nu'\} \cup \{(m, \alpha') : (m, \nu') \rightarrow (m-1, \nu) | m \in \mathbb{Z}, \alpha' : \nu \rightarrow \nu'\}$. The map $\tau : \mathbb{Z}Q \rightarrow \mathbb{Z}Q$ defined by $\tau(m, \nu) = (m-1, \nu)$ is called the translation of the quiver $\mathbb{Z}Q$. The pair $(\mathbb{Z}Q, \tau)$ is a *translation quiver*, for more details we refer to [4]. The translation quiver $\mathbb{Z}Q$ is stable because its translation is defined everywhere. We recall that an automorphism of a translation quiver is a quiver automorphism commuting with the translation τ .

The quiver $\Gamma(\mathcal{D}^d(\text{mod}KQ))$ consists of two types of components: the regular components and the transjective components, the latter are of the form $\mathbb{Z}Q$, see [14]. The *cluster category* \mathcal{C}_Q of Buan-Marsh-Reineke-Reiten-Todorov is defined to be the orbit category of $\mathcal{D}^d(\text{mod}KQ)$ under the action of the automorphism $\tau^{-1}[1]$, where τ is the Auslander-Reiten translation and $[1]$ is the shift of $\mathcal{D}^d(\text{mod}KQ)$, see [7]. Then \mathcal{C}_Q is also a triangulated Krull-Schmidt category having almost split triangles. Moreover, \mathcal{C}_Q is a 2-Calabi-Yau category [7]. The Auslander-Reiten quiver $\Gamma(\mathcal{C}_Q)$ of \mathcal{C}_Q is the quotient of $\Gamma(\mathcal{D}^d(\text{mod}KQ))$ under the action of the quiver automorphism induced by $\tau^{-1}[1]$. Then, the combinatorics of cluster variables is encoded in the cluster category. That is, if $\mathcal{A} = \mathcal{A}(X, Q)$ is a cluster algebra, with Q an acyclic quiver having $n = |Q_0|$ points, there exists a map $X_\tau : \mathcal{C}_Q \rightarrow \mathbb{Z}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ called the canonical *cluster character*, or the *Caldero-Chapoton map*. The map X_τ induces a bijection between the indecomposables M (up to isomorphism) in \mathcal{C}_Q which have no self-extensions and the cluster

variables X_M , see [8,9]. Under this bijection, clusters correspond to cluster-tilting objects in \mathcal{C}_Q .

It has been shown in [3] that, if \mathcal{A} is a cluster algebra with a seed (X, Q) such that the underlying graph of Q is a tree, then there exists an inverse cluster automorphism $\sigma \in \text{Aut}(\mathcal{A})$ of order two and therefore $\text{Aut}(\mathcal{A}) = \text{Aut}^+(\mathcal{A}) \rtimes \mathbb{Z}_2$. In this section we show that this statement is valid for any acyclic quiver which is mutation equivalent to its opposite quiver.

Let j be an integer, a subquiver of $\mathbb{Z}Q$ of the form (j, Q) is a full subquiver of $\mathbb{Z}Q$ whose set of vertices is $(j, Q)_0 = \{(j, \nu) | \nu \in Q_0\}$.

We denote by C^j the subquiver of the quiver $\mathbb{Z}Q$ of the form (j, Q) .

The notion of *local slice* was introduced in [1] for the connected components of Auslander-Reiten quivers of the categories of modules. For the repetitive quiver $\mathbb{Z}Q$, the notion of local slice coincides with the notion of *section* as defined in [4, p.302].

We consider the 3-dimensional Euclidean space \mathbb{R}^3 with coordinates (x, y, z) , where the frame of reference has a negative orientation as shown in Fig 1.

Our main result will be obtained as a corollary of the following theorem.

Theorem 3.1. *Let Q be a finite connected acyclic quiver such that the translation quiver $\mathbb{Z}Q$ contains two local slices Σ and $\tilde{\Sigma}$ whose quivers are respectively isomorphic to Q and Q^{op} . Then there is an embedding $\eta : \mathbb{Z}Q \rightarrow \mathbb{R}^3$ and an oblique reflection with respect to the plane (xOz) which maps $\eta(\Sigma)$ and $\eta(\tilde{\Sigma})$ to each other.*

Proof. The translation quiver $\mathbb{Z}Q$ contains two local slices Σ and $\tilde{\Sigma}$ whose quivers are isomorphic respectively to Q and Q^{op} . Since the translation quiver $\mathbb{Z}Q$ is stable, we can assume without loss of generality that Σ and $\tilde{\Sigma}$ are such that $\Sigma \cap \tilde{\Sigma} \neq \emptyset$ and Σ is chosen to be the copy of Q at zero, that is $\Sigma = (0, Q) = C^0$.

Because $\tilde{\Sigma}$ is a subquiver of $\mathbb{Z}Q$ and $\Sigma_0 \cap \tilde{\Sigma}_0 \subseteq C^0$, there exists a maximal nonnegative integer k , such that $C^{-j} \cap \tilde{\Sigma} \neq \emptyset$ for all j with $0 \leq j \leq k$. We use the following notation for the vertices of Σ : the vertex $(0, \nu_i)$ is denoted by $(0, \nu_{ji})$, where j is such that the vertex $(-j, \nu_i)$ belongs to $\tilde{\Sigma}$, for $0 \leq j \leq k$. We extend this notation to the translation quiver $\mathbb{Z}Q$ that is $(-m, \nu_{ji}) = \tau^m(0, \nu_{ji})$. We denote by s_j , where $0 \leq j \leq k$, the number of vertices $(0, \nu_{ji})$. Thus we have

$$\sum_{j=0}^k s_j = |Q_0|.$$

The proof is completed in the following three steps (a) (b) (c).

(a) If A_1, A_2, \dots, A_h are points of \mathbb{R}^3 , we denote by $[A_1 A_2 \dots A_h]$ the convex hull of the points A_1, A_2, \dots, A_h .




Fig.1

Let $n = |Q_0|$ and define \mathbb{P}_{nk} to be the convex hull of the points $O, D_0, D_k, E_k, E_0^n, E_k^n, D_0^n, D_k^n$ whose coordinates are given by:

$O = (0, 0, 0)$, $D_0 = (-1, 0, 0)$, $D_k = (-1, k, 0)$, $E_k = (0, k, 0)$, $E_0^n = (0, 0, n)$, $E_k^n = (0, k, n)$, $D_0^n = (-1, 0, n)$, $D_k^n = (-1, k, n)$; then \mathbb{P}_{nk} is a rectangular parallelepiped (see Fig.1).

We want to embed Σ in \mathbb{R}^3 in such a way that the vertices of the embedded quiver $\eta(\Sigma)$ belong to $\mathbb{P}_{nk} \setminus [D_0 D_k D_k^n D_0^n]$ that is, to the parallelepiped \mathbb{P}_{nk} from which we remove the face $D_0 D_k D_k^n D_0^n$. We set $H = (0, 0, n - 1)$ and $T = (-1, 0, n - 1)$. Let $(\mathcal{P}_m)_{m \in \mathbb{Z}}$ be the family of parallel planes in \mathbb{R}^3 defined by the equations $\mathcal{P}_m : y = m$. We decompose Σ into $k + 1$ non-intersecting subquivers Σ^j and construct the embedding of each subquiver separately.

The construction of $\eta(\Sigma)$ is thus done in three steps (a_i) , (a_{ii}) and (a_{iii}) as follows.

(a_i) Construction of $\eta(\Sigma^0)$, where $\Sigma^0 = \Sigma \cap \tilde{\Sigma}$. Then Σ^0 is the full subquiver of Σ whose set of vertices is $\Sigma_0^0 = \{(0, \nu_{0l}) | 1 \leq l \leq s_0\}$. We map the vertices $(0, \nu_{01}), (0, \nu_{02}), \dots, (0, \nu_{0s_0})$ of Σ , to distinct points $A_{01}, A_{02}, \dots, A_{0s_0}$ in $[E_0^n D_0^n T H] \setminus [T D_0^n]$ such that for all $i, j \in \{1, 2, \dots, s_0\}$ the line $(A_{0i} A_{0j})$ is not parallel to a line $(O D_0)$. Each arrow α of Σ from the vertex $(0, \nu_{0i})$ to the vertex $(0, \nu_{0j})$ is mapped to an arrow $\bar{\alpha}$ from A_{0i} to A_{0j} . In other words, we set $\eta(0, \nu_{0i}) = A_{0i}$ with $1 \leq i \leq s_0$ and $\eta(\alpha) = \bar{\alpha}$. The quiver $\eta(\Sigma^0)$ obtained in the plane \mathcal{P}_0 is isomorphic to Σ^0 , we call it an embedding of Σ^0 .

(a_{ii}) Now we construct $\eta(\Sigma^1)$, where Σ^1 is the full subquiver of Σ whose set of vertices is $\Sigma_0^1 = \{(0, \nu_{1l}) | 1 \leq l \leq s_1\}$.

We denote by $t_{\vec{v}}$ the translation in \mathbb{R}^3 by the vector $\vec{v} = (0, 1, -1)$.

Let Σ_0^{11} be the set of all vertices $(0, \nu_{1l})$ of Σ_0^1 such that there exists at least one arrow from a vertex $(0, \nu_{0i})$ in Σ^0 to the vertex $(-1, \nu_{1l})$. And denote by Σ_0^{12} the complement of Σ_0^{11} in Σ_0^1 ; then Σ_0^{12} is the set of the elements $(0, \nu_{1l})$ in Σ_0^1 such that there is no arrow from Σ^0 to $(-1, \nu_{1l})$. Since the quiver Σ is connected, we have $\Sigma_0^{11} \neq \emptyset$.

For the construction of the embedding of the set Σ_0^{11} , we divide Σ_0^{11} into two families of subsets as follows.

The first family consists of the non-intersecting subsets L_1, L_2, \dots, L_h of Σ_0^{11} such that $|L_j| \geq 2$ and for some $(0, \nu_{0i_j}) \in \Sigma_0^0$ there exists an arrow to $(0, \nu_{0i_j})$ from each element L_j . The second family is given by the remaining elements of Σ_0^{11} , the elements which do not belong to any set L_j with $1 \leq j \leq h$.

Now we construct first the embedding of the elements of the set $\Sigma_0^{11} \setminus (L_1 \cup L_2 \dots \cup L_h)$. For an element $(0, \nu_{1r})$ of $\Sigma_0^{11} \setminus (L_1 \cup L_2 \dots \cup L_h)$, we take any arrow originating in $(0, \nu_{1r})$ whose target is $(0, \nu_{0i_r}) \in \Sigma_0^0$. Then the vertex $(0, \nu_{1r})$ is mapped to the point $A_{1r} := t_{\bar{v}}(A_{0i_r})$.

For the construction of the embedding of the elements of the set L_j with $1 \leq j \leq h$, we choose an element $(0, \nu_{1l_j})$ in L_j and we map it to a point $A_{1l_j} := t_{\bar{v}}(A_{0i_j})$ with $A_{0i_j} = \eta(0, \nu_{0i_j})$. In order to define the embedding for the remaining elements of L_j , we proceed as follows. Let Δ_j be the line parallel to (Ox) passing through A_{1l_j} and let $[FG]$ be the intersection of Δ_j and the rectangle $[E_1 D_1 D_1^n E_1^n]$ (we recall that $A_{1l_j} \in [E_1 D_1 D_1^n E_1^n]$). The elements of $L_j \setminus \{(0, \nu_{1l_j})\}$ are mapped to arbitrary distinct points in $[GF] \setminus \{G, F, A_{1l_j}\}$. The elements of L_i , $i \neq j$, $1 \leq i \leq h$ are mapped to the distinct point of the line Δ_i constructed as above. We have $\Delta_i \cap \Delta_j = \emptyset$, for $i \neq j$.

It remains to construct the elements of the set Σ_0^{12} . We map the elements of Σ_0^{12} to distinct points (distinct from the points associated with the elements of Σ_0^{11}) in $t_{\bar{v}}([E_0^n D_0^n T H] \setminus [T D_0^n])$ such that any line containing at least two of them is not parallel to (OD_0) .

Each arrow α of Σ from the vertex $(0, \nu_{1r}) \in \Sigma_0^{12}$ to the vertex $(0, \nu_{1l}) \in \Sigma_0^1$ is mapped to an arrow $\bar{\alpha}$ from $A_{1r} = \eta(0, \nu_{1r})$ to $A_{1l} = \eta(0, \nu_{1l})$ and each arrow β of Σ from the vertex $(0, \nu_{1j}) \in \Sigma_0^{11}$ to the vertex $(0, \nu_{0i}) \in \Sigma_0^0$ is mapped to an arrow $\bar{\beta}$ from $A_{1j} = \eta(0, \nu_{1j})$ to $A_{0i} = \eta(0, \nu_{0i})$.

The obtained quiver $\eta(\Sigma^1)$ is an embedding of the subquiver Σ^1 into the plane $\mathcal{P}_1 : y = 1$. More precisely, $\eta(\Sigma^1)$ is constructed in the rectangle $\mathcal{P}_1 \cap \mathbb{P}_{nk}$.

Thus, we have constructed the quiver $\eta(\overline{\Sigma^0 \cup \Sigma^1})$, where $\overline{\Sigma^0 \cup \Sigma^1}$ is the full subquiver of Σ whose set of vertices is $\Sigma_0^0 \cup \Sigma_0^1$.

(*a_{iii}*) Construction of $\eta(\Sigma^j)$, where Σ^j , $2 \leq j \leq k$ is the full subquiver of Σ whose set of vertices is $\Sigma_0^j = \{(0, \nu_{jl}) | 1 \leq l \leq s_j\}$.

We construct $\eta(\Sigma^j)$ assuming that $\eta(\Sigma^{j-1})$ is constructed.

We reproduce the construction of step *a_{iii}* with Σ^j playing the role of Σ^1 and Σ^{j-1} playing the role of Σ^0 . The vertices of Σ^j are mapped by

η to the rectangle $[E_j D_j D_j^n E_j^n] \setminus [D_j D_j^n]$ and the arrows between two vertices ν and ζ of Σ_0 are mapped to arrows between the corresponding images $\eta(\nu)$ and $\eta(\zeta)$. Therefore, we construct the embedding $\eta(\Sigma^j)$ of Σ^j into the plane $\mathcal{P}_j : y = j$ for $0 \leq j \leq k$, more precisely $\eta(\Sigma^j) \in \mathcal{P}_j \cap \mathbb{P}_{nk}$. Thus, we have constructed the quiver $\eta(\overline{\Sigma^0 \cup \Sigma^1 \dots \cup \Sigma^j})$, where $\overline{\Sigma^0 \cup \Sigma^1 \dots \cup \Sigma^j}$ is the full subquiver of Σ whose set of vertices is $\Sigma_0^0 \cup \Sigma_0^1 \dots \cup \Sigma_0^j$. In other words, we have constructed the embedding of $\overline{\Sigma^0 \cup \Sigma^1 \dots \cup \Sigma^j}$ in $[OD_0 D_0^n E_0^n E_j^n E_j D_j D_j^n] \setminus [D_0 D_0^n D_j^n D_j]$.

Since Σ is finite, the process ends after k iterations and we find the required embedded quiver $\eta(\Sigma)$. By construction we have $\eta(\Sigma) = \eta(\overline{\Sigma^0 \cup \Sigma^1 \dots \cup \Sigma^k})$ and $|\eta(\Sigma)_0| = |\Sigma_0|$. Also, we have an isomorphism of quivers $\eta(\Sigma) \cong \Sigma$ and $\eta(\Sigma)$ is an embedding of Σ in $[OD_0 D_0^n E_0^n E_1^n E_1 D_1 D_1^n] \setminus [D_0 D_0^n D_1^n D_1]$.

(b) Construction of the quiver $\eta(\mathbb{Z}Q)$ in \mathbb{R}^3 . To define $\eta(\mathbb{Z}Q)$ in \mathbb{R}^3 , we need first to construct each $\eta(C^m)$, with $m \in \mathbb{Z}$, where C^m is a copy of the quiver Q .

Let $\vec{u} = (-1, -2, 0)$ and $t_{\vec{u}}$ be the translation in \mathbb{R}^3 by vector \vec{u} .

We extend η to each copy C^m , with $m \in \mathbb{Z}$, by $\eta(C^m) = t_{m\vec{u}}(\eta(\Sigma))$. We complete the construction of $\eta(\mathbb{Z}Q)$ as one would do for the repetitive quiver $\mathbb{Z}Q$ using that $\eta(\tau) = t_{\vec{u}}$. By the construction, there are no arrows which cross each other when considered as segments in \mathbb{R}^3 . Thus we have extended the embedding η to $\mathbb{Z}Q$.

(c) Finally we show that $\eta(\tilde{\Sigma})$ is the image of $\eta(\Sigma)$ under the oblique reflection S with respect to the plane \mathcal{P}_0 , and the direction given by \vec{u} .

If $A = (a, b, c)$ is a point of \mathbb{R}^3 , then the image of A under the oblique reflection S is the point $S(A) = (a - b, -b, c)$.

Let $A_{jr} = (a_{jr}, b_{jr}, c_{jr})$ be a point of $\eta(\Sigma)$ such that $\eta(0, \nu_{jr}) = A_{jr}$, with $(0, \nu_{jr}) \in \Sigma_0$. Since $A_{jr} \in \mathcal{P}_j$, then the second coordinate of A_{jr} is equal to j , and $S(A_{jr}) = (a_{jr} - j, -j, c_{jr})$.

By the definition of η , we have $\eta(\tau) = t_{\vec{u}}$ and $\eta(\tau^{-1}) = t_{-\vec{u}}$.

Now we want to show that $\eta(\tilde{\Sigma}) = S(\eta(\Sigma))$.

Let $N_r \in \tilde{\Sigma}_0$, then $N_r = (-j, \nu_{jr}) = \tau^j(0, \nu_{jr})$, with $1 \leq r \leq s_j$, where $\sum_{j=0}^k s_j = n$. We have


$$\begin{aligned} \eta(N_r) &= \eta(\tau^j(0, \nu_{jr})) \\ &= t_{j\vec{u}}(A_{jr}) \\ &= (a_{jr} - j, -j, c_{jr}) \\ &= S(A_{jr}). \end{aligned}$$

Therefore $\eta(\tilde{\Sigma}_0) \subseteq S(\eta(\Sigma_0))$. Since $|\eta(\tilde{\Sigma}_0)| = |S(\eta(\Sigma_0))|$, then $\eta(\tilde{\Sigma}_0) = S(\eta(\Sigma_0))$. Because the quivers Σ and $\tilde{\Sigma}$ are full subquivers of $\mathbb{Z}Q$, it follows that $S(\eta(\Sigma))$ and $\eta(\tilde{\Sigma})$ are also full subquivers of $\eta(\mathbb{Z}Q)$. Thus $S(\eta(\Sigma)) = \eta(\tilde{\Sigma})$. This completes the proof. \square


Definition 3.1. The quiver $\eta(\mathbb{Z}Q)$ is called a spatial representation of $\mathbb{Z}Q$ in \mathbb{R}^3 .

Remark 3.1. The reflection S defined in \mathbb{R}^3 in step (c) of the proof transforming $\eta(\Sigma)$ to $\eta(\Sigma^{op})$ does not induce necessarily a quiver morphism, as shown in the following example.


Example 3.1. Let Q be a quiver defined by:



The quiver $\mathbb{Z}Q$ contains a local slice $\tilde{\Sigma}$ isomorphic to Q^{op} and that intersects the local slice $\Sigma = (0, Q)$. The local slice Σ is given by



and the local slice $\tilde{\Sigma}$ is given by



We set $\eta(0, 3) = (0, 0, 2)$, $\eta(0, 2) = (0, 1, 1)$ and $\eta(0, 1) = (0, 1, 0)$. We have $S(\eta(0, 3)) = (0, 0, 2)$, $S(\eta(0, 2)) = \eta(-1, 2)$ and $S(\eta(0, 1)) = \eta(-1, 1)$. The reflection S maps the arrow $\eta(0, \alpha_{12}) : \eta(0, 1) \rightarrow \eta(0, 2)$ to the arrow $\eta(-1, \alpha_{12}) : \eta(-1, 1) \rightarrow \eta(-1, 2)$, with $S(\eta(0, 1)) = \eta(-1, 1)$ and $S(\eta(0, 2)) = \eta(-1, 2)$. Thus S does not induce an anti-morphism. The reflection S does not induce a quiver morphism, because for the arrow $\eta(0, \alpha_{23}) : \eta(0, 2) \rightarrow \eta(0, 3)$ of $\eta(\Sigma)$, we have $S(\eta(0, \alpha_{23})) = \eta(0, \alpha'_{23}) : \eta(0, 3) \rightarrow \eta(-1, 2)$.

4. INVOLUTIVE CLUSTER AUTOMORPHISMS

Before giving a first consequence of our result, we need to recall the notion of quiver with length appearing in [6, 10].

Let Q be a quiver, and let $\omega = \alpha_1^{\varepsilon_1} \alpha_2^{\varepsilon_2} \dots \alpha_s^{\varepsilon_s}$ be a walk, where $\alpha_i \in Q_1$ and $\varepsilon_i \in \{-1, 1\}$ for all i . We call *length* of ω the integer defined by $L(\omega) = \sum_{i=1}^s L(\alpha_i^{\varepsilon_i})$, where $L(\alpha_i) = 1$ and $L(\alpha_i^{-1}) = -1$ for all i . Two paths in Q having same source and target are called *parallel paths*. We say that Q is a *quiver with length* when any two parallel paths in Q have the same length. Let ν, ξ be two vertices of the quiver Q with

length and n an integer. The length $L(\nu, \xi)$ between ν and ξ is n if there exists a path from ν to ξ of length n .

Proposition 4.1. *Let Q be a finite connected acyclic quiver such that the translation quiver $\mathbb{Z}Q$ contains two local slices Σ and $\tilde{\Sigma}$ whose quivers are respectively isomorphic to Q and Q^{op} . If the quiver Q is a quiver with length, then the reflection S induces an anti-automorphism of $\mathbb{Z}Q$.*

Proof. We show that the reflection $S : \eta(\Sigma) \rightarrow \eta(\tilde{\Sigma})$ constructed in Theorem 3.1 is an anti-isomorphism of quivers. Consider the embedded quiver $\eta(\mathbb{Z}Q)$ defined in Theorem 3.1. Note that all sinks of Σ belong to the intersection $\Sigma \cap \tilde{\Sigma}$ and therefore are mapped to the plane \mathcal{P}_0 . As before Σ is decomposed into $k + 1$ non-intersecting subquivers Σ^j .

Because the point A_{ji} belongs to the plane $\mathcal{P}_j : y = j$, then $j = \rho(A_{ji}, \mathcal{P}_0)$ where $\rho(A_{ji}, \mathcal{P}_0)$ is the Euclidean distance from the point A_{ji} to the plane \mathcal{P}_0 . Note that by the definition of the reflection S , we have also $j = \rho(S(A_{ji}), \mathcal{P}_0)$. Because the quiver Q is a quiver with length, each subquiver Σ^l of Σ is discrete, that is, it contains no arrows. The only arrows of Σ are from a subquiver Σ^j to a subquiver Σ^{j-1} . Thus in spatial representation of $\eta(\mathbb{Z}Q)$ all arrows are between different planes and therefore are reversed by the reflection S , which shows that S is an anti-morphism. To make this reasoning formal, let $\bar{\alpha}_{il} : A_{ji} \rightarrow A_{j-1l}$ be an arrow of $\eta(\Sigma)$. Then the arrow $(0, \alpha_{il}) : (0, \nu_{ji}) \rightarrow (0, \nu_{j-1l})$ is an arrow of Σ and we have $\eta(0, \alpha_{il}) = \bar{\alpha}_{il}$. On the other hand, $(-j + 1, \alpha'_{il}) : (-j + 1, \nu_{j-1l}) \rightarrow (-j, \nu_{ji})$ is an arrow of $\tilde{\Sigma}$. Since $\eta(-j + 1, \nu_{j-1l}) = S(A_{j-1l})$ and $\eta(-j, \nu_{ji}) = S(A_{ji})$, then $\eta(-j + 1, \alpha'_{il}) = \bar{\alpha}'_{il}$ is an arrow from $S(A_{j-1l})$ to $S(A_{ji})$. Thus $S(\bar{\alpha}_{il}) = \bar{\alpha}'_{il}$. Hence S is an anti-isomorphism from the quiver Σ to the quiver $\tilde{\Sigma}$. The symmetry S is extended to an anti-automorphism of the quiver $\mathbb{Z}Q$. \square

Example 4.1. *Let Q be the Dynkin quiver of type \mathbb{A}_3 given by $1 \xrightarrow{\alpha_{12}} 2 \xrightarrow{\alpha_{23}} 3$. The quiver $\mathbb{Z}Q$ contains a local slice $\tilde{\Sigma}$ isomorphic to Q^{op} which intersects the local slice $\Sigma = (0, Q)$. The quiver Σ is given by*

$$(0, 1) \xrightarrow{(0, \alpha_{12})} (0, 2) \xrightarrow{(0, \alpha_{23})} (0, 3)$$

and the quiver $\tilde{\Sigma}$ is given by

$$(0, 3) \xrightarrow{(0, \alpha'_{23})} (-1, 2) \xrightarrow{(-1, \alpha'_{12})} (-2, 1).$$

We set $\eta(0, 3) = (0, 0, 3)$, $\eta(0, 2) = (0, 1, 1)$ and $\eta(0, 1) = (0, 2, 0)$. We have $S(\eta(0, 3)) = (0, 0, 3)$, $S(\eta(0, 2)) = \eta(-1, 2) = (-1, -1, 1)$ and $S(\eta(0, 1)) = \eta(-1, 1) = (-2, -2, 0)$.

We define the anti-morphism σ on the points by $\sigma(0, 0, 3) = (0, 0, 3)$, $\sigma(0, 1, 1) = (-1, -1, 1)$ and $\sigma(0, 2, 0) = (-2, -2, 0)$; and define on the arrows by $\sigma(\eta(0, \alpha_{23})) = \eta(0, \alpha'_{23})$ and $\sigma(\eta(0, \alpha_{12})) = \eta(-1, \alpha'_{12})$.

The map σ is an anti-isomorphism of quivers, which is extended to an anti-automorphism of quiver $\eta(\mathbb{Z}Q)$. The anti-automorphism σ is a reflection. Therefore, the reflection σ induces an involutive anti-automorphism of the quiver $\mathbb{Z}Q$.

We can now establish the existence of a particular automorphism of the translation quiver $\mathbb{Z}Q$.

Lemma 4.1. *Let Q be a finite connected acyclic quiver such that the translation quiver $\mathbb{Z}Q$ contains a local slice whose quiver is isomorphic to Q^{op} . Then there exists an anti-automorphism σ of $\mathbb{Z}Q$ such that $\sigma^2 = 1$.*

Proof. Let Q be a finite connected acyclic quiver.

If Q is a quiver with length, the reflection S induces an involutive anti-automorphism due to Proposition 4.1.

Suppose that Q is not a quiver with length. According to [10] the quiver Q contains a subquiver of type \tilde{A}_{pq} , with $p \neq q$. Let Σ and $\tilde{\Sigma}$ be two local slices of $\mathbb{Z}Q$ whose quivers are respectively isomorphic to Q and Q^{op} . Without loss of generality, we can choose Σ and $\tilde{\Sigma}$ such that $\Sigma_0 \cap \tilde{\Sigma}_0 \neq \emptyset$. Consider the spatial representation $\eta(\mathbb{Z}Q)$ of $\mathbb{Z}Q$ constructed in the proof of Theorem 3.1. In the case without length, there exist planes \mathcal{P}_j containing non-discrete quivers. It follows from Theorem 3.1 that $\eta(\Sigma)$ and $\eta(\tilde{\Sigma})$ are symmetric to each other with respect to the plane \mathcal{P}_0 of \mathbb{R}^3 via the reflection S . Let \tilde{Q} be the quiver defined by $\tilde{\Sigma} = (0, \tilde{Q})$ and denote by \tilde{C}^m the subquiver of $\mathbb{Z}\tilde{Q}$ of the form (m, \tilde{Q}) . Since $\tilde{\Sigma}$ is a local slice of $\mathbb{Z}Q$, then \tilde{C}^m is a local slice of $\mathbb{Z}Q$ for all integer m .

By the choice of Σ and $\tilde{\Sigma}$, each sink of $\eta(\Sigma)$ is a source of $\eta(\tilde{\Sigma})$. Because $\eta(\Sigma)$ is isomorphic to Q and $\eta(\tilde{\Sigma})$ is isomorphic to Q^{op} , then there exists an anti-isomorphism $\sigma : \eta(\Sigma) \rightarrow \eta(\tilde{\Sigma})$ such that every point of $\eta(\Sigma) \cap \eta(\tilde{\Sigma})$ is invariant under σ . The anti-isomorphism σ induces a permutation σ_j in S_{s_j} , where s_j is defined as in the proof of the Theorem 3.1. The permutation σ_j is viewed as the restriction of $S\sigma$ in $\mathcal{P}_j \cap \eta(\Sigma_0)$.

We define the map $\bar{\sigma} : \eta(\Sigma) \cup \eta(\tilde{\Sigma}) \rightarrow \eta(\Sigma) \cup \eta(\tilde{\Sigma})$ such that the restriction of $\bar{\sigma}$ on $\eta(\Sigma)$ is σ and the restriction of $\bar{\sigma}$ on $\eta(\tilde{\Sigma})$ is σ^{-1} . By the definition, the map $\bar{\sigma}$ is an anti-automorphism of the quiver $\eta(\Sigma) \cup \eta(\tilde{\Sigma})$ which has the following properties:

- (p₁) $\bar{\sigma}$ is an involution
- (p₂) $\bar{\sigma}$ maps \mathcal{P}_j on \mathcal{P}_{-j} , with $0 \leq j \leq k$.

We recall that $t_{\vec{u}}$ is the translation in \mathbb{R}^3 by vector $\vec{u} = (-1, -2, 0)$ defined on the step (b) of the proof of the Theorem 3.1. We extend $\bar{\sigma}$ to $\eta(\mathbb{Z}Q)$ by the map $\tilde{\sigma} : \eta(\mathbb{Z}Q) \rightarrow \eta(\mathbb{Z}Q)$ defined as follows: $\tilde{\sigma}(\eta(C^m)) = t_{m\vec{u}}\bar{\sigma}t_{m\vec{u}}(\eta(C^m))$ for all $m \in \mathbb{Z}$; because $\mathbb{Z}Q \cong \mathbb{Z}\tilde{Q}$ it is

equivalent to define $\tilde{\sigma}$ by $\tilde{\sigma}(\eta(\tilde{C}^m)) = t_{m\bar{u}}\tilde{\sigma}t_{m\bar{u}}(\eta(\tilde{C}^m))$ for all $m \in \mathbb{Z}$. We have $t_{m\bar{u}}(\eta(C^m)) = \eta(\Sigma)$ and $t_{-m\bar{u}}(\eta(C^{-m})) = \eta(\tilde{\Sigma})$ for m non-negative integer. We set $A_{mi} := t_{m\bar{u}}(A_{ji})$, where A_{ji} is defined as in the proof of the Theorem 3.1 and $\sigma_j \in S_{s_j}$ is the permutation induced by σ , with $0 \leq j \leq k$. To make the definition of $\tilde{\sigma}$ more specific, let A_{mi} be a point of $\eta(C^m)$ and A'_{mi} the image of A_{mi} under the reflection S , we have $\tilde{\sigma}(A_{mi}) = A'_{m\sigma_j(i)} = S(A_{m\sigma_j(i)})$ if m is non-negative integer, and $\tilde{\sigma}(A_{mi}) = A'_{m\sigma_j^{-1}(i)} = S(A_{m\sigma_j^{-1}(i)})$ otherwise, with $0 \leq j \leq k$.

It is enough to define $\tilde{\sigma}$ on the points of $\eta(\mathbb{Z}Q)$, because the definition of $\tilde{\sigma}$ on the arrows is induced by the one on the points. The map $\tilde{\sigma}$ is uniquely determined by its value on the points of $\eta(\mathbb{Z}Q)$ and thus extends in the unique way to an anti-automorphism of the quiver $\eta(\mathbb{Z}Q)$. Hence the extension $\tilde{\sigma}$ of σ is an anti-automorphism of the quiver $\eta(\mathbb{Z}Q)$ which maps C^m and \tilde{C}^{-m} to each other.

It remains to show that $\tilde{\sigma}$ is an involution. By the definition, it is sufficient to show this just for the points of $\eta(\mathbb{Z}Q)$. Let A_{mi} be a point of $\eta(\mathbb{Z}Q)$, if m is a non-negative integer then

$$\begin{aligned}\tilde{\sigma}^2(A_{mi}) &= \tilde{\sigma}(A'_{m\sigma_j(i)}) \\ &= A_{m\sigma_j^{-1}(\sigma_j(i))} \\ &= A_{mi};\end{aligned}$$


if m is a negative integer then

$$\begin{aligned}\tilde{\sigma}^2(A_{mi}) &= \tilde{\sigma}(A'_{m\sigma_j^{-1}(i)}) \\ &= A_{m\sigma_j(\sigma_j^{-1}(i))} \\ &= A_{mi}.\end{aligned}$$

The map $\tilde{\sigma}$ is an involutive anti-automorphism of $\eta(\mathbb{Z}Q)$. Thus $\tilde{\sigma}$ induces an involutive anti-automorphism of $\mathbb{Z}Q$. □

In order to illustrate the notations used in the proof of Lemma 4.1 we present an example.

Example 4.2. *We consider the quiver Q of the Example 3.1 defined by:*



We know that $\eta(0, 3) = (0, 0, 2)$, $\eta(0, 2) = (0, 1, 1)$ and $\eta(0, 1) = (0, 1, 0)$. We have $S(\eta(0, 3)) = (0, 0, 2)$, $S(\eta(0, 2)) = \eta(-1, 2)$ and $S(\eta(0, 1)) = \eta(-1, 1)$. We define σ by: the arrow $\eta(0, \alpha_{12}) : \eta(0, 1) \rightarrow \eta(0, 2)$ is mapped to the arrow $\eta(-1, \alpha_{12}) : \eta(-1, 1) \rightarrow \eta(-1, 2)$, that is $\sigma(\eta(0, 1)) = \eta(-1, 2)$ and $\sigma(\eta(0, 2)) = \eta(-1, 1)$; we have also $\sigma(\eta(0, 3)) =$

$\eta(0, 3)$ and $\sigma(\eta(0, \alpha_{13})) = \eta(0, \alpha'_{23})$ and $\sigma(\eta(0, \alpha_{23})) = \eta(0, \alpha'_{13})$. The map σ is extended to an involutive anti-automorphism of $\eta(\mathbb{Z}Q)$. Hence σ induces an involutive anti-automorphism in the transjective component $\Gamma_{tr} \cong \mathbb{Z}Q$ of the cluster category \mathcal{C}_Q .

To simplify the notations, let denote by σ the anti-automorphism of $\mathbb{Z}Q$ defined above. Let $X = \{x_1, x_2, x_3\}$ be the set of variables and \mathcal{A} the cluster algebra with initial seed (X, Q) . We define the map f from \mathcal{A} to \mathcal{A} by: $f(x_1) = \frac{x_1+x_3+x_2x_3^2}{x_1x_2}$, $f(x_2) = \frac{1+x_2x_3}{x_1}$ and $f(x_3) = x_3$. The above data defined an inverse cluster automorphism of \mathcal{A} . By observing that $f(x_1) = \frac{1+f(x_2)x_3}{x_2}$, we have $f^2(x_2) = \frac{1+f(x_2)x_3}{f(x_1)} = x_1$ and $f^2(x_1) = \frac{1+f^2(x_2)x_3}{f(x_2)} = x_2$; thus f is an involution.

One can show by using the categorification of the cluster algebra \mathcal{A} as in [7] that the involutive cluster automorphism f is induced by the involutive anti-automorphism σ .

It has been proved in [3, Theorem 3.11] that, if \mathcal{A} is a cluster algebra with a seed (X, Q) such that Q is a tree then the group of cluster automorphisms is the semidirect product $\text{Aut}\mathcal{A} = \text{Aut}^+\mathcal{A} \rtimes \mathbb{Z}_2$. Moreover this product is not direct. The following result shows that this holds for any finite connected acyclic quiver which is mutation equivalent to its opposite quiver.

Corollary 4.1. *Let Q be a finite connected acyclic quiver such that Q and Q^{op} are mutation equivalent. If \mathcal{A} is a cluster algebra with a seed (X, Q) then the group of cluster automorphisms is the semidirect product $\text{Aut}\mathcal{A} = \text{Aut}^+\mathcal{A} \rtimes \mathbb{Z}_2$. This product is usually not direct.*

Proof. Let Q be a finite connected acyclic quiver and \mathcal{A} a cluster algebra with a seed (X, Q) . We assume without loss of generality that (X, Q) is the initial seed of \mathcal{A} , where the quivers Q and Q^{op} are mutation equivalent. We may assume that Q is not Dynkin, for Dynkin type case we refer to [3, Lemma 3.10]. The transjective component of the cluster category is of the form $\Gamma_{tr} \cong \mathbb{Z}Q$ and $\Gamma_{tr} \cong \mathbb{Z}Q^{op}$. Then, there exist two local slices Σ and $\tilde{\Sigma}$ of Γ_{tr} whose associated quivers are respectively isomorphic to Q and Q^{op} . We may choose Σ and $\tilde{\Sigma}$ such that $\Sigma_0 \cap \tilde{\Sigma}_0 \neq \emptyset$. By [3, Lemma 3.10] the anti-automorphism σ of the Lemma 4.1 induces an inverse cluster automorphism f_σ of order two. Therefore [3, Corollary 2.12] implies that $\text{Aut}\mathcal{A} = \text{Aut}^+\mathcal{A} \rtimes \mathbb{Z}_2$. It was proved in [3, Theorem 3.11] that for the particular case where Q is a tree, this product is not direct. □

The existence of an involutive anti-automorphism of $\mathbb{Z}\Sigma$ give rise to a characterization of finite connected acyclic quiver Σ such that the translation quiver $\mathbb{Z}\Sigma$ contains a local slice whose quiver is isomorphic to Σ^{op} .

Let Q be a finite connected quiver and R a subquiver of Q . We call the *closure* of R and we denote by \overline{R} the smallest full subquiver of Q containing R . A *symmetric quiver* (Q, σ) is a pair consisting of a quiver $Q = (Q_0, Q_1, s, t)$ and an involution $\sigma : Q_0 \amalg Q_1 \longrightarrow Q_0 \amalg Q_1$ with $\sigma(Q_0) = Q_0$ and $\sigma(Q_1) = Q_1$ such that:

- (a) $s(\sigma(\alpha)) = \sigma(t(\alpha))$ and $t(\sigma(\alpha)) = \sigma(s(\alpha))$ for all $\alpha \in Q_1$
- (b) If $\sigma(t(\alpha)) = s(\alpha)$ then $\sigma(\alpha) = \alpha$.

In other words Q is a symmetric quiver if it has an involutive anti-automorphism. For more details about symmetric quivers, we refer to [15].

Definition 4.1. *Let Q be a connected quiver and $(Q^i)_{1 \leq i \leq m}$ a family of full subquivers of Q . We say that the family $(Q^i)_{1 \leq i \leq m}$ is a complete decomposition of Q if the following conditions are satisfied:*

(C1) $Q_0^i \cap Q_0^j = \emptyset$ if $i \neq j$

(C2) $Q = Q^1 \cup Q^2 \dots Q^n$.

(C3) *The only arrows between two connected components are from Q^i to Q^{i-1} for $1 \leq i \leq n - 1$*

Corollary 4.2. *Let Q be a finite connected acyclic quiver such that the translation quiver $\mathbb{Z}Q$ contains a local slice whose quiver is isomorphic to Q^{op} . Then Q possesses a complete decomposition into a family of symmetric subquivers.*

Proof. Let Q be a finite connected acyclic quiver, and Σ be a local slice in $\mathbb{Z}Q$ whose quiver is isomorphic to Q and $\tilde{\Sigma}$ be a local slice in $\mathbb{Z}Q$ whose quiver is isomorphic to Q^{op} . We consider the spatial representation of $\mathbb{Z}Q$. Because of Theorem 3.1 and the definition of the spatial representation of $\mathbb{Z}Q$, we assume that $\eta(\Sigma)$ intersects $k + 1$ planes $\mathcal{P}_m : y = m$, where $0 \leq m \leq k$. We have $\eta(\Sigma^l) = \eta(\Sigma) \cap \mathcal{P}_l$, where Σ^l is the full subquiver of $\eta(\Sigma)$ represented in the plane $\mathcal{P}_l : y = l$, via the embedding η of Theorem 3.1. By construction, the family $(\Sigma^l)_{0 \leq l \leq k}$ is a complete decomposition of Σ . It remains to prove that Σ^l is a symmetric quiver for $0 \leq l \leq k$. We have $\eta(\tilde{\Sigma}^l) = \eta(\tilde{\Sigma}) \cap \mathcal{P}_{-l}$, where $\eta(\tilde{\Sigma}^l)$ is the full subquiver of $\eta(\tilde{\Sigma})$ represented in the plane $\mathcal{P}_l : y = -l$, via the embedding η of Theorem 3.1. So the family $(\tilde{\Sigma}^l)_{0 \leq l \leq k}$ is a complete decomposition of $\tilde{\Sigma}$.

Since $\tilde{\Sigma}$ is a local slice of $\mathbb{Z}Q$, then by the Lemma 4.1, there exists an involutive anti-isomorphism $\sigma : \mathbb{Z}Q \longrightarrow \mathbb{Z}Q$ such that $\sigma(\tilde{\Sigma}) = \tilde{\Sigma}$.

Because $S(\eta(\Sigma^l))$ and $\eta(\tilde{\Sigma}^l)$ are two full subquivers of $\eta(\tilde{\Sigma})$ located in the same plane with $|S(\eta(\Sigma^l))| = |\eta(\tilde{\Sigma}^l)|$, we deduce that they coincide: $S(\eta(\Sigma^l)) = \eta(\tilde{\Sigma}^l)$. Since $S(\eta(\Sigma^l)) \cong \eta(\Sigma^l)$ and $\eta(\tilde{\Sigma}^l) \cong (\eta(\Sigma^l))^{op}$, we get that $\eta(\Sigma^l) \cong (\eta(\Sigma^l))^{op}$. Thus the restriction of σ on $\eta(\Sigma^l)$ is an involutive anti-automorphism. Hence Σ^l is a symmetric quiver for all l , with $0 \leq l \leq k$. This completes the proof of our assertion. \square

Corollary 4.3. *Let Q be a finite connected acyclic quiver and K an algebraically closed field. Then Q and Q^{op} are mutation equivalent if and only if KQ and KQ^{op} are derived equivalent.*

Proof. Let Q be a finite connected acyclic quiver and K an algebraically closed field, KQ the path algebra. Suppose that Q and Q^{op} are mutation equivalent. Then Corollary 4.1 implies that there exists an inverse automorphism of the cluster algebra \mathcal{A} denoted by σ . Then, because of [3, Corollary 3.1] σ induces a triangle equivalence $\sigma_{\mathcal{D}} : \mathcal{D}^b(\text{mod}KQ) \rightarrow \mathcal{D}^b(\text{mod}KQ^{op})$. Conversely, suppose that KQ and KQ^{op} are derived equivalent, the cluster category \mathcal{C}_Q of the finite connected acyclic quiver Q is an orbit category of $\mathcal{D}^b(\text{mod}KQ)$. It was proved by Keller in [16, Theorem 1] that the canonical projection $\pi : \mathcal{D}^b(\text{mod}KQ) \rightarrow \mathcal{C}_Q$ is a triangle functor. Therefore \mathcal{C}_Q and \mathcal{C}^{\vee}_Q are triangulated equivalent. Thus, according to [17], Q and Q^{op} are mutation equivalent. \square

REFERENCES

- [1] Assem, I., Brüstle, T., and Schiffler, R. *Cluster-tilted algebras without clusters*, J. Algebra 324 (2010), no. 9, 2475-2502
- [2] I. Assem, G. Dupont and R. Schiffler, *On a category of cluster algebras*, arXiv:1201.5986v1 [Math.RT], (2012)
- [3] Assem, I. Schiffler, R., and Shramchenko, V. *Cluster Automorphisms*, Proc. Lond. Math. Soc. (3) 104 (2012), no. 6, 1271-1302.
- [4] Assem, I., Simson, D., and Skowroński, A. *Elements of the representation theory of associative algebras. Vol. 1. Techniques of representation theory*. London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
- [5] Auslander, M., Reiten, I., and Smalø, S.O. *Representation theory of Artin algebras. Corrected reprint of the 1995 original. Cambridge Studies in Advanced Mathematics*, 36. Cambridge University Press, Cambridge, 1997.
- [6] Bongartz, K.; Gabriel, P. *Covering spaces in representation-theory.*, Invent. Math. 65 (1981/82), no. 3, 331-378.
- [7] Buan, A.B., Marsh, R.J., Reineke, M. Reiten, I., and Todorov, G. *Tilting theory and cluster combinatorics*, Adv. Math. 204(2006), 572-618.
- [8] Caldero, P., and Chapoton, F. *Cluster algebras as Hall algebras of quiver representations*, Comment. Math. Helv. 81 (2006), 595-616.
- [9] Caldero, P., and Keller, B. *From triangulated categories to cluster algebras*, Invent. Math. 172 (2008), 169-211.
- [10] Chaio, C., Platzeck, M. I., Trepode, S., *On the degree of irreducible morphisms.*, J. Algebra 281 (2004), no. 1, 200-224.
- [11] Fomin, S., and Zelevinsky, A. *Cluster algebras I: Foundations*, J. Amer. Math. Soc. 15 (2002), 497-529
- [12] Fomin, S., and Zelevinsky, A. *Cluster algebras II: Finite type classification*, Invent. Math. 154 (2003), 63-121
- [13] Gekhtman, M., Shapiro, M., and Vainshtein, A. *Cluster algebras and Poisson geometry*, Moscow Math. J. 3 (2003), 899-934.
- [14] Happel, D. *Triangulated categories in the representation theory of finite-dimensional algebras*. London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988.

- [15] Derksen, H. and Weyman, J. *Generalized quivers associated to reductive groups*, Colloquium Math., vol. 94 (2002), no. 2, 151-173.
- [16] Keller, B. *On triangulated orbit categories*, Doc. Math. 10 (2005), 551-581.
- [17] Keller, B., and Reiten, I. *Acyclic Calabi-Yau categories.*, Compos. Math. 144 (2008), no. 5, 1332-1348.
- [18] Lusztig, S., *Canonical bases arising from quantized enveloping algebras.*, J. Amer. Math. Soc. 3 (1990), no. 2, 447-498.

Département de Mathématiques, Université de Sherbrooke 2500, boul.
de l'Université, Sherbrooke, Québec, J1K2R1 Canada .
E-mail address: ndoune.ndoune@usherbrooke.ca