# Privileged Factors in the Thue-Morse Word – A Comparison of Privileged Words and Palindromes

Jarkko Peltomäki

jspelt@utu.fi

Turku Centre for Computer Science TUCS, 20520 Turku, Finland University of Turku, Department of Mathematics and Statistics, 20014 Turku, Finland

#### Abstract

In this paper we study the privileged complexity function of the Thue-Morse word. We prove a recursive formula describing this function, and using the formula we show that the function is unbounded and that the values of the function have arbitrarily large gaps of zeros. This demonstrates that the privileged complexity function of an infinite word can drastically differ from its palindromic complexity function, even though there are relations between these functions. Further we study the behavior of palindromes and privileged words in infinite words and the relation between rich words and privileged words.

Keywords: thue-morse word, palindrome, privileged word, return word, rich word

## 1 Introduction

In this paper we continue the study of so-called privileged words. The study of privileged words was initiated in the articles [KLS13] and [Pel13]. In [KLS13] privileged words were used as a technical tool in order to characterize aperiodic and minimal subshifts with bounded powers. This new class of words has also interest in its own right. For example the paper [Pel13] characterized Sturmian words using their privileged complexity function (the privileged complexity function of a word counts the number of factors of given length that are privileged words).

Privileged words also appear in [FPZ13]. The authors of this paper consider the palindromic length of a word, that is, the minimal number of palindromes needed to write a given word as a concatenation of palindromes. The authors observe that some of their arguments need only properties shared by both privileged words and palindromes, so results on the analogous privileged length of a word are obtained. Namely, it is proven that given an infinite *k*-power free word *w*, for every positive integer *P* there exists a prefix of *w* which cannot be expressed as a concatenation of at most *P* privileged words.

The motivation for defining privileged words comes from the theory of rich words [Gle+09]. Rich words are words containing maximal number of palindromes as factors, and they are characterized by the fact that in a rich word every palindrome is a complete first return to a shorter palindrome. By altering this condition slightly, we get the definition of privileged words: a word is privileged if it is a complete first return to a shorter privileged word; the shortest privileged words are the letters of the alphabet and the empty word. At first glance, privileged words have nothing to do with palindromes. In general, this is indeed true, but there are some connections to rich words. Namely in [KLS13] and [Pel13] it was observed that a word is rich if and only if its set of palindromes coincides with its set of privileged words. Every position in a word introduces exactly one new privileged factor and at most one new palindromic factor. This means that in

<sup>© 2015,</sup> Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International.

a sense privileged factors which are not palindromes measure the palindromic defect of a word. The palindromic defect of a word is the number of positions in the word that do not introduce a new palindrome; rich words are the words with zero defect. Other than this connection to richness, privileged words have other similarities with palindromes. For instance a privileged prefix of a privileged word occurs also as a suffix, just as a palindromic prefix of a palindrome occurs also as a suffix.

In general privileged words behave differently from palindromes. In this paper we show that the privileged complexity function of an infinite word may behave in a much more complex way than its palindromic complexity function. We investigate the privileged complexity function  $\mathcal{A}_t$ of the Thue-Morse word t and prove that  $\mathcal{A}_t$  is unbounded and that its values have arbitrarily long gaps of zeros. In contrast it has been proven in [DZ00] and [All+03] that the palindromic complexity function of fixed points of primitive morphisms is bounded. Also, as is easily seen, for any palindromic complexity function two consecutive zeros would imply that the function takes only finitely many nonzero values. To obtain the results we derive complete recursive formulas for computing  $\mathcal{A}_t$ . Various other kinds of complexity functions for the Thue-Morse word have been considered in the past. Recently in [GMS13] Goč, Mousavi and Shallit proved that there exists a system of recurrences for the number of unbordered factors of given length in the Thue-Morse word.

After defining the necessary notation and definitions, in Section 3 we conclude the study of the function  $\mathcal{A}_t$  initiated in [Pel13], and present the main results of this paper. We describe the relations between different types of privileged factors in the Thue-Morse word, and give a recursive formula for computing  $\mathcal{A}_t$ . We study the asymptotic behavior of this function, and prove that it is unbounded. We also verify a conjecture of [Pel13] stating that the values of the function  $\mathcal{A}_t$  contain arbitrarily long gaps of zeros.

In Section 4 we briefly study the privileged palindrome complexity function of the Thue-Morse word, that is, the function counting the number of privileged factors which are also palindromes. As in Section 3, we give a recursive formula for computing the values of this function, and study its asymptotic behavior and gaps of zeros.

In the last section we compare palindromes and privileged words. First we further sharpen the connection to palindromic richness mentioned above. Secondly we investigate when palindromic factors can form a proper subset of privileged factors in a word (remember: in rich words palindromes are exactly the privileged words).

### 2 Preliminaries

In this text, we let *A* denote a finite *alphabet*, which is a finite non-empty set of symbols. The elements of *A* are called *letters*. A (finite) *word* over *A* is a sequence of letters. To the empty sequence corresponds the *empty word*, denoted by  $\varepsilon$ . The set of all finite words over *A* is denoted by *A*<sup>\*</sup>. The set of non-empty words over *A* is the set  $A^+ := A^* \setminus {\varepsilon}$ . A natural operation on words is concatenation. Under this operation  $A^*$  is a free monoid over *A*. The concatenation of two words *u* and *v* is denoted by *uv*. The letters occurring in the word *w* form the *alphabet* of *w*. From now on we assume that binary words are over the alphabet  $\{0, 1\}$ . Given a finite word  $w = a_1a_2 \cdots a_n$  of *n* letters, we say that the *length* of *w*, denoted by |w|, is equal to *n*. By convention the length of the empty word is 0. The set of all words of length *n* over the alphabet *A* is denoted by  $A^n$ . A word *w* is *primitive* if *w* is of the form  $z^n$  if and only if n = 1. A word is *non-primitive* if it is not primitive.

An *infinite word* w over A is a function from the natural numbers to A. We consider such a function as a sequence indexed by the natural numbers with values in A. We write concisely  $w = a_1a_2a_3\cdots$  with  $a_i \in A$ . The set of infinite words is denoted by  $A^{\omega}$ . For an infinite word w we write  $|w| = \infty$ . An infinite word w is said to be *ultimately periodic* if we can write it in the form  $w = uv^{\omega} = uvvv \cdots$  for some words  $u, v \in A^*$ . If  $u = \varepsilon$ , then w is said to be *periodic*. An infinite word which is not ultimately periodic is *aperiodic*.

A finite word *u* is a *factor* of the finite or infinite word *w* if we can write that w = zuv for some

 $z \in A^*$  and  $v \in A^* \cup A^\omega$ . If  $z = \varepsilon$ , then the factor u is called a *prefix* of w. If  $v = \varepsilon$ , then we say that u is a *suffix* of w. If a word u is both a prefix and a suffix of w, then u is a *border* of w. The set of factors of w is denoted by  $\mathcal{F}(w)$ . The set  $\mathcal{F}_w(n)$  is defined to contain all factors of w of length n. We call u a central factor of w if there exists a factorization w = xuy with |x| = |y|. If  $w = a_1a_2 \cdots a_n$ , then we let  $w[i, j] = a_i \cdots a_j$  whenever the choices of positions i and j make sense. This notion is extended to infinite words in a natural way. An *occurrence* of u in w is a a position i such that w[i, i + |u| - 1] = u. If such a position exists, then we say that u occurs in w. By  $|w|_u$  we denote the number of occurrences of the factor u in the word w. If  $|w|_u = 1$ , then we say that u is unioccurrent in w. We say that a position i introduces a factor u if w[i - |u| + 1, i] = u and u is unioccurrent in w[1, i]. A complete first return to the word u is a word starting and ending with u and containing exactly two occurrences of u (occurrences of u may overlap). A word which is a complete first return word. A complete return factor is a factor of some word which is a complete return word.

An infinite word w is *recurrent* if each of its factors occurs in it infinitely often. The word w is called *uniformly recurrent* if each factor u occurs in it infinitely often, and the gap between two consecutive occurrences of u in w is bounded by a constant depending only on u. Equivalently w is uniformly recurrent if for each factor u there exists an integer R such that every factor of w of length R contains an occurrence of u.

Let *A* and *B* be two alphabets. A *morphism* from  $A^*$  to  $B^+$  is a mapping  $\varphi : A^* \to B^+$  such that  $\varphi(uv) = \varphi(u)\varphi(v)$  for all words  $u, v \in A^*$ . A morphism  $\varphi : A^* \to A^+$  is said to be prolongable on the letter *a* if  $\varphi(a)$  begins with letter *a* and  $|\varphi(a)| > 1$ . Then clearly  $\varphi^n(a)$  is a prefix of  $\varphi^{n+1}(a)$ , so we obtain a fixed point  $\varphi^{\omega}(a) := \lim_{n\to\infty} \varphi^n(a)$ . A morphism  $\varphi : A^* \to A^+$  is *primitive* if there exists  $n \ge 1$  such that for all  $a \in A$  the image  $\varphi^n(a)$  contains every letter of *A* at least once.

The word  $\partial_{i,j}(u)$ , where  $i + j \leq |u|$ , is obtained from the word u by deleting i letters from the beginning, and j letters from the end. Let  $\varphi$  be a morphism with fixed point  $w = \varphi^{\omega}(a)$ . We say that a factor u of w admits an *interpretation*  $s = (x_0x_1 \cdots x_{n+1}, i, j)$  by  $\varphi$  if  $u = \partial_{i,j}(\varphi(x_0x_1 \cdots x_{n+1}))$  where  $x_i$  are letters,  $0 \leq i < |\varphi(x_0)|$  and  $0 \leq j < |\varphi(x_{n+1})|$ . The word  $x_0x_1 \cdots x_{n+1}$  is called the ancestor of the interpretation s. In this paper we are focused on the Thue-Morse morphism, and in this particular case all sufficiently long factors of the fixed point have a unique interpretation (this is demonstrated in Section 3). Thus it is convenient to just talk about the interpretation and the ancestor of u by  $\varphi$ . In a factor u of w we often separate images of letters by bars. For example if  $\varphi : 0 \mapsto 01, 1 \mapsto 10$  is the Thue-Morse morphism, then the word 01100 has ancestor 010, and we place bars as follows: 01|10|0. If a factor has a unique interpretation, then there is only one way to place the bars in that factor, and conversely.

The *reversal*  $\tilde{w}$  of  $w = a_1 a_2 \cdots a_n$  is the word  $\tilde{w} = a_n \cdots a_2 a_1$ ; notation  $w^{\sim}$  is also used for  $\tilde{w}$ . If  $\tilde{w} = w$ , then we say that w is a *palindrome*. By convention the empty word is a palindrome. The set of palindromes of w is denoted by  $\mathcal{P}al(w)$ . Moreover we define  $\mathcal{P}al_w(n) = \mathcal{P}al(w) \cap \mathcal{F}_w(n)$ . We say that a word w is *closed under reversal* if for each  $u \in \mathcal{F}(w)$  it holds that  $\tilde{u} \in \mathcal{F}(w)$ . It is well-known that a finite word w contains at most |w| + 1 distinct palindromic factors (see Proposition 2 of [DJP01]). Equality is attained if each position of w introduces exactly one new palindrome. *The (palindromic) defect*  $\mathcal{D}(w)$  of w is the number  $|w| + 1 - |\mathcal{P}al(w)|$ . The defect measures the abundance of palindromes in w. If  $\mathcal{D}(w) = 0$  (i.e., it contains the maximum possible number of distinct palindromes), then w is called *rich*. The defect for infinite word w is defined as  $\mathcal{D}(w) = \sup{\mathcal{D}(u) : u}$  is a prefix of w}. The defect of a finite word w is equal to the number of prefixes u of w such that u does not have a unioccurrent longest palindromic suffix. For a good reference on defect and rich words see [Gle+09].

*Privileged words* were first defined in [KLS13] and further developed in [Pel13]. The set of privileged words over the alphabet A, denoted by Pri(A), is defined inductively as follows:

- $u \in Pri(A)$  if u is a complete first return to a shorter privileged word  $v \in Pri(A)$ ,
- the shortest privileged words are the letters of A and the empty word  $\varepsilon$ .

The set of privileged factors of a word w is denoted Pri(w). We define  $Pri_w(n) = Pri(w) \cap \mathcal{F}_w(n)$ . We set  $\mathcal{A}_w(n) = |Pri_w(n)|$ . The function  $\mathcal{A}_w$  is called the *privileged complexity function* of w. Next we list elementary properties of privileged words proved in [Pel13]. For completeness we provide full proofs.

**Lemma 2.1.** *Let w be a privileged word and u an arbitrary privileged prefix (respectively suffix) of w. Then u is a suffix (respectively prefix) of w.* 

*Proof.* If  $|w| \le 1$  or u = w, then the claim is clear. Suppose that  $|w| \ge 2$  and |u| < |w|. By definition w is a complete first return to a shorter privileged word v. If |v| < |u|, then by induction v is a suffix of u, and thus v would have at least three occurrences in w which is impossible. If u = v, then the claim is clear. Finally assume that |v| > |u|. Then by induction u is a suffix of v, and thus a suffix of w. The proof in the case that the roles of prefix and suffix are reversed is symmetric.

**Lemma 2.2.** Let *w* be a privileged word, and *u* its longest proper privileged prefix (or suffix). Then *w* is a complete first return to *u*.

*Proof.* If  $|w| \le 1$ , then there is nothing to prove. Suppose that  $|w| \ge 2$  and that w is a complete first return to privileged word v. Now if |u| > |v|, then v is a prefix of u, and thus by Lemma 2.1 also a suffix of u. Hence w has at least three occurrences of v, a contradiction. Therefore  $|u| \le |v|$ , and by the maximality of u, u = v, which proves the claim. The proof in the case that the roles of prefix and suffix are reversed is symmetric.

**Lemma 2.3.** Every position in a word introduces exactly one new privileged factor.

*Proof.* It is sufficient to show that appending a new letter *a* to a word *w* introduces exactly one new privileged factor. Since letters are privileged, the word *wa* has as a suffix a privileged word *u* of maximal length. Assume on the contrary that *u* occurs in *w*. Then *wa* has as a suffix a complete first return to *u*, denote it by *v*. By definition *v* is privileged. This contradicts the maximality of *u*, so indeed *u* does not occur in *w*. Finally if appending *a* introduced another privileged factor, say *z*, then by maximality of *u*, we would have that |z| < |u|. Thus *z* would be a suffix of *u*, and by Lemma 2.1 *z* would be a prefix of *u*. Consequently *z* would occur already in *w* contradicting the assumption that appending *a* introduced *z*.

## **3** The Privileged Complexity of the Thue-Morse Word

In this section we prove a recursive formula for the privileged complexity function of the Thue-Morse word. Moreover we study the asymptotic behavior of the function and the occurrences of zeros in its values.

Let  $t = 0110100110010110 \cdots$  be the infinite Thue-Morse word (see Chapter 2 of [Lot83]). The word *t* is a fixed point of the morphism  $\varphi$  and its square  $\theta = \varphi^2$ .

$$\varphi: \begin{array}{cc} 0 \mapsto 01 \\ 1 \mapsto 10 \end{array} \qquad \theta: \begin{array}{cc} 0 \mapsto 0110 \\ 1 \mapsto 1001 \end{array}$$

The word *t* has the following well-known property (*an overlap* is a factor of the form *auaua* where *a* is a letter):

**Theorem 3.1.** The Thue-Morse word t does not contain overlaps, that is, it is overlap-free.

*Proof.* For a proof see e.g. Theorem 2.2.3 of [Lot83].

By this theorem the longest privileged proper border u of a privileged factor w of t cannot overlap with itself in w, that is,  $|u| \le |w|/2$ . In what follows, we implicitly assume this fact. Using overlap-freeness, by mere inspection we obtain the following:

**Lemma 3.2.** Every factor of t of length at least 4 admits a unique interpretation by  $\varphi$ . Every factor of t of length at least 7 admits a unique interpretation by  $\theta$ .

Using this lemma we are able to prove the following important proposition:

**Proposition 3.3.** Let  $w, u \in \mathcal{F}(t)$  be such that  $|w| \ge |u| \ge 2$ . Then  $|\theta(w)|_{\theta(u)} = |w|_u$ .

*Proof.* Clearly always  $|\theta(w)|_{\theta(u)} \ge |w|_u$ . Say  $\theta(u)$  occurs in  $\theta(w)$ , so  $\theta(w) = \alpha \theta(u)\beta$  for some words  $\alpha$  and  $\beta$ . There must exist words  $\lambda$  and  $\mu$  such that  $\theta(\lambda) = \alpha$  and  $\theta(\mu) = \beta$ , since otherwise  $\theta(u)$  would admit two interpretations by  $\theta$ , which is impossible by Lemma 3.2 as  $|\theta(u)| \ge 8$ . Hence  $w = \lambda u \mu$ . This proves that  $|\theta(w)|_{\theta(u)} \le |w|_u$ .

The following interesting result can be inferred from Theorem 4.3. in [BPS07].

**Proposition 3.4.** [BPS07]. Every factor except 0 and 1 in the Thue-Morse word has exactly 4 complete returns.

Let *E* be the morphism defined by E(0) = 1 and E(1) = 0. As for every  $w \in \mathcal{F}(t)$  also  $E(w) \in \mathcal{F}(t)$ , we can focus primarily on factors beginning with 0. We let  $\mathcal{P}ri_u(n)$  denote the set  $\mathcal{P}ri_t(n) \cap u \cdot \{0,1\}^*$ , and we set  $\mathcal{A}_u(n) = |\mathcal{P}ri_u(n)|$ . As 111 is not a factor of *t*, the complete first returns to 0 are 00,010 and 0110. Clearly privileged factors beginning with letter 0 with length greater than one can be divided into three groups depending on the first four letters of the word. We have that

$$\mathcal{P}ri_0(n) = \mathcal{P}ri_{00}(n) \cup \mathcal{P}ri_{010}(n) \cup \mathcal{P}ri_{0110}(n)$$

for n > 1. Thus for the privileged complexity function  $\mathcal{A}_t$  of the Thue-Morse word we have that

$$\frac{1}{2}\mathcal{A}_t(n) = \mathcal{A}_{00}(n) + \mathcal{A}_{010}(n) + \mathcal{A}_{0110}(n)$$

for n > 1. Using overlap-freeness, we can easily see that  $Pri_0(1) = \{0\}$ ,  $Pri_0(2) = \{00\}$ ,  $Pri_0(3) = \{010\}$  and  $Pri_0(4) = \{0110\}$ . Hence  $\mathcal{A}_t(1) = \mathcal{A}_t(2) = \mathcal{A}_t(3) = \mathcal{A}_t(4) = 2$ . Next we state the main results of this paper.

**Theorem 3.5.** The privileged complexity function  $A_t$  of the Thue-Morse word satisfies

$$\begin{aligned} \mathcal{A}_{t}(0) &= 1, \mathcal{A}_{t}(1) = \mathcal{A}_{t}(2) = \mathcal{A}_{t}(3) = \mathcal{A}_{t}(4) = 2, \\ \frac{1}{2}\mathcal{A}_{t}(4n) &= 3\mathcal{A}_{00}(n) + \mathcal{A}_{010}(n) + \mathcal{A}_{010}(n+1) + \mathcal{A}_{0110}(n+1) & \text{for } n \geq 2, \\ \frac{1}{2}\mathcal{A}_{t}(4n-2) &= \mathcal{A}_{00}(4(n-1)) + \mathcal{A}_{010}(4n) + \mathcal{A}_{0110}(4n) & \text{for } n \geq 2, \\ \mathcal{A}_{t}(2n+1) &= 0 & \text{for } n \geq 2. \end{aligned}$$

Theorem 3.5 is directly implied by Proposition 3.9 and the following theorem which allows the computation of values of  $\mathcal{A}_t$ .

**Theorem 3.6.** The functions  $\mathcal{A}_{00}(n)$ ,  $\mathcal{A}_{010}(n)$  and  $\mathcal{A}_{0110}(n)$  satisfy

$$\begin{aligned} \mathcal{A}_{00}(4n) &= 2\mathcal{A}_{00}(n),\\ \mathcal{A}_{00}(4n-2) &= \mathcal{A}_{0110}(4n),\\ \mathcal{A}_{0110}(4n) &= \mathcal{A}_{00}(n) + \mathcal{A}_{010}(n),\\ \mathcal{A}_{0110}(4n-2) &= \mathcal{A}_{00}(4(n-1)),\\ \mathcal{A}_{010}(4n) &= \mathcal{A}_{010}(n+1) + \mathcal{A}_{0110}(n+1),\\ \mathcal{A}_{010}(4n-2) &= \mathcal{A}_{010}(4n) \end{aligned}$$

for all  $n \geq 2$ .

*Proof.* The claim follows from Corollaries 3.17, 3.20 and 3.23 proven below.

2-16	18-32	34-48	50-64	66-80	82-96	98-112	114-128
2	2	14	0	2	0	16	0
2	2	6	0	2	0	8	0
4	4	4	0	2	4	4	6
8	8	8	0	2	12	4	18
8	8	8	0	2	12	4	18
4	4	4	0	2	4	4	6
0	6	2	0	2	4	4	8
0	14	2	0	2	12	4	24

Table 1: Values  $\mathcal{A}_t(n)$  for  $n = 2, 4, 6, 8, \dots, 128$  (the even numbers from 2 to 128).

$\alpha_1 = 00101100$	$\beta_1 = 01011010$	$\gamma_1 = 01100110$
$\alpha_2 = 00110100$	$\beta_2 = 010110011010$	$\gamma_2 = 011010010110$
$\alpha_3 = 001100$	$\beta_3 = 010010$	$\gamma_3 = 0110010110$
$\alpha_4 = 0010110100$	$\beta_4 = 0100110010$	$\gamma_4 = 0110100110$

Table 2: The set  $\mathcal{R}$  of all complete first returns to 00,010 and 0110 in *t*.

Using the formulas of Theorems 3.5 and 3.6 values of  $A_t$  can be computed. See Table 1. It is interesting to compare the privileged complexity with the palindromic complexity:

**Theorem 3.7.** [All+03; BBL08] *The palindromic complexity function*  $\mathcal{P}_t(n)$  *of the Thue-Morse word satisfies* 

$$\begin{aligned} & \mathcal{P}_t(0) = 1, \mathcal{P}_t(1) = \mathcal{P}_t(2) = \mathcal{P}_t(3) = \mathcal{P}_t(4) = 2, \\ & \mathcal{P}_t(4n) = \mathcal{P}_t(4n-2) = \mathcal{P}_t(n) + \mathcal{P}_t(n+1) & \text{for } n \ge 2, \\ & \mathcal{P}_t(2n+1) = 0 & \text{for } n \ge 2. \end{aligned}$$

In Table 2 we list the set  $\mathcal{R}$  of all complete first returns to 00,010 and 0110. These words are needed later on. We leave it to the reader to verify that these words actually are factors of *t*. By Proposition 3.4 the list is exhaustive (this fact is also easily verified directly).

**Lemma 3.8.** If  $w \in Pri_0(n)$  with n > 4, then w begins with a word in  $\mathcal{R}$ .

*Proof.* Since |w| > 4, the word w has proper prefix u where u is one of the words 00,010 or 0110. Since w is privileged, the word u is also a suffix of w, so w must have a complete first return to u as a prefix. The claim follows.

We see that we have at least two privileged factors beginning with 0 of odd length, namely 0 and 010. It turns out that there are no more:

#### **Proposition 3.9.** [Pel13] $\mathcal{A}_t(2n+1) = 0$ for $n \ge 2$ .

*Proof.* We may focus on privileged factors beginning with 0. Let w, |w| > 4, be a privileged factor of *t* beginning with 0. Now *w* begins with one of the three privileged words 00, 010 or 0110. With respect to the morphism  $\varphi$  the bars must be placed as follows: 0|0, 01|0 or 0|10 and 01|10. Now if *w* begins with 00 (respectively 0110), then it also ends with 00 (respectively 0110), and by the placement of the bars we immediately see that *w* has even length. Assume then that *w* begins with 010. As |w| > 4, by Lemma 3.8 *w* has as a prefix one of the words  $\beta_1 = 01|01|10|10, \beta_2 = 01|01|10|01|10|10, \beta_3 = 0|10|01|0 \text{ or } \beta_4 = 0|10|01|10|01|0 \text{ (bars with respect to <math>\varphi$ ). If *w* begins with some  $\beta_i$ , then it also ends with  $\beta_i$ . From the placement of the bars we see that |w| is necessarily even.

We frequently need complete information on short privileged factors. The next lemma provides this knowledge. In what follows, we assume the lemma to be known. **Lemma 3.10.** Let  $w \in Pri_0(n)$  with  $n \le 12$ . Then  $w \in \{0, 010, 0110\} \cup \mathcal{R}$ .

*Proof.* It was already remarked that if  $|w| \le 4$ , then  $w \in \{0,010,0110\}$ . If |w| > 4, then by Lemma 3.8 the word w begins with a word in  $\mathcal{R}$ . If  $w \notin \mathcal{R}$ , then as w is privileged, w must have as a prefix a complete first return to some word in  $\mathcal{R}$ . This prefix u cannot overlap with itself. Since  $|w| \le 12$ , it thus follows that  $|u| \le 6$ . The words of minimal length in  $\mathcal{R}$  have length 6, so  $w = u^2$  and  $u \in \{\alpha_3, \beta_3\}$ . However, both  $\alpha_3^2$  and  $\beta_3^2$  contain a third power (that is, an overlap) yielding a contradiction. Thus  $w \in \mathcal{R}$ .

Next we characterize the different classes of privileged factors in the Thue-Morse word. In what follows, we say that a word w is  $\theta$ -invertible if there exists a word u such that  $\theta(u) = w$ . Recall the words  $\alpha_i$ ,  $\beta_i$  and  $\gamma_i$  from Table 2.

**Lemma 3.11.** Let  $w \in Pri_{00}(n)$  for some n > 2. Then

(i)  $4 \mid |w| \iff 1w110 \text{ or } 011w1 \text{ is a } \theta$ -invertible factor of  $t \iff w$  begins with  $\alpha_1$  or  $\alpha_2$ ,

(ii)  $4 \nmid |w| \iff 1w1$  is a  $\theta$ -invertible factor of  $t \iff w$  begins with  $\alpha_3$  or  $\alpha_4$ .

*Proof.* By Lemma 3.2 all factors of *t* of length at least 7 admit a unique interpretation by  $\theta$ , so in  $\alpha_1, \alpha_2$  and  $\alpha_4$  there is a unique way to place bars:  $\alpha_1 = 001|0110|0, \alpha_2 = 0|0110|100$  and  $\alpha_4 = 001|0110|100$ . For  $\alpha_3$  there are potentially two ways to place bars:  $\alpha_3 = 001|100$  and  $\alpha_3 = 0|0110|0$ . However the latter is not possible as  $(0110)^3$  is not a factor of *t*.

(*i*) Assume that 4 | |w|. If w begins with  $\alpha_4$ , then it also ends with  $\alpha_4$ . From the placement of the bars it can be seen that this is not possible; it would follow that  $4 \nmid |w|$ . Similarly w cannot begin with  $\alpha_3$ . Hence w must begin with  $\alpha_1$  or  $\alpha_2$ . On the other hand if w begins with  $\alpha_1$  or  $\alpha_2$ , then 1w110 or 011w1 must be  $\theta$ -invertible by the placement of the bars. Then clearly  $4 \mid |w|$ .

(*ii*) Assume that  $4 \nmid |w|$ . By (i) w has to begin with  $\alpha_3$  or  $\alpha_4$ . In either case 1w1 is a  $\theta$ -invertible factor of t. The other direction is also clear: if w begins with  $\alpha_3$  or  $\alpha_4$ , then by (i) it must be that  $4 \nmid |w|$ .

**Lemma 3.12.** Let  $w \in Pri_{010}(n)$  for some  $n \ge 1$ . Then

(i)  $4 \mid |w| \iff 10w01$  is a  $\theta$ -invertible factor of  $t \iff w$  begins with  $\beta_1$  or  $\beta_2$ ,

- (*ii*)  $4 \nmid |w|, 2 \mid |w| \iff 011w110$  *is a*  $\theta$ *-invertible factor of*  $t \iff w$  *begins with*  $\beta_3$  *or*  $\beta_4$ *,*
- (iii)  $4 \nmid |w|, 2 \nmid |w| \iff w = 010.$

*Proof.* From Proposition 3.9 it follows that (iii) holds.

As in the previous proof, we know the placements of bars in the words  $\beta_1, \beta_2, \beta_3$  and  $\beta_4$ :  $\beta_1 = 01|0110|10, \beta_2 = 01|0110|0110|10, \beta_3 = 0|1001|0$  and  $\beta_4 = 0|1001|1001|0$ .

(*i*) Assume that  $4 \mid |w|$ . As in the previous proof, from the placement of the bars we see that w cannot begin with  $\beta_3$  or  $\beta_4$ , and hence it must start with  $\beta_1$  or  $\beta_2$ . Thus 10w01 is  $\theta$ -invertible. Again the unique placement of the bars implies that the converse is also true.

(*ii*) By (i) it is enough to note that if w begins with  $\beta_3$  or  $\beta_4$ , then 011w110 is  $\theta$ -invertible.

**Lemma 3.13.** Let  $w \in Pri_{0110}(n)$  for some n > 4. Then

(i)  $4 \mid |w| \iff w$  is a  $\theta$ -invertible factor of  $t \iff w$  begins with  $\gamma_1$  or  $\gamma_2$ ,

(ii)  $4 \nmid |w| \iff 10w$  or w01 is a  $\theta$ -invertible factor of  $t \iff w$  begins with  $\gamma_3$  or  $\gamma_4$ .

*Proof.* The placement of bars is known:  $\gamma_1 = 0110|0110, \gamma_2 = 0110|1001|0110, \gamma_3 = 01|1001|0110$  and  $\gamma_4 = 0110|1001|10$ . As in the two previous proofs, by looking at the placements of the bars the claim straightforwardly follows.

The three previous lemmas allow us to prove the following useful result:

**Corollary 3.14.** Let  $w \in Pri_t(n)$  and u be its longest privileged proper prefix such that |u| > 4. Then 4 | |w| if and only if 4 | |u|.

*Proof.* Let  $S = \{\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2\}$ . Assume that  $4 \mid |w|$ . The lemmas 3.11, 3.12 and 3.13 imply that w begins with some  $v \in S$ . Since |u| > 4, it follows from Lemma 3.8 that u has as a prefix a word in  $\mathcal{R}$ . Since no word in the set  $\mathcal{R}$  is a proper prefix of any word in  $\mathcal{R}$ , it follows that u begins with v. The same three lemmas now imply that  $4 \mid |u|$ . On the other hand if  $4 \mid |u|$ , then u begins with a word in S. Consequently w begins with a word in S, so  $4 \mid |w|$ .

Let us then define the following functions  $f_1, g_1 : \{0,1\}^* \to \{0,1\}^* : f_1(w) = \partial_{1,3}(\theta(1w))$  and  $g_1(w) = \partial_{3,1}(\theta(w1))$ .

**Lemma 3.15.** Let  $n \ge 2$ . The function  $f_1$  is a bijection  $\operatorname{Pri}_{00}(n) \to \operatorname{Pri}_{\alpha_1}(4n)$  and the function  $g_1$  is a bijection  $\operatorname{Pri}_{00}(n) \to \operatorname{Pri}_{\alpha_2}(4n)$ .

*Proof.* We will first prove the claim for  $f_1$ . If n = 2, then  $Pri_{00}(2) = \{00\}$  and  $Pri_{\alpha_1}(8) = \{\alpha_1\}$ , so the claim indeed holds. The latter part of this proof shows that if  $Pri_{\alpha_1}(4n) \neq \emptyset$ , then also  $Pri_{00}(n) \neq \emptyset$ . Thus the claim holds also if n = 3, 4, 5, as then  $Pri_{00}(n) = \emptyset$ . Assume that n > 5.

Let  $w \in Pri_{00}(n)$ , and let v be its longest privileged proper prefix. Note that now  $|v| \ge 2$ . As v begins with 00 it follows by induction that  $f_1(v) \in Pri_{\alpha_1}(t)$ . As v is always preceded by the letter 1, w = vw'1v, and thus

$$f_1(w) = \partial_{1,3}(\theta(1v)\theta(w')\theta(1v)) = \partial_{1,3}(\theta(1v))110\theta(w')1\partial_{1,3}(\theta(1v)) = f_1(v)110\theta(w')1f_1(v).$$

By Lemma 3.11 the factor  $f_1(v)$  is always preceded by 1 and followed by 110. Thus if  $f_1(v)$  would occur more than twice in  $f_1(w)$ , then  $\theta(v) = 1f_1(v)110$  would occur more than twice in  $\theta(w)$  which is impossible by Proposition 3.3. Hence  $f_1(w)$  is a complete first return to the privileged word  $f_1(v)$ , and thus  $f_1(w) \in Pri_{\alpha_1}(4n)$ .

Assume then that  $w \in \mathcal{P}ri_{\alpha_1}(4n)$ . By Lemma 3.11 there exists  $z \in \mathcal{F}_t(n+1)$  such that  $\theta(z) = 1w110$ . Write  $u = \partial_{1,0}(z)$ . Then  $f_1(u) = w$ . Let v be the longest privileged proper prefix of w. The assumption n > 5 implies that |v| > 4 (the maximum length of a word in  $\mathcal{R}$  is 12), so by Corollary 3.14 we have that 4 ||v|. Thus by induction there exists  $s \in \mathcal{P}ri_{00}(t)$  such that  $f_1(s) = v$ . Thus  $f_1(u) = w = f_1(s) \cdots f_1(s)$ , and so u begins and ends with s. Now if s would occur more than twice in u, then as s is always preceded by 1,  $f_1(s) = v$  would occur more than twice in w which is impossible. Thus u is a complete first return to s, so  $u \in \mathcal{P}ri_{00}(n)$ .

Now the claim for the function  $g_1$  follows as  $f_1(w)^{\sim} = g_1(\tilde{w})$  and  $\tilde{\alpha}_1 = \alpha_2$ .

**Lemma 3.16.** Let  $n \ge 1$ . The function  $f_2 : Pri_{00}(4n-2) \to Pri_{1001}(4n)$ ,  $f_2(w) = 1w1$  is a bijection.

*Proof.* If n = 1, then  $\mathcal{P}ri_{00}(2) = \{00\}$  and  $\mathcal{P}ri_{1001}(4) = \{1001\}$ , so the claim holds. The cases n = 2, 3 are also clear:  $\mathcal{P}ri_{00}(6) = \{\alpha_3\}, \mathcal{P}ri_{1001}(8) = \{E(\gamma_1)\}, \mathcal{P}ri_{00}(10) = \{\alpha_4\}$  and  $\mathcal{P}ri_{1001}(12) = \{E(\gamma_2)\}$ ; see Lemma 3.10. Assume that n > 3.

Let  $w \in \mathcal{P}ri_{00}(4n-2)$ . As the factor 00 is always preceded and followed by the letter 1,  $1w1 \in \mathcal{F}(t)$ , and 1w1 begins and ends with 1001. Let v be the longest privileged proper prefix of w. The assumption n > 3 implies that |v| > 4, so by Corollary 3.14 it holds that  $4 \nmid |v|$ . Thus by induction the word 1v1 is privileged. The word 1w1 is a complete first return to 1v1, as otherwise w would contain more than two occurrences of v. Thus  $1w1 \in \mathcal{P}ri_{1001}(4n)$ .

Let then  $1w1 \in Pri_{1001}(4n)$ . Again by applying Corollary 3.14 and induction to the longest privileged proper prefix of 1w1 we get that  $w \in Pri_{00}(4n-2)$ .

**Corollary 3.17.** 
$$\mathcal{A}_{00}(4n) = 2\mathcal{A}_{00}(n)$$
 and  $\mathcal{A}_{00}(4n-2) = \mathcal{A}_{0110}(4n)$  for all  $n \ge 2$ .

*Proof.* As the ranges of the functions  $f_1 : \mathcal{P}ri_{00}(n) \to \mathcal{P}ri_{\alpha_1}(4n)$  and  $g_n : \mathcal{P}ri_{00}(n) \to \mathcal{P}ri_{\alpha_2}$  are disjoint, the claim follows since by Lemma 3.11  $\mathcal{P}ri_{00}(4n) = \mathcal{P}ri_{\alpha_1}(4n) \cup \mathcal{P}ri_{\alpha_2}(4n)$ . The other equality follows directly from Lemma 3.16 as  $\mathcal{A}_{1001}(m) = \mathcal{A}_{0110}(m)$  for all  $m \ge 4$ .

**Lemma 3.18.** Let  $n \ge 2$ . The function  $f_3 : Pri_{101}(n+1) \cup Pri_{1001}(n+1) \rightarrow Pri_{010}(4n), f_3(w) = \partial_{2,2}(\theta(w))$  is a bijection.

*Proof.* If n = 2, then  $\mathcal{P}ri_{101}(3) = \{101\}$ ,  $\mathcal{P}ri_{1001}(3) = \emptyset$  and  $\mathcal{P}ri_{010}(8) = \{\beta_1\}$ . If n = 3, then  $\mathcal{P}ri_{101}(4) = \emptyset$ ,  $\mathcal{P}ri_{1001}(4) = \{1001\}$  and  $\mathcal{P}ri_{0110}(12) = \{\beta_2\}$ . We may assume that n > 3.

Let  $w \in \mathcal{P}ri_{101}(n + 1) \cup \mathcal{P}ri_{1001}(n + 1)$ , and v its longest privileged proper prefix. By induction  $f_3(v) \in \mathcal{P}ri_{010}(t)$ . As v is a prefix and a suffix of w,  $f_3(w)$  starts and ends with  $f_3(v)$ . By Lemma 3.12 the word  $f_3(v)$  is always preceded by 10 and followed by 01. Thus if  $f_3(w)$  contained more than two occurrences of  $f_3(v)$ , then Proposition 3.3 would imply that w contains more than two occurrences of v which would be a contradiction. We conclude that  $f_3(w) \in \mathcal{P}ri_{010}(4n)$ .

Let then  $w \in \mathcal{P}ri_{010}(4n)$ . By Lemma 3.12 there is such a word u that  $f_3(u) = w$ . Let v be the longest privileged proper prefix of w. By the assumption n > 3, we have that |v| > 4. By Corollary 3.14 we can apply induction to obtain a word  $s \in \mathcal{P}ri_{101}(t) \cup \mathcal{P}ri_{1001}(t)$  such that  $f_3(s) = v$ . Therefore  $f_3(u) = w = f_3(s) \cdots f_3(s)$ . By Lemma 3.12  $f_3(s)$  is always preceded by 10 and followed by 01. Therefore u begins and ends with s. Now if u would contain a third occurrence of s, then w would contain a third occurrence of v, which is not possible. Hence  $u \in \mathcal{P}ri_{101}(n+1) \cup \mathcal{P}ri_{1001}(n+1)$ .

**Lemma 3.19.** Let  $n \ge 2$ . The function  $f_4 : Pri_{101}(4n-2) \to Pri_{010}(4n)$ ,  $f_4(w) = 0w0$  is a bijection.

*Proof.* If n = 2, then  $\mathcal{P}ri_{101}(6) = \{E(\beta_3)\}$  and  $\mathcal{P}ri_{010}(8) = \{\beta_1\}$ . If n = 3, then  $\mathcal{P}ri_{101}(10) = \{E(\beta_4)\}$ , and  $\mathcal{P}ri_{010}(12) = \{\beta_2\}$ . Thus it can be assumed that n > 3.

Let  $w \in \mathcal{P}ri_{101}(4n-2)$ , and v its longest privileged proper prefix. As n > 3, we have that |v| > 4. By Corollary 3.14 and induction  $f_4(v) \in \mathcal{P}ri_{010}(t)$ . By Lemma 3.12 the factor  $f_4(v)$  is always preceded and followed by letter 0. Thus it can be written that  $f_4(w) = f_4(v) \cdots f_4(v)$ . If there was a third occurrence of  $f_4(v)$  in  $f_4(w)$ , then in w there would be at least three occurrences of v, which is false. Therefore  $f_4(w) \in \mathcal{P}ri_{010}(4n)$ .

Let  $0w0 \in Pri_{010}(4n)$ . Again, by applying Corollary 3.14 and induction to the longest privileged proper prefix of 0w0, we get that  $w \in Pri_{101}(4n - 2)$ .

**Corollary 3.20.**  $\mathcal{A}_{010}(4n) = \mathcal{A}_{010}(n+1) + \mathcal{A}_{0110}(n+1)$  and  $\mathcal{A}_{010}(4n-2) = \mathcal{A}_{010}(4n)$  for all  $n \ge 2$ .

*Proof.* This follows directly from Lemmas 3.18 and 3.19 as  $\mathcal{A}_{101}(n) = \mathcal{A}_{010}(n)$  and  $\mathcal{A}_{1001}(n) = \mathcal{A}_{0110}(n)$  for all  $n \ge 0$ .

**Lemma 3.21.** Let  $n \ge 2$ . The function  $\theta$  :  $\operatorname{Pri}_{00}(n) \cup \operatorname{Pri}_{010}(n) \to \operatorname{Pri}_{0110}(4n)$  is a bijection.

*Proof.* If n = 2, then  $Pri_{00}(2) = \{00\}$ ,  $Pri_{010}(2) = \emptyset$  and  $Pri_{0110}(8) = \{\gamma_1\}$ . If n = 3, then  $Pri_{00}(3) = \emptyset$ ,  $Pri_{010}(3) = \{010\}$  and  $Pri_{0110}(12) = \{\gamma_2\}$ . By the argument at the end of the proof if  $Pri_{0110}(4n) \neq \emptyset$ , then  $Pri_{00}(n) \cup Pri_{010}(n) \neq \emptyset$ . As  $Pri_{00}(n) \cup Pri_{010}(n) = \emptyset$  when n = 4, it can be further assumed that n > 4.

Let  $w \in Pri_{00}(n) \cup Pri_{010}(n)$ , and v its longest privileged proper prefix. Now  $|v| \ge 2$  as n > 4. Once again by induction  $\theta(v) \in Pri_{0110}(t)$ . By Proposition 3.3 the word  $\theta(w)$  must be a complete first return to  $\theta(v)$ , that is,  $\theta(w) \in Pri_{0110}(4n)$ .

Let  $w \in \mathcal{P}ri_{0110}(4n)$ . By Lemma 3.13 there is  $u \in \mathcal{F}_t(n)$  such that  $\theta(u) = w$ . Again by Corollary 3.14 and induction there exists  $s \in \mathcal{P}ri_{00}(t) \cup \mathcal{P}ri_{010}(t)$  such that  $\theta(s) = v$  where v is the longest privileged proper prefix of w. It follows that u must be a complete first return to s, so  $u \in \mathcal{P}ri_{00}(n) \cup \mathcal{P}ri_{010}(n)$ .

**Lemma 3.22.** Let  $n \ge 2$ . The function  $f_4 : Pri_{11}(4n) \to Pri_{0110}(4n+2)$ ,  $f_4(w) = 0w0$  is a bijection.

*Proof.* If n = 2, then  $\mathcal{P}ri_{11}(8) = \{E(\alpha_1), E(\alpha_2)\}$  and  $\mathcal{P}ri_{0110}(10) = \{\gamma_3, \gamma_4\}$ . If n = 3, then  $\mathcal{P}ri_{11}(12) = \emptyset$  and  $\mathcal{P}ri_{0110}(14) = \emptyset$ . The latter set is empty because if  $w \in \mathcal{P}ri_{0110}(14)$ , then w has as a prefix and a suffix a word u which is a complete first return to 0110. Since  $|u| \ge 8$ , these two occurrences of u in w overlap. This is contradictory.

The rest of the proof is by induction along the lines of the proof of Lemma 3.16.

**Corollary 3.23.**  $\mathcal{A}_{0110}(4n) = \mathcal{A}_{00}(n) + \mathcal{A}_{010}(n)$  and  $\mathcal{A}_{0110}(4n-2) = \mathcal{A}_{00}(4(n-1))$  for  $n \ge 2$ .

*Proof.* The result follows from Lemmas 3.21 and 3.22 as  $\mathcal{A}_{00}(n) = \mathcal{A}_{11}(n)$  for all  $n \ge 0$ .

Proposition 3.9 and Corollaries 3.17, 3.20 and 3.23 together prove Theorem 3.5. In the following we characterize the non-primitive privileged factors:

**Proposition 3.24.** The only non-primitive privileged factors of t beginning with 0 are 00,  $\beta_3$ ,  $\gamma_1$  and  $\gamma_2^2$ .

*Proof.* Let *w* be a non-primitive privileged factor of *t* beginning with 0. Since *t* is overlap-free, it cannot contain third powers. Thus  $w = u^2$  for some privileged factor *u*. If |u| = 1, then w = 00. If |u| > 1, then *u* cannot begin with 00 as otherwise *w* would have  $0^4$  as a central factor. Hence *u* begins with 010 or 0110. If |u| = 3, 4, then  $w = \beta_3, \gamma_1$ . We may assume that |u| > 5. Then as |u| is even, we have that  $4 \mid |w|$ , so  $4 \mid |u|$  by Corollary 3.14. By Lemma 3.12 if *u* begins with 010, then *u* begins with  $\beta_1$  or  $\beta_2$ , and so *w* has  $\beta_1^2$  or  $\beta_2^2$  as a central factor. This is, however, impossible as neither  $\beta_1^2$  nor  $\beta_2^2$  is a factor of *t*. Thus *u* must begin with 0110, and begin with  $\gamma_1$  or  $\gamma_2$ . The word  $\gamma_1$  is non-primitive, and thus  $\gamma_1^2 \notin \mathcal{F}(t)$ . As  $\gamma_2$  is primitive, and  $\gamma_2^2$  is not, we may assume that *u* begins with  $\gamma_2$  and that  $u \neq \gamma_2$ . Thus *w* has  $\gamma_2^2 = \theta(010)^2$  as a central factor. By Lemma 3.21  $w = \theta(v)$  where *v* is a privileged word beginning with 010. As  $(010)^2$  must be preceded by 011 and followed by 110, we have that *w* has  $\theta(011010 \cdot 010110) = \theta^2(010)$  as a central factor. As 010 cannot be preceded and followed by the same letter, we have that *v* ends with  $\beta_1$  and begins with  $\beta_2$ , or symmetrically *v* ends with  $\beta_2$  and begins with  $\beta_1$ . Either case is impossible as *v* is privileged.

Next we study the asymptotic behavior of the function  $\mathcal{A}_t$ .

**Proposition 3.25.**  $\limsup_{n \to \infty} \mathcal{A}_t(n) = \infty$  and  $\liminf_{n \to \infty} \mathcal{A}_t(n) = 0$ .

*Proof.* The fact that the inferior limit is 0 already follows from Proposition 3.9. We will prove that when  $n \ge 6$ ,

$$\mathcal{A}_t(2^n) = \begin{cases} 3 \cdot 2^{\frac{1}{2}(n-1)}, & \text{if } n \text{ is odd,} \\ 0, & \text{if } n \text{ is even,} \end{cases}$$

which proves the claim.

As  $2^{n-2} + 1$  is odd, by Corollary 3.20 it holds for n > 3 that

$$\mathcal{A}_{010}(2^n) = \mathcal{A}_{010}(2^{n-2}+1) + \mathcal{A}_{0110}(2^{n-2}+1) = 0.$$

Thus by Theorem 3.5

$$\frac{1}{2}\mathcal{A}_t(2^n) = 3\mathcal{A}_{00}(2^{n-2}).$$

Now  $\mathcal{A}_{00}(2) = 1$  and  $\mathcal{A}_{00}(4) = 0$ . By Corollary 3.17  $\mathcal{A}_{00}(2^n) = 2\mathcal{A}_{00}(2^{n-2})$ , so for all  $n \ge 1$ ,

$$\mathcal{A}_{00}(2^n) = \begin{cases} 2^{\frac{1}{2}(n-1)}, & \text{if } n \text{ is odd,} \\ 0, & \text{if } n \text{ is even,} \end{cases}$$

proving the desired equality.

This result is very interesting. It can be proven that the palindromic complexity function of a fixed point of a primitive morphism is bounded (see [DZ00; All+03]). So as the Thue-Morse word is a fixed point of a primitive morphism, the above results demonstrate that in general the palindromic complexity function and the privileged complexity function of an infinite word can behave radically differently.

Our next aim is to show that there exist arbitrarily long gaps of zeros in the values of  $\mathcal{A}_t$ . For this we need several lemmas.

Let us define an integer sequence  $(a_n)$  as follows:  $a_1 = 14$  and  $a_n = 4(a_{n-1}-2) + 2(-1)^n$  for n > 1. The first few terms of the sequence are 14, 50, 190, 754, 3006, ... Note that  $a_n$  is always even, and not divisible by 4.

**Lemma 3.26.** If *n* is even, then  $\mathcal{A}_{00}(a_n - 2) = \mathcal{A}_{010}(a_n - 2) = 0$  and  $\mathcal{A}_{0110}(a_n - 2) = 1$ . If n > 1 is odd, then  $\mathcal{A}_{00}(a_n - 2) = \mathcal{A}_{0110}(a_n - 2) = 0$  and  $\mathcal{A}_{010}(a_n - 2) = 1$ . Moreover if n > 1, then  $\mathcal{A}_t(a_n - 2) = 2$ .

*Proof.* Using the formulas of Corollaries 3.17, 3.20 and 3.23 it is readily verified that the claim holds for n = 2.

Let n > 1 be odd. Then  $a_n = 4(a_{n-1} - 2) - 2$ . Using induction, Proposition 3.9 and the formulas of Corollaries 3.17, 3.20 and 3.23 we get that

$$\mathcal{A}_{00}(a_n-2) = 2\mathcal{A}_{00}(a_{n-1}-3) = 0,$$
  

$$\mathcal{A}_{010}(a_n-2) = \mathcal{A}_{010}(a_{n-1}-2) + \mathcal{A}_{0110}(a_{n-1}-2) = \mathcal{A}_{0110}(a_{n-1}-2) = 1 \text{ and}$$
  

$$\mathcal{A}_{0110}(a_n-2) = \mathcal{A}_{00}(a_{n-1}-3) + \mathcal{A}_{010}(a_{n-1}-3) = 0.$$

Let *n* be even, so  $a_n = 4(a_{n-1} - 2) + 2$ . Similarly as above

$$\begin{aligned} \mathcal{A}_{00}(a_n-2) &= 2\mathcal{A}_{00}(a_{n-1}-2) = 0, \\ \mathcal{A}_{010}(a_n-2) &= \mathcal{A}_{010}(a_{n-1}-1) + \mathcal{A}_{0110}(a_{n-1}-1) = 0 \text{ and} \\ \mathcal{A}_{0110}(a_n-2) &= \mathcal{A}_{00}(a_{n-1}-2) + \mathcal{A}_{010}(a_{n-1}-2) = \mathcal{A}_{010}(a_{n-1}-2) = 1. \end{aligned}$$

It clearly follows that  $\mathcal{A}_t(a_n - 2) = 2$  for n > 1.

By Lemma 3.10 it can be verified that  $\mathcal{A}_t(12) = 4$ , so  $\mathcal{A}_t(a_n - 2) \neq 0$  for all  $n \geq 1$ .

**Lemma 3.27.**  $\mathcal{A}_t(2^n + 2) = 2$  for all even  $n \neq 2$ .

*Proof.* We will prove that  $\mathcal{A}_{00}(2^n + 2) = \mathcal{A}_{0110}(2^n + 2) = 0$  and  $\mathcal{A}_{010}(2^n + 2) = 1$ . The claim follows from this. This claim is readily verified when n = 0. Let then n > 2 be even. By the formulas of Corollaries 3.17 and 3.23 we obtain that

$$\mathcal{A}_{00}(2^{n}+2) = \mathcal{A}_{0110}(2^{n}+4) = \mathcal{A}_{00}(2^{n-2}+1) + \mathcal{A}_{010}(2^{n-2}+1) = 0$$

as  $2^{n-2} + 1$  is odd and greater than 3 when n > 2 is even. Similarly using in addition Corollary 3.20 we get by applying induction that

$$\mathcal{A}_{010}(2^n+2) = \mathcal{A}_{010}(2^n+4) = \mathcal{A}_{010}(2^{n-2}+2) + \mathcal{A}_{0110}(2^{n-2}+2) = 1$$
 and  
 $\mathcal{A}_{0110}(2^n+2) = \mathcal{A}_{00}(2^n) = 0$ ,

where the last equality was proven already in the proof of Proposition 3.25.

Finally we have proven enough lemmas in order to prove the following result:

**Proposition 3.28.** For all  $n \ge 1$  if  $a_n - 1 \le k \le 2^{2(n+1)} + 1$ , then  $\mathcal{A}_t(k) = 0$ . Also for all  $n \ge 1$  $\mathcal{A}_t(a_n - 2) \ne 0$  and  $\mathcal{A}_t(2^{2(n+1)} + 2) \ne 0$ .

*Proof.* By inspection it can be verified that indeed if  $a_1 - 1 = 13 \le k \le 17 = 2^{2(1+1)} + 1$ , then  $\mathcal{A}_t(k) = 0$ . Let n > 1. Assume that  $a_n \le k \le 2^{2(n+1)}$ . If k is odd, then  $\mathcal{A}_t(k) = 0$ . Suppose that k is even. Assume that 4 divides k. Then it follows that  $a_{n-1} - 1 \le k/4 \le 2^{2n}$ , so by the induction hypothesis  $\mathcal{A}_t(k/4) = 0$ . By Theorem 3.5

$$\frac{1}{2}\mathcal{A}_t(k) = 3\mathcal{A}_{00}(k/4) + \mathcal{A}_{010}(k/4) + \mathcal{A}_{010}(k/4+1) + \mathcal{A}_{0110}(k/4+1) = 0$$

Assume that 4 does not divide *k*. Now  $a_{n-1} - 1 \le \frac{k+2}{4} \le 2^{2n}$  and  $a_{n-1} - 2 \le \frac{k-2}{4} \le 2^{2n}$ . Now using the already familiar formulas we get

$$\begin{split} \frac{1}{2}\mathcal{A}_{t}(k) &= \mathcal{A}_{00}(k-2) + \mathcal{A}_{010}(k+2) + \mathcal{A}_{0110}(k+2) \\ &= 2\mathcal{A}_{00}\left(\frac{k-2}{4}\right) + \mathcal{A}_{010}\left(\frac{k+2}{4}+1\right) + \mathcal{A}_{0110}\left(\frac{k+2}{4}+1\right) + \mathcal{A}_{00}\left(\frac{k+2}{4}\right) + \mathcal{A}_{010}\left(\frac{k+2}{4}\right) \\ &= 2\mathcal{A}_{00}\left(\frac{k-2}{4}\right), \end{split}$$

where the last equality follows from the induction hypothesis. Now if  $k \neq a_n$ , then  $a_{n-1} - 1 \leq \frac{k-2}{4}$ , so by the induction hypothesis  $\mathcal{A}_t(k) = 0$ . If  $k = a_n$ , then  $\frac{k-2}{4} = a_{n-1} - 2$ . Then however by Lemma 3.26  $\mathcal{A}_{00}(a_{n-1} - 2) = 0$ , so also  $\mathcal{A}_t(k) = 0$ .

The claim now follows as  $a_n - 1$  and  $2^{2(n+1)} + 1$  are odd. Earlier it was proved that  $\mathcal{A}_t(a_n - 2)$ ,  $\mathcal{A}_t(2^{2(n+1)} + 2) \neq 0$ .

Straightforwardly using induction it can be proved that for  $n \ge 3$  it holds that  $a_n < 2^{2n+1} + 2^{2n} < 2^{2(n+1)}$ , so in particular if for  $n \ge 3$  it holds that  $2^{2n+1} + 2^{2n} \le k \le 2^{2(n+1)}$ , then  $\mathcal{A}_t(k) = 0$ . This verifies the following result which was conjectured in [Pel13]:

**Corollary 3.29.** There exists arbitrarily long (but not infinite) gaps of zeros in the values of the privileged complexity function of the Thue-Morse word.  $\Box$ 

Corollary 3.29 raises a natural question: If the privileged complexity function of a word w contains arbitrarily large gaps of zeros, does it follow that  $\limsup_{n\to\infty} \mathcal{A}_w(n) = \infty$ ? It is conceivable that the large gaps force large values of  $\mathcal{A}_w(n)$  between the gaps. On the other hand the gaps could occur so sparsely that  $\mathcal{A}_w(n)$  is still bounded. The author was not able to answer this question.

The privileged complexity function of the Thue-Morse word is complicated. Even though the Thue-Morse morphism has really nice properties, finding the recursive formula for the function is a long task. On the other hand without the nice properties of the morphism, the work may not have been possible at all. Indeed if the morphism were not uniform, then it would have been harder to calculate the length of the privileged factors. Other crucial property of the morphism is its *circularity*: every image of a letter is uniquely determined by its first or last letter. The author thinks that it could be possible to obtain results on the privileged complexity of fixed points of primitive uniform circular morphisms other than Thue-Morse.

## 4 Privileged Palindrome Complexity

In this section we will focus on the privileged palindrome complexity function of the Thue-Morse word *t*. This function  $\mathcal{B}_t$  counts the number of *n*-length factors of *t* which are both privileged and palindromic. The arguments given are similar to those of Section 3.

Similar to Section 3 we write  $\mathcal{M}_u(n) = \mathcal{P}ri_t(n) \cap \mathcal{P}al_t(n) \cap u \cdot \{0,1\}^*$ , and  $\mathcal{B}_u(n) = |\mathcal{M}_u(n)|$ . Again it suffices to consider factors beginning with letter 0.

**Lemma 4.1.** Let  $w \in Pal(t)$ . Then  $w \in Pal_t(4n)$  for some  $n \ge 1$  if and only if w is  $\varphi$ -invertible (i.e., there exists a word  $u \in \mathcal{F}(t)$  such that  $\varphi(u) = w$ ).

*Proof.* The claim holds for all palindromes of length less than or equal to 4, they are: 0, 1, 00, 11, 010, 101, 0110 and 1001. Suppose that  $n \ge 2$ .

Let  $w \in \mathcal{Pal}_t(4n)$  be a shortest palindrome that is not  $\varphi$ -invertible. Suppose first that w begins with 00, that is, w = 001w'100. Now  $1w'1 \in \mathcal{Pal}(4(n-1))$ , so by the minimality of |w| the word 1w'1 is  $\varphi$ -invertible. As w begins with 00, the word 1w'1 must have two interpretations by  $\varphi$ . This is a contradiction with Lemma 3.2. Say w begins with 01 (the case where it begins with 10 is symmetric), so it can be written that w = 01w'10. As w is not  $\varphi$ -invertible neither is w' (otherwise w would have two interpretations by  $\varphi$ ) which is a contradiction with the minimality of |w|.

Suppose then that w is the shortest  $\varphi$ -invertible palindrome such that  $4 \nmid |w|$ . It can be written that w = 01w'10 (or symmetrically w = 10w'01), so w' is a palindrome of length |w| - 4 which is  $\varphi$ -invertible. This contradicts the choice of w.

Let us have a closer look at the function  $\theta$  and the functions

$$f_{2}: w \mapsto 1w1$$
  
$$f_{3}: w \mapsto \partial_{2,2}(\theta(w))$$
  
$$f_{4}: w \mapsto 0w0$$

defined in Lemmas 3.16, 3.18 and 3.19. Obviously all functions preserve palindromes. By Lemmas 3.16, 3.18, 3.19, 3.21 and 3.22 the functions  $f_2$ ,  $f_3$ ,  $f_4$  and  $\theta$  preserve privileged words. Hence we have that the following functions are bijections:

$$\begin{split} f_{2} &: \mathcal{M}_{00}(4n-2) \to \mathcal{M}_{1001}(4n), w \mapsto 1w1, \\ f_{3} &: \mathcal{M}_{101}(n+1) \cup \mathcal{M}_{1001}(n+1) \to \mathcal{M}_{010}(4n), w \mapsto \partial_{2,2}(\theta(w)), \\ f_{4} &: \mathcal{M}_{101}(4n-2) \to \mathcal{M}_{010}(4n), w \mapsto 0w0, \\ f_{4} &: \mathcal{M}_{11}(4n) \to \mathcal{M}_{0110}(4n+2), w \mapsto 0w0, \\ \theta &: \mathcal{M}_{00}(n) \cup \mathcal{M}_{010}(n) \to \mathcal{M}_{0110}(4n), w \mapsto \theta(w). \end{split}$$

We have thus proved the following formulas:

$$\begin{aligned} \mathcal{B}_{00}(4n-2) &= \mathcal{B}_{0110}(4n),\\ \mathcal{B}_{010}(4n-2) &= \mathcal{B}_{010}(4n),\\ \mathcal{B}_{0110}(4n-2) &= \mathcal{B}_{00}(4(n-1)),\\ \mathcal{B}_{010}(4n) &= \mathcal{B}_{010}(n+1) + \mathcal{B}_{0110}(n+1),\\ \mathcal{B}_{0110}(4n) &= \mathcal{B}_{00}(n) + \mathcal{B}_{010}(n), \end{aligned}$$

for  $n \ge 2$ . We are still missing a formula for  $\mathcal{B}_{00}(4n)$ . However  $\mathcal{M}_{00}(4n) = \emptyset$  by Lemma 4.1, so  $\mathcal{B}_{00}(4n) = 0$ . By putting together these formulas we get the following result:

**Theorem 4.2.** The privileged palindrome complexity function  $\mathcal{B}_t$  of the Thue-Morse word satisfies

$$\begin{aligned} &\mathcal{B}_{t}(0) = 1, \,\mathcal{B}_{t}(1) = \mathcal{B}_{t}(2) = \mathcal{B}_{t}(3) = \mathcal{B}_{t}(4) = 2, \\ &\frac{1}{2}\mathcal{B}_{t}(4n) = \mathcal{B}_{00}(n) + \mathcal{B}_{010}(n) + \mathcal{B}_{010}(n+1) + \mathcal{B}_{0110}(n+1) & \text{for } n \ge 2, \\ &\mathcal{B}_{t}(4n-2) = \mathcal{B}_{t}(4n) & \text{for } n \ge 2, \\ &\mathcal{B}_{t}(2n+1) = 0 & \text{for } n \ge 2. \end{aligned}$$

As in the previous section for the function  $\mathcal{A}_t$ , we study next the asymptotic behavior of the function  $\mathcal{B}_t$  and the occurrences of zeros in its values.

Let us define an integer sequence  $(b_n)$  as follows:  $b_1 = 6$  and  $b_n = 4b_{n-1} - 2$  for n > 1. The first few terms of the sequence are 6, 22, 86, 342, 1366, ... Note that  $b_n$  is always even, and not divisible by 4.

## **Lemma 4.3.** $\mathcal{B}_t(b_n) = 4$ for all $n \ge 1$ .

*Proof.* By inspection  $\mathcal{B}_{00}(6) = 1$ ,  $\mathcal{B}_{010}(6) = 1$  and  $\mathcal{B}_{0110}(6) = 0$ , so  $\mathcal{B}(6) = 4$ . We will prove that  $\mathcal{B}_{00}(b_n) = 2$  and  $\mathcal{B}_{010}(b_n) = \mathcal{B}_{0110}(b_n) = 0$  for all n > 1. The claim follows from this. Now

$$\mathcal{B}_{00}(b_n) = \mathcal{B}_{0110}(b_n + 2) = \mathcal{B}_{00}(b_{n-1}) + \mathcal{B}_{010}(b_{n-1})$$
  
$$\mathcal{B}_{010}(b_n) = \mathcal{B}_{010}(b_n + 2) = \mathcal{B}_{010}(b_{n-1} + 1) + \mathcal{B}_{0110}(b_{n-1} + 1),$$
  
$$\mathcal{B}_{0110}(b_n) = \mathcal{B}_{00}(b_n - 2) = \mathcal{B}_{00}(b_{n-1} - 1).$$

So the claim is indeed true.

**Proposition 4.4.** The function  $\mathcal{B}_t$  takes values in  $\{0, 1, 2, 4\}$ , and the values 0, 2 and 4 are attained infinitely often.

*Proof.* As  $\mathcal{B}_t(0) = 1$ ,  $\mathcal{B}_t(1) = \mathcal{B}_t(2) = \mathcal{B}_t(3) = \mathcal{B}_t(4) = 2$ , by Theorem 4.2 we need only to consider the values  $\mathcal{B}_t(4n)$ . If n = 2, then  $\mathcal{B}_t(8) = 4$ . Let n > 4 be even. Then by Theorem 4.2

$$\frac{1}{2}\mathcal{B}_t(4n) = \mathcal{B}_{00}(n) + \mathcal{B}_{010}(n).$$

It suffices to prove that if  $\mathcal{B}_{00}(n) \neq 0$  then  $\mathcal{B}_{010}(n) = 0$ . We are only interested in the case where n is not divisible by 4, as otherwise  $\mathcal{B}_{00}(n) = 0$ . Now

$$\mathcal{B}_{00}(n) = \mathcal{B}_{0110}(n+2) = \mathcal{B}_{00}\left(\frac{n+2}{4}\right) + \mathcal{B}_{010}\left(\frac{n+2}{4}\right) \text{ and}$$
$$\mathcal{B}_{010}(n) = \mathcal{B}_{010}(n+2) = \mathcal{B}_{010}\left(\frac{n+2}{4}+1\right) + \mathcal{B}_{0110}\left(\frac{n+2}{4}+1\right).$$

Clearly if  $\mathcal{B}_{00}(n) \neq 0$ , then (n+2)/4 is even, and thus  $\mathcal{B}_{010}(n) = 0$ .

Let *n* be odd. Then

$$\frac{1}{2}\mathcal{B}_t(4n) = \mathcal{B}_{010}(n+1) + \mathcal{B}_{0110}(n+1).$$

Again it suffices to prove that if  $\mathcal{B}_{010}(n) \neq 0$ , then  $\mathcal{B}_{0110}(n) = 0$ . As  $\mathcal{B}_{0110}(n) = 0$ , when *n* is not divisible by 4, we need to consider only the case where *n* is divisible by 4. Now

$$\mathcal{B}_{010}(n) = \mathcal{B}_{010}\left(\frac{n}{4}+1\right) + \mathcal{B}_{0110}\left(\frac{n}{4}+1\right) \text{ and}$$
$$\mathcal{B}_{0110}(n) = \mathcal{B}_{00}\left(\frac{n}{4}\right) + \mathcal{B}_{010}\left(\frac{n}{4}\right).$$

Obviously if  $\mathcal{B}_{010}(n) \neq 0$ , then n/4 is odd, and thus  $\mathcal{B}_{0110}(n) = 0$ .

By Lemma 4.3 the function  $\mathcal{B}_t$  takes value 4 infinitely often. Moreover the arguments of Lemmas 3.26 and 3.27 work if the function  $\mathcal{A}_t$  is replaced with the function  $\mathcal{B}_t$ . Thus the value 2 is also attained infinitely often.

Let us now consider the gaps of zeros in the values of  $\mathcal{B}_t$  like we did for the function  $\mathcal{A}_t$  in Proposition 3.28. It is clear by Proposition 3.28 that if  $a_n - 1 \le k \le 2^{2(n+1)} + 1$ , then  $\mathcal{B}_t(k) = 0$ . The arguments of Lemmas 3.26 and 3.27 work if the function  $\mathcal{A}_t$  is replaced with the function  $\mathcal{B}_t$ (in the proof of the latter lemma, the last equality follows now more easily as  $\mathcal{B}_{00}(4n) = 0$  for all  $n \ge 1$ ), so  $\mathcal{B}_t(a_n - 2)$ ,  $\mathcal{B}_t(2^{2(n+1)} + 2) \ne 0$  for all  $n \ge 1$ . Therefore the function  $\mathcal{B}_t$  has the same gaps as the function  $\mathcal{A}_t$  described by Proposition 3.28, the gaps do not widen.

#### 5 A Comparison of Palindromes and Privileged Words

In this section we compare the behavior of palindromes and privileged words and the behavior of the respective complexity functions in general.

Let *w* be an infinite word. In [Pel13] and [KLS13] the following relation between the sets  $\mathcal{P}al(w)$  and  $\mathcal{P}ri(w)$  was proved:

**Proposition 5.1.** [Pel13; KLS13] A finite or infinite word w is rich if and only if Pri(w) = Pal(w).

First we strengthen this result a bit. Now if a word is rich, then from the previous proposition it obviously follows that the palindromic and privileged complexity functions of w coincide. Next we prove the surprising fact that the converse is also true. We start with a lemma which is interesting in its own right.

**Lemma 5.2.** Let w be a finite or infinite word. If w is not rich, then there exists a shortest privileged factor u which is not a palindrome. Moreover  $\operatorname{Pal}_w(n) = \operatorname{Pri}_w(n)$  for all n such that  $0 \le n < |u|$  and  $\operatorname{Pal}_w(|u|) \subsetneq \operatorname{Pri}_w(|u|)$ .

*Proof.* If *w* is not rich, then there exists a position *n* such that no new palindrome in position *n* is introduced. However position *n* introduces a new privileged factor, which thus cannot be a palindrome. Hence there exists a shortest privileged factor *u* which is not a palindrome. By the minimality of |u| it follows that  $Pri_w(n) \subseteq Pal_w(n)$  for all *n* such that  $0 \le n < |u|$ . Let p, |p| > 1, be a minimal length palindrome which is not privileged. Let *q* be the longest proper palindromic

suffix of *p*. By minimality *q* is privileged. As *p* is not privileged, it has as a suffix a complete first return to *q*, say *v*. As *q* is the longest palindromic suffix of *p*, *v* is not a palindrome. By minimality of |u| we have that  $|p| > |v| \ge |u|$ , so  $\mathcal{Pal}_w(n) \subseteq \mathcal{Pri}_w(n)$  for all *n* such that  $0 \le n \le |u|$ .

**Theorem 5.3.** A finite or infinite word w is rich if and only if  $\mathcal{P}_w(n) = \mathcal{A}_w(n)$  for all n such that  $0 \le n \le |w|$ .

*Proof.* The fact that the condition is necessary follows from Proposition 5.1. Assume that  $\mathcal{P}_w(n) = \mathcal{A}_w(n)$  for all n such that  $0 \le n \le |w|$ . If w is not rich, then by Lemma 5.2 there exists such n that  $\mathcal{P}al_w(n) \subsetneq \mathcal{P}ri_w(n)$ , so  $\mathcal{P}_w(n) < \mathcal{A}_w(n)$ , which is a contradiction. Therefore w is rich.

For instance, the Thue-Morse word has, as factors, the word 00101100 which is privileged and not palindromic, and the palindrome 00101100110100 which is not privileged. Thus for a word w its possible that neither of the sets Pal(w) and Pri(w) is included in the other. By Lemma 5.2 it follows that if  $Pri(w) \subseteq Pal(w)$ , then Pri(w) = Pal(w), i.e., w is rich. Next it is natural to ask if there are examples of infinite words w such that Pal(w) is properly contained in Pri(w). It turns out that this is possible, but not in the case of uniformly recurrent words containing infinitely many palindromes. We begin with a simple observation.

#### **Lemma 5.4.** Let w be a recurrent infinite word. If $Pal(w) \subsetneq Pri(w)$ , then w has infinite defect.

*Proof.* As  $\mathcal{P}al(w) \subsetneq \mathcal{P}ri(w)$  there exists a privileged factor u which is not a palindrome. Consider any factor v which is a complete first return to u in  $\mathcal{F}(w)$  (such a factor exists as w is recurrent). Let z be a prefix of w having v as a suffix. Let p be the longest palindromic suffix of z. By the assumption  $\mathcal{P}al(w) \subsetneq \mathcal{P}ri(w)$  the palindrome p is also privileged. If |p| > |u|, then p has  $\tilde{u}$  as as a prefix. Since p has a privileged suffix u, it also has u as a prefix, so  $\tilde{u} = u$ , which is impossible. Thus |p| < |u|, so p is actually the longest palindromic suffix of u. As z contains two occurrences of u, it follows that the longest palindromic suffix of z is not unioccurrent. As w is recurrent, there are infinitely many prefixes of w having v as a suffix. Thus w has infinite defect.

Next we define an infinite binary word  $\kappa = \lim_{n\to\infty} u_n$  as the limit of the sequence  $u_0 = 00101100$ ,  $u_{n+1} = u_n 0^n u_n$ . It is clear that  $\kappa$  is recurrent and aperiodic, and contains infinitely many palindromes of the form  $0^n$ . The word  $\kappa$  is, however, not closed under reversal as  $(1011)^{\sim} = 1101$  is not a factor. We claim that  $\mathcal{P}al_{\kappa}(n) = \{0^n, 10^{n-2}1\}$  for  $n \ge 7$ . Let  $p \in \mathcal{P}al_{\kappa}(n)$  for  $n \ge 7$ , and m be minimal such that p occurs in  $u_m$ . As  $u_{m-1}$  starts and ends with 00101100, and 1101 is not a factor of  $\kappa$ , we conclude that p must be a central factor of  $u_m$ . There are thus only two possibilities,  $0^m$  or  $10^{m-2}1$ . By direct inspection the reader can verify that  $\mathcal{P}al_w(6) = \{0^6\}$ ,  $\mathcal{P}al_w(5) = \{0^5\}$ ,  $\mathcal{P}al_w(4) = \{0000, 0110\}$  and  $\mathcal{P}al_w(3) = \{000, 010, 101\}$ . Thus we have proved the following:

**Lemma 5.5.** There exists an infinite recurrent aperiodic binary word w having the following properties: w is not closed under reversal, w contains infinitely many palindromes and  $Pal(w) \subsetneq Pri(w)$ .

Now let us consider the *Chacon word*  $\lambda$ , the fixed point of the (non-primitive) morphism  $0 \mapsto 0010, 1 \mapsto 1$  [Fer95]. The word  $\lambda$  is aperiodic, and uniformly recurrent, as the letter 0 occurs in bounded gaps. By a direct verification one can show that the word  $\lambda$  does not contain palindromes of length 13 or 14. Therefore  $\mathcal{Pal}_{\lambda}(n) = \emptyset$  for all  $n \ge 13$ . There are total 23 palindromes in  $\lambda$ . Using the same brute-force approach one can show that all palindromes in  $\lambda$  are privileged. The Chacon word is not closed under reversal: for instance  $(100100)^{\sim} = 001001 \notin \mathcal{F}(\lambda)$  as 001001 cannot be properly factored over the set {0010, 1}. We have:

**Lemma 5.6.** There exists an infinite uniformly recurrent aperiodic binary word w having the following properties: w is not closed under reversal, w contains finitely many palindromes and  $Pal(w) \subsetneq Pri(w)$ .

Next we recall a construction of [Ber+09]. Consider the infinite word  $\mu = \lim_{n\to\infty} u_n$ , the limit of the sequence  $u_0 = 01$ ,  $u_{n+1} = u_n 23\tilde{u}_n$ . The word  $\mu$  is uniformly recurrent, aperiodic, closed

under reversal and contains only finitely many palindromes, namely only the letters 0, 1, 2 and 3. By applying the morphism

$$h: \begin{array}{c} 0 \mapsto 101, \\ 1 \mapsto 1001, \\ 2 \mapsto 10001, \\ 3 \mapsto 100001 \end{array}$$

to the word  $\mu$  the authors obtain a uniformly recurrent aperiodic binary word which is closed under reversal and contains only finitely many palindromes (the longest is of length 12). By direct inspection it can be verified that each palindrome in  $h(\mu)$  is privileged. Hence we have:

**Lemma 5.7.** There exists an infinite uniformly recurrent aperiodic binary word w having the following properties: w is closed under reversal, w contains finitely many palindromes and  $Pal(w) \subsetneq Pri(w)$ .

However it turns out that if a uniformly recurrent word contains infinitely many palindromes, then the inclusion can not be proper. Note that such a word is necessarily closed under reversal.

**Proposition 5.8.** Let w be a uniformly recurrent word containing infinitely many palindromes. If  $Pal(w) \subseteq Pri(w)$ , then Pal(w) = Pri(w), that is, w is rich.

*Proof.* Assume on the contrary that  $\mathcal{Pal}(w) \subseteq \mathcal{Pri}(w)$  and that w is not rich. Then there exists a privileged factor u which is not a palindrome. Since w is uniformly recurrent u is a factor of some palindrome p. Clearly u cannot be a central factor of p. Thus there exists a central factor q of p which begins with u and ends with  $\tilde{u}$  (or q begins with  $\tilde{u}$  and ends with u, but this case is symmetric). It is immediate that q is a palindrome. Thus by the assumption q is privileged. As q has as a prefix the privileged word u, the prefix u also occurs as a suffix of q. Hence u is a palindrome, a contradiction.

Note that in the proof uniform recurrence was only needed to establish that *u* is a factor of some palindrome. Thus to obtain the result it is only necessary to suppose that every privileged factor occurs in some palindrome.

#### 6 Acknowledgments

The author was supported by a FiDiPro grant (137991) from the Academy of Finland.

We thank the referees for carefully reading the manuscript. The comments greatly improved the presentation.

## References

- [All+03] J.-P. Allouche, M. Baake, J. Cassaigne, and D. Damanik. Palindrome Complexity. *Theoret. Comput. Sci.* 292 (2003), 9–31.
- [BPS07] L. Balková, E. Pelantová, and W. Steiner. Return Words in Fixed Points of Substitutions. Preprint arXiv:math/0608603v3. 2007.
- [Ber+09] J. Berstel, L. Boasson, O. Carton, and I. Fagnot. Infinite Words without Palindrome. Preprint arXiv:0903.2382. 2009.
- [BBL08] A. Blondin-Massé, S. Brlek, and S. Labbé. Palindromic Lacunas of the Thue-Morse Word. GAS-Com 2008. 6th International Conference on Random Generation of Combinatorial Structures. 2008, pp. 53–67.
- [DZ00] D. Damanik and D. Zare. Palindrome Complexity Bounds for Primitive Substitution Sequences. Discrete Math. 222 (2000), 259–261.
- [DJP01] X. Droubay, J. Justin, and G. Pirillo. Episturmian Words and Some Constructions of de Luca and Rauzy. *Theoret. Comput. Sci.* 225 (2001), 539–553.

- [Fer95] S. Ferenczi. Les transformations de Chacon: combinatoire, structure géométrique, lien avec les systèmes de complexité 2n+1. *Bull. Soc. Math. France* 123 (1995), 271–292.
- [FPZ13] A. E. Frid, S. Puzynina, and L. Q. Zamboni. On Palindromic Factorization of Words. *Adv. in Appl. Math.* 50 (2013), 737–748.
- [Gle+09] A. Glen, J. Justin, S. Widmer, and L. Q. Zamboni. Palindromic Richness. *Eur. J. Combin.* 30 (2009), 510–531.
- [GMS13] D. Goč, H. Mousavi, and J. Shallit. On the Number of Unbordered Factors. Language and Automata Theory and Applications. 7th International Conference, LATA 2013. Ed. by A.-H. Dediu, C. Martín-Vide, and B. Truthe. Lecture Notes in Computer Science 7810. Springer, 2013, pp. 299–310.
- [KLS13] J. Kellendonk, D. Lenz, and J. Savinien. A Characterization of Subshifts with Bounded Powers. Discrete Math. 313 (2013), 2881–2894.
- [Lot83] M. Lothaire. Combinatorics on Words. Encyclopedia of Mathematics and Its Applications 17. Addison-Wesley, 1983.
- [Pel13] J. Peltomäki. Introducing Privileged Words: Privileged Complexity of Sturmian Words. *Theoret. Comput. Sci.* 500 (2013), 57–67.