

# Stringlike structures in Kerr-Schild geometry: N=2 string, twistors and Calabi-Yau twofold

Alexander Burinskii,  
Theor. Phys. Lab., NSI, Russian Academy of Sciences,  
B. Tuskaya 52, Moscow 115191 Russia

June 4, 2018

## Abstract

The four-dimensional Kerr-Schild geometry contains two stringy structures. The first one is the closed string formed by the Kerr singular ring, and the second one is an open complex string with was obtained in the complex structure of the Kerr-Schild geometry. The real and complex Kerr strings form together a membrane source of the over-rotating Kerr-Newman solution without horizon,  $a = J/m \gg m$ . It has also been obtained recently that the principal null congruence of the Kerr geometry, induced by the complex Kerr string, is determined by the Kerr theorem as a quartic in the projective twistor space, which corresponds to embedding of the Calabi-Yau twofold in the bulk of the Kerr geometry. In this paper we describe this embedding in details and show that the four folds of the twistorial K3 surface represent an analytic extension of the Kerr congruence created by antipodal involution.

Key-words: Kerr-Schild geometry, complex shift, Kerr theorem, twistors, K3 surface, N=2 superstring

## 1 Introduction

It is now commonly accepted that black holes (BH) are to be associated with elementary particles. Physics of black holes is based on the complex analyticity and conformal field theory, which unites them with superstring theory and particle physics. In the previous paper [1] (hereafter referred as I), we considered emergence of the string-like structures in the four-dimensional Kerr-Schild

geometry without horizons. In the dimensionless units  $G = c = \hbar = 1$ , this happens when the Kerr rotational parameter  $a = J/m$  exceeds the mass parameter  $m$ . This case of the over-rotating Kerr geometry is very important for application to elementary particles. So far as masses of the elementary particles are very small, and the spin may be extreme high, one obtains that for spinning particles  $a$  exceeds  $m$  for many orders. In particular, mass of an electron in the dimensionless units is  $m \sim 10^{-22}$ , while spin  $J = \hbar/2 = 1/2$ . It yields  $a = J/m \sim 10^{22}$ , and therefore, the value  $a$  exceeds  $m$  for about 44 orders.

Principal peculiarity of the over-rotating Kerr geometry is presence of the naked Kerr singular ring, which is branch line of the Kerr space into two sheets, and therefore, it forms a stringlike topological defect of the space-time. The singular metric should be regularized to an almost flat space-time to get a correspondence with quantum theory and the experimentally negligible role of gravity in particle physics. This procedure was specified step by step by many authors during about four decades, and resulted in formation of the smooth and regular source of the Kerr-Newman solution in the form of a relativistically rotating and highly oblate vacuum bubble. Structure of this source and the corresponding references were discussed in I and [2].

The region near the Kerr singular ring is excised and replaced by a vacuum bubble, while the Kerr-Newman electromagnetic field is regularized and gets a finite maximal value at the edge rim of the oblate bubble, forming a stringlike loop around the bubble. The very old proposal that the Kerr singular ring plays the role of a closed string [3] was later supported by the systematic investigations of the singular string solutions of the low energy string theory as solitons [4]. It has been shown in [5] that the structure of fields around the Kerr singular ring is similar to structure of the fundamental string solutions obtained by Sen [6, 7].

One more stringy structure was obtained in the complex representation of the Kerr geometry, in which the Kerr solution is represented as a retarded-time field from the particle propagating along a complex world line [8, 9, 10]. However, the complex world line is really a world sheet [11], and therefore, it is equivalent to a string [12, 13]. Complex world sheet of his string is embedded in the real Kerr geometry, and as it discussed in I, the real and complex Kerr strings form together a membrane, which is analogous of the enhancon model of the string/M-theory unification, [14]. Recently, the both Kerr strings were mentioned in [15] by Adamo and Newman, and they reaction is worth quoting:

“...It would have been a cruel god to have layed down such a pretty scheme and not have it mean something deep.”

In addition to this stringy system, one more remarkable fact was obtained in the paper I, emergence of the Calabi-Yau twofold (K3 surface) in the projective twistor space of the Kerr geometry. It appears as a quartic on the projective twistor space  $CP^3$ , and determines the Kerr Principal Null Congruence (PNC) in accordance with the Kerr theorem. In paper I we arrive at the conclusion, that this parallelism is not accidental, and there may be a fundamental underlying structure lying beyond these relationships. In particular, it was supposed that this correspondence may be related with a mysterious complex N=2 string, which has critical real dimension four [16] and is close related with twistors [11, 17]. In this paper we extend and specify the given in I treatment of the Calabi-Yau space K3 embedded into four-dimensional Kerr-Schild (KS) geometry and give extra arguments in favor of the assumption that the source of the striking parallelism with superstring theory is the N=2 critical superstring, structure of which is consistent with twistorial structure of KS geometry. We work in the Kerr-Schild formalism [18], which is parallel to the Newman-Penrose formalism, but has the advantage, related with action of the Kerr theorem, which allows to describe the geodesic and shear-free congruences (GSF) on the auxiliary Minkowski background in terms of twistors.

## 2 Basics of the Kerr-Schild formalism

**Structure of the Kerr-Newman solution.** KN metric is represented in the Kerr-Schild (KS) form [18],

$$g_{\mu\nu} = \eta_{\mu\nu} + 2he^3_\mu e^3_\nu, \quad (1)$$

where  $\eta_{\mu\nu}$  is auxiliary Minkowski background in Cartesian coordinates  $x = x^\mu = (t, x, y, z)$  with signature  $(-+++)$ , and

$$h = P^2 \frac{mr - e^2/2}{r^2 + a^2 \cos^2 \theta}, \quad P = (1 + Y\bar{Y})/\sqrt{2}, \quad (2)$$

and  $e^3(x)$  is a tangent direction to a *Principal Null Congruence (PNC)*, which is determined by the form

$$e^3_\mu dx^\mu = du + \bar{Y}d\zeta + Yd\bar{\zeta} - Y\bar{Y}dv, \quad (3)$$

where

$$\zeta = (x + iy)/\sqrt{2}, \quad \bar{\zeta} = (x - iy)/\sqrt{2}, \quad u = (z + t)/\sqrt{2}, \quad v = (z - t)/\sqrt{2} \quad (4)$$

are the null Cartesian coordinates, and function  $Y(x^\mu)$  is obtained from *the Kerr theorem*, [19, 20, 21, 9, 22, 23, 10]. Kerr's oblate spheroidal coordinates  $r, \theta, \phi$  are related to Cartesian coordinates as follows

$$x + iy = (r + ia)e^{i\phi} \sin \theta, \quad z = r \cos \theta, \quad \rho = t - r. \quad (5)$$

The Kerr singular ring corresponds to caustic of PNC at  $r = \cos \theta = 0$ . One sees that the metric (1) and electromagnetic potential of the the KN solution,

$$A_\mu = -P^{-2} Re \frac{e}{(r + ia \cos \theta)} e_\mu^3, \quad (6)$$

are aligned with directions  $e_\mu^3$  of the Kerr PNC. By regularization of the KN solution, the Kerr singular ring is excised, forming a vacuum bubble, boundary of which is determined by the condition  $h = 0$  corresponding to Minkowski metric inside the bubble and at the boundary. From (2) one sees that this boundary corresponds to  $r = r_e = e^2/2m$ , and (5) shows that it forms a highly oblate ellipsoidal membrane. The KN electromagnetic field is concentrated at the sharp edge of the membrane and regularized by the cut-off at  $r = r_e$ .

**Kerr theorem.** Kerr theorem determines the shear free null congruences [18, 19] with tangent direction (3) via a complex function  $Y(x^\mu)$ , which is solution of the equation

$$F(T^A) = 0, \quad (7)$$

where  $F(T^A)$  is any holomorphic function of the projective twistor coordinates

$$T^A = \{Y, \quad \lambda^1 = \zeta - Yv, \quad \lambda^2 = u + Y\bar{\zeta}\}, \quad A = 1, 2, 3. \quad (8)$$

Using the Cartesian coordinates  $x^\mu$ , one can rearrange coordinates and reduce function  $F(T^A)$  to the form  $F(Y, x^\mu)$ , which allows one to get explicit solution of the equation (7) in the form  $Y(x^\mu)$ .

For the Kerr and KN solutions function  $F(Y, x^\mu)$  is quadratic in  $Y$ ,

$$F = A(x^\mu)Y^2 + B(x^\mu)Y + C(x^\mu), \quad (9)$$

and the equation (7) represents a *quadric* in the projective twistor space  $\mathbf{CP}^3$ . If determinant  $\Delta = (B^2 - 4AC)$  is non-degenerate, it corresponds to the Kerr complex radial distance [9, 10]

$$\tilde{r} \equiv r + ia \cos \theta = -\Delta^{1/2}, \quad (10)$$

and the solutions of (7),

$$Y^\pm(x^\mu) = (-B \mp \tilde{r})/2A, \quad (11)$$

determine two different Kerr's congruences via expression (3). Another expression for the complex radial distance,

$$\tilde{r} = -dF/dY, \quad (12)$$

may be obtained from the relations (9) and (11). It shows that the Kerr singular ring  $\tilde{r} = 0$  is caustic of the Kerr congruence,  $dF/dY = 0$ .

As a consequence of the Vieta's formulas, the quadratic in  $Y$  function (9) may also be expressed via the solutions  $Y^\pm(x^\mu)$  in the form

$$F(Y, x^\mu) = A(Y - Y^+(x^\mu))(Y - Y^-(x^\mu)). \quad (13)$$




Figure 1: Kerr's congruence of twistor null lines (PNC) is focused on the Kerr singular ring, forming a branch line of the KS space into two sheets.

**Stationary KS geometries** The Kerr and KN solutions are stationary solutions of the Einstein-Maxwell field equations. There are also more general stationary KS solutions for which

$$A_\mu = -P^{-2} \text{Re} \frac{\psi(Y)}{(r + ia \cos \theta)} e_\mu^3 \quad (14)$$

and

$$h = P^2 \frac{mr - |\psi(Y)|^2/2}{r^2 + a^2 \cos^2 \theta}, \quad (15)$$

where  $\psi(Y)$  is arbitrary analytic function. The poles of  $\psi(Y)$  correspond to singular beams supported by twistor null lines of the Kerr congruence, [24].

The stationary solutions are characterized by a constant real Killing direction  $K^\mu$ , which corresponds to invariance of the metric  $g_{\mu\nu}$  with respect to the action of the Killing operator  $\hat{K} = K^\mu \partial_\mu$ ,

$$\hat{K}g^{\mu\nu} = 0. \quad (16)$$

It imposes requirements on the Kerr congruence,  $\hat{K}e^3 = 0$ , on the functions  $Y^\pm(x)$  and  $\bar{Y}^\pm(x)$ ,

$$\hat{K}Y = \hat{K}\bar{Y} = 0, \quad (17)$$

and as a consequence, on the Kerr generating function  $F(Y, x^\mu)$  and on the coefficients  $\hat{K}A = \hat{K}B = \hat{K}C = 0$ .

### 3 Complex structure of the Kerr geometry

The complex radial distance  $\tilde{r} = r + ia \cos \theta$  takes in the Cartesian coordinates (5) the form  $\tilde{r} = \sqrt{x^2 + y^2 + (z + ia)^2}$ , and therefore, the scalar component of the vector potential (6) may be obtained from the Coulomb solution  $\phi(\vec{x}) = e/r = e/\sqrt{x^2 + y^2 + z^2}$  by a complex shift  $z \rightarrow z + ia$ , or by the shift of its singular origin  $\vec{x}_L = (0, 0, 0)$  in the complex direction  $\vec{x}_L \rightarrow (0, 0, -ia)$ . The complex shift was first considered by Appel in 1887 [25], who noticed that the Coulomb solution is invariant with respect to the real shift of coordinates, and due to linearity of the Laplace equation, it should also be invariant with respect to the complex shift of its origin, which is trivial gauge transformation. This procedure results in nontrivial consequences for the real section of the solution. In particular, singular point of the Coulomb solution  $\vec{x}_L = (0, 0, 0)$  turns into singular ring  $x^2 + y^2 + (z + ia)^2 = 0$  (intersection of the sphere  $x^2 + y^2 + z^2 = a^2$  and plane  $z = 0$ ), which becomes a branch line of the space creating a twosheeted space-time.

It has been shown [8, 26] that the KN electromagnetic field may be obtained by the complex shift from the Coulomb solution. and Lind and Newman suggested in [8] a retarded-time construction, in which the linearized KN solution was represented as field generated by a complex source propagating along a complex world line (CWL). Later on, it was obtained in [9, 10] that the retarded-time construction turns out to be exact representation in the Kerr-Schild formalism. This *exactness* appears as a consequence of the specific linearization of the KS gravity due to analyticity of the twistorial structure determined by the Kerr theorem.

Principal element of the complex retarded-time construction is a family of the complex light cones  $\mathcal{K}$  adjoined to each point of the CWL. Taking the Left CWL as a basis for treatment, one can represent the family of the adjoined complex light cones in spinor form

$$\mathcal{K}_L = \{x : x = x_L^i(\tau_L) + \psi_L^A \sigma_{A\dot{A}}^i \tilde{\psi}_R^{\dot{A}}\}. \quad (18)$$

The cones are split into two families of null planes: "Left" ( $\psi_L = \text{const}$ ;  $\tilde{\psi}_R$  - var.) and "Right" ( $\tilde{\psi}_R = \text{const}$ ;  $\psi_L$  - var.). The null rays of the Kerr congruence appear as a real slice of the Left (or Right) complex null planes on the KS auxiliary Minkowski background. The real null direction  $e_\mu^3$ , (3) is a projective form of the spinor expression  $\psi_L^A \sigma_{\mu A\dot{A}} \tilde{\psi}_R^{\dot{A}}$ , in which the spinor  $\psi_L^A$  is replaced by the first projective twistor coordinate  $T^1 \equiv Y = \psi_L^2 / \psi_L^1$ . The real form  $e^3$  is completed by two complex conjugate null forms

$$e^1 = d\zeta - Y dv, \quad e^2 = d\bar{\zeta} - \bar{Y} dv. \quad (19)$$

One sees that the second projective twistor coordinate  $T^2 \equiv \lambda^1 = \zeta - Yv = x^\mu e_\mu^1$  represents projection of the space-time point  $x^\mu$  on the null direction  $e^1$ , while the third projective twistor coordinate  $T^3 \equiv \lambda^2 = u + Y\bar{\zeta} = x^\mu (e_\mu^3 - \bar{Y} e_\mu^1)$  represents a linear combination of the projections on the null directions  $e^3$  and  $e^1$ . The determined by the Kerr theorem function  $Y(x)$  allows one to restore at each point  $x$  the remainder twistor coordinates  $\lambda^1$  and  $\lambda^2$ , and to fix the incident Left null plane spanned by the null directions  $e^3$  and  $e^1$ . It should be mentioned *analyticity* of this construction. The Left null plane connects the real points  $x^\mu$  with a complex point of the Left CWL  $x_L^\mu$ . It belongs simultaneously to the in- fold of the complex light cone adjoined to the point  $x^\mu$  and to the out-fold of the light cone emanating from a point of the Left CWL  $x_L^\mu$ . The point of intersection of the Left null plane with CWL  $x_L^\mu$  fixes the value of the Left retarded time  $\tau_L$ , which becomes an analytic complex scalar function on the real space-time  $\tau_L(x^\mu)$ . In [10] this incidence was called as L-projection.

Similar system of the Right null planes, spanned by null directions  $e^3$  and  $e^2$ , connects real points  $x^\mu$  with Right CWL and determines *anti-analytic* R-projection and the retarded times  $\tau_R(x^\mu)$ . For the real  $x^\mu$  functions  $Y(x^\mu)$  and  $\bar{Y}(x^\mu)$  are complex conjugate and determine the conjugate Right and Left null planes. Analyticity of the generating function of the Kerr theorem  $F(T^A)$  and the corresponding solutions  $Y^\pm(x^\mu)$  creates independence of the analytic and anti-analytic L- and R- structures, allowing the complex-analytic extension of

the real KS geometry. In accord to analyticity of the Kerr theorem, the L-projection foliates the complexified Minkowski space-time  $CM^4$  into the Left null planes. Due to algebraically specific structure of the KS metric (1) and the gauge field (6), the non-linear terms drop out, and the analytic Left foliation performs a specific *linearization of the KS background*. In particular, it results in a *twistorial version* of the Fourier transform (introduced by Witten in [27]), which acts on the null planes of the curved KS space-time performing a bridge between KS gravity and quantum theory.

The stationary KN solution is represented as a retarded-time field generated by a *complex source moving along a straight CWL*, [9, 10],

$$x_L^\mu(\tau_L) = x_0^\mu(0) + u^\mu \tau_L + \frac{ia}{2} \{k_L^\mu - k_R^\mu\}, \quad (20)$$

where  $u^\mu = (1, \vec{v})$  is a real 4-velocity,  $k_R = (1, 0, 0, 1)$ ,  $k_L = (1, 0, 0, -1)$ , and  $\tau_L = t_L + i\sigma_L$  is a complex retarded-time parameter. Index  $L$  labels the Left complex structure, and we should also add the complex conjugate Right one

$$x_R^\mu(\tau_R) = x_0^\mu(0) + u^\mu \tau_R - \frac{ia}{2} \{k_L^\mu - k_R^\mu\}. \quad (21)$$

Complex shift turns the hedgehog of the Schwarzschild radial directions  $\vec{n} = \vec{r}/|r|$  into twisted directions of the Kerr congruence, Fig.1.

## 4 Complex open string

The complex world line  $x_0^\mu(\tau)$ , parametrized by complex time  $\tau$ , represents a two-dimensional surface which takes an intermediate position between particles and strings [12, 11]. The corresponding "hyperbolic string" equation [11],

$$\partial_\tau \partial_{\bar{\tau}} x_0(t, \sigma) = 0, \quad (22)$$

corresponds to bosonic part of the complex N=2 string [16]. The general solution  $x_0(t, \sigma) = x_L(\tau) + x_R(\bar{\tau})$  is a sum of the analytic and anti-analytic modes  $x_L(\tau)$ ,  $x_R(\bar{\tau})$ , which are not necessarily complex conjugate. For each real point  $x^\mu$ , the parameters  $\tau = \tau_L$  and  $\bar{\tau} = \tau_R$  should be determined by a complex retarded-time construction, a complex analog of the real one. Complex source of the real KN solution corresponds to two complex conjugate *straight* world-lines, (20) and (21). The complex light cone (18) is split into the Left and Right complex null planes. In accord to the retarded-(advanced)-time equations  $\tau^\mp = t \mp \tilde{r}$ , intersections of the Left null planes with Left CWL, together



with the conjugate Right structure, determine four retarded-advanced complex time parameters [9, 10]

$$\tau_L^\mp = t \mp (r_L + ia \cos \theta_L) \quad (23)$$

$$\tau_R^\mp = t \mp (r_R + ia \cos \theta_R). \quad (24)$$

The real slice of Kerr geometry is formed by the complex conjugate roots




Figure 2: The complex conjugate Left and Right null planes generate four roots for the Left and Right retarded and advanced times.

$\tau_L = t_L + i\sigma_L$  and  $\tau_R = t_R + i\sigma_R$ , which sets the constraint  $\sigma_L = -\sigma_R = \sigma = a \cos \theta$  resulting in dependence of the angular directions of the null rays  $\theta$  on the world-sheet parameter  $\sigma$ . Since  $|\cos \theta| \leq 1$ , we obtain  $\sigma \in [-a, a]$ , and consequently, *the complex string should have the endpoints  $\sigma = \pm a$* , and is to be open. The complex endpoints are mapped to the ‘north’ and ‘south’ lines of the Kerr congruence, corresponding to  $\theta = 0, \pi$ , as it is shown in Figure 3.

**World sheet orientifold.** The complex open string boundary conditions require the *orientifold* structure [12, 16, 28] which turns the open string in a closed but folded one. The world-sheet parity transformation  $\sigma \rightarrow -\sigma$  reverses orientation of the world sheet, and covers it second time in mirror direction. Simultaneously, the Left and Right modes are to be exchanged. Two oriented copies of the interval  $\Sigma = [-a, a]$ ,  $\Sigma^+ = [-a, a]$ , and  $\Sigma^- = [-a, a]$  are joined, forming a circle  $S^1 = \Sigma^+ \cup \Sigma^-$ , parametrized by  $\theta$ , and the map  $\theta \rightarrow \sigma = a \cos \theta$  covers the world-sheet twice. The string modes  $x_L(\tau)$ ,  $x_R(\bar{\tau})$ , are

extended on the second half-cycle by the well known extrapolation, [16, 28]

$$x_L(\tau^+) = x_R(\tau^-); \quad x_R(\tau^+) = x_L(\tau^-), \quad (25)$$

which forms the folded string, in which the retarded and advanced modes are exchanged every half-cycle. Revers of  $\sigma$  acts on the complex radial coordinate  $\tilde{r} = r + ia \cos \theta$  as anti-analytic involution

$$C : r + ia \cos \theta \rightarrow r - ia \cos \theta. \quad (26)$$

Another anti-analytic involution is reflection of the real radial coordinate,  $R : r \rightarrow -r$ . Analyticity of the world sheet is restored by use of double anti-analytic involution  $\mathcal{T} = CR$ . Applying  $\mathcal{T}\tilde{r} = -\tilde{r}$  to the Left CWL, we obtain

$$\mathcal{T} : \tau_L^\pm \rightarrow \tau_L^\mp, \quad (27)$$

which sets orientifold parity between retarded and advanced times of the same Left CWL allowing to leave aside the anti-analytical Right structure.




Figure 3: Ends of the open complex string, associated with quantum numbers of quark-antiquark pair, are mapped onto the real half-infinite  $z^+$ ,  $z^-$  axial strings. Dotted lines indicate orientifold projection.

## 5 Embedding of the complex string in the real Kerr-Schild geometry

The complex world-sheet parameters  $\tau^\pm = t \pm r + ia \cos \theta$  are mapped in the real KS geometry as coordinates  $\rho^\pm = t \pm r$  and  $\theta$ , and therefore, the Kerr coordinate transformations (5) may be considered as *embedding* of the

world-sheet in the bulk of the *real* KS geometry. Simultaneously, it is the embedding of the twistorial coordinates of the adjoined Left null planes. The first twistor coordinate  $Y = \psi_L^1/\psi_L^0$  determines directions of the null rays  $Y = e^{i\phi} \tan \frac{\theta}{2}$  in accord to (3). The coordinate  $\phi$  adds extra dimension to the world-sheet, extending it to a membrane formed by merging of the real closed Kerr string with the complex open string, similar to the model of string/M-theory unification [28, 1].

The given by Kerr theorem two solutions  $Y^\pm(x)$  have different embeddings in the real KS geometry. For simplicity, we consider the Kerr source at rest in the standard orientation of angular momentum along the z-axis. Assuming that the complex source is positioned at the point  $\vec{x}_L = (0, 0, -ia)$ , and setting  $x_0^\mu(0) = 0$  and  $u^\mu = (1, 0, 0, 0)$  in (20), we rewrite the Left CWL and its time-derivatives in the null coordinates (4) as follows,

$$x_L(\tau_L) = \{u_L, v_L, \zeta_L, \bar{\zeta}_L\} = 2^{-\frac{1}{2}}\{(-ia + \tau_L), (-ia - \tau_L), 0, 0\}, \quad (28)$$

$$\dot{x}_L = \{\dot{u}_L, \dot{v}_L, \dot{\zeta}_L, \dot{\bar{\zeta}}_L\} = 2^{-\frac{1}{2}}\{1, -1, 0, 0\}. \quad (29)$$

Coefficients  $A$ ,  $B$ ,  $C$  were obtained for this case in [9, 10], and take the form

$$\begin{aligned} A_L &= (\bar{\zeta} - \bar{\zeta}_L(\tau))\dot{v}_L(\tau) - (v - v_L(\tau))\dot{\bar{\zeta}}_L(\tau); \\ B_L &= (u - u_L(\tau))\dot{v}_L(\tau) + (\zeta - \zeta_L(\tau))\dot{\bar{\zeta}}_L(\tau) - (\bar{\zeta} - \bar{\zeta}_L(\tau))\dot{\zeta}_L(\tau) - (v - v_L(\tau))\dot{u}_L(\tau); \\ C_L &= (\zeta - \zeta_L(\tau))\dot{u}_L(\tau) - (u - u_L(\tau))\dot{\zeta}_L(\tau), \end{aligned} \quad (30)$$

which yields for the Left CWL the expressions

$$A_L = -\frac{1}{2}(x - iy), \quad B_L = -z - ia, \quad C_L = \frac{1}{2}(x + iy), \quad (31)$$

and using generating function  $F_L(T^A) = A_L Y^2 + B_L Y + C_L$  we obtain

$$Y_L^\pm = \frac{-z - ia \pm \tilde{r}_L}{x - iy}, \quad (32)$$

where

$$\tilde{r}_L = (B_L^2 - 4A_L C_L)^{\frac{1}{2}} = [x^2 + y^2 + (z + ia)^2]^{\frac{1}{2}} \quad (33)$$

corresponds to the complex radial distance of the Appel source. We will further omit the suffix  $L$ , if we work only with the Left CWL. Representing the complex radial distance in the form  $\tilde{r} = r + i\eta$  and squaring (33), we obtain two real equations. The first of them,  $r\eta = az$ , may be treated as an analog of the relation  $z = r \cos \theta$ , and, setting  $\eta = a \cos \theta$ , we obtain

$$\tilde{r} = r + ia \cos \theta, \quad z = r \cos \theta. \quad (34)$$

The second equation, following from (33), may be combined with (34) and yields  $x^2 + y^2 = (r^2 + a^2) \sin^2 \theta$ , which may be split into four different linear relations  $x + iy = (r \pm ia)e^{\pm i\phi} \sin \theta$ , characterized by different combinations of the signs  $\pm ia$  and  $\pm i\phi$ . We obtain four sheets corresponding to four different embeddings of the world-sheet of the complex string.

**Orientifolding the bulk of the Kerr space.** Embedding of the world-sheet fixes also the map of the Left null planes into the real rays of the Kerr congruence in accord to (3). Considering the Kerr coordinate relations (5) as *analytic* one, we can represent the given by (32) solution  $Y^+$  in the form

$$Y^+ = \frac{e^{i\phi}(1 - \cos \theta)}{\sin \theta} = e^{i\phi} \tan \frac{\theta}{2}, \quad (35)$$

which is just the original form of  $Y$  given in [18]. In the same time, for  $Y^-$  we obtain a bizarre expression  $Y^- = \frac{-(r+ia)(1+\cos \theta)}{x-iy}$ , which shows that ‘analytic’ embedding (5) is inconsistent with  $Y^-$ , and consistency is related with ‘anti-analytic’ transformation

$$x + iy = (r - ia)e^{i\phi} \sin \theta, \quad z = r \cos \theta, \quad (36)$$

which results in the relation

$$Y^- = \frac{-e^{i\phi}(1 + \cos \theta)}{\sin \theta} = -e^{i\phi} \cot \frac{\theta}{2} = -1/\bar{Y}^+. \quad (37)$$

We reveal that the solutions  $Y^+$  and  $Y^-$  are related by antipodal involution

$$I^* : Y^+ \rightarrow Y^- = -1/\bar{Y}^+, \quad (38)$$

which is correlated with the crossing-symmetry of the Left and Right world-sheet structures (25), as it is shown in Fig.2. Antipodal involution

$$I^* : \theta \longrightarrow \tilde{\theta} = \pi - \theta, \quad \phi \longrightarrow \tilde{\phi} = \phi + \pi \quad (39)$$

was studied for the case of non-rotating black holes in [29], where authors considered formation of the non-orientable or single-exterior BH solutions.

**Calabi-Yau twofold from the Kerr theorem.** As follows from (32), coordinates  $x_L^\mu(\tau^-)$  and 4-velocity  $u^\mu(\tau^-)$  of the Left CWL determined by the Left complex retarded time  $\tau_L^-$  lead to two solutions  $Y^\pm(x^\mu, \tau_L^-)$ . One sees from (11) that  $Y^+$  and  $Y^-$  differ only in the sign of  $\tilde{r}$ . In accord to (27), this

difference corresponds to change the Left retarded time  $\tau_L^-$  to the Left advanced time  $\tau_L^+$ , and therefore, the second solution should be identified with the Left advanced one  $Y^-(x^\mu, \tau_L^+)$ . The both Left solutions are analytic, since the transfer to  $Y^-(x^\mu, \tau_L^+)$  may be represented as result of two anti-analytic transformations: complex conjugation  $C$  and antipodal map  $I^*$ , like the two-step transformation of the world-sheet (27). On the real slice, the Left and Right retarded solutions,  $Y^+(x^\mu, \tau_L^-)$  and  $\bar{Y}^+(x^\mu, \tau_R^-)$ , are complex conjugate, as well as the corresponding retarded times  $\tau_R^-$  and  $\tau_L^-$ . Antipodal map (38) connects the Left advanced solution  $Y^-(x^\mu, \tau_L^+)$  with the Right retarded one  $\bar{Y}^+(x^\mu, \tau_R^-)$ , forming a crossing-symmetry, as shown in Fig.2. On the real slice the Left retarded and advanced solutions are rigidly correlated. However, this correlation is destroyed when  $x^\mu$  escapes the real slice. Complex analyticity of the Kerr theorem is used by integration of the field equation [18], which assumes independence of the Left generating functions  $F_{ret} \equiv F\{T^A, \tau^+\}$  and  $F_{adv} \equiv F\{T^A, \tau^-\}$  and the corresponding solutions  $Y_{ret}^-(x^\mu) \equiv Y^-(x^\mu, \tau_L^-)$  and  $Y_{adv}^+ \equiv Y^+(x^\mu, \tau_L^+)$ .

Each of the functions  $F_{ret}\{T^A\}$  and  $F_{adv}\{T^A, \}$  creates two sheets of the Kerr-Schild geometry. The space-time, determined by two independent generating function turns out to be four-sheeted and should be determined by multi-particle version of the Kerr theorem considered in [22, 23]. In particular, the analytic bipartite generating function  $F_2(T^A)$  should has the form  $F_2(T^A) = F_{ret}(T^A)F_{adv}(T^A)$ . As a consequence of (13),

$$F_2(T^A) = A_{ret}A_{adv}(Y - Y_{ret}^+(x^\mu))(Y - Y_{ret}^-(x^\mu))(Y - Y_{adv}^+(x^\mu))(Y - Y_{adv}^-(x^\mu)), \quad (40)$$

and the equation (7) acquires the fourth degree in  $Y$  forming a *quartic in  $CP^3$* , which is a canonic form of the Calabi-Yau two-fold, [28, 16]. For each point  $x^\mu \in M^4$ , solutions  $Y(x^\mu)$  may be transformed to twistor coordinates  $T^A$  by means of (8) and (4).

The four solutions  $Y(x^\mu)$  correspond to different sheets of the K3 surface and perform different embeddings of the Kerr congruence in the real Kerr geometry, in accord to (3). Antipodal map acts on the real null directions  $e^{3\mu}$  as inversion of its spacelike part, and therefore, the congruences generated by functions  $F_{ret}$  and  $F_{adv}$  are not to be equivalent. In particular, the tangent to Kerr congruence normalized null vector

$$k^\mu(x) = e^{3\mu}(x)/|e^{30}|, \quad (41)$$




Figure 4: One sheet of the K3 for  $r > 0$  and  $\phi = const.$  Kerr congruence is tangent to singular ring at  $\theta = \pi/2$ .




Figure 5: Section of the K3 near the Kerr ring corresponding to  $\phi = const.$ . Two sheets ( $r > 0$  and  $r < 0$ ) form a Möbius strip covering the space-time twice.

is transformed as follows

$$I^* : k_{out}^\mu = \{k^0, \vec{k}\} \rightarrow k_{in}^\mu = \{k^0, -\vec{k}\}. \quad (42)$$

A slice of the K3 surface corresponding to  $\phi = const.$  and  $r > 0$  is presented in Fig.4. One see that this sheet is a ruled surface generated by twistor null lines parametrized by affine parameter  $r \in [0, \infty]$ . By variation of  $\theta \in [0, \phi]$ , the lines rotate on angle  $\theta$ . The lines of the equatorial plane,  $\theta = \pi/2$ , are tangent to the Kerr ring. Analytic extension of this surface  $\phi = const.$  to region of negative  $r \in [0, -\infty]$ , as shown in Fig.1 and Fig.5, gives the second sheet of the K3 which corresponds to ‘negative’ sheet of the Kerr geometry. The surface form a Möbius strip, since the boundary lines at  $\theta = 0$  and  $\theta = \pi$  have opposite orientation. Antipodal map  $I^*$  creates two extra sheets of the K3 which are advanced folds of the Kerr congruence. The global antipodal involution (39) transforms the surface  $\phi = const.$  into the surface  $\phi + \pi = const.$  with simultaneous involution of the null generators. Although the surfaces  $\phi = const.$  and  $\phi + \pi = const.$  are smoothly matched, the North and South boundary lines of these surfaces have opposite orientation. So far as the orientation is changed by the transfer from  $r > 0$  to  $r < 0$ , one sees that analytic gluing of the four sheets of the K3 are to be reached under a cross-symmetric matching of the boundaries of the retarded and advanced folds of the Kerr congruence. The four folds of the K3 surface form orientifold parity of the Kerr geometry, under which the principal null directions  $e^3$  are doubled by the joined in-going and out-going null rays.

*String excitations.* The Killing direction  $K^\mu$  may be expressed in Cartesian coordinates via coordinates of the CWL

$$K^\mu = \partial_t x_L^\mu(\tau), \quad (43)$$

and the coefficients  $A, B, C$  (31) turn out to be functions of the coordinates and 4-velocity of the CWL  $u^\mu$ . The string excitations are related with radiation and a recoil leading to non-stationarity of the solution. The exact non-stationary KS solutions are unknown, however the recoilless solutions were obtained in [24, 30]. A recoil assumes a deviation of the 4-velocity  $\delta x_L^\mu(\tau)|_{\tau^-}$  for the moments  $t > 0$ , which creates a difference between parameters  $A, B, C$ , determined for the retarded and advanced times  $\tau^-$  and  $\tau^+$ , resulting in two independent generating functions  $F_{ret}(T^A)$  and  $F_{adv}(T^A)$ . This duplications of the Kerr congruence generates the four sheets of the K3 surface. Independence of the retarded and advanced solutions creates breakdown of the orientifold parity of the Kerr congruence. The North and South boundaries of the Kerr sheets are decoupled, but the smooth pairwise matching of the four sheets is retained, resulting in analyticity of the K3 surface.

## 6 Outlook.

It has been discussed in I that the Kerr-Schild geometry displays striking parallelism with basic structures of the superstring theory. However, along with wonderful parallelism, the KN geometry displays also several peculiarities.

The principal of them is that the Kerr real and complex Kerr strings, and the related stringy structures such as orientifold and formation of the M2-brane, are living in the four-dimensional space-time. There appears a wonderful possibility that compactification of higher dimensions may be replaced by complexification.

Second peculiarity is related with characteristic parameter of the Kerr strings  $a = \hbar/m$ , which corresponds to Compton scale, and this new parameter makes this version of the superstring theory to be closer to particle physics.

Finally, we notice that among the consistent critical strings in dimensions  $d=26$  and  $d=10$ , there is also the complex  $N=2$  string [16, 17], which has the real critical dimension four and could be used as a basis of some four-dimensional string theory, [31]. The principal obstacle for its application emerges from its signature, which may only be  $(2,2)$  or  $(4,0)$ , which conflicted with its embedding in the real minkowskian space-time. Up to our knowledge,

this trouble was not resolved so far, and the initially enormous interest to this string seems to be dampened. We note that N=2 string has organic embedding in the complexified 4d Kerr geometry. Hermitian action for the Kerr's complex world line (CWL),  $S = -\frac{1}{2\pi} \int d\tau d\bar{\tau} g_{\mu\nu} \partial_\tau x^\mu \partial_{\bar{\tau}} \bar{x}^\nu$ , corresponds to bosonic part of the N=2 string action [11, 16, 17]. Generalization  $\tau$  to complex super-time  $\mathcal{T}$  turns CWL into super-world-line  $X^\mu(\mathcal{T})$ , [12, 32], and leads to action of the N=2 superstring [16]. The extra world-sheet spinor field  $\psi(\tau)$  corresponds to the Left null planes, generators of the K3 and Kerr congruence. Detailed treatment of this structure goes beyond the frame of this paper and will be given elsewhere.

## Acknowledgments

This work is supported under the RFBR grant 13-01-00602. Author thanks Theo M. Nieuwenhuizen for permanent interest to this work and useful conversations, and also thanks Dirk Bouwmeester for invitation to Leiden University, where these ideas were discussed in details with him and members of his group: Jan W. Dalhuisen, V.A.L. Thompson and J.M.S. Swearngin. Author is also thankful to Laur Jhärvi for useful discussion at the Tallinn conference "3Quantum" and given reference to the paper [14].

## References

- [1] A. Burinskii, *Adv.High Energy Phys.* 2013:509749, (2013).  
Doi:10.1155/2013/509749, [arXiv:1211.6021]
- [2] A. Burinskii, *J. Phys. A: Math. Theor.* **43** (2010) 392001 [arXiv:1003.2928].
- [3] D.D. Ivanenko and A.Ya. Burinskii, *Gravitational strings in the models of elementary particles, Izv. Vuz. Fiz.* **5** (1975) 135.
- [4] A. Dabholkar, J. P. Gauntlett, J. A. Harvey, D. Waldram *Nucl.Phys.* **B 474** 85-121 (1996), [arXiv:hep-th/9511053],
- [5] A. Burinskii, *Phys. Rev. D* **52** (1995) 5826, [arXiv:hep-th/9504139].
- [6] A. Sen *Macroscopic Charged Heterotic String Nucl.Phys.* **B 388** 457 (1992) [arXiv:hep-th/9206016].
- [7] A. Sen, *Phys. Rev. Lett.* **69** 1006 (1992).
- [8] R.W. Lind and E.T. Newman, *J. Math. Phys.* **15** 1103 (1974).
- [9] A. Burinskii and R.P. Kerr, *Nonstationary Kerr Congruences*, [arXiv:gr-qc/9501012].



- [10] A. Burinskii, *Phys. Rev.* **D 67**, (2003) 124024.
- [11] H. Ooguri and C. Vafa. *Nucl. Phys.* **B 361** (1991) 469; *ibid* *N = 2 heterotic strings.* **B 367** (1991) 83.
- [12] A.Ya. Burinskii, *String-like Structures in Complex Kerr Geometry*, In: “*Relativity Today*”, Edited by R.P.Kerr and Z.Perjés, Akadémiai Kiadó, Budapest, 1994, p.149, [arXiv:gr-qc/9303003].
- [13] A. Burinskii, *Phys. Rev.* **D 70** (2004) 086006, [arXiv:hep-th/0406063].
- [14] L.M. Dyson, L. Järv and C. V. Johnson, *Oblate, Toroidal, and Other Shapes for the Enhanceon*, *JHEP* **0205** (2002) 019 [arXiv:hep-th/0112132],
- [15] T.M. Adamo and E.T. Newman, *Phys. Rev.* **D 83** (2011) 044023
- [16] M.B. Green, J. Schwarz, and E. Witten, *Superstring Theory, v.I*, Cambridge Univ. Press, 1987.
- [17] J. Barrett., G.W. Gibbons., M.J. Perry., C.N.Pope and P.Ruback, *Int.J.Mod.Phys. A* **9** (1994) 1457, [hep-th/9302073].
- [18] G. C. Debney, R. P. Kerr and A. Schild, *J. Math. Phys.* **10** 1842 (1969).
- [19] D.Kramer, H.Stephani, E. Herlt, M.MacCallum, *Exact Solutions of Einstein’s Field Equations*, Cambridge Univ. Press, Cambridge 1980.
- [20] R. Penrose, *J. Math. Phys.* **8** (1967) 345.
- [21] R. Penrose and W. Rindler, *Spinors and Space-time, Vol. 2: Spinor and twistor methods in space-time geometry*, Cambridge University Press, Cambridge U.K. (1986), pg. 501.
- [22] A. Burinskii, *Grav. Cosmol*, **11** (2005)301 [arXiv:hep-th/0506006].
- [23] A. Burinskii, *Int.J.Geom.Meth.Mod.Phys.* **4** (2007) 437 [hep-th/0510246].
- [24] A. Burinskii, *Gen. Rel. Grav.* **41** (2009) 2281 [arXiv:0903.3162].
- [25] E.T. Whittacker and G.N. Watson, *A Course of Modern Analysis*, Cambridge Univ. Press London/New York,p.400, 1969.
- [26] A.Ya. Burinskii, *Soviet Phys. JETP* **39** (1974) 193.
- [27] E. Witten, *Comm. Math. Phys.* **252** (2004) 198 [arXiv:hep-th/0312171].
- [28] K. Becker, M. Becker and J. Schwarz, *String Theory and M-Theory - A Modern Introduction*, Cambridge University Press, 2007.
- [29] A. Chamblin and G.W. Gibbons, *Phys. Rev.***D 55** (1997) 2177.
- [30] A. Burinskii, *Theor. Math.Phys.* **163**(3), (2010) 782 [arXiv:0911.5499],
- [31] A. D’Adda and F. Lizzi, *Phys. Lett.* 191B (1987) 85.
- [32] A.Ya. Burinskii, *Phys. Rev.***D 57** (1998) 2392.