# Early thermalization of quark-gluon matter initially created in high-energy nucleus-nucleus collisions

Xiao-Ming Xu

Department of Physics, Shanghai University, Baoshan, Shanghai 200444, China E-mail: xmxu@mail.shu.edu.cn

**Abstract.** Elastic parton-parton-parton scattering is briefly reviewed and is included in transport equations of quark-gluon matter. We solve the transport equations and get thermal states from initially produced quark-gluon matter. Both gluon matter and quark matter take early thermalization, but gloun matter has a shorter thermalization time than quark matter.

#### 1. Introduction

A large amount of particles move randomly and momentum distribution functions in different directions are identical. The distribution depends only on the absolute value of the threedimensional momentum. The dependence indicates the existence of a thermal state and exposes the basic quantity temperature, which value is acquired from the analytic expression of the distribution (e.g., temperature is in the Boltzmann distribution). Unfortunately, the system of gluons, quarks and antiquarks produced in initial nucleus-nucleus collisions at the Relativistic Heavy Ion Collider (RHIC) energies does not possess temperature. This is because the distribution in the incoming nucleus beam direction is much larger than the distribution in the direction perpendicular to the beam direction [1]. However, the system evolves rapidly into a thermal state (a quark-gluon plasma) as concluded from the elliptic flow data of hadrons [2] and the corresponding explanation of hydrodynamic calculations which assume early thermalization and ideal relativistic fluid flow [3, 4]. The early thermalization or the rapid creation of temperature is generally interesting. Parton-parton scattering was employed to study thermalization of initially produced quark-gluon matter, but is not enough to explain the early thermalization [5, 6, 7, 8]. Elastic gluon-gluon-gluon scattering was proposed to explain the early thermalization of gluon matter [9]. Nevertheless, elastic quark-quark scattering and elastic quark-quark-guark scattering give a long thermalization time of quark matter [10]. Recently we have obtained that quark matter takes the early thermalization due to all types of elastic scattering that involves quarks [11]. In the next section I briefly introduce the elastic scattering of quark-quark-quark [10], quark-quark-antiquark [12], quark-antiquarkantiquark, antiquark-antiquark, gluon-quark-quark [13], gluon-quark-antiquark [13], gluon-antiquark-antiquark, gluon-gluon-quark [11], gluon-gluon-antiquark or gluon-gluon-gluon [9]. In section 3 we present transport equations that include all types of elastic parton-partonparton scattering and numerical results of the equations. Conclusions are in the last section.

**Table 1.** The number of diagrams for elastic parton-parton-parton scattering, the number of diagrams in a class, the number of the triple-gluon vertex and the number of the four-gluon vertex of a diagram.

| elastic                       | number of  | class  | number of  | number of     | number of   |
|-------------------------------|------------|--------|------------|---------------|-------------|
| scattering                    | diagrams   |        | diagrams   | the 3-gluon   | the 4-gluon |
| 0                             | for a type |        | in a class | vertex        | vertex      |
| guark-guark-guark 1           | 42         | 1      | 36         | 0             | 0           |
|                               |            | 2      | 6          | 1             | 0           |
| quark-quark-quark 2           | 14         | 1      | 12         | 0             | 0           |
|                               |            | 2      | 2          | 1             | 0           |
| quark-quark-quark 3           | 7          | 1      | 6          | 0             | 0           |
|                               |            | 2      | 1          | 1             | 0           |
| quark-quark-antiquark 1       | 58         | 1      | 52         | 0             | 0           |
|                               |            | 2      | 6          | 1             | 0           |
| quark-quark-antiquark 2       | 14         | 1      | 12         | 0             | 0           |
|                               |            | 2      | 2          | 1             | 0           |
| quark-quark-antiquark 3       | 29         | 1      | 26         | 0             | 0           |
|                               |            | 2      | 3          | 1             | 0           |
| quark-quark-antiquark 4       | 7          | 1      | 6          | 0             | 0           |
|                               |            | 2      | 1          | 1             | 0           |
| gluon-quark-quark 1           | 72         | 1      | 40         | 0             | 0           |
|                               |            | 2      | 24         | 1             | 0           |
|                               |            | 3      | 8          | 2             | 0           |
| or                            | 20         |        | 20         | 0             | 1           |
| gluon-quark-quark 2           | 36         | 1      | 20         | 0             | 0           |
|                               |            | 2      | 12         | 1             | 0           |
|                               |            | 3      | 4          | 2             | 0           |
| Or<br>gluon quark antiquark 1 | 76         | 1      | 40         | 0             | 1           |
| giuon-quark-antiquark 1       | 70         | 1      | 40         | 0             | 0           |
|                               |            | 2<br>2 | 20         | 1             | 0           |
| or                            |            | 5      | 0          |               | 0           |
| gluon-quark-antiquark 2       | 36         | 1      | 20         | 0             | 0           |
| giuon-quark-antiquark 2       | 50         | 2      | 20<br>12   | 1             | 0           |
|                               |            | 3      | 4          | 2             | 0           |
| or                            |            | 0      | 1          | $\frac{2}{0}$ | 1           |
| gluon-gluon-quark             | 123        | 1      | 24         | 0             | 0           |
| 0 1 0 1                       | -          | 2      | 36         | 1             | 0           |
|                               |            | 3      | 38         | 2             | 0           |
| or                            |            |        |            | 0             | 1           |
|                               |            | 4      | 25         | 3             | 0           |
| or                            |            |        |            | 1             | 1           |
| gluon-gluon-gluon             | 220        | 1      | 105        | 4             | 0           |
|                               |            | 2      | 96         | 2             | 1           |
|                               |            | 3      | 19         | 0             | 2           |





**Figure 1.** Elastic gluon-quark-antiquark scattering. The wiggly (solid) lines stand for gluons (quarks and antiquarks).

Figure 2. Same as figure 1.

## 2. Elastic parton-parton-parton scattering

Each type of elastic 3-to-3 scattering is divided into 2, 3 or 4 classes according to the number of the triple-gluon vertex and the number of the four-gluon vertex. In Table 1 we list the number of diagrams for each type or in a class, the number of the triple-gluon vertex and the number of the four-gluon vertex of a diagram. The number following the entry quark-quark-quark has the meanings: the three quarks are identical at 1; only two quarks are identical at 2; the three quarks are different at 3. The number following the entry quark-quark-antiquark has the meanings: the antiquark have the same flavor at 1; only the two quarks have the same flavor at 2; only one quark has the same flavor as the antiquark at 3; the three flavors of the two quarks and the antiquark are not identical at 4. The number following the entry gluon-quark-quark (gluon-quark-antiquark) has the meanings: the two quarks (the quark and the antiquark) have the same flavor at 1; the two quarks (the quark and the antiquark) have the same flavor at 2.

Feynman diagrams for the elastic quark-antiquark-antiquark scattering (antiquark-antiquarkantiquark, gluon-antiquark-antiquark, gluon-gluon-antiquark) are obtained from the Feynman diagrams for the elastic quark-quark-antiquark (quark-quark-quark, gluon-quark, gluongluon-quark) via the replacement of quark by antiquark and vice versa. To illustrate elastic 3-to-3 scattering, we show four Feynman diagrams in figures 1-4. In figures 1 and 2 quarkantiquark annihilation and creation occur. In both figure 3 and figure 4 two triple-gluon vertices and one four-gluon vertex are involved.

The elastic gluon-gluon-gluon scattering is complicated but rich in the process as shown by the 220 Feynman diagrams at the tree level [9]. To better understand the scattering, we sort out Feynman diagrams in every class. We select three triple-gluon vertices of a diagram to subclassify the diagrams in the first class and six subclasses are obtained according to the initial gluons and the final gluons possessed by the selected vertices. The subclasses are listed in Table 2. Every diagram in the second class has a four-gluon vertex and two triple-gluon vertices. We select the four-gluon vertex and a triple-gluon vertex to subclassify the diagrams in the second class and nine subclasses are obtained according to the initial gluons and the final gluons possessed by the four-gluon vertex and the selected triple-gluon vertex. The subclasses are listed in Table 3. Diagrams in the third class can not be subclassified.



**Figure 3.** Elastic gluon-gluon-gluon scattering. The wiggly lines stand for gluons.

Figure 4. Same as figure 3.

**Table 2.** Number of diagrams in a subclass identified by the three selected triple-gluon vertices.  $n_{1i}$  and  $n_{1f}$  ( $n_{2i}$  and  $n_{2f}$ ,  $n_{3i}$  and  $n_{3f}$ ) are numbers of initial and final gluons possessed by the first (second, third) triple-gluon vertex, respectively.

| (1, 1, 1, 1, 1) $(1, 1, 1)$ $(1, 1, 1)$ $(1, 1, 1)$ $(1, 1, 1)$ $(1, 1, 1)$ |           |                   |                   |          |                   |          |                   |
|---|-----------|-------------------|-------------------|----------|-------------------|----------|-------------------|
| subclass  | number of | $n_{1\mathrm{i}}$ | $n_{1\mathrm{f}}$ | $n_{2i}$ | $n_{2\mathrm{f}}$ | $n_{3i}$ | $n_{3\mathrm{f}}$ |
|   | diagrams  |                   |                   |          |                   |          |                   |
| 1   | 36        | 1                 | 1                 | 1        | 1                 | 1        | 0                 |
| 2   | 18        | 0                 | 2                 | 1        | 1                 | 1        | 0                 |
| 3   | 18        | 2                 | 0                 | 1        | 1                 | 0        | 1                 |
| 4   | 18        | 2                 | 0                 | 0        | 2                 | 1        | 0                 |
| 5   | 6         | 1                 | 1                 | 1        | 1                 | 0        | 0                 |
| 6   | 9         | 2                 | 0                 | 0        | 2                 | 0        | 0                 |
|   |           |                   | -                 |          | -                 | -        | -                 |

**Table 3.** Number of diagrams in a subclass identified by the four-gluon vertex and the selected triple-gluon vertex.  $n_{i4}$  and  $n_{f4}$  ( $n_{i3}$  and  $n_{f3}$ ) are numbers of initial and final gluons possessed by the 4-gluon vertex (the 3-gluon vertex), respectively.

|          |           | // 🔺     |             |          |             |
|----------|-----------|----------|-------------|----------|-------------|
| subclass | number of | $n_{i4}$ | $n_{ m f4}$ | $n_{i3}$ | $n_{ m f3}$ |
|          | diagrams  |          |             |          |             |
| 1        | 18        | 1        | 2           | 1        | 1           |
| 2        | 18        | 2        | 1           | 1        | 1           |
| 3        | 9         | 1        | 2           | 2        | 0           |
| 4        | 9         | 2        | 1           | 0        | 2           |
| 5        | 3         | 0        | 3           | 2        | 0           |
| 6        | 3         | 3        | 0           | 0        | 2           |
| 7        | 18        | 1        | 1           | 1        | 1           |
| 8        | 9         | 0        | 2           | 2        | 0           |
| 9        | 9         | 2        | 0           | 0        | 2           |

# 3. Transport equations and thermal states

We assume that quark-gluon matter consists of gluons, quarks and antiquarks with up and down flavors and the two flavors (the quark and the antiquark as well) have the same amount. Let the distribution functions for the gluon, the up-quark, the down-quark, the up-antiquark and the down-antiquark be  $f_{gi}$ ,  $f_{ui}$ ,  $f_{di}$ ,  $f_{\bar{u}i}$  and  $f_{\bar{d}i}$ , respectively, where *i* labels the *i*th parton in scattering, and  $f_{ui} = f_{di} = f_{\bar{u}i} = f_{qi}$ . The transport equation for gluon matter is

$$\begin{aligned} \frac{\partial f_{g1}}{\partial t} &+ \vec{v}_{1} \cdot \vec{\nabla}_{\vec{r}} f_{g1} &= -\frac{1}{2E_{1}} \int \frac{d^{3}p_{2}}{(2\pi)^{3}2E_{2}} \frac{d^{3}p_{3}}{(2\pi)^{3}2E_{3}} \frac{d^{3}p_{4}}{(2\pi)^{3}2E_{4}} (2\pi)^{4} \delta^{4}(p_{1}+p_{2}-p_{3}-p_{4}) \\ &\times \left\{ \frac{g_{G}}{2} \mid \mathcal{M}_{gg \rightarrow gg} \mid^{2} [f_{g1}f_{g2}(1+f_{g3})(1+f_{g4}) - f_{g3}f_{g4}(1+f_{g1})(1+f_{g2})] \right. \\ &+ g_{Q}(\mid \mathcal{M}_{gu \rightarrow gu} \mid^{2} + \mid \mathcal{M}_{gd \rightarrow gd} \mid^{2} \mid \mathcal{M}_{g\bar{u} \rightarrow g\bar{u}} \mid^{2} \mid \mathcal{M}_{g\bar{d} \rightarrow g\bar{d}} \mid^{2}) \\ &\times [f_{g1}f_{q2}(1+f_{g3})(1-f_{q4}) - f_{g3}f_{q4}(1+f_{g1})(1-f_{q2})]] \\ &- \frac{1}{2E_{1}} \int \frac{d^{3}p_{2}}{(2\pi)^{3}2E_{2}} \frac{d^{3}p_{3}}{(2\pi)^{3}2E_{3}} \frac{d^{3}p_{4}}{(2\pi)^{3}2E_{5}} \frac{d^{3}p_{5}}{(2\pi)^{3}2E_{5}} \frac{d^{3}p_{6}}{(2\pi)^{3}2E_{6}} \\ &\times (2\pi)^{4} \delta^{4}(p_{1}+p_{2}+p_{3}-p_{4}-p_{5}-p_{6}) \left\{ \frac{g_{G}^{2}}{12} \mid \mathcal{M}_{ggg \rightarrow ggg} \mid^{2} \\ &\times [f_{g1}f_{g2}f_{g3}(1+f_{g4})(1+f_{g5})(1+f_{g6}) - f_{g4}f_{g5}f_{g6}(1+f_{g1})(1+f_{g2})(1+f_{g3})] \\ &+ \frac{g_{G}g_{Q}}{2} (\mid \mathcal{M}_{ggu \rightarrow ggu} \mid^{2} + \mid \mathcal{M}_{ggd \rightarrow ggd} \mid^{2} + \mid \mathcal{M}_{gg\bar{u} \rightarrow gg\bar{u}} \mid^{2} + \mid \mathcal{M}_{gg\bar{d} \rightarrow gg\bar{d}} \mid^{2}) \\ &\times [f_{g1}f_{g2}f_{q3}(1+f_{g4})(1+f_{g5})(1-f_{q6}) - f_{g4}f_{g5}f_{q6}(1+f_{g1})(1+f_{g2})(1-f_{q3})] \\ &+ g_{Q}^{2} [\frac{1}{4} \mid \mathcal{M}_{guu \rightarrow guu} \mid^{2} + \frac{1}{2} (\mid \mathcal{M}_{g\bar{u}\bar{d} \rightarrow g\bar{u}\bar{d}} \mid^{2} + \mid \mathcal{M}_{gd\bar{u} \rightarrow gd\bar{u}} \mid^{2} + \mid \mathcal{M}_{gd\bar{u} \rightarrow gd\bar{u}} \mid^{2}) \\ &+ |\mathcal{M}_{g\bar{u}\bar{u} \rightarrow g\bar{u}\bar{u}} \mid^{2} + \frac{1}{2} (\mid \mathcal{M}_{g\bar{u}\bar{d} \rightarrow g\bar{u}\bar{d}} \mid^{2} + \mid \mathcal{M}_{gd\bar{u} \rightarrow gd\bar{u}} \mid^{2}) \\ &\times [f_{g1}f_{q2}f_{q3}(1+f_{g4})(1-f_{q5})(1-f_{q6}) - f_{g4}f_{q5}f_{q6}(1+f_{g1})(1-f_{q2})(1-f_{q3})]] \\ &+ g_{Q}^{2} [\frac{1}{4} \mid \mathcal{M}_{g\bar{u}\bar{u} \rightarrow g\bar{u}\bar{u}} \mid^{2} + \frac{1}{2} (\mid \mathcal{M}_{g\bar{u}\bar{d} \rightarrow g\bar{u}\bar{u}} \mid^{2} + \mid \mathcal{M}_{gd\bar{u} \rightarrow gd\bar{u}} \mid^{2}) \\ &\times [f_{g1}f_{q2}f_{q3}(1+f_{g4})(1-f_{q5})(1-f_{q6}) - f_{g4}f_{q5}f_{q6}(1+f_{g1})(1-f_{q2})(1-f_{q3})]] \\ &+ (f_{g1}f_{g2}f_{q3}(1+f_{g4})(1-f_{q5})(1-f_{q6}) - f_{g4}f_{q5}f_{q6}(1+f_{g1})(1-f_{q2})(1-f_{q3})]] \\ &+ (f_{g1}f_{g1}f_{g2}f_{g3}(1+f_{g4})(1-f_{q5})(1-f_{q6}) - f_{g4}f_{q5}f_{q6}(1+f_{g1})(1-f_{q2})(1-f_{q3})]] \\ &+ (f_{g1}f_{g1}f_{g2}f_{g3}(1+f_{g$$

and the transport equation for up-quark matter is

$$\begin{split} \frac{\partial f_{q1}}{\partial t} &+ \vec{v}_{1} \cdot \vec{\nabla}_{\vec{r}} f_{q1} &= -\frac{1}{2E_{1}} \int \frac{d^{3}p_{2}}{(2\pi)^{3}2E_{2}} \frac{d^{3}p_{3}}{(2\pi)^{3}2E_{3}} \frac{d^{3}p_{4}}{(2\pi)^{3}2E_{4}} (2\pi)^{4} \delta^{4}(p_{1}+p_{2}-p_{3}-p_{4}) \\ &\times \left\{ g_{G} \mid \mathcal{M}_{ug \rightarrow ug} \mid^{2} \left[ f_{q1}f_{g2}(1-f_{q3})(1+f_{g4}) - f_{q3}f_{g4}(1-f_{q1})(1+f_{g2}) \right] \right. \\ &+ g_{Q}(\frac{1}{2} \mid \mathcal{M}_{uu \rightarrow uu} \mid^{2} + \mid \mathcal{M}_{ud \rightarrow ud} \mid^{2} + \mid \mathcal{M}_{u\bar{u} \rightarrow u\bar{u}} \mid^{2} + \mid \mathcal{M}_{u\bar{d} \rightarrow u\bar{d}} \mid^{2}) \\ &\times \left[ f_{q1}f_{q2}(1-f_{q3})(1-f_{q4}) - f_{q3}f_{q4}(1-f_{q1})(1-f_{q2}) \right] \right\} \\ &- \frac{1}{2E_{1}} \int \frac{d^{3}p_{2}}{(2\pi)^{3}2E_{2}} \frac{d^{3}p_{3}}{(2\pi)^{3}2E_{3}} \frac{d^{3}p_{4}}{(2\pi)^{3}2E_{5}} \frac{d^{3}p_{6}}{(2\pi)^{3}2E_{6}} \\ &\times (2\pi)^{4} \delta^{4}(p_{1}+p_{2}+p_{3}-p_{4}-p_{5}-p_{6}) \left\{ \frac{g_{G}^{2}}{4} \mid \mathcal{M}_{ug \rightarrow ugg} \mid^{2} \\ &\times \left[ f_{q1}f_{g2}f_{g3}(1-f_{q4})(1+f_{g5})(1+f_{g6}) - f_{q4}f_{g5}f_{g6}(1-f_{q1})(1+f_{g2})(1+f_{g3}) \right] \\ &+ g_{Q}g_{G}(\frac{1}{2} \mid \mathcal{M}_{uug \rightarrow uug} \mid^{2} + \mid \mathcal{M}_{udg \rightarrow udg} \mid^{2} + \mid \mathcal{M}_{u\bar{u}\bar{g} \rightarrow u\bar{u}\bar{g}} \mid^{2} + \mid \mathcal{M}_{u\bar{d}\bar{g} \rightarrow u\bar{d}\bar{g}} \mid^{2}) \\ &\times \left[ f_{q1}f_{q2}f_{g3}(1-f_{q4})(1-f_{q5})(1+f_{g6}) - f_{q4}f_{q5}f_{g6}(1-f_{q1})(1-f_{q2})(1+f_{g3}) \right] \end{split}$$

$$+g_{Q}^{2}\left[\frac{1}{12} \mid \mathcal{M}_{uuu \to uuu} \mid^{2} + \frac{1}{4}\left(\mid \mathcal{M}_{uud \to uud} \mid^{2} + \mid \mathcal{M}_{udu \to udu} \mid^{2}\right) + \frac{1}{4} \mid \mathcal{M}_{udd \to udd} \mid^{2} \\ + \frac{1}{2} \mid \mathcal{M}_{uu\bar{u} \to uu\bar{u}} \mid^{2} + \frac{1}{2} \mid \mathcal{M}_{uu\bar{d} \to uu\bar{d}} \mid^{2} + \mid \mathcal{M}_{ud\bar{u} \to ud\bar{u}} \mid^{2} + \mid \mathcal{M}_{ud\bar{d} \to ud\bar{d}} \mid^{2} \\ + \frac{1}{4} \mid \mathcal{M}_{u\bar{u}\bar{u} \to u\bar{u}\bar{u}} \mid^{2} + \frac{1}{2}\left(\mid \mathcal{M}_{u\bar{u}\bar{d} \to u\bar{u}\bar{d}} \mid^{2} + \mid \mathcal{M}_{u\bar{d}\bar{u} \to ud\bar{u}} \mid^{2}\right) + \frac{1}{4} \mid \mathcal{M}_{u\bar{d}\bar{d} \to ud\bar{d}} \mid^{2} \\ \times \left[f_{q1}f_{q2}f_{q3}(1 - f_{q4})(1 - f_{q5})(1 - f_{q6}) - f_{q4}f_{q5}f_{q6}(1 - f_{q1})(1 - f_{q2})(1 - f_{q3})\right]\right\},$$

$$(2)$$

where  $\vec{v}_1$  is the parton velocity;  $g_G(g_Q)$  is the gluon (quark) color-spin degeneracy factor;  $p_i(i = 1, \dots, 6)$  denote the four-momenta of initial and final partons;  $E_i$  is the energy component of  $p_i$ . In equation (1)  $\mathcal{M}_{gd\bar{d}\to gd\bar{d}}$  is the amplitude for the elastic scattering of a gluon, a down-quark and a down-antiquark; in equation (2)  $\mathcal{M}_{ud\bar{u}\to ud\bar{u}}$  is the amplitude for the elastic scattering of an up-quark, a down-antiquark and an up-antiquark. Other notations can be similarly understood. The transport equations for down-quark matter, up-antiquark matter and down-antiquark matter can be established just as the equation for up-quark matter.

We take the following replacement in the transport equations

$$\begin{split} f_{g1}f_{g2}(1+f_{g3})(1+f_{g4}) - f_{g3}f_{g4}(1+f_{g1})(1+f_{g2}) &\rightarrow f_{g1}f_{g2} - f_{g3}f_{g4}, \\ f_{g1}f_{q2}(1+f_{g3})(1-f_{q4}) - f_{g3}f_{q4}(1+f_{g1})(1-f_{q2}) &\rightarrow f_{g1}f_{q2} - f_{g3}f_{q4}, \\ f_{g1}f_{g2}f_{g3}(1+f_{g4})(1+f_{g5})(1+f_{g6}) - f_{g4}f_{g5}f_{g6}(1+f_{g1})(1+f_{g2})(1+f_{g3}) &\rightarrow f_{g1}f_{g2}f_{g3} - f_{g4}f_{g5}f_{g6}, \\ f_{g1}f_{g2}f_{q3}(1+f_{g4})(1+f_{g5})(1-f_{q6}) - f_{g4}f_{g5}f_{q6}(1+f_{g1})(1+f_{g2})(1-f_{q3}) &\rightarrow f_{g1}f_{g2}f_{q3} - f_{g4}f_{g5}f_{q6}, \\ f_{g1}f_{q2}f_{q3}(1+f_{g4})(1-f_{q5})(1-f_{q6}) - f_{g4}f_{g5}f_{q6}(1+f_{g1})(1-f_{q2})(1-f_{q3}) &\rightarrow f_{g1}f_{g2}f_{q3} - f_{g4}f_{g5}f_{q6}, \\ f_{g1}f_{g2}(1-f_{q3})(1+f_{g4}) - f_{g3}f_{g4}(1-f_{q1})(1+f_{g2}) &\rightarrow f_{q1}f_{g2} - f_{q3}f_{g4}, \\ f_{q1}f_{q2}(1-f_{q3})(1-f_{q4}) - f_{q3}f_{q4}(1-f_{q1})(1-f_{q2}) &\rightarrow f_{q1}f_{q2} - f_{q3}f_{q4}, \\ f_{g1}f_{g2}f_{g3}(1-f_{g4})(1+f_{g5})(1+f_{g6}) - f_{g4}f_{g5}f_{g6}(1-f_{q1})(1+f_{g2})(1+f_{g3}) &\rightarrow f_{g1}f_{g2}f_{g3} - f_{g4}f_{g5}f_{g6}, \\ f_{g1}f_{g2}f_{g3}(1-f_{g4})(1+f_{g5})(1+f_{g6}) - f_{g4}f_{g5}f_{g6}(1-f_{q1})(1+f_{g2}) &\rightarrow f_{q1}f_{g2} - f_{q3}f_{g4}, \\ f_{g1}f_{g2}f_{g3}(1-f_{g4})(1+f_{g5})(1+f_{g6}) - f_{g4}f_{g5}f_{g6}(1-f_{q1})(1+f_{g2})(1+f_{g3}) &\rightarrow f_{g1}f_{g2}f_{g3} - f_{g4}f_{g5}f_{g6}, \\ f_{g1}f_{g2}f_{g3}(1-f_{g4})(1+f_{g5})(1+f_{g6}) - f_{g4}f_{g5}f_{g6}(1-f_{g1})(1+f_{g2})(1+f_{g3}) &\rightarrow f_{g1}f_{g2}f_{g3} - f_{g4}f_{g5}f_{g6}, \\ f_{g1}f_{g2}f_{g3}(1-f_{g1})(1+f_{g2})(1+f_{g2})(1$$

 $\begin{aligned} f_{q1}f_{g2}f_{g3}(1-f_{q4})(1+f_{g5})(1+f_{g6}) - f_{q4}f_{g5}f_{g6}(1-f_{q1})(1+f_{g2})(1+f_{g3}) &\rightarrow f_{q1}f_{g2}f_{g3} - f_{q4}f_{g5}f_{g6}, \\ f_{q1}f_{q2}f_{g3}(1-f_{q4})(1-f_{q5})(1+f_{g6}) - f_{q4}f_{q5}f_{g6}(1-f_{q1})(1-f_{q2})(1+f_{g3}) &\rightarrow f_{q1}f_{q2}f_{g3} - f_{q4}f_{q5}f_{g6}, \\ f_{q1}f_{q2}f_{q3}(1-f_{q4})(1-f_{q5})(1-f_{q6}) - f_{q4}f_{q5}f_{q6}(1-f_{q1})(1-f_{q2})(1-f_{q3}) &\rightarrow f_{q1}f_{q2}f_{q3} - f_{q4}f_{q5}f_{q6}, \\ f_{q1}f_{q2}f_{q3}(1-f_{q4})(1-f_{q5})(1-f_{q6}) - f_{q4}f_{q5}f_{q6}(1-f_{q1})(1-f_{q2})(1-f_{q3}) &\rightarrow f_{q1}f_{q2}f_{q3} - f_{q4}f_{q5}f_{q6}, \\ f_{q1}f_{q2}f_{q3}(1-f_{q4})(1-f_{q5})(1-f_{q6}) - f_{q4}f_{q5}f_{q6}(1-f_{q1})(1-f_{q2})(1-f_{q3}) &\rightarrow f_{q1}f_{q2}f_{q3} - f_{q4}f_{q5}f_{q6}. \end{aligned}$ 

Then, any Monte Carlo method can be applied to solve the equations. We generate 1500 gluons, 250 up-quarks, 250 down-quarks, 250 up-antiquarks and 250 down-antiquarks within -0.3 < z < 0.3 fm in the longitudinal direction and r < 6.4 fm in the transverse direction from HIJING for central Au-Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. The number of partons moving in the longitudinal direction is much larger than the number of partons moving in the transverse direction. The anisotropically distributed partons undergo elastic 2-to-2 scattering [14, 15] and elastic 3-to-3 scattering to eventually become isotropic in momentum space (i.e. become a thermal state) at a time. But from gluon matter to quark matter the time is different, i.e. the thermalization time of gluon matter differs from the thermalization time of quark matter. At t = 0.52 fm/c the gluon momentum distribution meets the Jüttner distribution

$$f_g(\vec{p}) = \frac{\lambda_g}{\mathrm{e}^{|\vec{p}|/T} - \lambda_g},\tag{3}$$

with the temperature T = 0.52 GeV and the gluon fugacity  $\lambda_g = 0.328$  that label the thermal state of gluon matter. The corresponding thermalization time is 0.32 fm/c. At t = 0.86 fm/c the quark momentum distribution meets the Jüttner distribution

$$f_q(\vec{p}) = \frac{\lambda_q}{\mathrm{e}^{|\vec{p}|/T} + \lambda_q},\tag{4}$$

with T = 0.46 GeV and the quark fugacity  $\lambda_q = 0.143$  that label the thermal state of quark matter. The corresponding thermalization time is 0.66 fm/c. The thermalization time of quark matter is larger than the one of gluon matter.

# 4. Conclusions

The elastic 2-to-2 scattering and the elastic 3-to-3 scattering lead to the rapid creation of temperature or the early thermalization of quark-gluon matter. Solutions of the transport equations offer the thermalization times and the thermal states of gluon matter and quark matter. Gluon matter thermalizes faster than quark matter.

### Acknowledgments

This work was supported by the National Natural Science Foundation of China under grant no 11175111.

### References

- [1] Xu X-M 2006 Eur. Phys. J. A 29 39
- [2] Ackermann K H et al (STAR Collaboration) 2001 Phys. Rev. Lett. 86 402
- [3] Kolb P F, Huovinen P, Heinz U and Heiselberg H 2001 Phys. Lett. B 500 232
- [4] Teaney D, Lauret J and Shuryak E V 2001 A hydrodynamic description of heavy ion collisions at the SPS and RHIC Preprint nucl-th/0110037
- [5] Shuryak E 1992 Phys. Rev. Lett. 68 3270
- [6] Wong S M H 1996 Phys. Rev. C 54 2588
- [7] Shin G R and Müller B 2003 J. Phys. G: Nucl. Part. Phys. 29 2485
- [8] Blaizot J-P, Gelis F, Liao J, McLerran L and Venugopalan R 2012 Thermalization and Bose-Einstein condensation in overpopulated plasmas *Preprint* arXiv:1210.6838
- [9] Xu X-M, Sun Y, Chen A-Q and Zheng L 2004 Nucl. Phys. A 744 347
- $[10]\,$  Xu X-M, Peng R and Weber H J 2005 Phys. Lett. B 62968
- [11] Xu X-M, Shen Z-Y, Ye Z-C and Xu W-J 2013 Phys. Rev. C in press
- [12] Xu X-M, Ma C-C, Chen A-Q and Weber H J 2007 Phys. Lett. B 645 146
- [13] Xu X-M and Xu L-S 2010 J. Phys. G: Nucl. Part. Phys. 37 115003
- [14] Cutler R and Sivers D 1978 Phys. Rev. D 17 196
- [15] Combridge B L, Kripfganz J and Ranft J 1977 Phys. Lett. B 70 234