TETRAHEDRON EQUATION AND QUANTUM R MATRICES FOR INFINITE DIMENSIONAL MODULES OF $U_q(A_1^{(1)})$ AND $U_q(A_2^{(2)})$

ATSUO KUNIBA AND MASATO OKADO

Abstract

From the q-oscillator solution to the tetrahedron equation associated with a quantized coordinate ring, we construct solutions to the Yang-Baxter equation by applying a reduction procedure formulated earlier by S. Sergeev and the first author. The results are identified with the quantum R matrices for the infinite dimensional modules of $U_q(A_1^{(1)})$ and $U_q(A_2^{(2)})$ corresponding to an affinization of Verma modules of their subalgebras isomorphic to $U_q(sl_2)$ and $U_{q^4}(sl_2)$.

1. INTRODUCTION

The tetrahedron equation [13] is a 3 dimensional (3d) extension of the Yang-Baxter equation [1]. Among its many formulations, the homogeneous version of vertex type is a quartic equation on the matrix \mathcal{R} acting on the tensor cube of a vector space F representing the "single spin" states. See (2.1). The tetrahedron equation possesses the structure that if one of the components in $\mathcal{R} \in \operatorname{End}(F^{\otimes 3})$ is evaluated away appropriately, the resulting object in $\operatorname{End}(F^{\otimes 2})$ satisfies the Yang-Baxter equation. The eliminated space, let us call it the 3rd component, serves as the internal degrees of freedom of the local spins in the resulting 2d vertex model. This reduction works naturally also for the compositions of \mathcal{R} 's in the 3rd component, which implies that a single solution to the tetrahedron equation generates an infinite sequence of solutions to the Yang-Baxter equation. They correspond to the optional size in the 3rd direction of the 3d lattice on which an integrable vertex model associated with \mathcal{R} is defined. This kind of connection between the 2d and 3d integrable systems has been studied from various viewpoints since [2] for example. In particular, it has been shown recently that the reduction scheme reproduces the conventional quantum R matrices for some reducible representations of $U_q(A_n^{(1)})$ [4] and the spin representations of $U_q(B_n^{(1)}), U_q(D_n^{(1)}), U_q(D_{n+1}^{(2)})$ [10]. In the former the reduction of the 3rd space is due to the trace, whereas in the latter it is done by taking the matrix elements between special "boundary vectors". It is an open problem to clarify the grand picture of such 2d-3d connections. See Section 5 for a unified formulation of the problem under which the preceding results [4, 10] are allocated and explained more precisely.

In this paper we add a further solution of the problem as a modest step toward a thorough understanding of the 2d-3d connection. We apply the reduction scheme based on the special boundary vectors [10] to the solution \mathcal{R} of the tetrahedron equation studied in [6, 4, 3, 12, 8]¹. It acts on the tensor cube of the *q*-oscillator Fock space $F = \bigoplus_{m\geq 0} \mathbb{Q}(q)|m\rangle$. It is the same \mathcal{R} as the one used in the trace construction in [4]. For simplicity we concentrate on the simplest "single-site" situation along the 3rd direction. Then the resulting solution to the Yang-Baxter equation is the linear operator acting on $F \otimes F$, which defines an integrable 2d vertex model accommodating the *q*-oscillator Fock states on each edge. Besides its explicit formula (Proposition 2), our main result is the identification with the quantum R matrices for certain infinite dimensional representations of $U_q(A_1^{(1)})$ and $U_q(A_2^{(2)})$ under some specialization (Theorem 10). These representations are constructed from Verma modules of the subalgebras isomorphic to $U_q(sl_2)$ and $U_{q^4}(sl_2)$, respectively. We find it

¹ See Section 2 for more accounts on the origin of the solution.

ATSUO KUNIBA AND MASATO OKADO

natural to encounter the rank 1 quantum affine algebras since our construction corresponds to the single-site situation. Nonetheless being able to capture the both of them is gratifying. It stems from the choices of the two kinds of boundary vectors to which our construction can be applied. A similar feature has been observed in the earlier work [10, Remark 7.2] where the quantum R matrices for $B_n^{(1)}, D_n^{(1)}$ and $D_{n+1}^{(2)}$ have been covered from a single 3d L operator. The layout of the paper is as follows. In Section 2 we recall the reduction procedure based on the

The layout of the paper is as follows. In Section 2 we recall the reduction procedure based on the boundary vectors [10] along the simplest one-site situation. An explicit formula of the new solution to the Yang-Baxter equation is given in Proposition 2. In Section 3 we give the infinite dimensional representations of $U_q(A_1^{(1)})$ and $U_q(A_2^{(2)})$ relevant to our issue. The quantum R matrices for them are described in terms of spectral decompositions. Section 4 contains the main result of the paper giving the precise relation between the solutions to the Yang-Baxter equation obtained in Sections 2 and 3. In Section 5 we formulate a general class of problems that include the relevant results of this paper and [4, 10] as a future direction of researches. Some further results are announced. A full solution of them will bring an important insight into the 2d-3d connection in integrable systems in a broad sense.

In this paper we shall explain the basic construction and only state the main results. The detail of the derivation and generalization to more general setting (see Remark 1 and Section 5) will be given elsewhere. Throughout the paper we assume that q is generic and use the following notations:

$$(z;q)_m = \prod_{j=1}^m (1 - zq^{j-1}), \quad (q)_m = (q;q)_m, \quad \binom{m}{n}_q = \frac{(q)_m}{(q)_n(q)_{m-n}} \text{ (0 unless } 0 \le n \le m),$$
$$[m] = [m]_q = \frac{q^m - q^{-m}}{q - q^{-1}}, \quad [m]_q! = \prod_{j=1}^m [j]_q, \quad \{x\} = \frac{x - x^{-1}}{q - q^{-1}}, \quad (n)_+ = \max(n,0).$$

2. Reducing the tetrahedron equation to the Yang-Baxter equation

Let F be a vector space and $\mathcal{R} \in \text{End}(F^{\otimes 3})$. Consider the tetrahedron equation, which is an equality in $\text{End}(F^{\otimes 6})$:

$$\mathcal{R}_{124}\mathcal{R}_{135}\mathcal{R}_{236}\mathcal{R}_{456} = \mathcal{R}_{456}\mathcal{R}_{236}\mathcal{R}_{135}\mathcal{R}_{124},\tag{2.1}$$

where \mathcal{R}_{ijk} acts as \mathcal{R} on the i, j, k th components from the left in the tensor product $F^{\otimes 6}$.

We recall the prescription in [10] that reduces a solution to the tetrahedron equation to the one for the Yang-Baxter equation. We restrict ourselves to the simplest "single-site" situation relevant to the present paper. See [10] for a more general treatment.

Suppose there are vectors

$$|\chi_s(x,y)\rangle = |\chi_s(x)\rangle \otimes |\chi_s(xy)\rangle \otimes |\chi_s(y)\rangle \in F \otimes F \otimes F,$$
(2.2)

where x, y are extra (spectral) parameters such that

$$\Re|\chi_s(x,y)\rangle = |\chi_s(x,y)\rangle.$$
(2.3)

The index s is a label of possibly more than one such vectors. Suppose also similar vectors exist in the dual space:

$$\langle \chi_s(x,y)| = \langle \chi_s(x)| \otimes \langle \chi_s(xy)| \otimes \langle \chi_s(y)| \in F^* \otimes F^* \otimes F^*,$$
(2.4)

with the property

$$\langle \chi_s(x,y) | \mathcal{R} = \langle \chi_s(x,y) |.$$
(2.5)

Then evaluating the tetrahedron equation (2.1) between $\langle \chi_s(x,y) |$ and $|\chi_t(1,1)\rangle^2$ on the 4,5,6 th components, one gets the Yang-Baxter equation

$$\mathcal{R}_{12}(x)\mathcal{R}_{13}(xy)\mathcal{R}_{23}(y) = \mathcal{R}_{23}(y)\mathcal{R}_{13}(xy)\mathcal{R}_{12}(x) \in \mathrm{End}(F\otimes F\otimes F),$$
(2.6)

²In general $|\chi_t(x', y')\rangle$ can be used. However, in our examples treated later such a freedom is absorbed into elsewhere.

where indices again signify the non trivially acting components in $F^{\otimes 3}$, and

$$\mathcal{R}_{12}(z) = \langle \chi_s(z) | \mathcal{R}_{123} | \chi_t(1) \rangle \in \operatorname{End}(F \otimes F) \otimes 1$$
(2.7)

for example. Here the bracket means the evaluation with respect to the 3rd component. Denoting (2.7) just by $\mathcal{R}(z) \in \operatorname{End}(F \otimes F)$, it is also convenient to introduce

$$\dot{\mathfrak{R}}(z) = \varrho(z) P \,\mathfrak{R}(z), \tag{2.8}$$

where $P(u \otimes v) = v \otimes u$ is the linear operator exchanging the components and $\rho(z)$ is an arbitrary scalar function. Then the Yang-Baxter equation takes another familiar form:

$$(\hat{\mathfrak{X}}(x)\otimes 1)(1\otimes \hat{\mathfrak{X}}(xy))(1\otimes \hat{\mathfrak{X}}(y)) = (1\otimes \hat{\mathfrak{X}}(y))(\hat{\mathfrak{X}}(xy)\otimes 1)(1\otimes \hat{\mathfrak{X}}(x)).$$
(2.9)

Note the degree of freedom to choose s and t in (2.7) although it has temporarily been suppressed in the notation. In fact it will allow us to cover the quantum affine algebras for $A_1^{(1)}$ and $A_2^{(2)}$ in our main Theorem 10.

Now we proceed to a concrete realization of the above scheme in this paper. We will always take F to be an infinite dimensional space $F = \bigoplus_{m\geq 0} \mathbb{Q}(q)|m\rangle$. The dual space will be denoted by $F^* = \bigoplus_{m\geq 0} \mathbb{Q}(q)\langle m|$ with the bilinear pairing $\langle m|n\rangle = (q^2)_m \delta_{m,n}^3$. For simplicity vectors like $|i\rangle \otimes |j\rangle \otimes |k\rangle \in F^{\otimes 3}$ and $\langle i| \otimes \langle j| \in (F^*)^{\otimes 2}$ etc. will be abbreviated to $|i, j, k\rangle$ and $\langle i, j|$ etc.

The solution \mathcal{R} of the tetrahedron equation we are concerned with is the one obtained as the intertwiner of the quantum coordinate ring $A_q(sl_3)$ [6]⁴, which was also found from a quantum geometry consideration in a different gauge including square roots [4, 3]. They were shown to be essentially the same object and to constitute the solution of the 3d reflection equation [8]. It can also be identified with the transition matrix of the PBW bases of the nilpotent subalgebra of $U_q(sl_3)$ [12, 9]. Here we simply call it 3d \mathcal{R} . It is given by

$$\Re|i,j,k\rangle = \sum_{a,b,c} \Re^{a,b,c}_{i,j,k}|a,b,c\rangle,$$
(2.10)

$$\mathcal{R}_{i,j,k}^{a,b,c} = \delta_{i+j}^{a+b} \delta_{j+k}^{b+c} \sum_{\lambda+\mu=b} (-1)^{\lambda} q^{i(c-j)+(k+1)\lambda+\mu(\mu-k)} \frac{(q^2)_{c+\mu}}{(q^2)_c} \binom{i}{\mu}_{q^2} \binom{j}{\lambda}_{q^2},$$
(2.11)

where $\delta_n^m = \delta_{m,n}$ just to save the space. The sum (2.11) is over $\lambda, \mu \geq 0$ satisfying $\lambda + \mu = b$, which is also bounded by the condition $\mu \leq i$ and $\lambda \leq j$. The formula (2.11) is taken from [8, eq.(2.20)], where a proof of the property $(q^2)_a(q^2)_b(q^2)_c \mathcal{R}_{i,j,k}^{a,b,c} = (q^2)_i(q^2)_j(q^2)_k \mathcal{R}_{a,b,c}^{i,j,k}$ was also included. Note that the elements of 3d \mathcal{R} are polynomials in q alone, and the spectral parameter z comes into the game only through the reduction (2.7). Thus the notations $\mathcal{R}(z)$ and $\check{\mathcal{R}}(z)$ automatically distinguish them from the 3d \mathcal{R} , and they should be understood as the solutions to the Yang-Baxter equation. We specify their matrix elements by

$$\Re(z)|i,j\rangle = \sum_{a,b} \Re(z)^{a,b}_{i,j}|a,b\rangle, \quad \check{\Re}(z)|i,j\rangle = \sum_{a,b} \check{\Re}(z)^{a,b}_{i,j}|a,b\rangle, \quad \check{\Re}(z)^{a,b}_{i,j} = \varrho(z)\Re(z)^{b,a}_{i,j}.$$
(2.12)

Let us turn to the vectors $|\chi_s(x,y)\rangle$ and $\langle\chi_s(x,y)|$ in (2.2)–(2.5). We use two such vectors obtained in [10]. In the present notation they read

$$|\chi_1(z)\rangle = \sum_{m\geq 0} \frac{z^m}{(q)_m} |m\rangle, \quad |\chi_2(z)\rangle = \sum_{m\geq 0} \frac{z^m}{(q^4)_m} |2m\rangle, \tag{2.13}$$

$$\langle \chi_1(z)| = \sum_{m \ge 0} \frac{z^m}{(q)_m} \langle m|, \quad \langle \chi_2(z)| = \sum_{m \ge 0} \frac{z^m}{(q^4)_m} \langle 2m|.$$
 (2.14)

We define the four solutions to the Yang-Baxter equation $\Re(z) = \Re^{s,t}(z) = \Re^{s,t}(z,q)$ (s, t = 1, 2) by the formula (2.7) in which (2.11), (2.13) and (2.14) are substituted. They are the matrices acting

 $^{^3}$ The dual space F^\ast and this pairing will only be used in this section and Section 5.

 $^{^4}$ The formula for it on p194 in [6] contains a misprint unfortunately. Eq. (2.11) here is a correction of it.

on $F \otimes F$ whose elements are given by

$$\mathcal{R}^{s,t}(z)_{i,j}^{a,b} = \sum_{c,k\geq 0} \frac{z^c(q^2)_{sc}}{(q^{s^2})_c(q^{t^2})_k} \mathcal{R}^{a,b,sc}_{i,j,tk}.$$
(2.15)

Due to (2.11) this is zero unless a + b = i + j and the sum is actually a single one due to the constraint b + sc = j + tk. In follows that $\mathcal{R}^{2,2}$ is decomposed as

$$\mathcal{R}^{2,2}(z) = \mathcal{R}^{+,+}(z) \oplus \mathcal{R}^{+,-}(z) \oplus \mathcal{R}^{-,+}(z) \oplus \mathcal{R}^{-,-}(z), \qquad (2.16)$$

$$\mathfrak{R}^{\epsilon_1,\epsilon_2}(z) \in \operatorname{End}(F^{\epsilon_1} \otimes F^{\epsilon_2}), \quad F^{\pm} = \bigoplus_{m \ge 0, (-1)^m = \pm 1} \mathbb{Q}(q) |m\rangle.$$
(2.17)

It implies $\check{\mathbb{X}}^{\epsilon_1,\epsilon_2}(z) : F^{\epsilon_1} \otimes F^{\epsilon_2} \to F^{\epsilon_2} \otimes F^{\epsilon_1}$. For example $\check{\mathbb{X}}^{+,-}(z)$ is just the submatrix of $(\check{\mathbb{X}}^{2,2}(z)_{i,j}^{a,b})$ with the indices a, j restricted to be odd and b, i to be even. Another notable fact is

$$\mathcal{R}^{2,1}(z)_{i,j}^{a,b} = \frac{(q^2)_i (q^2)_j}{(q^2)_a (q^2)_b} z^{\frac{j-b}{2}} \mathcal{R}^{1,2} (z^{\frac{1}{2}})_{a,b}^{i,j},$$
(2.18)

which can easily be derived from the property of $\mathcal{R}^{a,b,c}_{i,j,k}$ mentioned after (2.11). Henceforth we concentrate on $\mathcal{R}^{1,1}(z), \mathcal{R}^{2,2}(z)$ and $\mathcal{R}^{1,2}(z)$ in the rest of the paper.

Remark 1. The prescription [10] applied to the *n*-site setting leads to the four solutions of the Yang-Baxter equation (s, t = 1, 2) acting on $F^{\otimes n} \otimes F^{\otimes n}$ whose elements are given by

$$\mathcal{R}^{s,t}(z)_{\mathbf{i},\mathbf{j}}^{\mathbf{a},\mathbf{b}} = \sum_{c_0,\dots,c_n \ge 0} \frac{z^{c_n}(q^2)_{sc_n}}{(q^{s^2})_{c_n}(q^{t^2})_{c_0}} \prod_{l=1}^n \mathcal{R}^{a_l,b_l,c_l'}_{i_l,j_l,c_{l-1}'}, \qquad c_l' = \begin{cases} tc_0 & l = 0, \\ sc_n & l = n, \\ c_l & \text{otherwise} \end{cases}$$

where $\mathbf{a} = (a_1, \ldots, a_n)$ etc. The quantity (2.15) corresponds to the n = 1 case. In the more general problem formulated in Section 5, this corresponds to $\mathcal{R}^{s,t}(z|\varepsilon_1, \ldots, \varepsilon_n)$ (5.8) with $\varepsilon_1 = \cdots = \varepsilon_n = 0$.

By a direct calculation we have

Proposition 2. For $(s,t) \in \{(1,1), (2,2), (1,2)\}$, the following formula is valid:

$$\begin{aligned} \mathcal{R}^{s,t}(z)_{i,j}^{a,b} &= \delta_{i+j}^{a+b} \, \delta_{(-1)^{(s+1)b}}^{(-1)^{(s+1)b}} \, z^{\varepsilon + \frac{1}{s}(j-b-\varepsilon)_{+}}(1+\varepsilon q) \sum_{m,n,\lambda,\mu,\lambda+\mu=b} (-1)^{(\kappa-1)(t-1)m+n+\lambda} q^{\phi} \\ &\times \left(\frac{|b-j|-\varepsilon}{\kappa}\right)_{q^{\kappa^{2}}} \left(\min(b,j)-\lambda\right)_{q^{2}} \binom{i}{\mu}_{q^{2}} \binom{j}{\lambda}_{q^{2}} \frac{((-1)^{s} z^{\frac{t}{s}} q^{t(\lambda-\mu+i+\kappa m+2n+\varepsilon)+s}; q^{st})_{\infty}}{(z^{\frac{t}{s}} q^{t(\lambda-\mu+i+\kappa m+2n)}; q^{st})_{\infty}}, \end{aligned}$$
(2.19)
$$\phi &= \frac{m}{2} (\kappa^{2}m-\kappa+2) + n(n+2|b-j|+1) - i\min(b,j) + \mu^{2} + \lambda + (\lambda-\mu)(b-j)_{+} \\ &+ \varepsilon (4m+2n+\lambda-\mu+i), \end{aligned}$$
(2.20)

$$\kappa = \begin{cases} 2 & \text{if } (s,t) = (1,2), \ b \ge j, \\ s & \text{otherwise,} \end{cases} \qquad \varepsilon = \begin{cases} 1 & \text{if } (s,t) = (1,2), \ b-j \in 2\mathbb{Z}_{\ge 0} + 1, \\ 0 & \text{otherwise.} \end{cases}$$
(2.21)

The sum in (2.19) is over $m, n, \lambda, \mu \ge 0$ with the constraint $\lambda + \mu = b$. It is a finite sum due to the support property of the q-binomial coefficients. The second Kronecker delta postulates $b \equiv j \mod 2$ when s = 2, which guarantees $(j - b - \varepsilon)_+/s$, $(|b - j| - \varepsilon)/\kappa \in \mathbb{Z}$.

Denoting the scalar function $\varrho(z)$ in (2.8) and (2.12) for $\mathbb{R}^{s,t}(z)$ by $\varrho^{s,t}(z)$, we choose it as

$$\varrho^{s,t}(z) = \left(\frac{\left(z^{\frac{t}{s}}; q^{st}\right)_{\infty}}{\left((-1)^s q^s z^{\frac{t}{s}}; q^{st}\right)_{\infty}}\right)^{\epsilon_1 \epsilon_2},\tag{2.22}$$

where for (s,t) = (2,2), the signs ϵ_1, ϵ_2 are to be taken according to the four components in (2.16). For (s,t) = (1,1) and (1,2), the quantity $\epsilon_1 \epsilon_2$ is to be interpreted as +1. Then it follows from Proposition 2 that the matrix elements of $\check{\mathcal{R}}^{s,t}(z)$ are rational functions of q and z. They are useful for computer checks of the Yang-Baxter equation (2.9). **Example 3.** Let $M_d^{s,t} = (\check{\mathbb{R}}^{s,t}(z)_{j,d-j}^{i,d-i})_{0 \le i,j \le d}$ be the matrix where *i* and *j* are the row and the column indices taking (0,0) at the top left corner. One has $M_0^{s,t} = (1)$ for any s, t.

$$\begin{split} M_1^{1,1} &= \left(\begin{array}{c} \frac{(q+1)z}{qz+1} & \frac{1-z}{qz+1} \\ \frac{q(z-1)}{qz+1} & \frac{q+1}{qz+1} \end{array}\right), \ M_1^{1,2} &= \left(\begin{array}{c} \frac{(q+1)z}{qz+1} & \frac{1-z^2}{qz+1} \\ \frac{q(z-1)}{qz+1} & \frac{(q+1)z}{qz+1} \end{array}\right), \ M_1^{2,2} &= \left(\begin{array}{c} 0 & \frac{1}{1-z} \\ \frac{q}{z-1} & 0 \end{array}\right), \\ M_2^{1,1} &= \left(\begin{array}{c} \frac{(q+1)(q^2+1)z^2}{(qz+1)(zq^2+1)} & -\frac{(q+1)(z-1)z}{(qz+1)(zq^2+1)} & \frac{(q+1)(qz-1)}{(qz+1)(zq^2+1)} \\ \frac{q(q+1)(q^2+1)(zq^2+1)}{(qz+1)(zq^2+1)} & \frac{zq^3+2zq^2-q^2-z^2+q^2+2zq+z}{(qz+1)(zq^2+1)} & -\frac{(q+1)(q^2+1)(zq^2+1)}{(qz+1)(zq^2+1)} \\ \frac{q^2(z-1)(qz-1)}{(qz^2+1)(zq^2+1)} & \frac{q(q+1)(z-1)}{(qz^2+1)(zq^2+1)} & \frac{(q+1)(q^2+1)}{(qz^2+1)(zq^2+1)} \\ \frac{q^2(z-1)(qz-1)(qz+1)}{(qz^2+1)(zq^2+1)} & \frac{zq^4+zq^2+q^2-q^2-q^2+z^2+q^2-q^2+z^2+q^2}{(qz^2+1)(zq^2+1)} & -\frac{q(q+1)(qz+1)(qz+1)}{(qz^2+1)(zq^2+1)} \\ \frac{q^2(z-1)(z+1)(qz-1)(qz+1)}{(qz^2+1)(zq^2+1)} & \frac{q^2(q+1)(z-1)z(z+1)}{(qz^2+1)(zq^2+1)} & -\frac{q(q+1)(qz+1)(zq^2+1)}{(qz^2+1)(zq^2+1)} \\ \frac{q^2(z-1)(z+1)(qz-1)(qz+1)}{(qz^2+1)(zq^2+1)} & \frac{q^2(q+1)(z-1)z(z+1)}{(qz^2+1)(zq^2+1)} & -\frac{q(q+1)(q^2+1)(z-1)z(z+1)}{(qz^2+1)(zq^2+1)} \\ \frac{q^2(z-1)(z+1)(qz-1)(qz+1)}{(qz^2+1)(zq^2+1)} & \frac{q^2(q+1)(z-1)z(z+1)}{(qz^2+1)(zq^2+1)} & -\frac{q(q+1)(q^2+1)(z-1)z(z+1)}{(qz^2+1)(zq^2+1)} \\ M_2^{2,2} &= \left(\begin{array}{c} \frac{(q^2-1)z}{q^2z-1} & 0 & \frac{z^2-1}{q^2z-1} \\ 0 & \frac{q^2-z}{q^2z-1} & 0 \\ \frac{q^2(z-1)}{q^2z-1} & 0 & \frac{q^2-1}{(z-1)(q^4z-1)} \\ 0 & \frac{q^2(q^2-z)}{(z-1)(q^4z-1)} & 0 & \frac{(z-q^2)}{(z-1)(q^4z-1)} \\ \frac{q(q^2(z-1)}{(z-1)(q^4z-1)} & 0 & \frac{q(q^2-1)}{(z-1)(q^4z-1)} \\ \frac{q^3(q^2z-1)}{(z-1)(q^4z-1)} & 0 & \frac{q(q^2-1)}{(z-1)(q^4z-1)} \\ 0 & \frac{q^2(q^2-z)}{(z-1)(q^4z-1)} & 0 \\ \end{array}\right). \end{split}$$

3. Quantum R matrices for infinite dimensional modules

The quantum affine algebras (without derivation operator) $U_q(A_1^{(1)})$ and $U_q(A_2^{(2)})$ are the Hopf algebras generated by $e_i, f_i, k_i^{\pm 1}$ (i = 0, 1) satisfying the relations

$$\begin{aligned} k_i k_i^{-1} &= k_i^{-1} k_i = 1, \quad [k_i, k_j] = 0, \\ k_i e_j k_i^{-1} &= q_i^{a_{ij}} e_j, \quad k_i f_j k_i^{-1} = q_i^{-a_{ij}} f_j, \quad [e_i, f_j] = \delta_{ij} \frac{k_i - k_i^{-1}}{q_i - q_i^{-1}}, \\ \sum_{\nu=0}^{1-a_{ij}} (-1)^{\nu} e_i^{(1-a_{ij}-\nu)} e_j e_i^{(\nu)} &= 0, \quad \sum_{\nu=0}^{1-a_{ij}} (-1)^{\nu} f_i^{(1-a_{ij}-\nu)} f_j f_i^{(\nu)} = 0 \quad (i \neq j), \end{aligned}$$

where $e_i^{(\nu)} = e_i^{\nu}/[\nu]_{q_i}!, f_i^{(\nu)} = f_i^{\nu}/[\nu]_{q_i}!, (q_0, q_1) = (q, q)$ for $A_1^{(1)}$ and (q^4, q) for $A_2^{(2)}$. The $(a_{ij})_{0 \le i,j \le 1}$ is the Cartan matrix:

$$(a_{ij})_{0 \le i,j \le 1} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$$
 for $A_1^{(1)}$, $\begin{pmatrix} 2 & -1 \\ -4 & 2 \end{pmatrix}$ for $A_2^{(2)}$.

We use the coproduct of the form

$$\Delta k_i^{\pm 1} = k_i^{\pm 1} \otimes k_i^{\pm 1}, \quad \Delta e_i = 1 \otimes e_i + e_i \otimes k_i, \quad \Delta f_i = f_i \otimes 1 + k_i^{-1} \otimes f_i.$$

Let us introduce a $U_q(A_1^{(1)})$ -module structure on the space F.

Proposition 4. The following defines a $U_q(A_1^{(1)})$ -module structure on F.

$$e_{1}|m\rangle = -[m]\{\alpha q^{m-1}\}|m-1\rangle, \quad f_{1}|m\rangle = |m+1\rangle, \quad k_{1}|m\rangle = \alpha^{-1}q^{-2m}|m\rangle, e_{0} = xf_{1}, \quad f_{0} = x^{-1}e_{1}, \quad k_{0}^{\pm 1} = k_{1}^{\mp 1},$$
(3.1)

where α and x are nonzero parameters.

For generic α it is irreducible, which will be denoted by $V_x(\alpha)$.

Remark 5. As a module over $U_q(sl_2)$ generated by e_1, f_1, k_1 , the space F is identified with a Verma module, namely, it is generated by f_1 from an eigenvector $|0\rangle$ of k_1 killed by e_1 . Such a module has already been known in [7], although eigenvalues are special there.

Proposition 6. The following defines an irreducible $U_q(A_2^{(2)})$ -module structure on F.

$$e_{0}|m\rangle = -\varepsilon_{0} \frac{[2m][2m-2]}{[4]^{2}}|m-2\rangle, \quad f_{0}|m\rangle = |m+2\rangle, \quad k_{0}|m\rangle = \varepsilon_{0}q^{-4m-2}|m\rangle,$$

$$e_{1}|m\rangle = x|m+1\rangle, \quad f_{1}|m\rangle = \varepsilon_{1}(-1)^{m}x^{-1}\frac{[2m]}{[2]}|m-1\rangle, \quad k_{1}|m\rangle = \varepsilon_{1}(-1)^{m}q^{2m+1}|m\rangle$$
(3.2)

where $\varepsilon_0^2 = \varepsilon_1^2 = 1$ and x is a nonzero parameter.

Remark 7. As a module over $U_{q^4}(sl_2)$ generated by e_0, f_0, k_0 , the space F is decomposed into two components F^+ and F^- in (2.17). Both are Verma modules but with different eigenvalues.

In what follows we will be exclusively concerned with the case $\varepsilon_0 = \varepsilon_1 = 1$, which will be denoted by V_x . Note that we have assigned x (the spectral parameter) to the color 1 generators rather than color 0 for $U_q(A_2^{(2)})$. As the vector spaces, $V_x(\alpha)$ and V_x are the same as F.

The quantum R matrix $\check{R}(z) = \check{R}(z, q | \alpha, \beta)$ for our modules of $U_q = U_q(A_1^{(1)}), U_q(A_2^{(2)})$ is a linear operator

$$\check{R}(z): V_x(\alpha) \otimes V_y(\beta) \to V_y(\beta) \otimes V_x(\alpha) \quad (z = x/y),$$
(3.3)

where α, β dependence should be dropped for $U_q(A_2^{(2)})$. The *R* matrix is characterized by the intertwining relations [5]

$$\Delta(g)\check{R}(z) = \check{R}(z)\Delta(g) \quad (\forall g \in U_q)$$
(3.4)

and the normalization which we choose as $\tilde{R}(z)|0,0\rangle = |0,0\rangle$. Define its matrix elements by

$$\check{R}(z)|i,j\rangle = \sum_{a,b} \check{R}(z)^{a,b}_{i,j}|a,b\rangle, \qquad (3.5)$$

where the sum is over those $a, b \ge 0$ satisfying a + b = i + j due to the weight conservation implied by (3.4) with $g = k_0, k_1$.

We include a description of the quantum R matrices in terms of their spectral decomposition. For $U_q(A_1^{(1)})$ we set

$$v_{\alpha,\beta}^{(d)} = \sum_{j=0}^{d} (-q^{1-d}\beta^{-1})^j \binom{d}{j}_{q^2} \prod_{k=1}^{j} \frac{\{\alpha q^{d-k}\}}{\{\beta q^{k-1}\}} |d-j,j\rangle \in V_x(\alpha) \otimes V_y(\beta).$$
(3.6)

Up to an overall constant, this vector is characterized by the conditions

$$(\Delta k_1)v_{\alpha,\beta}^{(d)} = (\alpha\beta)^{-1}q^{-2d}v_{\alpha,\beta}^{(d)}, \qquad (\Delta e_1)v_{\alpha,\beta}^{(d)} = 0.$$
(3.7)

Proposition 8 $(U_q(A_1^{(1)}) \text{ case})$. The following direct sum decomposition holds as the module over $U_q(sl_2)$ mentioned in Remark 5:

$$V_x(\alpha) \otimes V_y(\beta) = \bigoplus_{d \ge 0} V^{(d)}, \quad V^{(d)} = \bigoplus_{r \ge 0} \mathbb{Q}(q) (\Delta f_1)^r v_{\alpha,\beta}^{(d)}.$$
(3.8)

On each subspace $V^{(d)}$, $\check{R}(z) = \check{R}(z, q | \alpha, \beta)$ acts diagonally as (z = x/y)

$$\check{R}(z)(\Delta f_1)^r v_{\alpha,\beta}^{(d)} = \sigma_{\alpha,\beta}^{(d)}(z)(\Delta f_1)^r v_{\beta,\alpha}^{(d)}, \quad \sigma_{\alpha,\beta}^{(d)}(z) = \left(\frac{\beta z}{\alpha}\right)^d \frac{(\alpha^2; q^2)_d (\alpha\beta z^{-1}; q^2)_d}{(\beta^2; q^2)_d (\alpha\beta z; q^2)_d}.$$
(3.9)

The R matrix satisfies the Yang-Baxter equation (common q-dependence is suppressed)

$$(\tilde{R}(y/z|\beta,\gamma)\otimes 1)(1\otimes \tilde{R}(x/z|\alpha,\gamma))(\tilde{R}(x/y|\alpha,\beta)\otimes 1) = (1\otimes \tilde{R}(x/y|\alpha,\beta))(\tilde{R}(x/z|\alpha,\gamma)\otimes 1)(1\otimes \tilde{R}(y/z|\beta,\gamma)),$$
(3.10)

which is an equality of the maps $V_x(\alpha) \otimes V_y(\beta) \otimes V_z(\gamma) \to V_z(\gamma) \otimes V_y(\beta) \otimes V_x(\alpha)$. The inversion relation $\check{\mathcal{R}}(z|\alpha,\beta)\check{\mathcal{R}}(z^{-1}|\beta,\alpha) = \mathrm{Id}$ is valid.

For $U_q(A_2^{(2)})$ we set

$$u_{\pm}^{(d)} = \sum_{j=0}^{d} (-1)^{j(j\pm 1)/2} q^{j(j-1)} {d \choose j}_{q^4} |j, d-j\rangle \in V_x \otimes V_y.$$
(3.11)

Thus $u^{(0)}_+ = u^{(0)}_- = |0,0\rangle$. They are characterized by the conditions

$$(\Delta k_0)u_{\pm}^{(d)} = q^{-4d-4}u_{\pm}^{(d)}, \quad (\Delta e_0)u_{\pm}^{(d)} = 0, \quad \check{R}(z)u_{\pm}^{(d)} = (\pm 1)^d \sigma^{(d)}(\pm z)u_{\pm}^{(d)}, \tag{3.12}$$

where $\sigma^{(d)}(z)$ is specified in (3.14) and z = x/y.

Proposition 9 $(U_q(A_2^{(2)}) \text{ case})$. The following direct sum decomposition holds as the module over $U_{q^4}(sl_2)$ mentioned in Remark 7 $(V_+^{(0)} = V_-^{(0)})$ is denoted by $V^{(0)}$:

$$V_x \otimes V_y = V^{(0)} \oplus \bigoplus_{d \ge 1} (V_+^{(d)} \oplus V_-^{(d)}), \quad V_{\pm}^{(d)} = \bigoplus_{r \ge 0} \mathbb{Q}(q) (\Delta f_0)^r u_{\pm}^{(d)},$$
(3.13)

On each subspace $V^{(d)}_{\pm}$, $\check{R}(z) = \check{R}(z,q)$ acts diagonally as (z = x/y)

$$\check{R}(z)(\Delta f_0)^r u_{\pm}^{(d)} = (\pm 1)^d \sigma^{(d)}(\pm z)(\Delta f_0)^r u_{\pm}^{(d)}, \quad \sigma^{(d)}(z) = \prod_{m=1}^d \frac{q^{2m-1} + (-1)^{m-1}z}{(-1)^{m-1} + q^{2m-1}z}.$$
(3.14)

The *R* matrix satisfies the Yang-Baxter equation (3.10) without dependence on α, β, γ . The inversion relation $\check{\mathbb{R}}(z)\check{\mathbb{R}}(z^{-1}) = \mathrm{Id}$ is valid.

4. MAIN THEOREM

In Section 2 we derived the solutions $\check{\mathbb{R}}^{s,t}(z) = \check{\mathbb{R}}^{s,t}(z,q)$ to the Yang-Baxter equation by a reduction of the 3d \mathscr{R} . In Section 3 the quantum R matrices of the rank 1 quantum affine algebras were determined in terms of their spectral decompositions. The both of these matrices act on the infinite dimensional space $F \otimes F$. Our main theorem presented below identifies them up to a scalar multiple and a similarity transformation.

Theorem 10. (i) The $\check{\mathbb{R}}^{1,1}(z)$ equals a similarity transformation of the $U_q(A_1^{(1)}) R$ matrix $\check{R}(z,q|\alpha,\beta)$ specialized as

$$\check{\mathcal{R}}^{1,1}(z,q^2)^{a,b}_{i,j} = (-iq)^{j-b}\check{R}(z,q|-iq,-iq)^{a,b}_{i,j},\tag{4.1}$$

where i in -iq means $\sqrt{-1}$ and is unrelated to the matrix indices.

(ii) The components $\check{\mathbb{R}}^{\pm,\pm}(z)$ and $\check{\mathbb{R}}^{\pm,\mp}(z)$ of $\check{\mathbb{R}}^{2,2}(z)$ are proportional to a similarity transformation of the $U_q(A_1^{(1)}) R$ matrix $\check{R}(z,q|\alpha,\beta)$ specialized as

$$\check{\mathfrak{R}}^{\epsilon_1,\epsilon_2}(z,q)^{a,b}_{i,j} = r^{\epsilon_1,\epsilon_2} q^{\bar{j}-\bar{b}} \check{R}(z,q^2|q^{2-\epsilon_1},q^{2-\epsilon_2})^{\bar{a},\bar{b}}_{\bar{i},\bar{j}} \quad (\epsilon_1,\epsilon_2=\pm 1),$$
(4.2)

$$r^{+,+} = 1, \ r^{+,-} = \frac{q}{z-1}, \ r^{-,+} = \frac{1}{1-z}, \ r^{-,-} = \frac{q^2-z}{q^2z-1},$$
 (4.3)

where \bar{n} denotes the largest integer not exceeding $\frac{n}{2}$.

(iii) The $\check{\mathbb{R}}^{1,2}(z)$ equals a similarity transformation of the $U_q(A_2^{(2)})$ R matrix $\check{R}(z,q)$ as

$$\check{R}^{1,2}(z,-q^2)^{a,b}_{i,j} = q^{j-b}\check{R}(z,q)^{a,b}_{i,j}.$$
(4.4)

In (ii) of the theorem, the general formula for $r^{\epsilon_1,\epsilon_2}$ is $\check{\mathbb{R}}^{2,2}(z,q)_{-\epsilon_1,-\epsilon_2}^{-\epsilon_2,-\epsilon_1}$, which can be found in Example 3. The factors of the form p^{j-b} is attributed to the similarity transformation by the operator $1 \otimes K_p$ with $K_p|m\rangle = p^m|m\rangle$ which does not spoil the Yang-Baxter equation. Combined with Proposition 2, Theorem 10 provides an explicit formula for the quantum R matrices of our infinite dimensional modules of the rank 1 quantum affine algebras.

Let $M_d^{s,t}(q)$ be the $(d+1) \times (d+1)$ matrix $M_d^{s,t}$ introduced in Example 3 exhibiting the q-dependence. We close the section with a corollary of Theorem 10 giving the eigenvalues of $M_d^{1,1}(q^2), M_{2d}^{2,2}(q)$ and $M_d^{1,2}(-q^2)^5$.

Corollary 11. (i) The eigenvalues of $M_d^{1,1}(q^2)$ are given by the specialization of $\sigma_{\alpha,\beta}^{(j)}(z)$ (3.9) as

$$\{1, \sigma_{-iq,-iq}^{(1)}(z), \dots, \sigma_{-iq,-iq}^{(d)}(z)\}, \quad \sigma_{-iq,-iq}^{(j)}(z) = \prod_{m=1}^{J} \frac{z+q^{2m}}{1+zq^{2m}}.$$
(4.5)

(ii) The eigenvalues of $M_{2d}^{2,2}(q)$ are given by the specialization of $\sigma_{\alpha,\beta}^{(j)}(z)$ (3.9) as

$$\{1, \tilde{\sigma}^{(1)}(z)^{\times 2}, \dots, \tilde{\sigma}^{(d)}(z)^{\times 2}\}, \quad \tilde{\sigma}^{(j)}(z) = \sigma^{(j)}_{\alpha, \beta}(z)|_{q \to q^2, \, \alpha = \beta = q} = \prod_{m=1}^{j} \frac{z - q^{4m-2}}{1 - zq^{4m-2}}, \tag{4.6}$$

where the superscript $\times 2$ stands for the two-fold degeneracy.

(iii) The eigenvalues of $M_d^{1,2}(-q^2)$ are given by $\sigma^{(j)}(z)$ (3.14) as

$$\{ \pm \sigma^{(1)}(\pm z), \pm \sigma^{(3)}(\pm z), \dots, \pm \sigma^{(d)}(\pm z) \} \quad if \ d \ is \ odd, \\ \{ 1, \sigma^{(2)}(\pm z), \sigma^{(4)}(\pm z), \dots, \sigma^{(d)}(\pm z) \} \quad if \ d \ is \ even.$$
 (4.7)

These results can be directly checked for small d by using Example 3.

5. Generalizations

The result in this paper is regarded as the solution of a special case of a more general problem, which we shall now explain. First we recall the 3d L operator [4] in the form adapted to the present context. Let $F = \bigoplus_{m\geq 0} \mathbb{Q}(q)|m\rangle$ be the Fock space and F^* be its dual as before. Set $V = \mathbb{Q}(q)v_0 \oplus \mathbb{Q}(q)v_1$. By the 3d L operator we mean the following:

$$\mathcal{L} = (\mathcal{L}_{\alpha,\beta}^{\gamma,\delta}) \in \operatorname{End}(V \otimes V \otimes F), \quad \mathcal{L}(v_{\alpha} \otimes v_{\beta} \otimes |m\rangle) = \sum_{\gamma,\delta} v_{\gamma} \otimes v_{\delta} \otimes \mathcal{L}_{\alpha,\beta}^{\gamma,\delta} |m\rangle, \tag{5.1}$$

where there are six nonzero $\mathcal{L}_{\alpha,\beta}^{\gamma,\delta} \in \operatorname{End}(F)$ given by

$$\mathcal{L}_{0,0}^{0,0} = \mathcal{L}_{1,1}^{1,1} = 1, \quad \mathcal{L}_{0,1}^{0,1} = \mathbf{k}, \quad \mathcal{L}_{1,0}^{1,0} = -q\mathbf{k}, \quad \mathcal{L}_{1,0}^{0,1} = \mathbf{a}^{-}, \quad \mathcal{L}_{0,1}^{1,0} = \mathbf{a}^{+}.$$
(5.2)

The operators $\mathbf{a}^{\pm}, \mathbf{k} \in \text{End}(F)$ are called *q*-oscillators and act on *F* by

$$\mathbf{a}^+|m\rangle = |m+1\rangle, \quad \mathbf{a}^-|m\rangle = (1-q^{2m})|m-1\rangle, \quad \mathbf{k}|m\rangle = q^m|m\rangle.$$
 (5.3)

In short, the 3d L operator \mathcal{L} represents a six-vertex model having the q-oscillator valued Boltzmann weights. It satisfies the tetrahedron equation [4]:

$$\mathcal{R}_{1,2,3}\mathcal{L}_{b,c,3}\mathcal{L}_{a,c,2}\mathcal{L}_{a,b,1} = \mathcal{L}_{a,b,1}\mathcal{L}_{a,c,2}\mathcal{L}_{b,c,3}\mathcal{R}_{1,2,3}.$$
(5.4)

This is an equality in $\operatorname{End}(\overset{a}{V} \otimes \overset{b}{V} \otimes \overset{c}{V} \otimes \overset{1}{F} \otimes \overset{2}{F} \otimes \overset{2}{F})$, where $\overset{a}{V}, \overset{b}{V}, \overset{c}{V}$ are the copies of V and $\overset{1}{F}, \overset{2}{F}, \overset{3}{F}$ are the ones for F. The indices of \mathcal{R} and \mathcal{L} signify the components of the tensor product on which these operators act non trivially. Viewed as an equation on \mathcal{R} , (5.4) is equivalent [8] to the intertwining relation of the irreducible representations of the quantized coordinate ring $A_q(sl_3)$ [6] in the sense that the both lead to the same solution given in (2.11) up to an overall normalization.

⁵ The eigenvalues of $M_d^{2,2}(q)$ with odd d is not directly derivable from (3.9) since the matrix consists of the sub-matrices $\check{\mathbb{R}}^{\epsilon_1,\epsilon_2}(z)$ with $\epsilon_1\epsilon_2 = -1$ which correspond, due to (4.2), to the situation $\alpha \neq \beta$ in (3.9).

Next we introduce the notation allowing us to teat \mathcal{R} and \mathcal{L} on an equal footing.

$$W^{(0)} = F, \quad W^{(1)} = V, \quad S^{(0)} = \mathcal{R}, \quad S^{(1)} = \mathcal{L}.$$
 (5.5)

The $S^{(r)}$ acts on $W^{(r)} \otimes W^{(r)} \otimes F$ for r = 0, 1. In what follows the copy of the Fock space F that plays the role analogous to the auxiliary space of the transfer matrices will be designated by "3". Define \mathbf{h}_3 acting on it by $\mathbf{h}_3 |m\rangle = m|m\rangle$.

Let *n* be any positive integer. Consider the copies of $W^{(\varepsilon_i)}$ denoted by $\overset{\alpha_i}{W}^{(\varepsilon_i)}$ and $\overset{\beta_i}{W}^{(\varepsilon_i)}$ for $i = 1, 2, \ldots, n$. We write their tensor product as

$$\mathbf{W}^{(\varepsilon)} = \overset{\alpha_1}{W}^{(\varepsilon_1)} \otimes \cdots \otimes \overset{\alpha_n}{W}^{(\varepsilon_n)} \text{ for } \varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n),$$
(5.6)

and similarly for $\mathbf{W}^{(\varepsilon)}$ and $\mathbf{W}^{(\varepsilon)}$. The labels α, β and γ of the copies are put just for distinction and these spaces are the same as $W^{(\varepsilon_1)} \otimes \cdots \otimes W^{(\varepsilon_n)}$ as vector spaces.

We introduce the five families of R matrices each consisting of the 2^n members labeled with $(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_n) \in \{0, 1\}^n$ as follows:

$$\mathcal{R}^{\mathrm{tr}}(z|\varepsilon_1,\varepsilon_2,\ldots,\varepsilon_n) = \mathrm{Tr}_3\big(z^{\mathbf{h}_3} \mathcal{S}_{\alpha_1,\beta_1,3}^{(\varepsilon_1)} \mathcal{S}_{\alpha_2,\beta_2,3}^{(\varepsilon_2)} \cdots \mathcal{S}_{\alpha_n,\beta_n,3}^{(\varepsilon_n)}\big),\tag{5.7}$$

$$\mathcal{R}^{s,t}(z|\varepsilon_1,\varepsilon_2,\ldots,\varepsilon_n) = \langle \chi_s(z)|\mathcal{S}^{(\varepsilon_1)}_{\alpha_1,\beta_1,3} \mathcal{S}^{(\varepsilon_2)}_{\alpha_2,\beta_2,3}\cdots \mathcal{S}^{(\varepsilon_n)}_{\alpha_n,\beta_n,3}|\chi_t(1)\rangle \quad (s,t=1,2).$$
(5.8)

Here the bracket is evaluated in the Fock space 3 by $\langle m|m'\rangle = \delta_{m,m'}(q^2)_m$ and the trace is by $\operatorname{Tr}_3(X) = \sum_{m\geq 0} \frac{\langle m|X|m\rangle}{(q^2)_m}$. They are linear operators acting on $\mathbf{W}^{(\varepsilon)} \otimes \mathbf{W}^{(\varepsilon)}$. By the construction and the tetrahedron equations (2.1) and (5.4) together with the properties (2.3) and (2.5), we have

Theorem 12. Denote any one of (5.7) and (5.8) with a fixed $(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_n)$ by $\mathcal{R}_{\alpha,\beta}(z)$. Then it satisfies the Yang-Baxter equation in End $(\overset{\alpha}{\mathbf{W}}^{(\varepsilon)} \otimes \overset{\beta}{\mathbf{W}}^{(\varepsilon)} \otimes \overset{\gamma}{\mathbf{W}}^{(\varepsilon)})$:

$$\mathfrak{R}_{\alpha,\beta}(x)\mathfrak{R}_{\alpha,\gamma}(xy)\mathfrak{R}_{\beta,\gamma}(y) = \mathfrak{R}_{\beta,\gamma}(y)\mathfrak{R}_{\alpha,\gamma}(xy)\mathfrak{R}_{\alpha,\beta}(x).$$
(5.9)

These solutions $\mathcal{R}_{\alpha,\beta}(z)$ to the Yang-Baxter equation are rational functions of the parameter q and the spectral parameter z up to an overall scalar function of z. Thus it is natural to seek their origin in the conventional quantum group theory. We formulate it as

Problem. Find the appropriate quantum affine algebra and its (possibly infinite dimensional) representation by which the $\mathcal{R}_{\alpha,\beta}(z)$ is characterized as the intertwiner of the tensor product up to a normalization. So far it has been studied in the following cases. (The two q's entering $\mathcal{R}_{\alpha,\beta}(z)$ and U_q are not necessarily the same.)

- (1) $\mathcal{R}^{tr}(z|, 0, 0, ..., 0)$ was claimed to be the direct sum $\bigoplus_{J,J' \ge 0} R_{J\omega_1, J'\omega_1}$ of the $U_q(A_{n-1}^{(1)})$ quantum R matrices for the symmetric representations (relative normalization of the summands left unspecified) [4].
- (2) $\mathcal{R}^{s,t}(z|1,1,\ldots,1)$ with (s,t) = (2,1), (2,2) and (1,1) were identified [10] with the quantum R matrices for the spin representations of $U_q(B_n^{(1)}), U_q(D_n^{(1)})$ [11] and $U_q(D_{n+1}^{(2)})$, respectively. It was also suggested [10] that the case (s,t) = (1,2) is the quantum R matrix of the spin representation of $U_q(B_n^{(1)})$ corresponding to the realization of $B_n^{(1)}$ as an affinization of its classical subalgebra D_n rather than the standard B_n .
- (3) $\mathcal{R}^{s,t}(z|0)$ with (s,t) = (1,1), (2,2) (resp. (1,2), (2,1)) are identified with the $U_q(A_1^{(1)})$ (resp. $U_q(A_2^{(2)})$) quantum R matrices for the infinite dimensional representations corresponding to an affinization of the Verma module of their classical subalgebras in this paper.

The full solution of the problem is a feasible task and will shed a valuable insight into the relation between 2d and 3d integrable systems. In fact the s = t = 1 case of the above result (3) can be extended to the *n*-site situation. We have proved that $\mathcal{R}^{1,1}(z|0,0,\ldots,0)$ is the quantum R matrix of $U_q(D_{n+1}^{(2)})$ associated with the affinization of the Verma (q-oscillator) module of its

classical subalgebra $U_q(B_n)$. It is consistent with (4.1) in view of $D_2^{(2)} = A_1^{(1)}$ and is also natural from [10, Remark 7.2].

It is interesting to investigate the effect of the mixture of \mathcal{R} and \mathcal{L} in (5.7) and (5.8) which is firstly formulated here explicitly. We finish by presenting an explicit formula of $\mathcal{R}^{tr}(z|0,1)$ as the simplest example.

Note the decomposition $\mathbf{W}^{(0,1)} = \bigoplus_{d \ge 0} W^{(d)}$, where $W^{(0)} = \mathbb{Q}(q)|0\rangle \otimes v_0$ and $W^{(d)} = \mathbb{Q}(q)|d\rangle \otimes v_0 \oplus \mathbb{Q}(q)|d-1\rangle \otimes v_1$ for $d \ge 1$. Accordingly $\mathcal{R}^{\mathrm{tr}}(z|0,1)$ splits into the direct sum of the matrices acting on $W^{(d)} \otimes W^{(d')}$. It turns out that they are zero unless d = d' = 0 or $dd' \ge 1$. Thus $\mathcal{R}^{\mathrm{tr}}(z|0,1)$ consists of a 1×1 matrix and infinitely many 4×4 matrices corresponding to $\mathrm{End}(W^{(d)} \otimes W^{(d')})$ with $dd' \ge 1$. Explicitly it is expressed as

$$\mathcal{R}^{\rm tr}(z|0,1) = \frac{1}{1-z} \mathrm{Id}_{0,0} \oplus \bigoplus_{d,d' \ge 1} \frac{z^{d'-1} (q^{d-d'+2} z^{-1}; q^2)_{d'-1}}{(q^{d-d'} z; q^2)_{d'+1}} (1 \otimes K^{-1}) \mathcal{R}_{q^d, q^{d'}}(z) (K \otimes 1), \quad (5.10)$$

where $\mathrm{Id}_{0,0}$ denotes the 1×1 identity matrix and $K(|m\rangle \otimes v_{\alpha}) = q^{1-2\alpha}|m\rangle \otimes v_{\alpha}$ causes just a gauge transformation. The $\mathcal{R}_{q^d,q^{d'}}(z) \in \mathrm{End}(W^{(d)} \otimes W^{(d')})$ is given by the specialization of

$$\mathcal{R}_{\mu,\nu}(z) = \begin{pmatrix} z - \mu\nu & 0 & 0 & 0\\ 0 & \nu - \mu z & (1 - \nu^2)z & 0\\ 0 & 1 - \mu^2 & \mu - \nu z & 0\\ 0 & 0 & 0 & 1 - \mu\nu z \end{pmatrix}.$$
(5.11)

By this we mean the linear operator acting as $\xi_0 \otimes \eta_0 \mapsto (z - \mu\nu)\xi_0 \otimes \eta_0$, $\xi_0 \otimes \eta_1 \mapsto (\nu - \mu z)\xi_0 \otimes \eta_1 + (1 - \mu^2)\xi_1 \otimes \eta_0$, etc. in terms of the basis $\xi_i \otimes \eta_j$ of $W^{(d)} \otimes W^{(d')}$ taken as $\xi_i = |d-i\rangle \otimes v_i$ and $\eta_j = |d'-j\rangle \otimes v_j$ for i, j = 0, 1. The case $\mu = \nu$ is known to be the intertwiner of the quantum affine super algebra $U_q(\widehat{sl}(1|1))$. The $\mathcal{R}_{\mu,\nu}(z)$ satisfies the Yang-Baxter equation $\mathcal{R}_{\lambda,\mu}(x)\mathcal{R}_{\lambda,\nu}(xy)\mathcal{R}_{\mu,\nu}(y) = \mathcal{R}_{\mu,\nu}(y)\mathcal{R}_{\lambda,\nu}(xy)\mathcal{R}_{\lambda,\mu}(x)$. Further results including the detailed derivation and the proof of this paper will appear elsewhere.

Acknowledgments

The authors thank Yasuhiko Yamada for collaboration in the previous work, Kailash C. Misra and Yoshihisa Saito for communications on literature and Tatsuya Toyoda for a careful reading of the manuscript. A.K. thanks Vladimir Bazhanov, Vladimir Mangazeev and Sergey Sergeev for kind interest during his stay in Canberra in March 2012. Especially he was benefited from the collaboration with S. Sergeev in [10]. This work is supported by Grants-in-Aid for Scientific Research No. 23340007, No. 24540203 and No. 23654007 from JSPS.

References

- [1] R. J. Baxter, Exactly solved models in statistical mechanics, Dover (2007).
- [2] V. V. Bazhanov and Yu. G. Stroganov, Conditions of commutativity of transfer matrices on a multidimensional lattice, Theor. Math. Phys. 52 685-691 (1982).
- [3] V. V. Bazhanov, V. V. Mangazeev and S. M. Sergeev, Quantum geometry of 3-dimensional lattices, J. Stat. Mech. P07004 (2008).
- [4] V. V. Bazhanov and S. M. Sergeev, Zamolodchikov's tetrahedron equation and hidden structure of quantum groups, J. Phys. A: Math. Theor. 39 3295–3310 (2006).
- [5] M. Jimbo, A q-difference analogue of U(g) and the Yang-Baxter equation, Lett. Math. Phys. 10 63-69 (1985).
- M. M. Kapranov and V. A. Voevodsky, 2-Categories and Zamolodchikov tetrahedron equations, Proc. Symposia in Pure Math. 56 177–259 (1994).
- [7] P. P. Kulish and E. V. Damaskinsky, On the q oscillator and the quantum algebra, $su_q(1, 1)$. J. Phys. A 23 L415–L419 (1990).
- [8] A. Kuniba and M. Okado, Tetrahedron and 3D reflection equations from quantized algebra of functions, J. Phys. A: Math.Theor. 45 (2012) 465206 (27pp).
- [9] A. Kuniba, M. Okado and Y. Yamada, A common structure in PBW bases of the nilpotent subalgebra of $U_q(\mathfrak{g})$ and quantized algebra of functions, SIGMA 9 (2013), 049, 23 pages.
- [10] A. Kuniba and S. Sergeev, Tetrahedron equation and quantum R matrices for spin representations of $B_n^{(1)}, D_n^{(1)}$ and $D_{n+1}^{(2)}$, Commun. Math. Phys. **324** 695-713 (2013).

- [11] M. Okado, Quantum R matrices related to the spin representations of B_n and D_n , Commun. Math. Phys. **134** 467-486 (1990).
- [12] S. M. Sergeev, Tetrahedron equations and nilpotent subalgebras of $\mathcal{U}_q(sl_n)$, Lett. Math. Phys. 83 231–235 (2008).
- [13] A. B. Zamolodchikov, Tetrahedra equations and integrable systems in three-dimensional space, Soviet Phys. JETP 79 641–664 (1980).

E-mail address: atsuo@gokutan.c.u-tokyo.ac.jp

Institute of Physics, Graduate School of Arts and Sciences, University of Tokyo, Komaba, Tokyo 153-8902, Japan

E-mail address: okado@sci.osaka-cu.ac.jp

DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, 3-3-138, SUGIMOTO, SUMIYOSHI-KU, OSAKA, 558-8585, JAPAN