

On conjugacy classes in the Lie group E_8

George Lusztig*


Based on a talk given at the AMS-RMS joint meeting in Alba Iulia, 6/28/2013.

The classification of simple Lie algebras (or Lie groups)/ \mathbf{C} (Killing 1879, a glory of 19-th century mathematics) leads to 4 ∞ series and 5 exceptional groups of which the largest one, E_8 , has dimension 248.

The quantity $\frac{\dim(G)}{\text{rk}(G)^2}$ defined for any simple Lie group G is bounded. It reaches its maximum $\frac{248}{8^2} \cong 4$ for G of type E_8 . This shows that the group of type E_8 is the most noncommutative of all simple Lie groups.

The Lie group E_8 can be obtained from the graph E_8 (to be described below) by a method of Chevalley (1955), simplified using theory of "canonical bases" (1990).

The graph E_8 has 8 vertices and 7 edges. It can be obtained by removing 5 edges from the 1-skeleton of a cube in \mathbf{R}^3 (which has 8 vertices and 12 edges).



Let I be the set of vertices of the graph E_8 . Let V be the free \mathbf{Z} -module with basis $\{\alpha_i; i \in I\}$ and bilinear form:

$$(\alpha_i, \alpha_i) = 2; (\alpha_i, \alpha_j) = -\#(\text{edges joining } i, j) \text{ if } i \neq j.$$

This bilinear form was defined by Korkine-Zolotarev (1873) before Killing and nonconstructively earlier by Smith (1867). The set $R = \{\alpha \in V, (\alpha, \alpha) = 2\}$ (roots) has exactly 240 elements. Its subset $R^+ = R \cap \sum_{i \in I} \mathbf{N}\alpha_i$ has exactly 120 elements.

Let M be the \mathbf{C} -vector with basis $X_\alpha (\alpha \in R)$, $t_i (i \in I)$. For $i \in I$, $\epsilon = 1, -1$, we define linear maps $E_{i,\epsilon} : M \rightarrow M$ by

$$E_{i,\epsilon} X_\alpha = X_{\alpha + \epsilon\alpha_i} \text{ if } \alpha \in R, \alpha + \epsilon\alpha_i \in R,$$

$$E_{i,\epsilon} X_\alpha = 0 \text{ if } \alpha \in R, \alpha + \epsilon\alpha_i \notin R \cup 0,$$

$$E_{i,\epsilon} X_{-\epsilon\alpha_i} = t_i, E_{i,\epsilon} t_j = |(\alpha_i, \alpha_j)| X_{\epsilon\alpha_i}.$$

Let $M^+ = \sum_{\alpha \in R^+} \mathbf{C}X_\alpha \subset M$. We have $\dim M = 248$, $\dim M^+ = 120$.

*Supported in part by National Science Foundation grant DMS-6927854.

Let $G = E_8(\mathbf{C})$ be the subgroup of $GL(M)$ generated by the automorphisms $\exp(\lambda E_{i,\epsilon})$, $i \in I$, $\epsilon = 1, -1$, $\lambda \in \mathbf{C}^*$. (The Lie group E_8 .) Replacing \mathbf{C} by a finite field F_q leads to a finite group $E_8(F_q)$ which, by Chevalley (1955), is simple of order $q^{248} +$ lower powers of q . (The denominators in the exponential do not cause troubles.)

The Weyl group is $W = \{g \in \text{Aut}(V); gR = R\}$. It can be viewed as a limit of $E_8(F_q)$ as $q \rightarrow 1$ hence we can denote it by $E_8(F_1)$. It is a finite group:

$$\sharp(E_8(F_1)) = \lim_{q \rightarrow 1} \frac{\sharp(E_8(F_q))}{(q-1)^8} = 4!6!8!.$$

For a group H , the *conjugacy classes* of H are the orbits of the action $x : g \mapsto xgx^{-1}$ of H on itself. They form a set $cl(H)$.

The main guiding principle of this talk:

The conjugacy classes of $G = E_8(\mathbf{C})$ should be organized according to the conjugacy classes of $W = E_8(F_1)$.

For $w \in W$ let $l(w) = \sharp(R^+ \cap w(R - R^+)) \in \mathbf{N}$ (length).

For $C \in cl(W)$ let $C_{min} = \{w \in C; l : C \rightarrow \mathbf{N}$ reaches minimum at $w\}$.

$C \in cl(W)$ is *elliptic* if $\{x \in V; wx = x\} = 0$ for some/any $w \in C$.

$cl(W)$ has been described by Carter (1972); we have $\sharp cl(W) = 112$, $\sharp\{C \in cl(W); C \text{ elliptic}\} = 30$.

The G -orbit of M^+ in the Grassmannian of 120-dimensional subspaces of M is a closed smooth subvariety \mathcal{B} of dimension 120, the *flag manifold*. The diagonal G -action on $\mathcal{B} \times \mathcal{B}$ has orbits in canonical bijection

$$\mathcal{O}_w \leftrightarrow w$$

with W . (Bruhat 1954, Harish Chandra 1956).

For $w \in W$ let $G_w = \{g \in G; (B, g(B)) \in \mathcal{O}_w \text{ for some } B \in \mathcal{B}\}$.

For $C \in cl(W)$ let $G_C = G_w$ where $w \in C_{min}$; one shows (using results of Geck-Pfeiffer 1993, simplified by He-Nie 2012) that G_C is independent of the choice of w in C_{min} ; also, $G_C \neq \emptyset$, G_C is a union of conjugacy classes. Let

$$\delta_C = \min_{\gamma \in cl(G); \gamma \subset G_C} \dim \gamma,$$

$$\boxed{G_C} = \cup_{\gamma \in cl(G); \gamma \subset G_C, \dim \gamma = \delta_C} \gamma.$$

Then $\boxed{G_C}$ is $\neq \emptyset$, a union of conjugacy classes of fixed dimension, δ_C .

Let $G^{un} = \{g \in G; \text{all eigenvalues of } g : M \rightarrow M \text{ are } 1\}$.

Theorem. (a) $\cup_{C \in cl(W)} \boxed{G_C} = G$.

(b) For any $C, C' \in cl(W)$, $\boxed{G_C}, \boxed{G_{C'}}$ are either equal or disjoint.

(c) For any $C \in cl(W)$, $\boxed{G_C} \cap G^{un}$ is either empty or a single conjugacy class.

(d) If $C \in cl(W)$ is elliptic then $\delta_C = 248 - l(w)$ for any $w \in C_{min}$ and $\boxed{G_C} \cap G^{un}$ is a single conjugacy class.

See arXiv:1305.7168.

The proof uses representation theory of $E_8(F_q)$ (a part of its character table known since 1980's) and computer calculation. (Help with programming in GAP was provided by Gongqin Li.)

The subsets $\boxed{G_C}$ partition G into the *strata* of G . If $\gamma, \gamma' \in cl(G)$ we say: $\gamma \sim \gamma'$ if γ, γ' are contained in the same stratum. If $C, C' \in cl(W)$ we say: $C \sim C'$ if $\boxed{G_C} = \boxed{G_{C'}}$. These are equivalence relations on $cl(G) = cl(E_8(\mathbf{C}))$, $cl(W) = cl(E_8(F_1))$; by the theorem we have canonically

$$cl(E_8(\mathbf{C}))/\sim \leftrightarrow \mathbf{cl}(\mathbf{E}_8(\mathbf{F}_1))/\sim.$$

Examples. If $C = \{1\}$ then $\boxed{G_C} = \{1\}$.

If $C \in cl(W)$ contains all w in W of length 1 then $\boxed{G_C}$ is a single conjugacy class (it has dimension 58.)

Let $C_{cox} = \{w \in W; w \text{ has order } 30\}$; it is a single conjugacy class in W , the Coxeter class. If $C = C_{cox}$ then $G_C = \boxed{G_C}$ is the union of all conjugacy classes of dimension 240 (Steinberg 1965).

For $n = 2, 3, 5$ let \mathbf{c}_n be the unique conjugacy class in G such that $\text{codim}_{\mathbf{c}_n} = 240/n$ and any element of \mathbf{c}_n has order n . One can show: $\mathbf{c}_n \subset \boxed{G_{C_{cox}^{30/n}}}$.

Let $cl_u(E_8(\mathbf{C})) = \{\gamma \in \mathbf{cl}(\mathbf{E}_8(\mathbf{C})); \gamma \in \mathbf{G}^{\text{un}}\}$. Let $cl_u(E_8(\bar{F}_p))$ be the analogous set with \mathbf{C} replaced by \bar{F}_p , p a prime number. We have $\sharp(cl_u(E_8(\mathbf{C}))) = 70$, (Dynkin, Kostant), $\sharp(cl_u(E_8(\bar{F}_p))) = 70 + n$ where $n = 4, 1, 0, 0$, for $p = 2, 3, 5, 7$, (Mizuno 1980).

We have a natural imbedding $j_p : cl_u(E_8(\mathbf{C})) \rightarrow \mathbf{cl}_{\mathbf{u}}(\mathbf{E}_8(\bar{F}_p))$ and $cl(E_8(\mathbf{C}))/\sim \leftrightarrow \cup_{p \text{ prime}} \mathbf{cl}_{\mathbf{u}}(\mathbf{E}_8(\bar{F}_p))$ (union taken using j_p). Hence the number of strata is 75.

The definitions above extend to any simple Lie group G .

If $G = S0_5(\mathbf{C})$, then $cl(W) = \{C_4, C_4^2, C', C'', \{1\}\}$ where C_4 consists of the elements of order 4;

$\boxed{G_{C_4}}$ is the union of classes of dimension 8,

$\boxed{G_{C_4^2}}$ is the union of classes of dimension 6,

$\boxed{G_{C'}}$, $\boxed{G_{C''}}$ are the two conjugacy classes of dimension 4,

$\boxed{G_{\{1\}}} = \{1\}$.