Possible Consistent Extra Time Dimensions in the Early Universe

Patrick L. Nash*

475 Redwood Street Unit 801, San Diego, CA 92103-5864 (Dated: November 16, 2021)

Abstract

Gravity cannot be quantized unless the quantized theory is cast on a manifold whose concomitant number of physical space dimensions and number of physical time dimensions correspond to physical reality, and not simply to the perception of reality. At present, the accepted number of physical time dimensions is dictated more by folklore than by science. True, there are theorems that restrict the number of possible time dimensions to one, but these only apply to rather special scenarios. In this paper we discuss a model of the early universe in which the number of physical time dimensions is four, and formulate Theorem[10.1], which underlies an explanation of why the extra time dimensions do not source unphysical effects.

In this paper we describe a new model of gravitational inflation that is driven by dark energy and "mediated" by a real massless scalar inflaton field φ whose potential is identically equal to zero. The coupled Einstein gravitational and inflaton field equations are formulated on an eightdimensional spacetime manifold of **four space** dimensions and **four time** dimensions. We find explicit solutions to these field equations that exhibit temporal exponential **deflation of three of the four time dimensions**, and then study the dynamics of a massive complex scalar field ψ that propagates on the background ground state Einstein gravitational field to determine whether its quantum fluctuations $\delta \psi$ are stable or unstable. We compute explicit approximate solutions to the $\delta \psi$ field equations that are **stable**, meaning that the quantum fluctuations $\delta \psi$ of the field ψ do not grow exponentially with time. **Instabilities** driven by the momenta associated to the three extra time dimensions do not appear in the physical solutions of the field equations of this model.

PACS numbers: 98.80.Cq, 04.62.+v, 04.50.Kd

Keywords: cosmology, extra time, inflation, deflation, dark energy

^{*}Electronic address: Patrick299Nash@gmail.com

1. INTRODUCTION

Recent Planck 2013 data analysis [1] is in remarkable accord with a flat Λ CDM model with inflation, based upon a spatially flat, expanding universe whose dynamics are governed by General Relativity and sourced by cold dark matter, a cosmological constant Λ , and a slow-roll scalar inflaton field [2–4]. The main predictions of inflationary cosmology are also consistent [5–8] with other recent observational data from important experiments such as WMAP [9] and the Sloan Digital Sky Survey [10–12], to name only two. Also, assuming that the large value of the tensor-to-scalar ratio in the cosmic microwave background radiation reported by the BICEP2 collaboration [13] is correct, then the energy scale of inflation is approximately 2×10^{16} GeV, which is only roughly two orders of magnitude below the Planck scale of $\frac{1.22 \times 10^{19}}{\sqrt{8\pi}}$ GeV.

However, parameterized ACDM inflationary cosmology may not represent fundamental physics. Here we discuss a radical new model of inflation/deflation that overcomes the well known issues of [1] the precise definition of the inflaton potential, [2] the precise definition of the inflaton mass (many models of inflation omit the Higgs field that presumably generates the inflaton mass, or do not explain why the Higgs field develops a non-zero vacuum expectation value at such a high energy) and [3] re-heating. This model is based on the idea that our universe has as many time dimensions as space dimensions.

In this paper we describe a new model of gravitational inflation that is driven by dark energy and "mediated" by a real massless scalar inflaton field φ whose potential is identically equal to zero. The coupled Einstein gravitational and inflaton field equations are formulated on an eight-dimensional spacetime manifold of **four space** dimensions and **four time** dimensions. For the case of a diagonal metric, two periodic solution classes ("ground" and "excited" classes) for the coupled Einstein field equations are obtained that exhibit temporal exponential **deflation of three of the four time dimensions** and temporal exponential inflation of three of the four space dimensions. Moreover, for the ground state solution class the universe does not cool during inflation. Reheating [14] is a non-issue. Moreover, we show that the extra time dimensions do not generally induce exponentially rapid growth of fluctuations in quantum fields that propagate on the ground state solution class.

Comoving coordinates for the two non-inflating/deflating dimensions are chosen to be (x^4, x^8) . The x^4 coordinate corresponds to our universe's observed physical time dimension,

while the x^8 coordinate corresponds to a new spatial dimension that is assumed to be compact with constant unit radius $R_8 \equiv 1$, i.e., the radius equals the Planck length. The value of R_8 can be adjusted to fit experimental data if need be. Dark energy is realized in terms of a cosmological constant Λ , whose values are determined to be quantized in terms of the constant unit radius of the compact spatial x^8 dimension. It is also interesting to note that although the inflaton potential is zero, an effective inflaton potential for the ground state is computed to be $V_{\text{eff}} = \frac{5}{4!} \Lambda \cot(\frac{1}{2}x^8 \sqrt{2\Lambda})^2$.

In this model, after "inflation" the observable physical macroscopic world appears to a classical observer to be a homogeneous, isotropic universe with three space dimensions and one time dimension. Also, under the assumption that an "arrow of time" exists for each timelike dimension, unphysical closed timelike curves are not observed in this model. We also prove that a well known so-called "single time" theorem **does not apply** to our model.

2. ARGUMENTS AGAINST PLURAL TIME DIMENSIONS

Everyone is familiar with arguments against the existence of extra time dimensions. While there are many aspects to this issue, we address two arguments against the existence of extra time dimensions which stand out most clearly[15]:

- A spacetime with extra time dimensions may not carry a spin structure ⇔ spinors cannot be defined on the spacetime;
- 2. Momenta corresponding to the extra time dimensions induce exponentially rapid growth of quantum fluctuations of the field; the universe is unstable. This instability is associated with the very largest momenta (shortest wavelengths).

Issue [1] is not applicable here; it is known that our model spacetime carries a spin structure [16]. The issues raised in [2] may be investigated by studying the field equation for the propagation of a massive complex scalar field ψ on the background gravitational field:

$$0 = \frac{1}{\sqrt{\det(g_{\alpha\beta})}} \partial_{\mu} \left[\sqrt{\det(g_{\alpha\beta})} g^{\mu\nu} \partial_{\nu} \psi \right] - \left(\Lambda m^2 + \zeta R \right) \psi, \qquad (2.1)$$

which is derivable from the Lagrangian

$$L_{\psi} = \sqrt{\det(g_{\alpha\beta})} \left[-g^{\mu\nu}\partial_{\mu}\psi^* \ \partial_{\nu}\psi \ -\psi^* \ \psi \ \left(\Lambda \ m^2 + \zeta \ R\right) \right]. \tag{2.2}$$

Here R is the Ricci scalar, and m and ζ are real input parameters; the factors of Λ are included for convenience. The quantum fluctuations $\delta \psi$ of ψ satisfy a field equation similar to Eq.[2.1], but generally with a different rest mass parameter.

It will be seen that Eq.[2.1] is not separable. Because Eq.[2.1] is not separable, there will not be a simple dispersion relation, per se, relating a physical-frequency ($\omega \leftrightarrow -i\frac{\partial}{\partial x^4} \ln \psi$) to the momentum wave vectors (\vec{k}, \vec{w}) that are defined two paragraphs below. In fact we find decaying quasi-normal mode approximate solutions to Eq.[2.1]. Modes of this class are well known in black hole physics [17–19]: the authors of Ref.[18] discuss the theory of quasi-normal modes of compact objects, including perturbations of (Schwarzschild, Reissner-Nordström, Kerr and Kerr-Newman) black holes; Ref.[17] provides a rigorous definition of quasi-normal modes for a rotating black hole; the author proves that the local energy of linear waves in certain black hole backgrounds decay exponentially once orthogonality to the zero resonance is imposed. In the present early universe model, decaying quasi-normal type modes appear as approximate solutions to Eq.[2.1]; however numerical experiments indicate that the contribution to the complex frequency that is responsible for decay gets smaller as the order of the approximation increases (at least for a finite percentage of the modes). The consequence of this for the unitarity of the quantum field theory is not presently known.

We shall discuss two types of solution to Eq.[2.1], the first with nonzero coupling to the Ricci scalar and the second with $\zeta = 0$. It should be emphasized that the expected Yukawa interaction coupling of the massive complex scalar field ψ with the massless real scalar inflaton field φ is not included here. We do not wish to confuse the production/annhilation of ψ "particles" due to its interaction with the inflaton field with instabilities in the ψ field that are sourced by the momenta associated to the extra time dimensions. Also, it is beyond the scope of the present paper to consider the one-loop effective potentials and concomitant renormalizations of both φ and ψ . These are important questions, but outside the scope of this work.

In Sections 5 and 11 we seek "plane wave" solutions to Eq.[2.1] in terms of comoving wave vectors and coordinates of the form

$$\psi = \Psi(x^4, x^8) e^{i(\vec{k} \cdot \vec{r} - \vec{w} \cdot \vec{R})}, \text{ where}$$

$$\vec{r} = (x^1, x^2, x^3)^T \text{ and } \vec{k} = (k_1, k_2, k_3)^T;$$

$$\vec{R} = (x^5, x^6, x^7)^T \text{ and } \vec{w} = (k_5, k_6, k_7)^T.$$
(2.3)

When discussing these solutions the following shorthand is employed:

$$\Lambda k^{2} = \vec{k} \cdot \vec{k} = k_{1}^{2} + k_{2}^{2} + k_{3}^{2};$$

$$\Lambda w^{2} = \vec{w} \cdot \vec{w} = k_{5}^{2} + k_{6}^{2} + k_{7}^{2}.$$
(2.4)

To summarize, in this paper we find explicit solutions of the coupled Einstein-inflaton field equations, which are then substituted into Eq.[2.1]; solutions to the latter equation are then determined for which the effects of multiple time dimensions are stable, meaning that quantum fluctuations $\delta\psi$ of the ψ field do not grow exponentially with time. Exponentially rapid growth of quantum fluctuations $\delta\psi$ of the ψ field cause perturbation theory to break down when the norm of the term(s) involving $\delta\psi$ become of order one.

2.1. Single-Time Theorem (STT)

The following "Single-Time" Theorem [20] has been asserted, incorrectly, to argue that extra time dimensions cannot exist in viable physical models of the universe. To demonstrate that this theorem does not apply to the model in this paper, we need only quote the statement of the STT and highlight two points in its proof that clearly are not satisfied in our model. We quote the author of the STT:

"Theorem 2 (Single-Time Theorem) [20]. Any BLACK HOLE solution with k > 0 contains precisely one coordinate t such that $g_{tt} = 0$ at the horizon."

"The starting point is ... the model action for D-dimensional gravity with several scalar dilatonic fields ϕ_a and antisymmetric n_s -forms F_s ... in a (pseudo-)Riemannian manifold $M = R_u \times M_0 \times \ldots \times M_n$ with the metric $ds^2 = g_{MN}dz^Mdz^N = we^{2\alpha(u)}du^2 +$ $\sum_{i=0}^n e^{2\beta^i(u)}ds_i^2$, $w = \pm 1$, where u is a selected coordinate ranging in $R_u \subseteq R; g^i = ds_i^2$ are metrics on d_i -dimensional factor spaces M_i of arbitrary signatures $\varepsilon_i = \text{sign } g^i; |g| =$ $|detg_{MN}|$ and similarly for subspaces; ... $M_i, i > 0$ are assumed to be Ricci-flat, while M_0 is allowed to be a space of constant curvature $K_0 = 0, \pm 1$." Two key steps in the proof of the STT, which have no counterpoint in our model, are the facts that the author of the STT : [1] employs the equations of motion of a test particle moving in the presence of a black hole (see STT Eq.[29]); and [2] employs the black hole potential (see STT Eq.[30]) in order to derive several equalities, from which he then finally concludes that more than one time dimension is not possible for his model.

However, in our present model the very early universe is modeled as *homogeneous* and isotropic. There are no black holes and no black hole potentials, so the STT clearly does not apply.

3. NOTATION AND CONVENTIONS

This model is cast on $\mathbb{X}_{4,4}$, which is an eight-dimensional pseudo-Riemannian manifold that is a spacetime of four space dimensions, with local comoving spatial coordinates (x^1, x^2, x^3, x^8) , and four time dimensions, with local comoving temporal coordinates (x^4, x^5, x^6, x^7) (employing the usual component notation in local charts). Greek indices run from 1 to 8. Each of the x^{α} has units (dimensions) of length. It is assumed that $-\infty < x^{\alpha} < \infty$ for $\alpha = 1, 2, ..., 7$.

The eight-dimensional pseudo-Riemannian manifold $\mathbb{X}_{4,4}$ may be decomposed as $\mathbb{X}_{4,4} = \mathbb{X}_{3,4} \times S^1_{(8)}$, where $S^1_{(8)}$ is a one dimensional compact space that is homeomorphic to the unit circle S^1 . We need at least two charts to cover $S^1_{(8)}$, which are chosen to be $x^8 : S^1_{(8)} \to \mathbb{R}$, $-\pi R_8 < x^8 < \pi R_8$ and $x'^8 : S^1_{(8)} \to \mathbb{R}$, $0 < x'^8 < 2\pi R_8$, where $R_8 \equiv 1$ is the Planck length. R_8 is introduced to give x^8 the correct dimension. R_8 is not associated to a dilatonic degree of freedom since it is constant, $R_8 \equiv 1$.

Let U be a subset of $\mathbb{X}_{4,4}$, $p \in U$ and $\phi: U \to \mathbb{R}^8$ be a local chart. $\frac{\partial}{\partial x^{\alpha}}$ denotes the natural basis of $T_p(\mathbb{X}_{4,4})$ associated with the coordinates $x^{\alpha} = [\phi(p)]^{\alpha}$, where $[.]^{\alpha}$ denotes the canonical projection of the α^{th} component of $x \in \mathbb{R}^8$. In the context of this paper a phrase such as " x^6 is a timelike dimension" means that the timelike vector $\frac{\partial}{\partial x^6}$ is tangent to the lines of constant spatial coordinates and constant (x^4, x^5, x^7) , i.e. the curves of $\mathbb{X}_{4,4}$ defined by $(x^1 = K_1, x^2 = K_2, x^3 = K_3, x^4 = K_4, x^5 = K_5, x^7 = K_7, x^8 = K_8)$, where $K_1, K_2, K_3, K_4, K_5, K_6, K_8$ are seven constants.

In this paper we obtain solutions to the coupled Einstein-inflaton field equations that exhibit temporal exponential deflation of three of the four time dimensions. More precisely, we record solutions that possess exponentially decreasing scale factors for three of the four timelike dimensions, where decreasing means decreasing with respect to the remaining fourth time coordinate, which is chosen to be x^4 . In effect we arbitrarily label the temporal direction/dimension that does not deflate (or inflate, for that matter) the x^4 - axis. The three remaining time dimensions are labeled with (x^5, x^6, x^7) and they undergo deflation. We assume that unknown quantum gravity effects sort things out shortly after the Universe is created so that the three mutually orthogonal deflating timelike directions and the remaining fourth mutually orthogonal non-inflating/non-deflating timelike direction (i.e., the temporal dimension whose scale factor is a constant) are the same for all values of (x^1, x^2, x^3, x^8) . This is an unsatisfying assumption, but a necessary hypothesis given our current knowledge of quantum gravity.

Let \mathbf{g} denote the pseudo-Riemannian metric tensor on $\mathbb{X}_{4,4}$. The signature of the metric \mathbf{g} is $(4,4) \leftrightarrow (+++---+)$. The covariant derivative with respect to the symmetric connection associated to the metric \mathbf{g} is denoted by a vertical double-bar. We employ the Landau-Lifshitz spacelike sign conventions [21]. $\mathbf{g} \leftrightarrow g_{\alpha\beta} = g_{\alpha\beta}(x^{\mu})$ is assumed to carry the Newton-Einstein gravitational degrees of freedom. It is moreover assumed that the ordinary Einstein field equations (on $\mathbb{X}_{4,4}$)

$$G_{\alpha\beta} + g_{\alpha\beta}\Lambda = 8 \pi \mathbb{G} T_{\alpha\beta}$$

$$(3.1)$$

are satisfied. Here $G_{\alpha\beta}$ denotes the Einstein tensor, Λ is the cosmological constant, \mathbb{G} denotes the Newtonian gravitational constant, and the reduced Planck mass is $M_{Pl} = [8\pi \mathbb{G}]^{-1/2}$. Natural units $c = 1 = \hbar = 8\pi \mathbb{G}$ are used throughout. Lastly, if $f = f(x^4, x^8)$ then

$$f^{(1,0)} = \frac{\partial}{\partial x^4} f(x^4, x^8), \quad f^{(0,1)} = \frac{\partial}{\partial x^8} f(x^4, x^8), \quad f^{(1,1)} = \frac{\partial^2}{\partial x^4 \partial x^8} f(x^4, x^8),$$

$$f^{(2,0)} = \frac{\partial^2}{\partial x^{42}} f(x^4, x^8), \quad f^{(0,2)} = \frac{\partial^2}{\partial x^{82}} f(x^4, x^8), \quad \text{etc.}$$
(3.2)

4. EINSTEIN FIELD EQUATIONS

4.1. Preview

We seek solutions to the Einstein field equations Eq.[3.1] for which the observable physical macroscopic early universe appears to a classical observer, after inflation ends, to be a homogeneous, isotropic universe with three space dimensions and one time dimension. Accordingly the three space dimensions (x^1, x^2, x^3) are assumed to possess equal scale factors, denoted $a = a(x^4, x^8)$, and the three extra time dimensions (x^5, x^6, x^7) are assumed to possess equal scale factors, denoted $b = b(x^4, x^8)$.

We find two periodic (in x^8) solution classes (E_0, E_1) to the Einstein field equations Eq.[3.1] on $\mathbb{X}_{4,4}$ that exhibit inflation/deflation and describe a universe that is spatially flat throughout the inflation era. Moreover it is found that Λ is quantized in terms of R_8 . For (E_0, E_1) , during "inflation", the scale factor $a = a(x^4, x^8)$ for the three space dimensions (x^1, x^2, x^3) exponentially inflates as a function of x^4 , and the scale factor $b = b(x^4, x^8)$ for the three extra time dimensions (x^5, x^6, x^7) exponentially deflates as a function of x^4 ; moreover, the scale factors for the x^4 and x^8 dimensions are constants equal to one.

We define the cosmological *relative expansion rate* ℓ as

$$\ell = \frac{1}{2} \frac{\partial}{\partial x^4} \ln\left(\frac{b}{a}\right). \tag{4.1}$$

In Section 5 it is shown that ℓ is independent of time and equal to $\ell_0 = -\sqrt{\frac{1}{18}\Lambda}$ for solutions in the first class E_0 ; this solution corresponds to pure exponential time inflation of the scale factor $a(x^4, x^8)$ and pure exponential time deflation of the scale factor $b(x^4, x^8)$. For this case $|\ell_0|$ coincides with the Hubble parameter H.

For the class E_0 solution we find that Λ is "quantized" in terms of the compact spatial x^8 unit radius $R_8 \equiv 1$, and given by

$$\Lambda_0 = \frac{1}{2} \frac{1}{R_8^2},\tag{4.2}$$

and therefore is one half the square of the Planck mass.

Therefore ℓ_0 verifies

$$R_8 \left| \ell_0 \right| = \frac{1}{6},\tag{4.3}$$

The two relationships are reminiscent of semiclassical quantization relationships.

For solutions in the second class E_1 , which describes a transition from pure exponential "inflation," $\ell = \ell_1$ is generally time dependent. In this case the "quantized" value of Λ is

$$\Lambda_1 = \frac{6}{5}\Lambda_0. \tag{4.4}$$

We refer to Λ_0 as the ground state value of the cosmological constant and Λ_1 as its first excited state.

Mathematical universes that possess positive relative expansion rate $\ell > 0$ for $x^4 > 0$ exist and are reported below. However they are analogous to collapsing universe solutions in the usual Friedmann cosmological model, or to universes with multiple macroscopic time dimensions, and hence are not directly relevant to the physical processes discussed in this paper.

4.2. Line element

The line element for inflation/deflation is assumed to be given by

$$\{ds\}^{2} = \{a(x^{4}, x^{8})\}^{2} \left[(dx^{1})^{2} + (dx^{2})^{2} + (dx^{3})^{2} \right] - (dx^{4})^{2} - \{b(x^{4}, x^{8})\}^{2} \left[(dx^{5})^{2} + (dx^{6})^{2} + (dx^{7})^{2} \right] + (dx^{8})^{2} = a^{2} \left[(dx^{1})^{2} + (dx^{2})^{2} + (dx^{3})^{2} \right] - b^{2} \left[(dx^{5})^{2} + (dx^{6})^{2} + (dx^{7})^{2} \right] - (dx^{4})^{2} + (dx^{8})^{2} ;$$

$$(4.5)$$

where $a = a(x^4, x^8)$ and $b = b(x^4, x^8)$ carry the metric degrees of freedom in this model. The real massless scalar inflaton field is $\varphi = \varphi(x^4, x^8)$. The action for the metric and inflaton degrees of freedom is assumed to be given by

$$S = \int \left[\frac{1}{16 \pi \mathbb{G}} \left(R - 2\Lambda \right) - \frac{1}{2} g^{\mu\nu} \partial_{\mu} \varphi \ \partial_{\nu} \varphi - \xi R \varphi^{2} + \mathfrak{L}_{\mathrm{SM}} \right]^{2} \sqrt{\det(g_{\alpha\beta})} \ d^{8}x.$$
 (4.6)

Here Λ is the cosmological constant. The inflaton potential is zero; its action is purely kinematic, although $\varphi^{(0,1)}(x^4, x^8)^2$ may be regarded as contributing to an effective inflaton potential. For the remainder of the paper we assume minimal coupling of the inflaton with gravity and set $\xi = 0$. We also neglect the standard model degrees of freedom and set \mathfrak{L}_{SM} = 0.

4.3. Canonical stress-energy tensor

The canonical stress-energy tensor for the real massless scalar inflaton field is $T_{\mu\nu} = -\frac{2}{\sqrt{\det(g_{\alpha\beta})}} \frac{\partial}{\partial g^{\mu\nu}} L_{\varphi}, \ L_{\varphi} = \sqrt{\det(g_{\alpha\beta})} \left[-\frac{1}{2} g^{\mu\nu} \partial_{\mu} \varphi \ \partial_{\nu} \varphi \right]$. The distinct components are

$$T_{33} = \frac{1}{2} \mathbf{a} (x^4, x^8)^2 \left[\varphi^{(1,0)} (x^4, x^8)^2 - \varphi^{(0,1)} (x^4, x^8)^2 \right], \qquad (4.7)$$

$$T_{44} = \frac{1}{2} \left[\varphi^{(1,0)}(x^4, x^8)^2 + \varphi^{(0,1)}(x^4, x^8)^2 \right], \qquad (4.8)$$

(which clearly satisfies the weak energy condition),

$$T_{55} = \frac{1}{2} \mathbf{b}(x^4, x^8)^2 \left(\varphi^{(0,1)}(x^4, x^8)^2 - \varphi^{(1,0)}(x^4, x^8)^2\right), \tag{4.9}$$

$$T_{88} = \frac{1}{2}\varphi^{(0,1)}(x^4, x^8)^2 + \frac{1}{2}\varphi^{(1,0)}(x^4, x^8)^2, \qquad (4.10)$$

and

$$T_{48} = T_{84} = \varphi^{(0,1)}(x^4, x^8)\varphi^{(1,0)}(x^4, x^8), \qquad (4.11)$$

which is non-zero, in general. Due to the (x^4, x^8) dependence of the metric the (4, 8) and (8, 4) components of the Einstein tensor $G_{48} = G_{84}$ are also, in general, non-zero.

Let $\rho = \rho(x^4, x^8)$ denote the effective energy density of the (classical) inflaton field and $p = p(x^4, x^8)$ denote the effective pressure. ρ and p may be associated to $T_{\mu\nu}$ according to

$$T^{\mu\nu} = \left[\sum_{j=4}^{j=7} u^{\mu}_{(j)} u^{\nu}_{(j)}\right] (\rho+p) + p g^{\mu\nu} + \left[\delta^{\mu}_{4}\delta^{\nu}_{8} + \delta^{\mu}_{8}\delta^{\nu}_{4}\right] T^{48}$$
(4.12)

where $u^{\mu}_{(j)} = \delta^{\mu}_{j} \ j = 4, 5, 6, 7$ in this comoving frame. This implies that

$$T_{44} = \rho = \frac{1}{2} \left[\varphi^{(1,0)}(x^4, x^8)^2 + \varphi^{(0,1)}(x^4, x^8)^2 \right]$$

$$T_{33} = g_{33} p = \frac{1}{2} \mathbf{a} (x^4, x^8)^2 \left[\varphi^{(1,0)}(x^4, x^8)^2 - \varphi^{(0,1)}(x^4, x^8)^2 \right].$$
(4.13)

Therefore

$$w = \frac{p}{\rho} = \frac{T_{33}}{g_{33}T_{44}} = \frac{\varphi^{(1,0)}(x^4, x^8)^2 - \varphi^{(0,1)}(x^4, x^8)^2}{\varphi^{(1,0)}(x^4, x^8)^2 + \varphi^{(0,1)}(x^4, x^8)^2}.$$
(4.14)

When $\varphi^{(1,0)}(x^4, x^8) = 0$, as is the case for the Class E_0 solution of Subsection[5], then

$$w = -1.$$
 (4.15)

In this case the effective inflaton potential may be defined as

$$V_{\rm eff} = \frac{1}{2} \varphi^{(0,1)} (x^4, x^8)^2.$$
(4.16)

Therefore for the Class E_0 solution of Section[5],

$$V_{\rm eff} = \frac{5}{4!} \Lambda \cot\left(\frac{1}{2}x^8 \sqrt{2\Lambda}\right)^2.$$
 (4.17)

4.4. Field Equations

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8 \pi \mathbb{G} T_{\mu\nu} \tag{4.18}$$

The distinct field equation components may be written as

$$G_{48} = G_{84} = -\frac{3a^{(1,1)}}{a} - \frac{3b^{(1,1)}}{b} = 8\pi \mathbb{G} \varphi^{(0,1)} \varphi^{(1,0)}$$
(4.19)

$$G_{33} = \frac{1}{b^2} \left[3ab \left(2a^{(0,1)}b^{(0,1)} - 2a^{(1,0)}b^{(1,0)} + a \left(b^{(0,2)} - b^{(2,0)} \right) \right) + \left(a^{(0,1)^2} - a^{(1,0)^2} + 2a \left(a^{(0,2)} - a^{(2,0)} \right) \right) b^2 + 3a^2 \left(b^{(0,1)^2} - b^{(1,0)^2} \right) \right] = 4\pi \mathbb{G}a^2 \left(-\varphi^{(0,1)^2} + \varphi^{(1,0)^2} \right)$$
(4.20)

$$G_{44} = -\frac{3}{a^2b^2} \left[b^2 \left(a^{(0,1)2} - a^{(1,0)2} + aa^{(0,2)} \right) + ab \left(3a^{(0,1)}b^{(0,1)} - 3a^{(1,0)}b^{(1,0)} + ab^{(0,2)} \right) \right]$$

+ $a^2 \left(b^{(0,1)2} - b^{(1,0)2} \right) \right]$
= $4\pi \mathbb{G} \left(\varphi^{(0,1)2} + \varphi^{(1,0)2} \right)$ (4.21)

$$G_{55} = \frac{1}{a^2} \left[2ab \left(-3a^{(0,1)}b^{(0,1)} + 3a^{(1,0)}b^{(1,0)} + a \left(b^{(2,0)} - b^{(0,2)} \right) \right) + 3 \left(-a^{(0,1)2} + a^{(1,0)2} + a \left(a^{(2,0)} - a^{(0,2)} \right) \right) b^2 + a^2 \left(b^{(1,0)2} - b^{(0,1)2} \right) \right] = 4\pi \mathbb{G}b^2 \left(\varphi^{(0,1)2} - \varphi^{(1,0)2} \right)$$

$$(4.22)$$

$$G_{88} = -\frac{3}{a^2b^2} \left[a^2 \left(b^{(1,0)2} - b^{(0,1)2} \right) + \left(-a^{(0,1)2} + a^{(1,0)2} + aa^{(2,0)} \right) b^2 + ab \left(-3a^{(0,1)}b^{(0,1)} + 3a^{(1,0)}b^{(1,0)} + ab^{(2,0)} \right) \right] = 4\pi \mathbb{G} \left(\varphi^{(0,1)2} + \varphi^{(1,0)2} \right)$$
(4.23)

The components of $T^{\mu}_{||\mu \alpha}$ that are not identically zero must satisfy

$$a b T^{\mu}_{||\mu 4} = 0 = 3a \left(-b^{(1,0)} \left(\varphi^{(1,0)2}\right) + b^{(0,1)} \varphi^{(0,1)} \varphi^{(1,0)} + b^{(0,1)}\right) + b \left(-3a^{(1,0)} \left(\varphi^{(1,0)2}\right) + 3a^{(0,1)} \varphi^{(0,1)} \varphi^{(1,0)} + a \left(\varphi^{(1,0)} \left(\varphi^{(0,2)} - \varphi^{(2,0)}\right)\right)\right) a b T^{\mu}_{||\mu 8} = 0 = 3a \left(b^{(0,1)} \left(\varphi^{(0,1)2}\right) - b^{(1,0)} \varphi^{(0,1)} \varphi^{(1,0)}\right) - b \left(-3a^{(0,1)} \varphi^{(0,1)2} + 3a^{(1,0)} \varphi^{(1,0)} \varphi^{(0,1)} - a \left(\varphi^{(0,1)} \left(\varphi^{(0,2)} - \varphi^{(2,0)}\right)\right)\right) .$$

$$(4.24)$$

Let $L_1 = \frac{\mathbf{a}^{(1,0)}(x^4, x^8)}{\mathbf{a}^{(x^4, x^8)}}$, $L_3 = \frac{\mathbf{a}^{(0,1)}(x^4, x^8)}{\mathbf{a}^{(x^4, x^8)}}$, $L_2 = \frac{\mathbf{b}^{(1,0)}(x^4, x^8)}{\mathbf{b}^{(x^4, x^8)}}$ and $L_4 = \frac{\mathbf{b}^{(0,1)}(x^4, x^8)}{\mathbf{b}^{(x^4, x^8)}}$. The Euler-Lagrange equation for the inflaton field yields

$$\varphi^{(2,0)}(x^4, x^8) - \varphi^{(0,2)}(x^4, x^8) + 3\varphi^{(1,0)}(x^4, x^8) (L_1 + L_2) - 3\varphi^{(0,1)}(x^4, x^8) (L_3 + L_4)$$

= 0, (4.25)

or, equivalently,

$$\varphi^{(2,0)}(x^4, x^8) + 3\varphi^{(1,0)}(x^4, x^8) (L_1 + L_2) + \mu^2 \varphi(x^4, x^8)$$

= $\varphi^{(0,2)}(x^4, x^8) + 3\varphi^{(0,1)}(x^4, x^8) (L_3 + L_4) + \mu^2 \varphi(x^4, x^8),$ (4.26)

where μ^2 is arbitrary.

Lastly, the field equations demand that the constraint equation

$$L_{1}^{2} + 3L_{1}L_{2} + L_{2}^{2} - \frac{4}{3}\pi \mathbb{G} \varphi^{(1,0)}(x^{4}, x^{8})^{2} - \left[L_{3}^{2} + 3L_{3}L_{4} + L_{4}^{2} - \frac{4}{3}\pi \mathbb{G} \varphi^{(0,1)}(x^{4}, x^{8})^{2}\right] = \frac{2}{9}\Lambda, \qquad (4.27)$$

be satisfied.

To solve the field equations we use the fact that $\mathbb{P} = \ln(a \times b)$, $\mathbb{Q} = \ln(\frac{b}{a})$ and φ satisfy, respectively, uncoupled, linear and linear field equations of the form

$$f^{(0,2)}(x^4, x^8) - f^{(2,0)}(x^4, x^8) = S + + 3 \mathbb{P}^{(1,0)}(x^4, x^8) f^{(1,0)}(x^4, x^8) - 3 \mathbb{P}^{(0,1)}(x^4, x^8) f^{(0,1)}(x^4, x^8), \qquad (4.28)$$

where $f = \mathbb{P}, \mathbb{Q}$ or φ and S = 0 unless $f = \mathbb{P}$ in which case $S = -\frac{2}{3}\Lambda$. General solutions to these equations are substituted into the constraint Eq[4.27] and the remaining field equations, which are then solved; this procedure yields the solutions in the classes E_0 and E_1 that are described below.

5. CLASS E_0 SOLUTION: EXACT TEMPORAL EXPONENTIAL INFLA-TION/DEFLATION

Using the technique described above we find that this model admits the following solutions, which are exponential in x^4 , periodic in x^8 and that are parameterized by Λ : The scale factors are

$$a = a(x^{4}, x^{8}) = a_{0}e^{\pm \frac{1}{3}\sqrt{\frac{\Lambda}{2}}x^{4}} \sqrt[12]{\sin^{2}\left(x^{8}\sqrt{2\Lambda}\right)}$$

$$b = b(x^{4}, x^{8}) = b_{0}e^{\mp \frac{1}{3}\sqrt{\frac{\Lambda}{2}}x^{4}} \sqrt[12]{\sin^{2}\left(x^{8}\sqrt{2\Lambda}\right)},$$
 (5.1)

where a_0 and b_0 are constants. The relative expansion rate and the inflaton field are given by

$$\ell_0 = \mp \frac{1}{3} \sqrt{\frac{\Lambda}{2}}$$

$$\varphi = \pm \frac{1}{2} \sqrt{\frac{5}{6}} \ln \left[\tan^2 \left(\frac{1}{2} \sqrt{2\Lambda} x^8 \right) \right].$$
(5.2)

For this case the scale factors (a, b) have a spatial period equal to $\frac{1}{2} C_8$, while the three functions (a, b, φ) possess a common spatial periodicity in the x^8 coordinate whose nonzero minimum value is equal to the x^8 -dimension spatial period

$$C_8 = 2\pi R_8 = \pi \sqrt{\frac{2}{\Lambda}} = \frac{\pi}{3|\ell_0|},$$
(5.3)

$$\Lambda = \Lambda_0 = \frac{1}{2} \frac{1}{R_8^2} \tag{5.4}$$

For this case $|\ell_0|$ coincides with the Hubble parameter H.

Since a universe with many macroscopic times is not observed, one may identify the physical solution for inflation by choosing the solution with the - sign in the equation for b. This solution then predicts the exponential deflation, with respect to time, of the scale factor b associated with the three extra time dimensions. This coincides with the exponential inflation, with respect to time, of the scale factor a associated with the three observed spatial dimensions.

6. TEMPERATURE HISTORY OF INFLATION IN THE CLASS E_0 BACK-GROUND

The reduced volume element $d\Omega$ on the hypersurface $x^4 = x_0^4 = \text{constant}$ is

$$d\Omega = d\Omega(x^{\alpha}) = \left[\sqrt{\det\left(\mathbf{g}\right)}\right]_{x^4 = x_0^4} d^7\tau$$
(6.1)

where

$$d^{7}\tau = |dx^{1} \wedge dx^{2} \wedge dx^{3} \wedge dx^{5} \wedge dx^{6} \wedge dx^{7} \wedge dx^{8}|.$$
(6.2)

For the class E_0 solution

$$d\Omega = d^7 \tau \left| \sin \left(x^8 \sqrt{2\Lambda} \right) \right|, \tag{6.3}$$

which clearly does not inflate. The fact that $d\Omega$ vanishes on a set of measure zero is discussed in the *Conclusion*.

To compute the temperature history $T = T(x^4)$ during inflation we assume that before standard model particles are created the chemical potential of the universe is zero. We also assume that the entropy density, energy density, and pressure describe only Λ and the inflaton φ field, and that their averages over x^8 are functions $s(T), \rho(T)$, and p(T) of the temperature T alone.

In analogy with the conventional definition of thermal equilibrium [see, for example, reference [22], p.150, Eq. (3.1.1)], we assume that the x^8 -average entropy in a co-moving volume element is fixed:

$$s(T)\left\langle \frac{d\Omega}{d^6\tau} \right\rangle_{x^8} = \int_{-\pi}^{\pi} s(T, x^8) \left| \sin\left(x^8 \sqrt{2\Lambda}\right) \right| \, dx^8 = \text{ constant}, \tag{6.4}$$

where $d^6\tau = |dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^5 \wedge dx^6 \wedge dx^7|$. Differentiating Eq.[6.4] with respect to x^4 yields $\frac{d}{dx^4} \left[s(T(x^4)) \frac{d\Omega}{d^6\tau} \right] = \frac{dT(x^4)}{dx^4} s'(T) \frac{d\Omega}{d^6\tau} + s(T) \frac{d}{dx^4} \left[\frac{d\Omega}{d^6\tau} \right] = \dot{T}(x^4) s'(T) \frac{d\Omega}{d^6\tau} + 0 = 0.$ Assuming that $s'(T) \neq 0$ gives

$$\frac{d}{dx^4}T(x^4) = 0. (6.5)$$

For this case the universe does not cool during inflation. Reheating [14] is a non-issue.

7. THE PROBLEM OF THE STABILITY OF A MASSIVE COMPLEX SCALAR FIELD PROPAGATING ON THE CLASS E_0 BACKGROUND

If a massive complex scalar field ψ satisfies a linear field equation, as does the ψ field under discussion, then a quantum fluctuation $\delta \psi$ of ψ satisfies a linear field equation that is of the same form. Therefore the question of the possible existence of instabilities of the massive complex scalar ψ field that are sourced by the momenta \vec{w} associated to the extra time dimensions, Issue [2] raised in Section[1], may be investigated by studying the propagation of the complex scalar field ψ on the background gravitational field, but which is not coupled to the inflaton field φ . In this paper the field equation for this scalar field is given in Eq.[2.1]. Substituting the background gravitational field of Eq.[5.1] into Eq.[2.1] and using Eq.[2.3] yields

$$0 = -\Psi^{(2,0)}(x^{4}, x^{8}) + \Psi^{(0,2)}(x^{4}, x^{8}) + \sqrt{2\Lambda} \cot\left(\sqrt{2\Lambda}x^{8}\right)\Psi^{(0,1)}(x^{4}, x^{8}) + \Lambda\left(w^{2}e^{\frac{1}{3}\sqrt{2\Lambda}x^{4}} - k^{2}e^{-\frac{1}{3}\sqrt{2\Lambda}x^{4}}\right)\sqrt[6]{\csc^{2}\left(\sqrt{2\Lambda}x^{8}\right)}\Psi(x^{4}, x^{8}) - \Lambda\left[m^{2} + \frac{1}{3}\zeta\left(8 + 5\csc^{2}\left(\sqrt{2\Lambda}x^{8}\right)\right)\right]\Psi(x^{4}, x^{8}),$$
(7.1)

which is a non-separable linear wave equation. Here $\vec{k} = (k_1, k_2, k_3)^T$, $\vec{w} = (k_5, k_6, k_7)^T$, $\Lambda k^2 = \vec{k} \cdot \vec{k} = k_1^2 + k_2^2 + k_3^2$ and $\Lambda w^2 = \vec{w} \cdot \vec{w} = k_5^2 + k_6^2 + k_7^2$; we have also used the fact that the Ricci scalar is $\frac{1}{3}\Lambda \left(5\csc^2\left(\sqrt{2\Lambda}x^8\right) + 8\right)$ for the class E_0 background gravitational field. In Eq.[7.1] we put $\Psi(x^4, x^8) = \psi \left(\sqrt{2\Lambda}x^4, \sqrt{2\Lambda}x^8\right)$, and then for notational simplicity set $x = \sqrt{2\Lambda}x^8$ (where $-\pi \leq x \leq \pi$), and $t = \sqrt{2\Lambda}x^4$. Since $\Lambda = \frac{1}{2}$ for this case, $x = x^8$. This yields a wave equation for the massive complex scalar $\psi(t, x)$ field that is given by

$$0 = -\psi^{(2,0)}(t, x) + \psi^{(0,2)}(t, x) + \cot(x)\psi^{(0,1)}(t, x) + \frac{1}{2}\left(w^2 e^{\frac{1}{3}t} - k^2 e^{-\frac{1}{3}t}\right) \sqrt[6]{\csc^2(x)} \psi(t, x) - \frac{1}{2}\left[m^2 + \frac{1}{3}\zeta\left(8 + 5\csc^2(x)\right)\right]\psi(t, x).$$
(7.2)

Note that

$$\frac{1}{2}\left(w^{2}e^{t/3} - k^{2}e^{-t/3}\right) = kw\frac{1}{2}\left(\frac{w}{k}e^{t/3} - \frac{k}{w}e^{-t/3}\right) = kw\sinh\left[\frac{t}{3} + \ln\left(\frac{w}{k}\right)\right].$$
 (7.3)

Eq.[7.2] is non-separable. Obviously one cannot write $\psi(t, x) = T(t)X(x)$ then separate functions of x from functions of t. On the other hand, for a separable linear partial differential equation one may separate variables, then introduce a separation constant and in principle solve the ODE for eigenmodes, subject to given boundary conditions, and deduce a dispersion relation that directly relates a physical frequency ω to, say, the momentum

wave vectors (\vec{k}, \vec{w}) . Eq.[7.2] does not admit a simple dispersion relation that relates a physical-frequency ω to the momentum wave vectors (\vec{k}, \vec{w}) .

We study two special cases of Eq.[7.2], the first with $\zeta = 3/10$ and the second with $\zeta = 0$. The solution for $\zeta = 3/10$ is expanded into a series of Fourier modes $\sum_{n_8=-\infty}^{\infty} f_{n_8}(t) \exp(i n_8 x)$. The second solution with $\zeta = 0$ is expanded into a series of (two) Legendre polynomial modes, each of the form $\sum_{n_8=0}^{\infty} h_{n_8}(t) \left[\sqrt{(n_8 + \frac{1}{2})}P_{n_8}(\cos x)\right]$. There is an expansion for each of the intervals $-\pi \leq x \leq 0$ and $0 \leq x \leq \pi$. The two Legendre expansions must be matched at x = 0 and $x = \pm \pi$.

7.0.1. Wave equation with $\zeta = 3/10$

We take advantage of the coupling of the massive scalar field to the Ricci scalar to simplify the wave equation. In Eq.[7.2] we first put $\psi(t, x) = \sin^{-\frac{1}{2}}(x) F(t, x)$. This yields

$$0 = F^{(2,0)}(t,x) - F^{(0,2)}(t,x) + \frac{1}{2} \left(k^2 e^{-t/3} - w^2 e^{t/3} \right) \sqrt[6]{\csc^2(x)} F(t,x) + \frac{1}{12} \left(16\zeta + 6m^2 - 3 \right) F(t,x) + \frac{1}{12} (10\zeta - 3) \csc^2(x) F(t,x).$$
(7.4)

In order to eliminate the $\frac{1}{12}(10\zeta - 3)\csc^2(x)F(t,x)$ term from this equation we henceforth assume in this section that $\zeta = 3/10$.

Substituting a Fourier decomposition

 $F(t,x) = \sum_{n_8=-\infty}^{\infty} f_{n_8}(t) \exp(i n_8 x)$ into Eq.[7.4], with $\zeta = 3/10$, and then multiplying by $\frac{1}{2\pi} \exp(-i m_8 x) dx$ and integrating from $(-\pi, \pi)$ yields

$$0 = \frac{d^2}{dt^2} f_{m_8}(t) + m_{\text{eff}}^2 f_{m_8}(t) - \frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3} \right) \sum_{n_8 = -\infty}^{\infty} \left\{ c_{m_8 n_8} f_{n_8}(t) \right\}$$
$$= \frac{d^2}{dt^2} f_{m_8}(t) + m_{\text{eff}}^2 f_{m_8}(t) - k w \sinh\left[\frac{t}{3} + \ln\left(\frac{w}{k}\right)\right] \sum_{n_8 = -\infty}^{\infty} \left\{ c_{m_8 n_8} f_{n_8}(t) \right\}, \quad (7.5)$$

where the c_{m_8,n_8} are defined in Eq.[7.7] and

$$m_{\rm eff}^2 = \frac{m^2}{2} + \frac{3}{20} + m_8^2$$
(7.6)

As expected, the square of the effective mass $m_{\rm eff}$ of the scalar field receives contributions from the spatial x^8 Fourier modes. The square of the effective mass $m_{\rm eff}$ also receives a contribution $\zeta = 3/10$ from the coupling with the Ricci scalar. The matrix elements c_{m_8,n_8} of $\sqrt[6]{\csc^2(x)}$ in the Fourier basis are

$$c_{m_8,n_8} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sqrt[6]{\csc^2(x)} \exp[i(n_8 - m_8)x] dx$$

= $\sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} \begin{cases} (-1)^N \frac{\Gamma(\frac{5}{6})^2}{\Gamma(\frac{5}{6} - N)\Gamma(\frac{5}{6} + N)} & \text{if } n_8 - m_8 = 2N \text{ is even} \\ 0 & \text{if } n_8 - m_8 \text{ is odd} \end{cases}$. (7.7)

We note that the $(-1)^N \frac{\Gamma(\frac{5}{6})^2}{\Gamma(\frac{5}{6}-N)\Gamma(\frac{5}{6}+N)}$ are rational numbers,

$$\gamma_{N} = (-1)^{N} \frac{\Gamma(\frac{5}{6})^{2}}{\Gamma(\frac{5}{6} - N) \Gamma(\frac{5}{6} + N)}$$

$$= \left\{ 1, \frac{1}{5}, \frac{7}{55}, \frac{91}{935}, \frac{1729}{21505}, \frac{8645}{124729}, \frac{7657}{124729}, \frac{283309}{5113889}, \frac{12182287}{240352783}, \frac{596932063}{12738697499}, \frac{2984660315}{68325741131}, \frac{2800988911}{68325741131}, \frac{187666257037}{4851127620301}, \frac{1957090966243}{53362403823311}, \frac{154610186333197}{4429079517334813}, \frac{773050931665985}{23187533943694021} \dots \right\}$$

$$(7.8)$$

for N = 0, 1, 2, ..., 15, ... The c_{m_8,n_8} are the matrix elements of a Toeplitz matrix and verify $c_{m_8,n_8} = c_{m_8-n_8,0} = c_{0,n_8-m_8}$. Accordingly, Eq.[7.5] may be written as

$$0 = \frac{d^2}{dt^2} f_{m_8}(t) + \left\{ m_{\text{eff}}^2 - k \, w \, \sinh\left[\frac{t}{3} + \ln\left(\frac{w}{k}\right)\right] c_{0,0} \right\} f_{m_8}(t) - k \, w \, \sinh\left[\frac{t}{3} + \ln\left(\frac{w}{k}\right)\right] \sum_{n_8=1}^{\infty} c_{2n_8,0} \left[f_{2n_8+m_8}(t) + f_{-2n_8+m_8}(t)\right]$$
(7.9)

or

$$0 = \frac{d^2}{dt^2} f_{m_8}(t) + \left\{ m_{\text{eff}}^2 - \sqrt{Y} \, 2 \, \sinh(\tau) \right\} f_{m_8}(t) - \sqrt{Y} \, 2 \, \sinh(\tau) \sum_{n_8=1}^{\infty} (-1)^{n_8} \frac{\Gamma(\frac{5}{6})^2}{\Gamma(\frac{5}{6} - n_8) \, \Gamma(\frac{5}{6} + n_8)} \left[f_{2n_8 + m_8}(t) + f_{-2n_8 + m_8}(t) \right],$$
(7.10)

where

$$\tau = \tau \left(t \right) = \frac{1}{3}t + \log\left(\frac{w}{k}\right), \tag{7.11}$$

$$\sqrt{Y} = \frac{1}{2} k \, w \, \sqrt[3]{2} \, \frac{\Gamma\left(\frac{2}{3}\right)}{\Gamma\left(\frac{5}{6}\right)^2} \tag{7.12}$$

and

$$m_{\rm eff}^2 = \frac{m^2}{2} + \frac{3}{20} + m_8^2.$$
 (7.13)

Because $\frac{\Gamma(\frac{5}{6})^2}{\Gamma(\frac{5}{6}-N)\Gamma(\frac{5}{6}+N)} \propto \left(\frac{1}{|N|}\right)^{2/3}$ as $N \to \pm \infty$, the series expansions in Eq.[7.5] and Eq.[7.10] exhibit a very long range coupling between modes. In general the infinite sums in Eq.[7.5] and Eq.[7.10] will converge if $|f_{n_8}(t)| < \left(\frac{1}{|n_8|}\right)^{1/3}$ as $|n_8| \to \infty$.

In order to estimate the error of an approximate numerical solution to Eq.[7.10] an explicit approximate form for $\sqrt[6]{\csc^2(x)}$ in terms of its Fourier components $c_{m_8 n_8}$ is needed. We find that

$$\sqrt[6]{\operatorname{csc}^{2}(x)} = \sum_{n_{8} = -\infty}^{\infty} \left\{ c_{m_{8}n_{8}} e^{i n_{8}x} \right\}
= \frac{\sqrt[3]{2}\Gamma\left(\frac{2}{3}\right)}{\Gamma\left(\frac{5}{6}\right)^{2}} \sum_{m_{8} = -\infty}^{\infty} e^{2i m_{8}x} \left(\frac{(-1)^{m_{8}}\Gamma\left(\frac{5}{6}\right)^{2}}{\Gamma\left(\frac{5}{6} - m_{8}\right)\Gamma\left(\frac{5}{6} + m_{8}\right)} \right)
= \frac{2\sqrt{\frac{\pi}{3}}}{\Gamma\left(\frac{2}{3}\right)\Gamma\left(\frac{5}{6}\right)} \left({}_{2}F_{1}\left(\frac{1}{6}, 1; \frac{5}{6}; e^{2ix}\right) + {}_{2}F_{1}\left(\frac{1}{6}, 1; \frac{5}{6}; e^{-2ix}\right) - 1 \right) \quad (7.14)$$

almost everywhere, except $x = 0, \pm \pi$. A plot of the difference

 $\int \sqrt[6]{\csc^2(x)} - \frac{2\sqrt{\frac{\pi}{3}}}{\Gamma(\frac{2}{3})\Gamma(\frac{5}{6})} \left[{}_2F_1\left(\frac{1}{6}, 1; \frac{5}{6}; e^{2ix}\right) + {}_2F_1\left(\frac{1}{6}, 1; \frac{5}{6}; e^{-2ix}\right) - 1 \right]$ is given in Figure[1]. Let $n \in \mathbb{N}$, the natural numbers (excluding 0). Given an approximate numerical solution that contains only a finite number of Fourier modes $-n \leq n_8 \leq n$ one may make the substitution based on Eq.[7.14]

$$\sqrt[6]{\operatorname{csc}^2(x)} \to \frac{\sqrt[3]{2}\Gamma\left(\frac{2}{3}\right)}{\Gamma\left(\frac{5}{6}\right)^2} \sum_{m8=-\left[\frac{n}{2}\right]}^{\left[\frac{n}{2}\right]} \frac{e^{i(2m8)x}\left((-1)^{m8}\Gamma\left(\frac{5}{6}\right)^2\right)}{\Gamma\left(\frac{5}{6}-m8\right)\Gamma\left(m8+\frac{5}{6}\right)}$$
(7.15)

when computing the numerical error. Here, $\left[\frac{n}{2}\right]$ gives the greatest integer less than or equal to $\frac{n}{2}$.

We call the first line of Eq.[7.10]

$$\mathcal{D}[f](k, w; m_8; t) = \frac{d^2}{dt^2} f_{m_8}(t) + \left[m_{\text{eff}}^2 - \frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3} \right) c_{0,0} \right] f_{m_8}(t)$$
(7.16)

the diagonal contribution to Eq.[7.10]. Here $c_{0,0} = \sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2}$.

7.0.2. Special solutions to $\mathcal{D}[f](k, w; m_8; t) = 0$

 $\mathcal{D}[f](k, w; m_8; t) = 0$ possesses simple mathematical solutions for the two special cases k = 0 and w = 0, which provide much insight into this problem, but which are not physical because the long range coupling with the other modes is completely neglected. These solutions are discussed in the next section.

Let $J_n(z)$ denote an ordinary Bessel function of the first type, $Y_n(z)$ denote an ordinary Bessel function of the second type (Neumann function), $I_{\lambda}(z)$ denote the modified Bessel function of the first type and order λ , and $K_{\lambda}(z)$ denote a modified Bessel function of the second type (also known as the modified Bessel function of the third kind).

If k = 0 then the general solution to $\mathcal{D}[f](k = 0, w; m_8; t) = 0$ is

$$f_{m_8}(t) = c_1 K_{6\,i\,m_{\text{eff}}}\left(w\,e^{t/6}\,\alpha\right) + c_2 I_{6\,i\,m_{\text{eff}}}\left(w\,e^{t/6}\,\alpha\right),\tag{7.17}$$

where

$$\alpha = \sqrt{18 c_{0,0}} = 3 \times 2^{2/3} \frac{\sqrt{\Gamma\left(\frac{2}{3}\right)}}{\Gamma\left(\frac{5}{6}\right)}$$
(7.18)

and the (c_1, c_2) are arbitrary constants. Asymptotically, for large argument |z|, the modified Bessel functions behave as $K_{\lambda}(z) \sim e^{-z} \sqrt{\frac{\pi}{2z}} + \cdots$ and $I_{\lambda}(z) \sim e^{z} \sqrt{\frac{1}{2\pi z}} + \cdots$ [23]. Therefore $f_{m_8}(t)$ is a stable solution as $w \to \infty$ if $c_2 = 0$. Moreover, $K_{i\nu}(z)$ possesses a well known integral representation

$$K_{i\nu}(z) = \int_0^\infty \cos(\nu\,\xi) \, e^{-z\,\cosh(\xi)} \, d\,\xi, \tag{7.19}$$

valid for $\arg z < \frac{\pi}{2}$, which enables a simple demonstration that $f_{m_8}(t)$ is stable as $w \to \infty$.

A stable solution to $\mathcal{D}[f](k = 0, w; m_8; t) = 0$ may be obtained by choosing an initial condition for $\frac{f'_{m_8}(t)}{f_{m_8}(t)}$ so that c_2 vanishes. The required initial condition is

$$\frac{f_{m_8}'(0)}{f_{m_8}(0)} = \left\{ \frac{\frac{\partial}{\partial t} K_{6\,i\,m_{\text{eff}}}\left(w\,e^{t/6}\,\alpha\right)}{K_{6\,i\,m_{\text{eff}}}\left(w\,e^{t/6}\,\alpha\right)} \right\}_{t=0} = \left\{ \frac{1}{6}\,W\,\frac{\partial\log(K_{6\,i\,m_{\text{eff}}}(W))}{\partial W} \right\}_{W=w\,\alpha} \,, \qquad (7.20)$$

where α is defined in Eq.[7.18]. To make this solution also well-behaved as $w \to 0$ one can, for example, put $c_1 = \frac{w}{1+w}C_1$.

If one attempts to numerically integrate $\mathcal{D}[f](k = 0, w; m_8; t) = 0$ using a numerical quadrature algorithm then at each time step t_n of the process one must maintain tight control of the error

$$\left\{\frac{f'_{m_8}(t_n)}{f_{m_8}(t_n)}\right\}_{\text{numerical}} - \left\{\frac{\frac{\partial}{\partial t}K_{6\,i\,m_{\text{eff}}}\left(w\,e^{t/6}\,\alpha\right)}{K_{6\,i\,m_{\text{eff}}}\left(w\,e^{t/6}\,\alpha\right)}\right\}_{t=t_n},\tag{7.21}$$

because a non-zero value of this difference will source a contribution from $I_{6im_{\text{eff}}}(w e^{t/6} \alpha)$ in the numerical solution, which may eventually exponentially overwhelm the true analytical solution. This problem is stiff.

We note in passing that the behavior of the modified Bessel function of the second kind with pure imaginary order, $K_{i\nu}(z)$, has been investigated by Balogh [24] and others. Balogh has proved that $K_{i\nu}(\frac{\nu}{p})$ is a positive monotone decreasing convex function of p for 0and it oscillates boundedly for <math>p > 1, having a countably infinite number of zeros. Balogh gives asymptotic expansions of the zeros of $K_{i\nu}(z)$ and its derivative that are uniform with respect to the enumeration of the zeros.

If w = 0 then the stable solution to $\mathcal{D}[f](k, w = 0; m_8; t) = 0$ is

$$f_{m_8}(t) = c_3 J_{6\,i\,m_{\text{eff}}}\left(k\,e^{-t/6}\,\alpha\right) + c_4 Y_{6\,i\,m_{\text{eff}}}\left(k\,e^{-t/6}\,\alpha\right),\tag{7.22}$$

where the (c_3, c_4) are arbitrary constants.

7.1. Wave equation with $\zeta = 0$

We put $\zeta = 0$ in Eq.[7.2] to obtain the wave equation for a minimally coupled massive complex scalar field propagating through the background class E_0 gravitational field:

$$0 = -\psi^{(2,0)}(t, x) + \psi^{(0,2)}(t, x) + \cot(x)\psi^{(0,1)}(t, x) - \frac{1}{2}m^{2}\psi(t, x) + \frac{1}{2}\left(w^{2}e^{\frac{1}{3}t} - k^{2}e^{-\frac{1}{3}t}\right)\sqrt[6]{\csc^{2}(x)}\psi(t, x) = -\psi^{(2,0)}(t, x) + \frac{1}{\sin(x)}\frac{\partial}{\partial x}\left[\sin(x)\frac{\partial}{\partial x}\psi(t, x)\right] - \frac{1}{2}m^{2}\psi(t, x) + \frac{1}{2}\left(w^{2}e^{\frac{1}{3}t} - k^{2}e^{-\frac{1}{3}t}\right)\sqrt[6]{\csc^{2}(x)}\psi(t, x).$$
(7.23)

Recall that $-\pi \leq x \leq \pi$ and not $0 \leq x \leq \pi$. In order to obtain a solution to Eq.[7.23] we solve for $\psi(t, x)$ on each of the two intervals $\mathcal{I}_{-} = -\pi \leq x < 0$ and $\mathcal{I}_{+} = 0 \leq x \leq \pi$ and then enforce appropriate continuity and boundary conditions: We assume that $\psi(t, x)$ and $\frac{\partial}{\partial x}\psi(t, x)$ are continuous, and that $\psi(t, -\pi) = \psi(t, \pi)$, since x may be regarded as a canonical angular coordinate θ for the boundary of the unit circle. Note that, with $X = \cos x = \cos \theta, \ x = \theta = \pm \arccos X$, Eq.[7.23] maps to

$$- \psi^{(2,0)}(t,X) + (1-X^2) \psi^{(0,2)}(t,X) - 2X\psi^{(0,1)}(t,X) - \frac{m^2}{2}\psi(t,X) + \frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3}\right) \left(\sqrt[6]{\frac{1}{1-X^2}}\right) \psi(t,X) = 0.$$
(7.24)

Let U(x) denote the unit step function

$$U(x) = \begin{cases} 0 & \text{for } x < 0\\ \frac{1}{2} & \text{for } x = 0\\ 1 & \text{for } x > 0; \end{cases}$$
(7.25)

accordingly, $\psi(t, x)$ may be expressed as

$$\psi(t, x) = \psi(t, x) \rfloor_{\mathcal{I}_{-}} U(-x) + \psi(t, x) \rfloor_{\mathcal{I}_{+}} U(x)$$

= $\psi_{-}(t, x) U(-x) + \psi_{+}(t, x) U(x),$ (7.26)

and $\frac{\partial}{\partial x}\psi(t, x)$ is given by

$$\psi^{(0,1)}(t, x) \equiv \frac{\partial}{\partial x} \psi(t, x)$$

= $U(-x) \frac{\partial}{\partial x} \psi_{-}(t, x) + U(x) \frac{\partial}{\partial x} \psi_{+}(t, x)$
+ $\delta(x) [\psi_{+}(t, 0) - \psi_{-}(t, 0)].$ (7.27)

As stated above, we assume that $\psi(t, x)$ and $\frac{\partial}{\partial x}\psi(t, x)$ are continuous, and that $\psi(t, -\pi) = \psi(t, \pi)$. These conditions imply that

$$\psi(t,0) = \frac{1}{2} \left[\psi_{-}(t,0) + \psi_{+}(t,0) \right] = \begin{cases} \lim_{x \to 0^{+}} \psi_{-}(t,x) \\ \lim_{x \to 0^{-}} \psi_{+}(t,x) \end{cases},$$
(7.28)

whence

$$\psi_{-}(t, 0) = \psi_{+}(t, 0), \qquad (7.29)$$

and, using Eq.[7.29] in Eq.[7.27],

$$\psi^{(0,1)}(t,0) = \frac{1}{2} \left[\psi^{(0,1)}_{-}(t,0) + \psi^{(0,1)}_{+}(t,0) \right] = \begin{cases} \lim_{x \to 0^+} \psi^{(0,1)}_{-}(t,x) \\ \lim_{x \to 0^-} \psi^{(0,1)}_{+}(t,x) . \end{cases}$$
(7.30)

This implies that

$$\psi_{-}^{(0,1)}(t,0) = \psi_{+}^{(0,1)}(t,0)$$
(7.31)

For simplicity, in order to satisfy the boundary and continuity conditions we henceforth assume for the remainder of this paper that

$$\psi^{(0,1)}(t,0) = 0 = \psi^{(0,1)}_{-}(t,0) = \psi^{(0,1)}_{+}(t,0)$$
(7.32)

and that when $x \in \mathcal{I}_{-}, \ -\pi \leq x < 0$,

$$\psi_{-}(t, x) = \psi_{+}(t, -x), \ x \in \mathcal{I}_{-}.$$
(7.33)

Next, let us expand $\psi(t, x)$ on each of the two intervals $\mathcal{I}_{-} = -\pi \leq x < 0$ and $\mathcal{I}_{+} = 0 \leq x \leq \pi$ in terms of a series of Legendre polynomials of the form

$$\psi_{\pm}(t, x) \equiv \psi(t, x) \rfloor_{\mathcal{I}_{\pm}} = \sum_{n_8=0}^{\infty} h_{n_8}^{(\pm)}(t) \left[\sqrt{\left(n_8 + \frac{1}{2}\right)} P_{n_8}(\cos x) \right].$$
(7.34)

In virtue of Eq.[7.33], $h_{n_8}^{(-)}(t) = h_{n_8}^{(+)}(t)$. For brevity we put $h_{n_8}^{(+)}(t) = h_{n_8}(t)$. Before substituting Eq.[7.34] into Eq.[7.23] and simplifying we record the following definitions.

Let \mathbb{N} denote the non-negative integers $0, 1, 2, 3, \ldots$. Define, $\forall m_8, n_8 \in \mathbb{N}$,

$$d_{m_8 n_8} = d_{m_8, n_8}$$

$$= \sqrt{\left(m_8 + \frac{1}{2}\right)} \sqrt{\left(n_8 + \frac{1}{2}\right)} \int_0^{\pi} P_{m_8}(\cos x) P_{n_8}(\cos x) \sin^{\frac{2}{3}}(x) dx$$

$$= \sqrt{\left(m_8 + \frac{1}{2}\right)} \sqrt{\left(n_8 + \frac{1}{2}\right)} \int_{-1}^{1} P_{m_8}(X) P_{n_8}(X) \frac{1}{\sqrt[6]{1 - X^2}} dX$$

$$= \pi \Gamma\left(\frac{5}{6}\right)^2 \sqrt{\left(m_8 + \frac{1}{2}\right)} \left(n_8 + \frac{1}{2}\right)$$

$$\sum_{\substack{j = |n_8 - m_8|, \text{ step } 2}}^{m_8 + n_8} \left\{ \frac{(2j+1)(-1)^{j+m_8 + n_8}}{\Gamma\left(\frac{1}{2} - \frac{j}{2}\right)\Gamma\left(\frac{5}{6} - \frac{j}{2}\right)\Gamma\left(\frac{j}{2} + 1\right)\Gamma\left(\frac{j}{2} + \frac{4}{3}\right)} \frac{\Gamma(-j + m_8 + n_8 + 1)\Gamma(j - m_8 + n_8 + 1)\Gamma(j + m_8 - n_8 + 1)}{\Gamma(j + m_8 + n_8 + 2)} \left[\frac{\Gamma\left(\frac{1}{2}(j + m_8 + n_8 + 2)\right)}{\Gamma\left(\frac{1}{2}(j - j + m_8 + n_8 + 2)\right)\Gamma\left(\frac{1}{2}(j - m_8 + n_8 + 2)\right)\Gamma\left(\frac{1}{2}(j + m_8 - n_8 + 2)\right)} \right]^2 \right\}$$

$$(7.35)$$

(of course, $\sin^{\frac{2}{3}}(x) = \sqrt[6]{\csc^2(x)} \sin x$ on $0 \le x \le \pi$). One observes that

$$d_{2m_8+1,2n_8} = 0 = d_{2m_8,2n_8+1} \ \forall \ m_8, n_8 \in \mathbb{N}.$$
(7.36)

Substituting Eq.[7.34] into Eq.[7.23], multiplying by $\left[\sqrt{\left(m_8 + \frac{1}{2}\right)}P_{m_8}(\cos x)\right]\sin x \, dx, m_8 \in \mathbb{N}$, and integrating over the interval $0 \leq x \leq \pi$ yields

$$\ddot{h}_{m_8}(t) + \left[\frac{1}{2}m^2 + m_8(m_8 + 1)\right]h_{m_8}(t) + \frac{1}{2}\left(k^2 e^{-\frac{1}{3}t} - w^2 e^{\frac{1}{3}t}\right)\sum_{n_8=0}^{\infty} d_{m_8n_8}h_{n_8}(t) = 0. \quad (7.37)$$

For this case the effective mass $m_{\rm eff}$ is given by

$$m_{\rm eff}^2 = \frac{1}{2}m^2 + m_8(m_8 + 1)$$
 (7.38)

We find that $d_{m_8 n_8} \propto \left(\frac{1}{n_8}\right)^{2/3}$ for $n_8 \gg m_8$ as $n_8 \to \pm \infty$. the series expansion in Eq.[7.37] also exhibits a very long range coupling between modes. The infinite sum in Eq.[7.37] will converge if $|h_{n_8}(t)| < \left(\frac{1}{n_8}\right)^{1/3}$ as $n_8 \to \pm \infty$.

The diagonal contribution to Eq.[7.37] is

$$\mathcal{D}[h](k, w; m_8; t) = \ddot{h}_{m_8}(t) + \left[m_{\text{eff}}^2 - \frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3}\right) d_{m_8 m_8}\right] h_{m_8}(t),$$
(7.39)

which has the same form as the diagonal contribution to Eq.[7.10].

Setting to zero the diagonal contribution to either Eq.[7.10] or Eq.[7.37] yields an equation of the form

$$-\ddot{Y}_{m_8}(t) + B \frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3} \right) Y_{m_8}(t) = m_{\text{eff}}^2 Y_{m_8}(t), \qquad (7.40)$$
where
$$\int m_{\text{eff}}^2 = \frac{3}{2} + \frac{1}{2} m^2 + m_8^2 \qquad B = c_{0.0} \qquad \text{for } \zeta = \frac{3}{2}$$

$$\begin{cases} m_{\text{eff}}^{-} = \frac{1}{20} + \frac{1}{2}m^{2} + m_{8}^{2}, & B = c_{0,0}, & \text{for } \zeta = \frac{1}{10}, \\ m_{\text{eff}}^{-} = \frac{1}{2}m^{2} + m_{8}(m_{8} + 1), & B = B(m_{8}) = d_{m_{8}m_{8}}, & \text{for } \zeta = 0 \end{cases}$$

$$(7.41)$$

8. EXISTENCE OF STABLE SOLUTIONS

We recall the content of Issue [2], Momenta corresponding to the extra time dimensions induce exponentially rapid growth of quantum fluctuations of the field; the universe is unstable. This instability is associated with the very largest momenta (shortest wavelengths).

In this paper we are concerned with exponentially rapid growth of quantum fluctuations of the field. This causes perturbation theory to break down since $|\delta\psi|$ becomes of order one faster than some positive power of t. Exponentially rapid growth of the field ψ itself is not an instability. Instability of the field ψ means the norm $|\psi|$ of the field goes to infinity in finite time. This is not the issue under discussion. "Stable" means that solutions $f_{m_8} = \delta\psi_{m_8}$ of Eq.[7.10] and $h_{m_8} = \delta\psi_{m_8}$ of Eq.[7.37] exist that possess a norm that is non-exponentially increasing as w increases. Then if $|\delta\psi|$ is initially small then $|\delta\psi|$ does not become of order one exponentially with time.

A formal argument can be made that demonstrates that stable solutions $f_{m_8} = \delta \psi_{m_8}$ to Eq.[7.10] and $h_{m_8} = \delta \psi_{m_8}$ to Eq.[7.37] always exist. Since both Eq.[7.10] and Eq.[7.37] have similar forms, our discussion will focus on Eq.[7.10]; our conclusions will be valid for both equations.

The phrase stable solution means, most importantly, that the norm of the solution is finite, for fixed time t and k, as $w \to \infty$ [recall that $\vec{k} = (k_1, k_2, k_3)^T$, $\vec{w} = (k_5, k_6, k_7)^T$, $\Lambda k^2 = \vec{k} \cdot \vec{k} = k_1^2 + k_2^2 + k_3^2$ and $\Lambda w^2 = \vec{w} \cdot \vec{w} = k_5^2 + k_6^2 + k_7^2$].

The argument begins as follows. We affinely transform the physical time t to a new time

coordinate τ that is defined by setting

$$t = 3\tau + 3\log\left(\frac{k}{w}\right) \tag{8.1}$$

As we have seen above, under this transformation

$$\frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3} \right) = k w \frac{1}{2} \left(\frac{w}{k} e^{t/3} - \frac{k}{w} e^{-t/3} \right)$$

= $k w \sinh\left[\frac{t}{3} + \ln\left(\frac{w}{k}\right)\right] = k w \sinh(\tau).$ (8.2)

In terms of the new time coordinate τ , $\psi = \psi(\tau, x)$ satisfies the wave equation Eq.[8.10], below. Since the quantum fluctuations of ψ satisfy a field equation similar to Eq.[8.10], we limit our discussion of stability to solutions of Eq.[8.10] and its relatives.

The inverse transformation to the map of the time coordinate t defined in Eq.[8.1] is

$$\tau = \frac{1}{3}t + \log\left(\frac{w}{k}\right). \tag{8.3}$$

For fixed t and k, as $w \to \infty$ then $\tau \to \infty$. For fixed t and k a solution $\delta\psi(\tau, x)$ is called *stable* if the norm $\|\delta\psi\|$ of the solution to the wave equation Eq.[8.10], below, satisfies $0 \leq \|\delta\psi\| < \infty$ as $\tau \to \infty$. Note that an effective frequency-like parameter ϖ , with $\varpi^2 = 2\sqrt{Y} \propto kw$ appears in Eq.[8.4] and Eq.[8.10], below. For fixed w, as $k \to \infty$ then $\varpi \to \infty$ and for fixed k, as $w \to \infty$ then $\varpi \to \infty$. The dependence of $\delta\psi(\tau, x)$ on ϖ as $\varpi \to \infty$ does not enter into our definition of stability. $\delta\psi(\tau, x)$ oscillates at higher and higher frequencies as $\varpi \to \infty$, but does not become unstable. Both non-resonant and resonant oscillations do not drive $|\delta\psi|$ to order one faster than some positive power t^{γ} of t.

Under the change of time coordinate defined in Eq.[8.1], Eq.[7.10] maps to

$$0 = \frac{1}{9} \frac{d^2}{d\tau^2} h_{n_8}(\tau) + \left\{ m_{\text{eff}}^2 - \sqrt{Y} \, 2 \, \sinh(\tau) \right\} h_{n_8}(\tau) - \sqrt{Y} \, 2 \, \sinh(\tau) \sum_{m_8=1}^{\infty} (-1)^{m_8} \frac{\Gamma(\frac{5}{6})^2}{\Gamma(\frac{5}{6} - m_8) \, \Gamma(\frac{5}{6} + m_8)} \left[h_{2m_8 + n_8}(\tau) + h_{-2m_8 + n_8}(\tau) \right],$$
(8.4)

where $h_{n_8}(\tau) = f_{n_8}(t) = \delta \psi_{n_8}(t)$, $m_{\text{eff}}^2 = \frac{m^2}{2} + \frac{3}{20} + n_8^2$ (see Eq.[7.13]) and $\sqrt{Y} = \frac{1}{2}k w \sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2}$ (see Eq.[7.12]); the independent 'modes' are labelled by n_8 . From the form of this equation one sees that only the product k w, and not k and w independently, govern

the behavior of the solutions. Note that Eq.[8.4] has the property that the even modes and odd modes evolve independently of each other (because $\sqrt[6]{\csc^2(x)}$ is an even function of x).

According to the theory of systems of linear differential equations a solution associated to Eq.[8.4] exists that is expressible in the form

$$h_{n_8}(\tau) = \begin{cases} \sum_{j=-\infty}^{\infty} h_{n_8;j} \exp\left(\left(-\sqrt{a_2} + j\right)\tau\right) & \text{for } n_8 \text{ even} \\ \sum_{j=-\infty}^{\infty} h_{n_8;j} \exp\left(\left(-\sqrt{a_1} + j\right)\tau\right) & \text{for } n_8 \text{ odd} \end{cases},$$
(8.5)

where the $h_{n_8;j}$, a_1 , a_2 are constants to be determined (see, for example, Ref.[25] Equation[7], page 177). We substitute Eq.[8.5] into Eq.[8.4] and then demand that the coefficient of each $e^{j\tau}$, $j = -\infty \dots \infty$, vanish, where $a = a_1$ if n_8 is odd and $a = a_2$ if n_8 is even. This yields the system of linear equations for the $h_{n_8;j}$ given by

$$0 = -h_{n_{8};j} \left[\frac{3}{20} + \frac{m^{2}}{2} + n_{8}^{2} + \frac{1}{9} \left(j - \sqrt{a} \right)^{2} \right] + \sqrt{Y} \left(h_{n_{8};j-1} - h_{n_{8};j+1} \right) + \sqrt{Y} \sum_{m_{8}=1}^{\infty} \frac{\left(-1 \right)^{m_{8}} \Gamma \left(\frac{5}{6} \right)^{2}}{\Gamma \left(\frac{5}{6} - m_{8} \right) \Gamma \left(\frac{5}{6} + m_{8} \right)} \left(h_{n_{8}-2m_{8};j-1} - h_{n_{8}-2m_{8};j+1} + h_{n_{8}+2m_{8};j-1} - h_{n_{8}+2m_{8};j+1} \right)$$

$$(8.6)$$

where $a = a_1$ if n_8 is odd and $a = a_2$ if n_8 is even.

,

Note that if the $h_{n_8;j}$ satisfy the three term recurrence relation

$$\sqrt{Y}\left(h_{n_{8};j-1} - h_{n_{8};j+1}\right) = \beta_{n_{8}} \left[\frac{3}{20} + \frac{m^{2}}{2} + n_{8}^{2} + \frac{1}{9}\left(j - \sqrt{a}\right)^{2}\right] h_{n_{8};j}, \quad (8.7)$$

where the β_{n_8} are also to be determined, then

$$0 = \sum_{m_8 = -\infty}^{\infty} \frac{(-1)^{m_8} \Gamma\left(\frac{5}{6}\right)^2}{\Gamma\left(\frac{5}{6} - m_8\right) \Gamma\left(\frac{5}{6} + m_8\right)} \left[\frac{3}{20} + \frac{m^2}{2} + (n_8 - 2m_8)^2 + \frac{1}{9} \left(j - \sqrt{a}\right)^2\right] \times (\beta_{n_8 - 2m_8} - \delta_{m_8,0}) h_{n_8 - 2m_8;j},$$
(8.8)

which is block diagonal in the temporal index j.

Except at the irregular singular points $x = 0 \pm \pi$, both the massive complex scalar wave function ψ and its quantum fluctuation $\delta \psi$ are expressible in the form given by

$$\Psi(\tau, x) = C_1 \Psi_{\text{odd}}(\tau, x) + C_2 \Psi_{\text{even}}(\tau, x)$$

$$\Psi_{\text{even}}(\tau, x) = \sum_{n_8 = -\infty, n_8}^{\infty} \sum_{\text{even } j = -\infty}^{\infty} h_{n_8;j} \exp\left(\left(-\sqrt{a_2} + j\right)\tau + in_8 x\right)$$

$$\Psi_{\text{odd}}(\tau, x) = \sum_{n_8 = -\infty, n_8}^{\infty} \sum_{\text{odd } j = -\infty}^{\infty} h_{n_8;j} \exp\left(\left(-\sqrt{a_1} + j\right)\tau + in_8 x\right), \quad (8.9)$$

where the $h_{n_8;j}$ satisfy Eq.[8.6]. Of course, the constants C_1 and C_2 are redundant and employed only for emphasis. The scalar wave function $\Psi(\tau, x)$ satisfies

$$0 = -\Psi^{(2,0)}(\tau, x) + \frac{1}{9}\Psi^{(0,2)}(\tau, x) + \left(\frac{1}{2}m^2 + \frac{3}{20} - 2\sqrt{Y}\sinh(\tau)\sqrt[6]{\csc^2(x)}\right)\Psi(\tau, x).$$
(8.10)

The mass parameter m is in general different for $\Psi = \psi$ and $\Psi = \delta \psi$.

A sequence of approximate solutions to Eq.[8.6] may be defined as follows: Let $n \in \mathbb{N}$, the natural numbers (excluding 0). n will be called the index of the approximation. For each positive integer $n \in \mathbb{N} = 1, 2, \ldots$ we define an approximate solution to Eq.[8.6] in terms of $(2n+1)^2$ mode coefficients ${}^{(n)}h_{n_8;j}$, $-n \leq n_8, j \leq n$ that, in the limit $n \to \infty$, will satisfy Eq.[8.6] exactly. Eq.[8.6] defines a system of linear equations for all of the $h_{n_8;j}$ mode coefficients. In order to define a finite approximate solution set, we set all of the mode coefficients $h_{n_8;j} = 0$ for $-\infty < n_8, j < -n$ and $n < n_8, j < \infty$. This results in two homogeneous systems of linear equations for the $(2n+1)^2$ mode coefficients $^{(n)}h_{n_8;j}$, $-n \leq n_8, j \leq n$, one system for the even n_8 modes and one system for the odd n_8 modes (since the even n_8 modes and odd n_8 modes evolve independently of each other). To obtain non-trivial solutions to these two systems we solve for the complex values of a_1 and a_2 that make the determinants of their respective coefficient matrices vanish. Two dispersion relations thereby arise: one that relates n and both m and kw, the product of the magnitudes of the momentum wave vectors (\vec{k}, \vec{w}) , to the complex parameter a_1 for the odd n_8 modes, and one that relates n and (m, kw) to the complex parameter a_2 for the even n_8 modes.

Recall that in this paper "stable" means that solutions $\Psi = \delta \psi$ of Eq.[8.10] possess a norm that is non-exponentially increasing as τ increases. Stable modes exist when $-\text{Real}(\sqrt{a_1}) + n$ and $-\text{Real}(\sqrt{a_2}) + n$ are both negative, or zero. To see this first define the corresponding approximate solution ${}^{(n)}\Psi(\tau, x)$ to Eq.[8.10] as

$${}^{(n)}\Psi_{\text{odd}}(\tau,x) = \sum_{n_8=-n+evn(n), \text{ step 2}}^n \sum_{j=-n}^n {}^{(n)}h_{n_8;j} e^{[(-\sqrt{a_1}+j)\tau + in_8x]}$$

$${}^{(n)}\Psi_{\text{even}}(\tau,x) = \sum_{n_8=-n+odd(n), \text{ step 2}}^n \sum_{j=-n}^n {}^{(n)}h_{n_8;j} e^{[(-\sqrt{a_2}+j)\tau + in_8x]}$$

$$\cdot \qquad (8.11)$$

Here we have defined two simple functions on \mathbb{N} , which for $n \in \mathbb{N}$, are given by

$$evn(n) = \begin{cases} 1 & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases}, \quad odd(n) = \begin{cases} 0 & \text{if } n \text{ is even} \\ 1 & \text{if } n \text{ is odd} \end{cases}.$$
 (8.12)

One sees that, of all of the $(2n+1)^2$ wave modes, each of the form

 ${}^{(n)}h_{n_8;j}e^{\left[(-\sqrt{a}+j)\tau+in_8x\right]}$, the j=n waves increase at the greatest rate for given a. If $-\operatorname{Real}(\sqrt{a_1})+n$ and $-\operatorname{Real}(\sqrt{a_2})+n$ are both negative, or zero, then the j=n waves have non-increasing norms as time τ increases (either because w increases, t increases or k decreases), so that the norm of ${}^{(n)}\Psi_{\rm odd}(\tau,x)$ and ${}^{(n)}\Psi_{\rm even}(\tau,x)$ also have non-increasing norms as time τ increases (here $\Psi = \delta \psi$). These stable modes are associated to so-called "stable zeros" (a_1, a_2) of the determinants of the coefficient matrices for the odd n_8 labeled modes and the even n_8 labeled modes, respectively. The "unstable zeros" (a_1, a_2) of the determinants of the coefficient matrices driven by the momenta associated to the three extra time dimensions, and do not yield physical solutions of the field equations.

In passing we remark that the error of an approximation may be provisionally defined to be

$$\operatorname{error}[{}^{(n)}\Psi(\tau,x)] = -{}^{(n)}\Psi^{(2,0)}(\tau,x) + \frac{1}{9}{}^{(n)}\Psi^{(0,2)}(\tau,x) + \left(\frac{1}{2}m^2 + \frac{3}{20} - 2\sqrt{Y}\sinh(\tau)\sqrt[6]{\operatorname{csc}^2(x)}\right){}^{(n)}\Psi(\tau,x). \quad (8.13)$$

However since ${}^{(n)}\Psi(\tau, x)$ contains only a finite number of Fourier modes it is reasonable to make a substitution based on Eq.[7.15], $\sqrt[6]{\csc^2(x)} \rightarrow \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} \sum_{m8=-[\frac{n}{2}]}^{[\frac{n}{2}]} \frac{e^{i(2m8)x}\left((-1)^{m8}\Gamma(\frac{5}{6})^2\right)}{\Gamma(\frac{5}{6}-m8)\Gamma(m8+\frac{5}{6})}$, in Eq.[8.13] when computing the numerical error $\left([\frac{n}{2}]\right)$ gives the greatest integer less than or equal to $\frac{n}{2}$). Since the error is expected to be large for small n we defer error analysis.

9. COEFFICIENT MATRICES FOR THE ${}^{(n)}h_{n_8\,;j}$ MODE COEFFICIENTS IN EQ.[8.6]

Let

$$M = \frac{1}{2}m^2 + \frac{3}{20},\tag{9.1}$$

and recall that $\sqrt{Y} = \frac{1}{2}k w \sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2}$. Fix $n \in \mathbb{N}$; we group the $(2n+1)^2 = n (2n+1) + (n+1) (2n+1)$ mode coefficients ${}^{(n)}h_{n_8;j}$, $-n \leq n_8, j \leq n$, into two vectors $\overrightarrow{h}_{\text{even}}$ and

 \overrightarrow{h}_{odd} according to whether n_8 is even or odd. When n is an odd positive integer then \overrightarrow{h}_{odd} has dimension (n+1)(2n+1) = (n + odd(n))(2n+1) and $\overrightarrow{h}_{even}$ has dimension n(2n+1) = (n + evn(n))(2n+1). When n is an even positive integer then \overrightarrow{h}_{odd} has dimension n(2n+1) = (n + odd(n))(2n+1) and $\overrightarrow{h}_{even}$ has dimension (n+1)(2n+1) = (n + evn(n))(2n+1). Note that \overrightarrow{h}_{odd} always has an even number of components and that $\overrightarrow{h}_{even}$ always has an odd number of components.

Explicitly, the i^{th} component of the vector \overrightarrow{h} associated to ${}^{(n)}h_{n_8;j}$ has index

$$i = \frac{1}{2}(2n+1)(n+n_8 - evn(n) odd(n_8) - odd(n) evn(n_8)) + j + n + 1$$
(9.2)

so that

$$h_{\text{even}}\left(\frac{1}{2}(2n+1)\left(n+n_8 - odd(n)\right) + j + n + 1\right) = {}^{(n)}h_{n_8;j}, \text{ where } n_8 \text{ is even}$$
(9.3)

and

$$h_{\text{odd}}\left(\frac{1}{2}(2n+1)\left(n+n_8 - evn(n)\right) + j + n + 1\right) = {}^{(n)}h_{n_8;j}, \text{ where } n_8 \text{ is odd.}$$
(9.4)

Here $h(i) = \left\{ \overrightarrow{h} \right\}_i = i^{\text{th}} \text{ component of } \overrightarrow{h}$.

The inverse map $(n; i, parity) \mapsto (n_8, j)$ is defined as follows: Let

 $\mathbb{N} \ni parity = \begin{cases} 1 & \text{if indices are extracted from } \overrightarrow{h}_{\text{odd}} \\ 2 & \text{if indices are extracted from } \overrightarrow{h}_{\text{even}} \end{cases}.$

The inverse map $(n; i, \text{parity}) \mapsto (n_8, j)$ is given by $j = (i-1) \mod (2n+1) - n$ and $n_8 = 2\left[\frac{i-1}{2n+1}\right] - n + odd(parity)evn(n) + evn(parity)odd(n)$, where $\left[\frac{i-1}{2n+1}\right]$ gives the greatest integer less than or equal to $\frac{i-1}{2n+1}$.

Let μ denote the minimum value of n_8 for either $\overrightarrow{h}_{even}$ or \overrightarrow{h}_{odd} (thus, μ is either equal to -n or -n + 1). The components of these two vectors are ordered according to $((n_8 = \mu, j = -n), (n_8 = \mu, j = -n + 1), \dots, (n_8 = \mu, j = n), (n_8 = \mu + 2, j = -n), (n_8 = \mu + 2, j = -n + 1), \dots, (n_8 = \mu + 2, j = -n), (n_8 = -\mu, j = -n), (n_8 = -\mu, j = -n + 1), \dots, (n_8 = -\mu, j = -n), (n_8 = -\mu, j = -n + 1), \dots, (n_8 = -\mu, j = -n), (n_8 = -\mu, j = -n + 1), \dots, (n_8 = -\mu, j = -n), (n_8 = -\mu, j = -n + 1), \dots, (n_8 = -\mu, j = -n), (n_8 = -\mu, j = -n + 1), \dots, (n_8 = -\mu, j = -n), (n_8 = -\mu, j = -$

With respect the obvious canonical bases each coefficient matrix is a square matrix and is equal to the sum D + A of a diagonal matrix D, which carries the a and M dependence, and an antisymmetric matrix $\sqrt{Y} A$, where $\widetilde{A} = -A$ (the tilde denotes the transpose). For a given $n \in \mathbb{N}$, the coefficient matrix ${}^{(n)}D_{\text{even}} + {}^{(n)}A_{\text{even}}$ corresponding to the even n_8 mode components has dimensions equal to $[(n + even(n)) (2n + 1)] \times [(n + even(n)) (2n + 1)],$ and thus always has an odd number of rows and columns. The coefficient matrix

 ${}^{(n)}D_{\text{odd}} + {}^{(n)}A_{\text{odd}}$ corresponding to the odd n_8 mode components has dimensions equal to $[(n + odd(n)) (2n + 1)] \times [(n + odd(n)) (2n + 1)]$, and thus always has an even number of rows and columns.

The diagonal matrix ${}^{(n)}D$ has non-zero matrix elements $-\left(M + n_8^2 + \frac{1}{9}(j - \sqrt{a})^2\right)$, arranged (as described above) along the main diagonal in blocks, which are labelled by n_8 , of the 2n + 1 possible j values.

Because ${}^{(n)}A_{\text{even}}$ always has an odd number of rows and columns it possesses a vanishing determinant, det ${}^{(n)}A_{\text{even}} \equiv 0$. The antisymmetric ${}^{(n)}A_{\text{odd}}$ corresponding to the odd n_8 mode components always has an even number of rows and columns, and hence a possibly non-vanishing determinant. However for this case the rows

 $^{(n)}A_{\text{odd}}(i), i = 1, 2, \dots, [n + odd(n)] \times (2n + 1)$ of $^{(n)}A_{\text{odd}}$ are not linearly independent and satisfy $^{(n+odd(n))(2n+1)}$

$${}^{(n)}A_{\rm odd}(1) + {}^{(n)}A_{\rm odd}(3) + \dots + {}^{(n)}A_{\rm odd}(2n+1) = \underbrace{(0,0,\dots,0)}^{(n+{\rm odd}(n))(2n+1)}.$$
(9.5)

Hence for this case A_{odd} also has a vanishing determinant, det $\binom{(n)}{A_{\text{odd}}} = 0$.

The antisymmetric matrix ${}^{(n)}A$ is related to an antisymmetric Toeplitz matrix ${}^{(n)}T = -{}^{(n)}T$ whose non-zero upper triangular matrix elements are given by ${}^{(n)}T_{i,i+\ell(2n+1)+1} = -\gamma_{\ell}, \ \ell = 0, 1, \dots, \text{ and}$ ${}^{(n)}T_{i,i+\ell(2n+1)-1} = \gamma_{\ell}, \ \ell = 1, \dots, \text{ where the } \gamma_{\ell} \text{ are defined in Eq.}[7.8] \text{ and } i + \ell \ (2n+1) + 1 \leq [(n + odd(n)) \ (2n + 1)] \text{ for } {}^{(n)}T \text{ corresponding to the } {}^{(n)}A_{\text{odd}} \text{ associated to the odd } n_8 \text{ mode} \text{ components and}$ $i + \ell \ (2n + 1) + 1 \leq [(n + evn(n)) \ (2n + 1)] \text{ for } {}^{(n)}T \text{ corresponding to the } {}^{(n)}A_{\text{even}} \text{ associated to the odd } n_8 \text{ mode} \text{ components and}$

to the even n_8 mode components. The remaining matrix elements of ${}^{(n)}T$ are either dictated by antisymmetry, or are zero.

The matrix elements of ${}^{(n)}A$ equal the matrix elements of ${}^{(n)}T$ except that they are punctuated by additional zeros along the super diagonals of ${}^{(n)}T$. ${}^{(n)}A_{\text{even}}$, $n \geq 2$, has $N_{\text{additional zeros}} = N_{AZ} = 2([n + evn(n)]^2 - 1)$ replacements of non-zero matrix elements of ${}^{(n)}T$ by zero along its super diagonals. ${}^{(n)}A_{\text{odd}}$, $n \geq 1$, has $N_{AZ} = 2([n + odd(n)]^2 - 1)$ zero replacements of non-zero matrix elements of ${}^{(n)}T$ by zero along its super diagonals.

The specification of the row and column indices (r, c) of a particular matrix element ${}^{(n)}A_{rc}$ of ${}^{(n)}A$ that is zero, when ${}^{(n)}T_{rc}$ is not zero, is somewhat involved, but may be

described in stages. Let the dimensions of ${}^{(n)}A$ be $L \times L$. Define a mapping of the twodimensional array ${}^{(n)}A_{rc}$, r, c = 1, ..., L to a one-dimensional array

 ${}^{(n)}B_i, i = 1, \dots, L^2$ by setting ${}^{(n)}B_i = {}^{(n)}B_{(r-1)L+c} = {}^{(n)}A_{rc}$. Formally, i = (r-1)L+c. The inverse map $(n; i, \text{parity}) \mapsto (r, c)$ is given by $r = (i-1) \mod (L) + 1$ and $c = \left[\frac{i-1}{L}\right] + 1$, where L = (2n+1) (n + evn(parity) evn(n) + odd(parity) odd(n)).

Assume that the row and column indices $\{(r,c)_j\}_{j=1}^{N_{AZ}}$ of the N_{AZ} matrix elements whose non-zero ${}^{(n)}T_{r_j c_j}$ values are to be replaced by zero are known, and that their corresponding $\{i_j\}_{j=1}^{N_{AZ}}$ values have been computed; sort the $\{i_j\}$ values in ascending order. Define the relative displacement coordinates $\{I_j^{\text{RDC}}\}_{j=1}^{N_{AZ}}$ by $I_1^{\text{RDC}} = i_1$, $I_2^{\text{RDC}} = i_2 - i_1$, ..., $I_j^{\text{RDC}} = i_j - i_{j-1}$, $j = 2, \ldots, N_{AZ}$. The inverse map is simply $i_j = \sum_{h=1}^{j} I_h^{\text{RDC}}$.

We find that the $\{I_j^{\text{RDC}}\}_{j=1}^{N_{AZ}}$ are given by two sequences, one for the odd n_8 mode components and one for the even n_8 mode components. Let $N_j = 2n + 1$ denote the number of *j*-values in the *n*th-order approximation. Let $\{N_j\}$ [*h*] denote the finite sequence $\{N_j\}$ [*h*] = $\{N_j, N_j, \ldots, N_j\}$ with *h* elements, and let $\{S\}$ [*h*] denote a finite sequence repeated *h* times.

For the case of the odd n_8 mode components let

N = n - evn(n) and $\alpha = \frac{1}{2} \left(2n \left(2n \left(2n - (-1)^n + 1\right) - 1\right) + (-1)^n + 1\right)$; the sequence $\left\{I_j^{\text{RDC}}\right\}_{j=1}^{N_{AZ}}$ for the odd n_8 mode components is

$$\left\{ I_{\text{odd }j}^{\text{RDC}} \right\}_{j=1}^{N_{AZ}} = \left\{ \left\{ N_j \right\} [N], \alpha + N_j, \left\{ \left\{ N_j \right\} [N], 1 + 4n, \left\{ N_j \right\} [N], \alpha \right\} [N - 1], \\ \left\{ N_j \right\} [N], \alpha + N_j, \left\{ N_j \right\} [N] \right\}.$$

$$(9.6)$$

For the case of the even n_8 mode components let

N = n - odd(n) and $\beta = \frac{1}{2} \left(2n \left(2n \left(2n + (-1)^n + 1\right) - 1\right) - (-1)^n + 1\right)$; the sequence $\left\{I_j^{\text{RDC}}\right\}_{j=1}^{N_{AZ}}$ for the even n_8 mode components is

$$\left\{ I_{\text{even } j}^{\text{RDC}} \right\}_{j=1}^{N_{AZ}} = \left\{ \left\{ N_{j} \right\} [N], \beta + N_{j}, \left\{ \left\{ N_{j} \right\} [N], 1 + 4n, \left\{ N_{j} \right\} [N], \beta \right\} [N - 1], \\ \left\{ N_{j} \right\} [N], \beta + N_{j}, \left\{ N_{j} \right\} [N] \right\}.$$

$$(9.7)$$

Only the first N_{AZ} sequence elements are to be employed when evaluating the the relative displacement coordinates from the previous two sequence formulas.

10. THEOREM: EXISTENCE OF STABLE SOLUTIONS

We record the

Theorem 10.1. Let $n \in \mathbb{N}$ be the index of an approximation. Define an approximate solution as described above to Eq.[8.6] in terms of $(2n+1)^2$ mode coefficients ${}^{(n)}h_{n_8;j}$, $-n \leq n_8, j \leq n$.

Let ${}^{(n)}D_{n_8 \ odd}(a_1, Y, M)$ denote the determinant of the coefficient matrix of the ${}^{(n)}h_{n_8;j}$ for the odd n_8 labeled modes and ${}^{(n)}D_{n_8 \ even}(a_2, Y, M)$ denote the determinant of the coefficient matrix of the ${}^{(n)}h_{n_8;j}$ for the even n_8 labeled modes.

Let n_{a_1} denote the number of stable zeros of ${}^{(n)}D_{n_8 \ odd}(a_1, Y, M)$ and n_{a_2} denote the number of stable zeros of ${}^{(n)}D_{n_8 \ even}(a_2, Y, M)$.

Then

- 1. ${}^{(n)}D_{n_8 \ odd}(a_1, Y, M)$ is a trivariate polynomial in (a_1, Y, M) with rational coefficients;
- 2. ${}^{(n)}D_{n_8}$ even (a_2, Y, M) is a trivariate polynomial in (a_2, Y, M) with rational coefficients;
- 3. For each positive integer $n \in \mathbb{N} = 1, 2, ...,$ stable zeros (a_1, a_2) of the the determinants of the coefficient matrices ${}^{(n)}D_{n_8} {}_{odd}(a_1, Y, M)$ and ${}^{(n)}D_{n_8} {}_{even}(a_2, Y, M)$ exist that satisfy $-Real(\sqrt{a_1}) + n \leq 0$ and $-Real(\sqrt{a_2}) + n \leq 0$;
- 4. if n is an odd positive integer then
 - (a) ${}^{(n)}D_{n_8 \ odd}(a_1, Y, M)$ is of degree $(n+1)(2 \ n+1)$ in a_1 and M, and of degree n(n+1)in Y
 - (b) ${}^{(n)}D_{n_8 even}(a_2, Y, M)$ is of degree n(2 n+1) in a_2 and M, and of degree n^2 in Y
 - If Y = 1 then

 n_{a1} = 2 n+2
 n_{a2} = 2 n

 If Y ≠ 1 then

 n_{a1} ≥ 2n + 2
 n_{a2} ≥ 2n
- 5. if n is an even positive integer then

- (a) ${}^{(n)}D_{n_8 \ odd}(a_1, Y, M)$ is of degree $n(2 \ n+1)$ in a_1 and M, and of degree n^2 in Y
- (b) ${}^{(n)}D_{n_8 even}(a_2, Y, M)$ is of degree (n+1)(2 n+1) in a_2 and M, and of degree n(n+1) in Y
 - If Y = 1 then
 - *i.* $n_{a_1} = 2 n$
 - *ii.* $n_{a_2} = 2 n + 2$
 - If $Y \neq 1$ then
 - *i.* $n_{a_1} \ge 2n$
 - *ii.* $n_{a_2} \ge 2n+2$

This relationship holds in the limit $n \to \infty$ yielding exact stable solutions to Eq.[8.6], and therefore to Eq.[8.10].

Partial Proof of Theorem[10.1]: For n = 1, ..., 9 the proof has been accomplished by direct calculation and is summarized in Table[10.2].

In general, parts [1] and [2] may be proven by employing Theorem 13.7.3. of [26], which gives the expansion for the determinant |D + B| of the sum of two $n \times n$ matrices D and B for the special case where one of the matrices is diagonal. We quote Theorem 13.7.3. [26]: Let B represent an $n \times n$ matrix, and let D represent an $n \times n$ diagonal matrix whose diagonal elements are d_1, \ldots, d_n . Then,

$$|D+B| = \sum_{\{i_1, i_2, \dots, i_r\}} d_{i_1} \cdot d_{i_2} \cdots \cdot d_{i_r} |B^{\{i_1, i_2, \dots, i_r\}}|.$$
(10.1)

where $\{i_1, i_2, \ldots, i_r\}$ is a subset of the first n positive integers $1, \ldots, n$ (and the summation is over all 2^n such subsets) and where $B^{\{i_1, i_2, \ldots, i_r\}}$ is the $(n-r) \times (n-r)$ principal submatrix of B obtained by striking out the i_1, i_2, \ldots, i_r th rows and columns. [The term in the sum [10.1] corresponding to the empty set is to be interpreted as |B|, and the term corresponding to the set $\{1, 2, \ldots, n\}$ is to be interpreted as $|D| = d_1 \cdot d_2 \cdots \cdot d_n$.]

In our case the matrix elements of D and B are defined in Section [9]. $B = \sqrt{Y} A$ is antisymmetric. Note that all principal submatrices $B^{\{i_1, i_2, \dots, i_r\}}$ of the antisymmetric matrix $B = \sqrt{Y} A$ are also antisymmetric.

The general proof of this theorem is a work in progress. Thus, for all other cases than those explicitly noted, the Theorem devolves to a Conjecture. Note that according to Theorem[10.1], for given n, the degrees of the $\{a_1, a_2\}$ equal the numbers of independent symmetric and anti-symmetric matrix elements of a $(2n + 1) \times (2n + 1)$ real matrix. The sum of the degrees of the (a_1, a_2) equals $(2n + 1)^2$.

Table [10.2] summarizes the degrees of the polynomials ${}^{(n)}D_{n_8 \text{ odd}}(a_1, Y, M)$ and ${}^{(n)}D_{n_8 \text{ even}}(a_2, Y, M)$ found by direct calculation for $n = 1, \ldots, 9$. The degree of $M = \frac{1}{2}m^2 + \frac{3}{20}$ is found to always equal the degree of a_1 for ${}^{(n)}D_{n_8 \text{ odd}}(a_1, Y, M)$ and always equal the degree of a_2 for ${}^{(n)}D_{n_8 \text{ even}}(a_2, Y, M)$, and therefore is not displayed in this table.

0			1			
	${}^{(n)}D_{n_8 \text{ odd}}(a_1, Y, M)$		$^{(n)}D_{n_8 \text{ even}}(a_2, Y, M)$		# STABLE	
Polynomial	$a_1{}^{\alpha}$	Y^{β}	$a_2{}^{\gamma}$	Y^{δ}	for Y=1	
Degree n	α	β	γ	δ	n_{a_1}	<i>n</i> _{<i>a</i>₂}
1	6	2	3	1	4	2
2	10	4	15	6	4	6
3	28	12	21	9	8	6
4	36	16	45	20	8	10
5	66	30	55	25	12	10
6	78	36	91	42	12	14
7	120	56	105	49	16	14
8	136	64	153	72	16	18
9	190	90	171	81	20	18
	CONJECTURE					
odd n	(n+1) (2 n+1)	n (n+1)	n (2 n+1)	n^2	2 n+2	2 n
even n	n (2 n+1)	n^2	(n+1) (2 n+1)	n (n+1)	2 n	2 n+2

(10.2)

The significance of this theorem is that each pair of "stable" zeros (a_1, a_2) defines a pair of stable propagation modes for a quantum fluctuation $\delta\psi$. The numbers of these modes increases with n as $n \to \infty$ yielding a possibly complete set of decaying quasi-normal modes that one may employ in cosmological perturbation calculations. Quantitative calculations of the quantum fluctuations that occur during inflation are deferred until we have an analytical approximation for a complete set of decaying quasi-normal mode functions for the massive complex scalar field.

10.1. Examples for Y = 1, m = 0

The results of an example for n = 6, m = 0 and $\sqrt{Y} = \frac{1}{2}k w \sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 1$ are displayed in Figures [2] and [3], which present pairs of plots (τ dependence, for fixed x) and (xdependence, for fixed τ) of the stable scalar wave functions ${}^{(6)}\psi(\tau, x)$ of Eq.[8.11]. For n = 6there are $169 = (2n+1)^2$ mode coefficients, 78 with odd n_8 and 91 with even n_8 . Roughly 352 digits of precision in the solved-for values of a_1 and a_2 are required to make the determinants of the coefficient matrices vanish to double precision accuracy. The relevant values (rounded to 9 digits) of a_1 and a_2 for each plot are listed in the yellow box associated to the plot; space limitations do not permit their display in a table (if one is reading the pdf version of the manuscript then zooming-in may be required, unless you have remarkable eyesight). Red curves correspond to the real part of the solutions and blue curves and correspond to the imaginary part.

The results of an example for n = 7, m = 0 and $\sqrt{Y} = \frac{1}{2}k w \sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 1$ are displayed in Figures [4] and [5], which also present pairs of plots (τ dependence, for fixed x) and (xdependence, for fixed τ) of the stable scalar wave functions ${}^{(7)}\psi(\tau, x)$ of Eq.[8.11]. For n = 7there are $225 = (2n + 1)^2$ mode coefficients, 120 with odd n_8 and 105 with even n_8 . In this case, about 400 digits of precision in the solved-for values of a_1 and a_2 are required to make the determinants of the coefficient matrices vanish to double precision accuracy. The relevant values of a_1 and a_2 for a plot are again listed in the yellow box (zooming-in may also be required, unless you have remarkable eyesight). Red curves correspond to the real part of the solutions and blue curves and correspond to the imaginary part.

The results of an example for n = 8, m = 0 and $\sqrt{Y} = \frac{1}{2}k w \sqrt[3]{2} \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 1$ are displayed in Figures [6] and [7], which present pairs of plots (τ dependence, for fixed x) and (xdependence, for fixed τ) of the stable scalar wave functions ${}^{(8)}\psi(\tau, x)$ of Eq.[8.11]. For n = 8there are $361 = (2n + 1)^2$ mode coefficients, 190 with odd n_8 and 171 with even n_8 . In this case, about 400 digits of precision in the solved-for values of a_1 and a_2 are required to make the determinants of the coefficient matrices vanish to double precision accuracy. The relevant values of a_1 and a_2 for a plot are again listed in the yellow box (zooming-in may also be required, unless you have remarkable eyesight). Red curves correspond to the real part of the solutions and blue curves and correspond to the imaginary part.

Qualitatively, corresponding n = 6, n = 7 and n = 8 plots agree, although there are

quantitative differences due to the low orders of approximation. The overall number of modes increases by four for each increment in n. In the limit $n \to \infty$ it is reasonable to hope that this procedure generates a "complete" set of stable modes, in the sense that any **stable** wavefunction can be expressed as a linear combination of these modes.

We arrive at a picture of stable quasi-normal modes that decay during inflation. According to Theorem[10.1] this model possesses an infinite number of stable quasi-normal modes, such that the momenta associated with the extra time dimensions do not create instability.

11. CLASS E_1 SOLUTIONS OF THE EINSTEIN EQUATIONS

The field equations admit a second class of solutions that is periodic in x^8 and parameterized by ξ , $-\frac{1}{\sqrt{5}} < \xi < \frac{1}{\sqrt{5}}$, and which has a time dependent ℓ . In this case the scale factors are

$$a = a_{1} \left[\cosh\left(x^{4}\sqrt{\frac{\Lambda}{3}}\right) \right]^{\frac{1}{6}(1+5\xi)} \\ \times \left[\tan^{2}\left(\frac{1}{2}\sqrt{\frac{5}{3}}\sqrt{\Lambda}x^{8}\right) \right]^{\pm\sqrt{\frac{1}{30}}\sqrt{1-5\xi^{2}}} \left[\sin^{2}\left(\sqrt{\frac{5}{3}}\sqrt{\Lambda}x^{8}\right) \right]^{\frac{1}{12}(1-\xi)} \\ b = a_{1} \left[\cosh\left(x^{4}\sqrt{\frac{\Lambda}{3}}\right) \right]^{\frac{1}{6}(1-5\xi)} \\ \times \left[\tan^{2}\left(\frac{1}{2}\sqrt{\frac{5}{3}}\sqrt{\Lambda}x^{8}\right) \right]^{\mp\sqrt{\frac{1}{30}}\sqrt{1-5\xi^{2}}} \left[\sin^{2}\left(\sqrt{\frac{5}{3}}\sqrt{\Lambda}x^{8}\right) \right]^{\frac{1}{12}(1+\xi)},$$
(11.1)

where a_1 and b_1 are constants. Periodicity in x^8 requires that Λ be quantized,

$$\Lambda = \Lambda_1 = \frac{3}{5} = \frac{6}{5}\Lambda_0 \tag{11.2}$$

We also find that

$$\varphi = \frac{\sqrt{1 - 5\xi^2} \ln \left[\cosh^5 \left(\frac{\sqrt{\Lambda}x^4}{\sqrt{3}} \right) \csc^2 \left(\sqrt{\frac{5}{3}} \sqrt{\Lambda}x^8 \right) \right]}{2\sqrt{30}} \mp \xi \ln \left[\tan^2 \left(\frac{1}{2} \sqrt{\frac{5}{3}} \sqrt{\Lambda}x^8 \right) \right]$$
$$\ell_1 = \mp \frac{5\sqrt{\Lambda}\xi \tanh \left(\frac{\sqrt{\Lambda}x^4}{\sqrt{3}} \right)}{3\sqrt{3}} = \pm 5\xi \sqrt{\frac{2}{3}} \ell_0 \tanh \left(x^4 \sqrt{\frac{\Lambda}{3}} \right). \tag{11.3}$$

To exclude collapsing universe solutions, which correspond to $\frac{\partial}{\partial x^4} \ln(a) < 0$, and solutions for universes with multiple macroscopic times, which correspond to $\frac{\partial}{\partial x^4} \ln(b) \ge 0$, from the set of physical solutions we restrict ξ to the interval $\frac{1}{5} < \xi < \frac{1}{\sqrt{5}}$.

For this case the volume element $d\Omega = d\tau \cosh\left(x^4\sqrt{\frac{\Lambda}{3}}\right) \left|\sin\left(x^8\sqrt{\frac{5}{3}}\Lambda\right)\right|$, is independent of both ξ and the choice of \pm signs. The temperature history during inflation is non-trivial in this case. This and the possible stability of quantum fluctuations are of interest, but not investigated further in this paper.

12. CONCLUSION

We have presented a model of the very early universe that possesses equal numbers of space and time dimensions. In this model, after "inflation" the observable physical macroscopic world appears to a classical observer to be a homogeneous, isotropic universe with three space dimensions and one time dimension. Notwithstanding the often-expressed concern that the momenta associated to extra time dimensions source destabilizing quantum fluctuations, we have shown that the physical solutions $\delta \psi$ to Eq.[8.10] that propagate on the class E_0 solution to the coupled Einstein-inflaton equations possess decaying quasi-normal modes, and have no instabilities that are sourced by the momenta \vec{w} associated to the extra time dimensions.

This model and the class E_0 solution to the coupled Einstein-inflaton field equations provide simple answers to at least three of the important questions about the early universe, namely, how to the define the inflaton potential [27], how to the define the inflaton mass and how to resolve the issue of reheating [14]. At the classical level the answers to the first two questions are zero inflaton potential and zero inflaton mass. Computable quantum fluctuations may shift the classical $V_{\text{inflaton}} = 0$ value and the classical $m_{\text{inflaton}} = 0$ value. How these values behave under renormalization are important questions, but beyond the scope of the present paper. The answer to the last question is that there is no obvious need for reheating, since inflation/deflation is isothermal in this model for the class E_0 solution (the ground state solution to the Einstein-inflaton field equations).

We have also proved that a well known so-called "single time" theorem [20] **does not** apply to our model.

Do closed timelike curves exist in this model? If we assume that an "arrow of time" exists

for each timelike dimension then this model does not admit unphysical closed timelike curves. The reason is simply that the tangent vector to any such curve is not everywhere futuredirected, if it is timelike. Let $x^{\alpha} = x^{\alpha}(\xi)$, $0 \leq \xi \leq 1$, with $x^{\alpha}(0) = x^{\alpha}(1)$, parameterize a timelike closed curve in $\mathbb{X}_{4,4}$; then $\frac{\partial}{\partial \xi} x^{\alpha}(\xi)$ for $0 \leq \xi \leq 1$ is not everywhere either future-directed timelike, null, or spacelike. For example, a closed timelike curve within the (x^4, x^5, x^6, x^7) 4-manifold that traverses the x^4 dimension is not physical classically because such a curve must move both forward and backward in physical time x^4 .

If we do not assume that an "arrow of time" exists for each extra timelike dimension then, if deflation for the scale factor associated to the extra time dimensions proceeded for sufficiently long Δx^4 interval(s), then observationally the effects of closed timelike curves might be described as a difficulty in assigning classical spacetime coordinates to certain events. A discussion of this is beyond the scope of this paper.

In this model as presently formulated, inflation also goes on forever. This "feature" obviously must be eliminated from the model. This model must be generalized in a way that leads to inflaton decay and also terminates inflation. Possible generalizations include: [1] modifing the inflaton Lagrangian to include a self-coupling; and [2] coupling the inflaton to a Higgs doublet (before the Higgs develops a non-zero vacuum expectation value, in order to keep the inflaton massless) through a Yukawa interaction, while also generalizing the total Lagrangian in the model to incorporate Standard Model quarks, leptons and gauge bosons.

12.1. The scale factors vanish on a set of measure zero

Typically and approximately, inflation scenarios inflate a scale of the size of one billionth the present radius of a proton to the size of the present radius of a marble or a grapefruit in about 10^{-32} seconds. In virtue of the Heisenberg Uncertainty Principle, and because the comoving (x^1, x^2, x^3) dimensions have undergone inflation while the x^8 dimension has not, present epoch quantum fields that are functions of $(x^1, x^2, x^3, x^4, x^8)$ are expected to almost uniformly sample the region of the x^8 dimension that they occupy. The spatial x^8 average of functions of x^8 are expected to appear in effective four dimensional spacetime theories. The fact that the scale factors vanish on a set of measure zero may be handled in a straightforward manner by employing spatial x^8 averages in physical calculations.

Appendix A: Solution of the diagonal contribution to the mode equations (neglecting mode coupling)

For both the $\zeta = 3/10$ case and the second case with $\zeta = 0$, one may solve the diagonal mode equations obtained from Eq.[7.10] and Eq.[7.37] using Mathieu functions. However complex manipulations involving such mode functions is limited by the state of Mathieu function science. Instead, for both cases, to solve the uncoupled mode function equations we make use of the following identity, which is easily verified using the recursion relations for the Bessel functions. First, let us define several quantities.

Let $m_{\text{eff}}^2 > 0, B > 0, \nu, \alpha$ and β be given constants, and $n \in \mathbb{Z}$; let $J_n = J_n \left(3 k \sqrt{2B} \exp\left(-\frac{t}{6}\right) \right), \quad K_\mu = K_\mu \left(3 w \sqrt{2B} \exp\left(+\frac{t}{6}\right) \right)$ and $I_\mu = I_\mu \left(3 w \sqrt{2B} \exp\left(+\frac{t}{6}\right) \right)$. Then using the recursion relations for the Bessel functions [23] it is straightforward to show that

$$0 = \frac{\partial^{2}}{\partial t^{2}} \left[\left(\alpha K_{n+i\nu} + \beta I_{n+i\nu} \right) J_{n} \right] + \left[m_{\text{eff}}^{2} - B \frac{1}{2} \left(w^{2} e^{t/3} - k^{2} e^{-t/3} \right) \right] \left[\left(\alpha K_{n+i\nu} + \beta I_{n+i\nu} \right) J_{n} \right] - \left[m_{\text{eff}}^{2} + \frac{1}{9} \left(n + \frac{i\nu}{2} \right)^{2} \right] \left[\left(\alpha K_{n+i\nu} + \beta I_{n+i\nu} \right) J_{n} \right] + \frac{1}{2} k w B \left(-\alpha K_{n-1+i\nu} + \beta I_{n-1+i\nu} \right) J_{n-1} + \frac{1}{2} k w B \left(-\alpha K_{n+1+i\nu} - \beta I_{n+1+i\nu} \right) J_{n+1}.$$
(A.1)

Employing this identity to solve the uncoupled mode function equations has its roots in a closely related technique due to Dougall [25], Section[15], pages 191-193, with appropriate modifications that account for the asymmetric "potential" $\frac{1}{2} \left(w^2 e^{t/3} - k^2 e^{-t/3} \right)$ in this problem. Dougall gives the solution of Mathieu's modified differential equation as a series of products of Bessel functions.

The general solution to Eq.[7.40],

$$0 = \ddot{Y}_{m_8}(t) + \left[m_{\text{eff}}^2 - B\frac{1}{2}\left(w^2 e^{t/3} - k^2 e^{-t/3}\right)\right] Y_{m_8}(t),$$

may be obtained by introducing coefficient sets $\{c_n, d_n\}$, $n = -\infty, ..., \infty$ and then setting $\alpha = (-i)^n c_n$ and $\beta = (i)^n d_n$ in Eq.[A.1]. Next we sum over n from $n = -\infty, ..., \infty$ and

then re-label indices so that we may identify coefficients of $\sum [(\alpha K_{n+i\nu} + \beta I_{n+i\nu}) J_n]$ plus a remainder, which we require to vanish. This generates a three term recursion relation. We find that a general solution to Eq.[7.40] is given by

$$Y_{m_8}(t) = \sum_{n=-\infty}^{\infty} \left\{ i^n J_n \left(e^{-t/6} \, 3 \, k \, \sqrt{2 \, B} \right) \left[d_n I_{n+i\nu} \left(e^{t/6} \, 3 \, w \, \sqrt{2 \, B} \right) \right. + \left. (-1)^n c_n K_{n+i\nu} \left(e^{t/6} \, 3 \, w \, \sqrt{2 \, B} \right) \right] \right\}.$$
(A.2)

The coefficient sets $\{c_n, d_n\}, n = -\infty, \dots, \infty$ are solutions of the same recurrence relation

$$\frac{1}{2}i\,k\,w\,B\left(C_{n-1}+C_{n+1}\right) - \left[m_{\text{eff}}^2 + \frac{1}{9}\left(n + \frac{i\,\nu}{2}\right)^2\right]C_n = 0,\tag{A.3}$$

but with possibly distinct initial values, since a general solution to the three-term recurrence relation Eq.[A.3] possesses two arbitrary constants.

Eq.[A.2] may find application in calculating initial values in a numerical simulation, or in the computation of the approximate cross section for the creation of particle/anti-particle pairs of ψ particles through the annihilation of φ quanta, if one generalizes this model to include an interaction of ψ with the inflaton φ of the form $\lambda \psi^* \psi \varphi$.

- Planck Collaboration, P. A. R. Ade., and et al. Planck 2013 results. i. overview of products and scientific results. Astronomy and Astrophysics, submitted, 2013.
- [2] A. H. Guth. The inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D*, 23:347, 1981.
- [3] A. D. Linde. A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems. *Phys. Lett. B*, 108:389, 1982.
- [4] A. Albrecht and P. J. Steinhardt. Cosmology for grand unified theories with radiatively induced symmetry breaking. *Phys. Rev. Lett.*, 48:1220, 1982.
- [5] J. Dunkley [WMAP Collaboration]. Five-year wilkinson microwave anisotropy probe (wmap) observations: Likelihoods and parameters from the wmap data. Astrophys. J. Suppl., 180:306, 2009.
- [6] E. Komatsu [WMAP Collaboration]. Five-year wilkinson microwave anisotropy probe observations: Cosmological interpretation. Astrophys. J. Suppl., 180:330, 2009.
- [7] E. Komatsu and K. M. Smith. Seven-year wilkinson microwave anisotropy probe (wmap) observations: Cosmological interpretation. *The Astrophysical Journal Supplement Series*, 192(2):18, 2011.
- [8] A. Melchiorri W. H. Kinney, E. W. Kolb and A. Riotto. Latest inflation model constraints from cosmic microwave background measurements. *Phys. Rev. D*, 78:087302, 2008.
- G.F Hinshaw. Nine-year wilkinson microwave anisotropy probe (wmap) observations: Cosmology results. *submitted Astrophys. J. Suppl.*, 2013.
- [10] Max Tegmark, Michael A. Strauss, and et al. Cosmological parameters from sdss and wmap. Phys. Rev. D, 69(10):103501, May 2004.
- [11] Daniel J. Eisenstein, Idit Zehavi, David W. Hogg, and Roman Scoccimarro. Detection of the baryon acoustic peak in the large-scale correlation function of sdss luminous red galaxies. *The Astrophysical Journal*, 633(2):560, 2005.
- [12] Max Tegmark, Daniel J. Eisenstein, Michael A. Strauss, David H. Weinberg, and et al. Cosmological constraints from the sdss luminous red galaxies. *Phys. Rev. D*, 74(12):123507, Dec 2006.
- [13] P.A.R. Ade et al. Bicep2 i: Detection of b-mode polarization at degree angular scales. TBD,

2014.

- [14] Lev Kofman, Andrei Linde, and Alexei A. Starobinsky. Towards the theory of reheating after inflation. Phys. Rev. D, 56:3258–3295, Sep 1997.
- [15] A.D. Linde. Particle Physics and Inflationary Cosmology. Harwood Academic Publishers, Chur, Switzerland, 2005,1990.
- [16] Patrick L. Nash. Second gravity. Journal of Mathematical Physics, 51:042501-1 042501-27, 2010.
- [17] Quasi-normal modes and exponential energy decay for the kerr-de sitter black hole. Communications in Mathematical Physics, 306(1), 2011.
- [18] Kostas D. Kokkotas and Bernd Schmidt. Quasi-normal modes of stars and black holes. Living Reviews in Relativity, 2(2), 1999.
- [19] LI-HUI XUE, ZAI-XIONG SHEN, BIN WANG, and RU-KENG SU. Numerical simulation of quasi-normal modes in time-dependent background. *Modern Physics Letters A*, 19(03):239– 252, 2004.
- [20] K. A. Bronnikov. Gravitating brane systems: Some general theorems. Journal of Mathematical Physics, 40(2):924–938, 1999.
- [21] C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. San Francisco: W.H. Freeman and Co., 1973, 1973.
- [22] S. Weinberg. Cosmology. OUP Oxford, 2008.
- [23] G. N. Watson. A Treatise on the Theory of Bessel Functions. Cambridge University Press, Cambridge, England, 1944.
- [24] Charles Bla Balogh. Asymptotic expansions and general behavior of the modified bessel function of the third kind of imaginary order. Ph. D. Thesis, Oregon State University, 1965-04-08.
- [25] John Dougall. The solution of mathteu's differential equation. Proceedings of the Edinburgh Mathematical Society, 34(1):176–196, 1915.
- [26] David A. Harville. Matrix algebra from a statistician's perspective. Springer, New York, 1997.
- [27] James E. Lidsey, Andrew R. Liddle, Edward W. Kolb, Edmund J. Copeland, Tiago Barreiro, and Mark Abney. Reconstructing the inflaton potentialan overview. *Rev. Mod. Phys.*, 69:373– 410, Apr 1997.



FIG. 1: Difference of $\sqrt[6]{\csc^2(x)}$ and its realization in terms of a Fourier series that may be evaluated in closed form using hypergeometric functions. [Color online]



FIG. 2: Stable approximate modes FOR ODD n_8 ; pairs of plots (τ dependence, for fixed x and x dependence, for fixed τ). Total number of modes = 169 = 91 EVEN + 78 ODD n. $n = 6, m = 0, kw \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 2\sqrt{Y} = 2$. The rows and columns are delimited by distinct values of a_1 . $[a_1$ corresponds to odd n_8 ; a_2 corresponds to even n_8]. [Color online]



FIG. 3: Stable approximate modes FOR EVEN n_8 ; pairs of plots (τ dependence, for fixed x and x dependence, for fixed τ). Total number of modes = 169 = 91 EVEN + 78 ODD n. $n = 6, m = 0, kw \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 2\sqrt{Y} = 2$. The rows and columns are delimited by distinct values of a_2 . $[a_1$ corresponds to odd n_8 ; a_2 corresponds to even n_8]. [Color online]



FIG. 4: Stable approximate modes FOR ODD n_8 ; pairs of plots (τ dependence, for fixed x and x dependence, for fixed τ). Total number of modes = 225 = 105 EVEN + 120 ODD n. $n = 7, m = 0, kw \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 2\sqrt{Y} = 2$. The rows and columns are delimited by distinct values of a_1 . $[a_1$ corresponds to odd n_8 ; a_2 corresponds to even n_8]. [Color online]



FIG. 5: Stable approximate modes FOR EVEN n_8 ; pairs of plots (τ dependence, for fixed x and x dependence, for fixed τ). Total number of modes = 225 = 105 EVEN + 120 ODD n. $n = 7, m = 0, kw \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 2\sqrt{Y} = 2$. The rows and columns are delimited by distinct values of a_2 . [a_1 corresponds to odd n_8 ; a_2 corresponds to even n_8]. [Color online]



FIG. 6: Stable approximate modes FOR ODD n_8 ; pairs of plots (τ dependence, for fixed x and x dependence, for fixed τ). Total number of modes = 289 = 153 EVEN + 136 ODD n. $n = 8, m = 0, kw \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 2\sqrt{Y} = 2$. The rows and columns are delimited by distinct values of a_1 . $[a_1$ corresponds to odd n_8 ; a_2 corresponds to even n_8]. [Color online]



FIG. 7: Stable approximate modes FOR EVEN n_8 ; pairs of plots (τ dependence, for fixed x and x dependence, for fixed τ). Total number of modes = 289 = 153 EVEN + 136 ODD n. $n = 8, m = 0, kw \frac{\sqrt[3]{2}\Gamma(\frac{2}{3})}{\Gamma(\frac{5}{6})^2} = 2\sqrt{Y} = 2$. The rows and columns are delimited by distinct values of a_2 . [a_1 corresponds to odd n_8 ; a_2 corresponds to even n_8]. [Color online]