

# A NEW GENERALIZATION OF FERMAT'S LAST THEOREM

TIANXIN CAI AND DEYI CHEN

ABSTRACT. In this paper, we consider some hybrid Diophantine equations of addition and multiplication. We first improve a result on new Hilbert-Waring problem. Then we consider the equation

$$\begin{cases} A + B = C \\ ABC = D^n \end{cases} \quad (1)$$

where  $A, B, C, D, n \in \mathbb{Z}_+$  and  $n \geq 3$ , which may be regarded as a generalization of Fermat's equation  $x^n + y^n = z^n$ . When  $\gcd(A, B, C) = 1$ , (1) is equivalent to Fermat's equation, which means it has no positive integer solutions. We discuss several cases for  $\gcd(A, B, C) = p^k$  where  $p$  is an odd prime. In particular, for  $k = 1$  we prove that (1) has no nonzero integer solutions when  $n = 3$  and we conjecture that it is also true for any prime  $n > 3$ . Finally, we raise two conjectures on other hybrid Diophantine equations related with primes.

## 1. Introduction

In this paper, we consider some hybrid Diophantine equations of addition and multiplication. First of all,

$$n = x_1 + x_2 + \cdots + x_s$$

such that

$$x_1 x_2 \cdots x_s = x^k,$$

for  $n, x_i, x, k \in \mathbb{Z}_+$ , which is a new variant of Waring's problem:

$$n = x_1^k + x_2^k + \cdots + x_s^k.$$

We denote by  $g'(k)$  (resp.  $G'(k)$ ) the least positive integer such that every integer (resp. all sufficiently large integer) can be represented as a sum of at most  $g'(k)$  (resp.  $G'(k)$ ) positive integers, and the product of the  $g'(k)$  (resp.  $G'(k)$ ) integers is a  $k$ -th power. We show [1] that

$$g'(k) = 2k - 1; \quad G'(p) \leq p + 1; \quad G'(2p) \leq 2p + 2 \quad (p \geq 3); \quad G'(4p) \leq 4p + 2 \quad (p \geq 7);$$

where  $k$  is a positive integer and  $p$  is prime. In this paper, we improve the above results as follow.

**Theorem 1.** *For any composite number  $k$ ,  $G'(k) \leq k + 2$ .*

Next, we consider Fermat's Last Theorem. In 1637, Fermat claimed that the Diophantine equation

$$x^n + y^n = z^n$$

has no positive integer solutions for any integer  $n \geq 3$ . This was proved finally by Andrew Wiles in 1995 [13][11].

There are several generalizations of Fermat's Last Theorem, e.g., Fermat-Catalan conjecture, which states that the equation  $a^m + b^n = c^k$  has only finitely many

---

2010 Mathematics Subject Classification. Primary 11D41; Secondary 11D72.

Project supported by the National Natural Science Foundation of China 10871169.

solutions  $(a, b, c, m, n, k)$ , where  $a, b, c$  are positive coprime integers and  $m, n, k$  are positive integers, satisfying  $\frac{1}{m} + \frac{1}{n} + \frac{1}{k} < 1$ . So far there are only 10 solutions found [3][9]. Meanwhile, Beal's conjecture [7] states that the equation  $A^x + B^y = C^z$  has no solution in positive integers  $A, B, C, x, y$  and  $z$  with  $x, y$  and  $z$  at least 3 and  $A, B$  and  $C$  coprime. Beal has offered a prize of one million dollars for a proof of his conjecture or a counterexample [14]. Obviously, there are only finite solutions for Beal's equation under Fermat-Catalan conjecture. Meanwhile, it's known that both FLT and Fermat-Catalan conjecture are the consequences of the abc-conjecture, the latter was claimed to be proved in 2012 but not confirmed yet by Japanese mathematician Shinichi Mochizuki [8].

Now we expand the same idea to Fermat's equation as we did before to Hilbert-Waring problem. We consider a new Diophantine equation

$$\begin{cases} A + B = C \\ ABC = D^n \end{cases} \quad (1)$$

where  $A, B, C, D, n \in \mathbb{Z}_+$  and  $n \geq 3$ .

It is easy to see that if  $\gcd(A, B, C) = 1$  then  $A, B, C$  are pairwise coprime. Therefore,

$$\boxed{\text{Equation (1) has no positive integer solutions for } \gcd(A, B, C) = 1 \iff \text{FLT}} \quad (2)$$

In view of (2), we may ask some natural questions concerning (1) :

1. Is it possible to have a solution for all  $n \geq 3$ ? If so, is it possible to have infinitely many solutions ?
2. Is it possible to have a solution when  $\gcd(A, B, C) = p^k$ , where  $p$  is a prime,  $k \in \mathbb{Z}_+$  ?

In this paper, we answer the first question affirmatively by proving the following result.

**Theorem 2.** *For any  $n \not\equiv 0 \pmod{3}, n \geq 3$ , (1) has infinitely many positive integer solutions; for any  $n \equiv 0 \pmod{3}, n \geq 3$ , (1) has no positive integer solutions.*

For the second question, we discuss the special cases  $n = 4, 5$  and obtain the following

**Theorem 3.** *If  $\gcd(A, B, C) = p^k$  where  $k \in \mathbb{Z}_+$ ,  $p$  is odd prime and  $p \equiv 3 \pmod{8}$ , then the equation*

$$\begin{cases} A + B = C \\ ABC = D^4 \end{cases} \quad (3)$$

*has no positive integer solutions.*

For  $p = 2$  and some  $p \equiv 1, 5, 7 \pmod{8}$ , it is possible for (3) to have a positive integer solution when  $\gcd(A, B, C) = p$ . For example

$$\begin{cases} 2 + 2 = 4 & \begin{cases} 17 + 272 = 289 \\ 17 \times 272 \times 289 = 34^4, \end{cases} & \begin{cases} 5 + 400 = 405 \\ 5 \times 400 \times 405 = 30^4, \end{cases} \\ 2 \times 2 \times 4 = 2^4, & & \\ \begin{cases} 47927607119 + 1631432881 = 49559040000 \\ 47927607119 \times 1631432881 \times 49559040000 = 44367960^4, \end{cases} & & \end{cases}$$

where  $p = \gcd(A, B, C) = 2$  and  $p = \gcd(A, B, C) = 17, 5, 239$ ,  $p \equiv 1, 5, 7 \pmod{8}$  respectively.

**Theorem 4.** *If  $\gcd(A, B, C) = p^k$  where  $k \in \mathbb{Z}_+$ ,  $p$  is odd prime and  $p \not\equiv 1 \pmod{10}$ , then the equation*

$$\begin{cases} A + B = C \\ ABC = D^5 \end{cases} \quad (4)$$

*has no positive integer solutions.*

In general, we have the follows:

**Conjecture 1.** *If  $n \geq 3$  is prime,  $\gcd(A, B, C) = p^k$  where  $k \in \mathbb{Z}_+$ ,  $p$  is odd prime and  $p \not\equiv 1 \pmod{2n}$ , then (1) has no positive integer solutions.*

Finally, if  $n > 3$  is prime, we construct special prime  $p$  such that (1) has positive integer solutions for  $\gcd(A, B, C) = p^k$  as following

**Theorem 5.** *If  $n > 3$  is prime,  $n \equiv r \pmod{3}$ ,  $1 \leq r \leq 2$ ,  $a, b, m \in \mathbb{Z}$ ,  $m \neq 0$ , such that  $\frac{a^n + b^n}{a + b} = p$  is an odd prime and  $a + b = m^n$ , then  $p \equiv 1 \pmod{2n}$  and (1) has positive integer solutions for  $\gcd(A, B, C) = p^k$  where  $k \equiv \frac{rn-1}{3} \pmod{n}$ ,  $k \in \mathbb{Z}_+$ .*

Let  $a = 2, b = -1, m = 1$ , we obtain

**Corollary 1.** *If  $n > 3$  is prime,  $n \equiv r \pmod{3}$ ,  $1 \leq r \leq 2$  and  $p = 2^n - 1$  is Mersenne prime, then (1) has positive integer solutions for  $\gcd(A, B, C) = p^k$  where  $k \equiv \frac{rn-1}{3} \pmod{n}$ ,  $k \in \mathbb{Z}_+$ .*

Moreover, we have the following

**Conjecture 2.** *If  $n > 3$  is prime,  $n \equiv r \pmod{3}$ ,  $1 \leq r \leq 2$  and  $\gcd(A, B, C) = p^k$  where  $p$  is prime and  $k \not\equiv \frac{rn-1}{3} \pmod{n}$ , then (1) has no positive integer solutions.*

In particular, if  $n > 3$  is prime, then  $4 \not\equiv rn \pmod{n}$ , so  $1 \not\equiv \frac{rn-1}{3} \pmod{n}$ , it follows that we have a special case of Conjecture 2 when  $k = 1$ :

**Conjecture 3.** *If  $n$  is odd prime,  $\gcd(A, B, C) = p$  is prime, then (1) has no positive integer solutions.*

Finally, we consider another hybrid Diophantine equation of addition and multiplication. If we write  $n = a + b$ , then  $ab$  is not prime unless  $n = p + 1$  where  $p$  is prime, but the situation becomes interesting when we consider  $ab \pm 1$ . More precisely, we raise the following two conjectures ( we check for  $n$  up to  $10^7$ ), which seems deeper than Bertrand's postulate.

**Conjecture 4.** *For any positive integer  $n > 1$ , there exist  $a, b \in \mathbb{Z}_+$  such that*

$$\begin{cases} n = a + b \\ ab - 1 = p, \end{cases}$$

*where  $p$  is prime.*

**Conjecture 5.** *For any positive integer  $n > 1$ ,  $n \neq 6, 30, 54$ , there exist  $a, b \in \mathbb{Z}_+$  such that*

$$\begin{cases} n = a + b \\ ab + 1 = p, \end{cases}$$

*where  $p$  is prime.*

## 2. Preliminaries

**Lemma 1.** [2, Proposition 6.5.6] *Let  $c$  be a nonzero integer. The equation  $x^4 - y^4 = cz^2$  has a rational solution with  $xyz \neq 0$  if and only if  $|c|$  is a congruent number.*

**Lemma 2.** [12, Proposition 5] *Let  $p$  be a prime congruent to 3 modulo 8, then  $p$  is not a congruent number.*

**Lemma 3.** [4] *Let  $A > 2$  be a positive integer and has no prime divisors of the form  $10k + 1$ , then the equation*

$$\begin{cases} x^5 + y^5 = Az^5 \\ \gcd(x, y) = 1 \end{cases} \quad (5)$$

*has no nonzero integer solution. If  $A = 2$ , the solutions of (5) are  $(x, y, z) = \pm(1, 1, 1)$ .*

This lemma was first conjectured by V. A. Lebesgue [6] in 1843 and proved by E. Halberstadt and A. Kraus [4] in 2004.

**Lemma 4.** *For any prime  $p$ , integer  $n \geq 2$ , if  $\gcd(A, B, C) = p^k$  and  $k \equiv 0 \pmod{n}$ ,  $k \in \mathbb{Z}_+$ , then (1) has no nonzero integer solutions.*

*Proof.* Let  $A_1 = \frac{A}{p^k}$ ,  $B_1 = \frac{B}{p^k}$ ,  $C_1 = \frac{C}{p^k}$ , in view of  $\gcd(A, B, C) = p^k$  and (1), we obtain that  $A_1, B_1, C_1$  are pairwise coprime and (1) can be changed into

$$\begin{cases} A_1 + B_1 = C_1 \\ p^{3k} A_1 B_1 C_1 = D^n. \end{cases}$$

But  $k \equiv 0 \pmod{n}$ ,  $A_1, B_1, C_1$  are pairwise coprime, so  $A_1 = x^n$ ,  $B_1 = y^n$ ,  $C_1 = z^n$  and  $x^n + y^n = z^n$ . By Fermat's Last Theorem, we deduce that  $xyz = 0$ , so  $ABC = 0$ , a contradiction!  $\square$

## 3. Proofs of the Theorems

**Proof of Theorem 1.** For every positive integer  $n$ , let  $n = km + r$  where  $0 \leq r \leq k - 1$ .

If  $r = 0$ , when  $n > 2k^{2k}$ , we have  $m = \frac{n}{k} > 2k^{2k-1}$  and

$$n = km = \underbrace{(m - 2k^{2k-1}) + \dots + (m - 2k^{2k-1})}_k + k^{2k} + k^{2k}.$$

If  $0 < r \leq k - 1$ , when  $n > k^{2k-1} + k$ , we have  $m = \frac{n-r}{k} > \frac{k^{2k-1}}{k} > k^{k-1}r^{k-1}$  and

$$n = km + r = \underbrace{(m - k^{k-1}r^{k-1}) + \dots + (m - k^{k-1}r^{k-1})}_k + k^k r^{k-1} + r.$$

$\square$

**Proof of Theorem 2.** If  $n \not\equiv 0 \pmod{3}$ , then there exists  $k \in \mathbb{Z}_+$  such that  $3k + 2 \equiv 0 \pmod{n}$ . It is well known that there exist infinitely many  $(a, b, c) \in \mathbb{Z}_+^3$  such that  $a^2 + b^2 = c^2$ . Let

$$\begin{cases} A = a^{k+2} b^k c^k \\ B = a^k b^{k+2} c^k \\ C = a^k b^k c^{k+2}. \end{cases}$$

So we have infinitely many positive solutions  $(A, B, C)$  satisfying (1), where  $D = (abc)^{\frac{3k+2}{n}}$ .

If  $n \equiv 0 \pmod{3}$ , suppose  $(A, B, C)$  is a positive solution of (1). Let  $d = \gcd(A, B, C)$ . Then  $\frac{A}{d}, \frac{B}{d}$  and  $\frac{C}{d}$  are pairwise coprime and  $d^3 | D^n$ . But  $3|n$ , so we have  $d | D^{\frac{n}{3}}$  and

$$\begin{cases} \frac{A}{d} + \frac{B}{d} = \frac{C}{d} \\ \frac{A}{d} \cdot \frac{B}{d} \cdot \frac{C}{d} = \left(\frac{D^{\frac{n}{3}}}{d}\right)^3. \end{cases}$$

Since  $\frac{A}{d}, \frac{B}{d}$  and  $\frac{C}{d}$  are pairwise coprime, so  $\frac{A}{d} = x^3, \frac{B}{d} = y^3, \frac{C}{d} = z^3$  and  $x^3 + y^3 = z^3$ . By Fermat's Last Theorem, we deduce that  $xyz = 0$  and  $ABC = 0$ , a contradiction!  $\square$

**Proof of Theorem 3.** In view of Lemma 4, we only need to discuss  $k \not\equiv 0 \pmod{4}$ . Suppose odd prime  $p \equiv 3 \pmod{8}$  and (3) has a positive integer solution  $(A, B, C)$ . In view of  $\gcd(A, B, C) = p^k$ , (3) can be changed into

$$\begin{cases} A_1 + B_1 = C_1 \\ p^{3k} A_1 B_1 C_1 = D^4. \end{cases} \quad (6)$$

where  $A_1 = \frac{A}{p^k}, B_1 = \frac{B}{p^k}$  and  $C_1 = \frac{C}{p^k}$  are pairwise coprime. But  $k \not\equiv 0 \pmod{4}$ , we deduce that only one of  $A_1, B_1, C_1$  is congruent to 0 modulo  $p$ . Let  $3k \equiv r \pmod{4}$ , then  $1 \leq r \leq 3$ . In view of (6) and reorder  $A_1, B_1$  if necessary, we obtain

$$\begin{cases} A_1 = x^4 \\ B_1 = y^4, \\ C_1 = p^{4-r} z^4, \end{cases} \quad (7)$$

or

$$\begin{cases} A_1 = p^{4-r} z^4, \\ B_1 = y^4, \\ C_1 = x^4, \end{cases} \quad (8)$$

where  $x, y, pz$  are pairwise coprime.

If  $A_1, B_1$  and  $C_1$  satisfy (7), then

$$x^4 + y^4 = p^{4-r} z^4. \quad (9)$$

But  $\gcd(y, p) = 1$ , we have integer  $s \not\equiv 0 \pmod{p}$  such that  $sy \equiv 1 \pmod{p}$ . By (9) we deduce that

$$(xs)^4 \equiv -1 \pmod{p},$$

which implies that  $-1$  is a square modulo  $p$ , this can only hold for  $p = 2$  and  $p \equiv 1 \pmod{4}$ , in contradiction with  $p \equiv 3 \pmod{8}$ .

If  $A_1, B_1$  and  $C_1$  satisfy (8), then

$$x^4 - y^4 = p^{4-r} z^4. \quad (10)$$

When  $r = 2$ , we have  $x^4 - y^4 = (pz^2)^2$ , but it is well known that the equation  $X^4 - Y^4 = Z^2$  has no nonzero integer solutions. So  $r = 1$  or  $r = 3$  and (10) can be changed into

$$x^4 - y^4 = p(pz^2)^2 \quad (11)$$

or

$$x^4 - y^4 = p(z^2)^2 \quad (12)$$

respectively. By Lemma 1 and Lemma 2, we obtain that (11) and (12) have no positive integer solutions when  $p \equiv 3 \pmod{8}$ .  $\square$

**Proof of Theorem 4.** In view of Lemma 4, we only need to discuss  $k \not\equiv 0 \pmod{5}$ . In view of  $\gcd(A, B, C) = p^k$ , (4) can be changed into

$$\begin{cases} A_1 + B_1 = C_1 \\ p^{3k} A_1 B_1 C_1 = D^5. \end{cases} \quad (13)$$

where  $A_1 = \frac{A}{p^k}$ ,  $B_1 = \frac{B}{p^k}$  and  $C_1 = \frac{C}{p^k}$  are pairwise coprime. But  $k \not\equiv 0 \pmod{5}$ , we deduce that only one of  $A_1, B_1, C_1$  congruent to 0 modulo  $p$ . Let  $3k \equiv r \pmod{5}$ , then  $1 \leq r \leq 4$ . In view of (13) and reorder  $A_1, B_1, C_1$  if necessary, we obtain

$$\begin{cases} A_1 = x^5 \\ B_1 = y^5, \\ C_1 = p^{5-r} z^5, \end{cases} \quad (14)$$

where  $x, y$  and  $pz$  are pairwise coprime. From (13) and (14) we deduce that

$$x^5 + y^5 = p^{5-r} z^5. \quad (15)$$

But  $p$  is an odd prime and  $p \not\equiv 1 \pmod{10}$ , by Lemma 3, (15) has no positive integer solutions.  $\square$

**Proof of Theorem 5.** First of all, we prove that  $f(x, y) = \frac{x^k + y^k}{x + y} > 0$  for any odd positive integer  $k$ , where  $(x, y) \in \mathbb{R}^2$ ,  $x + y \neq 0$ . It is obvious that  $f(x, y) > 0$  when  $xy \geq 0$ . So we only need to prove the case  $xy < 0$ . If  $k = 1$ ,  $f(x, y) = \frac{x + y}{x + y} = 1 > 0$ . Suppose  $\frac{x^k + y^k}{x + y} > 0$  where  $x + y \neq 0$ . Because  $xy < 0$  and  $x + y \neq 0$ , we obtain

$$\frac{x^{k+2} + y^{k+2}}{x + y} = \frac{x^k + y^k}{x + y} (-xy) + x^{k+1} + y^{k+1} > 0.$$

By mathematical induction, we deduce that  $f(x, y) = \frac{x^k + y^k}{x + y} > 0$  for any odd positive integer  $k$ , where  $(x, y) \in \mathbb{R}^2$ ,  $x + y \neq 0$ . By condition of Theorem 5, we have a prime  $p$  such that

$$0 < p = \frac{a^n + b^n}{a + b} = a^{n-1} - a^{n-2}b + a^{n-3}b^2 - \dots + b^{n-1} = \sum_{j=0}^{n-1} a^{n-1-j} (-b)^j. \quad (16)$$

Since  $n \geq 3$ , we deduce from (16) that

$$\gcd(p, a) = \gcd(p, b) = \gcd(a, b) = 1. \quad (17)$$

So

$$\gcd(a, a + b) = \gcd(b, a + b) = 1. \quad (18)$$

Let

$$\begin{cases} A = p^{\frac{rn-1}{3} + tn} a^n \\ B = p^{\frac{rn-1}{3} + tn} b^n \\ C = p^{\frac{rn+2}{3} + tn} (a + b) \end{cases} \quad (19)$$

where  $r$  is defined in the condition of Theorem 5 and integer  $t \geq 0$ .

By (17), (18), (19) and  $a + b = m^n$ , we obtain that  $A, B$  and  $C$  satisfy (1) where  $D = p^{r+3t} abm$  and

$$\gcd(A, B, C) = p^k,$$

where  $k = \frac{rn-1}{3} + tn \equiv \frac{rn-1}{3} \pmod{n}$ .

Finally, we want to prove that  $p \equiv 1 \pmod{2n}$ . We only need to prove that  $p \equiv 1 \pmod{n}$  because both  $p$  and  $n$  are odd primes.

Since  $p = \frac{a^n+b^n}{a+b}$ ,  $n \geq 3$  is prime, by Fermat's little Theorem, we have

$$p(a+b) = a^n + b^n \equiv a + b \pmod{n}.$$

Now we only need to prove that  $a+b \not\equiv 0 \pmod{n}$ . Suppose  $a+b \equiv 0 \pmod{n}$ , then  $-b \equiv a \pmod{n}$  and

$$p = \frac{a^n + b^n}{a+b} = a^{n-1} - a^{n-2}b + a^{n-3}b^2 - \dots + b^{n-1} \equiv na^{n-1} \equiv 0 \pmod{n}.$$

But  $p$  and  $n$  are two primes, so we deduce that  $p = n$ .

If  $ab < 0$ , We may assume that  $a > 0, b < 0$ , then

$$p = \frac{a^n + b^n}{a+b} = a^{n-1} - a^{n-2}b + a^{n-3}b^2 - \dots + b^{n-1} > \underbrace{1+1+\dots+1}_n = n = p.$$

A contradiction!

If  $ab \geq 0$ , we may assume that  $a \geq 0, b \geq 0$ , but  $p = \frac{a^n+b^n}{a+b}$  is prime, so  $a \neq b$  and  $ab \neq 0$ . Reorder  $a, b$  (if necessary), we may assume that  $a \geq 1, b \geq 2$ .

If  $a = 1, b \geq 2$ , then  $p = \frac{1+b^n}{1+b} = \frac{1+b^p}{1+b}$ . Let  $f(b) = (b^p + 1) - p(b + 1)$ , then

$$f'(b) = pb^{p-1} - p = p(b^{p-1} - 1) > 0,$$

so

$$f(b) \geq f(2) = (2^p + 1) - 3p > 0$$

for odd prime  $p$ , from which we deduce that  $\frac{1+b^p}{1+b} > p$ , a contradiction!

If  $a \geq 2, b \geq 2$ , then  $p = \frac{a^n+b^n}{a+b} = \frac{a^p+b^p}{a+b} > \frac{pa+pb}{a+b} = p$ , once again a contradiction!  $\square$

Finally, we list some primes  $p$  satisfying the condition of Theorem 5 when  $n = 5, 7$ .

TABLE 1. Primes  $p < 10^7$  which satisfy the condition of Theorem 5 when  $n = 5$ .

$p$	$a$	$b$	$m$	1641301	35	-3	2
31	2	-1	1	1803001	25	-24	1
211	3	-2	1	2861461	28	-27	1
4651	6	-5	1	4329151	31	-30	1
61051	11	-10	1	4925281	32	-31	1
132661	11	21	2	5754901	45	-13	2
202981	9	23	2	7086451	35	-34	1
371281	17	-16	1	7944301	36	-35	1
723901	20	-19	1	8782981	49	-17	2

TABLE 2. Primes  $p < 10^{11}$  which satisfy the condition of Theorem 5 when  $n = 7$ .

$p$	$a$	$b$	$m$				
127	2	-1	1	1928294551	26	-25	1
14197	4	-3	1	8258704609	33	-32	1
543607	7	-6	1	14024867221	36	-35	1
1273609	8	-7	1	22815424087	39	-38	1
2685817	9	-8	1	30914273881	41	-40	1
5217031	10	-9	1	77617224511	59	69	2
16344637	12	-11	1	91154730577	49	-48	1
141903217	17	-16	1	98201826199	55	73	2

#### 4. Remarks on Conjectures 3-5

**Remark 1.** If  $n = 3$ , Conjecture 3 is true by Theorem 2. If abc-conjecture is true, then Conjecture 3 should be true for fixed prime  $p$  and sufficiently large  $n$  where  $n$  need not be prime.

*Proof.* Because  $\gcd(A, B, C) = p$ , then  $\gcd(\frac{A}{p}, \frac{B}{p}, \frac{C}{p}) = 1$ . By (1) we have

$$\begin{cases} \frac{A}{p} + \frac{B}{p} = \frac{C}{p} \\ ABC = D^n. \end{cases}$$

So  $\text{rad}\left(\frac{ABC}{p^3}\right) = \text{rad}\left(\frac{D}{p^3}\right) \leq \text{rad}(D)$  and  $C > D^{\frac{n}{3}}$ . For any  $n \geq 7$  and  $0 < \epsilon < \frac{1}{3}$ , we deduce

$$p \leq D = D^{\frac{7}{3}-1-\frac{1}{3}} \leq D^{\frac{7}{3}-1-\epsilon}$$

and

$$q\left(-\frac{A}{p}, -\frac{B}{p}, \frac{C}{p}\right) = \frac{\log\left(\frac{C}{p}\right)}{\log\left(\text{rad}\left(\frac{ABC}{p^3}\right)\right)} \geq \frac{\frac{n}{3} \log D - \left(\frac{7}{3} - 1 - \epsilon\right) \log D}{\log D} \geq 1 + \epsilon.$$

By abc-conjecture, there exist only finitely many triples  $\left(-\frac{A}{p}, -\frac{B}{p}, \frac{C}{p}\right)$ . Let  $A_1 = \frac{A}{p}, B_1 = \frac{B}{p}, C_1 = \frac{C}{p}$ , then  $\gcd(A_1, B_1, C_1) = 1, p^3 A_1 B_1 C_1 = D^n$ . Let  $M$  be the greatest  $m$  such that there is prime  $q$  satisfying  $q^m | A_1 B_1 C_1$ . Then, if  $n > M + 3$ , there is no solution for  $p^3 A_1 B_1 C_1 = D^n$ . Hence, when  $\gcd(A, B, C) = p$ , (1) has no positive integer solutions for sufficiently large  $n$ .  $\square$

However, we could not deduce Conjectures 1-3 from abc-conjecture.

**Remark 2.** Now we give some notes on Conjecture 4 and Conjecture 5. Obviously,

$$\begin{cases} n = a + b \\ ab \pm 1 = p \end{cases} \iff \begin{cases} a + b = n \\ ab = p \mp 1. \end{cases}$$

That is to say,  $a, b$  are the solutions of  $x^2 - nx + (p \mp 1) = 0$ ,

$$x = \frac{n \pm \sqrt{n^2 - 4(p \mp 1)}}{2} \in \mathbb{Z}_+. \quad (20)$$

Hence, there exists  $m \in \mathbb{Z}_+$  such that

$$n^2 - 4(p \mp 1) = m^2. \quad (21)$$

If  $n = 2k$ , then  $m = 2s$ , by (21) we have

$$k^2 \pm 1 = p + s^2. \quad (22)$$

If  $n = 2k + 1$ , then  $m = 2s + 1$ , by (21) we have

$$k^2 + k \pm 1 = p + s(s + 1). \quad (23)$$

Equation (22) is a corollary of Hardy-Littlewood conjecture [5]:

**Hardy-Littlewood Conjecture H.** *Every large integer is either a square or a sum of a prime and a square.*

For any odd prime  $p$ , if we define  $\frac{p-1}{2}$  as a half-prime, then the twin-prime conjecture could be expressed as: *There are infinitely many pair of consecutive half-primes* and the Goldbach's conjecture is expressed as: *Every positive integer  $n > 1$  is a sum of two half-primes.*

Equation (23) is a corollary of :

**Conjecture 6.** *Every positive integer  $n > 1$  is a sum of a half-prime and a triangle number.*

Conjecture 6 is equal to a conjecture of Sun's [10] : *Every odd integer  $n \geq 5$  can be written in the form  $p + x(x + 1)$  where  $p$  is a prime and  $x$  a positive integer.*

#### REFERENCES

1. T. X. Cai, D. Y. Chen, A new variant of the Hilbert-Waring Problem, Math. Comp. 82 (2013), 2333-2341.
2. H. Cohen, Number Theory, Volume I: Tools and diophantine equations, Graduate Texts in Mathematics, vol. 239, Springer-Verlag, New York, 2007.
3. H. Darmon and A. Granville, On the equations  $z^m = F(x, y)$  and  $Ax^p + By^q = cZ^r$ , Bull. London Math. Soc. 27(1995), 513-543.
4. E. Halberstadt and A. Kraus, Une conjecture de V. A. Lebesgue, J. London Math. Soc. (2) 69 (2004), no. 2, 291-302.
5. G. H. Hardy and J. E. Littlewood, Some problems of partitio numerorum III: on the expression of a large number as a sum of primes, Acta Math. 44 (1923), 1-70.
6. V. A. Lebesgue, Théorèmes nouveaux sur l'équation indéterminée  $x^5 + y^5 = az^5$ , J. Math. Pures Appl. 8 (1843) 49-70.
7. R. D. Mauldin, A Generalization of Fermat's Last Theorem: The Beal Conjecture and Prize Problem. Notices of the AMS 44 (11) (1997): 1436-1439.
8. S. Mochizuki, Inter-Universal Teichmüller Theory IV: Log-Volume Computations and Set-Theoretic Foundations.
9. C. Pomerance, Computational Number Theory, Leader, Imre, The Princeton Companion to Mathematics, Princeton University Press, pp. 361-362, June 2008.
10. Z. W. Sun, On sums of primes and triangular numbers, Journal of Combinatorics and Number Theory 1(2009), no. 1, 65-76.
11. R. Taylor and A. Wiles, Ring-theoretic properties of certain Hecke algebras. Annals of Mathematics, (1995), 553-572.
12. J. B. Tunnell, A classical Diophantine problem and modular forms of weight 3/2. Invent. Math., 72 (2) (1983), 323-334.
13. A. Wiles, Modular elliptic curves and Fermat's last theorem. Annals of Mathematics, (1995), 443-551.
14. The Beal Prize, AMS, <http://www.ams.org/profession/prizes-awards/ams-supported/beal-prize>

Department of Mathematics, Zhejiang University, Hangzhou, 310027, China  
**Email address:** [txcai@zju.edu.cn](mailto:txcai@zju.edu.cn)

Department of Mathematics, Zhejiang University, Hangzhou, 310027, China  
**Email address:** [chendeyi1986@126.com](mailto:chendeyi1986@126.com)