

CELL DECOMPOSITIONS OF DOUBLE BOTT-SAMELSON VARIETIES

VICTOR MOUQUIN

ABSTRACT. Let G be a connected complex semisimple Lie group. Webster and Yakimov have constructed partitions of the double flag variety $G/B \times G/B_-$, where (B, B_-) is a pair of opposite Borel subgroups of G , generalizing the Deodhar decompositions of G/B . We show that these partitions can be better understood by constructing cell decompositions of a product of two Bott-Samelson varieties $Z_{\mathbf{u}, \mathbf{v}}$, where $\mathbf{u}$ and $\mathbf{v}$ are sequences of simple reflections. We construct coordinates on each cell of the decompositions and in the case of a positive subexpression, we relate these coordinates to regular functions on a particular open subset of $Z_{\mathbf{u}, \mathbf{v}}$.

The geometry of the cell decompositions of $Z_{\mathbf{u}, \mathbf{v}}$ will be used in a forthcoming article about a Poisson structure on $Z_{\mathbf{u}, \mathbf{v}}$.

1. INTRODUCTION

Let G be a connected complex semisimple Lie group and B a Borel subgroup of G . In [5], V. Deodhar defined a family of decompositions of the flag variety G/B into pieces each of which is isomorphic to $\mathbb{C}^k \times (\mathbb{C}^*)^m$ for some integers $k, m \geq 0$. The Deodhar decomposition has been used to study the Kazdan-Lusztig polynomials associated to G and total positivity in G/B (see [5, 13]). Let B_- be a Borel subgroup of G opposite to B and consider the *double flag variety*

$$(G \times G)/(B \times B_-) \cong G/B \times G/B_-.$$

In [16], Webster and Yakimov generalised the decompositions of Deodhar to the double flag variety and showed that each piece of the decompositions is coisotropic with respect to a naturally defined Poisson structure on $G/B \times G/B_-$.

The Deodhar decomposition of G/B can be better understood through a natural cell decomposition on Bott-Samelson varieties (see [6, 9]). Motivated by the problem of better understanding the decompositions by Webster-Yakimov, we introduce *double Bott-Samelson varieties* $Z_{\mathbf{u}, \mathbf{v}}$, where $\mathbf{u} = (s_1, \dots, s_l)$ and $\mathbf{v} = (s_{l+1}, \dots, s_n)$ are any two sequences of simple reflections in the Weyl group W of G . We construct cell decompositions of $Z_{\mathbf{u}, \mathbf{v}}$ which give rise to the Deodhar-type decompositions of $G/B \times G/B_-$ by Webster-Yakimov when $\mathbf{u}$ and $\mathbf{v}$ are reduced. Coordinates are constructed on each piece of the decompositions, and these coordinates will be used in an upcoming article to study a naturally defined Poisson structure on $Z_{\mathbf{u}, \mathbf{v}}$.

1.1. Cell decompositions of $Z_{\mathbf{u}, \mathbf{v}}$ associated to shuffles. Throughout this article, if K is a Lie group, Q a closed subgroup of K , and $K_1, \dots, K_n$ submanifolds of K invariant under left and right multiplications by elements in Q , then

$$K_1 \times_Q \cdots \times_Q K_n / Q$$

denotes the quotient of $K_1 \times \cdots \times K_n$ by the right action of Q^n defined by

$$(1) \quad (k_1, \dots, k_n) \cdot (q_1, \dots, q_n) = (k_1 q_1, q_1^{-1} k_2 q_2, \dots, q_{n-1}^{-1} k_n q_n), \quad k_i \in K_i, q_j \in Q.$$

This article consists of the first part of the author's doctoral thesis.

For two sequences $\mathbf{u} = (s_1, \dots, s_l)$, $\mathbf{v} = (s_{l+1}, \dots, s_n)$ of simple reflections in the Weyl group W of G , let $Z_{\mathbf{u}}$ and $Z_{-\mathbf{v}}$ be the Bott-Samelson varieties

$$Z_{\mathbf{u}} = P_{s_1} \times_B \cdots \times_B P_{s_l}/B, \quad Z_{-\mathbf{v}} = P_{-s_{l+1}} \times_{B_-} \cdots \times_{B_-} P_{-s_n}/B_-,$$

where $P_s = B \cup BsB$ and $P_{-s} = B_- \cup B_-sB_-$ for a simple reflection s . Let

$$Z_{\mathbf{u}, \mathbf{v}} = Z_{\mathbf{u}} \times Z_{-\mathbf{v}}$$

and call $Z_{\mathbf{u}, \mathbf{v}}$ the *double Bott-Samelson variety* associated to $(\mathbf{u}, \mathbf{v})$. The multiplication in $G \times G$ induces a morphism (see (8))

$$\theta_{\mathbf{u}, \mathbf{v}} : Z_{\mathbf{u}, \mathbf{v}} \rightarrow G/B \times G/B_-.$$

Of particular interest is the open subset

$$(2) \quad \mathcal{O}^{\mathbf{u}, \mathbf{v}} = (Bs_1B \times_B \cdots \times_B Bs_lB/B) \times (B_{-s_{l+1}}B_- \times_{B_-} \cdots \times_{B_-} B_{-s_n}B_-/B_-)$$

of $Z_{\mathbf{u}, \mathbf{v}}$. When $\mathbf{u}, \mathbf{v}$ are reduced, $\theta_{\mathbf{u}, \mathbf{v}}$ induces an isomorphism between $\mathcal{O}^{\mathbf{u}, \mathbf{v}}$ and the product Schubert cell

$$(Bs_1 \cdots s_l B)/B \times (B_{-s_{l+1}} \cdots s_n B_-)/B_- \subset G/B \times G/B_-$$

and thus also a birational isomorphism between $Z_{\mathbf{u}, \mathbf{v}}$ and the product Schubert variety

$$\overline{(Bs_1 \cdots s_l B)/B} \times \overline{(B_{-s_{l+1}} \cdots s_n B_-)/B_-} \subset G/B \times G/B_-.$$

One of our aims is to demonstrate that double Bott-Samelson varieties not only provide resolutions of singularities of Schubert varieties, they also have interesting geometry of their own.

Recall that an (l, n) -shuffle is an element of the symmetric group S_n such that

$$\sigma(1) < \sigma(2) < \cdots < \sigma(l), \quad \sigma(l+1) < \sigma(l+2) < \cdots < \sigma(n).$$

Let σ be an (l, n) -shuffle. By a σ -*shuffled subexpression* of $(\mathbf{u}, \mathbf{v})$ we mean a sequence $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_n)$, where for each $1 \leq j \leq n$, $\gamma_j = s_{\sigma^{-1}(j)}$ or e , the identity element of W . Let $\Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$ be the set of all σ -shuffled subexpressions of $(\mathbf{u}, \mathbf{v})$. Our first construction, presented in Section 3, is a decomposition

$$(3) \quad Z_{\mathbf{u}, \mathbf{v}} = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}} C_{\sigma}^{\gamma}$$

associated to each (l, n) -shuffle σ , where for each $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$,

$$C_{\sigma}^{\gamma} \cong \mathbb{C}^{n-|J(\gamma)|}$$

for a subset $J(\gamma)$ of $\{1, 2, \dots, n\}$ associated to γ . More precisely, for each (l, n) -shuffle σ , we define an embedding of $Z_{\mathbf{u}, \mathbf{v}}$ into the quotient variety

$$(4) \quad DF_n = (G \times G) \times_{B \times B_-} \cdots \times_{B \times B_-} (G \times G)/(B \times B_-).$$

Let $Z_{\mathbf{u}, \mathbf{v}}^{\sigma}$ be the image of $Z_{\mathbf{u}, \mathbf{v}}$ in DF_n . Using certain combinatorial data associated to points in DF_n we first arrive at the set-theoretical decomposition

$$(5) \quad Z_{\mathbf{u}, \mathbf{v}}^{\sigma} = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}} C^{\gamma}.$$

We introduce an open affine covering $\{\mathcal{O}^{\gamma} \mid \gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}\}$ of $Z_{\mathbf{u}, \mathbf{v}}^{\sigma}$ and coordinates $\{z_1, z_2, \dots, z_n\}$ on each $\mathcal{O}^{\gamma}$, and we then show that the combinatorially defined set C^{γ} coincides with the subset

of $\mathcal{O}^\gamma$ defined by $z_j = 0$ for $j \in J(\gamma)$.

The image of $\mathcal{O}^{\mathbf{u}, \mathbf{v}} \subset Z_{\mathbf{u}, \mathbf{v}}$ under the embedding of $Z_{\mathbf{u}, \mathbf{v}}$ into DF_n is $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$, where $\sigma(\mathbf{u}, \mathbf{v}) = (s_{\sigma^{-1}(1)}, s_{\sigma^{-1}(2)}, \dots, s_{\sigma^{-1}(n)})$. The decomposition in (5) then gives rise to the decomposition

$$(6) \quad \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}^d} C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})},$$

where $\Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}^d$ denotes the set of *distinguished σ -shuffled subexpressions* of $(\mathbf{u}, \mathbf{v})$ (see Definition 3.21). We show that for each $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}^d$,

$$C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} \cong \mathbb{C}^k \times (\mathbb{C}^*)^m$$

for some integers $k, m \geq 0$ (see Proposition 3.25). When $\mathbf{u}$ and $\mathbf{v}$ are reduced, we show in Section 3.8 that the decomposition in (6) corresponds to that of

$$(Bs_1 \cdots s_l B)/B \times (B_{-s_{l+1}} \cdots s_n B_{-})/B_{-} \subset G/B \times G/B_{-}$$

by Webster and Yakimov under the isomorphisms

$$\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} \cong \mathcal{O}^{\mathbf{u}, \mathbf{v}} \cong (Bs_1 \cdots s_l B)/B \times (B_{-s_{l+1}} \cdots s_n B_{-})/B_{-}.$$

As in [16] for the case when $\mathbf{u}$ and $\mathbf{v}$ are reduced, for each $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}^d$, we describe in Section 4 the subset $C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ of $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ using a set of regular functions $\{\psi_{\gamma, j}\}_{j=1, \dots, n}$ on $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ which are defined in terms of generalized minors [8]. When γ is *positive* (see Definition 3.11), we show that the coordinate functions $\{z_j \mid j \notin J(\gamma)\}$ on C^γ and the regular functions $\psi_{\gamma, j}$ are related by monomial transformations.

1.2. General notation. If an upper case letter denotes a Lie group, its Lie algebra will be denoted by the corresponding lower case gothic letter.

The identity element in any group will be denoted by e .

For integers $k \leq n$, we denote by $[k, n]$ the set of integers j such that $k \leq j \leq n$.

If a set X has a right action by a group L , and $p : X \rightarrow Y = X/L$ is the projection, we write $[x]_Y \in Y$ for $x \in X$.

Acknowledgments. The author would like to thank gratefully Jiang-Hua Lu for her help and encouragements. This work was completed while the author was supported by a University of Hong Kong Post-graduate Studentship and a grant from the department of Mathematics of the University of Hong Kong.

2. LIE THEORY BACKGROUND

Here we set up the notation from Lie theory, recall the definition of Bott-Samelson varieties and basic facts about the double flag variety of a complex semisimple Lie group.

2.1. Notation. Recall that G is a connected complex semisimple Lie group, and that we have fixed a pair (B, B_-) of opposite Borel subgroups. Let $H = B \cap B_-$ be the maximal torus defined by B and B_- . Let N and N_- be respectively the unipotent radicals of B and B_- . Let Δ be the root system defined by H , and let Δ_+ and Γ be respectively the sets of positive and simple roots defined by B . Let $\mathfrak{g} = \mathfrak{h} \oplus \sum_{\alpha \in \Delta} \mathfrak{g}_\alpha$ be the root space decomposition of $\mathfrak{g}$. Let $\langle, \rangle_{\mathfrak{g}}$ be a fixed nonzero multiple of the Killing form on $\mathfrak{g}$. Recall that the restriction of $\langle, \rangle_{\mathfrak{g}}$ to $\mathfrak{h}$ is nondegenerate and defines a nondegenerate symmetric bilinear form on $\mathfrak{h}^*$, which will be denoted by $\langle, \rangle$. For $\alpha \in \Delta_+$, let $h_\alpha \in \mathfrak{h}$ be the unique element in $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$ such that $\alpha(h_\alpha) = 2$. Fix root vectors $e_\alpha \in \mathfrak{g}_\alpha$ and $e_{-\alpha} \in \mathfrak{g}_{-\alpha}$ such that $[e_\alpha, e_{-\alpha}] = h_\alpha$. Let $\phi_\alpha : \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{g}$ be the Lie algebra homomorphism defined by

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \mapsto e_\alpha, \text{ and } \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \mapsto e_{-\alpha}.$$

The corresponding Lie group homomorphism from $SL(2, \mathbb{C})$ to G will also be denoted by ϕ_α . For $z \in \mathbb{C}$, let

$$x_\alpha(z) = \phi_\alpha \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix}, \quad x_{-\alpha}(z) = \phi_\alpha \begin{pmatrix} 1 & 0 \\ z & 1 \end{pmatrix}, \text{ and} \\ \bar{s}_\alpha = \phi_\alpha \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Let $N_G(H)$ be the normaliser subgroup of H in G , and let $W = N_G(H)/H$ be the Weyl group of G . Let $S = \{s_\alpha = \bar{s}_\alpha H \mid \alpha \in \Gamma\}$. It is well known that the pair (W, S) forms a Coxeter system. In particular, W is generated by the simple reflections s_α , $\alpha \in \Gamma$. We denote the action of W on H by conjugation as a right action by $h^w = \dot{w}^{-1}h\dot{w}$, where $h \in H$, $w \in W$ and $\dot{w}$ is any representative of w .

Let $X^*(H) = \text{Hom}(H, \mathbb{C}^*)$ and $X_*(H) = \text{Hom}(\mathbb{C}^*, H)$ be respectively the lattices of characters and co-characters of H . We write the action of $\lambda \in X^*(H)$ on H by $h^\lambda \in \mathbb{C}^*$, $h \in H$. There is a natural embedding of $X^*(H)$ in $\mathfrak{h}^*$ and $X_*(H)$ in $\mathfrak{h}$, and the natural pairing $(,) : \mathfrak{h}^* \times \mathfrak{h} \rightarrow \mathbb{C}$ restricts to a perfect pairing of free abelian groups between $X^*(H)$ and $X_*(H)$ which we also denote by $(,)$. Recall that for $\alpha \in \Delta_+$ the coroot $\check{\alpha} \in X_*(H)$ is given by

$$\check{\alpha}(z) = \phi_\alpha \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix}, \quad z \in \mathbb{C}^*.$$

An element $x \in N_- H N$ can be uniquely written $x = [x]_- [x]_0 [x]_+$, with $[x]_- \in N_-$, $[x]_0 \in H$, and $[x]_+ \in N$.

When G is simply connected, denote by $\lambda_\alpha \in X^*(H)$ the fundamental weight associated to $\alpha \in \Gamma$. Recall that λ_α is defined by $(\lambda_\alpha, \check{\beta}) = \delta_{\alpha, \beta}$, $\beta \in \Gamma$.

The Weyl group W of G is naturally endowed with a partial ordering $\leq$, called the *Bruhat order*, defined by

$$v \leq w, \text{ if } BvB \subset \overline{BwB}, \quad v, w \in W,$$

where $\overline{BuB}$ is the Zariski closure of BuB in G , $u \in W$. For $w \in W$, let $l(w)$ be the smallest integer such that w can be written as a product of simple reflections. The function $l : W \rightarrow \mathbb{N}$

is called the *length function* of W . If $w = s_1 \cdots s_{l(w)}$, with $s_j \in S$, one calls the finite sequence $(s_1, \dots, s_{l(w)})$ a *reduced expression* for w . Let $R(w)$ be the set of all reduced expressions for w . If $(s_1, \dots, s_k) \in R(w)$, define $\bar{w} = \bar{s}_1 \cdots \bar{s}_k \in N_G(H)$. It is well known (see [15, Proposition 9.3.2]) that $\bar{w}$ does not depend on the choice of the reduced expression. Let $v, w \in W$. If $(s_1, \dots, s_k)$ is a reduced expression for w , then $v \leq w$ if and only if $v = s_{i_1} \cdots s_{i_l}$, where $(i_1 < \cdots < i_l)$ is a subsequence of $(1, \dots, k)$.

Recall that a *monoid* is a pair $(M, *)$, where M is a set, and $*$: $M \times M \rightarrow M$ a binary operation which is associative, and admits an identity element $e \in M$. The Weyl group has a natural monoid product $*$, given by

$$s_\alpha * w = \max\{s_\alpha w, w\} = \begin{cases} s_\alpha w, & \text{if } w < s_\alpha w \\ w, & \text{if } s_\alpha w < w, \end{cases} \quad \alpha \in \Gamma, w \in W,$$

and

$$u * w = s_1 * (s_2 * (\cdots s_l * w)), \quad u, w \in W,$$

where $(s_1, \dots, s_l)$ is any reduced expression for u .

In [10] two operations on W are introduced. For $w \in W$ and a simple root s_α , define

$$s_\alpha \triangleright w = \min\{s_\alpha w, w\} = \begin{cases} s_\alpha w, & \text{if } s_\alpha w < w \\ w, & \text{if } w < s_\alpha w, \end{cases}$$

and

$$w \triangleleft s_\alpha = \min\{ws_\alpha, w\} = \begin{cases} ws_\alpha, & \text{if } ws_\alpha < w \\ w, & \text{if } w < ws_\alpha. \end{cases}$$

For $u \in W$ define

$$\begin{aligned} u \triangleright w &= s_1 \triangleright (s_2 \triangleright (\cdots s_n \triangleright w)), \\ w \triangleleft u &= ((w \triangleleft s_1) \triangleleft s_2) \cdots \triangleleft s_n, \end{aligned}$$

where $(s_1, \dots, s_n)$ is a reduced expression for u . These two definitions are independent of the choice of the reduced expression for u . The operation $\triangleleft$ (resp. $\triangleright$) is a right (resp. left) monoidal action of $(W, *)$ on W . See [10] for more details.

2.2. Bott-Samelson varieties. Bott-Samelson varieties have their origins in the papers [3] of Bott-Samelson and [4] of Demazure. For more details, see [1, 2]. For any integer $n \geq 1$, let

$$(7) \quad F_n = G \times_B \cdots \times_B G/B, \quad F_{-n} = G \times_{B_-} \cdots \times_{B_-} G/B_-.$$

Recall that if s is a simple reflection, then $P_s = B \cup BsB$, and $P_{-s} = B_- \cup B_-sB_-$. Let $\mathbf{u} = (s_1, \dots, s_l)$ be a sequence of simple reflections. Then one has

$$\begin{aligned} Z_{\mathbf{u}} &= P_{s_1} \times_B \cdots \times_B P_{s_l}/B \subset F_l, \\ Z_{-\mathbf{u}} &= P_{-s_1} \times_{B_-} \cdots \times_{B_-} P_{-s_l}/B_- \subset F_{-l}. \end{aligned}$$

Both $Z_{\mathbf{u}}$ and $Z_{-\mathbf{u}}$ are smooth projective varieties of complex dimension l . Let $\theta_{\mathbf{u}} : Z_{\mathbf{u}} \rightarrow G/B$ and $\theta_{-\mathbf{u}} : Z_{-\mathbf{u}} \rightarrow G/B_-$ be the multiplication maps

$$\begin{aligned} \theta_{\mathbf{u}}([p_1, \dots, p_l]_{Z_{\mathbf{u}}}) &= p_1 \cdots p_l \cdot B, \\ \theta_{-\mathbf{u}}([p_{-1}, \dots, p_{-l}]_{Z_{-\mathbf{u}}}) &= p_{-1} \cdots p_{-l} \cdot B_-, \end{aligned}$$

where $p_i \in P_{s_i}$ and $p_{-i} \in P_{-s_i}$. The image of $\theta_{\mathbf{u}}$ is the Schubert variety $\overline{BuB/B}$, where $u = s_1 * \cdots * s_l$. If $(s_1, \dots, s_l) \in R(u)$, then it is well known that

$$\theta_{\mathbf{u}} : Z_{\mathbf{u}} \rightarrow \overline{BuB/B}$$

is a proper, surjective, birational isomorphism.

2.3. The double flag variety and double Bott-Samelson varieties. The double flag variety

$$DF_1 = (G \times G)/(B \times B_-) \cong G/B \times G/B_-$$

has the natural left $G \times G$ action given by

$$(g_1, g_2) \cdot (h_1.B, h_2.B_-) = (g_1 h_1.B, g_2 h_2.B_-), \quad g_i, h_i \in G.$$

Denote by G_{diag} the diagonal subgroup of $G \times G$, and for any $w \in W$, let $G_{\text{diag}}(w)$ be the G_{diag} -orbit passing through the point (wB, B_-) . It is well known that the map $w \mapsto G_{\text{diag}}(w)$ gives a parametrization of the G_{diag} -orbits in DF_1 by W . Identifying G_{diag} with G , the stabilizer subgroup of (wB, B_-) is $B_- \cap wBw^{-1}$. Thus

$$\dim_{\mathbb{C}}(G_{\text{diag}}(w)) = |\Delta| - l(w).$$

For $h_1, h_2 \in G$, $(h_1.B, h_2.B_-)$ lies in $G_{\text{diag}}(w)$ if and only if $h_2^{-1}h_1 \in B_-wB$.

For $u, v \in W$, let $\mathcal{O}^{u,v}$ be the orbit of $B \times B_-$ passing through the point (uB, vB_-) . The map $(u, v) \mapsto \mathcal{O}^{u,v}$ is a parametrization of the $B \times B_-$ -orbits in DF_1 by $W \times W$. One has

$$\dim_{\mathbb{C}}(\mathcal{O}^{u,v}) = l(u) + l(v).$$

For $u, v, w \in W$, define $\mathcal{O}_w^{u,v} = \mathcal{O}^{u,v} \cap G_{\text{diag}}(w)$.

Proposition 2.1. [16] *Let $u, v, w \in W$. Then $\mathcal{O}_w^{u,v}$ is nonempty if and only if $w \leq v^{-1} * u$.*

If $\mathbf{u} = (s_1, \dots, s_l)$ and $\mathbf{v} = (s_{l+1}, \dots, s_n)$ are two sequences of simple reflections, recall that we have defined the double Bott Samelson variety associated to $(\mathbf{u}, \mathbf{v})$ as

$$Z_{\mathbf{u}, \mathbf{v}} = Z_{\mathbf{u}} \times Z_{-\mathbf{v}}.$$

Let $\theta_{\mathbf{u}, \mathbf{v}} : Z_{\mathbf{u}, \mathbf{v}} \rightarrow DF_1$ be the map defined by

$$(8) \quad \theta_{\mathbf{u}, \mathbf{v}}([p_1, \dots, p_l]_{Z_{\mathbf{u}}}, [p_{-(l+1)}, \dots, p_{-n}]_{Z_{-\mathbf{v}}}) = (p_1 \cdots p_l.B, p_{-(l+1)} \cdots p_{-n}.B_-)$$

where $p_i \in P_{s_i}$, and $p_{-j} \in P_{-s_j}$. The image of $\theta_{\mathbf{u}, \mathbf{v}}$ is the product Schubert cell

$$\overline{Bs_1 * \cdots * s_l B/B} \times \overline{B_{-s_{l+1}} * \cdots * s_n B_-/B_-}.$$

If $\mathbf{u}$ and $\mathbf{v}$ are reduced, then $\theta_{\mathbf{u}, \mathbf{v}}$ restricts to an isomorphism

$$\theta_{\mathbf{u}, \mathbf{v}}|_{\mathcal{O}^{\mathbf{u}, \mathbf{v}}} : \mathcal{O}^{\mathbf{u}, \mathbf{v}} \rightarrow \mathcal{O}^{u, v},$$

where $u = s_1 \cdots s_l$, $v = s_{l+1} \cdots s_n$, and $\mathcal{O}^{\mathbf{u}, \mathbf{v}}$ has been defined in (2).

3. AFFINE CHARTS AND CELL DECOMPOSITIONS OF $Z_{\mathbf{u}, \mathbf{v}}$ ASSOCIATED TO SHUFFLES

If $\mathbf{u} = (s_1, \dots, s_l)$, $\mathbf{v} = (s_{l+1}, \dots, s_n)$ are two sequences of simple reflections, we construct for any (l, n) -shuffle an embedding of $Z_{\mathbf{u}, \mathbf{v}}$ into DF_n . Using some combinatorial data associated to points in DF_n , we give a set-theoretic decomposition

$$Z_{\mathbf{u}, \mathbf{v}}^\sigma = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}} C^\gamma,$$

of the image $Z_{\mathbf{u}, \mathbf{v}}^\sigma$ of $Z_{\mathbf{u}, \mathbf{v}}$ in DF_n . We then cover $Z_{\mathbf{u}, \mathbf{v}}^\sigma$ with 2^n open subsets each of which is isomorphic to $\mathbb{C}^n$, and use this covering to prove that each C^γ is isomorphic to $\mathbb{C}^k$, for some $k \in [1, n]$.

In Section 3.8, we discuss the relation between our decompositions of $Z_{\mathbf{u}, \mathbf{v}}$ and the Deodhar-type decompositions of $G/B \times G/B_-$ constructed in [16].

3.1. Double flag varieties DF_n . Let $n \geq 1$ and recall

$$DF_n = (G \times G) \times_{B \times B_-} \cdots \times_{B \times B_-} (G \times G) / (B \times B_-)$$

from (4). Then DF_n is naturally isomorphic to $F_n \times F_{-n}$ via the isomorphism

$$(9) \quad \varphi_n : DF_n \rightarrow F_n \times F_{-n}, [(g_1, h_1), \dots, (g_n, h_n)]_{DF_n} \mapsto ([g_1, \dots, g_n]_{F_n}, [h_1, \dots, h_n]_{F_{-n}}).$$

Moreover, DF_n is naturally endowed with a map to DF_1 ,

$$(10) \quad \theta_n : DF_n \rightarrow DF_1, [(g_1, h_1), \dots, (g_n, h_n)]_{DF_n} \mapsto (g_1 \cdots g_n \cdot B, h_1 \cdots h_n \cdot B_-),$$

and for $j \in [1, n-1]$, one has the projection

$$(11) \quad \rho_{j,n} : DF_n \rightarrow DF_j, [(g_1, h_1), \dots, (g_n, h_n)]_{DF_n} \mapsto [(g_1, h_1), \dots, (g_j, h_j)]_{DF_j}.$$

Let $x \in DF_n$. Let $w_n(x)$ be the unique Weyl group element such that $\theta_n(x) \in G_{\text{diag}}(w_n)$, and for $j \in [1, n-1]$, denote by $w_j(x) \in W$ the unique element such that $\theta_j(\rho_{j,n}(x)) \in G_{\text{diag}}(w_j)$. One thus obtains a map

$$(12) \quad \Phi_n : DF_n \rightarrow W^n$$

by sending x to $\Phi_n(x) = (w_1(x), \dots, w_n(x))$.

3.2. Embeddings of $Z_{\mathbf{u}, \mathbf{v}}$ into DF_n using shuffles.

Definition 3.1. Let $n \geq 1$ and $n \geq l \geq 0$ be integers and denote by $S_{l,n}$ the set of all (l, n) -shuffles. To any $\sigma \in S_{l,n}$ we associate a sequence

$$\epsilon(\sigma) = (\epsilon(\sigma)_1, \dots, \epsilon(\sigma)_n)$$

of elements in $\{1, -1\}$ by setting

$$\epsilon(\sigma)_{\sigma(j)} = \begin{cases} -1, & \text{if } j \in [1, l] \\ 1, & \text{if } j \in [l+1, n]. \end{cases}$$

For notational simplicity, once σ is fixed, we will write $\epsilon = (\epsilon_1, \dots, \epsilon_n)$ instead of $\epsilon(\sigma) = (\epsilon(\sigma)_1, \dots, \epsilon(\sigma)_n)$. Note that there is a one to one correspondance between $S_{l,n}$ and the set of sequences $\epsilon \in \{1, -1\}^n$, with -1 appearing exactly l times.

Let $\mathbf{u} = (s_1, \dots, s_l)$ and $\mathbf{v} = (s_{l+1}, \dots, s_n)$ be two sequences of simple reflections and let $\sigma \in S_{l,n}$. Define the sequence

$$(13) \quad \sigma(\mathbf{u}, \mathbf{v}) = (\delta_1, \dots, \delta_n)$$

by letting $\delta_{\sigma(j)} = s_j$, $j \in [1, n]$.

Let σ be an (l, n) -shuffle. Define maps $\mu_\sigma^- : G^n \rightarrow G^l$ and $\mu_\sigma^+ : G^n \rightarrow G^{n-l}$ by

$$(14) \quad \begin{aligned} \mu_\sigma^-(g_1, \dots, g_n) &= (g_1 \cdots g_{\sigma(1)}, g_{\sigma(1)+1} \cdots g_{\sigma(2)}, \dots, g_{\sigma(l-1)+1} \cdots g_{\sigma(l)}) \\ \mu_\sigma^+(g_1, \dots, g_n) &= (g_1 \cdots g_{\sigma(l+1)}, g_{\sigma(l+1)+1} \cdots g_{\sigma(l+2)}, \dots, g_{\sigma(n-1)+1} \cdots g_{\sigma(n)}). \end{aligned}$$

Both μ_σ^- and μ_σ^+ descend to maps $[\mu_\sigma^-] : F_n \rightarrow F_l$ and $[\mu_\sigma^+] : F_{-n} \rightarrow F_{-(n-l)}$ given by

$$(15) \quad \begin{aligned} [\mu_\sigma^-]([g_1, \dots, g_n]_{F_n}) &= [g_1 \cdots g_{\sigma(1)}, g_{\sigma(1)+1} \cdots g_{\sigma(2)}, \dots, g_{\sigma(l-1)+1} \cdots g_{\sigma(l)}]_{F_l} \\ [\mu_\sigma^+]([g_1, \dots, g_n]_{F_{-n}}) &= [g_1 \cdots g_{\sigma(l+1)}, g_{\sigma(l+1)+1} \cdots g_{\sigma(l+2)}, \dots, g_{\sigma(n-1)+1} \cdots g_{\sigma(n)}]_{F_{-(n-l)}}. \end{aligned}$$

For $j \in [1, n]$, define

$$(16) \quad G_j^\sigma = \begin{cases} G \times B_-, & \text{if } \epsilon_j = -1 \\ B \times G, & \text{if } \epsilon_j = 1, \end{cases}$$

and

$$(17) \quad F_{l,n}^\sigma = G_1^\sigma \times_{B \times B_-} \cdots \times_{B \times B_-} G_n^\sigma / (B \times B_-) \subset DF_n$$

For each $j \in [1, n]$, let $G_{j,L}^\sigma$ and $G_{j,R}^\sigma$ be respectively the first and second component of G_j^σ . Then the diffeomorphism φ_n , recall (9), restricts to a diffeomorphism

$$\varphi_n|_{F_{l,n}^\sigma} : F_{l,n}^\sigma \rightarrow F_L^\sigma \times F_R^\sigma,$$

where

$$F_L^\sigma = G_{1,L}^\sigma \times_B \cdots \times_B G_{n,L}^\sigma / B \subset F_n, \quad F_R^\sigma = G_{1,R}^\sigma \times_{B_-} \cdots \times_{B_-} G_{n,R}^\sigma / B_- \subset F_{-n}.$$

By composing with $[\mu_\sigma^-] \times [\mu_\sigma^+]$, one obtains the diffeomorphism

$$(18) \quad \varphi^\sigma = ([\mu_\sigma^-] \times [\mu_\sigma^+]) \circ \left(\varphi_n|_{F_{l,n}^\sigma} \right) : F_{l,n}^\sigma \rightarrow F_l \times F_{-(n-l)}.$$

Explicitly, φ^σ is given by

$$\begin{aligned} \varphi^\sigma([(p_1, q_1), \dots, (p_n, q_n)]_{DF_n}) &= \left([(p_1 \cdots p_{\sigma(1)}, p_{\sigma(1)+1} \cdots p_{\sigma(2)}, \dots, p_{\sigma(l-1)+1} \cdots p_{\sigma(l)}]_{F_l} \right. \\ &\quad \left. [q_1 \cdots q_{\sigma(l+1)}, q_{\sigma(l+1)+1} \cdots q_{\sigma(l+2)}, \dots, q_{\sigma(n-1)+1} \cdots q_{\sigma(n)}]_{F_{-(n-l)}} \right), \end{aligned}$$

where $(p_j, q_j) \in G_j^\sigma$.

Now, let $\mathbf{u} = (s_1, \dots, s_l)$ and $\mathbf{v} = (s_{l+1}, \dots, s_n)$ be two sequences of simple reflections and let $\sigma(\mathbf{u}, \mathbf{v}) = (\delta_1, \dots, \delta_n)$. For $j \in [1, n]$ define

$$(19) \quad P_j^\sigma = \begin{cases} P_{\delta_j} \times B_-, & \text{if } \epsilon_j = -1 \\ B \times P_{-\delta_j}, & \text{if } \epsilon_j = 1, \end{cases}$$

and let

$$(20) \quad Z_{\mathbf{u}, \mathbf{v}}^\sigma = P_1^\sigma \times_{B \times B_-} \cdots \times_{B \times B_-} P_n^\sigma / (B \times B_-) \subset F_{l,n}^\sigma.$$

Then φ^σ restricts to a diffeomorphism $\varphi^\sigma|_{Z_{\mathbf{u},\mathbf{v}}^\sigma}: Z_{\mathbf{u},\mathbf{v}}^\sigma \rightarrow Z_{\mathbf{u},\mathbf{v}}$. One thus has an embedding

$$(21) \quad I_{\mathbf{u},\mathbf{v}}^\sigma = (\varphi^\sigma|_{Z_{\mathbf{u},\mathbf{v}}^\sigma})^{-1} : Z_{\mathbf{u},\mathbf{v}} \hookrightarrow DF_n.$$

3.3. Shuffled subexpressions and the subvarieties C^γ . Let $\mathbf{u}, \mathbf{v}$ and σ be as in Definition 3.1. Recall the map $\Phi_n : DF_n \rightarrow W^n$ from (12), and the sequence $\sigma(\mathbf{u}, \mathbf{v})$ from (13).

Lemma 3.2. *Let $\mathbf{w} = (w_1, \dots, w_n) \in W^n$. If $\mathbf{w} \in \Phi_n(Z_{\mathbf{u},\mathbf{v}}^\sigma)$, then $w_1 \in \{e, \delta_1\}$, and for $j \in [2, n]$,*

$$w_j \in \begin{cases} \{w_{j-1}, w_{j-1}\delta_j\}, & \text{if } \epsilon_j = -1 \\ \{w_{j-1}, \delta_j w_{j-1}\}, & \text{if } \epsilon_j = 1. \end{cases}$$

Proof. Let $x \in Z_{\mathbf{u},\mathbf{v}}^\sigma$, and $\mathbf{w} = \Phi_n(x)$. For $j \in [1, n-1]$, we write $x_j = \rho_{j,n}(x) \in DF_j$ and set $x_n = x$. If $\epsilon_1 = -1$, then $x_1 \in \mathcal{O}^{e,e} \cup \mathcal{O}^{s_1,e}$. Similarly, if $\epsilon_1 = 1$, $x_1 \in \mathcal{O}^{e,e} \cup \mathcal{O}^{e,s_{l+1}}$. By Proposition 2.1, the assertion is true for $j = 1$. Thus assume $j \geq 2$, and write $(g.B, h.B_-) = \theta_{j-1}(x_{j-1})$, so that one has $h^{-1}g \in B_-w_{j-1}B$.

Case $\epsilon_j = -1$: One has $\theta_j(x_j) = (gp.B, h.B_-)$, where $p \in P_{\delta_j}$. Thus $h^{-1}gp \in B_-w_jB$, and so

$$B_-w_jB \subset B_-w_{j-1}BP_{\delta_j} = B_-w_{j-1}B \cup B_-w_{j-1}B\delta_jB.$$

One knows by [10, Appendix A] that

$$B_-w_{j-1}B\delta_jB \subset B_-w_{j-1}\delta_jB \cup B_-w_{j-1}B,$$

which proves the assertion.

Case $\epsilon_j = 1$: One has $\theta_j(x_j) = (g.B, hp.B_-)$, where $p \in P_{-\delta_j}$. Thus $p^{-1}h^{-1}g \in B_-w_jB$, and

$$B_-w_jB \subset P_{-\delta_j}B_-w_{j-1}B = B_-w_{j-1}B \cup B_-w_{j-1}B\delta_jB.$$

Again, by using [10, Appendix A] one concludes that $w_j \in \{w_{j-1}, \delta_j w_{j-1}\}$.

Q.E.D.

Let $W^{\mathbf{u},\mathbf{v},\sigma}$ be the elements of W^n satisfying the condition in Lemma 3.2. Thus $\Phi_n(Z_{\mathbf{u},\mathbf{v}}^\sigma) \subset W^{\mathbf{u},\mathbf{v},\sigma}$. We will show in Proposition 3.8 that $\Phi_n(Z_{\mathbf{u},\mathbf{v}}^\sigma) = W^{\mathbf{u},\mathbf{v},\sigma}$. Lemma 3.2 motivates the following definition.

Definition 3.3. Let $\mathbf{u} = (s_1, \dots, s_l)$ be a sequence of simple reflections. We say that a sequence $\gamma = (\gamma_1, \dots, \gamma_l)$ is a *subexpression of $\mathbf{u}$* if $\gamma_j \in \{e, s_j\}$ for all $j \in [1, l]$. Denote by $\Upsilon_{\mathbf{u}}$ the set of all subexpressions of $\mathbf{u}$. Let $\mathbf{v} = (s_{l+1}, \dots, s_n)$ be another sequence, and let $\sigma \in S_{l,n}$. We say that $\gamma = (\gamma_1, \dots, \gamma_n)$ is a *subexpression of $(\mathbf{u}, \mathbf{v})$ shuffled by σ* , or a *σ -shuffled subexpression of $(\mathbf{u}, \mathbf{v})$* , if $\gamma \in \Upsilon_{\sigma(\mathbf{u},\mathbf{v})}$.

Fix $(\mathbf{u}, \mathbf{v}, \sigma)$ as in Definition 3.1, and let $\gamma \in \Upsilon_{\sigma(\mathbf{u},\mathbf{v})}$. Construct the sequence $(\gamma^0, \gamma^1, \dots, \gamma^n)$ of Weyl group elements as follows. Let $\gamma^0 = e$, and for any $j \in [1, n]$, define

$$(22) \quad \gamma^j = \begin{cases} \gamma^{j-1}\gamma_j, & \text{if } \epsilon_j = -1 \\ \gamma_j\gamma^{j-1}, & \text{if } \epsilon_j = 1. \end{cases}$$

In other words, $(\gamma^1, \dots, \gamma^n) \in W^{\mathbf{u},\mathbf{v},\sigma}$. Conversely, it is easily seen that any sequence $(w_0, \dots, w_n)$ of elements in W satisfying $w_0 = e$ and $(w_1, \dots, w_n) \in W^{\mathbf{u},\mathbf{v},\sigma}$ uniquely defines a $\gamma \in \Upsilon_{\sigma(\mathbf{u},\mathbf{v})}$ such that $(\gamma^1, \dots, \gamma^n) = (w_1, \dots, w_n)$. Hence the map

$$\Upsilon_{\sigma(\mathbf{u},\mathbf{v})} \rightarrow W^{\mathbf{u},\mathbf{v},\sigma}, \quad \gamma = (\gamma_1, \dots, \gamma_n) \mapsto (\gamma^1, \dots, \gamma^n)$$

is a bijection.

Definition 3.4. For any $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$, let

$$C^\gamma = \Phi_n^{-1}(\gamma^1, \dots, \gamma^n) \cap Z_{\mathbf{u}, \mathbf{v}}^\sigma \quad \text{and} \quad C_\sigma^\gamma = \varphi^\sigma(C^\gamma) \subset Z_{\mathbf{u}, \mathbf{v}}.$$

One thus has the two disjoint unions

$$(23) \quad Z_{\mathbf{u}, \mathbf{v}}^\sigma = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}} C^\gamma \quad \text{and} \quad Z_{\mathbf{u}, \mathbf{v}} = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}} C_\sigma^\gamma.$$

3.4. Affine charts and coordinates on $Z_{\mathbf{u}, \mathbf{v}}^\sigma$. We introduce an atlas of 2^n affine charts on $Z_{\mathbf{u}, \mathbf{v}}^\sigma$ parametrized by $\Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$. As a first application, we show that all the C^γ 's are isomorphic to affine spaces.

Fix $(\mathbf{u}, \mathbf{v}, \sigma)$ as in Definition 3.1, and let $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$. We fix representatives in $N_G(H)$ of the Weyl group elements γ^j . Set $\widetilde{\gamma}^0 = e$, and for $j \in [1, n]$, let

$$(24) \quad \widetilde{\gamma}^j = \begin{cases} \widetilde{\gamma^{j-1}} \widetilde{\gamma}_j, & \text{if } \epsilon_j = -1 \\ \widetilde{\gamma}_j \gamma^{j-1}, & \text{if } \epsilon_j = 1. \end{cases}$$

Recall from (13) that $\sigma(\mathbf{u}, \mathbf{v}) = (\delta_1, \dots, \delta_n)$. Denote by α_j the simple root such that $\delta_j = s_{\alpha_j}$. For $j \in [1, n]$ and $z \in \mathbb{C}$, let

$$(25) \quad u_{\gamma, j}(z) = (p_{\gamma, j}(z), q_{\gamma, j}(z)) \in P_j^\sigma,$$

where

$$(26) \quad p_{\gamma, j}(z) = \begin{cases} x_{-\gamma_j \alpha_j}(z) \widetilde{\gamma}_j, & \text{if } \epsilon_j = -1 \\ [\widetilde{\gamma^{j-1}}^{-1} x_{\gamma_j \alpha_j}(z) \widetilde{\gamma^{j-1}}]_+, & \text{if } \epsilon_j = 1, \end{cases}$$

and

$$(27) \quad q_{\gamma, j}(z) = \begin{cases} [\widetilde{\gamma^{j-1}} x_{-\gamma_j \alpha_j}(z) \widetilde{\gamma^{j-1}}^{-1}]_-, & \text{if } \epsilon_j = -1 \\ x_{\gamma_j \alpha_j}(z) \widetilde{\gamma}_j^{-1}, & \text{if } \epsilon_j = 1. \end{cases}$$

Define

$$(28) \quad \mathcal{O}^\gamma = \{[u_{\gamma, 1}(z_1), \dots, u_{\gamma, n}(z_n)]_{Z_{\mathbf{u}, \mathbf{v}}^\sigma} \mid z = (z_1, \dots, z_n) \in \mathbb{C}^n\}.$$

The fact that the map

$$(29) \quad u_\gamma : \mathbb{C}^n \rightarrow \mathcal{O}^\gamma, \quad z \mapsto u_\gamma(z) = [u_{\gamma, 1}(z_1), \dots, u_{\gamma, n}(z_n)]_{Z_{\mathbf{u}, \mathbf{v}}^\sigma}$$

is an isomorphism is a consequence of applying inductively the following Lemma 3.5.

Lemma 3.5. *Let $\alpha \in \Gamma$, and let $B \times B_-$ act on $P_{s_\alpha} \times B_-$ by right multiplication. For any $\dot{w} \in N_G(H)$, the sets*

$$\{(x_\alpha(z) \bar{s}_\alpha, [\dot{w} x_\alpha(z) \dot{w}^{-1}]_-) \mid z \in \mathbb{C}\}, \quad \text{and} \quad \{(x_{-\alpha}(z), [\dot{w} x_{-\alpha}(z) \dot{w}^{-1}]_-) \mid z \in \mathbb{C}\}$$

intersect each $(B \times B_-)$ -orbits in exactly one point. Consequently, the two maps

$$\begin{aligned} (x_\alpha(z) \bar{s}_\alpha.B, B_-) &\mapsto (x_\alpha(z) \bar{s}_\alpha, [\dot{w} x_\alpha(z) \dot{w}^{-1}]_-), \\ (x_{-\alpha}(z).B, B_-) &\mapsto (x_{-\alpha}(z), [\dot{w} x_{-\alpha}(z) \dot{w}^{-1}]_-), \quad z \in \mathbb{C} \end{aligned}$$

are sections over open subsets of the fiber bundle

$$P_{s_\alpha} \times B_- \rightarrow (P_{s_\alpha} \times B_-)/(B \times B_-).$$

Proof. Suppose that $(x_\alpha(z)\bar{s}_\alpha, [\dot{w}x_\alpha(z)\dot{w}^{-1}]_-(b, b_-)) = (x_\alpha(z')\bar{s}_\alpha, [\dot{w}x_\alpha(z')\dot{w}^{-1}]_-)$ for some $z, z' \in \mathbb{C}$, $b \in B$, $b_- \in B_-$. Looking at the first component, one has $b = e$ and $z = z'$. Thus $b_- = e$. Proceed similarly for the other case.

Q.E.D.

Moreover, one sees that $Z_{\mathbf{u}, \mathbf{v}}^\sigma$ is covered by the open subsets $\mathcal{O}^\gamma$, $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$.

Definition 3.6. For any $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$, let

$$J(\gamma) = \{j \in [1, n] \mid (\gamma^j)^{-\epsilon_j} \alpha_j < 0\}.$$

Lemma 3.7. Let $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$. Let $(z_1, \dots, z_n) \in \mathbb{C}^n$ be such that $z_j = 0$ if $j \in J(\gamma)$. Then for any $j \in [1, n]$,

$$q_{\gamma, j}(z_j)^{-1} \cdots q_{\gamma, 1}(z_1)^{-1} p_{\gamma, 1}(z_1) \cdots p_{\gamma, j}(z_j) = \widetilde{\gamma}^j.$$

Proof. Let $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ with $z_j = 0$ if $j \in J(\gamma)$. Denote by $F_{\gamma, j}(z)$ the product on the left hand side, and set $F_{\gamma, 0} = e \in G$. Thus $F_{\gamma, 0}(z) = \widetilde{\gamma}^0$. So assume now that $j \in [1, n]$ and that $F_{\gamma, j-1}(z) = \widetilde{\gamma}^{j-1}$.

Case $\epsilon_j = -1$: Using the induction hypothesis, one has

$$\begin{aligned} F_{\gamma, j}(z) &= q_{\gamma, j}(z_j)^{-1} F_{\gamma, j-1}(z) p_{\gamma, j}(z_j) \\ &= [\widetilde{\gamma}^{j-1} x_{-\gamma_j \alpha_j}(-z_j) \widetilde{\gamma}^{j-1}]^{-1} \widetilde{\gamma}^{j-1} x_{-\gamma_j \alpha_j}(z_j) \widetilde{\gamma}^j \end{aligned}$$

If $j \notin J(\gamma)$, then $\gamma^j \alpha_j = \gamma^{j-1} \gamma_j \alpha_j > 0$, and so $F_{\gamma, j}(z) = \widetilde{\gamma}^j$. If $j \in J(\gamma)$, then $z_j = 0$, so $F_{\gamma, j}(z) = \widetilde{\gamma}^j$.

Case $\epsilon_j = 1$: One has

$$\begin{aligned} F_{\gamma, j}(z) &= q_{\gamma, j}(z_j)^{-1} F_{\gamma, j-1}(z) p_{\gamma, j}(z_j) \\ &= \widetilde{\gamma}^j x_{\gamma_j \alpha_j}(-z_j) \widetilde{\gamma}^{j-1} [\widetilde{\gamma}^{j-1}]^{-1} x_{\gamma_j \alpha_j}(z_j) \widetilde{\gamma}^{j-1} \end{aligned}$$

If $j \notin J(\gamma)$, then $(\gamma^j)^{-1} \alpha_j = (\gamma^{j-1})^{-1} \gamma_j \alpha_j > 0$, and so $F_{\gamma, j}(z) = \widetilde{\gamma}^j$. If $j \in J(\gamma)$, then $z_j = 0$, so $F_{\gamma, j}(z) = \widetilde{\gamma}^j$.

Q.E.D.

The following Proposition 3.8 shows that each C^γ in the combinatorially defined decomposition of $Z_{\mathbf{u}, \mathbf{v}}^\sigma$ in (23) is isomorphic to an affine space.

Proposition 3.8. For any $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$, one has

$$C^\gamma = \{u_\gamma(z) \in \mathcal{O}^\gamma \mid z_j = 0, \text{ if } j \in J(\gamma)\}.$$

Proof. Recall that C^γ has been introduced in Definition 3.4. Denote by $\tilde{C}^\gamma$ the set on the right hand side. Recall that an element

$$[(g_1, h_1), \dots, (g_n, h_n)]_{DF_n}$$

lies in C^γ if and only if for every $j \in [1, n]$, $(g_1 \cdots g_j.B, h_1 \cdots h_j.B_-) \in G_{\text{diag}}(\gamma^j)$, which is equivalent to

$$(30) \quad (h_1 \cdots h_j)^{-1} g_1 \cdots g_j \in B_- \gamma^j B, \quad j \in [1, n].$$

By Lemma 3.7, one has $\tilde{C}^\gamma \subset C^\gamma$, so we need to prove

$$(31) \quad C^\gamma \subset \tilde{C}^\gamma.$$

For $k \in [1, n]$, let

$$Z_k = P_1^\sigma \times_{B \times B_-} \cdots \times_{B \times B_-} P_k^\sigma / (B \times B_-) \subset DF_k.$$

Define the following two subsets of Z_k .

$$\begin{aligned} C^{(\gamma_1, \dots, \gamma_k)} &= \{[(g_1, h_1), \dots, (g_k, h_k)]_{DF_k} \in Z_k \mid (30) \text{ holds for } j \in [1, k]\} \\ \tilde{C}^{(\gamma_1, \dots, \gamma_k)} &= \{[u_{\gamma,1}(z_1), \dots, u_{\gamma,k}(z_k)]_{DF_k} \in Z_k \mid z_i = 0, \text{ if } i \in [1, k] \cap J(\gamma)\}. \end{aligned}$$

Fix $x = [(g_1, h_1), \dots, (g_n, h_n)]_{DF_n} \in C^\gamma$, i.e x satisfies (30). Write $x_k = \rho_{k,n}(x)$ for $k \in [1, n-1]$ and set $x_n = x$. Then $x_k \in C^{(\gamma_1, \dots, \gamma_k)}$ for every $k \in [1, n]$. We show by induction on k that $x_k \in \tilde{C}^{(\gamma_1, \dots, \gamma_k)}$ for every $k \in [1, n]$. Then (31) is the statement for $k = n$.

Assume that $\epsilon_1 = -1$, the case $\epsilon_1 = 1$ being similar. So $h_1 \in B_-$ and $g_1 \in P_{\delta_1}$. Then $g_1 \in B_- \gamma_1 B$. Hence one can write $g_1 = x_{-\gamma_1 \alpha_1}(z_1) \bar{\gamma}_1 b_1$ with $b_1 \in B$ and $z_1 = 0$ if $\gamma_1 = \delta_1$, that is if $1 \in J(\gamma)$. Thus

$$(g_1.B, h_1.B_-) = (x_{-\gamma_1 \alpha_1}(z_1) \bar{\gamma}_1.B, [x_{-\gamma_1 \alpha_1}(z_1)]_-.B_-) \in \tilde{C}^{(\gamma_1)}.$$

Let now $k \in [1, n-1]$, and assume that $C^{(\gamma_1, \dots, \gamma_k)} \subset \tilde{C}^{(\gamma_1, \dots, \gamma_k)}$. Once again the two cases of $\epsilon_{k+1} = 1$ or $\epsilon_{k+1} = -1$ can be treated similarly, so we can assume that $\epsilon_{k+1} = -1$. So $\gamma^{k+1} = \gamma^k \gamma_{k+1}$. By the induction hypothesis, $x_k \in \tilde{C}^{(\gamma_1, \dots, \gamma_k)}$, thus there exist $g'_{k+1} \in P_{\delta_{k+1}}$, $h'_{k+1} \in B_-$ and $(z_1, \dots, z_k) \in \mathbb{C}^k$ with $z_j = 0$ if $j \in [1, k] \cap J(\gamma)$, such that

$$x_{k+1} = [u_{\gamma,1}(z_1), \dots, u_{\gamma,k}(z_k), (g'_{k+1}, h'_{k+1})]_{DF_{k+1}}.$$

Since $x_{k+1} \in C^{(\gamma_1, \dots, \gamma_k)}$, we have

$$h'_{k+1}{}^{-1} q_{\gamma,k}(z_k)^{-1} \cdots q_{\gamma,1}(z_1)^{-1} p_{\gamma,1}(z_1) \cdots p_{\gamma,k}(z_k) g'_{k+1} = h'_{k+1}{}^{-1} \widetilde{\gamma^k} g'_{k+1} \in B_- \gamma^{k+1} B.$$

Hence $\widetilde{\gamma^k} g'_{k+1} \in B_- \gamma^{k+1} B$. If $\gamma_{k+1} = \delta_{k+1}$, then $\gamma^{k+1} = \gamma^k \delta_{k+1} \neq \gamma^k$ and one cannot have $g'_{k+1} \in B$. Similarly, if $\gamma_{k+1} = e$, then one cannot have $g'_{k+1} \in \delta_{k+1} B$. Thus g'_{k+1} can be written as $x_{-\gamma_{k+1} \alpha_{k+1}}(z_{k+1}) \bar{\gamma}_{k+1} b_{k+1}$ for some $z_{k+1} \in \mathbb{C}$ and $b_{k+1} \in B$. Furthermore, if $k+1 \in J(\gamma)$, that is if $\gamma^k \gamma_{k+1} \alpha_{k+1} < 0$, then one must have $z_{k+1} = 0$. If $k+1 \notin J(\gamma)$, then $\widetilde{\gamma^k} g'_{k+1} \in B_- \gamma^{k+1} B$ for any value of z_{k+1} . Moreover one can write $h'_{k+1} = [\widetilde{\gamma^k} x_{-\gamma_{k+1} \alpha_{k+1}}(z_{k+1}) \widetilde{\gamma^k}^{-1}]_{-b_{-(k+1)}}$, for some $b_{-(k+1)} \in B_-$. Hence

$$x_{k+1} = [u_{\gamma,1}(z_1), \dots, u_{\gamma,k}(z_k), u_{\gamma,k+1}(z_{k+1})]_{Z_{k+1}} \in \tilde{C}^{(\gamma_1, \dots, \gamma_{k+1})}.$$

Q.E.D.

3.5. The H -action on $\mathcal{O}^\gamma$ in the z -coordinates. Fix again $(\mathbf{u}, \mathbf{v}, \sigma)$ as in Definition 3.3. Let H act diagonally on $Z_{\mathbf{u}, \mathbf{v}}$. This action corresponds, before applying φ^σ , to the action on $Z_{\mathbf{u}, \mathbf{v}}^\sigma$ given by

$$h \cdot [(g_1, h_1), \dots, (g_n, h_n)]_{Z_{\mathbf{u}, \mathbf{v}}^\sigma} = [(hg_1, hh_1), (g_2, h_2), \dots, (g_n, h_n)]_{Z_{\mathbf{u}, \mathbf{v}}^\sigma}, \quad h \in H, (g_i, h_i) \in P_i^\sigma.$$

Let $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$. We show in this section that $\mathcal{O}^\gamma$ is invariant under the H -action, and compute the action of H in the coordinates $(z_1, \dots, z_n)$. For any $j \in [1, n]$, one can write $\gamma^j = (\gamma_{\mathbf{v}}^j)^{-1} \gamma_{\mathbf{u}}^j$, with

$$(32) \quad \gamma_{\mathbf{v}}^j = \prod_{1 \leq k \leq j, \epsilon_k = 1} \gamma_k, \quad \gamma_{\mathbf{u}}^j = \prod_{1 \leq k \leq j, \epsilon_k = -1} \gamma_k,$$

where the index in both products is increasing. Set $\gamma_{\mathbf{u}}^0 = \gamma_{\mathbf{v}}^0 = e$. In particular, one can write $\gamma^n = \gamma_{\mathbf{v}}^{-1} \gamma_{\mathbf{u}}$, with $\gamma_{\mathbf{v}} = \gamma_{\mathbf{v}}^n$, $\gamma_{\mathbf{u}} = \gamma_{\mathbf{u}}^n$.

Proposition 3.9. *Let $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$. Then $\mathcal{O}^\gamma$ is invariant under H , and each coordinate function z_j , $j \in [1, n]$, is a weight function for H , with*

$$h \cdot z_j = \begin{cases} h^{-\gamma_{\mathbf{u}}^j \alpha_j} z_j, & \epsilon_j = -1 \\ h^{\gamma_{\mathbf{v}}^j \alpha_j} z_j, & \epsilon_j = 1. \end{cases}$$

Proof. Let $h_1, h_2 \in H$. Let $\alpha \in \Gamma$, and $\delta \in \{e, s_\alpha\}$. For any $\dot{w} \in N_G(H)$ and $z \in \mathbb{C}$, one has

$$(33) \quad (h_1, h_2)(x_{-\delta\alpha}(z)\bar{\delta}, [\dot{w}x_{-\delta\alpha}(z)\dot{w}^{-1}]_-) = (x_{-\delta\alpha}(h_1^{-\delta\alpha}z)\bar{\delta}, [\dot{w}x_{-\delta\alpha}(h_2^{-w\delta\alpha}z)\dot{w}^{-1}]_-)(h_1^\delta, h_2)$$

$$(34) \quad (h_1, h_2)([\dot{w}^{-1}x_{\delta\alpha}(z)\dot{w}]_+, x_{\delta\alpha}(z)\bar{\delta}^{-1}) = ([\dot{w}^{-1}x_{\delta\alpha}(h_1^{w^{-1}\delta\alpha}z)\dot{w}]_+, x_{\delta\alpha}(h_2^{\delta\alpha}z)\bar{\delta}^{-1})(h_1, h_2^\delta)$$

Let $h \in H$ and $j \in [1, n]$. If $\epsilon_j = -1$, using (33), one gets

$$\begin{aligned} (h^{\gamma_{\mathbf{u}}^{j-1}}, h^{\gamma_{\mathbf{v}}^{j-1}})(p_{\gamma, j}(z_j), q_{\gamma, j}(z_j)) &= (p_{\gamma, j}(h^{-\gamma_{\mathbf{u}}^j \alpha_j} z_j), q_{\gamma, j}(h^{-\gamma_{\mathbf{v}}^{j-1} \gamma_j^{-1} \gamma_j \alpha_j} z_j))(h^{\gamma_{\mathbf{u}}^j}, h^{\gamma_{\mathbf{v}}^{j-1}}) \\ &= (p_{\gamma, j}(h^{-\gamma_{\mathbf{u}}^j \alpha_j} z_j), q_{\gamma, j}(h^{-\gamma_{\mathbf{u}}^j \alpha_j} z_j))(h^{\gamma_{\mathbf{u}}^j}, h^{\gamma_{\mathbf{v}}^j}). \end{aligned}$$

The same can be proved using (34) if $\epsilon_j = 1$. Thus an induction on j yields the result.

Q.E.D.

Definition 3.10. We denote by $\nu_{\gamma, j}$ the weight of the coordinate z_j for the action of H . That is

$$\nu_{\gamma, j} = \begin{cases} -\gamma_{\mathbf{u}}^j \alpha_j, & \text{if } \epsilon_j = -1 \\ \gamma_{\mathbf{v}}^j \alpha_j, & \text{if } \epsilon_j = 1. \end{cases}$$

3.6. Positive and distinguished σ -shuffled subexpressions. We now prove a few facts on the combinatorics of shuffled subexpressions that will be useful later. In particular, we introduce positive and distinguished shuffled subexpressions.

Fix $(\mathbf{u}, \mathbf{v}, \sigma)$ as in Definition 3.1, and let $\sigma(\mathbf{u}, \mathbf{v}) = (\delta_1, \dots, \delta_n)$ be as in (13).

Definition 3.11. Let $\gamma \in \Upsilon_{\mathbf{u}}$. We say that γ is *positive*, if

$$\gamma_1 \cdots \gamma_{j-1} = (\gamma_1 \cdots \gamma_j) \triangleleft s_j,$$

for any $j \in [1, l]$. We write $\Upsilon_{\mathbf{u}}^+$ for the set of all positive subexpressions of $\mathbf{u}$. We say that $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$ is σ -*positive*, or that γ is a *positive σ -shuffled subexpression of $(\mathbf{u}, \mathbf{v})$* , if for any $j \in [1, n]$,

$$\gamma^{j-1} = \begin{cases} \gamma^j \triangleleft \delta_j, & \text{if } \epsilon_j = -1 \\ \delta_j \triangleright \gamma^j, & \text{if } \epsilon_j = 1. \end{cases}$$

Denote by $\Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$ the set of all positive σ -shuffled subexpressions of $(\mathbf{u}, \mathbf{v})$.

Remark 3.12. Recall that $\gamma^j \triangleleft \delta_j = \min\{\gamma^j, \gamma^j \delta_j\}$, and $\delta_j \triangleright \gamma^j = \min\{\gamma^j, \delta_j \gamma^j\}$. So one sees that γ is positive if and only if for every $j \in [1, n]$,

$$\begin{cases} \gamma^{j-1} < \gamma^{j-1} \delta_j, & \text{when } \epsilon_j = -1 \\ \gamma^{j-1} < \delta_j \gamma^{j-1}, & \text{when } \epsilon_j = 1. \end{cases}$$

The notion of positivity of subexpressions has previously been studied in [13, 16]. Our definition of positivity for shuffled subexpressions is a generalization of the notion of “positive double subexpression” found in [16]. Indeed, if $\mathbf{u}$ and $\mathbf{v}$ are reduced, the two notions coincide. See Section 3.8 for more details.

Example 3.13. In the root system A_4 , let

$$\mathbf{u} = (\mathfrak{s}_3, \mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_2, \mathfrak{s}_3), \text{ and } \mathbf{v} = (\mathfrak{s}_4, \mathfrak{s}_2, \mathfrak{s}_1, \mathfrak{s}_4)$$

(We follow the Bourbaki convention for the labelling of simple roots in simple root systems).

Let

$$\begin{aligned} \epsilon(\sigma) &= (-1, -1, 1, -1, 1, 1, -1, -1, 1) \\ \sigma(\mathbf{u}, \mathbf{v}) &= (\mathfrak{s}_3, \mathfrak{s}_2, \mathfrak{s}_4, \mathfrak{s}_3, \mathfrak{s}_2, \mathfrak{s}_1, \mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4), \end{aligned}$$

and consider the shuffled subexpressions

$$\gamma_1 = (e, e, e, \mathfrak{s}_3, e, e, \mathfrak{s}_2, \mathfrak{s}_3, \mathfrak{s}_4) \text{ and } \gamma_2 = (e, e, \mathfrak{s}_4, \mathfrak{s}_3, \mathfrak{s}_2, \mathfrak{s}_1, \mathfrak{s}_2, \mathfrak{s}_3, e).$$

Then γ_1 is positive, but γ_2 is not.

Let

$$(35) \quad u = s_1 * \cdots * s_l, \quad v = s_{l+1} * \cdots * s_n.$$

We wish to show that elements $w \in W$ with $w \leq v^{-1} * u$ are in 1-1 correspondance with positive σ -shuffled subexpressions.

Definition 3.14. For $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$, let

$$\begin{aligned} \gamma_{[1, l]} &= (\gamma_{\sigma(1)}, \dots, \gamma_{\sigma(l)}) \in \Upsilon_{\mathbf{u}}, \\ \gamma_{[l+1, n]} &= (\gamma_{\sigma(l+1)}, \dots, \gamma_{\sigma(n)}) \in \Upsilon_{\mathbf{v}}. \end{aligned}$$

Lemma 3.15. For any $w \in W$ with $w \leq v^{-1} * u$, there is a unique $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$ such that $\gamma^n = w$. Moreover γ has the following properties.

- 1) For any $j \in [1, n]$, $l(\gamma^j) = l(\gamma_{\mathbf{v}}^j) + l(\gamma_{\mathbf{u}}^j)$;
- 2) $\gamma_{[1, l]}$ is a positive subexpression of $\mathbf{u}$, and $\gamma_{[l+1, n]}$ is a positive subexpression of $\mathbf{v}$.

Proof. Let $w \in W$ such that $w \leq v^{-1} * u$. Set $w_n = w$, and for any $j \in [1, n]$, let

$$w_{j-1} = \begin{cases} w_j \triangleleft \delta_j, & \text{if } \epsilon_j = -1 \\ \delta_j \triangleright w_j, & \text{if } \epsilon_j = 1. \end{cases}$$

Then, by [10, Lemma A.3], one has $w_0 = v \triangleright w \triangleleft u^{-1} = e$. Let $\gamma_j = w_{j-1}^{\epsilon_j} w_j^{-\epsilon_j}$. Then $\gamma = (\gamma_1, \dots, \gamma_n)$ is a positive σ -shuffled subexpression, and is clearly the unique one satisfying $\gamma^n = w$. We now prove that

$$(36) \quad l(\gamma^j) = l(\gamma_{\mathbf{v}}^j) + l(\gamma_{\mathbf{u}}^j), \quad j \in [1, n],$$

for any $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$. One clearly has $l(\gamma^1) = l(\gamma_{\mathbf{v}}^1) + l(\gamma_{\mathbf{u}}^1)$. Thus suppose $j \geq 2$, and that $l(\gamma^{j-1}) = l(\gamma_{\mathbf{v}}^{j-1}) + l(\gamma_{\mathbf{u}}^{j-1})$. If $\gamma_j = e$, then (36) holds. So assume $\gamma_j = \delta_j$. Suppose that $\epsilon_j = -1$, the other case being similar. Then $\gamma^j = \gamma^{j-1} \delta_j > \gamma^{j-1}$ and one has $\gamma_{\mathbf{u}}^j = \gamma_{\mathbf{u}}^{j-1} \delta_j > \gamma_{\mathbf{u}}^{j-1}$. Thus

$$l(\gamma^j) = l(\gamma_{\mathbf{v}}^{j-1}) + l(\gamma_{\mathbf{u}}^{j-1}) + 1 = l(\gamma_{\mathbf{v}}^j) + l(\gamma_{\mathbf{u}}^j).$$

This proves 1). Let now $j \in [1, n]$, and suppose that $\epsilon_j = 1$. One has $\delta_j \gamma^{j-1} > \gamma^{j-1}$. Since $l(\gamma^{j-1}) = l(\gamma_{\mathbf{v}}^{j-1}) + l(\gamma_{\mathbf{u}}^{j-1})$, this implies that $\delta_j (\gamma_{\mathbf{v}}^{j-1})^{-1} > (\gamma_{\mathbf{v}}^{j-1})^{-1}$. Or in other words, $(\gamma_{\mathbf{v}}^{j-1})^{-1} = \delta_j \triangleright (\gamma_{\mathbf{v}}^j)^{-1}$, which is equivalent, by [10, Lemma A3], to

$$\gamma_{\mathbf{v}}^j \triangleleft \delta_j = \gamma_{\mathbf{v}}^{j-1}.$$

This shows that $(\gamma_{\sigma(l+1)}, \dots, \gamma_{\sigma(n)})$ is a positive subexpression of $\mathbf{v}$. If $\epsilon_j = -1$, a similar computation shows that $(\gamma_{\sigma(1)}, \dots, \gamma_{\sigma(l)})$ is a positive subexpression of $\mathbf{u}$.

Q.E.D.

Remark 3.16. Recall the Weyl group representatives $\tilde{\gamma}^j$ from (24). If γ is positive, it follows from Lemma 3.15 that $\tilde{\gamma}^j = \overline{\gamma^j}$, for all $j \in [1, n]$.

Definition 3.17. If $w \leq u^{-1} * v$, we denote by $\gamma_w \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$ the unique positive σ -shuffled subexpression satisfying $\gamma_w^n = w$.

On the other hand, for any $\gamma \in \Upsilon_{\sigma(\mathbf{u}), \mathbf{v}}$, one has $\gamma^n = \gamma_{\mathbf{v}}^{-1} \gamma_{\mathbf{u}} \leq v^{-1} * u$. Thus we have shown the following.

Corollary 3.18. *The map*

$$\{w \in W \mid w \leq v^{-1} * u\} \rightarrow \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+, \quad w \mapsto \gamma_w$$

is a bijection. The inverse is given by $\gamma \mapsto \gamma^n$.

Lemma 3.19. *For any $\gamma \in \Upsilon_{\sigma(\mathbf{u}), \mathbf{v}}$, one has $l(\gamma^n) \leq |J(\gamma)|$, with equality if and only if γ is positive.*

Proof. We proceed by induction. The result is obvious when $n = 1$, so let $n \geq 2$. We assume that $\epsilon_n = 1$, the case $\epsilon_n = -1$ being treated similarly. Then one has $\sigma(n) = n$. Let $\mathbf{v}' = (s_{l+1}, \dots, s_{n-1})$, and $\gamma' = (\gamma_1, \dots, \gamma_{n-1})$. Then $\gamma' \in \Upsilon_{\sigma'(\mathbf{u}), \mathbf{v}'}$, where $\sigma'(j) = \sigma(j)$, for all $j \in [1, n-1]$. By the induction hypothesis, the result holds for γ' . Recall that we denote by α_n the simple root such that $\delta_n = s_{\alpha_n}$.

Case 1: $(\gamma^n)^{-1} \alpha_n > 0$ and $\gamma_n = e$.

Then $n \notin J(\gamma)$, and $|J(\gamma)| = |J(\gamma')| \geq l(\gamma^{n-1}) = l(\gamma^n)$. Since $(\gamma^{n-1})^{-1}\alpha_n > 0$ one has $\delta_n \gamma^{n-1} > \gamma^{n-1}$. Hence γ is positive if and only if γ' is positive. So we are done.

Case 2: $(\gamma^n)^{-1}\alpha_n > 0$ and $\gamma_n = \delta_n$.

Then $n \notin J(\gamma)$, and $|J(\gamma)| = |J(\gamma')| \geq l(\gamma^{n-1}) = l(\gamma^n) + 1$. Since $(\gamma^{n-1})^{-1}\alpha_n < 0$ one has $\delta_n \gamma^{n-1} < \gamma^{n-1}$, so γ is not positive.

Case 3: $(\gamma^n)^{-1}\alpha_n < 0$, and $\gamma_n = e$.

Then $n \in J(\gamma)$, and $|J(\gamma)| = |J(\gamma')| + 1 \geq l(\gamma^{n-1}) + 1 = l(\gamma^n) + 1$. Since $(\gamma^{n-1})^{-1}\alpha_n < 0$, γ is not positive.

Case 4: $(\gamma^n)^{-1}\alpha_n < 0$ and $\gamma_n = \delta_n$.

Then $n \in J(\gamma)$, and $|J(\gamma)| = |J(\gamma')| + 1 \geq l(\gamma^{n-1}) + 1 = l(\gamma^n)$. Since $(\gamma^{n-1})^{-1}\alpha_n > 0$, γ is positive if and only if γ' is positive. So we are done.

Q.E.D.

Remark 3.20. If $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$, then $j \in J(\gamma)$ if and only if $\gamma_j = \delta_j$. Indeed, $j \in J(\gamma)$ if and only if $\delta_j(\gamma^j)^{\epsilon_j} < (\gamma^j)^{\epsilon_j}$. But since γ is positive, this happens exactly when $\gamma_j = \delta_j$.

Definition 3.21. Let $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$. We say that γ is *distinguished*, if $C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ is non-empty. Denote by $\Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^d$ the set of all distinguished σ -shuffled subexpressions.

The open subset $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} \subset Z_{\mathbf{u}, \mathbf{v}}^\sigma$ has an intrinsic description. Indeed, for $j \in [1, n]$, let

$$B_j^\sigma = \begin{cases} B\delta_j B \times B_-, & \epsilon_j = -1 \\ B \times B_- \delta_j B_-, & \epsilon_j = 1. \end{cases}$$

Then $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} = B_1^\sigma \times_{B \times B_-} \cdots \times_{B \times B_-} B_n^\sigma / (B \times B_-)$. Note that we have $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} = I_{\mathbf{u}, \mathbf{v}}^\sigma(\mathcal{O}^{\mathbf{u}, \mathbf{v}})$, recall (2) and (21).

Definition 3.22. For any $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$, let

$$I(\gamma) = \{j \in [1, n] \mid \gamma_j = \delta_j\}.$$

Lemma 3.23. A σ -shuffled subexpression $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$ is distinguished if and only if $J(\gamma) \subset I(\gamma)$.

Proof. Recall (25). Notice that if $j \in I(\gamma)$, then $u_{\gamma, j}(z_j) \in B_j^\sigma$ for any $z_j \in \mathbb{C}$. And if $j \notin I(\gamma)$, then $u_{\gamma, j}(z_j) \in B_j^\sigma$ if and only if $z_j \neq 0$.

Suppose that $J(\gamma) \subset I(\gamma)$. Let $z = (z_1, \dots, z_n) \in \mathbb{C}^n$, with $z_j = 0$ if $j \in J(\gamma)$, and $z_j \neq 0$ if $j \notin J(\gamma)$. Then by Proposition 3.8, $u_\gamma(z) \in C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$. Conversely, if $J(\gamma)$ is not contained in $I(\gamma)$, let $j \in J(\gamma)$ such that $j \notin I(\gamma)$. If $u_\gamma(z) \in C^\gamma$, by Proposition 3.8, one must have $z_j = 0$. But then $u_\gamma(z) \notin \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$.

Q.E.D.

Remark 3.24. By Remark 3.20, any positive σ -shuffled subexpression γ is distinguished. Conversely, a distinguished σ -shuffled subexpression is positive if and only if $J(\gamma) = I(\gamma)$.

Let $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^d$. Then $u_\gamma(z) \in \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ if and only if $z_j \neq 0$ when $j \notin I(\gamma)$. Combining this with Proposition 3.8, one obtains the following

Proposition 3.25. *Let $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^d$ and let $(z_1, \dots, z_n) \in \mathbb{C}^n$ with*

$$z_j \in \begin{cases} \{0\}, & \text{if } j \in J(\gamma) \\ \mathbb{C}, & \text{if } I(\gamma) \setminus J(\gamma) \\ \mathbb{C}^*, & \text{if } j \notin I(\gamma). \end{cases}$$

Then the map

$$\mathbb{C}^{|I(\gamma) \setminus J(\gamma)|} \times (\mathbb{C}^*)^{|I(\gamma)^c|} \rightarrow C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})},$$

where $I(\gamma)^c = [1, n] \setminus I(\gamma)$, sending z to $u_\gamma(z)$ is an isomorphism.

3.7. The subvarieties A_w . Let $\mathbf{u} = (s_1, \dots, s_l)$ and $\mathbf{v} = (s_{l+1}, \dots, s_n)$ be two sequences of simple reflections. Consider the double flag variety $Z_{\mathbf{u}, \mathbf{v}}$, and recall the map $\theta_{\mathbf{u}, \mathbf{v}}$ from (8).

Definition 3.26. For $w \in W$, let

$$A_w = \theta_{\mathbf{u}, \mathbf{v}}^{-1}(G_{\text{diag}}(w)) \subset Z_{\mathbf{u}, \mathbf{v}},$$

and if σ is a (l, n) -shuffle, let

$$A_w^\sigma = I_{\mathbf{u}, \mathbf{v}}^\sigma(A_w) \subset Z_{\mathbf{u}, \mathbf{v}}^\sigma,$$

recall (21).

Let u, v be as in (35), and let $w \in W$ with $w \leq v^{-1} * u$. Then for any $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$ with $\gamma^n = w$, by definition of C^γ one has $C^\gamma \subset A_w^\sigma$.

Conversely, suppose that A_w is non-empty. Since $\theta_{\mathbf{u}, \mathbf{v}}(Z_{\mathbf{u}, \mathbf{v}}) = \overline{\mathcal{O}^{u, v}}$, Proposition 2.1 implies that $w \leq v^{-1} * u$. Thus one gets

Proposition 3.27. *Let $w \in W$. Then A_w is non-empty if and only if $w \leq v^{-1} * u$. Moreover, for any (l, n) -shuffle σ , one has*

$$A_w^\sigma = \bigsqcup_{\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})} : \gamma^n = w} C^\gamma.$$

Proposition 3.28. *For any $w \in W$ with $w \leq v^{-1} * u$, A_w is a smooth, irreducible subvariety of $Z_{\mathbf{u}, \mathbf{v}}$ of codimension $l(w)$ in $Z_{\mathbf{u}, \mathbf{v}}$.*

Proof. The image of $\theta_{\mathbf{u}, \mathbf{v}}$ can be written as

$$\theta_{\mathbf{u}, \mathbf{v}}(Z_{\mathbf{u}, \mathbf{v}}) = \overline{\mathcal{O}^{u, v}} = \bigsqcup_{u_1 \leq u, v_1 \leq v} \mathcal{O}^{u_1, v_1}.$$

On the other hand, it is known by [14, Theorem 1.4] that $\mathcal{O}^{u_1, v_1}$ and $G_{\text{diag}}(w)$ intersect transversally, for any $u_1, v_1, w \in W$. Hence $\theta_{\mathbf{u}, \mathbf{v}}$ is transverse to $G_{\text{diag}}(w)$, and so A_w is smooth, and every component of A_w has codimension

$$\dim_{\mathbb{C}} Z_{\mathbf{u}, \mathbf{v}} - \dim_{\mathbb{C}} A_w = \dim_{\mathbb{C}} DF_1 - \dim_{\mathbb{C}} G_{\text{diag}}(w) = l(w).$$

Choose now any $\sigma \in S_{l, n}$, and consider $\gamma_w \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$. By Lemma 3.19, C^{γ_w} has codimension $l(w)$ in $Z_{\mathbf{u}, \mathbf{v}}^\sigma$, and so is open in A_w^σ . Furthermore, by Proposition 3.27, the complement of C^{γ_w} in A_w^σ is a finite union of subvarieties of lower dimension. Hence C^{γ_w} is dense in A_w^σ , and so A_w^σ is irreducible in the Zariski topology.

Q.E.D.

Theorem 3.29. *Let $w \in W$ with $w \leq v^{-1} * u$. Then*

$$C^{\gamma w} = \mathcal{O}^{\gamma w} \cap A_w^\sigma = \mathcal{O}^{\gamma w} \cap \overline{A_w^\sigma}$$

where $\overline{A_w^\sigma}$ is the Zariski closure of A_w^σ .

Proof. One has the inclusions $C^{\gamma w} \subseteq \mathcal{O}^{\gamma w} \cap A_w^\sigma \subseteq \mathcal{O}^{\gamma w} \cap \overline{A_w^\sigma}$. On the other hand, both $C^{\gamma w}$ and $\mathcal{O}^{\gamma w} \cap \overline{A_w^\sigma}$ are closed, irreducible varieties of $\mathcal{O}^{\gamma w}$ of the same dimension. Hence they must be equal.

Q.E.D.

3.8. Relation to the Webster-Yakimov decompositions. Let $u, v, w \in W$ with $w \leq v^{-1} * u$, and consider $\mathcal{O}_w^{u,v} \subset DF_1$. In [16], Webster and Yakimov introduced decompositions of $\mathcal{O}_w^{u,v}$ into subvarieties isomorphic to $\mathbb{C}^{n_1} \times (\mathbb{C}^*)^{n_2}$ for some integers n_1, n_2 . We show in this section that the decomposition of A_w^σ in Proposition 3.27 recovers the decompositions in [16] when $\mathbf{u}$ and $\mathbf{v}$ are reduced.

We start by recalling the notations used in [16]. The set of simple roots of G is labelled by $[1, r]$, and $W \times W$ is considered as a Coxeter group with simple reflections s_{-i}, s_i , for $i \in [1, r]$. Let $(u, v) \in W \times W$, and let $\mathbf{i} = (s_{i_1}, \dots, s_{i_n})$, with $i_k \in [1, r] \cup [-r, -1]$, be a reduced expression for (u, v) . We let $\epsilon(k) = 1$, if $i_k \in [1, r]$, and $\epsilon(k) = -1$, if $i_k \in [-r, -1]$. A sequence $\mathbf{w} = (w_{(0)}, \dots, w_{(n)})$ of elements in W is called a *double subexpression of $\mathbf{i}$* , if $w_{(0)} = e$, and for $k \in [1, n]$,

$$w_{(k)} \in \begin{cases} \{w_{(k-1)}, w_{(k-1)}s_{|i_k|}\}, & \epsilon(k) = 1, \\ \{w_{(k-1)}, s_{|i_k|}w_{(k-1)}\}, & \epsilon(k) = -1. \end{cases}$$

Webster and Yakimov call $\mathbf{w}$ *double distinguished* if $w_{(k)}^{\epsilon(k)} = w_{(k-1)}^{\epsilon(k)}s_{|i_k|}$ for all $k \in [1, n]$ such that $w_{(k-1)}^{\epsilon(k)}s_{|i_k|} < w_{(k-1)}^{\epsilon(k)}$. For each double subexpression $\mathbf{w}$, we let

$$\begin{aligned} J_{\mathbf{w}}^0 & \text{ be the set of indices } k \in [1, n] \text{ such that } w_{(k-1)} = w_{(k)}, \\ J_{\mathbf{w}}^+ & \text{ be the set of indices } k \in [1, n] \text{ such that } w_{(k-1)} < w_{(k)}, \\ J_{\mathbf{w}}^- & \text{ be the set of indices } k \in [1, n] \text{ such that } w_{(k-1)} > w_{(k)}. \end{aligned}$$

A double distinguished subexpression is called *positive*, if $J_{\mathbf{w}}^-$ is empty.

We translate now the notations of [16] into ours. Let

$$\mathbf{u} = (s_{-i_{k_1}}, \dots, s_{-i_{k_l}}), \quad \mathbf{v} = (s_{i_{m_1}}, \dots, s_{i_{m_{n-l}}}),$$

where $\{k_1 < \dots < k_l\}$ is the set of indices in $[1, n]$ with $\epsilon(k_j) = -1$, $j \in [1, l]$, and $\{m_1 < \dots < m_{n-l}\}$ the set of indices in $[1, n]$ with $\epsilon(m_t) = 1$, $t \in [1, n-l]$. Then $\mathbf{u}$ and $\mathbf{v}$ are reduced expressions. Let σ be the unique (l, n) -shuffle such that $\sigma(\mathbf{u}, \mathbf{v}) = (s_{|i_1|}, \dots, s_{|i_n|})$. It is clear that $\mathbf{w}$ is a double subexpression of $\mathbf{i}$ if and only if there exists $\gamma \in \Upsilon_{\sigma(\mathbf{u}, \mathbf{v})}$ such that $w_{(k)}^{-1} = \gamma^k$, for any $k \in [1, n]$. Moreover, $\mathbf{w}$ is double distinguished if and only if γ is distinguished, and in this case

$$J_{\mathbf{w}}^0 = [1, n] \setminus I(\gamma), \quad J_{\mathbf{w}}^+ = J(\gamma), \quad J_{\mathbf{w}}^- = I(\gamma) \setminus J(\gamma).$$

Then $\mathbf{w}$ is positive if and only if γ is positive.

Since $\mathbf{u}$ and $\mathbf{v}$ are reduced, the restriction $\theta_n|_{\mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})}}: \mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})} \rightarrow \mathcal{O}^{u,v}$ is an isomorphism, recall (10). One has

$$\mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})} = \bigsqcup_{\gamma \in \Upsilon_{\mathbf{u},\mathbf{v},\sigma}^d} \mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})} \cap C^\gamma.$$

The subvariety $\theta_n(\mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})} \cap C^\gamma) \subset \mathcal{O}^{u,v}$ is called $\mathcal{P}_{\mathbf{u},\mathbf{v}}^{\mathbf{w}}$ in [16, Section 5], Thus one has

$$\mathcal{O}^{u,v} = \bigsqcup_{\mathbf{w}} \mathcal{P}_{\mathbf{u},\mathbf{v}}^{\mathbf{w}},$$

where $\mathbf{w}$ runs over all double distinguished subexpressions. The “piece” $\mathcal{P}_{\mathbf{u},\mathbf{v}}^{\mathbf{w}}$ is given in [16, Theorem 5.2] the following parametrization. Let $\gamma \in \Upsilon_{\mathbf{u},\mathbf{v},\sigma}^d$ be such that $(w_{(0)}^{-1}, \dots, w_{(n)}^{-1}) = (\gamma^0, \dots, \gamma^n)$. For $k \in [1, n]$, let α_k be the simple root such that $s_{|i_k|} = s_{\alpha_k}$. For $z \in \mathbb{C}$, let

$$g_k(z) = \begin{cases} \overline{\gamma_{\mathbf{v}}^k} x_{\alpha_k}(z) \overline{\gamma_{\mathbf{v}}^k}^{-1}, & \epsilon_k = 1, k \notin J(\gamma) \\ \overline{\gamma_{\mathbf{u}}^k} x_{-\alpha_k}(z) \overline{\gamma_{\mathbf{u}}^k}^{-1}, & \epsilon_k = -1, k \notin J(\gamma) \\ e, & k \in J(\gamma), \end{cases}$$

recall (32), and $g_{\mathbf{w}}(z_1, \dots, z_n) = g_1(z_1) \cdots g_n(z_n)$, where

$$(37) \quad z_k \in \begin{cases} \{0\}, & k \in J(\gamma), \\ \mathbb{C}^*, & k \notin I(\gamma), \\ \mathbb{C}, & k \in I(\gamma) \setminus J(\gamma). \end{cases}$$

Then $(z_1, \dots, z_n) \mapsto g_{\mathbf{w}}(z_1, \dots, z_n) \cdot (\gamma_{\mathbf{u}} \cdot B, \gamma_{\mathbf{v}} \cdot B_-)$ is an isomorphism between $(\mathbb{C}^*)^{|J_{\mathbf{w}}^0|} \times \mathbb{C}^{|J_{\mathbf{w}}^-|}$ and $\mathcal{P}_{\mathbf{u},\mathbf{v}}^{\mathbf{w}}$.

Lemma 3.30. *Let $z = (z_1, \dots, z_n)$ satisfying (37). There exist signs $\varepsilon_1, \dots, \varepsilon_n \in \{\pm 1\}$ such that*

$$\theta_n(u_\gamma(z)) = g_{\mathbf{w}}(\varepsilon_1 z_1, \dots, \varepsilon_n z_n) \cdot (\gamma_{\mathbf{u}} \cdot B, \gamma_{\mathbf{v}} \cdot B_-).$$

Proof. By Lemma 3.7, one has

$$\begin{aligned} \theta_n(u_\gamma(z)) &= (p_{\gamma,1}(z_1) \cdots p_{\gamma,n}(z_n) \cdot B, q_{\gamma,1}(z_1) \cdots q_{\gamma,n}(z_n) \cdot B_-) \\ &= p_{\gamma,1}(z_1) \cdots p_{\gamma,n}(z_n) \overline{\gamma_{\mathbf{u}}^j}^{-1} (\gamma_{\mathbf{u}} \cdot B, \gamma_{\mathbf{v}} \cdot B_-) \\ &= q_{\gamma,1}(z_1) \cdots q_{\gamma,n}(z_n) \overline{\gamma_{\mathbf{v}}^j}^{-1} (\gamma_{\mathbf{u}} \cdot B, \gamma_{\mathbf{v}} \cdot B_-). \end{aligned}$$

Let $H^{(2)} = \{h \in H \mid h^2 = 1\}$. It is enough to prove the existence of a $h_j \in H^{(2)}$ and $\varepsilon_1, \dots, \varepsilon_j \in \{\pm 1\}$ such that

$$(38) \quad g_1(z_1) \cdots g_j(z_j) = p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j}(\varepsilon_j z_j) \overline{\gamma_{\mathbf{u}}^j}^{-1} h_j,$$

for all $j \in [1, n]$. If $j = 1$, (38) is easily verified. So assume $j \geq 2$, and that (38) holds for $j - 1$. Suppose first that $\epsilon_j = -1$. If $j \in J(\gamma)$, then $g_j(z_j) = e$ and $p_{\gamma,j}(z_j) = s_{|i_j|}$. So

$$\begin{aligned} g_1(z_1) \cdots g_j(z_j) &= g_1(z_1) \cdots g_{j-1}(z_{j-1}) = p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j-1}(\varepsilon_{j-1} z_{j-1}) \overline{\gamma_{\mathbf{u}}^{j-1}}^{-1} h_{j-1} \\ &= p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j-1}(\varepsilon_{j-1} z_{j-1}) \overline{s_{|i_j|} \gamma_{\mathbf{u}}^j}^{-1} h_j, \end{aligned}$$

for some $h_j \in H^{(2)}$, so we are done. If $j \notin J(\gamma)$, then

$$\begin{aligned} g_1(z_1) \cdots g_j(z_j) &= p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j-1}(\varepsilon_{j-1} z_{j-1}) \overline{\gamma_{\mathbf{u}}^{j-1}}^{-1} h_{j-1} \overline{\gamma_{\mathbf{u}}^j} x_{-\alpha_j}(z_j) \overline{\gamma_{\mathbf{u}}^j}^{-1} \\ &= p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j-1}(\varepsilon_{j-1} z_{j-1}) \bar{\gamma}_j x_{-\alpha_j}(\varepsilon'_j z_j) \overline{\gamma_{\mathbf{u}}^j}^{-1} h' \\ &= p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j-1}(\varepsilon_{j-1} z_{j-1}) x_{-\gamma_j \alpha_j}(\varepsilon_j z_j) \bar{\gamma}_j \overline{\gamma_{\mathbf{u}}^j}^{-1} h_j \\ &= p_{\gamma,1}(\varepsilon_1 z_1) \cdots p_{\gamma,j}(\varepsilon_j z_j) \overline{\gamma_{\mathbf{u}}^j}^{-1} h_j, \end{aligned}$$

for some $h', h_j \in H^{(2)}$, and $\varepsilon', \varepsilon_j \in \{\pm 1\}$. Thus we are done. Suppose now that $\varepsilon_j = 1$. Observe that for any $k \in [1, n]$,

$$p_{\gamma,1}(z_1) \cdots p_{\gamma,k}(z_k) \overline{\gamma_{\mathbf{u}}^k}^{-1} = q_{\gamma,1}(z_1) \cdots q_{\gamma,k}(z_k) \overline{\gamma_{\mathbf{v}}^k}^{-1} h,$$

for some $h \in H^{(2)}$. Hence one can proceed similarly by proving

$$g_1(z_1) \cdots g_j(z_j) = q_{\gamma,1}(\varepsilon_1 z_1) \cdots q_{\gamma,j}(\varepsilon_j z_j) \overline{\gamma_{\mathbf{u}}^j}^{-1} h_j, \quad j \in [1, n],$$

for some $h_j \in H^{(2)}$ and $\varepsilon_1, \dots, \varepsilon_j \in \{\pm 1\}$.

Q.E.D.

We conclude that our parametrization of C^γ recovers the one of $\mathcal{P}_{\mathbf{u},\mathbf{v}}^{\mathbf{w}}$ when both $\mathbf{u}$ and $\mathbf{v}$ are reduced.

4. REGULAR FUNCTIONS ON $\mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})}$ DEFINED BY MINORS

Let $(\mathbf{u}, \mathbf{v}, \sigma)$ be as in Definition 3.1. We introduce, for each distinguished σ -shuffled subexpression γ , a family of regular functions on $\mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})}$, and describe $\mathcal{O}^{\sigma(\mathbf{u},\mathbf{v})} \cap C^\gamma$ using these functions. When γ is positive, we relate this family of functions to the coordinates on $\mathcal{O}^\gamma$.

4.1. Generalised minors. We recall the notion of generalised minors, and some of their basic properties that will be needed. See [8] for further details. We assume from now on that G is simply connected.

The set N_-HN is Zariski open in G . Moreover, recall that any element $x \in N_-HN$ is uniquely written as $x = [x]_-[x]_0[x]_+$, with $[x]_- \in N_-$, $[x]_0 \in H$, $[x]_+ \in N$. Let $\alpha \in \Gamma$. For $x \in N_-HN$, define

$$\Delta_{\lambda_\alpha}(x) = [x]_0^{\lambda_\alpha}.$$

The functions Δ_{λ_α} extend to regular functions on G . For $G = SL(n, \mathbb{C})$, these are simply the principal $i \times i$ minor of a matrix x , $i \in [1, n-1]$.

An equivalent definition of Δ_{λ_α} is the following. Let V_α be the irreducible representation of G of highest weight λ_α , and let $v_\alpha \in V_\alpha$ be a highest weight vector. For any $a \in V$, let $\xi_\alpha(a)$ be the coefficient of v_α in the expansion of a in any basis consisting of v_α and weight vectors. Then

$$\Delta_{\lambda_\alpha}(x) = \xi_\alpha(x \cdot v_\alpha), \quad x \in G.$$

For any $u, v \in W$, the corresponding *generalised minor* is the regular function on G given by

$$\Delta_{u\lambda_\alpha, v\lambda_\alpha}(x) = \Delta_{\lambda_\alpha}(\bar{u}^{-1}x\bar{v}).$$

It can be checked that $\Delta_{u\lambda_\alpha, v\lambda_\alpha}$ depends only on $u\lambda_\alpha$ and $v\lambda_\alpha$, thus making the notation consistent.

If $G = SL(n, \mathbb{C})$, the Weyl group of G is naturally identified with the symmetric group S_n . Then $\Delta_{u\lambda_\alpha, v\lambda_\alpha}(x)$ is the minor of x formed by lines $u([1, i])$ and columns $v([1, i])$, where α is the i 'th simple root.

Lemma 4.1. *[8, Proposition 2.4] Let $\alpha \in \Gamma$. The zero locus of Δ_{λ_α} is precisely $\overline{B_{-s_\alpha B}}$.*

Lemma 4.2. *Let $\alpha \in \Gamma$. Then for any $z \in \mathbb{C}$ and any $g \in G$, one has*

$$\begin{aligned}\Delta_{\lambda_\alpha}(x_\alpha(z)g) &= \Delta_{\lambda_\alpha}(g) + z\Delta_{s_\alpha\lambda_\alpha, \lambda_\alpha}(g) \\ \Delta_{\lambda_\alpha}(gx_{-\alpha}(z)) &= \Delta_{\lambda_\alpha}(g) + z\Delta_{\lambda_\alpha, s_\alpha\lambda_\alpha}(g).\end{aligned}$$

Proof. The two formulas can be proved similarly, so we will only prove the first one. Let V be the irreducible representation of G with highest weight λ_α , and let $v \in V$ be a highest weight vector. For any $g \in G$, one has

$$g \cdot v = \sum_{w\lambda_\alpha | w \in W} \Delta_{w\lambda_\alpha, \lambda_\alpha}(g) \bar{w} \cdot v + R,$$

where R lies in the direct sum of non-extremal weight spaces. Thus for $z \in \mathbb{C}$,

$$x_\alpha(z)g \cdot v = \sum_{w\lambda_\alpha | w \in W} \Delta_{w\lambda_\alpha, \lambda_\alpha}(g) x_\alpha(z) \bar{w} \cdot v + x_\alpha(z)R.$$

For any $a \in V$, let $\xi(a)$ be the coefficient of v in the expansion of a in any basis consisting of v and weight vectors. Then

$$\begin{aligned}\Delta_{\lambda_\alpha}(x_\alpha(z)g) &= \xi(x_\alpha(z)g \cdot v) = \sum_{w\lambda_\alpha | w \in W} \Delta_{w\lambda_\alpha, \lambda_\alpha}(g) \xi(x_\alpha(z) \bar{w} \cdot v) + \xi(x_\alpha(z)R) \\ &= \Delta_{\lambda_\alpha}(g) + z\Delta_{s_\alpha\lambda_\alpha, \lambda_\alpha}(g).\end{aligned}$$

Q.E.D.

4.2. Families of regular functions on $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ defined by minors. Let $(\mathbf{u}, \mathbf{v}, \sigma)$ be as in Definition 3.1 and γ be a σ -shuffled subexpression of $(\mathbf{u}, \mathbf{v})$. Recall that γ is *distinguished* if $C^\gamma \cap \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})} \neq \emptyset$. This is equivalent, by Lemma 3.23, to $J(\gamma) \subset I(\gamma)$, recall Definitions 3.6 and 3.22. If γ is distinguished, we denote

$$(39) \quad K(\gamma) = I(\gamma) \setminus J(\gamma).$$

Let $(z_1, \dots, z_n)$ be the coordinates on $\mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})}$ defined in (29). To lighten the notation we write

$$\mathcal{O} = \mathcal{O}^{\sigma(\mathbf{u}, \mathbf{v})},$$

and for $j \in [1, n]$,

$$p_j(z_j) = p_{\sigma(\mathbf{u}, \mathbf{v}), j}(z_j), \quad q_j(z_j) = q_{\sigma(\mathbf{u}, \mathbf{v}), j}(z_j), \quad u(z) = u_{\sigma(\mathbf{u}, \mathbf{v})}(z).$$

In particular, one has

$$p_j(z_j) \begin{cases} = x_{\alpha_j}(z_j) \bar{\delta}_j, & \text{if } \epsilon_j = -1 \\ \in N, & \text{if } \epsilon_j = 1, \end{cases} \quad q_j(z_j) \begin{cases} \in N_-, & \text{if } \epsilon_j = -1 \\ = x_{-\alpha_j}(z_j) \bar{\delta}_j^{-1}, & \text{if } \epsilon_j = 1. \end{cases}$$

Definition 4.3. For $j \in [1, n]$, let

$$g_j(z_1, \dots, z_j) = \begin{cases} (q_1(z_1) \cdots q_{j-1}(z_{j-1}))^{-1} p_1(z_1) \cdots p_j(z_j) \in G, & \text{if } \epsilon_j = -1 \\ (q_1(z_1) \cdots q_j(z_j))^{-1} p_1(z_1) \cdots p_{j-1}(z_{j-1}) \in G, & \text{if } \epsilon_j = 1. \end{cases}$$

For every $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^d$, and $j \in [1, n]$, define the regular function on $\mathcal{O}$,

$$\psi_{\gamma, j}(z_1, \dots, z_j) = \begin{cases} \Delta_{\lambda_j} \left(\overline{\gamma^{j-1}}^{-1} g_j(z_1, \dots, z_j) \right), & \text{if } \epsilon_j = -1 \\ \Delta_{\lambda_j} \left(g_j(z_1, \dots, z_j) \overline{\gamma^{j-1}}^{-1} \right), & \text{if } \epsilon_j = 1, \end{cases}$$

where $\lambda_j = \lambda_{\alpha_j}$ for $j \in [1, n]$.

Proposition 4.4. Let $u(z) \in \mathcal{O}$ and $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^d$. Then $u(z) \in C^\gamma$ if and only if

$$(40) \quad \begin{cases} \psi_{\gamma, j}(z_1, \dots, z_j) = 0, & \text{if } j \in J(\gamma) \\ \psi_{\gamma, j}(z_1, \dots, z_j) \neq 0, & \text{if } j \notin I(\gamma). \end{cases}$$

Proof. By definition of C^γ (recall Definition 3.4), $u(z) \in C^\gamma$ if and only if

$$(q_1(z_1) \cdots q_j(z_j))^{-1} p_1(z_1) \cdots p_j(z_j) \in B_- \gamma^j B, \quad j \in [1, n],$$

which is equivalent to

$$(41) \quad g_j(z_1, \dots, z_j) \in B_- \gamma^j B,$$

for any $j \in [1, n]$. Suppose that $u(z) \in C^\gamma$, and let $j \notin I(\gamma)$. Then $\gamma^j = \gamma^{j-1}$, and so

$$\begin{cases} \overline{\gamma^{j-1}}^{-1} g_j(z_1, \dots, z_j) \\ g_j(z_1, \dots, z_j) \overline{\gamma^{j-1}}^{-1} \end{cases} \in N_- H N.$$

Thus one has $\psi_{\gamma, j}(z_1, \dots, z_j) \neq 0$. Let now $j \in J(\gamma)$. Since γ is distinguished, this implies that $\gamma_j = \delta_j$, thus $(\gamma^{j-1})^{-\epsilon_j} \alpha_j > 0$. Assume that $\epsilon_j = -1$, the case $\epsilon_j = 1$ being similar. One has

$$g_j(z_1, \dots, z_j) \in N_- \gamma^j B = (N_- \cap \gamma^j N_- (\gamma^j)^{-1}) \gamma^{j-1} \delta_j B.$$

It follows that

$$\overline{\gamma^{j-1}}^{-1} g_j(z_1, \dots, z_j) \in ((\gamma^{j-1})^{-1} N_- \gamma^{j-1} \cap \delta_j N_- \delta_j) \delta_j B \subset N_- \delta_j B.$$

Thus by Lemma 4.1, one must have $\psi_{\gamma, j}(z_1, \dots, z_j) = 0$.

Conversely, suppose that $u(z) \in \mathcal{O}$ satisfies (40). We need to show that (41) holds for every $j \in [1, n]$. Let $j = 1$. Since $\gamma^0 = e$, one has

$$\psi_{\gamma, 1}(z_1) = \Delta_{\lambda_1}(x_{\alpha_1}(z_1) \bar{\delta}_1) = z_1.$$

If $\gamma_1 = e$, one has $z_1 \neq 0$, hence $g_1(z_1) \in B_- B = B_- \gamma^1 B$. If $\gamma_1 = \delta_1$, then $1 \in J(\gamma)$, so $z_1 = 0$. Thus $g_1(z_1) \in B_- \delta_1 B = B_- \gamma^1 B$. So assume that $j \geq 2$, and that (41) holds for $j-1$. Once again both cases $\epsilon_j = -1$ and $\epsilon_j = 1$ are similar, so we will assume that $\epsilon_j = -1$.

Case $j \notin I(\gamma)$:

Since γ is distinguished, $j \notin J(\gamma)$, so $\gamma^j \alpha_j > 0$, which means $\gamma^j \delta_j > \gamma^j$. Then

$$\begin{aligned} g_j(z_1, \dots, z_j) &= (q_1(z_1) \cdots q_{j-1}(z_{j-1}))^{-1} p_1(z_1) \cdots p_{j-1}(z_{j-1}) p_j(z_j) \\ &\in B_- \gamma^{j-1} B \delta_j B = B_- \gamma^{j-1} B \cup B_- \gamma^{j-1} \delta_j B. \end{aligned}$$

If $g_j(z_1, \dots, z_j) \in B_- \gamma^{j-1} \delta_j B$, then $\overline{\gamma^{j-1}}^{-1} g_j(z_1, \dots, z_j) \in B_- \delta_j B$. This implies, by Lemma 4.1, that $\psi_{\gamma,j}(z_1, \dots, z_j) = 0$, which is a contradiction. Thus

$$g_j(z_1, \dots, z_j) \in B_- \gamma^{j-1} B = B_- \gamma^j B.$$

Case $j \in J(\gamma)$:

Since γ is distinguished, $\gamma_j = \delta_j$. Thus $\gamma^{j-1} \delta_j > \gamma^{j-1}$. One gets

$$\begin{aligned} g_j(z_1, \dots, z_j) &= (q_1(z_1) \cdots q_{j-1}(z_{j-1}))^{-1} p_1(z_1) \cdots p_{j-1}(z_{j-1}) p_j(z_j) \\ &\in B_- \gamma^{j-1} B \delta_j B = B_- \gamma^{j-1} B \cup B_- \gamma^{j-1} \delta_j B. \end{aligned}$$

If $g_j(z_1, \dots, z_j) \in B_- \gamma^{j-1} B$, then $\overline{\gamma^{j-1}}^{-1} g_j(z_1, \dots, z_j) \in B_- B$. This implies that $\psi_{\gamma,j}(z_1, \dots, z_j) \neq 0$, which is a contradiction. Thus

$$g_j(z_1, \dots, z_j) \in B_- \gamma^{j-1} \delta_j B = B_- \gamma^j B.$$

Case $j \in K(\gamma)$:

Since γ is distinguished, $\gamma^j \alpha_j > 0$, so $\gamma^{j-1} \delta_j < \gamma^{j-1}$. Hence

$$\begin{aligned} g_j(z_1, \dots, z_j) &= (q_1(z_1) \cdots q_{j-1}(z_{j-1}))^{-1} p_1(z_1) \cdots p_{j-1}(z_{j-1}) p_j(z_j) \\ &\in B_- \gamma^{j-1} B \delta_j B = B_- \gamma^{j-1} \delta_j B = B_- \gamma^j B. \end{aligned}$$

Thus (41) holds for all $j \in [1, n]$.

Q.E.D.

Lemma 4.5. *Let $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^d$ and $j \in [1, n]$. There exist polynomials $L_j, M_j \in \mathbb{C}[z_1, \dots, z_{j-1}]$ such that*

$$\psi_{\gamma,j}(z_1, \dots, z_j) = L_j + z_j M_j.$$

Moreover, M_j vanishes nowhere on C^γ .

Proof. If $j = 1$, then $\psi_{\gamma,1}(z_1) = z_1$. So assume now that $j \geq 2$, and that $\epsilon_j = -1$, the other case being similar. One has $g_j(z_1, \dots, z_j) = x_{j-1} x_{\alpha_j}(z_j) \bar{\delta}_j$, where

$$x_{j-1} = (q_1(z_1) \cdots q_{j-1}(z_{j-1}))^{-1} p_1(z_1) \cdots p_{j-1}(z_{j-1})$$

depends only on $z_1, \dots, z_{j-1}$. Then, using Lemma 4.2,

$$\begin{aligned} \psi_{\gamma,j}(z_1, \dots, z_j) &= \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} x_{j-1} \bar{\delta}_j x_{-\alpha_j}(-z_j)) \\ &= \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} x_{j-1} \bar{\delta}_j) + z_j \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} x_{j-1}). \end{aligned}$$

Thus one lets

$$L_j = \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} x_{j-1} \bar{\delta}_j) \quad \text{and} \quad M_j = \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} x_{j-1}).$$

Suppose now that $u(z) \in C^\gamma$. Then in particular, $x_{j-1} \in B_- \gamma^{j-1} B$, so that $\overline{\gamma^{j-1}}^{-1} x_{j-1} \in B_- B$. Hence $M_j(z_1, \dots, z_{j-1}) \neq 0$.

Q.E.D.

Corollary 4.6. *The map*

$$\mathcal{O} \cap C^\gamma \rightarrow \mathbb{C}^{|K(\gamma)|} \times (\mathbb{C}^*)^{|I(\gamma)^c|}, \quad u(z) \mapsto ((\psi_{\gamma,j})_{j \in K(\gamma)}, (\psi_{\gamma,j})_{j \notin I(\gamma)}),$$

where $I(\gamma)^c = [1, n] \setminus I(\gamma)$, is an isomorphism.

Proof. Indeed by Lemma 4.5, for any value

$$((x_j)_{j \in K(\gamma)}, (x_j)_{j \notin I(\gamma)}) \in \mathbb{C}^{|K(\gamma)|} \times (\mathbb{C}^*)^{|I(\gamma)^c|},$$

one can uniquely solve $\psi_{\gamma,j}(z_1, \dots, z_j) = x_j$.

Q.E.D.

Fix now a positive σ -shuffled subexpression γ . Let $\xi = (\xi_1, \dots, \xi_n)$ be the coordinates on $\mathcal{O}^\gamma$ defined in (29). By Proposition 3.25, $u_\gamma(\xi) \in \mathcal{O} \cap C^\gamma$ if and only if $\xi_j = 0$ when $j \in J(\gamma)$, and $\xi_j \neq 0$ when $j \notin J(\gamma)$. Hence

$$\mathcal{O} \cap C^\gamma \rightarrow (\mathbb{C}^*)^{|J(\gamma)^c|}, \quad u_\gamma(\xi) \mapsto (\xi_j)_{j \notin J(\gamma)},$$

is an isomorphism. On the other hand, by Corollary 4.6,

$$\mathcal{O} \cap C^\gamma \rightarrow (\mathbb{C}^*)^{|J(\gamma)^c|}, \quad u(z) \mapsto (\psi_{\gamma,j})_{j \notin J(\gamma)}$$

is another set of coordinates on $\mathcal{O} \cap C^\gamma$. We show now that these two systems of coordinates are related by a triangular matrix of monomials.

Definition 4.7. For any $1 \leq k < j \leq n$, define

$$\mathbf{u}_{(k,j]} = \prod_{k < i \leq j, \epsilon_i = -1} \delta_i, \quad \mathbf{v}_{(k,j]} = \prod_{k < i \leq j, \epsilon_i = 1} \delta_i,$$

where the index in both products is increasing, and it is understood that $\mathbf{u}_{(k,j]}$ or $\mathbf{v}_{(k,j]}$ = e , if the set over which the product is defined is empty.

Theorem 4.8. *Let $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$, and $j \notin J(\gamma)$. Then over $\mathcal{O} \cap C^\gamma$, one has*

$$\psi_{\gamma,j} = \prod_{1 \leq k \leq j, k \notin J(\gamma)} \xi_k^{-m_{j,k}},$$

where

$$m_{j,k} = \begin{cases} (\mathbf{v}_{(k,j]} \gamma^j \lambda_j, \check{\alpha}_k), & \text{if } \epsilon_j = -1, \epsilon_k = 1 \\ (\mathbf{u}_{(k,j]} \lambda_j, \check{\alpha}_k), & \text{if } \epsilon_j = -1, \epsilon_k = -1 \\ (\mathbf{v}_{(k,j]} \lambda_j, \check{\alpha}_k), & \text{if } \epsilon_j = 1, \epsilon_k = 1 \\ (\mathbf{u}_{(k,j]} (\gamma^j)^{-1} \lambda_j, \check{\alpha}_k), & \text{if } \epsilon_j = 1, \epsilon_k = -1. \end{cases}$$

Proof. Let $z = (z_1, \dots, z_n), \xi = (\xi_1, \dots, \xi_n) \in \mathbb{C}^n$ and suppose that $u(z) = u_\gamma(\xi)$. Then there exist elements $b_j \in B$, $b_{-j} \in B_-$, for $j \in [1, n]$, such that

$$\begin{cases} p_{\gamma,1}(\xi_1) &= p_1(z_1)b_1 \\ b_1 p_{\gamma,2}(\xi_2) &= p_2(z_2)b_2 \\ \vdots & \\ b_{n-1} p_{\gamma,n}(\xi_n) &= p_n(z_n)b_n, \end{cases} \quad \begin{cases} q_{\gamma,1}(\xi_1) &= q_1(z_1)b_{-1} \\ b_{-1} q_{\gamma,2}(\xi_2) &= q_2(z_2)b_{-2} \\ \vdots & \\ b_{-(n-1)} q_{\gamma,n}(\xi_n) &= q_n(z_n)b_{-n}. \end{cases}$$

Write

$$b_j = t_j a_j, \quad b_{-j} = t_{-j} a_{-j},$$

with $t_j, t_{-j} \in H$, $a_j \in N$, and $a_{-j} \in N_-$, $j \in [1, n]$. Let now $j \notin J(\gamma)$ and suppose first that $\epsilon_j = -1$. If $u(z) = u_\gamma(\xi) \in C^\gamma$, then by Lemma 3.7, and Remark 3.16,

$$\begin{aligned} g_j(z_1, \dots, z_j) &= q_j(z_j)(q_1(z_1) \cdots q_j(z_j))^{-1} p_1(z_1) \cdots p_j(z_j) \\ &= q_j(z_j) \left(q_{\gamma,1}(\xi_1) \cdots q_{\gamma,j}(\xi_j) b_{-j}^{-1} \right)^{-1} p_{\gamma,1}(\xi_1) \cdots p_{\gamma,n}(\xi_n) b_j^{-1} \\ &= q_j(z_j) b_{-j} \overline{\gamma^j} b_j^{-1}. \end{aligned}$$

Similarly, if $\epsilon_j = 1$, then

$$g_j(z_1, \dots, z_j) = b_{-j} \overline{\gamma^j} b_j^{-1} p_j(z_j)^{-1}.$$

Recall that since γ is positive, one has $\gamma^j = \gamma^{j-1}$. So if $\epsilon_j = -1$,

$$\begin{aligned} \psi_{\gamma,j} &= \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} g_j(z_1, \dots, z_j)) = \Delta_{\lambda_j}(\overline{\gamma^{j-1}}^{-1} q_j(z_j) b_{-j} \overline{\gamma^j} b_j^{-1}) \\ &= \Delta_{\lambda_j}(\overline{\gamma^j}^{-1} q_j(z_j) b_{-j} \overline{\gamma^j} b_j^{-1}) \\ (42) \quad &= t_{-j}^{\gamma^j \lambda_j} t_j^{-\lambda_j}. \end{aligned}$$

Similarly, if $\epsilon_j = 1$, one gets

$$(43) \quad \psi_{\gamma,j} = t_{-j}^{\lambda_j} t_j^{-(\gamma^j)^{-1} \lambda_j}.$$

Suppose now that $\epsilon_1 = -1$. Since $q_{\gamma,1}(\xi_1)$ and $q_1(z_1)$ both lie in N_- , one has $t_{-1} = e$. If $\gamma_1 = \delta_1$, one has $1 \in J(\gamma)$, and $p_{\gamma,1}(\xi_1) = \bar{\delta}_1$. Thus $t_1 = e$. If $\gamma_1 = e$, then

$$p_{\gamma,1}(\xi_1) = x_{-\alpha_1}(\xi_1) = x_{\alpha_1}(\xi_1^{-1}) \bar{\delta}_1 \check{\alpha}_1(\xi_1) x_{\alpha_1}(\xi_1^{-1}),$$

and so $t_1 = \check{\alpha}_1(\xi_1)$. Similarly, if $\epsilon_1 = 1$, then $t_1 = e$, and

$$t_{-1} = \begin{cases} e, & \text{if } \gamma_1 = \delta_1 \\ \check{\alpha}_1(\xi_1^{-1}), & \text{if } \gamma_1 = e. \end{cases}$$

We claim now that for all $j \in [2, n]$ and $u(z) = u_\gamma(\xi) \in \mathcal{O} \cap C^\gamma$,

$$(44) \quad t_j = \begin{cases} t_{j-1}^{\delta_j}, & \text{if } \epsilon_j = -1, j \in J(\gamma) \\ \check{\alpha}_j(\xi_j) t_{j-1}^{\delta_j}, & \text{if } \epsilon_j = -1, j \notin J(\gamma) \\ t_{j-1}, & \text{if } \epsilon_j = 1. \end{cases}$$

and

$$(45) \quad t_{-j} = \begin{cases} t_{-(j-1)}^{\delta_j}, & \text{if } \epsilon_j = 1, j \in J(\gamma) \\ \check{\alpha}_j(\xi_j^{-1}) t_{-(j-1)}^{\delta_j}, & \text{if } \epsilon_j = 1, j \notin J(\gamma) \\ t_{-(j-1)}, & \text{if } \epsilon_j = -1. \end{cases}$$

Both (44) and (45) arise from similar calculations, so we will concentrate on (44). Suppose that $\epsilon_j = -1$ and $j \in J(\gamma)$. Then $p_{\gamma,j}(\xi_j) = \bar{\delta}_j$, so

$$b_{j-1} p_{\gamma,j}(\xi_j) = t_{j-1} a_{j-1} \bar{\delta}_j = x_{\alpha_j}(z_j) \bar{\delta}_j t_{j-1}^{\delta_j} a_j.$$

Hence $t_j = t_{j-1}^{\delta_j}$. Suppose now that $j \notin J(\gamma)$. Write $a_{j-1} = x_{\alpha_j}(\xi'_j)a'_{j-1}$, with $\xi'_j \in \mathbb{C}$ and $a'_{j-1} \in N \cap \delta_j N \delta_j$. Then

$$\begin{aligned} b_{j-1}p_{\gamma,j}(\xi_j) &= t_{j-1}a_{j-1}x_{-\alpha_j}(\xi_j) \\ &= t_{j-1}x_{\alpha_j}(\xi'_j)x_{-\alpha_j}(\xi_j) (x_{-\alpha_j}(-\xi_j)a'_{j-1}x_{-\alpha_j}(\xi_j)) \\ &= t_{j-1}x_{\alpha_j}((1 + \xi_j\xi'_j)/\xi_j)\bar{\delta}_j\check{\alpha}_j(\xi_j)x_{\alpha}(\xi_j^{-1}) (x_{-\alpha_j}(-\xi_j)a'_{j-1}x_{-\alpha_j}(\xi_j)) \\ &= x_{\alpha_j}(z_j)\bar{\delta}_jt_{j-1}^{\delta_j}\check{\alpha}_j(\xi_j)a_j, \end{aligned}$$

so $t_j = \check{\alpha}_j(\xi_j)t_{j-1}^{\delta_j}$. Suppose now that $\epsilon_j = 1$. Then since $p_{\gamma,j}(\xi_j)$ and $p_j(z_j)$ lie in N , one has $t_j = t_{j-1}$.

Now (44) and (45) give

$$\begin{aligned} t_j &= \prod_{k=1, \epsilon_k=-1, k \notin J(\gamma)}^j \check{\alpha}_k(\xi_k)^{\mathbf{u}_{(k,j]}} \\ t_{-j} &= \prod_{k=1, \epsilon_k=1, k \notin J(\gamma)}^j \check{\alpha}_k(\xi_k^{-1})^{\mathbf{v}_{(k,j]}}. \end{aligned}$$

Substituting in (42) and (43) yields the result.

Q.E.D.

Definition 4.9. For any $\gamma \in \Upsilon_{\mathbf{u}, \mathbf{v}, \sigma}^+$, let $M_\gamma = (m_{j,k})_{j,k \notin J(\gamma)}$ be the $|J(\gamma)^c| \times |J(\gamma)^c|$ lower triangular matrix where $m_{j,k}$ is as in Theorem 4.8, for $k \leq j$, $j, k \notin J(\gamma)$.

Notice that M_γ has diagonal entries 1. Informally, we write Theorem 4.8 as

$$\psi_\gamma = \xi^{M_\gamma}.$$

4.3. On the inverse of M_γ . Let L_γ be the inverse of M_γ . The relation

$$\xi = \psi_\gamma^{L_\gamma}$$

can be thought as an analogous of the inverse factorization problem of Fomin and Zelevinsky, see [8]. That is, the local coordinates $(\xi_k)_{k \notin J(\gamma)}$ on $\mathcal{O} \cap C^\gamma$ are being written in terms of regular functions $(\psi_{\gamma,j})_{j \notin J(\gamma)} \in \mathbb{C}[\mathcal{O}]$.

The following Lemma 4.10 provides an inductive formula to express the entries of L_γ .

Lemma 4.10. *Let V be a vector space over a field $\mathbb{K}$, $v_1, \dots, v_n \in V$, and $\varphi_1, \dots, \varphi_n \in V^*$. For $1 \leq k < j \leq n$, Suppose given operators $M_{jk} \in \text{End}(V)$ and let $(m_{jk})_{j,k=1, \dots, n} \in \mathfrak{gl}(n, \mathbb{K})$ be the lower triangular matrix with diagonal entries 1 and*

$$m_{jk} = (M_{jk}v_j, \varphi_k), \quad 1 \leq k < j \leq n.$$

Let $(l_{jk})_{j,k=1, \dots, n}$ be the inverse of $(m_{jk})_{j,k=1, \dots, n}$. Then there exist operators $L_{ji} \in \text{End}(V)$ such that

$$l_{jk} = (L_{jk}v_j, \varphi_k), \quad 1 \leq k < j \leq n.$$

Proof. For $j - k = 1$, one lets $L_{jk} = -M_{jk}$. Thus assume $j - k > 1$ and the operators $L_{j'k'}$ defined for $j' - k' < j - k$. Then

$$\begin{aligned} l_{jk} &= - \left(\sum_{i=k+1}^{j-1} l_{ji} m_{ik} \right) - m_{jk} \\ &= - \left(\sum_{i=k+1}^{j-1} ((L_{ji} v_j, \varphi_i) M_{ik} v_i, \varphi_k) \right) - m_{jk} \\ &= \left(- \left(\sum_{i=k+1}^{j-1} (L_{ji} v_j, \varphi_i) M_{ik} v_i \right) - M_{jk} v_j, \varphi_k \right). \end{aligned}$$

Thus set

$$L_{jk} x = - \left(\sum_{i=k+1}^{j-1} (L_{ji} x, \varphi_i) M_{ik} v_i \right) - M_{jk} x, \quad x \in V.$$

Q.E.D.

By Lemma 4.10, one can inductively define operators $L_{jk} \in \text{End}(\mathfrak{h}^*)$, for $k < j$, $j, k \notin J(\gamma)$, such that

$$(46) \quad (L_\gamma)_{j,k} = (L_{jk} \lambda_j, \check{\alpha}_k).$$

In the case when $\mathbf{v} = \emptyset$, that is one only has a single Bott Samelson variety $Z_{\mathbf{u}}$, the operators in (46) can be expressed in simple way.

Definition 4.11. For any $\alpha \in \Gamma$, define $r_\alpha \in \text{End}(\mathfrak{h}^*)$ by

$$r_\alpha x = x - (x, \check{\alpha}) \lambda_\alpha, \quad x \in \mathfrak{h}^*.$$

Proposition 4.12. [11] Suppose that $\mathbf{v} = \emptyset$. For $i \in [1, n]$, let $r_i = r_{\alpha_i}$ and

$$\tilde{\gamma}_i = \begin{cases} s_i, & \text{if } i \in J(\gamma) \\ r_i, & \text{if } i \notin J(\gamma). \end{cases}$$

For $k < j$ and $k, j \notin J(\gamma)$, one has

$$(L_\gamma)_{k,j} = -(\tilde{\gamma}_{k+1} \cdots \tilde{\gamma}_{j-1} s_j \lambda_j, \check{\alpha}_k).$$

REFERENCES

- [1] Brion M., *Lectures on the geometry of flag varieties*, Topics in cohomological studies of algebraic varieties, 33–85, Trends Math., Birkhäuser, Basel, 2005.
- [2] Brion M., Kumar S., *Frobenius splitting methods in geometry and representation theory*, Progress in Mathematics, 231. Birkhäuser Boston, Inc., Boston, MA, 2005. x+250 pp. ISBN: 0-8176-4191-2.
- [3] Bott R., Samelson H., *Application of the theory of Morse to symmetric spaces*, Amer. J. Math. 80 1958 964–1029.
- [4] Demazure M., *Désingularisation des variétés de Schubert généralisées*, Ann. Sci. École Norm. Sup. (4) 7 (1974), 53–88.
- [5] Deodhar V., *On some geometric aspects of Bruhat orderings. I. A finer decomposition of Bruhat cells*, Invent. Math. 79 (1985), no. 3, 499–511.
- [6] Dudas O., *Note on the Deodhar decomposition of a double Schubert cell*, arXiv:0807.2198 [math.AG].
- [7] Evens S., Lu J-H., *On the variety of Lagrangian subalgebras II*, Ann. Sci. École Norm. Sup. (4) 39 (2006), no. 2, 347–379.
- [8] Fomin S., Zelevinsky A., *Double Bruhat cells and Total Positivity*, J. Amer. Math. Soc. 12 (1999), no. 2, 335–380.
- [9] Härterich M., *The T -equivariant Cohomology of Bott-Samelson varieties*, arXiv:math/0412337v1 [math.AG].
- [10] He X-H, Lu J-H, *On intersections of certain partitions of a group compactification*, Int. Math. Res. Not. IMRN 2011, no. 11, 2534–2564.
- [11] Lu J-H, *On a Poisson structure on Bott-Samelson varieties*, in preparation.
- [12] Lu J-H, Mouquin V., *A Poisson structure on product of flag varieties*, in preparation.
- [13] Marsh R. J., Rietsch K., *Parametrizations of flag varieties*, Represent. Theory 8 (2004), 212–242.
- [14] Richardson R., *Intersections of double cosets in algebraic groups*, Indag. Math. (N.S.) 3 (1992), no. 1, 69–77.
- [15] Springer T.A., *Linear Algebraic Groups, Second edition*, Progress in Mathematics, 9. Birkhäuser Boston, Inc., Boston, MA, 2009. xvi+334 pp. ISBN: 978-0-8176-4839-8.
- [16] Webster B., Yakimov M., *A Deodhar-type stratification on the double flag variety*, Transform. Groups 12 (2007), no. 4, 769–785.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG
E-mail address: victor.mouquin@gmail.com