

HERMITE-HADAMARD AND SIMPSON-LIKE TYPE INEQUALITIES FOR DIFFERENTIABLE HARMONICALLY CONVEX FUNCTIONS

İMDAT İŞCAN

ABSTRACT. In this paper, a new identity for differentiable functions is derived. A consequence of the identity is that the author establishes some new general inequalities containing all of the Hermite-Hadamard and Simpson-like type for functions whose derivatives in absolute value at certain power are harmonically convex. Some applications to special means of real numbers are also given.

1. INTRODUCTION

Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The following inequality

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$$

holds. This double inequality is known in the literature as Hermite-Hadamard integral inequality for convex functions. Note that some of the classical inequalities for means can be derived from (1.1) for appropriate particular selections of the mapping f . Both inequalities hold in the reversed direction if f is concave.

Following inequality is well known in the literature as Simpson inequality:

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a four times continuously differentiable mapping on (a, b) and $\|f^{(4)}\|_{\infty} = \sup_{x \in (a, b)} |f^{(4)}(x)| < \infty$. Then the following inequality holds:*

$$\left| \frac{1}{3} \left[\frac{f(a) + f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{1}{2880} \|f^{(4)}\|_{\infty} (b-a)^4.$$

For some results which generalize, improve and extend the Hermite-Hadamard and Simpson inequalities, we refer the reader to the recent papers (see [1, 2, 3, 4, 6, 7, 8]).

In [5], the author introduced the concept of harmonically convex functions and established some results connected with the right-hand side of new inequalities similar to the inequality (1.1) for these classes of functions. Some applications to special means of positive real numbers are also given.

2000 *Mathematics Subject Classification.* Primary 26A51; Secondary 26D15.

Key words and phrases. Harmonically convex function, Hermite-Hadamard type inequality.

Definition 1. Let $I \subset \mathbb{R} \setminus \{0\}$ be a real interval. A function $f : I \rightarrow \mathbb{R}$ is said to be harmonically convex, if

$$(1.2) \quad f\left(\frac{xy}{tx + (1-t)y}\right) \leq tf(y) + (1-t)f(x)$$

for all $x, y \in I$ and $t \in [0, 1]$. If the inequality in (1.2) is reversed, then f is said to be harmonically concave.

The following result of the Hermite-Hadamard type holds.

Theorem 2. Let $f : I \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be a harmonically convex function and $a, b \in I$ with $a < b$. If $f \in L[a, b]$ then the following inequalities hold

$$(1.3) \quad f\left(\frac{2ab}{a+b}\right) \leq \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \leq \frac{f(a) + f(b)}{2}.$$

The above inequalities are sharp.

Some results connected with the right part of (1.3) was given in [5] as follows:

Theorem 3. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° , $a, b \in I$ with $a < b$, and $f' \in L[a, b]$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q \geq 1$, then

$$(1.4) \quad \left| \frac{f(a) + f(b)}{2} - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \lambda_1^{1-\frac{1}{q}} [\lambda_2 |f'(a)|^q + \lambda_3 |f'(b)|^q]^{\frac{1}{q}},$$

where

$$\begin{aligned} \lambda_1 &= \frac{1}{ab} - \frac{2}{(b-a)^2} \ln \left(\frac{(a+b)^2}{4ab} \right), \\ \lambda_2 &= \frac{-1}{b(b-a)} + \frac{3a+b}{(b-a)^3} \ln \left(\frac{(a+b)^2}{4ab} \right), \\ \lambda_3 &= \frac{1}{a(b-a)} - \frac{3b+a}{(b-a)^3} \ln \left(\frac{(a+b)^2}{4ab} \right) \\ &= \lambda_1 - \lambda_2. \end{aligned}$$

Theorem 4. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° , $a, b \in I$ with $a < b$, and $f' \in L[a, b]$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, then

$$(1.5) \quad \left| \frac{f(a) + f(b)}{2} - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \left(\frac{1}{p+1} \right)^{\frac{1}{p}} (\mu_1 |f'(a)|^q + \mu_2 |f'(b)|^q)^{\frac{1}{q}},$$

where

$$\begin{aligned}\mu_1 &= \frac{[a^{2-2q} + b^{1-2q} [(b-a)(1-2q) - a]]}{2(b-a)^2(1-q)(1-2q)}, \\ \mu_2 &= \frac{[b^{2-2q} - a^{1-2q} [(b-a)(1-2q) + b]]}{2(b-a)^2(1-q)(1-2q)}.\end{aligned}$$

In this paper, we shall give some general integral inequalities connected with the left and right parts of (1.3), as a result of this, we shall obtained some new midpoint, trapezoid and Simpson like-type inequalities for differentiable harmonically convex functions.

2. MAIN RESULTS

In order to prove our main results we need the following lemma:

Lemma 1. *Let $f : I \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I$ with $a < b$. If $f' \in L[a, b]$ then for $\lambda \in [0, 1]$ we have the equality*

$$\begin{aligned}& (1-\lambda)f\left(\frac{2ab}{a+b}\right) + \lambda\left(\frac{f(a)+f(b)}{2}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \\ &= \frac{ab(b-a)}{2} \left[\int_0^{1/2} \frac{\lambda-2t}{A_t^2} f'\left(\frac{ab}{A_t}\right) dt + \int_{1/2}^1 \frac{2-\lambda-2t}{A_t^2} f'\left(\frac{ab}{A_t}\right) dt \right],\end{aligned}$$

where $A_t = tb + (1-t)a$.

Proof. It suffices to note that

$$\begin{aligned}I_1 &= ab(b-a) \int_0^1 \frac{\lambda-2t}{A_t^2} f'\left(\frac{ab}{A_t}\right) dt \\ &= (2t-\lambda)f\left(\frac{ab}{A_t}\right) \Big|_0^{1/2} - 2 \int_0^{1/2} f\left(\frac{ab}{A_t}\right) dt \\ &= (1-\lambda)f\left(\frac{2ab}{a+b}\right) + \lambda f(b) - 2 \int_0^{1/2} f\left(\frac{ab}{A_t}\right) dt.\end{aligned}$$

Setting $x = \frac{ab}{A_t}$ and $dx = \frac{-ab(b-a)}{A_t^2} dt$, which gives

$$I_1 = (1-\lambda)f\left(\frac{2ab}{a+b}\right) + \lambda f(b) - \frac{2ab}{b-a} \int_{2ab/(a+b)}^b \frac{f(x)}{x^2} dx.$$

Similarly, we can show that

$$\begin{aligned} I_2 &= ab(b-a) \int_{1/2}^1 \frac{2-\lambda-2t}{A_t^2} f' \left(\frac{ab}{A_t} \right) dt \\ &= \lambda f(a) + (1-\lambda) f \left(\frac{2ab}{a+b} \right) - \frac{2ab}{b-a} \int_a^{2ab/(a+b)} \frac{f(x)}{x^2} dx. \end{aligned}$$

Thus,

$$\frac{I_1 + I_2}{2} = (1-\lambda) f \left(\frac{2ab}{a+b} \right) + \lambda \left(\frac{f(a) + f(b)}{2} \right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx$$

which is required. \square

Theorem 5. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° , $a, b \in I$ with $a < b$, and $f' \in L[a, b]$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q \geq 1$ and then we have the following inequality for $\lambda \in [0, 1]$

$$\begin{aligned} (2.1) \quad & \left| (1-\lambda) f \left(\frac{2ab}{a+b} \right) + \lambda \left(\frac{f(a) + f(b)}{2} \right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\ & \leq \frac{ab(b-a)}{2} \left\{ C_1^{1-\frac{1}{q}}(\lambda; a, b) [C_2(\lambda; a, b) |f'(a)|^q + C_3(\lambda; a, b) |f'(b)|^q]^{\frac{1}{q}} \right. \\ & \quad \left. + C_1^{1-\frac{1}{q}}(\lambda; b, a) [C_3(\lambda; b, a) |f'(a)|^q + C_2(\lambda; b, a) |f'(b)|^q]^{\frac{1}{q}} \right\}, \end{aligned}$$

where

$$\begin{aligned} C_1(\lambda; u, \vartheta) &= \frac{1}{(\vartheta - u)^2} \\ & \times \left[-4 + \frac{[\lambda(\vartheta - u) + 2u](3u + \vartheta)}{u(u + \vartheta)} + 2 \ln \left(\frac{2u(u + \vartheta)}{(2u + \lambda(\vartheta - u))^2} \right) \right], \end{aligned}$$

$$\begin{aligned} C_2(\lambda; u, \vartheta) &= \frac{1}{(\vartheta - u)^3} \\ & \times \left\{ [\lambda(\vartheta - u) + 4u] \ln \left(\frac{[\lambda(\vartheta - u) + 2u]^2}{2u(u + \vartheta)} \right) \right. \\ & \quad \left. - \frac{[\lambda(\vartheta - u) + 2u](5u + 3\vartheta)}{u + \vartheta} + 7u + \vartheta \right\}, \end{aligned}$$

and

$$C_3(\lambda; u, \vartheta) = C_1(\lambda; u, \vartheta) - C_2(\lambda; u, \vartheta), \quad u, \vartheta > 0.$$

Proof. Let $A_t = tb + (1-t)a$. From Lemma 1 and using the power mean inequality, we have

$$\begin{aligned}
& \left| (1-\lambda) f\left(\frac{2ab}{a+b}\right) + \lambda \left(\frac{f(a)+f(b)}{2}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\
& \leq \frac{ab(b-a)}{2} \left[\int_0^{1/2} \frac{|\lambda-2t|}{A_t^2} \left| f'\left(\frac{ab}{A_t}\right) \right| dt + \int_{1/2}^1 \frac{|2-\lambda-2t|}{A_t^2} \left| f'\left(\frac{ab}{A_t}\right) \right| dt \right] \\
& \leq \frac{ab(b-a)}{2} \left\{ \left(\int_0^{1/2} \frac{|\lambda-2t|}{A_t^2} dt \right)^{1-\frac{1}{q}} \left(\int_0^{1/2} \frac{|\lambda-2t|}{A_t^2} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left(\int_{1/2}^1 \frac{|2-\lambda-2t|}{A_t^2} dt \right)^{1-\frac{1}{q}} \left(\int_{1/2}^1 \frac{|2-\lambda-2t|}{A_t^2} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \right)^{\frac{1}{q}} \right\}.
\end{aligned}$$

Hence, by harmonically convexity of $|f'|^q$ on $[a, b]$, we have

$$\begin{aligned}
& \left| (1-\lambda) f\left(\frac{2ab}{a+b}\right) + \lambda \left(\frac{f(a)+f(b)}{2}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\
& \times \left\{ \left(\int_0^{1/2} \frac{|\lambda-2t|}{A_t^2} dt \right)^{1-\frac{1}{q}} \left(\int_0^{1/2} \frac{|\lambda-2t| [t|f'(a)|^q + (1-t)|f'(b)|^q]}{A_t^2} dt \right)^{\frac{1}{q}} \right. \\
& \left. + \left(\int_{1/2}^1 \frac{|2-\lambda-2t|}{A_t^2} dt \right)^{1-\frac{1}{q}} \left(\int_{1/2}^1 \frac{|2-\lambda-2t| [t|f'(a)|^q + (1-t)|f'(b)|^q]}{A_t^2} dt \right)^{\frac{1}{q}} \right\} \\
& \leq \frac{ab(b-a)}{2} C_1^{1-\frac{1}{q}}(\lambda; a, b) \left\{ [C_2(\lambda; a, b) |f'(a)|^q + C_3(\lambda; a, b) |f'(b)|^q]^{\frac{1}{q}} \right. \\
& \quad \left. [C_3(\lambda; b, a) |f'(a)|^q + C_2(\lambda; b, a) |f'(b)|^q]^{\frac{1}{q}} \right\}.
\end{aligned}$$

It is easily check that

$$\begin{aligned}
& \int_0^{1/2} \frac{|\lambda-2t|}{A_t^2} dt = C_1(\lambda; a, b) = \frac{1}{(b-a)^2} \\
& \times \left[-4 + \frac{[\lambda(b-a) + 2a](3a+b)}{a(a+b)} + 2 \ln \left(\frac{2a(a+b)}{(2a + \lambda(b-a))^2} \right) \right],
\end{aligned}$$

$$\begin{aligned}
& \int_0^{1/2} \frac{|\lambda - 2t| t}{A_t^2} dt = C_2(\lambda; a, b) = \frac{1}{(b-a)^3} \\
& \times \left\{ [\lambda(b-a) + 4a] \ln \left(\frac{[\lambda(b-a) + 2a]^2}{2a(a+b)} \right) \right. \\
& \quad \left. - \frac{[\lambda(b-a) + 2a](5a + 3b)}{a+b} + 7a + b \right\}, \\
& \int_0^{1/2} \frac{|\lambda - 2t| (1-t)}{A_t^2} dt = C_3(\lambda; a, b) = C_1(\lambda; a, b) - C_2(\lambda; a, b), \\
& \int_{1/2}^1 \frac{|2 - \lambda - 2t|}{A_t^2} dt = C_1(\lambda; b, a), \quad \int_{1/2}^1 \frac{|2 - \lambda - 2t| (1-t)}{A_t^2} dt = C_2(\lambda; b, a),
\end{aligned}$$

and

$$\int_{1/2}^1 \frac{|2 - \lambda - 2t| t}{A_t^2} dt = C_3(\lambda; b, a) = C_1(\lambda; b, a) - C_2(\lambda; b, a).$$

This concludes the proof. \square

Corollary 1. *Under the assumptions Theorem 5 with $\lambda = 0$, we have*

$$\begin{aligned}
& \left| f \left(\frac{2ab}{a+b} \right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\
& \times \left\{ C_1^{1-\frac{1}{q}}(0; a, b) [C_2(0; a, b) |f'(a)|^q + C_3(0; a, b) |f'(b)|^q]^{\frac{1}{q}} \right. \\
& \quad \left. C_1^{1-\frac{1}{q}}(0; b, a) [C_3(0; b, a) |f'(a)|^q + C_2(0; b, a) |f'(b)|^q]^{\frac{1}{q}} \right\},
\end{aligned}$$

where

$$\begin{aligned}
C_1(0; u, \vartheta) &= \frac{2}{(\vartheta - u)^2} \left[\ln \left(\frac{u + \vartheta}{2u} \right) - \frac{\vartheta - u}{u + \vartheta} \right], \\
C_2(0; u, \vartheta) &= \frac{1}{(\vartheta - u)^3} \left[\frac{(3u + \vartheta)(\vartheta - u)}{u + \vartheta} + 4u \ln \left(\frac{2u}{u + \vartheta} \right) \right], \\
C_3(0; u, \vartheta) &= \frac{1}{(\vartheta - u)^2} \left[\frac{2(u + \vartheta)}{\vartheta - u} \ln \left(\frac{u + \vartheta}{2u} \right) - \frac{u + 3\vartheta}{u + \vartheta} \right], \quad u, \vartheta > 0.
\end{aligned}$$

Corollary 2. *Under the assumptions Theorem 5 with $\lambda = 1$, we have*

$$\left| \frac{f(a) + f(b)}{2} - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2}$$

$$\times \left\{ C_1^{1-\frac{1}{q}}(1; a, b) [C_2(1; a, b) |f'(a)|^q + C_3(1; a, b) |f'(b)|^q]^{\frac{1}{q}} \right. \\ \left. C_1^{1-\frac{1}{q}}(1; b, a) [C_3(1; b, a) |f'(a)|^q + C_2(1; b, a) |f'(b)|^q]^{\frac{1}{q}} \right\},$$

where

$$\begin{aligned} C_1(1; u, \vartheta) &= \frac{1}{(\vartheta - u)^2} \left[\frac{\vartheta - u}{u} + 2 \ln \left(\frac{2u}{u + \vartheta} \right) \right], \\ C_2(1; u, \vartheta) &= \frac{1}{(\vartheta - u)^3} \left[(3u + \vartheta) \ln \left(\frac{u + \vartheta}{2u} \right) - 2(\vartheta - u) \right], \\ C_3(1; u, \vartheta) &= \frac{1}{(\vartheta - u)^2} \left[\frac{u + \vartheta}{u} - \frac{u + 3\vartheta}{\vartheta - u} \ln \left(\frac{u + \vartheta}{2u} \right) \right], \quad u, \vartheta > 0. \end{aligned}$$

Corollary 3. Under the assumptions Theorem 5 with $\lambda = 1/3$, we have

$$\left| \frac{1}{3} \left[\frac{f(a) + f(b)}{2} + 2f \left(\frac{2ab}{a+b} \right) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\ \times \left\{ C_1^{1-\frac{1}{q}}(1/3; a, b) [C_2(1/3; a, b) |f'(a)|^q + C_3(1/3; a, b) |f'(b)|^q]^{\frac{1}{q}} \right. \\ \left. C_1^{1-\frac{1}{q}}(1/3; b, a) [C_3(1/3; b, a) |f'(a)|^q + C_2(1/3; b, a) |f'(b)|^q]^{\frac{1}{q}} \right\},$$

where

$$\begin{aligned} C_1(1/3; u, \vartheta) &= \frac{1}{(\vartheta - u)^2} \left[\frac{(\vartheta - u)(\vartheta - 3u)}{3u(u + \vartheta)} + 2 \ln \left(\frac{18u(u + \vartheta)}{(5u + \vartheta)^2} \right) \right], \\ C_2(1/3; u, \vartheta) &= \frac{1}{(\vartheta - u)^3} \left[\left(\frac{11u + \vartheta}{3} \right) \ln \left(\frac{(5u + \vartheta)^2}{18u(u + \vartheta)} \right) + \frac{4u(\vartheta - u)}{3(u + \vartheta)} \right], \\ C_3(1/3; u, \vartheta) &= \frac{1}{(\vartheta - u)^2} \left[\frac{\vartheta^2 - 4u\vartheta - u^2}{3u(u + \vartheta)} + \frac{5u + 7\vartheta}{3(\vartheta - u)} \ln \left(\frac{18u(u + \vartheta)}{(5u + \vartheta)^2} \right) \right], \quad u, \vartheta > 0. \end{aligned}$$

Theorem 6. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° , $a, b \in I$ with $a < b$, and $f' \in L[a, b]$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q > 1$ and then we have the following inequality for $\lambda \in [0, 1]$

$$(2.2) \quad \left| (1 - \lambda) f \left(\frac{2ab}{a+b} \right) + \lambda \left(\frac{f(a) + f(b)}{2} \right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\ \times \left\{ C_1^{\frac{1}{p}}(\lambda, p; a, b) \left(\frac{|f'(\frac{2ab}{a+b})|^q + |f'(b)|^q}{4} \right)^{\frac{1}{q}} + C_1^{\frac{1}{p}}(\lambda, p; b, a) \left(\frac{|f'(\frac{2ab}{a+b})|^q + |f'(a)|^q}{4} \right)^{\frac{1}{q}} \right\},$$

where

$$C_1(\lambda, p; u, \vartheta) = \int_0^{1/2} \frac{|\lambda - 2t|^p}{(tb + (1-t)a)^{2p}} dt, \quad u, \vartheta > 0.$$

and $1/p + 1/q = 1$.

Proof. Let $A_t = tb + (1-t)a$. Using Lemma 1 and Hölder's integral inequality, we deduce

$$\begin{aligned}
 & \left| (1-\lambda) f\left(\frac{2ab}{a+b}\right) + \lambda \left(\frac{f(a)+f(b)}{2}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\
 & \leq \frac{ab(b-a)}{2} \left[\int_0^{1/2} \frac{|\lambda-2t|}{A_t^2} \left| f'\left(\frac{ab}{A_t}\right) \right| dt + \int_{1/2}^1 \frac{|2-\lambda-2t|}{A_t^2} \left| f'\left(\frac{ab}{A_t}\right) \right| dt \right] \\
 (2.3) \quad & \leq \frac{ab(b-a)}{2} \left\{ \left(\int_0^{1/2} \frac{|\lambda-2t|^p}{A_t^{2p}} dt \right)^{\frac{1}{p}} \left(\int_0^{1/2} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \right)^{\frac{1}{q}} \right. \\
 & \quad \left. + \left(\int_{1/2}^1 \frac{|2-\lambda-2t|^p}{A_t^{2p}} dt \right)^{\frac{1}{p}} \left(\int_{1/2}^1 \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \right)^{\frac{1}{q}} \right\}.
 \end{aligned}$$

Using the harmonically convexity of $|f'|^q$, we obtain the following inequalities from inequality (1.3):

$$\begin{aligned}
 \int_0^{1/2} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt & \leq \frac{1}{2} \left(\frac{2ab}{b-a} \int_a^b \frac{|f'(x)|^q}{x^2} dx \right) \\
 (2.4) \quad & \leq \frac{\left| f'\left(\frac{2ab}{a+b}\right) \right|^q + |f'(b)|^q}{4}
 \end{aligned}$$

and

$$\begin{aligned}
 \int_{1/2}^1 \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt & \leq \frac{1}{2} \left(\frac{2ab}{b-a} \int_a^b \frac{|f'(x)|^q}{x^2} dx \right) \\
 (2.5) \quad & \leq \frac{\left| f'\left(\frac{2ab}{a+b}\right) \right|^q + |f'(a)|^q}{4}.
 \end{aligned}$$

A combination of (2.3)-(2.5) gives the required inequality (2.2). \square

Corollary 4. Under the assumptions Theorem 6 with $\lambda = 0$, we have

$$\begin{aligned}
 & \left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\
 & \times \left\{ C_1^{\frac{1}{p}}(0, p; a, b) \left(\frac{\left| f'\left(\frac{2ab}{a+b}\right) \right|^q + |f'(b)|^q}{4} \right)^{\frac{1}{q}} + C_1^{\frac{1}{p}}(0, p; b, a) \left(\frac{\left| f'\left(\frac{2ab}{a+b}\right) \right|^q + |f'(a)|^q}{4} \right)^{\frac{1}{q}} \right\}.
 \end{aligned}$$

Corollary 5. *Under the assumptions Theorem 6 with $\lambda = 1$, we have*

$$\left| \frac{f(a) + f(b)}{2} - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\ \times \left\{ C_1^{\frac{1}{p}}(1, p; a, b) \left(\frac{|f'(\frac{2ab}{a+b})|^q + |f'(b)|^q}{4} \right)^{\frac{1}{q}} + C_1^{\frac{1}{p}}(1, p; b, a) \left(\frac{|f'(\frac{2ab}{a+b})|^q + |f'(a)|^q}{4} \right)^{\frac{1}{q}} \right\}.$$

Corollary 6. *Under the assumptions Theorem 6 with $\lambda = 1/3$, we have*

$$\left| \frac{1}{3} \left[\frac{f(a) + f(b)}{2} + 2f\left(\frac{2ab}{a+b}\right) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{2} \\ \times \left\{ C_1^{\frac{1}{p}}\left(\frac{1}{3}, p; a, b\right) \left(\frac{|f'(\frac{2ab}{a+b})|^q + |f'(b)|^q}{4} \right)^{\frac{1}{q}} + C_1^{\frac{1}{p}}\left(\frac{1}{3}, p; b, a\right) \left(\frac{|f'(\frac{2ab}{a+b})|^q + |f'(a)|^q}{4} \right)^{\frac{1}{q}} \right\}$$

Theorem 7. *Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° , $a, b \in I$ with $a < b$, and $f' \in L[a, b]$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q > 1$ and then we have the following inequality for $\lambda \in [0, 1]$*

$$(2.6) \quad \left| (1-\lambda)f\left(\frac{2ab}{a+b}\right) + \lambda\left(\frac{f(a) + f(b)}{2}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{4}$$

$$\times \frac{C_4^{1/p}(\lambda, p)}{\left[(1-q)(1-2q)(b-a)^2\right]^{1/q}} \left\{ (C_5(q; a, b)|f'(a)|^q + C_6(q; a, b)|f'(b)|^q)^{\frac{1}{q}} \right. \\ \left. + (C_6(q; b, a)|f'(a)|^q + C_5(q; b, a)|f'(b)|^q)^{\frac{1}{q}} \right\},$$

where

$$C_4(\lambda, p) = \frac{\lambda^{p+1} + (1-\lambda)^{p+1}}{p+1},$$

$$C_5(q; u, \vartheta) = \left[\left(\frac{u + \vartheta}{2} \right)^{1-2q} \left[\frac{\vartheta - 3u}{2} - q(\vartheta - u) \right] + u^{2-2q} \right],$$

$$C_6(q; u, \vartheta) = \left[\left(\frac{u + \vartheta}{2} \right)^{1-2q} \left[\frac{3\vartheta - u}{2} - q(\vartheta - u) \right] + u^{1-2q} [u - 2\vartheta + 2q(\vartheta - u)] \right], \quad u, \vartheta > 0$$

and $1/p + 1/q = 1$.

Proof. Let $A_t = tb + (1-t)a$. Using Lemma 1 and Hölder's integral inequality, we deduce

$$\begin{aligned}
 & \left| (1-\lambda) f\left(\frac{2ab}{a+b}\right) + \lambda \left(\frac{f(a)+f(b)}{2}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\
 & \leq \frac{ab(b-a)}{2} \left[\int_0^{1/2} \frac{|\lambda-2t|}{A_t^{2q}} \left| f'\left(\frac{ab}{A_t}\right) \right| dt + \int_{1/2}^1 \frac{|2-\lambda-2t|}{A_t^{2q}} \left| f'\left(\frac{ab}{A_t}\right) \right| dt \right] \\
 (2.7) \quad & \leq \frac{ab(b-a)}{2} \left\{ \left(\int_0^{1/2} |\lambda-2t|^p dt \right)^{\frac{1}{p}} \left(\int_0^{1/2} \frac{1}{A_t^{2q}} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \right)^{\frac{1}{q}} \right. \\
 & \quad \left. + \left(\int_{1/2}^1 |2-\lambda-2t|^p dt \right)^{\frac{1}{p}} \left(\int_{1/2}^1 \frac{1}{A_t^{2q}} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \right)^{\frac{1}{q}} \right\}.
 \end{aligned}$$

Using the harmonically convexity of $|f'|^q$, we obtain

$$\begin{aligned}
 & \int_0^{1/2} \frac{1}{A_t^{2q}} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \leq \int_0^{1/2} \frac{t|f'(a)|^q + (1-t)|f'(b)|^q}{A_t^{2q}} dt = \frac{1}{2(1-q)(1-2q)(b-a)^2} \\
 (2.8) \quad & \times \left\{ \left[\left(\frac{a+b}{2}\right)^{1-2q} \left[\frac{b-3a}{2} - q(b-a) \right] + a^{2-2q} \right] |f'(a)|^q \right. \\
 & \quad \left. + \left[\left(\frac{a+b}{2}\right)^{1-2q} \left[\frac{3b-a}{2} - q(b-a) \right] + a^{1-2q} [a-2b+2q(b-a)] \right] |f'(b)|^q \right\}
 \end{aligned}$$

and

$$\begin{aligned}
 & \int_{1/2}^1 \frac{1}{A_t^{2q}} \left| f'\left(\frac{ab}{A_t}\right) \right|^q dt \leq \int_{1/2}^1 \frac{t|f'(a)|^q + (1-t)|f'(b)|^q}{A_t^{2q}} dt = \frac{1}{2(1-q)(1-2q)(b-a)^2} \\
 (2.9) \quad & \times \left\{ \left[b^{1-2q} [b-2a-2q(b-a)] + \left(\frac{a+b}{2}\right)^{1-2q} \left[\frac{3a-b}{2} + q(b-a) \right] \right] |f'(a)|^q \right. \\
 & \quad \left. + \left[\left(\frac{a+b}{2}\right)^{1-2q} \left[\frac{a-3b}{2} + q(b-a) \right] + b^{2-2q} \right] |f'(b)|^q \right\}
 \end{aligned}$$

Further, we have

$$(2.10) \quad \int_0^{1/2} |\lambda-2t|^p dt = \int_{1/2}^1 |2-\lambda-2t|^p dt = \frac{\lambda^{p+1} + (1-\lambda)^{p+1}}{2(p+1)}$$

A combination of (2.7)-(2.10) gives the required inequality (2.6). \square

Corollary 7. *Under the assumptions Theorem 7 with $\lambda = 0$, we have*

$$\begin{aligned} & \left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{4(p+1)^{1/p}} \\ & \times \frac{1}{\left[(1-q)(1-2q)(b-a)^2\right]^{1/q}} \left\{ (C_5(q; a, b) |f'(a)|^q + C_6(q; a, b) |f'(b)|^q)^{\frac{1}{q}} \right. \\ & \left. + (C_6(q; b, a) |f'(a)|^q + C_5(q; b, a) |f'(b)|^q)^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 8. *Under the assumptions Theorem 7 with $\lambda = 1$, we have*

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{4(p+1)^{1/p}} \\ & \times \frac{1}{\left[(1-q)(1-2q)(b-a)^2\right]^{1/q}} \left\{ (C_5(q; a, b) |f'(a)|^q + C_6(q; a, b) |f'(b)|^q)^{\frac{1}{q}} \right. \\ & \left. + (C_6(q; b, a) |f'(a)|^q + C_5(q; b, a) |f'(b)|^q)^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 9. *Under the assumptions Theorem 7 with $\lambda = 1/3$, we have*

$$\begin{aligned} & \left| \frac{1}{3} \left[\frac{f(a) + f(b)}{2} + 2f\left(\frac{2ab}{a+b}\right) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq \frac{ab(b-a)}{4(3^{p+1}(p+1))^{1/p}} \\ & \times \frac{1 + 2^{p+1}}{\left[(1-q)(1-2q)(b-a)^2\right]^{1/q}} \left\{ (C_5(q; a, b) |f'(a)|^q + C_6(q; a, b) |f'(b)|^q)^{\frac{1}{q}} \right. \\ & \left. + (C_6(q; b, a) |f'(a)|^q + C_5(q; b, a) |f'(b)|^q)^{\frac{1}{q}} \right\}. \end{aligned}$$

3. SOME APPLICATIONS FOR SPECIAL MEANS

Let us recall the following special means of two nonnegative number a, b with $b > a$:

(1) The arithmetic mean

$$A = A(a, b) := \frac{a+b}{2}.$$

(2) The geometric mean

$$G = G(a, b) := \sqrt{ab}.$$

(3) The harmonic mean

$$H = H(a, b) := \frac{2ab}{a+b}.$$

(4) The Logarithmic mean

$$L = L(a, b) := \frac{b-a}{\ln b - \ln a}.$$

(5) The p -Logarithmic mean

$$L_p = L_p(a, b) := \left(\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right)^{\frac{1}{p}}, \quad p \in \mathbb{R} \setminus \{-1, 0\}.$$

(6) the Identric mean

$$I = I(a, b) = \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{\frac{1}{b-a}}.$$

These means are often used in numerical approximation and in other areas. However, the following simple relationships are known in the literature:

$$H \leq G \leq L \leq I \leq A.$$

It is also known that L_p is monotonically increasing over $p \in \mathbb{R}$, denoting $L_0 = I$ and $L_{-1} = L$.

Proposition 1. *Let $0 < a < b$ and $\lambda \in [0, 1]$. Then we have the following inequality*

$$\left| (1-\lambda)H + \lambda A - \frac{G^2}{L} \right| \leq \frac{ab(b-a)}{2} \{C_1(\lambda; a, b) + C_1(\lambda; b, a)\},$$

where C_1 is defined as in Theorem 5.

Proof. The assertion follows from the inequality (2.1) in Theorem 5, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x$. \square

Proposition 2. *Let $0 < a < b$ and $\lambda \in [0, 1]$. Then we have the following inequality*

$$\left| (1-\lambda)H + \lambda A - \frac{G^2}{L} \right| \leq \frac{ab(b-a)}{2^{1+1/q}} \left\{ C_1^{\frac{1}{p}}(\lambda, p; a, b) + C_1^{\frac{1}{p}}(\lambda, p; b, a) \right\},$$

where $q > 1$, $1/p + 1/q = 1$ and C_1 is defined as in Theorem 6.

Proof. The assertion follows from the inequality (2.2) in Theorem 6, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x$. \square

Proposition 3. *Let $0 < a < b$ and $\lambda \in [0, 1]$. Then we have the following inequality*

$$\begin{aligned} \left| (1-\lambda)H + \lambda A - \frac{G^2}{L} \right| &\leq \frac{ab(b-a)C_4^{1/p}(\lambda, p)}{4 \left[(1-q)(1-2q)(b-a)^2 \right]^{1/q}} \\ &\times \left\{ (C_5(q; a, b) + C_6(q; a, b))^{\frac{1}{q}} + (C_6(q; b, a) + C_5(q; b, a))^{\frac{1}{q}} \right\}, \end{aligned}$$

where $q > 1$, $1/p + 1/q = 1$ and C_4, C_5 and C_6 are defined as in Theorem 7.

Proof. The assertion follows from the inequality (2.6) in Theorem 7, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x$. \square

Proposition 4. *Let $0 < a < b$, $\lambda \in [0, 1]$ and $q \geq 1$. Then we have the following inequality*

$$\begin{aligned} &|(1-\lambda)H^2 + \lambda A(a^2, b^2) - G^2| \\ &\leq ab(b-a) \left\{ C_1^{1-\frac{1}{q}}(\lambda; a, b) [C_2(\lambda; a, b)a^q + C_3(\lambda; a, b)b^q]^{\frac{1}{q}} \right. \\ &\quad \left. + C_1^{1-\frac{1}{q}}(\lambda; b, a) [C_3(\lambda; b, a)a^q + C_2(\lambda; b, a)b^q]^{\frac{1}{q}} \right\}, \end{aligned}$$

where C_1, C_2 and C_3 are defined as in Theorem 5.

Proof. The assertion follows from the inequality (2.1) in Theorem 5, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^2$. \square

Proposition 5. Let $0 < a < b$ and $\lambda \in [0, 1]$. Then we have the following inequality

$$\begin{aligned} |(1-\lambda)H^2 + \lambda A(a^2, b^2) - G^2| &\leq \frac{ab(b-a)}{2^{1/q}} \\ &\times \left\{ C_1^{\frac{1}{p}}(\lambda, p; a, b) A^{\frac{1}{q}}(H^q, b^q) + C_1^{\frac{1}{p}}(\lambda, p; b, a) A^{\frac{1}{q}}(a^q, H^q) \right\}, \end{aligned}$$

where $q > 1$ and $1/p + 1/q = 1$.

Proof. The assertion follows from the inequality (2.2) in Theorem 6, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^2$. \square

Proposition 6. Let $0 < a < b$ and $\lambda \in [0, 1]$. Then we have the following inequality

$$\begin{aligned} |(1-\lambda)H^2 + \lambda A(a^2, b^2) - G^2| &\leq \frac{ab(b-a)C_4^{1/p}(\lambda, p)}{2 \left[(1-q)(1-2q)(b-a)^2 \right]^{1/q}} \\ &\times \left\{ (C_5(q; a, b)a^q + C_6(q; a, b)b^q)^{\frac{1}{q}} + (C_6(q; b, a)a^q + C_5(q; b, a)b^q)^{\frac{1}{q}} \right\}, \end{aligned}$$

where $q > 1$, $1/p + 1/q = 1$ and C_4, C_5 and C_6 are defined as in Theorem 7.

Proof. The assertion follows from the inequality (2.6) in Theorem 7, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^2$. \square

Proposition 7. Let $0 < a < b$, $n \in (-1, \infty) \setminus \{0\}$, $\lambda \in [0, 1]$ and $q \geq 1$. Then we have the following inequality

$$\begin{aligned} &|(1-\lambda)H^{n+2} + \lambda A(a^{n+2}, b^{n+2}) - G^2.L_n^n| \\ &\leq \frac{ab(b-a)(n+2)}{2} \left\{ C_1^{1-\frac{1}{q}}(\lambda; a, b) \left[C_2(\lambda; a, b)a^{(n+1)q} + C_3(\lambda; a, b)b^{(n+1)q} \right]^{\frac{1}{q}} \right. \\ &\quad \left. + C_1^{1-\frac{1}{q}}(\lambda; b, a) \left[C_3(\lambda; b, a)a^{(n+1)q} + C_2(\lambda; b, a)b^{(n+1)q} \right]^{\frac{1}{q}} \right\}, \end{aligned}$$

where C_1, C_2 and C_3 are defined as in Theorem 5.

Proof. The assertion follows from the inequality (2.1) in Theorem 5, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^{n+2}$, $n \in (-1, \infty) \setminus \{0\}$. \square

Proposition 8. Let $0 < a < b$ and $n \in (-1, \infty) \setminus \{0\}$. Then we have the following inequality

$$\begin{aligned} &|(1-\lambda)H^{n+2} + \lambda A(a^{n+2}, b^{n+2}) - G^2.L_n^n| \leq \frac{ab(b-a)(n+2)}{2^{1+1/q}} \\ &\times \left\{ C_1^{1-\frac{1}{q}}(\lambda; a, b) A^{\frac{1}{q}}(H^{(n+1)q}, b^{(n+1)q}) + C_1^{1-\frac{1}{q}}(\lambda; b, a) A^{\frac{1}{q}}(a^{(n+1)q}, H^{(n+1)q}) \right\}, \end{aligned}$$

where $q > 1$, $1/p + 1/q = 1$ and C_1 is defined as in Theorem 6.

Proof. The assertion follows from the inequality (2.2) in Theorem 6, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^{n+2}$, $n \in (-1, \infty) \setminus \{0\}$. \square

Proposition 9. Let $0 < a < b$, $\lambda \in [0, 1]$. and $n \in (-1, \infty) \setminus \{0\}$. Then we have the following inequality

$$\begin{aligned} & |(1 - \lambda) H^{n+2} + \lambda A(a^{n+2}, b^{n+2}) - G^2.L_n^n| \\ & \leq \frac{ab(b-a)(n+2)C_4^{1/p}(\lambda, p)}{4 \left[(1-q)(1-2q)(b-a)^2 \right]^{1/q}} \left\{ \left(C_5(q; a, b) a^{(n+1)q} + C_6(q; a, b) b^{(n+1)q} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(C_6(q; b, a) a^{(n+1)q} + C_5(q; b, a) b^{(n+1)q} \right)^{\frac{1}{q}} \right\}, \end{aligned}$$

where $q > 1$, $1/p + 1/q = 1$ and C_5 and C_6 are defined as in Theorem 7.

Proof. The assertion follows from the inequality (2.6) in Theorem 7, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^{n+2}$, $n \in (-1, \infty) \setminus \{0\}$. \square

Proposition 10. Let $0 < a < b$, $\lambda \in [0, 1]$ and $q \geq 1$. Then we have the following inequality

$$\begin{aligned} & |(1 - \lambda) H^2 \ln H + \lambda A(a^2 \ln a, b^2 \ln b) - G^2 \ln I| \leq ab(b-a) \\ & \times \left\{ C_1^{1-\frac{1}{q}}(\lambda; a, b) [C_2(\lambda; a, b) G^{2q}(a, A(1, \ln a)) + C_3(\lambda; a, b) G^{2q}(b, A(1, \ln b))]^{\frac{1}{q}} \right. \\ & \quad \left. + C_1^{1-\frac{1}{q}}(\lambda; b, a) [C_3(\lambda; b, a) G^{2q}(a, A(1, \ln a)) + C_2(\lambda; b, a) G^{2q}(b, A(1, \ln b))]^{\frac{1}{q}} \right\}, \end{aligned}$$

where C_1, C_2 and C_3 are defined as in Theorem 5.

Proof. The assertion follows from the inequality (2.1) in Theorem 5, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^2 \ln x$. \square

Proposition 11. Let $0 < a < b$, and $\lambda \in [0, 1]$. Then we have the following inequality

$$\begin{aligned} & |(1 - \lambda) H^2 \ln H + \lambda A(a^2 \ln a, b^2 \ln b) - G^2 \ln I| \leq \frac{ab(b-a)}{2^{1/q}} \\ & \times \left\{ C_1^{\frac{1}{p}}(0, p; a, b) A^{\frac{1}{q}}(G^{2q}(H, A(1, \ln H)), G^{2q}(b, A(1, \ln b))) \right. \\ & \quad \left. + C_1^{\frac{1}{p}}(0, p; b, a) A^{\frac{1}{q}}(G^{2q}(H, A(1, \ln H)), G^{2q}(a, A(1, \ln a))) \right\}, \end{aligned}$$

where $q > 1$, $1/p + 1/q = 1$ and C_1 is defined as in Theorem 6.

Proof. The assertion follows from the inequality (2.2) in Theorem 6, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^2 \ln x$. \square

Proposition 12. Let $0 < a < b$, and $\lambda \in [0, 1]$. Then we have the following inequality

$$|(1 - \lambda) H^2 \ln H + \lambda A(a^2 \ln a, b^2 \ln b) - G^2 \ln I| \leq \frac{ab(b-a)}{2}$$

$$\frac{C_4^{1/p}(\lambda, p)}{\left[(1-q)(1-2q)(b-a)^2\right]^{1/q}} \left\{ \left(C_5(q; a, b)G^{2q}(a, A(1, \ln a)) + C_6(q; a, b)G^{2q}(b, A(1, \ln b)) \right)^{\frac{1}{q}} \right. \\ \left. + \left(C_6(q; b, a)G^{2q}(a, A(1, \ln a)) + C_5(q; b, a)G^{2q}(b, A(1, \ln b)) \right)^{\frac{1}{q}} \right\},$$

where $q > 1$, $1/p + 1/q = 1$ and C_5 and C_6 are defined as in Theorem 7.

Proof. The assertion follows from the inequality (2.6) in Theorem 7, for $f : (0, \infty) \rightarrow \mathbb{R}$, $f(x) = x^2 \ln x$. \square

REFERENCES

- [1] S.S. Dragomir and C.E.M. Pearce, Selected Topics on Hermite-Hadamard Inequalities and Applications, RGMIA Monographs, Victoria University, 2000.
- [2] I. Iscan, New estimates on generalization of some integral inequalities for (α, m) -convex functions, Contemp. Anal. Appl. Math. 1(2) (2013) 253-264.
- [3] I. Iscan, New estimates on generalization of some integral inequalities for s -convex functions and their applications, Int. J. Pure Appl. Math. 86(4) (2013) 727-746.
- [4] I. Iscan, Generalization of different type integral inequalities via fractional integrals for functions whose second derivatives absolute values are quasi-convex, Konuralp J. Math. 1(2) (2013) 67-79.
- [5] I. Iscan, Hermite-Hadamard type inequalities for harmonically convex functions, Hacettepe Journal of Mathematics and Statistics. Accepted for publication.
- [6] U.S. Kirmaci, Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, Appl. Math. Comput. 147 (2004) 137-146
- [7] B.-Y. Xi and F. Qi, Some Integral Inequalities of Hermite-Hadamard Type for Convex Functions with Applications to Means, J. Funct. Spaces Appl. 2012 (2012), 14 pages. Article ID 980438. doi:10.1155/2012/980438.
- [8] G.-S. Yang, D.-Y. Hwnag and K.-L. Tseng, Some Inequalities for Differentiable Convex and Concave Mappings, Comput. Math. Appl. 47 (2004) 207-216

DEPARTMENT OF MATHEMATICS, FACULTY OF ARTS AND SCIENCES,, GIRESUN UNIVERSITY, 28100, GIRESUN, TURKEY.

E-mail address: imdati@yahoo.com, imdat.iscan@giresun.edu.tr