

TRIVIALITY AND SPLIT OF VECTOR BUNDLES ON RATIONAL CONNECTED VARIETIES

XUANYU PAN

ABSTRACT. In this paper, we give a very simple proof of a triviality criterion due to I.Biswas and J.Pedro and P.Dos Santos. We also prove a vector bundle on a homogenous space is trivial if and only if the restrictions of the vector bundle to Schubert lines are trivial. Using this result and Chern classes of vector bundles, we give a general criterion of a uniform vector bundle on a homogenous space to be splitting. As an application, we prove a uniform vector bundle on classical Grassmannians and quadrics of low rank is splitting.

CONTENTS

1. Introduction	1
2. Triviality of Vector Bundles and Families of Rational Curves	2
3. Triviality of Vector Bundles on Homogenous Spaces	4
4. A Splitting Criterion of Vector Bundles on Homogenous Spaces	6
5. Applications	8
References	9

1. INTRODUCTION

The study of uniform vector bundles on homogenous space starts from the papers [Sat76], [VdV72], and [Bal81]. Recently, [MOSC12] give a more systematical approach. The proofs usually come with a triviality criterion for a vector bundle.

In the preprint [IJS10], it provides a beautiful triviality criterion for a vector bundle on a separably rationally connected variety. In this paper, we give a simple proof of this theorem by a technique from [Kol08]. Namely,

Theorem 1.1. *Let X be a smooth projective variety over an algebraic closed field. If X is separably rationally connected, then a vector bundle on X is trivial if and only if the restriction of V to every rational curve on X is trivial.*

We also use the technique and irreducibility of some moduli space of some rational chains on homogenous spaces to give a triviality criterion for vector bundles on homogenous spaces, namely,

Theorem 1.2. *For a homogenous space G/P , a vector bundle V on G/P is trivial if and only if the restriction of V to every Schubert line is trivial.*

Use this triviality criterion and the theory of Chern Classes, we give a splitting criterion for low rank uniform vector bundles on homogenous spaces.

Date: July 22, 2022.

Theorem 1.3. *Let G/P be a homogenous space of Picard number one. If its VRMT has good divisibility up to degree r , then a uniform vector bundle V on G/P of rank at most r is splitting.*

As an application, we apply this theorem to classical Grassmannians and quadrics.

Theorem 1.4. *A uniform vector bundle on classical Grassmannian or quadrics X of rank at most $s(X)$ is splitting, for the corresponding X and $s(X)$, see Theorem 5.6.*

Acknowledgments. I am very grateful for my advisor Prof. A. de Jong , Prof. J. Starr, Xue Hang , Yang Yanhong and Yi Zhu for useful conversations.

2. TRIVIALITY OF VECTOR BUNDLES AND FAMILIES OF RATIONAL CURVES

We start from a very general triviality criterion for a vector bundle on a smooth projective variety to be trivial.

Proposition 2.1. *Let X be a smooth projective variety over an algebraic closed field k of $\text{char}(k) = 0$. A vector bundle E of rank n on X is trivial if and only if the restriction of E on every curve $C \subseteq X$ is trivial.*

Proof. We prove this proposition by induction on the dimension. For $\dim(X) = 1$, it is trivial. By the existence of Lefschetz pencil, see [FK88], we always have the following blow-up diagram,

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{f} & X \\ \downarrow \pi & & \\ \mathbb{P}^1 & & \end{array}$$

where f is a blowing up map along a smooth projective subvariety in X of codimension 2, the map π is a Lefschetz pencil. Therefore, we have $f^*\mathcal{O}_{\tilde{X}} = \mathcal{O}_X$. By induction, the restriction to a smooth fiber $f^*V|_{X_s}$ is trivial bundle for every $s \in \mathbb{P}^1$. Suppose \tilde{X}_s is a singular fiber with one double singularity. We blow up the singularity and get a resolution

$$g : \tilde{X}_s^{Bl} \rightarrow \tilde{X}_s$$

with $g_*(\mathcal{O}_{\tilde{X}_s^{Bl}}) = \mathcal{O}_{\tilde{X}_s}$. By induction, we have

$$g^*(f^*V|_{\tilde{X}_s}) = \mathcal{O}_{\tilde{X}_s^{Bl}}^{\oplus n}.$$

By projection formula, it implies that

$$f^*V|_{\tilde{X}_s} = g_*g^*(f^*V|_{\tilde{X}_s}) = \mathcal{O}_{\tilde{X}_s}^{\oplus n}$$

Since π is flat, there exists a vector bundle W on \mathbb{P}_1^k such that $\pi^*W = f^*V$ by base change theorem. We can choose a smooth projective curve C such that

$$\begin{array}{ccccc} C & \xrightarrow{h} & \tilde{X} & \xrightarrow{f} & X \\ & \searrow p & \downarrow \pi & & \\ & & \mathbb{P}^1 & & \end{array}$$

where p is a finite map of degree N , hence, it induces a trace map

$$\text{tr} : p_*p^*(\mathcal{O}_{\mathbb{P}^1}) = p_*(\mathcal{O}_C) \rightarrow \mathcal{O}_{\mathbb{P}^1}.$$

The natural map $i : \mathcal{O}_{\mathbb{P}^1} \rightarrow p_*p^*(\mathcal{O}_{\mathbb{P}^1})$ is induced by the adjointness. The composition $tr \circ i$ is just an isomorphism of $\mathcal{O}_{\mathbb{P}^1}$. In fact, the map is multiplication by N . Therefore, we have

$$\begin{array}{ccc} W & \longrightarrow & p_*p^*W \xrightarrow{tr} W \\ & \parallel & \parallel \\ & W \otimes p_*(\mathcal{O}_C) & \xrightarrow{tr} W \otimes \mathcal{O}_{\mathbb{P}^1} \end{array}$$

By the induction, we have $p^*W = h^*\pi^*W = h^*f^*V$ is trivial, so the last trace map above is just

$$tr^{\oplus n} : p_*(\mathcal{O}_C)^{\oplus n} \rightarrow (\mathcal{O}_{\mathbb{P}^1})^{\oplus n}$$

Hence, we have $W = (\mathcal{O}_{\mathbb{P}^1})^{\oplus n}$. It completes the proof. \square

For a separable rationally connected variety, we can choose the test curves to be rational curves.

Lemma 2.2. *Let X is a normal scheme and B, C are Artin stacks. Suppose we have the following diagram,*

$$\begin{array}{ccc} & C & \xrightarrow{f} X \\ \sigma \nearrow & \downarrow \pi & \\ & B & \end{array}$$

A vector bundle V on X is trivial if we have the following hypothesis,

- (1) the morphism π is representable and flat with fibers of arithmetic genus 0,
- (2) the image of $f \circ \sigma$ is a point,
- (3) the morphism f is proper and its general fibers are nonempty, connected and generic reduced.
- (4) C is irreducible and generic reduced.

If the restrictions $f^*V|_{C_s}$ are trivial on fibers C_s of π , then the vector bundle V is trivial.

Proof. Since a general fiber $f^{-1}(p)$ is irreducible and generic reduced, i.e, hypothesis (3) and (4), by Stein factorization for stacks [LMB00] or [Ols07], we have $f_*\mathcal{O}_C = \mathcal{O}_X$. Since $f^*V|_{C_s}$ are trivial and fibers C_s are of arithmetic genus 0, we have the following properties by base change theorem, see [LMB00],

- (1) $\pi^*\pi_*(f^*V) \rightarrow f^*V$,
- (2) $\pi_*(f^*V)$ is an isomorphism.

In other words, there exists a vector bundle W on B such that $\pi^*W = f^*V$. Hence, we have that $W = \sigma^*\pi^*W = \sigma^*f^*V = E$ where E is a trivial bundle since $\sigma \circ f$ is trivial. It implies $f^*V = \pi^*W$ is trivial bundle. By projection formula, see [LMB00], we have

$$V = V \otimes f_*\mathcal{O}_C = f_*(f^*V) = (f_*\mathcal{O}_C)^{\oplus n} = \mathcal{O}_X^{\oplus n}$$

\square

Lemma 2.3. [Kol08, Theorem 8.10] *For a separably rationally connected variety X , there is a family of very free rational curves*

$$\begin{array}{ccc} B \times \mathbb{P}^1 & \xrightarrow{f} & X \\ \sigma \uparrow & & \downarrow \pi \\ & & B \end{array}$$

such that $f \circ \sigma$ is a point and the general fibers of f are smooth and irreducible.

The following theorem is due to [IJS10], we give a simple proof of this beautiful theorem.

Theorem 2.4. *Let X be a smooth projective variety over an algebraic closed field. If X is separably rationally connected, then a vector bundle on X is trivial if and only if the restriction of V to every rational curves on X is trivial.*

Proof. By Lemma 2.3, we have a morphism

$$h : B \rightarrow \overline{\mathcal{M}}_{0,0}(X, e),$$

where $\overline{\mathcal{M}}_{0,0}(X, e)$ is the Kontsevich moduli space parameterizing stable maps of arithmetic genus zero and of degree e . Since it is obvious that a vector bundle of a chain of rational curves is trivial if and only if its restriction to every component of the chain is trivial, take B be $\overline{Im}(h)$, then Lemma 2.2 implies the theorem immediately. \square

3. TRIVIALITY OF VECTOR BUNDLES ON HOMOGENOUS SPACES

From this section on, we take some notations from the paper [Zhu10] and assume k be an algebraically closed field of characteristic zero. Let $X = G/P$ be a projective homogeneous space under a linear algebraic k -group G . By Bruhat decomposition, the Picard lattice of X is freely generated by the line bundles associated to the Schubert varieties of codimension one, denoted by L_1, \dots, L_r . Dually, the curve classes are generated by Schubert curves, denoted by C_1, \dots, C_r .

Definition 3.1. *For numbers m and $i_1, \dots, i_m \in 1, \dots, r$, we define a moduli space of chains of Schubert curves with two pointed points and of the pattern (i_1, \dots, i_m) . More precisely,*

$$Chn_2(X, i_1, \dots, i_m) = \overline{\mathcal{M}}_{0,2}(X, [C_{i_1}]) \times_X \dots \times_X \overline{\mathcal{M}}_{0,2}(X, [C_{i_m}])$$

where the fiber product in the definition is induced by evaluation maps

$$ev_k : \overline{\mathcal{M}}_{0,2}(X, [C_{i_j}]) \rightarrow X,$$

($k = 1, 2$), in a natural way. Moreover, we have an obvious evaluation map induced by the first pointed point and last pointed point of the chains, this moduli space is the following

$$ev_{1,m+1} : Chn_2(X, i_1, \dots, i_m) \rightarrow X \times X.$$

see the figure below.




Figure 0

Lemma 3.2. *The moduli space $Chn_2(X, i_1, \dots, i_m)$ is a smooth irreducible and rationally connected variety.*

Proof. By [Zhu10, Proposition 10.1], the fibers of $ev : \overline{\mathcal{M}}_{0,1}(X, [C_i]) \rightarrow X$ are smooth irreducible and rationally connected varieties. Since we have the following cartesian diagram

$$\begin{array}{ccccc} ev_1^{-1}(p) & \longrightarrow & \overline{\mathcal{M}}_{0,2}(X, [C_i]) & \longrightarrow & X \times X \\ \downarrow & & \downarrow & & \downarrow pr_1 \\ ev^{-1}(p) & \longrightarrow & \overline{\mathcal{M}}_{0,1}(X, [C_i]) & \longrightarrow & X \end{array} \quad \square$$

where the middle vertical arrow is universal bundle over $\overline{\mathcal{M}}_{0,1}(X, [C_i])$, in particular, it is smooth (\mathbb{P}^1 bundle). Hence, the fiber $ev_1^{-1}(p)$ is \mathbb{P}^1 bundle over $ev^{-1}(p)$ which is a smooth projective and rationally connected variety. So $ev_1^{-1}(p)$ is a smooth projective rationally connected variety, by symmetry, so $ev_2^{-1}(p)$ is. By induction on m , it is easy to see

$$Chn_2(X, i_1, \dots, i_m) = \overline{\mathcal{M}}_{0,2}(X, [C_{i_1}]) \times_X \dots \times_X \overline{\mathcal{M}}_{0,2}(X, [C_{i_m}])$$

is smooth, irreducible and rationally connected. \square

Lemma 3.3. *There exists m and $i_1, \dots, i_m \in 1, \dots, r$ such that the fiber of the evaluation map $ev_{1,m+1}$ is irreducible.*

Proof. To prove this lemma, by Lemma 3.2 and [dJHS11, Lemma 15.6], it is sufficient to prove the evaluation map is surjective. In fact, it is what the proof of [Zhu10, Proposition 10.6] did. \square

Theorem 3.4. *For a homogenous space G/P , under the natural embedding, a vector bundle V on G/P is trivial if and only if the restriction of V to every Schubert line is trivial.*

Proof. By previous lemma, we have an evaluation map

$$ev_{1,m+1} : Chn_2(X, i_1, \dots, i_m) \rightarrow X \times X$$

which is surjective with integral fibers. Suppose $ev_{1,m+1} = (ev_1, ev_{m+1})$. Let C be $ev_1^{-1}(p)$. It is the universal space of a moduli space B parameterizing chains of Schubert lines passing through a point. For instance, we take forgetful functor to forget the second point of $Chn_2(X, i_1, \dots, i_m)$. In other words, we have

$$\begin{array}{ccc} & C & \xrightarrow{ev_{m+1}} X \\ \sigma \nearrow & \downarrow \pi & \\ & B & \end{array}$$

where $ev_{m+1} \circ \sigma$ is a constant map with image p . Since the fibers of π are chain of lines, in particular arithmetic genus 0. Since it is obvious that a vector bundle of a chain of rational curves is trivial if and only if its restriction to every component of the chain is trivial. Apply Lemma 2.2, we get the proof. \square

Remark 3.5. *The invertible sheaf $L = L_1 + \dots + L_r$ is ample. Since X is simply connected and homogeneous, by Stein factorization, the invertible sheaf L is in fact very ample. Therefore, using this embedding, Schubert curves are lines in some projective space \mathbb{P}^N , that is why we call it Schubert lines.*

4. A SPLITTING CRITERION OF VECTOR BUNDLES ON HOMOGENOUS SPACES

If a homogenous space G/P is of Picard number one, then the ample generator of its Picard group is very ample. Under the embedding induced by this very ample line bundle, we can talk about lines on G/P and the variety of minimal rational tangents (VRMT for short) on G/P . We refer to [Hwa01] for a complete account on the VMRT. In this case, VMRT is just the fiber of the evaluation map

$$ev : \overline{\mathcal{M}}_{0,1}(G/P, 1) \rightarrow G/P.$$

In principle, we can describe the VMRT of a homogenous space by the root data, see [LM03, Section 4] for the details.

Definition 4.1. *We call a smooth projective variety X has a good divisibility up to degree r if $x \cdot y = 0$, then we have $x = 0$ or $y = 0$, where $x \in CH^i(X)$, $y \in CH^j(X)$ and $i + j \leq r$.*

Theorem 4.2. *Let G/P be a homogenous space of Picard number one over complex numbers \mathbb{C} . If its VRMT has good divisibility up to degree r , then a uniform vector bundle V on G/P of rank at most r is splitting.*

Proof. Since we can twist the vector bundle V with the ample generator of $Pic(G/P)$, by Theorem 3.4, we can assume the vector bundle V has type (a_1, a_2, \dots, a_r) when restricting on Schubert lines where

$$a_1 = a_2 = \dots = a_k = 0 \text{ and } 0 > a_{k+1} \geq a_{k+2} \geq \dots \geq a_r.$$

Since we have

$$\begin{array}{ccc} C & \xrightarrow{ev} & G/P \\ \downarrow p & & \\ \overline{\mathcal{M}}_{0,0}(G/P, 1) & & \end{array}$$

where C is the universal bundle of $\overline{\mathcal{M}}_{0,0}(G/P, 1)$, i.e, it is $\overline{\mathcal{M}}_{0,1}(G/P, 1)$. Under our assumption, we have

$$H^0(C_x, ev^*V|_{C_x}) = H^0(\mathbb{P}^1, (\oplus^k \mathcal{O}_{\mathbb{P}^1}) \oplus (\oplus_{i=k+1}^r \mathcal{O}_{\mathbb{P}^1}(a_i))) = \mathbb{C}^k,$$

where $x \in \overline{\mathcal{M}}_{0,0}(G/P, 1)$. Since the dimension does not change when x varies, the push forward $p_* ev^*V$ is a vector bundle of rank k on $\overline{\mathcal{M}}_{0,0}(G/P, 1)$. We claim there exists a short exact sequence of vector bundles as following

$$0 \rightarrow H = p^* p_* ev^*V \rightarrow ev^*V \rightarrow Q \rightarrow 0.$$

In fact, if we restrict to C_x , then we get

$$H|_{C_x} = \mathcal{O}_{C_x}^{\oplus k} \subseteq ev^*V|_{C_x} = \mathcal{O}_{C_x}^{\oplus k} \oplus (\oplus_{i=k+1}^r \mathcal{O}_{C_x}(a_i)).$$

In particular, the quotient $Q|_{C_x}$ is a vector bundle $\oplus_{i=k+1}^r \mathcal{O}_{C_x}(a_i)$, hence, the claim is clear.

The second claim is that there exists vector bundles V_1 and V_2 on G/P such that

$$H \cong ev^*V_1 \text{ and } Q \cong ev^*V_2.$$

Since the evaluation map is smooth and $VRMT = ev^{-1}(p)$, it is enough to prove that

$$H|_{VRMT} \text{ and } Q|_{VRMT} \text{ are trivial bundles.}$$

Since the restriction of ev^* to $VRMT = ev^{-1}(p)$ is trivial, we get a short exact sequence of vector bundles as following

$$0 \rightarrow H|_{VRMT} \rightarrow \mathbb{C}^{\oplus r} \rightarrow Q|_{VRMT} \rightarrow 0.$$

Therefore, we have $Q|_{VRMT}$ and $H^\vee|_{VRMT}$ are global generated. Therefore, to prove the claim, it is sufficient to prove the first Chern Classes are vanishing, i.e.,

$$c_1(H^\vee|_{VRMT}) = c_1(Q|_{VRMT}) = 0.$$

By the Whitney formula of Chern Classes and the above short exact sequence, we have $c(H|_{VRMT}) \cdot c(Q|_{VRMT}) = a$, i.e.,

$$(c_k + c_{k-1} + \dots + 1)(\tilde{c}_{r-k} + \tilde{c}_{r-k-1} + \dots + 1) = 1,$$

where we have

$$c(H|_{VRMT}) = c_k + c_{k-1} + \dots + 1 \text{ and } c(Q|_{VRMT}) = \tilde{c}_{r-k} + \tilde{c}_{r-k-1} + \dots + 1.$$

Since c_i and \tilde{c}_j are in $CH^*(VRMT)$ and $VRMT$ has good divisibility up to degree r , expanding the above equality of Chern Classes and comparing both sides, we get $c_1(H^\vee|_{VRMT}) = c_1(Q|_{VRMT}) = 0$. We prove the claim.

Therefore, we have the following exact sequence

$$0 \rightarrow ev^*V_1 \rightarrow ev^*V \rightarrow ev^*V_2 \rightarrow 0$$

Since we know $H|_{C_x}$ is trivial, by Theorem 3.4, the vector bundle V_1 is trivial. Since $VRMT$ is rationally connected, the cohomology group $H^1(VRMT, \mathcal{O})$ is trivial. Therefore, apply the functor ev_* to this short exact sequence, we have the following short exact sequence by projection formula and vanishing of the first derived functor of ev_* ,

$$0 \rightarrow V_1 \rightarrow V \rightarrow V_2 \rightarrow 0.$$

By induction on the rank of vector bundles, it implies V_2 is splitting vector bundle. Since $H^1(G/P, L) = 0$ for any line bundle on G/P if $\dim(G/P)$ is at least 2, we have that

$$Ext^1(V_2, V_1) = H^1(G/P, (V_2)^\vee) = 0.$$

Therefore, we have $V = V_1 \oplus V_2$, it completes the proof. For $\dim(G/P) = 1$, it is \mathbb{P}^1 , the theorem is trivial. □

5. APPLICATIONS

In the paper [MOSC12], Muñoz, Roberto and Occhetta, Gianluca and Solá Conde prove uniform vector bundles of low rank on Grassmannians and quadrics are splitting in two different approach. In this section, we apply Theorem 4.2 to provide a uniform proof for Grassmannians and quadrics. We also get more examples which can not be deduced from their approach, namely, Classical Grassmannians.

Lemma 5.1. *Projective space \mathbb{P}^m has good divisibility up to degree m .*

Proof. It is obvious since we know $CH^*(\mathbb{P}^m) = \mathbb{Z}[u]/(u^{m+1})$, see [Ful98]. \square

Lemma 5.2. *Let X and Y be projective varieties. If we have*

- (1) X (.resp Y) *has good divisibility up to degree s (r),*
- (2) $CH^*(X \times Y) = CH^*(X) \otimes CH^*(Y)$,
- (3) $CH^*(X)$ and $CH^*(Y)$ *are free \mathbb{Z} -module,*

then $X \times Y$ has good divisibility up to $\min(s, t)$.

Proof. Let $A \in CH^r(X \times Y)$ (.resp $B \in CH^h(X \times Y)$) be

$$\sum_{k+l=r} a_k \otimes b_l \text{ (.resp } \sum_{p+q=h} c_p \otimes d_q)$$

where $r + h \leq \min(s, t)$ and l, k, p, q are nonnegative. Suppose $A \cdot B = 0$. If A and B are nonzero, then we can assume

- $a_{k_1} \otimes b_{l_1} \neq 0$ where $k_1 = \min\{k | a_k \otimes b_l \neq 0\}$,
- $c_{p_1} \otimes d_{q_1} \neq 0$ where $p_1 = \min\{p | c_p \otimes d_l \neq 0\}$.

It implies that $a_{k_1} \cdot c_{p_1} = 0$ or $b_{l_1} \cdot d_{q_1} = 0$. Since we have

- $k_1 + p_1 \leq r + h \leq \min(s, t) \leq s$,
- $l_1 + q_1 \leq r + h \leq \min(s, t) \leq t$,

it implies that $a_{k_1} = 0$ or $c_{p_1} = 0$ or $b_{l_1} = 0$ or $d_{q_1} = 0$. It is a contradiction. \square

Lemma 5.3. [Rei72, Theorem 1.12, Page 32] *If $m = 2n + 1$ and X is a m -dimensional quadric, then*

$$1, b, \dots, b^{n-1}, a, ba, \dots, b^{n+1}a$$

generate $H^{2i}(X, \mathbb{Z})$ and $b^n = 2a$ where $a \in H^{2n}(X, \mathbb{Z})$. In particular, the quadric X has good divisibility up to m .

Lemma 5.4. [Rei72, Theorem 1.13, Page 33] *If $m = 2n$ and X is a m -dimensional quadric, then*

$$1, c, \dots, c^{n-1}, a \text{ and } b, ca, \dots, c^n a$$

generate $H^{2i}(X, \mathbb{Z})$ and $ca = cb$ where a and $b \in H^{2n}(X, \mathbb{Z})$. Moreover,

- (1) $a^2 = b^2 = 1$ and $ab = 0$ if n is even,
- (2) $a^2 = b^2 = 0$ and $ab = 1$ if n is odd,

In particular, the quadric X has good divisibility up to $m - 1$.

Proof. Since quadrics are homogenous spaces, by Bruhat decomposition, we know $H^*(X) = CH^*(X)$, see [Ful98]. \square

Lemma 5.5. *If X is a nonsingular variety and has good divisibility up to r and V is a vector bundle of rank n on X , then $\mathbb{P}(V)$ has good divisibility up to $\min(r, n - 1)$.*

Proof. By [Ful98], we have

$$CH^*(\mathbb{P}(V)) = CH^*(X)[u]/(u^n + c_1(V)u + \dots + c_n(V)).$$

The proof is similar to the proof of Lemma 5.2. \square

Theorem 5.6. *A uniform vector bundle on X of rank at most $s(X)$ is splitting, where X and $s(X)$ is the following*

X	$s(X)$	$V.R.M.T$
\mathbb{P}^n	$n-1$	\mathbb{P}^{n-1}
\mathbb{Q}^n	$n-2$ if n is odd $n-3$ if n is even	\mathbb{Q}^{n-2}
$\mathbb{G}(k, n)$	$\min(k, n-k-1)$	$\mathbb{P}^k \times \mathbb{P}^{n-k-1}$
$\mathbb{OG}(k, 2m-1)$	$\min(k, 2m-2k-3)$	$\mathbb{P}^k \times \mathbb{Q}^{2m-2k-3}$ for $1 \leq k \leq m-2$
$\mathbb{SG}(k, 2m-1)$	$\min(m, 2m-2k-2)$	$\mathbb{P}_{\mathbb{P}^m}(\mathcal{O}(2) \oplus \mathcal{O}(1)^{2m-2k-2})$ for $1 \leq k \leq m-2$
$\mathbb{SG}(m-1, 2m-1)$	$m-1$	$v_2(\mathbb{P}^{m-1})$

Proof. By the paper [LM03] or the survey [Muñ72], we know the *VRMT* for these examples. Apply Theorem 4.2 and the lemmas in this section.

For \mathbb{P}^n , we apply Lemma 5.1.

For \mathbb{Q}^n , we apply Lemma 5.3 and 5.4.

For $\mathbb{G}(k, n)$, we apply Lemma 5.2.

For $\mathbb{OG}(k, 2m-1)$, we apply Lemma 5.2 and 5.3.

For $\mathbb{SG}(k, 2m-1)$, we apply Lemma 5.5.

For $\mathbb{SG}(m-1, 2m-1)$, we apply Lemma 5.1. \square

Question 5.7. *Classify the vector bundles on X of rank $s(X) + 1$?*

See the paper [MOSC12] for some answers.

REFERENCES

- [Bal81] Edoardo Ballico. Uniform vector bundles on quadrics. *Ann. Univ. Ferrara Sez. VII (N.S.)*, 27:135–146 (1982), 1981.
- [dJHS11] A. J. de Jong, Xuhua He, and Jason Michael Starr. Families of rationally simply connected varieties over surfaces and torsors for semisimple groups. *Publ. Math. Inst. Hautes Études Sci.*, (114):1–85, 2011.
- [FK88] Eberhard Freitag and Reinhardt Kiehl. *Étale cohomology and the Weil conjecture*, volume 13 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1988. Translated from the German by Betty S. Waterhouse and William C. Waterhouse, With an historical introduction by J. A. Dieudonné.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.

- [Hwa01] Jun-Muk Hwang. Geometry of minimal rational curves on Fano manifolds. In *School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000)*, volume 6 of *ICTP Lect. Notes*, pages 335–393. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001.
- [IJS10] I. Biswas, J. Pedro, and P. Dos Santos. Triviality criterion for vector bundles over rationally connected varieties. *Preprint*, 2010.
- [Kol08] János Kollár. Looking for rational curves on cubic hypersurfaces. In *Higher-dimensional geometry over finite fields*, volume 16 of *NATO Sci. Peace Secur. Ser. D Inf. Commun. Secur.*, pages 92–122. IOS, Amsterdam, 2008. Notes by Ulrich Derenthal.
- [LM03] Joseph M. Landsberg and Laurent Manivel. On the projective geometry of rational homogeneous varieties. *Comment. Math. Helv.*, 78(1):65–100, 2003.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.
- [MOSC12] Roberto Muñoz, Gianluca Occhetta, and Luis E. Solá Conde. Uniform vector bundles on fano manifolds and applications. *J. Reine Angew. Math.*, 664:141–162, 2012.
- [Muñ72] Roberto Muñoz. *Uniform vector bundles on Fano manifolds*. 1972. Survey, KIAS, Seoul, April 6th, 2011.
- [Ols07] Martin Olsson. Sheaves on Artin stacks. *J. Reine Angew. Math.*, 603:55–112, 2007.
- [Rei72] Miles Reid. *The Complete Intersection of two or more Quadrics*. 1972. Thesis (Ph.D.)—Trinity College, Cambridge.
- [Sat76] Ei-ichi Sato. Uniform vector bundles on a projective space. *J. Math. Soc. Japan*, 28(1):123–132, 1976.
- [VdV72] A. Van de Ven. On uniform vector bundles. *Math. Ann.*, 195:245–248, 1972.
- [Zhu10] Yi Zhu. Homogeneous fibrations over surfaces. *Preprint*, 2010.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10025
E-mail address: `pan@math.columbia.edu`