

Special functions and reversible three-term recurrence formula (R3TRF)

YOON SEOK CHOUN ^{1 2}
Baruch College, The City University of New York,
Natural Science Department, A506,
17 Lexington Avenue,
New York, NY 10010

October 4, 2018

¹The first series was ‘Special functions and three-term recurrence formula (3TRF).’ It is available as arXiv. This book is a second series.

²Yoon.Choun@baruch.cuny.edu; ychoun@gc.cuny.edu; ychoun@gmail.com

Contents

1	Generalization of the reversible three-term recurrence formula (R3TRF) and its applications	5
1.1	Introduction	6
1.2	Mathieu equation, Frobenius method and reversible three term recurrence relation	7
1.3	Infinite series	8
1.4	Polynomial which makes A_n term terminated	14
1.5	Summary	24
	Bibliography	24
2	Heun function using reversible three-term recurrence formula	27
2.1	Introduction	27
2.2	Power series	30
2.2.1	Polynomial of type 2	30
2.2.2	Infinite series	34
2.3	Integral Formalism	38
2.3.1	Polynomial of type 2	38
2.3.2	Infinite series	46

2.4	Summary	52
Appendices		53
.1	Power series expansion of 192 Heun functions	55
.2	Integral formalism of 192 Heun functions	75
	Bibliography	92
3	The generating function for Heun polynomial using reversible three-term recurrence formula	95
3.1	Introduction	95
3.2	Generating function for Heun polynomial of type 2	98
3.3	Summary	117
Appendices		119
.1	Generating functions for 192 Heun polynomials of type 2	121
	Bibliography	139
4	Confluent Heun function using three-term recurrence formula	143
4.1	Introduction	143
4.1.1	Non-symmetrical canonical form of the CHE	143
4.1.2	Generalized spheroidal equation (GSE)	144
4.1.3	Generalized spheroidal equation in the Leaver version	147
4.2	CHF with a regular singular point at zero	147
4.2.1	Power series	149
	Polynomial of type 1	149
	Infinite series	153

4.2.2	Integral formalism	155
	Polynomial of type 1	156
	Infinite series	162
4.2.3	Generating function for the CHP of type 1	168
4.3	Summary	180
	Bibliography	182
5	Confluent Heun function using reversible three-term recurrence formula	185
5.1	Non-symmetrical canonical form of Confluent Heun Equation	185
5.2	CHF with a regular singular point at 0	187
5.2.1	Power series	188
	Polynomial of type 2	188
	Infinite series	193
5.2.2	Integral formalism	196
	Polynomial of type 2	196
	Infinite series	203
5.2.3	Generating function for the CHP of type 2	208
5.3	Summary	225
	Bibliography	226
6	Grand Confluent Hypergeometric function using reversible three-term recurrence formula	229
6.1	Introduction	229
6.2	GCH equation about regular singular point at zero	231
6.2.1	Power series	232

Polynomial of type 2	232
Infinite series	235
6.2.2 Integral formalism	238
Polynomial of type 2	238
Infinite series	244
6.2.3 Generating function for the GCH polynomial of type 2	250
6.3 GCH equation about irregular singular point at infinity	260
6.3.1 Power series for polynomial of type 1	262
6.3.2 Integral formalism for polynomial of type 1	264
6.3.3 Generating function of the GCH polynomial of type 1	271
6.4 Summary	282
Bibliography	283
7 Mathieu function using reversible three-term recurrence formula	289
7.1 Introduction	289
7.2 Mathieu equation about regular singular point at zero	293
7.2.1 Power series	293
Polynomial of type 2	293
Infinite series	297
7.2.2 Integral formalism	300
Polynomial of type 2	301
Infinite series	307
7.2.3 Generating function for the Mathieu polynomial of type 2	313
7.3 Mathieu equation about regular singular point at one	327

7.3.1	Mathieu equation about $x = 1$ using 3TRF	328
	Power series for infinite series	328
	Integral formalism for infinite series	329
7.3.2	Mathieu equation about $x = 1$ using R3TRF	329
	Power series	329
	Integral formalism	332
	Generating function for the Mathieu polynomial of type 2	335
7.4	Mathieu equation about irregular singular point at infinity	338
7.4.1	Mathieu equation about $x = \infty$	338
7.4.2	Mathieu equation about $\zeta = \infty$	339
7.5	Summary	339
	Bibliography	341
8	Lame function in the algebraic form using reversible three-term recurrence formula	347
8.1	Introduction	347
8.2	Power series	350
8.2.1	Polynomial of type 2	350
8.2.2	Infinite series	355
8.3	Integral formalism	358
8.3.1	Polynomial of type 2	359
8.3.2	Infinite series	366
8.4	Generating function for the Lame polynomial of type 2	371
8.5	Summary	388
	Bibliography	389

9	Lame function in Weierstrass's form using reversible three-term recurrence formula	393
9.1	Introduction	394
9.2	Power series	397
9.2.1	Polynomial of type 2	397
9.2.2	Infinite series	401
9.3	Integral formalism	403
9.3.1	Polynomial of type 2	405
9.3.2	Infinite series	407
9.4	Generating function for the Lame polynomial of type 2	409
9.5	Summary	415
	Appendices	419
.1	Power series	422
.2	Integral representation	440
.3	Generating function for the type 2 polynomial	458
	Bibliography	475

Preface

For the past 350 years, we only have constructed the power series expansion in closed forms using the two term recurrence relation in linear differential equation. However, currently the analytic solution and its integral forms for more than three term cases are unknown.

For the three term case, we assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda}$$

On the above, λ is an indicial root, $y(x)$ is its analytic solution for the linear differential equation and x is an independent variable for its solution. When we substitute the above equation into any linear ordinary differential equation having three different coefficients in their power series expansion, the recurrence relation for frobenius solution is

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1$$

where

$$c_1 = A_0 c_0$$

On the above, A_n and B_n are themselves polynomials of degree n : for the second ordinary differential equations, numerator or/and denominator of A_n and B_n is/are usually equal or less than polynomials of degrees 2. For the case of three term recurrence in its power series expansion, all other differential equations having no analytic solution in closed forms can be described as in the above recurrence relation equation.¹

In the first series “Special functions and three term recurrence formula (3TRF)”, I showed how to construct the power series expansions in closed forms, its integral representations and its generating functions for linear ordinary differential equations

¹For more details, look at arXiv:1303.0806 in “Special functions and three term recurrence formula (3TRF).”

(infinite series and polynomial which makes B_n term terminated) such as Heun, Mathieu, Lamé, Grand Confluent Hypergeometric (GCH) equations.² There are three types polynomial in their three term recurrence relation: (1) polynomial which makes B_n term terminated, (2) polynomial which makes A_n term terminated, (3) polynomial which makes A_n and B_n terms terminated at same time. However, by using 3TRF, there are no way to construct the power series in closed forms indeed its integral representation for the type 2 and 3 polynomials.

In this series natural numbers $\mathbb{N}_0$ means $\{0, 1, 2, 3, \dots\}$. Pochhammer symbol $(x)_n$ is used to represent the rising factorial: $(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)}$. And I will show how to obtain the formula for the type 2 polynomial and infinite series of its power series expansion. In the future series I will show you for the type 3 case; the power series, integral formalism and generating function such as Heun, Confluent Heun, Lamé and GCH equations will be constructed analytically.

Structure of book

In chapter 1, I will generalize the three term recurrence relation in linear differential equation in a backward for the infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's.

In chapters 2–9, I will apply reversible three term recurrence formula to (1) the power series expansion in closed forms, (2) its integral representation and (3) generating functions for Heun, Confluent Heun, GCH, Lamé and Mathieu equations that consist of three term recursion relation for the infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's. The short descriptions of all 9 chapters are as follows.

Chapter 1. “Generalization of the reversible three-term recurrence formula and its applications”– generalize the three term recurrence relation in linear differential equation in a backward. Obtain the exact solution of the three term recurrence relation for infinite series and polynomial which makes A_n term terminated

Chapter 2. “Heun function using reversible three-term recurrence formula”– apply the reversible three term recurrence formula to the power series expansion in closed forms and its integral forms of Heun functions (infinite series and polynomial which makes A_n

²These 4 linear differential equations have three different coefficients in their power series expansion. Bi-confluent Heun equation is the special case of GCH equation. Look at arXiv:1303.0813 and arXiv:1303.0819 in “Special functions and three term recurrence formula (3TRF).”

term terminated) including all higher terms of B_n 's.

Chapter 3. “The generating function for Heun polynomial using reversible three-term recurrence formula”– apply the reversible three term recurrence formula and derive generating function for Heun polynomial which makes A_n term terminated including all higher terms of B_n 's.

Chapter 4. “Confluent Heun function using three-term recurrence formula”– (1) apply the three term recurrence formula to the power series expansion in closed forms and its integral forms of the Confluent Heun functions (infinite series and polynomial which makes B_n term terminated) including all higher terms of A_n 's. (2) apply the three term recurrence formula and derive the generating function for the Confluent Heun polynomial which makes B_n term terminated including all higher terms of A_n 's.

Chapter 5. “Confluent Heun function using reversible three-term recurrence formula”– (1) apply the reversible three term recurrence formula to the power series expansion in closed forms and its integral forms of the Confluent Heun functions (infinite series and polynomial which makes A_n term terminated) including all higher terms of B_n 's. (2) apply three term recurrence formula and derive the generating function for the Confluent Heun polynomial which makes A_n term terminated including all higher terms of B_n 's.

Chapter 6. “Grand Confluent Hypergeometric function using reversible three-term recurrence formula”– (1) apply the reversible three term recurrence formula and formulate the analytic solution for the power series expansion and its integral forms of grand confluent hypergeometric function (infinite series and polynomial which makes A_n term terminated) including all higher terms of B_n 's. Replacing μ and $\varepsilon\omega$ by 1 and $-q$ transforms the grand confluent hypergeometric function into the Biconfluent Heun function. (2) apply the reversible three term recurrence formula and construct the generating function for grand confluent hypergeometric polynomial which makes A_n term terminated including all higher terms of B_n 's.

Chapter 7. “Mathieu function using reversible three-term recurrence formula”– apply the reversible three term recurrence formula, and analyze the power series expansion in closed forms of Mathieu function and its integral forms (infinite series and polynomial which makes A_n term terminated) including all higher terms of B_n 's. Construct the generating function for Mathieu polynomial which makes A_n term terminated.

Chapter 8. “Lame function in the algebraic form using reversible three-term recurrence formula”– (1) apply the reversible three term recurrence formula, and analyze the power series expansion in closed forms and integral forms of Lamé function in the algebraic form (infinite series and polynomial which makes A_n term terminated) including all higher

terms of B_n 's. (2) apply the reversible three term recurrence formula and construct the generating function for Lamé polynomial which makes A_n term terminated in the algebraic form including all higher terms of B_n 's.

Chapter 9. “Lamé function in Weierstrass’s form using reversible three-term recurrence formula” – (1) apply the reversible three term recurrence formula, and analyze the power series expansion in closed forms of Lamé function in Weierstrass’s form and its integral forms (infinite series and polynomial which makes A_n term terminated) including all higher terms of B_n 's. (2) apply the reversible three term recurrence formula and construct the generating function for Lamé polynomial which makes A_n term terminated in Weierstrass’s form including all higher terms of B_n 's.

Acknowledgements

I thank Bogdan Nicolescu. The endless discussions I had with him were of great joy.

Chapter 1

Generalization of the reversible three-term recurrence formula (R3TRF) and its applications

In “Generalization of the three-term recurrence formula and its applications” [1], I generalize the three term recurrence relation in a linear differential equation for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n ’s.¹

In this chapter, I generalize the three term recurrence relation in a linear ordinary differential equation in a backward for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n ’s.²

The rest of chapters in this series show how to apply reversible three-term recurrence formula (R3TRF) into linear differential equations having three term recursion relations and go on producing analytic solutions of some of the well known special functions including: Heun, Confluent Heun, GCH (Biconfluent Heun), Lamé and Mathieu equations.

¹“ higher terms of A_n ’s” means at least two terms of A_n ’s.

²“ higher terms of B_n ’s” means at least two terms of B_n ’s.

1.1 Introduction

The history of linear differential equations is over 350 years. By using Frobenius method and putting the power series expansion into linear differential equations, the recursive relation of coefficients starts to appear. There can be between two and infinity number of coefficients in the recurrence relation in the power series expansion. During this period mathematicians developed analytic solutions of only two term recursion relation in closed forms. Currently the analytic solution of three term recurrence relation is unknown.

In “Generalization of the three-term recurrence formula and its applications” [1], I show how to obtain the Frobenius solutions for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n ’s using three term recurrence formula (3TRF) in a linear ordinary differential equation having a relation between three different coefficients. The sequence c_n consists of combinations A_n and B_n . I observed the term inside parenthesis of sequence c_n which includes zero term of A_n ’s, one term of A_n ’s, two terms of A_n ’s, three terms of A_n ’s, etc in order to construct general expressions of the power series in closed forms for infinite series and polynomial which makes B_n term terminated.

Any problems having three term recurrence relation in their linear differential equations in modern physics (E & M, Newtonian mechanics, quantum mechanic, QCD, supersymmetric field theories, string theories, general relativity, etc) mostly involve the quantum mechanical energy in A_n or B_n term. In general, the energy is quantized in quantum mechanical points of a view. The power series expansion of a eigen-function is the polynomial. In Ref.[1] I construct the Frobenius solution in closed forms of three term recurrence relation in a linear differential equation for polynomial which makes B_n term terminated. Also we need to derive the analytic solution of three term recurrence relation in a linear differential equation for polynomial which makes A_n term terminated. Because in some modern physics problems the eigenvalue (quantum mechanical energy) is involved in A_n term: Mathieu equation, Heun equation and its confluent forms (black hole problems [11, 12]) , Lamé equation in the Weierstrass’s form [13, 23, 15, 16], etc.

In this chapter I construct three term recurrence relation in a linear ordinary differential equation in a backward to obtain the analytic solution of the three term recurrence relation for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n ’s. I designate the mathematical formulas of infinite series and polynomial which makes A_n terminated of a linear differential equation having three different coefficients in its power series expansion as “reversible three-term recurrence formula (R3TRF)” that will be expressed below.

In chapters 2–9, using the R3TRF, I show how to obtain analytic solutions of infinite

series and polynomial which makes A_n term terminated (1) in the power series expansion, (2) in the integral formalism and (3) in the generating function for any linear ordinary differential equations having three term recurrence relation.

1.2 Mathieu equation, Frobenius method and reversible three term recurrence relation

In some rare cases, there are no such solutions of the power series expansion for polynomial which makes B_n term terminated in a linear ordinary differential equation having three different coefficients. One of examples is Mathieu differential equation in the algebraic form. Mathieu functions[2], is an example of three term recurrence relation appears in physical problems involving elliptical shapes[4] or periodic potentials, were introduced by Mathieu (1868)[11] when he investigated the vibrating elliptical drumhead.

Mathieu ordinary differential equation is of Fuchsian types with the two regular and one irregular singularities. In contrast, Heun equation of Fuchsian types has the three regular and one irregular singularities [1]. Heun equation has the four kind of confluent forms: (1) Confluent Heun (two regular and one irregular singularities) [2, 21], (2) Doubly confluent Heun (two irregular singularities), (3) Biconfluent Heun (one regular and one irregular singularities) [17], (4) Triconfluent Heun equations (one irregular singularity). Mathieu equation in algebraic forms is also derived from the Confluent Heun equation by changing all coefficients $\delta = \gamma = \frac{1}{2}$, $\beta = 0$, $\alpha = \frac{q}{0}$ and $q = \frac{\lambda+2q}{4}$: see chapters 4 and 5.

Mathieu equation is

$$\frac{d^2 y}{dz^2} + (\lambda - 2q \cos 2z) y = 0 \quad (1.2.1)$$

where λ and q are parameters. This is an equation with periodic-function coefficient. Mathieu equation also can be described in algebraic forms putting $x = \cos^2 z$:

$$4x(1-x)\frac{d^2 y}{dx^2} + 2(1-2x)\frac{dy}{dx} + (\lambda + 2q - 4qx)y = 0 \quad (1.2.2)$$

This equation has two regular singularities: $x = 0$ and $x = 1$; the other singularity $x = \infty$ is irregular. Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\nu} \quad \text{where } \nu = \text{indicial root} \quad (1.2.3)$$

Plug (1.2.3) into (1.2.2):

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (1.2.4)$$

where

$$A_n = \frac{4(n + \nu)^2 - (\lambda + 2q)}{2(n + 1 + \nu)(2(n + \nu) + 1)} \quad (1.2.5a)$$

$$B_n = \frac{4q}{2(n + 1 + \nu)(2(n + \nu) + 1)} \quad (1.2.5b)$$

$$c_1 = A_0 c_0 \quad (1.2.5c)$$

We have two indicial roots which are $\nu = 0$ and $\frac{1}{2}$. As we see (1.2.5b), there is no way to make B_n term terminate at certain index value of n , because the numerator of (1.2.5b) is just consist of constant q parameter.³ So there are only two kind of power series expansions: infinite series and the polynomial which makes A_n term terminated.

In “The power series expansion of Mathieu function and its integral formalism” [3], I construct the analytic solution of Mathieu equation for infinite series by using 3TRF [1]. In this chapter I generalize the three term recurrence relation in a linear differential equation in a backward in order to construct the analytic solution for polynomial which makes A_n term terminated.

1.3 Infinite series

Assume that

$$c_1 = A_0 c_0 \quad (1.3.1)$$

(1.3.1) is a necessary boundary condition. The three term recurrence relation in all differential equations having no analytic solution in closed forms follow (1.3.1).

$$\prod_{n=a_i}^{a_i-1} B_n = 1 \quad \text{where } a_i \text{ is positive integer including } 0 \quad (1.3.2)$$

(1.3.2) is also a necessary condition. Every differential equations having no analytic solution in closed forms also take satisfied with (1.3.2).

My definition of $B_{i,j,k,l}$ refer to $B_i B_j B_k B_l$. Also, $A_{i,j,k,l}$ refer to $A_i A_j A_k A_l$. For

³Whenever index n increase in (1.2.5b), the B_n term never terminated with a fixed constant parameter q .

$n = 0, 1, 2, 3, \dots$, (1.2.4) gives

$$\begin{aligned}
c_0 & \\
c_1 &= A_0 c_0 \\
c_2 &= (A_{0,1} + B_1) c_0 \\
c_3 &= (A_{0,1,2} + B_1 A_2 + B_2 A_0) c_0 \\
c_4 &= (A_{0,1,2,3} + B_1 A_{2,3} + B_2 A_{0,3} + B_3 A_{0,1} + B_{1,3}) c_0 \\
c_5 &= (A_{0,1,2,3,4} + B_1 A_{2,3,4} + B_2 A_{0,3,4} + B_3 A_{0,1,4} + B_4 A_{0,1,2} \\
&\quad + B_{1,3} A_4 + B_{1,4} A_2 + B_{2,4} A_0) c_0 \\
c_6 &= (A_{0,1,2,3,4,5} + B_1 A_{2,3,4,5} + B_2 A_{0,3,4,5} + B_3 A_{0,1,4,5} + B_4 A_{0,1,2,5} + B_5 A_{0,1,2,3} \\
&\quad + B_{1,3} A_{4,5} + B_{1,4} A_{2,5} + B_{2,4} A_{0,5} + B_{1,5} A_{2,3} + B_{2,5} A_{0,3} + B_{3,5} A_{0,1} + B_{1,3,5}) c_0 \\
c_7 &= (A_{0,1,2,3,4,5,6} + B_1 A_{2,3,4,5,6} + B_2 A_{0,3,4,5,6} + B_3 A_{0,1,4,5,6} + B_4 A_{0,1,2,5,6} + B_5 A_{0,1,2,3,6} \\
&\quad + B_6 A_{0,1,2,3,4} + B_{1,4} A_{2,5,6} + B_{1,3} A_{4,5,6} + B_{2,4} A_{0,5,6} + B_{1,5} A_{2,3,6} + B_{2,5} A_{0,3,6} \\
&\quad + B_{3,5} A_{0,1,6} + B_{1,6} A_{2,3,4} + B_{2,6} A_{0,3,4} + B_{3,6} A_{0,1,4} + B_{4,6} A_{0,1,2} \\
&\quad + B_{1,3,6} A_4 + B_{1,3,5} A_6 + B_{1,4,6} A_2 + B_{2,4,6} A_0) c_0 \\
c_8 &= (A_{0,1,2,3,4,5,6,7} + B_1 A_{2,3,4,5,6,7} + B_2 A_{0,3,4,5,6,7} + B_3 A_{0,1,4,5,6,7} + B_4 A_{0,1,2,5,6,7} + B_5 A_{0,1,2,3,6,7} \\
&\quad + B_6 A_{0,1,2,3,4,7} + B_7 A_{0,1,2,3,4,5} + B_{1,4} A_{2,5,6,7} + B_{2,4} A_{0,5,6,7} + B_{1,5} A_{2,3,6,7} + B_{1,3} A_{4,5,6,7} \\
&\quad + B_{2,5} A_{0,3,6,7} + B_{3,5} A_{0,1,6,7} + B_{1,6} A_{2,3,4,7} + B_{2,6} A_{0,3,4,7} + B_{3,6} A_{0,1,4,7} + B_{4,6} A_{0,1,2,7} \\
&\quad + B_{1,7} A_{2,3,4,5} + B_{3,7} A_{0,1,4,5} + B_{2,7} A_{0,3,4,5} + B_{4,7} A_{0,1,2,5} + B_{5,7} A_{0,1,2,3} \\
&\quad + B_{1,4,6} A_{2,7} + B_{2,4,6} A_{0,7} + B_{1,3,7} A_{4,5} + B_{1,4,7} A_{2,5} + B_{1,3,6} A_{4,7} + B_{2,4,7} A_{0,5} \\
&\quad + B_{1,3,5} A_{6,7} + B_{1,5,7} A_{2,3} + B_{2,5,7} A_{0,3} + B_{3,5,7} A_{0,1} + B_{1,3,5,7}) c_0 \\
&\vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.3.3}$$

In Ref.[1] I construct the Frobenius solution for infinite series and polynomial which makes B_n term terminated in three term recurrence relation of linear ordinary differential equation by letting A_n in sequence c_n is the leading term in the analytic function $y(x)$. In this chapter I construct the power series for infinite series and polynomial which makes A_n term terminated in three term recurrence relation by letting B_n in the sequence c_n is the leading term in $y(x)$.

In (1.3.3) the number of individual sequence c_n follows Fibonacci sequence: 1,1,2,3,5,8,13,21,34,55,... The sequence c_n consists of combinations A_n and B_n in (1.3.3). First observe the term inside parentheses of sequence c_n which does not include

any B_n 's: c_n with every index $(c_0, c_1, c_2, \dots)$.

(a) Zero term of B_n 's

$$\begin{aligned}
 c_0 & \\
 c_2 &= A_0 c_0 \\
 c_4 &= A_{0,1} c_0 \\
 c_6 &= A_{0,1,2} c_0 \\
 c_8 &= A_{0,1,2,3} c_0 \\
 \vdots & \quad \vdots
 \end{aligned} \tag{1.3.4}$$

When a function $y(x)$, analytic at $x = 0$, is expanded in a power series, we write

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} = \sum_{m=0}^{\infty} y_m(x) = y_0(x) + y_1(x) + y_2(x) + \dots \tag{1.3.5}$$

where

$$y_m(x) = \sum_{l=0}^{\infty} c_l^m x^{l+\lambda} \tag{1.3.6}$$

λ is the indicial root. $y_m(x)$ is sub-power series that have sequence c_n including m term of B_n 's in (1.3.3). For example $y_0(x)$ has sequences c_n including zero term of B_n 's in (1.3.3), $y_1(x)$ has sequences c_n including one term of B_n 's in (1.3.3), $y_2(x)$ has sequences c_n including two term of B_n 's in (1.3.3), etc. Substitute (1.3.4) in (1.3.6) putting $m = 0$.

$$y_0(x) = c_0 \sum_{n=0}^{\infty} \left\{ \prod_{i_1=0}^{n-1} A_{i_1} \right\} x^{n+\lambda} \tag{1.3.7}$$

Observe the terms inside parentheses of sequence c_n which include one term of B_n 's in (1.3.3): c_n with every index except c_0 and c_1 ($c_2, c_3, c_4, \dots$).

(b) One term of B_n 's

$$\begin{aligned}
c_2 &= B_1 c_0 \\
c_3 &= \left\{ B_1 \cdot 1 \cdot \left(\frac{A_2}{1} \right) + B_2 A_0 \left(\frac{A_2}{A_2} \right) \right\} c_0 \\
c_4 &= \left\{ B_1 \cdot 1 \cdot \left(\frac{A_{2,3}}{1} \right) + B_2 A_0 \left(\frac{A_{2,3}}{A_2} \right) + B_3 A_{0,1} \left(\frac{A_{2,3}}{A_{2,3}} \right) \right\} c_0 \\
c_5 &= \left\{ B_1 \cdot 1 \cdot \left(\frac{A_{2,3,4}}{1} \right) + B_2 A_0 \left(\frac{A_{2,3,4}}{A_2} \right) + B_3 A_{0,1} \left(\frac{A_{2,3,4}}{A_{2,3}} \right) + B_4 A_{0,1,2} \left(\frac{A_{2,3,4}}{A_{2,3,4}} \right) \right\} c_0 \\
c_6 &= \left\{ B_1 \cdot 1 \cdot \left(\frac{A_{2,3,4,5}}{1} \right) + B_2 A_0 \left(\frac{A_{2,3,4,5}}{A_2} \right) + B_3 A_{0,1} \left(\frac{A_{2,3,4,5}}{A_{2,3}} \right) + B_4 A_{0,1,2} \left(\frac{A_{2,3,4,5}}{A_{2,3,4}} \right) \right. \\
&\quad \left. + B_5 A_{0,1,2,3} \left(\frac{A_{2,3,4,5}}{A_{2,3,4,5}} \right) \right\} c_0 \\
c_7 &= \left\{ B_1 \cdot 1 \cdot \left(\frac{A_{2,3,4,5,6}}{1} \right) + B_2 A_0 \left(\frac{A_{2,3,4,5,6}}{A_2} \right) + B_3 A_{0,1} \left(\frac{A_{2,3,4,5,6}}{A_{2,3}} \right) + B_4 A_{0,1,2} \left(\frac{A_{2,3,4,5,6}}{A_{2,3,4}} \right) \right. \\
&\quad \left. + B_5 A_{0,1,2,3} \left(\frac{A_{2,3,4,5,6}}{A_{2,3,4,5}} \right) + B_6 A_{0,1,2,3,4} \left(\frac{A_{2,3,4,5,6}}{A_{2,3,4,5,6}} \right) \right\} c_0 \\
&\vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.3.8}$$

(1.3.8) is simply

$$c_{n+2} = \sum_{i_1=0}^n \left\{ B_{i_1+1} \prod_{i_2=0}^{i_1-1} A_{i_2} \prod_{i_3=i_1}^{n-1} A_{i_3+2} \right\} c_0 \tag{1.3.9}$$

Substitute (1.3.9) in (1.3.6) putting $m = 1$.

$$y_1(x) = c_0 \sum_{n=0}^{\infty} \left\{ \sum_{i_1=0}^n \left\{ B_{i_1+1} \prod_{i_2=0}^{i_1-1} A_{i_2} \prod_{i_3=i_1}^{n-1} A_{i_3+2} \right\} \right\} x^{n+2+\lambda} \tag{1.3.10}$$

Observe the terms inside parentheses of sequence c_n which include two terms of B_n 's in (1.3.3): c_n with every index except c_0 – c_3 (c_4 , c_5 , $c_6, \dots$).

(c) Two terms of B_n 's

$$\begin{aligned}
c_4 &= B_{1,3}c_0 \\
c_5 &= \left\{ B_1 \cdot 1 \cdot \left\{ B_3 \left(\frac{1}{1} \right) \left(\frac{A_4}{1} \right) + B_4 \left(\frac{A_2}{1} \right) \left(\frac{A_4}{A_4} \right) \right\} + B_2 A_0 \left\{ B_4 \left(\frac{A_2}{A_2} \right) \left(\frac{A_4}{A_4} \right) \right\} \right\} c_0 \\
c_6 &= \left\{ B_1 \cdot 1 \cdot \left\{ B_3 \left(\frac{1}{1} \right) \left(\frac{A_{4,5}}{1} \right) + B_4 \left(\frac{A_2}{1} \right) \left(\frac{A_{4,5}}{A_4} \right) + B_5 \left(\frac{A_{2,3}}{1} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \right\} \right. \\
&\quad \left. + B_2 A_0 \left\{ B_4 \left(\frac{A_2}{A_2} \right) \left(\frac{A_{4,5}}{A_4} \right) + B_5 \left(\frac{A_{2,3}}{A_2} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \right\} + B_3 A_{0,1} \left\{ B_5 \left(\frac{A_{2,3}}{A_{2,3}} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \right\} \right\} c_0 \\
c_7 &= \left\{ B_1 \cdot 1 \cdot \left\{ B_3 \left(\frac{1}{1} \right) \left(\frac{B_{4,5,6}}{1} \right) + B_4 \left(\frac{A_2}{1} \right) \left(\frac{A_{4,5,6}}{A_4} \right) + B_5 \left(\frac{A_{2,3}}{1} \right) \left(\frac{A_{4,5,6}}{A_{4,5}} \right) \right. \right. \\
&\quad \left. \left. + B_6 \left(\frac{A_{2,3,4}}{1} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \right\} \right. \\
&\quad + B_2 A_0 \left\{ B_4 \left(\frac{A_2}{A_2} \right) \left(\frac{A_{4,5,6}}{A_4} \right) + B_5 \left(\frac{A_{2,3}}{A_2} \right) \left(\frac{A_{4,5,6}}{A_{4,5}} \right) + B_6 \left(\frac{A_{2,3,4}}{A_2} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \right\} \\
&\quad + B_3 A_{0,1} \left\{ B_5 \left(\frac{A_{2,3}}{A_{2,3}} \right) \left(\frac{A_{4,5,6}}{A_{4,5}} \right) + B_6 \left(\frac{A_{2,3,4}}{A_{2,3}} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \right\} \\
&\quad \left. + B_4 A_{0,1,2} \left\{ B_6 \left(\frac{A_{2,3,4}}{A_{2,3,4}} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \right\} \right\} c_0 \\
&\quad \vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.3.11}$$

(1.3.11) is simply

$$c_{n+4} = \sum_{i_1=0}^n \left\{ B_{i_1+1} \sum_{i_2=i_1}^n \left\{ B_{i_2+3} \prod_{i_3=0}^{i_1-1} A_{i_3} \prod_{i_4=i_1}^{i_2-1} A_{i_4+2} \prod_{i_5=i_2}^{n-1} A_{i_5+4} \right\} \right\} c_0 \tag{1.3.12}$$

Substitute (1.3.12) in (1.3.6) putting $m = 2$.

$$y_2(x) = c_0 \sum_{n=0}^{\infty} \left\{ \sum_{i_1=0}^n \left\{ B_{i_1+1} \sum_{i_2=i_1}^n \left\{ B_{i_2+3} \prod_{i_3=0}^{i_1-1} A_{i_3} \prod_{i_4=i_1}^{i_2-1} A_{i_4+2} \prod_{i_5=i_2}^{n-1} A_{i_5+4} \right\} \right\} \right\} x^{n+4+\lambda} \tag{1.3.13}$$

Observe the terms inside parentheses of sequence c_n which include three terms of B_n 's in (1.3.3): c_n with every index except c_0 - c_5 ($c_6, c_7, c_8, \dots$).

(d) Three terms of B_n 's

$$c_6 = B_{1,3,5} c_0$$

$$c_7 = \left\{ B_1 \left\{ B_3 \cdot 1 \cdot \left[B_5 \left(\frac{1}{1} \right) \left(\frac{1}{1} \right) \left(\frac{A_6}{1} \right) + B_6 \left(\frac{1}{1} \right) \left(\frac{A_4}{1} \right) \left(\frac{A_6}{A_6} \right) \right] \right. \right. \\ \left. \left. + B_4 \cdot 1 \cdot \left[B_6 \left(\frac{A_2}{1} \right) \left(\frac{A_4}{A_4} \right) \left(\frac{A_6}{A_6} \right) \right] \right\} + B_2 \left\{ B_4 A_0 \left[B_6 \left(\frac{A_2}{A_2} \right) \left(\frac{A_4}{A_4} \right) \left(\frac{A_6}{A_6} \right) \right] \right\} \right\} c_0$$

$$c_8 = \left\{ B_1 \left\{ B_3 \cdot 1 \cdot \left[B_5 \left(\frac{1}{1} \right) \left(\frac{1}{1} \right) \left(\frac{A_{6,7}}{1} \right) + B_6 \left(\frac{1}{1} \right) \left(\frac{A_4}{1} \right) \left(\frac{A_{6,7}}{A_6} \right) + B_7 \left(\frac{1}{1} \right) \left(\frac{A_{4,5}}{1} \right) \left(\frac{A_{6,7}}{A_{6,7}} \right) \right] \right. \right. \\ \left. \left. + B_4 \cdot 1 \cdot \left[B_6 \left(\frac{A_2}{1} \right) \left(\frac{A_4}{A_4} \right) \left(\frac{A_{6,7}}{A_6} \right) + B_7 \left(\frac{A_2}{1} \right) \left(\frac{A_{4,5}}{A_4} \right) \left(\frac{A_{6,7}}{A_{6,7}} \right) \right] + B_5 \cdot 1 \cdot \left[B_7 \left(\frac{A_{2,3}}{1} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \left(\frac{A_{6,7}}{A_{6,7}} \right) \right] \right\} \right. \\ \left. + B_2 \left\{ B_4 A_0 \left[B_6 \left(\frac{A_2}{A_2} \right) \left(\frac{A_4}{A_4} \right) \left(\frac{A_{6,7}}{A_6} \right) + B_7 \left(\frac{A_2}{A_2} \right) \left(\frac{A_{4,5}}{A_4} \right) \left(\frac{A_{6,7}}{A_{6,7}} \right) \right] \right. \right. \\ \left. \left. + B_5 A_0 \left[B_7 \left(\frac{A_{2,3}}{A_2} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \left(\frac{A_{6,7}}{A_{6,7}} \right) \right] \right\} + B_3 \left\{ B_5 A_{0,1} \left[B_7 \left(\frac{A_{2,3}}{A_{2,3}} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \left(\frac{A_{6,7}}{A_{6,7}} \right) \right] \right\} \right\} c_0$$

$$c_9 = \left\{ B_1 \left\{ B_3 \cdot 1 \cdot \left[B_5 \left(\frac{1}{1} \right) \left(\frac{1}{1} \right) \left(\frac{A_{6,7,8}}{1} \right) + B_6 \left(\frac{1}{1} \right) \left(\frac{A_4}{1} \right) \left(\frac{A_{6,7,8}}{A_6} \right) + B_7 \left(\frac{1}{1} \right) \left(\frac{A_{4,5}}{1} \right) \left(\frac{A_{6,7,8}}{A_{6,7}} \right) + B_8 \left(\frac{1}{1} \right) \left(\frac{A_{4,5,6}}{1} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right. \right. \\ \left. \left. + B_4 \cdot 1 \cdot \left[B_6 \left(\frac{A_2}{1} \right) \left(\frac{A_4}{A_4} \right) \left(\frac{A_{6,7,8}}{A_6} \right) + B_7 \left(\frac{A_2}{1} \right) \left(\frac{A_{4,5}}{A_4} \right) \left(\frac{A_{6,7,8}}{A_{6,7}} \right) + B_8 \left(\frac{A_2}{1} \right) \left(\frac{A_{4,5,6}}{A_4} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right. \right. \\ \left. \left. + B_5 \cdot 1 \cdot \left[B_7 \left(\frac{A_{2,3}}{1} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \left(\frac{A_{6,7,8}}{A_{6,7}} \right) + B_8 \left(\frac{A_{2,3}}{1} \right) \left(\frac{A_{4,5,6}}{A_{4,5}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] + B_6 \cdot 1 \cdot \left[B_8 \left(\frac{A_{2,3,4}}{1} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right\} \right. \\ \left. + B_2 \left\{ B_4 A_0 \left[B_6 \left(\frac{A_2}{A_2} \right) \left(\frac{A_4}{A_4} \right) \left(\frac{A_{6,7,8}}{A_6} \right) + B_7 \left(\frac{A_2}{A_2} \right) \left(\frac{A_{4,5}}{A_4} \right) \left(\frac{A_{6,7,8}}{A_{6,7}} \right) + B_8 \left(\frac{A_2}{A_2} \right) \left(\frac{A_{4,5,6}}{A_4} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right. \right. \\ \left. \left. + B_5 A_0 \left[B_7 \left(\frac{A_{2,3}}{A_2} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \left(\frac{A_{6,7,8}}{A_{6,7}} \right) + B_8 \left(\frac{A_{2,3}}{A_2} \right) \left(\frac{A_{4,5,6}}{A_{4,5}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] + B_6 A_0 \left[B_8 \left(\frac{A_{2,3,4}}{A_2} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right\} \right. \\ \left. + B_3 \left\{ B_5 A_{0,1} \left[B_7 \left(\frac{A_{2,3}}{A_{2,3}} \right) \left(\frac{A_{4,5}}{A_{4,5}} \right) \left(\frac{A_{6,7,8}}{A_{6,7}} \right) + B_8 \left(\frac{A_{2,3}}{A_{2,3}} \right) \left(\frac{A_{4,5,6}}{A_{4,5}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right. \right. \\ \left. \left. + B_6 A_{0,1} \left[B_8 \left(\frac{A_{2,3,4}}{A_{2,3}} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right\} + B_4 \left\{ B_6 A_{0,1,2} \left[B_8 \left(\frac{A_{2,3,4}}{A_{2,3,4}} \right) \left(\frac{A_{4,5,6}}{A_{4,5,6}} \right) \left(\frac{A_{6,7,8}}{A_{6,7,8}} \right) \right] \right\} \right\} c_0$$

$$\vdots$$

$$\vdots$$

(1.3.14)

(1.3.14) is simply

$$c_{n+6} = \sum_{i_1=0}^n \left\{ B_{i_1+1} \sum_{i_2=i_1}^n \left\{ B_{i_2+3} \sum_{i_3=i_2}^n \left\{ B_{i_3+5} \prod_{i_4=0}^{i_1-1} A_{i_4} \prod_{i_5=i_1}^{i_2-1} A_{i_5+2} \prod_{i_6=i_2}^{i_3-1} A_{i_6+4} \prod_{i_7=i_3}^{n-1} A_{i_7+6} \right\} \right\} \right\} c_0 \quad (1.3.15)$$

Substitute (1.3.15) in (1.3.6) putting $m = 3$.

$$\begin{aligned}
 y_3(x) = & c_0 \sum_{n=0}^{\infty} \left\{ \sum_{i_1=0}^n \left\{ B_{i_1+1} \sum_{i_2=i_1}^n \left\{ B_{i_2+3} \sum_{i_3=i_2}^n \left\{ B_{i_3+5} \prod_{i_4=0}^{i_1-1} A_{i_4} \prod_{i_5=i_1}^{i_2-1} A_{i_5+2} \right. \right. \right. \right. \\
 & \times \left. \left. \prod_{i_6=i_2}^{i_3-1} A_{i_6+4} \prod_{i_7=i_3}^{n-1} A_{i_7+6} \right\} \right\} \right\} \Big\} x^{n+6+\lambda} \quad (1.3.16)
 \end{aligned}$$

By repeating this process for all higher terms of B_n 's, we obtain every $y_m(x)$ terms where $m \geq 4$. Substitute (1.3.7), (1.3.10), (1.3.13), (1.3.16) and including all $y_m(x)$ terms where $m \geq 4$ into (1.3.5).

Theorem 1.3.1 *The general expression of $y(x)$ for infinite series is*

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 \left\{ \sum_{n=0}^{\infty} \left(\prod_{i_1=0}^{n-1} A_{i_1} \right) x^{n+\lambda} + \sum_{n=0}^{\infty} \left\{ \sum_{i_1=0}^n \left\{ B_{i_1+1} \prod_{i_2=0}^{i_1-1} A_{i_2} \prod_{i_3=i_1}^{n-1} A_{i_3+2} \right\} \right\} x^{n+2+\lambda} \right. \\
 &+ \sum_{N=2}^{\infty} \left\{ \sum_{n=0}^{\infty} \left\{ \prod_{k=1}^{N-1} \left(\sum_{i_k=i_{k-1}}^n B_{i_k+2k-1} \right) \right. \right. \\
 &\times \left. \left. \sum_{i_N=i_{N-1}}^n \left\{ B_{i_N+2N-1} \prod_{l=0}^{N-1} \left(\prod_{i_{l+1}+N=i_l}^{i_{l+1}-1} A_{i_{l+1}+N+2l} \right) \prod_{i_{2N+1}=i_N}^{n-1} A_{i_{2N+1}+2N+1} \right\} \right\} \right\} x^{n+2N+\lambda} \Big\} \\
 &\text{where, } i_0 = 0 \quad (1.3.17)
 \end{aligned}$$

1.4 Polynomial which makes A_n term terminated

Now let's investigate the polynomial case of $y(x)$. Assume that A_n is terminated at certain value of n . Then each $y_i(x)$ where $i = 0, 1, 2, \dots$ will be polynomial. Examples of these are Heun's equation, GCH function[6], Lamé function, etc. First A_k is terminated at certain value of k . I choose eigenvalue α_0 which A_k is terminated where $\alpha_0 = 0, 1, 2, \dots$. A_{k+2} is terminated at certain value of k . I choose eigenvalue α_1 which A_{k+2} is terminated where $\alpha_1 = 0, 1, 2, \dots$. Also A_{k+4} is terminated at certain value of k . I choose eigenvalue α_2 which A_{k+4} is terminated where $\alpha_2 = 0, 1, 2, \dots$. By repeating this process I obtain

$$A_{\alpha_i+2i} = 0 \quad \text{where } i, \alpha_i = 0, 1, 2, \dots \quad (1.4.1)$$

In general, the two term recurrence relation for polynomial has only one eigenvalue: for example, the Laguerre function, confluent hypergeometric function, Legendre function, etc. In “Generalization of the three-term recurrence formula and its applications” [1], polynomial which makes B_n term terminated has infinite eigenvalues which is β_i where $i, \beta_i = 0, 1, 2, \dots$. In this chapter polynomial which makes A_n term terminated also has infinite eigenvalues which is α_i .

First observe the term in sequence c_n which does not include any B_n 's in (1.3.4): c_n with every index ($c_0, c_1, c_2, \dots$).

(a) As $\alpha_0=0$, then $A_0=0$ in (1.3.4).

$$c_0 \tag{1.4.2}$$

(b) As $\alpha_0=1$, then $A_1=0$ in (1.3.4).

$$\begin{aligned} c_0 \\ c_1 = A_0 c_0 \end{aligned} \tag{1.4.3}$$

(c) As $\alpha_0=2$, then $A_2=0$ in (1.3.4).

$$\begin{aligned} c_0 \\ c_1 = A_0 c_0 \\ c_2 = A_{0,1} c_0 \end{aligned} \tag{1.4.4}$$

Substitute (1.4.2), (1.4.3) and (1.4.4) in (1.3.6) putting $m = 0$.

$$y_0(x) = c_0 \sum_{n=0}^{\alpha_0} \left\{ \prod_{i_1=0}^{n-1} A_{i_1} \right\} x^{n+\lambda} \tag{1.4.5}$$

Observe the terms inside curly brackets of sequence c_n which include one term of B_n 's in (1.3.8): c_n with every index except c_0 and c_1 ($c_1, c_3, c_5, \dots$).

(a) As $\alpha_0=0$, then $A_0=0$ in (1.3.8).

$$\begin{aligned} c_2 &= B_1 c_0 \\ c_3 &= B_1 \cdot 1 \cdot A_2 c_0 \\ c_4 &= B_1 \cdot 1 \cdot A_{2,3} c_0 \\ c_5 &= B_1 \cdot 1 \cdot A_{2,3,4} c_0 \\ c_6 &= B_1 \cdot 1 \cdot A_{2,3,4,5} c_0 \\ &\vdots \quad \vdots \end{aligned} \tag{1.4.6}$$

As $i=1$ in (1.4.1),

$$A_{\alpha_1+2} = 0 \quad \text{where } \alpha_1 = 0, 1, 2, \dots \quad (1.4.7)$$

Substitute (1.4.6) in (1.3.6) putting $m = 1$ by using (1.4.7).

$$y_1^0(x) = c_0 B_1 \sum_{n=0}^{\alpha_1} \left\{ \prod_{i_1=0}^{n-1} A_{i_1+2} \right\} x^{n+2+\lambda} \quad (1.4.8)$$

In (1.4.8) $y_1^0(x)$ is sub-power series, having sequences c_n including one term of B_n 's in (1.3.3) as $\alpha_0=0$, for the polynomial case in which makes A_n term terminated.

(b) As $\alpha_0=1$, then $A_1=0$ in (1.3.8).

$$\begin{aligned} c_2 &= B_1 c_0 \\ c_3 &= \{B_1 \cdot 1 \cdot A_2 + B_2 A_0 \cdot 1\} c_0 \\ c_4 &= \{B_1 \cdot 1 \cdot A_{2,3} + B_2 A_0 A_3\} c_0 \\ c_5 &= \{B_1 \cdot 1 \cdot A_{2,3,4} + B_2 A_0 A_{3,4}\} c_0 \\ c_6 &= \{B_1 \cdot 1 \cdot A_{2,3,4,5} + B_2 A_0 A_{3,4,5}\} c_0 \\ &\vdots \end{aligned} \quad (1.4.9)$$

The first term in curly brackets of sequence c_n in (1.4.9) is same as (1.4.6). Then, its solution is equal to (1.4.8). Substitute (1.4.7) into the second term in curly brackets of sequence c_n in (1.4.9). Its power series expansion including the first and second terms in curly brackets of sequence c_n in (1.4.9), analytic at $x = 0$, is

$$y_1^1(x) = c_0 \left\{ B_1 \sum_{n=0}^{\alpha_1} \left\{ \prod_{i_1=0}^{n-1} A_{i_1+2} \right\} + B_2 A_0 \sum_{n=1}^{\alpha_1} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+2} \right\} \right\} x^{n+2+\lambda} \quad (1.4.10)$$

In (1.4.10) $y_1^1(x)$ is sub-power series, having sequences c_n including one term of B_n 's in (1.3.3), as $\alpha_0=1$ for the polynomial case in which makes A_n term terminated.

(c) As $\alpha_0=2$, then $A_2=0$ in (1.3.8).

$$\begin{aligned} c_2 &= B_1 c_0 \\ c_3 &= \{B_1 \cdot 1 \cdot A_2 + B_2 A_0 \cdot 1\} c_0 \\ c_4 &= \{B_1 \cdot 1 \cdot A_{2,3} + B_2 A_0 A_3 + B_3 A_{0,1} \cdot 1\} c_0 \\ c_5 &= \{B_1 \cdot 1 \cdot A_{2,3,4} + B_2 A_0 A_{3,4} + B_3 A_{0,1} A_4\} c_0 \\ c_6 &= \{B_1 \cdot 1 \cdot A_{2,3,4,5} + B_2 A_0 A_{3,4,5} + B_3 A_{0,1} A_{4,5}\} c_0 \\ c_7 &= \{B_1 \cdot 1 \cdot A_{2,3,4,5,6} + B_2 A_0 A_{3,4,5,6} + B_3 A_{0,1} A_{4,5,6}\} c_0 \\ &\vdots \end{aligned} \quad (1.4.11)$$

The first and second term in curly brackets of sequence c_n in (1.4.11) is same as (1.4.9). Then its power series expansion is same as (1.4.10). Substitute (1.4.7) into the third term in curly brackets of sequence c_n in (1.4.11). Its power series expansion including the first, second and third terms in curly brackets of sequence c_n in (1.4.11), analytic at $x = 0$, is

$$\begin{aligned} y_1^2(x) = & c_0 \left\{ B_1 \sum_{n=0}^{\alpha_1} \left\{ \prod_{i_1=0}^{n-1} A_{i_1+2} \right\} + B_2 A_0 \sum_{n=1}^{\alpha_1} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+2} \right\} \right. \\ & \left. + B_3 A_{0,1} \sum_{n=2}^{\alpha_1} \left\{ \prod_{i_1=2}^{n-1} A_{i_1+2} \right\} \right\} x^{n+2+\lambda} \end{aligned} \quad (1.4.12)$$

In (1.4.12) $y_1^2(x)$ is sub-power series, having sequence c_n including one term of B_n 's in (1.3.3) as $\alpha_0=2$, for the polynomial case in which makes A_n term terminated. By repeating this process for all $\alpha_0 = 3, 4, 5, \dots$, I obtain every $y_1^j(x)$ terms where $j \geq 3$. According to (1.4.8), (1.4.10), (1.4.12) and every $y_1^j(x)$ where $j \geq 3$, the general expression of $y_1(x)$ for all α_0 is

$$y_1(x) = c_0 \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right\} \right\} x^{i_2+2+\lambda} \text{ where } \alpha_0 \leq \alpha_1 \quad (1.4.13)$$

Observe the terms of sequence c_n which include two terms of B_n 's in (1.3.11): c_n with every index except c_0-c_3 ($c_4, c_5, c_6, \dots$).

(a) As $\alpha_0=0$, then $A_0=0$ in (1.3.11).

$$\begin{aligned} c_4 &= B_{1,3} c_0 \\ c_5 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_4 + B_4 \cdot 1 \cdot A_2 \cdot 1 \} c_0 \\ c_6 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_{4,5} + B_4 \cdot 1 \cdot A_2 A_5 + B_5 \cdot 1 \cdot A_{2,3} \cdot 1 \} c_0 \\ c_7 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6} + B_4 \cdot 1 \cdot A_2 A_{5,6} + B_5 \cdot 1 \cdot A_{2,3} A_6 + B_6 \cdot 1 \cdot A_{2,3,4} \cdot 1 \} c_0 \\ c_8 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6,7} + B_4 \cdot 1 \cdot A_2 A_{5,6,7} + B_5 \cdot 1 \cdot A_{2,3} A_{6,7} + B_6 \cdot 1 \cdot A_{2,3,4} A_7 \\ &\quad + B_7 \cdot 1 \cdot A_{2,3,4,5} \cdot 1 \} c_0 \\ &\vdots \qquad \qquad \qquad \vdots \end{aligned} \quad (1.4.14)$$

(i) As $\alpha_1=0$, then $A_2=0$ in (1.4.14).

$$\begin{aligned}
c_4 &= B_{1,3} c_0 \\
c_5 &= B_1 B_3 \cdot 1 \cdot 1 \cdot A_4 c_0 \\
c_6 &= B_1 B_3 \cdot 1 \cdot 1 \cdot A_{4,5} c_0 \\
c_7 &= B_1 B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6} c_0 \\
c_8 &= B_1 B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6,7} c_0 \\
&\vdots \qquad \qquad \vdots
\end{aligned} \tag{1.4.15}$$

As $i=2$ in (1.4.1),

$$A_{\alpha_2+4} = 0 \quad \text{where } \alpha_2 = 0, 1, 2, \dots \tag{1.4.16}$$

Substitute (1.4.15) in (1.3.6) putting $m = 2$ by using (1.4.16).

$$y_2^{0,0}(x) = c_0 B_1 B_3 \sum_{n=0}^{\alpha_2} \left\{ \prod_{i_1=0}^{n-1} A_{i_1+4} \right\} x^{n+4+\lambda} \tag{1.4.17}$$

In (1.4.17) $y_2^{0,0}(x)$ is sub-power series, having sequences c_n including two term of B_n 's in (1.3.3) as $\alpha_0=0$ and $\alpha_1=0$, for the polynomial case in which makes A_n term terminated.

(ii) As $\alpha_1=1$, then $A_3=0$ in (1.4.14).

$$\begin{aligned}
c_4 &= B_{1,3} c_0 \\
c_5 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_4 + B_4 \cdot 1 \cdot A_2 \cdot 1 \} c_0 \\
c_6 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_{4,5} + B_4 \cdot 1 \cdot A_2 \cdot A_5 \} c_0 \\
c_7 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6} + B_4 \cdot 1 \cdot A_2 \cdot A_{5,6} \} c_0 \\
c_8 &= B_1 \{ B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6,7} + B_4 \cdot 1 \cdot A_2 \cdot A_{5,6,7} \} c_0 \\
&\vdots \qquad \qquad \vdots
\end{aligned} \tag{1.4.18}$$

The first term in curly brackets of sequence c_n in (1.4.18) is same as (1.4.15). Then its power series expansion is equal to (1.4.17). Substitute (1.4.16) into the second term in curly brackets of sequence c_n in (1.4.18). Its power series expansion including the first and second terms in curly brackets of sequence c_n , analytic at $x = 0$, is

$$y_2^{0,1}(x) = c_0 B_1 \left\{ B_3 \sum_{n=0}^{\alpha_2} \left\{ \prod_{i_1=0}^{n-1} A_{i_1+4} \right\} + B_4 A_2 \sum_{n=1}^{\alpha_2} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+4} \right\} \right\} x^{n+4+\lambda} \tag{1.4.19}$$

In (1.4.19) $y_2^{0,1}(x)$ is sub-power series, having sequences c_n including two term of B_n 's in (1.3.3) as $\alpha_0=0$ and $\alpha_1=1$, for the polynomial case in which makes A_n term terminated.

(iii) As $\alpha_1=2$, then $A_4=0$ in (1.4.14).

$$\begin{aligned}
c_4 &= B_{1,3}c_0 \\
c_5 &= B_1\{B_3 \cdot 1 \cdot 1 \cdot A_4 + B_4 \cdot 1 \cdot A_2 \cdot 1\}c_0 \\
c_6 &= B_1\{B_3 \cdot 1 \cdot 1 \cdot A_{4,5} + B_4 \cdot 1 \cdot A_2 \cdot A_5 + B_5 \cdot 1 \cdot A_{2,3} \cdot 1\}c_0 \\
c_7 &= B_1\{B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6} + B_4 \cdot 1 \cdot A_2 \cdot A_{5,6} + B_5 \cdot 1 \cdot A_{2,3} \cdot A_6\}c_0 \\
c_8 &= B_1\{B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6,7} + B_4 \cdot 1 \cdot A_2 \cdot A_{5,6,7} + B_5 \cdot 1 \cdot A_{2,3} \cdot A_{6,7}\}c_0 \\
&\vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.4.20}$$

The first and second term in curly brackets of sequence c_n in (1.4.20) is same as (1.4.18). Then its power series expansion is same as (1.4.19). Substitute (1.4.16) into the third term in curly brackets of sequence c_n in (1.4.20). Its power series expansion including the first, second and third terms in curly brackets of sequence c_n in (1.4.20), analytic at $x = 0$, is

$$\begin{aligned}
y_2^{0,2}(x) &= c_0 B_1 \left\{ B_3 \sum_{n=0}^{\alpha_2} \left\{ \prod_{i_1=0}^{n-1} A_{i_1+4} \right\} + B_4 A_2 \sum_{n=1}^{\alpha_2} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+4} \right\} \right. \\
&\quad \left. + B_5 A_{2,3} \sum_{n=2}^{\alpha_2} \left\{ \prod_{i_1=2}^{n-1} A_{i_1+4} \right\} \right\} x^{n+4+\lambda}
\end{aligned} \tag{1.4.21}$$

In (1.4.21) $y_2^{0,2}(x)$ is sub-power series, having sequences c_n including two term of B_n 's in (1.3.3) as $\alpha_0=0$ and $\alpha_1=2$, for the polynomial case in which makes A_n term terminated. By repeating this process for all $\alpha_1 = 3, 4, 5, \dots$, we obtain every $y_2^{0,j}(x)$ terms where $j \geq 3$. According to (1.4.17), (1.4.19), (1.4.21) and every $y_2^{0,j}(x)$ where $j \geq 3$, the general expression of $y_2^0(x)$ for the case of $\alpha_0 = 0$ replacing the index n by i_0 is

$$y_2^0(x) = c_0 B_1 \sum_{i_0=0}^{\alpha_1} \left\{ B_{i_0+3} \prod_{i_1=0}^{i_0-1} A_{i_1+2} \sum_{i_2=i_0}^{\alpha_2} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+4} \right\} \right\} x^{i_2+4+\lambda} \tag{1.4.22}$$

In (1.4.22) $y_2^0(x)$ is sub-power series, having sequences c_n including two term of B_n 's in (1.3.3) as $\alpha_0=0$, for the polynomial case in which makes A_n term terminated.

(b) As $\alpha_0=1$, then $A_1=0$ in (1.3.11).

$$\begin{aligned}
c_4 &= B_{1,3}c_0 \\
c_5 &= \left\{ B_1 \left[B_3 \cdot 1 \cdot 1 \cdot A_4 + B_4 \cdot 1 \cdot A_2 \cdot 1 \right] + B_2 \left[B_4 A_0 \cdot 1 \cdot 1 \right] \right\} c_0 \\
c_6 &= \left\{ B_1 \left[B_3 \cdot 1 \cdot 1 \cdot A_{4,5} + B_4 \cdot 1 \cdot A_2 A_5 + B_5 \cdot 1 \cdot A_{2,3} \cdot 1 \right] \right. \\
&\quad \left. + B_2 \left[B_4 A_0 \cdot 1 \cdot A_5 + B_5 A_0 A_3 \cdot 1 \right] \right\} c_0 \\
c_7 &= \left\{ B_1 \left[B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6} + B_4 \cdot 1 \cdot A_2 A_{5,6} + B_5 \cdot 1 \cdot A_{2,3} \cdot A_6 + B_6 \cdot 1 \cdot A_{2,3,4} \cdot 1 \right] \right. \\
&\quad \left. + B_2 \left[B_4 A_0 \cdot 1 \cdot A_{5,6} + B_5 A_0 A_3 A_6 + B_6 A_0 A_{3,4} \cdot 1 \right] \right\} c_0 \\
c_8 &= \left\{ B_1 \left[B_3 \cdot 1 \cdot 1 \cdot A_{4,5,6,7} + B_4 \cdot 1 \cdot A_2 A_{5,6,7} + B_5 \cdot 1 \cdot A_{2,3} \cdot A_{6,7} \right. \right. \\
&\quad \left. + B_6 \cdot 1 \cdot A_{2,3,4} \cdot A_7 + B_7 \cdot 1 \cdot A_{2,3,4,5} \cdot 1 \right] \\
&\quad \left. + B_2 \left[B_4 A_0 \cdot 1 \cdot A_{5,6,7} + B_5 A_0 A_3 A_{6,7} + B_6 A_0 A_{3,4} \cdot A_7 + B_7 A_0 A_{3,4,5} \cdot 1 \right] \right\} c_0 \\
&\quad \vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.4.23}$$

The first square brackets including B_1 inside curly brackets in sequence c_n in (1.4.23) is same as (1.4.14). Then its power series expansion is same as (1.4.22). Observe the second square brackets including B_2 inside curly brackets in sequence c_n in (1.4.23).

(i) As $\alpha_1=1$, then $A_3=0$ in the second square brackets including B_2 inside curly brackets in sequence c_n in (1.4.23).

$$\begin{aligned}
c_4 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot 1 \right\} c_0 \\
c_5 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot A_5 \right\} c_0 \\
c_6 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot A_{5,6} \right\} c_0 \\
c_7 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot A_{5,6,7} \right\} c_0 \\
&\quad \vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.4.24}$$

Its power series expansion of (1.4.24) by using (1.4.16), analytic at $x = 0$, is

$$y_2^{1,1}(x) = c_0 A_0 B_{2,4} \sum_{n=1}^{\alpha_2} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+4} \right\} x^{n+4+\lambda} \tag{1.4.25}$$

In (1.4.25) $y_2^{1,1}(x)$ is sub-power series, for the second square brackets inside curly brackets in sequence c_n including two term of B_n 's in (1.4.23) as $\alpha_0=1$ and $\alpha_1=1$, for the polynomial case in which makes A_n term terminated.

(ii) As $\alpha_1=2$, then $A_4=0$ in second square brackets inside curly brackets in sequence c_n including B_2 in (1.4.23).

$$\begin{aligned}
c_5 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot 1 \right\} c_0 \\
c_6 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot A_5 + B_5 A_3 \cdot 1 \right\} c_0 \\
c_7 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot A_{5,6} + B_5 A_3 \cdot A_6 \right\} c_0 \\
c_8 &= B_2 A_0 \left\{ B_4 \cdot 1 \cdot A_{5,6,7} + B_5 A_3 \cdot A_{6,7} \right\} c_0 \\
&\vdots \qquad \qquad \qquad \vdots
\end{aligned} \tag{1.4.26}$$

The first term in curly brackets of sequence c_n in (1.4.26) is same as (1.4.24). Then its power series expansion is same as (1.4.25). Substitute (1.4.16) into the second term in curly brackets of sequence c_n in (1.4.26). Its power series expansion including the first and second terms in curly brackets of sequence c_n in (1.4.26), analytic at $x = 0$, is

$$y_2^{1,2}(x) = c_0 B_2 A_0 \left\{ B_4 \sum_{n=1}^{\alpha_2} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+4} \right\} + B_5 A_3 \sum_{n=2}^{\alpha_2} \left\{ \prod_{i_1=2}^{n-1} A_{i_1+4} \right\} \right\} x^{n+4+\lambda} \tag{1.4.27}$$

In (1.4.27) $y_2^{1,2}(x)$ is sub-power series, for the second square brackets inside curly brackets in sequence c_n including two term of B_n 's in (1.4.23) as $\alpha_0=1$ and $\alpha_1=2$, for the polynomial case which makes A_n term terminated.

By using similar process as I did before, the solution for $\alpha_1=3$ with $A_5=0$ for the second square brackets inside curly brackets in sequence c_n including B_2 in (1.4.23) is

$$\begin{aligned}
y_2^{1,3}(x) &= c_0 B_2 A_0 \left\{ B_4 \sum_{n=1}^{\alpha_2} \left\{ \prod_{i_1=1}^{n-1} A_{i_1+4} \right\} + B_5 A_3 \sum_{n=2}^{\alpha_2} \left\{ \prod_{i_1=2}^{n-1} A_{i_1+4} \right\} \right. \\
&\quad \left. + B_6 A_{3,4} \sum_{n=3}^{\alpha_2} \left\{ \prod_{i_1=3}^{n-1} A_{i_1+4} \right\} \right\} x^{n+4+\lambda}
\end{aligned} \tag{1.4.28}$$

By repeating this process for all $\alpha_1 = 4, 5, 6, \dots$, we obtain every $y_2^{1,j}(x)$ terms where $j \geq 4$ for the second square brackets inside curly brackets in sequence c_n including two term of B_n 's in (1.4.23). According to (1.4.22), (1.4.25), (1.4.27), (1.4.28) and every

$y_2^{1,j}(x)$ where $j \geq 4$, the general expression of $y_2^1(x)$ for all $\alpha_0 = 1$ replacing the index n by i_0 is

$$y_2^1(x) = c_0 \left\{ B_1 \sum_{i_0=0}^{\alpha_1} \left\{ B_{i_0+3} \prod_{i_1=0}^{i_0-1} A_{i_1+2} \sum_{i_2=i_0}^{\alpha_2} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+4} \right\} \right\} \right. \\ \left. + B_2 A_0 \sum_{i_0=1}^{\alpha_1} \left\{ B_{i_0+3} \prod_{i_1=1}^{i_0-1} A_{i_1+2} \sum_{i_2=i_0}^{\alpha_2} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+4} \right\} \right\} \right\} x^{i_2+4+\lambda} \quad (1.4.29)$$

(c) As $\alpha_0=2$, then $A_2=0$ in (1.3.11).

$$\begin{aligned} c_4 &= B_{1,3} c_0 \\ c_5 &= \left\{ B_1 \left[B_3 \cdot 1 \cdot 1 \cdot A_4 + B_4 \cdot 1 \cdot A_2 \cdot 1 \right] + B_2 \left[B_4 A_0 \cdot 1 \cdot 1 \right] \right\} c_0 \\ c_6 &= \left\{ B_1 \cdot 1 \left[B_3 \cdot 1 \cdot A_{4,5} + B_4 A_2 A_5 + B_5 A_{2,3} \cdot 1 \right] + B_2 A_0 \left[B_4 \cdot 1 \cdot A_5 + B_5 A_3 \cdot 1 \right] \right. \\ &\quad \left. + B_3 A_{0,1} \left[B_5 \cdot 1 \cdot 1 \right] \right\} c_0 \\ c_7 &= \left\{ B_1 \cdot 1 \left[B_3 \cdot 1 \cdot A_{4,5,6} + B_4 A_2 A_{5,6} + B_5 A_{2,3} A_6 + B_6 A_{2,3,4} \cdot 1 \right] \right. \\ &\quad \left. + B_2 A_0 \left[B_4 \cdot 1 \cdot A_{5,6} + B_5 A_3 A_6 + B_6 A_{3,4} \cdot 1 \right] + B_3 A_{0,1} \left[B_5 \cdot 1 \cdot A_6 + B_6 A_4 \cdot 1 \right] \right\} c_0 \\ c_8 &= \left\{ B_1 \cdot 1 \left[B_3 \cdot 1 \cdot A_{4,5,6,7} + B_4 A_2 A_{5,6,7} + B_5 A_{2,3} A_{6,7} + B_6 A_{2,3,4} A_7 + B_7 A_{2,3,4,5} \cdot 1 \right] \right. \\ &\quad \left. + B_2 A_0 \left[B_4 \cdot 1 \cdot A_{5,6,7} + B_5 A_3 A_{6,7} + B_6 A_{3,4} A_7 + B_7 A_{3,4,5} \cdot 1 \right] \right. \\ &\quad \left. + B_3 A_{0,1} \left[B_5 \cdot 1 \cdot A_{6,7} + B_6 A_4 A_7 + B_7 A_{4,5} \cdot 1 \right] \right\} c_0 \\ &\vdots \qquad \qquad \qquad \vdots \end{aligned} \quad (1.4.30)$$

By repeating similar process from the above, the general expression of $y_2^2(x)$ for all $\alpha_0 = 2$ in (1.4.30) is

$$y_2^2(x) = c_0 \left\{ B_1 \sum_{i_0=0}^{\alpha_1} \left\{ B_{i_0+3} \prod_{i_1=0}^{i_0-1} A_{i_1+2} \sum_{i_2=i_0}^{\alpha_2} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+4} \right\} \right\} \right. \\ + B_2 A_0 \sum_{i_0=1}^{\alpha_1} \left\{ B_{i_0+3} \prod_{i_1=1}^{i_0-1} A_{i_1+2} \sum_{i_2=i_0}^{\alpha_2} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+4} \right\} \right\} \\ \left. + B_3 A_{0,1} \sum_{i_0=2}^{\alpha_1} \left\{ B_{i_0+3} \prod_{i_1=2}^{i_0-1} A_{i_1+2} \sum_{i_2=i_0}^{\alpha_2} \left\{ \prod_{i_3=i_0}^{i_2-1} A_{i_3+4} \right\} \right\} \right\} x^{i_2+4+\lambda} \quad (1.4.31)$$

Again by repeating this process for all $\alpha_0 = 3, 4, 5, \dots$, I obtain every $y_2^j(x)$ terms where $j \geq 3$. Then I have general expression $y_2(x)$ for all α_0 of two term of B_n 's according (1.4.22), (1.4.29), (1.4.31) and $y_2^j(x)$ terms where $j \geq 3$.

$$y_2(x) = c_0 \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left\{ B_{i_2+3} \prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \sum_{i_4=i_2}^{\alpha_2} \left\{ \prod_{i_5=i_2}^{i_4-1} A_{i_5+4} \right\} \right\} \right\} x^{i_4+4+\lambda}$$

where $\alpha_0 \leq \alpha_1 \leq \alpha_2$

(1.4.32)

By using similar process for the previous cases of zero, one and two terms of B_n 's, the function $y_3(x)$ for the case of three term of B_n 's is

$$y_3(x) = c_0 \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left\{ B_{i_2+3} \prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right. \right.$$

$$\left. \times \sum_{i_4=i_2}^{\alpha_2} \left\{ B_{i_4+5} \prod_{i_5=i_2}^{i_4-1} A_{i_5+4} \sum_{i_6=i_4}^{\alpha_3} \left\{ \prod_{i_7=i_4}^{i_6-1} A_{i_7+6} \right\} \right\} \right\} \left. \right\} x^{i_6+6+\lambda}$$

where $\alpha_0 \leq \alpha_1 \leq \alpha_2 \leq \alpha_3$

(1.4.33)

By repeating this process for all higher terms of B_n 's, I obtain every $y_m(x)$ terms where $m > 3$. Substitute (1.4.5), (1.4.13), (1.4.32), (1.4.33) and including all $y_m(x)$ terms where $m > 3$ into (1.3.5).

Theorem 1.4.1 *The general expression of $y(x)$ for the polynomial case which makes A_n term terminated in the three term recurrence relation is*

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots$$

$$= c_0 \left\{ \sum_{i_0=0}^{\alpha_0} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right.$$

$$+ \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\alpha_k} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right.$$

$$\left. \times \sum_{i_{2N}=i_{2(N-1)}}^{\alpha_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \left. \right\} x^{i_{2N}+2N+\lambda} \right\}$$
(1.4.34)

In the above, $\alpha_i \leq \alpha_j$ only if $i \leq j$ where $i, j, \alpha_i, \alpha_j \in \mathbb{N}_0$

Theorem 1.4.2 *For infinite series, replacing $\alpha_0, \alpha_1, \alpha_k$ and α_N by ∞ in (1.4.34)*

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (1.4.35)
\end{aligned}$$

(1.4.35) is equivalent to (1.3.17). (1.4.35) is the another general expression of $y(x)$ for the infinite series.

1.5 Summary

In Ref.[1] I showed how to obtain the general expression of the power series for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's using 3TRF in a linear ordinary differential equation having a relation between three different coefficients. This was done by letting A_n in sequence c_n is the leading term in the analytic function $y(x)$: the sequence c_n consists of combinations A_n and B_n .

In this chapter, I show how to construct the Frobenius solution in closed forms for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's using R3TRF in a linear ordinary differential equation having a relation between three different coefficients. This is done by letting B_n in sequence c_n is the leading term in the analytic function $y(x)$.

The infinite series in this chapter are equivalent to the infinite series in Ref.[1]. The former is the letting B_n in sequence c_n is the leading term in the analytic function $y(x)$. And the latter is the letting A_n in sequence c_n is the leading term in the analytic function $y(x)$.

In chapters 2–9 I apply R3TRF to the power series expansions, integral forms and generating functions for Heun, Confluent Heun, Mathieu, etc equations; local solutions of these equations for infinite series and polynomial which makes A_n term terminated are constructed analytically.

Bibliography

- [1] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806
- [2] Choun, Y.S., “Asymptotic behavior of Heun function and its integral formalism,” arXiv:1303.0876
- [3] Choun, Y.S., “The power series expansion of Mathieu function and its integral formalism,” arXiv:1303.0820.
- [4] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische Annalen* 33, (1889)161.
- [5] Mathieu, E., “Mémoire sur Le Mouvement Vibratoire d’une Membrane de forme Elliptique,” *J. Math. Pure Appl.* 13, (1868), p. 137–203.
- [6] Choun, Y.S., “Approximative solution of the spin free Hamiltonian involving only scalar potential for the $q - \bar{q}$ system,” arXiv:1302.7309
- [7] Gutierrez-Vega, J. C., Rodriguez-Dagnino, R. M., Meneses-Nava, M. A. and Chavez-Cerda, S., “Mathieu functions, a visual approach,” *Amer. J. Phys.* 71(3), (2003), p. 233–242.
- [8] Troesch, B. A. and Troesch, H. R., “Eigenfrequencies of an elliptic membrane,” *Math. Comp.* 27(124), (1973), p. 755–765.
- [9] Ronveaux, A., *Heuns Differential Equations*, Oxford University Press, (1995).
- [10] Decarreau, A., Dumont-Lepage, M.-C., Maroni, P., Robert, A., Ronveaux, A., “Formes canoniques des equations confluentes de l’equation de Heun,” *Ann. Soc. Sci. Bruxelles Vol.92*, (1978), p. 53–78.
- [11] Suzuki, H., Takasugi, E., Umetsu, H., “Perturbations of Kerr-de Sitter Black Hole and Heun’s Equation,” *Prog. Theor. Phys.* 100, (1998)491-505.

- [12] Suzuki, H., Takasugi, E., Umetsu, H., “Analytic solution of Teukolsky Equation in Kerr-de Sitter and Kerr-Newman-de Sitter Geometries,” Prog. Theor. Phys. 102, (1999)253-272.
- [13] Lamé, G., “Sur les surfaces isothermes dans les corps homogènes en équilibre de température,” J. Math. Pures Appl. 2, (1837)147188.
- [14] Bateman, H., Erdelyi, A.: *Higher transcendental functions 3. Automorphic functions*, (McGraw-Hill, 1955).
- [15] Hobson, E.W.: *The theory of spherical and ellipsoidal harmonics* , (Cambridge Univ. Press, 1931).
- [16] Whittaker, E.T. and Watson, G.N.: *A course of modern analysis* , (Cambridge Univ. Press, 1952).
- [17] Maroni, P., “Sur quelques relations intégrales entre les solutions de l’équation biconfluente de Heun,” Ann. Inst. H.P. 30, (1979), p. 315–332.

Chapter 2

Heun function using reversible three-term recurrence formula

In Ref.[2, 3], I construct the power series expansions in closed forms of Heun equation and its integral forms for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's¹ by applying three term recurrence formula (3TRF).

In this chapter, I will apply reversible three term recurrence formula to the Frobenius solutions in closed forms of Heun equation and its integral forms for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's².

The next chapter out of all 9 chapters describes the generating function of Heun polynomial which makes A_n term terminated including all higher terms of B_n 's by using R3TRF.

Nine examples of 192 local solutions of the Heun equation (Maier, 2007) are provided in the appendices 1 and 2. For each example, I construct the power series expansions of Heun equation and its integral forms by applying R3TRF.

2.1 Introduction

The Heun function generalizes all well-known special functions such as Spheroidal Wave, Lamé, Mathieu, and hypergeometric ${}_2F_1$, ${}_1F_1$ and ${}_0F_1$ functions. The Heun function,

¹“ higher terms of A_n 's” means at least two terms of A_n 's.

²“ higher terms of B_n 's” means at least two terms of B_n 's.

having three term recurrence relations in its power series, is the most outstanding special functions in among every analytic functions.

Recently Heun function started to appear in theoretical modern physics. For example the Heun function comes out in the hydrogen-molecule ion[14], in the Schrödinger equation with doubly anharmonic potential[10] (its solution is the confluent forms of Heun equation), in the Stark effect[12], in perturbations of the Kerr metric[15, 16, 17, 18, 19], in crystalline materials[13], in Collogero-Moser-Sutherland systems[20], etc., just to mention a few.[6, 7, 8, 9] Traditionally, we have explained all physical phenomenons by only using two term recursion relation in the power series expansion until 19th century. However, since modern physics (quantum gravity, SUSY, general relativity, etc) come out of the world, we have at least three or four recurrence relations in power series expansions. Furthermore these type of problems can not be reduced to two term recurrence relations by changing independent variables and coefficients.[11]

Due to its complexity Heun function was neglected for almost 100 years[4]. According to Whittaker's hypothesis, 'The Heun function can not be described in form of contour integrals of elementary functions even if it is the simplest class of special functions.' Because Heun equation consists of three recurrence relations in its power series expansion. Heun equation could not be described in the form of a definite or contour integral of any elementary function for the last 100 years. The three different coefficients in its power series expansion create mathematical difficulty to be described it into direct integral representation. Instead in Fredholm integral equations, Heun equation is obtained by integral equations; such integral relationships express one analytic solution in terms of another analytic solution. More precisely, in earlier literature the integral representations of Heun equation were constructed by using two types of relations: (1) Linear relations using Fredholm integral equations. [21, 12] (2) Non-linear relation (Malurkar-type integral relations) including Fredholm integral equations using two variables. [23, 24, 25, 26]

In previous papers I show the analytic solutions of Heun equation for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's by applying 3TRF [2, 3]; (1) the power series expansion of Heun equation, (2) its integral representation and its asymptotic behavior including the boundary condition for an independent variable x . Indeed, according to analyzing Heun function into its combined direct and contour integral forms resulting in a precise, we are able to observe how Heun function relate to other well-known special functions such as Mathieu function, Lamé function, confluent forms of Heun function and etc. Because a ${}_1F_1$ or ${}_2F_1$ function recurs in each of sub-integral forms of all these analytic functions. Also, an orthogonal relation of Heun function can be obtained from its integral representation.

In this chapter, by applying R3TRF, I construct the power series expansion in closed forms of Heun equation for infinite series and polynomial which makes A_n term

terminated including all higher terms of B_n 's analytically and its integral form. Heun's differential equation is a second-order linear ordinary differential equation of the form [4].

$$\frac{d^2 y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0 \quad (2.1.1)$$

With the condition $\epsilon = \alpha + \beta - \gamma - \delta + 1$. The parameters play different roles: $a \neq 0$ is the singularity parameter, $\alpha, \beta, \gamma, \delta, \epsilon$ are exponent parameters, q is the accessory parameter which in many physical applications appears as a spectral parameter. Also, α and β are identical to each other. The total number of free parameters is six. It has four regular singular points which are $0, 1, a$ and ∞ with exponents $\{0, 1 - \gamma\}$, $\{0, 1 - \delta\}$, $\{0, 1 - \epsilon\}$ and $\{\alpha, \beta\}$. Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} \quad \text{where } \lambda = \text{indicial root} \quad (2.1.2)$$

Plug (2.1.2) into (2.1.1).

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (2.1.3)$$

where,

$$\begin{aligned} A_n &= \frac{(n+\lambda)(n-1+\gamma+\epsilon+\lambda+a(n-1+\gamma+\lambda+\delta))+q}{a(n+1+\lambda)(n+\gamma+\lambda)} \\ &= \frac{(n+\lambda)(n+\alpha+\beta-\delta+\lambda+a(n+\delta+\gamma-1+\lambda))+q}{a(n+1+\lambda)(n+\gamma+\lambda)} \\ &= \frac{(1+a)}{a} \frac{\left(n - \frac{-(\varphi+2(1+a)\lambda)-\sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \right) \left(n - \frac{-(\varphi+2(1+a)\lambda)+\sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \right)}{(n+\lambda+1)(n+\lambda+\gamma)} \end{aligned} \quad (2.1.4a)$$

and

$$\varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1)$$

$$\begin{aligned} B_n &= -\frac{(n-1+\lambda)(n+\gamma+\delta+\epsilon-2+\lambda)+\alpha\beta}{a(n+1+\lambda)(n+\gamma+\lambda)} \\ &= -\frac{(n-1+\lambda+\alpha)(n-1+\lambda+\beta)}{a(n+1+\lambda)(n+\gamma+\lambda)} \end{aligned} \quad (2.1.4b)$$

$$c_1 = A_0 c_0 \quad (2.1.4c)$$

We have two indicial roots which are $\lambda = 0$ and $1 - \gamma$.

2.2 Power series

2.2.1 Polynomial of type 2

There are three types of polynomials in three term recurrence relation of a linear ordinary differential equation: (1) polynomial which makes B_n term terminated: A_n term is not terminated, (2) polynomial which makes A_n term terminated: B_n term is not terminated, (3) polynomial which makes A_n and B_n terms terminated at the same time. In general Heun polynomial is defined as type 3 polynomial where A_n and B_n terms terminated. Heun polynomial comes from Heun equation that has a fixed integer value of α or β , just as it has a fixed value of q . In three term recurrence relation, polynomial of type 3 I categorize as complete polynomial. In future papers I will derive type 3 Heun polynomial. In this part I construct the power series expansion and an integral forms for Heun polynomial of type 2: I treat α , β , γ and δ as free variables and the accessory parameter q as a fixed value. In my next papers I will work on the generating functions for Heun polynomial of type 2.³

In chapter 1, the general expression of power series of $y(x)$ for polynomial of type 2 is

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 \left\{ \sum_{i_0=0}^{\alpha_0} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
 &\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\alpha_k} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
 &\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\alpha_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (2.2.1)
 \end{aligned}$$

In the above, $\alpha_i \leq \alpha_j$ only if $i \leq j$ where $i, j, \alpha_i, \alpha_j \in \mathbb{N}_0$.

For a polynomial, we need a condition, which is:

$$A_{\alpha_i+2i} = 0 \quad \text{where } i, \alpha_i = 0, 1, 2, \dots \quad (2.2.2)$$

Put $n = \alpha_i + 2i$ in (2.1.4a) and use the condition $A_{\alpha_i+2i} = 0$.

$$\alpha_i + 2i = \frac{-(\varphi + 2(1+a)\lambda) \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)}$$

³If A_n and B_n terms are not terminated, it turns to be infinite series.

In the above, α_i is an eigenvalue that makes A_n term terminated at certain value of index n . (2.2.2) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (2.2.1).

The case of $\sqrt{\varphi^2 - 4(1+a)q} = -\{\varphi + 2(1+a)(q_i + 2i + \lambda)\}$ In (2.1.4a)-(2.1.4c) replace $\sqrt{\varphi^2 - 4(1+a)q}$ by $-\{\varphi + 2(1+a)(q_i + 2i + \lambda)\}$. In (2.2.2) replace index α_i by q_i . Take the new (2.1.4a)-(2.1.4c), (2.2.2) and put them in (2.2.1). After the replacement process, the general expression of power series of Heun's equation for polynomial of type 2 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}}{(-q_1)_{i_0} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}} \frac{(3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \eta^{i_1} \Bigg\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \lambda + \alpha)(i_k + 2k + \lambda + \beta)}{(i_k + 2(k+1) + \lambda)(i_k + 2k + 1 + \gamma + \lambda)} \right. \\
&\quad \times \frac{(-q_k)_{i_k} \left(q_k + 4k + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_k}}{(-q_k)_{i_{k-1}} \left(q_k + 4k + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_{k-1}}} \frac{(2k+1+\lambda)_{i_{k-1}} (2k+\gamma+\lambda)_{i_{k-1}}}{(2k+1+\lambda)_{i_k} (2k+\gamma+\lambda)_{i_k}} \Bigg\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} \left(q_n + 4n + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_n}}{(-q_n)_{i_{n-1}} \left(q_n + 4n + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_{n-1}}} \frac{(2n+1+\lambda)_{i_{n-1}} (2n+\gamma+\lambda)_{i_{n-1}}}{(2n+1+\lambda)_{i_n} (2n+\gamma+\lambda)_{i_n}} \eta^{i_n} \Bigg\} z^n \Bigg\} \quad (2.2.3)
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ q = -(q_j + 2j + \lambda)\{\varphi + (1+a)(q_j + 2j + \lambda)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

The case of $\sqrt{\varphi^2 - 4(1+a)q} = \varphi + 2(1+a)(q_i + 2i + \lambda)$ In (2.1.4a)-(2.1.4c) replace $\sqrt{\varphi^2 - 4(1+a)q}$ by $\varphi + 2(1+a)(q_i + 2i + \lambda)$. In (2.2.2) replace index α_i by q_i . Take the new (2.1.4a)-(2.1.4c), (2.2.2) and put them in (2.2.1). After the replacement process, its solution is equivalent to (2.2.3).

Put $c_0 = 1$ as $\lambda=0$ for the first kind of independent solutions of Heun equation and $c_0 = \left(\frac{1+a}{a}\right)^{1-\gamma}$ as $\lambda = 1 - \gamma$ for the second one in (2.2.3).

Remark 2.2.1 *The power series expansion of Heun equation of the first kind for polynomial of type 2 about $x = 0$ as*

$q = -(q_j + 2j)\{\alpha + \beta - \delta + a(\delta + \gamma - 1) + (1+a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned} y(x) &= HF_{q_j}^R \left(q_j = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j, \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_1 = \frac{\varphi}{(1+a)} \right. \\ &\quad \left. ; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\ &= \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega_1)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \\ &+ \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega_1)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega_1)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega_1)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\ &+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega_1)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\ &\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha)(i_k + 2k + \beta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \frac{(-q_k)_{i_k} (q_k + 4k + \Omega_1)_{i_k} (2k + 1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega_1)_{i_{k-1}} (2k + 1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\ &\times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega_1)_{i_n} (2n + 1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega_1)_{i_{n-1}} (2n + 1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \end{aligned} \quad (2.2.4)$$

Remark 2.2.2 The power series expansion of Heun equation of the second kind for polynomial of type 2 about $x = 0$ as

$q = -(q_j + 2j + 1 - \gamma)\{\alpha + \beta + 1 - \gamma - (1 - a)\delta + (1 + a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
 y(x) &= HS_{q_j}^R \left(q_j = \frac{-\{\varphi + 2(1+a)(1-\gamma)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j \right. \\
 &\quad \left. , \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_2 = \frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
 &= \eta^{1-\gamma} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega_2)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \eta^{i_0} \right. \\
 &\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + 1 - \gamma + \alpha)(i_0 + 1 - \gamma + \beta)}{(i_0 + 3 - \gamma)(i_0 + 2)} \frac{(-q_0)_{i_0} (q_0 + \Omega_2)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \right. \\
 &\quad \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega_2)_{i_1} (4-\gamma)_{i_0} (3)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega_2)_{i_0} (4-\gamma)_{i_1} (3)_{i_1}} \eta^{i_1} \Big\} z \\
 &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + 1 - \gamma + \alpha)(i_0 + 1 - \gamma + \beta)}{(i_0 + 3 - \gamma)(i_0 + 2)} \frac{(-q_0)_{i_0} (q_0 + \Omega_2)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \right. \\
 &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + 1 - \gamma + \alpha)(i_k + 2k + 1 - \gamma + \beta)}{(i_k + 2(k+1) + 1 - \gamma)(i_k + 2(k+1))} \right. \\
 &\quad \times \frac{(-q_k)_{i_k} (q_k + 4k + \Omega_2)_{i_k} (2(k+1) - \gamma)_{i_{k-1}} (2k+1)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega_2)_{i_{k-1}} (2(k+1) - \gamma)_{i_k} (2k+1)_{i_k}} \Big\} \\
 &\quad \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega_2)_{i_n} (2(n+1) - \gamma)_{i_{n-1}} (2n+1)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega_2)_{i_{n-1}} (2(n+1) - \gamma)_{i_n} (2n+1)_{i_n}} \eta^{i_n} \right\} z^n \Big\} \quad (2.2.5)
 \end{aligned}$$

In Ref.[2, 3] I treat α and/or β as a fixed value and γ, δ, q as free variables to construct Heun polynomial of type 1: (1) if $\alpha = -2\alpha_j - j$ and/or $\beta = -2\beta_j - j$ where $j, \alpha_j, \beta_j \in \mathbb{N}_0$, an analytic solution of Heun equation turns to be the first kind of independent solution of Heun polynomial which makes B_n term terminated. (2) if $\alpha = -2\alpha_j - j - 1 + \gamma$ and/or $\beta = -2\beta_j - j - 1 + \gamma$, an analytic solution of Heun equation turns to be the second kind of independent solution of Heun polynomial of type 1.

In this chapter I treat q as a fixed value and $\alpha, \beta, \gamma, \delta$ as free variables to construct Heun polynomial of type 2: (1) if $q = -(q_j + 2j)\{\alpha + \beta - \delta + a(\delta + \gamma - 1) + (1 + a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$, an analytic solution of Heun equation turns to be the first kind of independent solution of Heun polynomial of type 2. (2) if

$q = -(q_j + 2j + 1 - \gamma)\{\alpha + \beta + 1 - \gamma - (1 - a)\delta + (1 + a)(q_j + 2j)\}$, an analytic solution of Heun equation turns to be the second kind of independent solution of Heun polynomial of type 2.

2.2.2 Infinite series

In chapter 1, the general expression of power series of $y(x)$ for infinite series is

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (2.2.6)
\end{aligned}$$

Substitute (2.1.4a)-(2.1.4c) into (2.2.6). The general expression of power series of Heun equation for infinite series about $x = 0$ is

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)(i_0+\lambda+\beta)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)(i_0+\lambda+\beta)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+\lambda+\alpha)(i_k+2k+\lambda+\beta)}{(i_k+2(k+1)+\lambda)(i_k+2k+1+\gamma+\lambda)} \right. \\
&\times \left. \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1+\lambda)_{i_{k-1}} (2k+\gamma+\lambda)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1+\lambda)_{i_k} (2k+\gamma+\lambda)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1+\lambda)_{i_{n-1}} (2n+\gamma+\lambda)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} 3(2n+1+\lambda)_{i_n} (2n+\gamma+\lambda)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (2.2.7)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\{\varphi+2(1+a)\lambda\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_1^\pm = \frac{\{\varphi+2(1+a)(\lambda+2)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_k^\pm = \frac{\{\varphi+2(1+a)(\lambda+2k)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_n^\pm = \frac{\{\varphi+2(1+a)(\lambda+2n)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \end{cases}$$

Put $c_0 = 1$ as $\lambda=0$ for the first kind of independent solutions of Heun equation and $c_0 = \left(\frac{1+a}{a}\right)^{1-\gamma}$ as $\lambda = 1 - \gamma$ for the second one in (2.2.7).

Remark 2.2.3 *The power series expansion of Heun equation of the first kind for infinite*

series about $x = 0$ using R3TRF is

$$\begin{aligned}
y(x) &= HF^R \left(\varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1); \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
&= \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \alpha)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0}} \frac{(3)_{i_0} (2 + \gamma)_{i_0}}{(3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \alpha)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \alpha)(i_k + 2k + \beta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}}} \frac{(2k+1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(2k+1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}}} \frac{(2n+1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(2n+1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \tag{2.2.8}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(1+a)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(1+a)k\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(1+a)n\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

Remark 2.2.4 The power series expansion of Heun equation of the second kind for

infinite series about $x = 0$ using R3TRF is

$$\begin{aligned}
y(x) &= HS^R \left(\varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1); \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
&= \eta^{1-\gamma} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+1-\gamma+\alpha)(i_0+1-\gamma+\beta)}{(i_0+3-\gamma)(i_0+2)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (4-\gamma)_{i_0} (3)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (4-\gamma)_{i_1} (3)_{i_1}} \eta^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+1-\gamma+\alpha)(i_0+1-\gamma+\beta)}{(i_0+3-\gamma)(i_0+2)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+1-\gamma+\alpha)(i_k+2k+1-\gamma+\beta)}{(i_k+2k+3-\gamma)(i_k+2(k+1))} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2(k+1)-\gamma)_{i_{k-1}} (2k+1)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2(k+1)-\gamma)_{i_k} (2k+1)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2(n+1)-\gamma)_{i_{n-1}} (2n+1)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2(n+1)-\gamma)_{i_n} (2n+1)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (2.2.9)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\{\varphi+2(1+a)(1-\gamma)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_1^{\pm} = \frac{\{\varphi+2(1+a)(3-\gamma)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_k^{\pm} = \frac{\{\varphi+2(1+a)(1+2k-\gamma)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_n^{\pm} = \frac{\{\varphi+2(1+a)(1+2n-\gamma)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \end{cases}$$

It's required that $\gamma \neq 0, -1, -2, \dots$ for the first kind of independent solution of Heun function for all cases. Because if it does not, its solution will be divergent. And it's required that $\gamma \neq 2, 3, 4, \dots$ for the second kind of independent solution of Heun function for all cases.

The infinite series in this chapter are equivalent to the infinite series in Ref.[2]. In this chapter B_n is the leading term in sequence c_n in the analytic function $y(x)$. In Ref.[2] A_n is the leading term in sequence c_n in the analytic function $y(x)$.

2.3 Integral Formalism

2.3.1 Polynomial of type 2

Now let's investigate the integral formalism for the polynomial of type 2 at certain eigenvalue. There is a generalized hypergeometric function which is

$$\begin{aligned}
 I_l &= \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} \left(q_l + 4l + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} \left(q_l + 4l + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} \eta^{i_l} \\
 &= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+2l+\lambda, j+1) B(i_{l-1}+2l-1+\gamma+\lambda, j+1)}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-1+\gamma+\lambda)^{-1}} \\
 &\quad \times \frac{(i_{l-1}-q_l)_j \left(i_{l-1}+q_l+4l + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_j}{(1)_j j!} \eta^j
 \end{aligned} \tag{2.3.1}$$

By using integral form of beta function,

$$B(i_{l-1}+2l+\lambda, j+1) = \int_0^1 dt_l t_l^{i_{l-1}+2l-1+\lambda} (1-t_l)^j \tag{2.3.2a}$$

$$B(i_{l-1}+2l-1+\gamma+\lambda, j+1) = \int_0^1 du_l u_l^{i_{l-1}+2(l-1)+\gamma+\lambda} (1-u_l)^j \tag{2.3.2b}$$

Substitute (2.3.2a) and (2.3.2b) into (2.3.1). And divide $(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)$ into I_l .

$$\begin{aligned}
 K_l &= \frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)} \\
 &\quad \times \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} \left(q_l + 4l + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} \left(q_l + 4l + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} \eta^{i_l} \\
 &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} (\eta t_l u_l)^{i_{l-1}} \\
 &\quad \times \sum_{j=0}^{\infty} \frac{(i_{l-1}-q_l)_j \left(i_{l-1}+q_l+4l + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_j}{(1)_j j!} [\eta(1-t_l)(1-u_l)]^j
 \end{aligned}$$

The integral form of Gauss hypergeometric function is

$$\begin{aligned}
{}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n (n!)} z^n \\
&= -\frac{1}{2\pi i} \frac{\Gamma(1-\alpha)\Gamma(\gamma)}{\Gamma(\gamma-\alpha)} \oint dv_l (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \quad (2.3.3) \\
&\quad \text{where } \operatorname{Re}(\gamma-\alpha) > 0
\end{aligned}$$

Replace α, β, γ and z by $i_{l-1} - q_l, i_{l-1} + q_l + 4l + \frac{\varphi + 2(1+a)\lambda}{(1+a)}, 1$ and $\eta(1-t_l)(1-u_l)$ in (2.3.3).

$$\begin{aligned}
&\sum_{j=0}^{\infty} \frac{(i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + \frac{\varphi + 2(1+a)\lambda}{(1+a)})_j}{(1)_j j!} [\eta(1-t_l)(1-u_l)]^j \\
&= \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \frac{1}{1 - \eta(1-t_l)(1-u_l)v_l} \right)^{q_l} (1 - \eta(1-t_l)(1-u_l)v_l)^{-\left(4l + \frac{\varphi + 2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \left(\frac{v_l}{v_l - 1} \frac{1}{1 - \eta(1-t_l)(1-u_l)v_l} \right)^{i_{l-1}} \quad (2.3.4)
\end{aligned}$$

Substitute (2.3.4) into K_l .

$$\begin{aligned}
K_l &= \frac{1}{(i_{l-1} + 2l + \lambda)(i_{l-1} + 2l - 1 + \gamma + \lambda)} \\
&\quad \times \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} \left(q_l + 4l + \frac{\varphi + 2(1+a)\lambda}{(1+a)} \right)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \gamma + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} \left(q_l + 4l + \frac{\varphi + 2(1+a)\lambda}{(1+a)} \right)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \gamma + \lambda)_{i_l}} \eta^{i_l} \\
&= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \frac{1}{1 - \eta(1-t_l)(1-u_l)v_l} \right)^{q_l} \\
&\quad \times (1 - \eta(1-t_l)(1-u_l)v_l)^{-\left(4l + \frac{\varphi + 2(1+a)\lambda}{(1+a)}\right)} \left(\frac{v_l}{v_l - 1} \frac{\eta t_l u_l}{1 - \eta(1-t_l)(1-u_l)v_l} \right)^{i_{l-1}} \quad (2.3.5)
\end{aligned}$$

Substitute (2.3.5) into (2.2.3) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(x)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(x)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(x)$, etc.⁴

⁴ $y_1(x)$ means the sub-power series in (2.2.3) contains one term of $B'_n s$, $y_2(x)$ means the sub-power series in (2.2.3) contains two terms of $B'_n s$, $y_3(x)$ means the sub-power series in (2.2.3) contains three terms of $B'_n s$, etc.

Theorem 2.3.1 *The general representation in the form of integral of Heun polynomial of type 2 is given by*

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\left(4(n-k) + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta+\lambda} \left. \right\} \\
&\quad \times \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \left. \right\} z^n \Bigg\} \tag{2.3.6}
\end{aligned}$$

where

$$\overleftrightarrow{w}_{i,j} = \begin{cases} \frac{v_i}{(v_i-1)} \frac{\overleftrightarrow{w}_{i+1,j} t_i u_i}{1 - \overleftrightarrow{w}_{i+1,j} v_i (1-t_i)(1-u_i)} & \text{where } i \leq j \\ \eta & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof According to (2.2.3),

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \tag{2.3.7}$$

In the above, sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of Heun polynomial which makes A_n term terminated about $x = 0$ are

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \tag{2.3.8a}$$

$$\begin{aligned}
y_1(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}}{(-q_1)_{i_0} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}} \frac{(3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \eta^{i_1} \right\} z \quad (2.3.8b)
\end{aligned}$$

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{q_1} \frac{(i_1 + 2 + \lambda + \alpha)(i_1 + 2 + \lambda + \beta)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(-q_1)_{i_1} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}}{(-q_1)_{i_0} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}} \frac{(3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\
&\quad \times \left. \sum_{i_2=i_1}^{q_2} \frac{(-q_2)_{i_2} \left(q_2 + 8 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_2}}{(-q_2)_{i_1} \left(q_2 + 8 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}} \frac{(5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} \eta^{i_2} \right\} z^2 \quad (2.3.8c)
\end{aligned}$$

$$\begin{aligned}
y_3(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{q_1} \frac{(i_1 + 2 + \lambda + \alpha)(i_1 + 2 + \lambda + \beta)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(-q_1)_{i_1} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}}{(-q_1)_{i_0} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}} \frac{(3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\
&\quad \times \sum_{i_2=i_1}^{q_2} \frac{(i_2 + 4 + \lambda + \alpha)(i_2 + 4 + \lambda + \beta)}{(i_2 + \lambda + 6)(i_2 + \lambda + 5 + \gamma)} \frac{(-q_2)_{i_2} \left(q_2 + 8 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_2}}{(-q_2)_{i_1} \left(q_2 + 8 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}} \frac{(5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} \\
&\quad \times \left. \sum_{i_3=i_2}^{q_3} \frac{(-q_3)_{i_3} \left(q_2 + 12 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_3}}{(-q_3)_{i_2} \left(q_2 + 12 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_2}} \frac{(7 + \lambda)_{i_2} (6 + \gamma + \lambda)_{i_2}}{(7 + \lambda)_{i_3} (6 + \gamma + \lambda)_{i_3}} \eta^{i_3} \right\} z^3 \quad (2.3.8d)
\end{aligned}$$

Put $l = 1$ in (2.3.5). Take the new (2.3.5) into (2.3.8b).

$$\begin{aligned}
y_1(x) &= \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - \eta(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \\
&\quad \times (1 - \eta(1 - t_1)(1 - u_1)v_1)^{-\left(4 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\beta+\lambda} \\
&\quad \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right\} z
\end{aligned} \tag{2.3.9}$$

where

$$\overleftrightarrow{w}_{1,1} = \frac{v_1}{v_1 - 1} \frac{\eta t_1 u_1}{1 - \eta(1 - t_1)(1 - u_1)v_1}$$

Put $l = 2$ in (2.3.5). Take the new (2.3.5) into (2.3.8c).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - \eta(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\
&\quad \times (1 - \eta(1 - t_2)(1 - u_2)v_2)^{-\left(8 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha-\beta} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2+\beta+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}}{(-q_1)_{i_0} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}} \frac{(3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \overleftrightarrow{w}_{2,2}^{i_1} \right\} z^2
\end{aligned} \tag{2.3.10}$$

where

$$\overleftrightarrow{w}_{2,2} = \frac{v_2}{v_2 - 1} \frac{\eta t_2 u_2}{1 - \eta(1 - t_2)(1 - u_2)v_2}$$

Put $l = 1$ and $\eta = \overleftarrow{w}_{2,2}$ in (1.4.15). Take the new (1.4.15) into (1.4.24).

$$\begin{aligned}
y_2(x) = & \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - \eta(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\
& \times (1 - \eta(1 - t_2)(1 - u_2)v_2)^{-\left(8 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,2} \partial \overleftarrow{w}_{2,2} \right) \overleftarrow{w}_{2,2}^{\alpha-\beta} \left(\overleftarrow{w}_{2,2} \partial \overleftarrow{w}_{2,2} \right) \overleftarrow{w}_{2,2}^{2+\beta+\lambda} \\
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \\
& \times (1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1)^{-\left(4 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,2} \partial \overleftarrow{w}_{1,2} \right) \overleftarrow{w}_{1,2}^{\alpha-\beta} \left(\overleftarrow{w}_{1,2} \partial \overleftarrow{w}_{1,2} \right) \overleftarrow{w}_{1,2}^{\beta+\lambda} \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,2}^{i_0} \right\} z^2 \tag{2.3.11}
\end{aligned}$$

where

$$\overleftarrow{w}_{1,2} = \frac{v_1}{v_1 - 1} \frac{\overleftarrow{w}_{2,2} t_1 u_1}{1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1}$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3 - 1}{v_3} \frac{1}{1 - \eta(1 - t_3)(1 - u_3)v_3} \right)^{q_3} \\
& \times (1 - \eta(1 - t_3)(1 - u_3)v_3)^{-\left(12 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{3,3}^{-(4+\alpha+\lambda)} \left(\overleftarrow{w}_{3,3} \partial \overleftarrow{w}_{3,3} \right) \overleftarrow{w}_{3,3}^{\alpha-\beta} \left(\overleftarrow{w}_{3,3} \partial \overleftarrow{w}_{3,3} \right) \overleftarrow{w}_{3,3}^{4+\beta+\lambda} \\
& \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - \overleftarrow{w}_{3,3}(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\
& \times (1 - \overleftarrow{w}_{3,3}(1 - t_2)(1 - u_2)v_2)^{-\left(8 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{2,3}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,3} \partial \overleftarrow{w}_{2,3} \right) \overleftarrow{w}_{2,3}^{\alpha-\beta} \left(\overleftarrow{w}_{2,3} \partial \overleftarrow{w}_{2,3} \right) \overleftarrow{w}_{2,3}^{2+\beta+\lambda}
\end{aligned}$$

$$\begin{aligned}
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - \overleftrightarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
& \times (1 - \overleftrightarrow{w}_{2,3}(1-t_1)(1-u_1)v_1)^{-\left(4 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftrightarrow{w}_{1,3}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,3} \partial_{\overleftrightarrow{w}_{1,3}} \right) \overleftrightarrow{w}_{1,3}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,3} \partial_{\overleftrightarrow{w}_{1,3}} \right) \overleftrightarrow{w}_{1,3}^{\beta+\lambda} \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,3}^{i_0} \right\} z^3 \tag{2.3.12}
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{w}_{3,3} = \frac{v_3}{v_3-1} \frac{\eta t_3 u_3}{1-\eta(1-t_3)(1-u_3)v_3} \\ \overleftrightarrow{w}_{2,3} = \frac{v_2}{v_2-1} \frac{\overleftrightarrow{w}_{3,3} t_2 u_2}{1-\overleftrightarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \\ \overleftrightarrow{w}_{1,3} = \frac{v_1}{v_1-1} \frac{\overleftrightarrow{w}_{2,3} t_1 u_1}{1-\overleftrightarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, I obtain every integral forms of $y_m(x)$ terms. Since we substitute (2.3.8a), (2.3.9), (2.3.11), (2.3.12) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (2.3.7), We obtain (2.3.6). ■

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of Heun equation and $c_0 = \left(\frac{1+a}{a} \right)^{1-\gamma}$ as $\lambda = 1 - \gamma$ for the second one in (2.3.6).

Remark 2.3.2 *The integral representation of Heun equation of the first kind for polynomial of type 2 about $x = 0$ as*

$q = -(q_j + 2j)\{\alpha + \beta - \delta + a(\delta + \gamma - 1) + (1 + a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= HF_{q_j}^R \left(q_j = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j, \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1) \right. \\
&\quad \left. , \Omega_1 = \frac{\varphi}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
&= {}_2F_1(-q_0, q_0 + \Omega_1; \gamma; \eta) \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega_1)} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta} \Big\} \\
&\quad \left. \times {}_2F_1(-q_0, q_0 + \Omega_1; \gamma; \overleftrightarrow{w}_{1,n}) \right\} z^n \tag{2.3.13}
\end{aligned}$$

Remark 2.3.3 The integral representation of Heun equation of the second kind for polynomial of type 2 about $x = 0$ as

$q = -(q_j + 2j + 1 - \gamma)\{\alpha + \beta + 1 - \gamma - (1-a)\delta + (1+a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= HS_{q_j}^R \left(q_j = \frac{-\{\varphi + 2(1+a)(1-\gamma)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j \right. \\
&\quad \left. , \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_2 = \frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
&= \eta^{1-\gamma} \left\{ {}_2F_1(-q_0, q_0 + \Omega_2; 2 - \gamma; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega_2)} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\gamma} \Big\} \\
&\quad \left. \times {}_2F_1(-q_0, q_0 + \Omega_2; 2 - \gamma; \overleftrightarrow{w}_{1,n}) \right\} z^n \Big\} \tag{2.3.14}
\end{aligned}$$

2.3.2 Infinite series

Let's consider the integral representation of Heun equation about $x = 0$ for infinite series by applying R3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned} M_l &= \sum_{i_l=i_{l-1}}^{\infty} \frac{(\Delta_l^-)_{i_l} (\Delta_l^+)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(\Delta_l^-)_{i_{l-1}} (\Delta_l^+)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} \eta^{i_l} \\ &= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+2l+\lambda, j+1) B(i_{l-1}+2l-1+\gamma+\lambda, j+1) (\Delta_l^- + i_{l-1})_j (\Delta_l^+ + i_{l-1})_j}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-1+\gamma+\lambda)^{-1} (1)_j j!} \eta^j \end{aligned} \quad (2.3.15)$$

where

$$\Delta_l^{\pm} = \frac{\varphi + 2(1+a)(\lambda+2l) \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)}$$

Substitute (2.3.2a) and (2.3.2b) into (2.3.15). Divide $(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)$ into the new (2.3.15).

$$\begin{aligned} V_l &= \frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\Delta_l^-)_{i_l} (\Delta_l^+)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(\Delta_l^-)_{i_{l-1}} (\Delta_l^+)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} (\eta t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(\Delta_l^- + i_{l-1})_j (\Delta_l^+ + i_{l-1})_j (\eta(1-t_l)(1-u_l))^j}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-1+\gamma+\lambda)^{-1} (1)_j j!} \end{aligned}$$

The hypergeometric function is defined by

$$\begin{aligned} {}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n (n!)} z^n \\ &= \frac{1}{2\pi i} \frac{\Gamma(1+\alpha-\gamma)}{\Gamma(\alpha)} \int_0^{(1+)} dv_l (-1)^\gamma (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \quad (2.3.16) \\ &\text{where } \gamma - \alpha \neq 1, 2, 3, \dots, \operatorname{Re}(\alpha) > 0 \end{aligned}$$

Replace α, β, γ and z by $\Delta_l^- + i_{l-1}, \Delta_l^+ + i_{l-1}, 1$ and $\eta(1-t_l)(1-u_l)$ in (2.3.16). Take the new (2.3.16) into V_l .

$$\begin{aligned} V_l &= \frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\Delta_l^-)_{i_l} (\Delta_l^+)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(\Delta_l^-)_{i_{l-1}} (\Delta_l^+)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l-1}{v_l} \right)^{-\Delta_l^-} \\ &\quad \times (1-\eta(1-t_l)(1-u_l)v_l)^{-\Delta_l^+} \left(\frac{v_l}{v_l-1} \frac{\eta t_l u_l}{1-\eta(1-t_l)(1-u_l)v_l} \right)^{i_{l-1}} \end{aligned} \quad (2.3.17)$$

Substitute (2.3.17) into (2.2.7) where $l = 1, 2, 3, \dots$; apply V_1 into the second summation of sub-power series $y_1(x)$, apply V_2 into the third summation and V_1 into the second summation of sub-power series $y_2(x)$, apply V_3 into the forth summation, V_2 into the third summation and V_1 into the second summation of sub-power series $y_3(x)$, etc.⁵

Theorem 2.3.4 *The general representation in the form of integral of Heun equation for infinite series about $x = 0$ using R3TRF is given by*

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k}) v_{n-k})^{-\Delta_{n-k}^+} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta+\lambda} \left. \right\} \\
&\quad \times \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \left. \right\} z^n \Bigg\} \tag{2.3.18}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi+2(1+a)\lambda \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_{n-k}^\pm = \frac{\varphi+2(1+a)(\lambda+2(n-k)) \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \end{cases}$$

In the above, the first sub-integral form contains one term of $B'_n s$, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof In (2.2.7) sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of Heun equation for infinite series about $x = 0$ using R3TRF are given by

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \tag{2.3.19a}$$

⁵ $y_1(x)$ means the sub-power series in (2.2.7) contains one term of $B'_n s$, $y_2(x)$ means the sub-power series in (2.2.7) contains two terms of $B'_n s$, $y_3(x)$ means the sub-power series in (2.2.7) contains three terms of $B'_n s$, etc.

$$\begin{aligned}
y_1(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
& \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \eta^{i_1} \right\} z \quad (2.3.19b)
\end{aligned}$$

$$\begin{aligned}
y_2(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + 2 + \lambda + \alpha)(i_1 + 2 + \lambda + \beta)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\
& \times \left. \sum_{i_2=i_1}^{\infty} \frac{(\Delta_2^-)_{i_2} (\Delta_2^+)_{i_2} (5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(\Delta_2^-)_{i_1} (\Delta_2^+)_{i_1} (5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} \eta^{i_2} \right\} z^2 \quad (2.3.19c)
\end{aligned}$$

$$\begin{aligned}
y_3(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + 2 + \lambda + \alpha)(i_1 + 2 + \lambda + \beta)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\
& \times \sum_{i_2=i_1}^{\infty} \frac{(i_2 + 4 + \lambda + \alpha)(i_2 + 4 + \lambda + \beta)}{(i_2 + \lambda + 6)(i_2 + \lambda + 5 + \gamma)} \frac{(\Delta_2^-)_{i_2} (\Delta_2^+)_{i_2} (5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(\Delta_2^-)_{i_1} (\Delta_2^+)_{i_1} (5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} \\
& \times \left. \sum_{i_3=i_2}^{\infty} \frac{(\Delta_3^-)_{i_3} (\Delta_3^+)_{i_3} (7 + \lambda)_{i_2} (6 + \gamma + \lambda)_{i_2}}{(\Delta_3^-)_{i_2} (\Delta_3^+)_{i_2} (7 + \lambda)_{i_3} (6 + \gamma + \lambda)_{i_3}} \eta^{i_3} \right\} z^3 \quad (2.3.19d)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi+2(1+a)\lambda \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_1^\pm = \frac{\varphi+2(1+a)(\lambda+2) \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_2^\pm = \frac{\varphi+2(1+a)(\lambda+4) \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_3^\pm = \frac{\varphi+2(1+a)(\lambda+6) \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \end{cases}$$

Put $l = 1$ in (2.3.17). Take the new (2.3.17) into (2.3.19b).

$$\begin{aligned}
y_1(x) &= \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \right)^{-\Delta_1^-} \\
&\quad \times (1 - \eta(1 - t_1)(1 - u_1)v_1)^{-\Delta_1^+} \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\beta+\lambda} \\
&\quad \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right\} z
\end{aligned} \tag{2.3.20}$$

where

$$\overleftrightarrow{w}_{1,1} = \frac{v_1}{v_1 - 1} \frac{\eta t_1 u_1}{1 - \eta(1 - t_1)(1 - u_1)v_1}$$

Put $l = 2$ in (2.3.17). Take the new (2.3.17) into (2.3.19c).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \right)^{-\Delta_2^-} \\
&\quad \times (1 - \eta(1 - t_2)(1 - u_2)v_2)^{-\Delta_2^+} \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha-\beta} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2+\beta+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)(i_0 + \lambda + \beta)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \overleftrightarrow{w}_{2,2}^{i_1} \right\} z^2
\end{aligned} \tag{2.3.21}$$

where

$$\overleftrightarrow{w}_{2,2} = \frac{v_2}{v_2 - 1} \frac{\eta t_2 u_2}{1 - \eta(1 - t_2)(1 - u_2)v_2}$$

Put $l = 1$ and $\eta = \overleftrightarrow{w}_{2,2}$ in (2.3.17). Take the new (2.3.17) into (2.3.21).

$$\begin{aligned}
y_2(x) &= \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \right)^{-\Delta_2^-} \\
&\quad \times (1 - \eta(1 - t_2)(1 - u_2)v_2)^{-\Delta_2^+} \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha-\beta} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2+\beta+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \right)^{-\Delta_1^-} \\
&\quad \times (1 - \overleftrightarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1)^{-\Delta_1^+} \overleftrightarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\beta+\lambda} \\
&\quad \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right\} z^2
\end{aligned} \tag{2.3.22}$$

where

$$\overleftrightarrow{w}_{1,2} = \frac{v_1}{v_1 - 1} \frac{\overleftrightarrow{w}_{2,2} t_1 u_1}{1 - \overleftrightarrow{w}_{2,2} (1 - t_1) (1 - u_1) v_1}$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned} y_3(x) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3 - 1}{v_3} \right)^{-\Delta_3^-} \\ & \times (1 - \eta(1 - t_3)(1 - u_3)v_3)^{-\Delta_3^+} \overleftrightarrow{w}_{3,3}^{-(4+\alpha+\lambda)} \left(\overleftrightarrow{w}_{3,3} \partial_{\overleftrightarrow{w}_{3,3}} \right) \overleftrightarrow{w}_{3,3}^{\alpha-\beta} \left(\overleftrightarrow{w}_{3,3} \partial_{\overleftrightarrow{w}_{3,3}} \right) \overleftrightarrow{w}_{3,3}^{4+\beta+\lambda} \\ & \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \right)^{-\Delta_2^-} \\ & \times (1 - \overleftrightarrow{w}_{3,3}(1 - t_2)(1 - u_2)v_2)^{-\Delta_2^+} \overleftrightarrow{w}_{2,3}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,3} \partial_{\overleftrightarrow{w}_{2,3}} \right) \overleftrightarrow{w}_{2,3}^{\alpha-\beta} \left(\overleftrightarrow{w}_{2,3} \partial_{\overleftrightarrow{w}_{2,3}} \right) \overleftrightarrow{w}_{2,3}^{2+\beta+\lambda} \\ & \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \right)^{-\Delta_1^-} \\ & \times (1 - \overleftrightarrow{w}_{2,3}(1 - t_1)(1 - u_1)v_1)^{-\Delta_1^+} \overleftrightarrow{w}_{1,3}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,3} \partial_{\overleftrightarrow{w}_{1,3}} \right) \overleftrightarrow{w}_{1,3}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,3} \partial_{\overleftrightarrow{w}_{1,3}} \right) \overleftrightarrow{w}_{1,3}^{\beta+\lambda} \\ & \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,3}^{i_0} \right\} z^3 \end{aligned} \quad (2.3.23)$$

where

$$\begin{cases} \overleftrightarrow{w}_{3,3} = \frac{v_3}{v_3-1} \frac{\eta t_3 u_3}{1-\eta(1-t_3)(1-u_3)v_3} \\ \overleftrightarrow{w}_{2,3} = \frac{v_2}{v_2-1} \frac{\overleftrightarrow{w}_{3,3} t_2 u_2}{1-\overleftrightarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \\ \overleftrightarrow{w}_{1,3} = \frac{v_1}{v_1-1} \frac{\overleftrightarrow{w}_{2,3} t_1 u_1}{1-\overleftrightarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, I obtain every integral forms of $y_m(x)$ terms. Since we substitute (2.3.19a), (2.3.20), (2.3.22), (2.3.23) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (2.2.7), we obtain (2.3.18).⁶

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of Heun equation and

$$c_0 = \left(\frac{1+a}{a} \right)^{1-\gamma} \text{ as } \lambda = 1 - \gamma \text{ for the second one in (2.3.18).}$$

⁶Or replace the finite summation with an interval $[0, q_0]$ by infinite summation with an interval $[0, \infty]$ in (2.3.6). Replace q_0 and q_{n-k} by $\frac{-(\varphi+2(1+a)\lambda)+\sqrt{\varphi^2-4(1+a)q}}{2(1+a)}$ and $\frac{-(\varphi+2(1+a)(\lambda+2(n-k)))+\sqrt{\varphi^2-4(1+a)q}}{2(1+a)}$ into the new (2.3.6). Its solution is also equivalent to (2.3.18).

Remark 2.3.5 *The integral representation of Heun equation of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
y(x) &= HF^R \left(\varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1); \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
&= {}_2F_1 \left(\Delta_0^-, \Delta_0^+; \gamma; \eta \right) \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-\Delta_{n-k}^+} \\
&\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)+\beta} \Big\} \\
&\quad \times {}_2F_1 \left(\Delta_0^-, \Delta_0^+; \gamma; \overleftarrow{w}_{1,n} \right) \Big\} z^n \tag{2.3.24}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_{n-k}^{\pm} = \frac{\varphi + 4(1+a)(n-k) \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

Remark 2.3.6 *The integral representation of Heun equation of the second kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
y(x) &= HS^R \left(\varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1); \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
&= \eta^{1-\gamma} \left\{ {}_2F_1 \left(\Delta_0^-, \Delta_0^+; 2 - \gamma; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-\Delta_{n-k}^+} \\
&\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\gamma} \Big\} \\
&\quad \times {}_2F_1 \left(\Delta_0^-, \Delta_0^+; 2 - \gamma; \overleftarrow{w}_{1,n} \right) \Big\} \Big\} z^n \tag{2.3.25}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi + 2(1+a)(1-\gamma) \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_{n-k}^{\pm} = \frac{\varphi + 2(1+a)(2(n-k)+1-\gamma) \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

2.4 Summary

In mathematical definition, Heun polynomial is the polynomial of type 3: for α or $\beta = -N - \lambda$ where $N \in \mathbb{N}_0$ and $\lambda = 0$ or $1 - \gamma$ there will be a polynomial equation of degree $N + 1$ for the determination q about $x = 0$.

In Ref.[2, 3] I show how to obtain the power series expansion in closed forms and its integral forms of Heun functions (infinite series and polynomial of type 1) including all higher terms of A_n 's by applying 3TRF. This was done by letting A_n in sequence c_n is the leading term in the analytic function $y(x)$: the sequence c_n consists of combinations A_n and B_n . For polynomial of type 1, I treat γ , δ and q as free variables and fixed values of α and/or β .

In this chapter I show how to construct the Frobenius solutions in closed forms of Heun equation and its integral forms for infinite series and polynomial of type 2 including all higher terms of B_n 's by applying R3TRF. This is done by letting B_n in sequence c_n is the leading term in the analytic function $y(x)$. For polynomial of type 2, I treat α , β , γ and δ as free variables and a fixed value of q .

The power series expansion of Heun equation and its integral form for infinite series about $x = 0$ in this chapter are equivalent to infinite series of Heun equation in Ref.[2, 3]. In this chapter B_n is the leading term in sequence c_n in the analytic function $y(x)$. In Ref.[2, 3] A_n is the leading term in sequence c_n in the analytic function $y(x)$.

In Ref.[2, 3] as we see the power series expansions of Heun equation for all cases of infinite series and polynomial, the denominators and numerators in all B_n terms of Heun equation about $x = 0$ arise with Pochhammer symbol. Again in this chapter the denominators and numerators in all A_n terms of Heun function about $x = 0$ arise with Pochhammer symbol. Since we construct the power series expansions with Pochhammer symbols in numerators and denominators, we are able to describe integral forms of Heun equation in simple way. As we observe representations in closed form integrals of Heun equation by applying either 3TRF or R3TRF, a ${}_2F_1$ function recurs in each of sub-integral forms of Heun function. We can transform Heun function into any special functions with two recursive coefficients because of ${}_2F_1$ functions in integral forms of Heun function. After we replace ${}_2F_1$ functions in integral forms of Heun function to other special functions, we can rebuild the power series expansion of Heun function in a backward.

Appendices

.1 Power series expansion of 192 Heun functions

In this chapter, the fundamental power series expansion in closed forms and its integral forms of Heun function about $x = 0$ is constructed by using R3TRF. A machine-generated list of 192 (isomorphic to the Coxeter group of the Coxeter diagram D_4) local solutions of the Heun equation was obtained by Robert S. Maier(2007) [5]. In appendix of Ref.[3], applying 3TRF, I analyzed the power series expansions and its integral forms of Heun function (infinite series and polynomial of type 1) of nine out of the 192 local solutions of Heun function in Table 2 [5].

In these appendices 1 and 2, replacing coefficients in the general expression of power series expansion and integral representation of Heun function for polynomial and infinite series, I derive the analytic solutions for the power series expansions and its integrals (infinite series and polynomial of type 2) of the previous nine examples of the 192 local solutions of Heun function in Ref.[3].⁷

⁷I treat α , β , γ and δ as free variables and a fixed value of q to construct the polynomial of type 2 and its integral formalism for all nine examples of the 192 local solutions of Heun function.

$$\mathbf{.1.1} \quad (1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x)$$

Polynomial of type 2

Replacing coefficients $q, \alpha, \beta, \delta, c_0$ and λ by $q - (\delta-1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$ and zero into (2.2.3). Multiply $(1-x)^{1-\delta}$ and the new (2.2.3) together.

$$\begin{aligned}
& (1-x)^{1-\delta} y(x) \\
&= (1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x) \\
&= (1-x)^{1-\delta} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + 1 + \alpha - \delta)(i_0 + 1 + \beta - \delta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
&\times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + 1 + \alpha - \delta)(i_0 + 1 + \beta - \delta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + 1 + \alpha - \delta)(i_k + 2k + 1 + \beta - \delta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \right. \\
&\times \left. \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.1)
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - \delta + a(\gamma - \delta + 1) \\ \Omega = \frac{\varphi}{(1+a)} \\ q = (\delta - 1)\gamma a - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $q, \alpha, \beta, \delta, c_0$ and λ by $q - (\delta - 1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$ and zero into (2.2.7). Multiply $(1 - x)^{1-\delta}$ and the new (2.2.7) together.

$$\begin{aligned}
& (1 - x)^{1-\delta} y(x) \\
&= (1 - x)^{1-\delta} Hl(a, q - (\delta - 1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x) \\
&= (1 - x)^{1-\delta} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \alpha - \delta)(i_0 + 1 + \beta - \delta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \alpha - \delta)(i_0 + 1 + \beta - \delta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + 1 + \alpha - \delta)(i_k + 2k + 1 + \beta - \delta)}{(i_k + 2(k + 1))(i_k + 2k + 1 + \gamma)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.2)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(1+a)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(1+a)k\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(1+a)n\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

.1.2

$$x^{1-\gamma} (1 - x)^{1-\delta} Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x)$$

Polynomial of type 2

Replacing coefficients $q, \alpha, \beta, \gamma, \delta, c_0$ and λ by

$$q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$$

and zero into (2.2.3). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and the new (2.2.3) together.

$$\begin{aligned}
& x^{1-\gamma}(1-x)^{1-\delta}y(x) \\
= & x^{1-\gamma}(1-x)^{1-\delta}Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2 \\
& , \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x) \\
= & x^{1-\gamma}(1-x)^{1-\delta} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (2 - \gamma)_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + 2 + \alpha - \gamma - \delta)(i_0 + 2 + \beta - \gamma - \delta)}{(i_0 + 2)(i_0 + 3 - \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (2 - \gamma)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_0} (4 - \gamma)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (4 - \gamma)_{i_1}} \eta^{i_1} \Big\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + 2 + \alpha - \gamma - \delta)(i_0 + 2 + \beta - \gamma - \delta)}{(i_0 + 2)(i_0 + 3 - \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (2 - \gamma)_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + 2 + \alpha - \gamma - \delta)(i_k + 2k + 2 + \beta - \gamma - \delta)}{(i_k + 2(k+1))(i_k + 2k + 3 - \gamma)} \right. \\
& \times \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + 2 - \gamma)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + 2 - \gamma)_{i_k}} \Big\} \\
& \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + 2 - \gamma)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + 2 - \gamma)_{i_n}} \eta^{i_n} \Big\} z^n \Big\} \quad (.1.3)
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - 2\gamma - \delta + 2 + a(3 - \gamma - \delta) \\ \Omega = \frac{\varphi}{(1+a)} \\ q = (\gamma + \delta - 2)a + (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1) \\ \quad - (q_j + 2j)\{\varphi + (1 + a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $q, \alpha, \beta, \gamma, \delta, c_0$ and λ by

$$q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$$

and zero into (2.2.7). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and the new (2.2.7) together.

$$\begin{aligned}
& x^{1-\gamma}(1-x)^{1-\delta}y(x) \\
= & x^{1-\gamma}(1-x)^{1-\delta}Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2 \\
& , \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x) \\
= & x^{1-\gamma}(1-x)^{1-\delta} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (2 - \gamma)_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 2 + \alpha - \gamma - \delta)(i_0 + 2 + \beta - \gamma - \delta)}{(i_0 + 2)(i_0 + 3 - \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (2 - \gamma)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (4 - \gamma)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (4 - \gamma)_{i_1}} \eta^{i_1} \right\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 2 + \alpha - \gamma - \delta)(i_0 + 2 + \beta - \gamma - \delta)}{(i_0 + 2)(i_0 + 3 - \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (2 - \gamma)_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + 2 + \alpha - \gamma - \delta)(i_k + 2k + 2 + \beta - \gamma - \delta)}{(i_k + 2(k + 1))(i_k + 2k + 3 - \gamma)} \right. \\
& \times \left. \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + 2 - \gamma)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + 2 - \gamma)_{i_k}} \right\} \\
& \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + 2 - \gamma)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + 2 - \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.4)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(1+a)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(1+a)k\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(1+a)n\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

$$\mathbf{.1.3} \quad Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x)$$

Polynomial of type 2

Replacing coefficients $a, q, \gamma, \delta, x, c_0$ and λ by $1-a, -q + \alpha\beta, \delta, \gamma, 1-x, 1$ and zero into (2.2.3).

$$\begin{aligned}
y(\xi) &= Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x) \\
&= \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \eta^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_1} (2 + \delta)_{i_1}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (2 + \delta)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha)(i_k + 2k + \beta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \delta)} \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + \delta)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + \delta)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + \delta)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + \delta)_{i_n}} \eta^{i_n} \right\} z^n \quad (.1.5)
\end{aligned}$$

where

$$\begin{cases} \xi = 1 - x \\ z = \frac{-1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = \alpha + \beta - \delta + (1-a)(\delta + \gamma - 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = \alpha\beta + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \gamma, \delta, x, c_0$ and λ by $1-a, -q+\alpha\beta, \delta, \gamma, 1-x, 1$ and zero into (2.2.7).

$$\begin{aligned}
y(\xi) &= Hl(1-a, -q+\alpha\beta; \alpha, \beta, \delta, \gamma; 1-x) \\
&= \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \eta^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\alpha)(i_0+\beta)}{(i_0+2)(i_0+1+\delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2+\delta)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2+\delta)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\alpha)(i_0+\beta)}{(i_0+2)(i_0+1+\delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+\alpha)(i_k+2k+\beta)}{(i_k+2(k+1))(i_k+2k+1+\delta)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1)_{i_{k-1}} (2k+\delta)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1)_{i_k} (2k+\delta)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1)_{i_{n-1}} (2n+\delta)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1)_{i_n} (2n+\delta)_{i_n}} \eta^{i_n} \right\} z^n \quad (.1.6)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(2-a)\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(2-a)k\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(2-a)n\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \end{cases}$$

$$.1.4 \quad (1-x)^{1-\delta} Hl(1-a, -q+(\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x)$$

Polynomial of type 2

Replacing coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0$ and λ by $1-a, -q+(\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma, 1-x, 1$ and zero

into (2.2.3). Multiply $(1-x)^{1-\delta}$ and the new (2.2.3) together.

$$\begin{aligned}
& (1-x)^{1-\delta} y(\xi) \\
= & (1-x)^{1-\delta} Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1 \\
& , \beta-\delta+1, 2-\delta, \gamma; 1-x) \\
= & (1-x)^{1-\delta} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (2-\delta)_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0+1+\alpha-\delta)(i_0+1+\beta-\delta)}{(i_0+2)(i_0+3-\delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (2-\delta)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1+4+\Omega)_{i_1} (3)_{i_0} (4-\delta)_{i_0}}{(-q_1)_{i_0} (q_1+4+\Omega)_{i_0} (3)_{i_1} (4-\delta)_{i_1}} \eta^{i_1} \Big\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0+1+\alpha-\delta)(i_0+1+\beta-\delta)}{(i_0+2)(i_0+3-\delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (2-\delta)_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k+2k+1+\alpha-\delta)(i_k+2k+1+\beta-\delta)}{(i_k+2(k+1))(i_k+2k+3-\delta)} \right. \\
& \times \frac{(-q_k)_{i_k} (q_k+4k+\Omega)_{i_k} (2k+1)_{i_{k-1}} (2k+2-\delta)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k+4k+\Omega)_{i_{k-1}} (2k+1)_{i_k} (2k+2-\delta)_{i_k}} \Big\} \\
& \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n+4n+\Omega)_{i_n} (2n+1)_{i_{n-1}} (2n+2-\delta)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n+4n+\Omega)_{i_{n-1}} (2n+1)_{i_n} (2n+2-\delta)_{i_n}} \eta^{i_n} \Big\} z^n \Big\} \quad (.1.7)
\end{aligned}$$

where

$$\begin{cases} \xi = 1-x \\ z = \frac{-1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = \alpha + \beta - \gamma - 2\delta + 2 + (1-a)(\gamma - \delta + 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1) \\ \quad + (q_j + 2j) \{ \varphi + (2-a)(q_j + 2j) \} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0$ and λ by $1 - a, -q + (\delta - 1)\gamma a + (\alpha - \delta + 1)(\beta - \delta + 1), \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, \gamma, 1 - x, 1$ and zero into (2.2.7). Multiply $(1 - x)^{1-\delta}$ and the new (2.2.7) together.

$$\begin{aligned}
 & (1 - x)^{1-\delta} y(\xi) \\
 = & (1 - x)^{1-\delta} Hl(1 - a, -q + (\delta - 1)\gamma a + (\alpha - \delta + 1)(\beta - \delta + 1); \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, \gamma; 1 - x) \\
 = & (1 - x)^{1-\delta} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (2 - \delta)_{i_0}} \eta^{i_0} \right. \\
 & + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \alpha - \delta)(i_0 + 1 + \beta - \delta)}{(i_0 + 2)(i_0 + 3 - \delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (2 - \delta)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (4 - \delta)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (4 - \delta)_{i_1}} \eta^{i_1} \right\} z \\
 & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \alpha - \delta)(i_0 + 1 + \beta - \delta)}{(i_0 + 2)(i_0 + 3 - \delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (2 - \delta)_{i_0}} \right. \\
 & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + 1 + \alpha - \delta)(i_k + 2k + 1 + \beta - \delta)}{(i_k + 2(k + 1))(i_k + 2k + 3 - \delta)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + 2 - \delta)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + 2 - \delta)_{i_k}} \right\} \\
 & \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + 2 - \delta)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + 2 - \delta)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (1.8)
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(2-a)\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(2-a)k\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(2-a)n\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

$$\mathbf{.1.5} \quad x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1, \alpha - \beta + 1, \delta; \frac{1}{x} \right)$$

Polynomial of type 2

Replacing coefficients $a, q, \beta, \gamma, x, c_0$ and λ by $\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}, \alpha - \gamma + 1, \alpha - \beta + 1, \frac{1}{x}, 1$ and zero into (2.2.3). Multiply $x^{-\alpha}$ and the new (2.2.3) together.

$$\begin{aligned}
& x^{-\alpha} y(\xi) \\
= & x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1, \alpha - \beta + 1, \delta; \frac{1}{x} \right) \\
= & x^{-\alpha} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (1 + \alpha - \beta)_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + 1 + \alpha - \gamma)}{(i_0 + 2)(i_0 + 2 + \alpha - \beta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (1 + \alpha - \beta)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_0} (3 + \alpha - \beta)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (3 + \alpha - \beta)_{i_1}} \eta^{i_1} \Big\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + 1 + \alpha - \gamma)}{(i_0 + 2)(i_0 + 2 + \alpha - \beta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (1 + \alpha - \beta)_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha)(i_k + 2k + 1 + \alpha - \gamma)}{(i_k + 2(k+1))(i_k + 2k + 2 + \alpha - \beta)} \right. \\
& \times \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + 1 + \alpha - \beta)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + 1 + \alpha - \beta)_{i_k}} \Big\} \\
& \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + 1 + \alpha - \beta)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + 1 + \alpha - \beta)_{i_n}} \eta^{i_n} \right\} z^n \Big\} \quad (.1.9)
\end{aligned}$$

where

$$\begin{cases} \xi = \frac{1}{x} \\ z = -a\xi^2 \\ \eta = (1+a)\xi \\ \varphi = 2\alpha - \gamma - \delta + 1 + \frac{1}{a}(\alpha - \beta + \delta) \\ \Omega = \frac{a\varphi}{(1+a)} \\ q = -\alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta] \\ \quad - a(q_j + 2j)\{\varphi + (1 + 1/a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \beta, \gamma, x, c_0$ and λ by $\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}, \alpha-\gamma+1, \alpha-\beta+1, \frac{1}{x}, 1$ and zero into (2.2.7). Multiply $x^{-\alpha}$ and the new (2.2.7) together.

$$\begin{aligned} & x^{-\alpha}y(\xi) \\ &= x^{-\alpha}Hl\left(\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}; \alpha, \alpha-\gamma+1, \alpha-\beta+1, \delta; \frac{1}{x}\right) \\ &= x^{-\alpha}\left\{\sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(1+\alpha-\beta)_{i_0}} \eta^{i_0}\right. \\ &+ \left\{\sum_{i_0=0}^{\infty} \frac{(i_0+\alpha)(i_0+1+\alpha-\gamma)}{(i_0+2)(i_0+2+\alpha-\beta)} \frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(1+\alpha-\beta)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1}(\Delta_1^+)_{i_1}(3)_{i_0}(3+\alpha-\beta)_{i_0}}{(\Delta_1^-)_{i_0}(\Delta_1^+)_{i_0}(3)_{i_1}(3+\alpha-\beta)_{i_1}} \eta^{i_1}\right\} z \\ &+ \sum_{n=2}^{\infty} \left\{\sum_{i_0=0}^{\infty} \frac{(i_0+\alpha)(i_0+1+\alpha-\gamma)}{(i_0+2)(i_0+2+\alpha-\beta)} \frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(1+\alpha-\beta)_{i_0}}\right. \\ &\times \prod_{k=1}^{n-1} \left\{\sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+\alpha)(i_k+2k+1+\alpha-\gamma)}{(i_k+2(k+1))(i_k+2k+2+\alpha-\beta)} \frac{(\Delta_k^-)_{i_k}(\Delta_k^+)_{i_k}(2k+1)_{i_{k-1}}(2k+1+\alpha-\beta)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}}(\Delta_k^+)_{i_{k-1}}(2k+1)_{i_k}(2k+1+\alpha-\beta)_{i_k}}\right\} \\ &\times \left.\sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n}(\Delta_n^+)_{i_n}(2n+1)_{i_{n-1}}(2n+1+\alpha-\beta)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}}(\Delta_n^+)_{i_{n-1}}(2n+1)_{i_n}(2n+1+\alpha-\beta)_{i_n}} \eta^{i_n}\right\} z^n \Bigg\} \end{aligned} \quad (1.10)$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi \pm \sqrt{\varphi^2 - 4(1+1/a)q}}{2(1+1/a)} \\ \Delta_1^\pm = \frac{\{\varphi + 4(1+1/a)\} \pm \sqrt{\varphi^2 - 4(1+1/a)q}}{2(1+1/a)} \\ \Delta_k^\pm = \frac{\{\varphi + 4(1+1/a)k\} \pm \sqrt{\varphi^2 - 4(1+1/a)q}}{2(1+1/a)} \\ \Delta_n^\pm = \frac{\{\varphi + 4(1+1/a)n\} \pm \sqrt{\varphi^2 - 4(1+1/a)q}}{2(1+1/a)} \end{cases}$$

$$\mathbf{.1.6} \quad \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replacing coefficients a, q, α, x, c_0 and λ by $1 - a, -q + \gamma\beta, -\alpha + \gamma + \delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.2.3). Multiply $(1 - \frac{x}{a})^{-\beta}$ and the new (2.2.3) together.

$$\begin{aligned} & \left(1 - \frac{x}{a}\right)^{-\beta} y(\xi) \\ = & \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right) \\ = & \left(1 - \frac{x}{a}\right)^{-\beta} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \right. \\ & + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 - \alpha + \gamma + \delta)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0}} \frac{(3)_{i_0} (2 + \gamma)_{i_0}}{(3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\ & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 - \alpha + \gamma + \delta)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\ & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k - \alpha + \gamma + \delta)(i_k + 2k + \beta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}}} \frac{(2k+1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(2k+1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\ & \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}}} \frac{(2n+1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(2n+1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \end{aligned} \quad (.1.11)$$

where

$$\begin{cases} \xi = \frac{(1-a)x}{x-a} \\ z = -\frac{1}{1-a}\xi^2 \\ \eta = \frac{2-a}{1-a}\xi \\ \varphi = -\alpha + \beta + \gamma + (1-a)(\gamma + \delta - 1) \\ \Omega = \frac{a\varphi}{(2-a)} \\ q = \gamma\beta + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients a, q, α, x, c_0 and λ by $1-a, -q + \gamma\beta, -\alpha + \gamma + \delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.2.7). Multiply $(1 - \frac{x}{a})^{-\beta}$ and the new (2.2.7) together.

$$\begin{aligned} & \left(1 - \frac{x}{a}\right)^{-\beta} y(\xi) \\ &= \left(1 - \frac{x}{a}\right)^{-\beta} Hl\left(1-a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right) \\ &= \left(1 - \frac{x}{a}\right)^{-\beta} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \right. \\ &+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 - \alpha + \gamma + \delta)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\ &+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 - \alpha + \gamma + \delta)(i_0 + \beta)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\ &\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k - \alpha + \gamma + \delta)(i_k + 2k + \beta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\ &\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \end{aligned} \quad (.1.12)$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_1^\pm = \frac{\{\varphi + 4(2-a)\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_k^\pm = \frac{\{\varphi + 4(2-a)k\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_n^\pm = \frac{\{\varphi + 4(2-a)n\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \end{cases}$$

.1.7

$$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl \left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma, 2 - \delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replacing coefficients $a, q, \alpha, \beta, \delta, x, c_0$ and λ by $1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1], -\alpha + \gamma + 1, \beta - \delta + 1, 2 - \delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.2.3). Multiply

$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}$ and the new (2.2.3) together.

$$\begin{aligned}
 & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} y(\xi) \\
 = & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl \left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma \right. \\
 & \left. , 2 - \delta; \frac{(1-a)x}{x-a} \right) \\
 = & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \right. \\
 & + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 - \alpha + \gamma + 1)(i_0 + \beta - \delta + 1)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
 & \times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\
 & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 - \alpha + \gamma + 1)(i_0 + \beta - \delta + 1)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
 & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k - \alpha + \gamma + 1)(i_k + 2k + \beta - \delta + 1)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \right. \\
 & \times \left. \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\
 & \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (1.13)
 \end{aligned}$$

where

$$\begin{cases} \xi = \frac{(1-a)x}{x-a} \\ z = -\frac{1}{1-a} \xi^2 \\ \eta = \frac{(2-a)}{1-a} \xi \\ \varphi = -\alpha + \beta + \gamma + (1-a)(\gamma - \delta + 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = \gamma[(\delta-1)a + \beta - \delta + 1] + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \alpha, \beta, \delta, x, c_0$ and λ by $1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1], -\alpha + \gamma + 1, \beta - \delta + 1, 2 - \delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.2.7). Multiply $(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}$ and the new (2.2.7) together.

$$\begin{aligned}
& (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} y(\xi) \\
= & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl\left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma, 2 - \delta; \frac{(1-a)x}{x-a}\right) \\
= & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 - \alpha + \gamma + 1)(i_0 + \beta - \delta + 1)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} \eta^{i_1} \right\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 - \alpha + \gamma + 1)(i_0 + \beta - \delta + 1)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k - \alpha + \gamma + 1)(i_k + 2k + \beta - \delta + 1)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\
& \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1)_{i_n} (2n + \gamma)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.14)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(2-a)\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(2-a)k\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(2-a)n\} \pm \sqrt{\varphi^2 - 4(2-a)q}}{2(2-a)} \end{cases}$$

$$\mathbf{.1.8} \quad x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}; \alpha, \alpha-\gamma+1, \delta, \alpha-\beta+1; \frac{x-1}{x} \right)$$

Polynomial of type 2

Replacing coefficients $a, q, \beta, \gamma, \delta, x, c_0$ and λ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha-\gamma+1, \delta, \alpha-\beta+1, \frac{x-1}{x}, 1$ and zero into (2.2.3). Multiply $x^{-\alpha}$ and the new (2.2.3) together.

$$\begin{aligned} & x^{-\alpha} y(\xi) \\ = & x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}; \alpha, \alpha-\gamma+1, \delta, \alpha-\beta+1; \frac{x-1}{x} \right) \\ = & x^{-\alpha} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \eta^{i_0} \right. \\ & + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + \alpha - \gamma + 1)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_0} (2 + \delta)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (2 + \delta)_{i_1}} \eta^{i_1} \right\} z \\ & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 + \alpha - \gamma + 1)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \right. \\ & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha)(i_k + 2k + \alpha - \gamma + 1)}{(i_k + 2(k+1))(i_k + 2k + 1 + \delta)} \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + \delta)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + \delta)_{i_k}} \right\} \\ & \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + \delta)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + \delta)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \end{aligned} \quad (.1.15)$$

where

$$\begin{cases} \xi = \frac{x-1}{x} \\ z = \frac{-a}{a-1} \xi^2 \\ \eta = \frac{2a-1}{a-1} \xi \\ \varphi = \alpha + \beta - \gamma + \frac{a-1}{a} (\alpha - \beta + \delta) \\ \Omega = \frac{a\varphi}{(2a-1)} \\ q = \alpha(\delta a + \beta - \delta) + (q_j + 2j) \{a\varphi + (2a-1)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \beta, \gamma, \delta, x, c_0$ and λ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha-\gamma+1, \delta, \alpha-\beta+1, \frac{x-1}{x}, 1$ and zero into (2.2.7). Multiply $x^{-\alpha}$ and the new (2.2.7) together.

$$\begin{aligned}
& x^{-\alpha} y(\xi) \\
&= x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}; \alpha, \alpha-\gamma+1, \delta, \alpha-\beta+1; \frac{x-1}{x} \right) \\
&= x^{-\alpha} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\alpha)(i_0+\alpha-\gamma+1)}{(i_0+2)(i_0+1+\delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2+\delta)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2+\delta)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\alpha)(i_0+\alpha-\gamma+1)}{(i_0+2)(i_0+1+\delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+\alpha)(i_k+2k+\alpha-\gamma+1)}{(i_k+2(k+1))(i_k+2k+1+\delta)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1)_{i_{k-1}} (2k+\delta)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1)_{i_k} (2k+\delta)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1)_{i_{n-1}} (2n+\delta)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1)_{i_n} (2n+\delta)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.16)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{a\varphi \pm a\sqrt{\varphi^2 - 4\frac{(2a-1)}{a}q}}{2(2a-1)} \\ \Delta_1^{\pm} = \frac{a\left\{\varphi + 4\frac{(2a-1)}{a}\right\} \pm a\sqrt{\varphi^2 - 4\frac{(2a-1)}{a}q}}{2(2a-1)} \\ \Delta_k^{\pm} = \frac{a\left\{\varphi + 4\frac{(2a-1)}{a}k\right\} \pm a\sqrt{\varphi^2 - 4\frac{(2a-1)}{a}q}}{2(2a-1)} \\ \Delta_n^{\pm} = \frac{a\left\{\varphi + 4\frac{(2a-1)}{a}n\right\} \pm a\sqrt{\varphi^2 - 4\frac{(2a-1)}{a}q}}{2(2a-1)} \end{cases}$$

$$.1.9 \quad \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right)$$

Polynomial of type 2

Replacing coefficients $q, \beta, \gamma, \delta, x, c_0$ and λ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$ and zero into (2.2.3). Multiply $\left(\frac{x-a}{1-a} \right)^{-\alpha}$ and the new (2.2.3) together.

$$\begin{aligned} & \left(\frac{x-a}{1-a} \right)^{-\alpha} y(\xi) \\ = & \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right) \\ = & \left(\frac{x-a}{1-a} \right)^{-\alpha} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \eta^{i_0} \right. \\ & + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)(i_0 - \beta + \gamma + \delta)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(-q_0)_{i_0} (q_0 + \Omega)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \Omega)_{i_1} (3)_{i_0} (2 + \delta)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \Omega)_{i_0} (3)_{i_1} (2 + \delta)_{i_1}} \eta^{i_1} \right\} z \\ & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha)(i_k + 2k - \beta + \gamma + \delta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \delta)} \frac{(-q_k)_{i_k} (q_k + 4k + \Omega)_{i_k} (2k + 1)_{i_{k-1}} (2k + \delta)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \Omega)_{i_{k-1}} (2k + 1)_{i_k} (2k + \delta)_{i_k}} \right\} \\ & \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \Omega)_{i_n} (2n + 1)_{i_{n-1}} (2n + \delta)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \Omega)_{i_{n-1}} (2n + 1)_{i_n} (2n + \delta)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \end{aligned} \quad (.1.17)$$

where

$$\begin{cases} \xi = \frac{a(x-1)}{x-a} \\ z = -\frac{1}{a}\xi^2 \\ \eta = \frac{(1+a)}{a}\xi \\ \varphi = \alpha - \beta + \delta + a(\gamma + \delta - 1) \\ \Omega = \frac{\varphi}{(1+a)} \\ q = (\beta - \delta)\alpha - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $q, \beta, \gamma, \delta, x, c_0$ and λ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$ and zero into (2.2.7). Multiply $\left(\frac{x-a}{1-a}\right)^{-\alpha}$ and the new (2.2.7) together.

$$\begin{aligned}
& \left(\frac{x-a}{1-a}\right)^{-\alpha} y(\xi) \\
&= \left(\frac{x-a}{1-a}\right)^{-\alpha} Hl\left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a}\right) \\
&= \left(\frac{x-a}{1-a}\right)^{-\alpha} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \alpha)(i_0 - \beta + \gamma + \delta)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2 + \delta)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2 + \delta)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \alpha)(i_0 - \beta + \gamma + \delta)}{(i_0 + 2)(i_0 + 1 + \delta)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\delta)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \alpha)(i_k + 2k - \beta + \gamma + \delta)}{(i_k + 2(k+1))(i_k + 2k + 1 + \delta)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \delta)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \delta)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \delta)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \delta)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.18)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_1^{\pm} = \frac{\{\varphi + 4(1+a)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_k^{\pm} = \frac{\{\varphi + 4(1+a)k\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \\ \Delta_n^{\pm} = \frac{\{\varphi + 4(1+a)n\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} \end{cases}$$

.2 Integral formalism of 192 Heun functions

.2.1 $(1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x)$

Polynomial of type 2

Replacing coefficients $q, \alpha, \beta, \delta, c_0$ and λ by $q - (\delta-1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$ and zero into (2.3.6). Multiply $(1-x)^{1-\delta}$ and the new (2.3.6) together.

$$\begin{aligned}
& (1-x)^{1-\delta} y(x) \\
= & (1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x) \\
= & (1-x)^{1-\delta} \left\{ {}_2F_1(-q_0, q_0 + \Omega; \gamma; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
& \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
& \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\delta)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \Big\} \\
& \times {}_2F_1(-q_0, q_0 + \Omega; \gamma; \overleftrightarrow{w}_{1,n}) \Big\} z^n \Big\} \tag{.2.1}
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - \delta + a(\gamma - \delta + 1) \\ \Omega = \frac{\varphi}{(1+a)} \\ q = (\delta-1)\gamma a - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $q, \alpha, \beta, \delta, c_0$ and λ by $q - (\delta - 1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$ and zero into (2.3.18). Multiply $(1 - x)^{1-\delta}$ and the new (2.3.18) together.

$$\begin{aligned}
& (1 - x)^{1-\delta} y(x) \\
&= (1 - x)^{1-\delta} Hl(a, q - (\delta - 1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x) \\
&= (1 - x)^{1-\delta} \left\{ {}_2F_1(-\Pi_0^+, -\Pi_0^-; \gamma; \eta) \right. \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\delta)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \Big\} \\
&\times {}_2F_1(-\Pi_0^+, -\Pi_0^-; \gamma; \overleftrightarrow{w}_{1,n}) \Big\} z^n \Big\} \tag{.2.2}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)(q - (\delta - 1)\gamma a)}}{2(1+a)} \\ \Pi_{n-k}^\pm = \frac{-(\varphi + 4(1+a)(n-k)) \pm \sqrt{\varphi^2 - 4(1+a)(q - (\delta - 1)\gamma a)}}{2(1+a)} \end{cases}$$

.2.2

$$x^{1-\gamma} (1 - x)^{1-\delta} Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x)$$

Polynomial of type 2

Replacing coefficients $q, \alpha, \beta, \gamma, \delta, c_0$ and λ by

$$q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$$

and zero into (2.3.6). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and the new (2.3.6) together.

$$\begin{aligned}
& x^{1-\gamma}(1-x)^{1-\delta}y(x) \\
= & x^{1-\gamma}(1-x)^{1-\delta}Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2 \\
& , \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x) \\
= & x^{1-\gamma}(1-x)^{1-\delta} \left\{ {}_2F_1(-q_0, q_0 + \Omega; 2 - \gamma; \eta) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\gamma} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)+\alpha-\gamma-\delta)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)+\beta-\gamma-\delta} \left. \right\} \\
& \times {}_2F_1(-q_0, q_0 + \Omega; 2 - \gamma; \overleftarrow{w}_{1,n}) \left. \right\} z^n \quad (.2.3)
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - 2\gamma - \delta + 2 + a(3 - \gamma - \delta) \\ \Omega = \frac{\varphi}{(1+a)} \\ q = (\gamma + \delta - 2)a + (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1) \\ \quad - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $q, \alpha, \beta, \gamma, \delta, c_0$ and λ by

$$q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$$

and zero into (2.3.18). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and the new (2.3.18) together.

$$\begin{aligned}
& x^{1-\gamma}(1-x)^{1-\delta}y(x) \\
= & x^{1-\gamma}(1-x)^{1-\delta}Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2 \\
& , \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x) \\
= & x^{1-\gamma}(1-x)^{1-\delta} \left\{ {}_2F_1(-\Pi_0^+, -\Pi_0^-; 2 - \gamma; \eta) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\gamma} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)+\alpha-\gamma-\delta)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)+\beta-\gamma-\delta} \left. \right\} \\
& \times {}_2F_1(-\Pi_0^+, -\Pi_0^-; 2 - \gamma; \overleftarrow{w}_{1,n}) \left. \right\} z^n \Bigg\} \tag{.2.4}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)(q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1))}}{2(1+a)} \\ \Pi_{n-k}^\pm = \frac{-(\varphi + 4(1+a)(n-k)) \pm \sqrt{\varphi^2 - 4(1+a)(q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1))}}{2(1+a)} \end{cases}$$

$$\mathbf{.2.3} \quad Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x)$$

Polynomial of type 2

Replacing coefficients $a, q, \gamma, \delta, x, c_0$ and λ by $1-a, -q + \alpha\beta, \delta, \gamma, 1-x, 1$ and zero into (2.3.6).

$$\begin{aligned}
y(\xi) &= Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x) \\
&= {}_2F_1(-q_0, q_0 + \Omega; \delta; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta} \Big\} \\
&\times {}_2F_1(-q_0, q_0 + \Omega; \delta; \overleftrightarrow{w}_{1,n}) \Big\} z^n \tag{.2.5}
\end{aligned}$$

where

$$\begin{cases} \xi = 1-x \\ z = \frac{-1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = \alpha + \beta - \delta + (1-a)(\delta + \gamma - 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = \alpha\beta + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \gamma, \delta, x, c_0$ and λ by $1-a, -q+\alpha\beta, \delta, \gamma, 1-x, 1$ and zero into (2.3.18).

$$\begin{aligned}
y(\xi) &= Hl(1-a, -q+\alpha\beta; \alpha, \beta, \delta, \gamma; 1-x) \\
&= {}_2F_1(-\Pi_0^+, -\Pi_0^-; \delta; \eta) \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)+\beta} \Big\} \\
&\times {}_2F_1(-\Pi_0^+, -\Pi_0^-; \delta; \overleftarrow{w}_{1,n}) \Big\} z^n \tag{.2.6}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(2-a)(-q+\alpha\beta)}}{2(2-a)} \\ \Pi_{n-k}^\pm = \frac{-(\varphi + 4(2-a)(n-k)) \pm \sqrt{\varphi^2 - 4(2-a)(-q+\alpha\beta)}}{2(2-a)} \end{cases}$$

$$\textbf{.2.4} \quad (1-x)^{1-\delta} Hl(1-a, -q+(\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x)$$

Polynomial of type 2

Replacing coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0$ and λ by $1-a, -q+(\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma, 1-x, 1$ and zero

into (2.3.6). Multiply $(1-x)^{1-\delta}$ and the new (2.3.6) together.

$$\begin{aligned}
& (1-x)^{1-\delta} y(\xi) \\
&= (1-x)^{1-\delta} Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x) \\
&= (1-x)^{1-\delta} \left\{ {}_2F_1(-q_0, q_0 + \Omega; 2-\delta; \eta) \right. \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\delta} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\times (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\delta)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \Big\} \\
&\times {}_2F_1(-q_0, q_0 + \Omega; 2-\delta; \overleftrightarrow{w}_{1,n}) \Big\} z^n \Big\} \tag{.2.7}
\end{aligned}$$

where

$$\left\{ \begin{array}{l} \xi = 1-x \\ z = \frac{-1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = \alpha + \beta - \gamma - 2\delta + 2 + (1-a)(\gamma - \delta + 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1) \\ \quad + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{array} \right.$$

Infinite series

Replacing coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0$ and λ by $1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma, 1-x, 1$ and zero

into (2.3.18). Multiply $(1-x)^{1-\delta}$ and the new (2.3.18) together.

$$\begin{aligned}
& (1-x)^{1-\delta} y(\xi) \\
= & (1-x)^{1-\delta} Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, \\
& 2-\delta, \gamma; 1-x) \\
= & (1-x)^{1-\delta} \left\{ {}_2F_1(-\Pi_0^+, -\Pi_0^-; 2-\delta; \eta) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\delta} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
& \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\delta)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \Big\} \\
& \times \left. {}_2F_1(-\Pi_0^+, -\Pi_0^-; 2-\delta; \overleftrightarrow{w}_{1,n}) \right\} z^n \Big\} \tag{.2.8}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-(\varphi+4(2-a)(n-k)) \pm \sqrt{\varphi^2-4(2-a)(-q+(\delta-1)\gamma a+(\alpha-\delta+1)(\beta-\delta+1))}}{2(2-a)} \\ \Pi_{n-k}^\pm = \frac{-(\varphi+4(2-a)(n-k)) \pm \sqrt{\varphi^2-4(2-a)(-q+(\delta-1)\gamma a+(\alpha-\delta+1)(\beta-\delta+1))}}{2(2-a)} \end{cases}$$

$$\textbf{.2.5} \quad x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1, \alpha - \beta + 1, \delta; \frac{1}{x} \right)$$

Polynomial of type 2

Replacing coefficients $a, q, \beta, \gamma, x, c_0$ and λ by $\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}, \alpha - \gamma + 1, \alpha - \beta + 1, \frac{1}{x}, 1$ and zero into (2.3.6). Multiply $x^{-\alpha}$ and the new (2.3.6) together.

$$\begin{aligned}
& x^{-\alpha} y(\xi) \\
&= x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1, \alpha - \beta + 1, \delta; \frac{1}{x} \right) \\
&= x^{-\alpha} \left\{ {}_2F_1(-q_0, q_0 + \Omega; 1 + \alpha - \beta; \eta) \right. \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1+\alpha-\beta} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\times (1 - \overleftrightarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\gamma-1} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \Big\} \\
&\times {}_2F_1(-q_0, q_0 + \Omega; 1 + \alpha - \beta; \overleftrightarrow{w}_{1,n}) \Big\} z^n \Big\} \tag{.2.9}
\end{aligned}$$

where

$$\begin{cases} \xi = \frac{1}{x} \\ z = -a\xi^2 \\ \eta = (1+a)\xi \\ \varphi = 2\alpha - \gamma - \delta + 1 + \frac{1}{a}(\alpha - \beta + \delta) \\ \Omega = \frac{a\varphi}{(1+a)} \\ q = -\alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta] \\ \quad -a(q_j + 2j)\{\varphi + (1 + 1/a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \beta, \gamma, x, c_0$ and λ by $\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}, \alpha-\gamma+1, \alpha-\beta+1, \frac{1}{x}, 1$ and zero into (2.3.18). Multiply $x^{-\alpha}$ and the new (2.3.18) together.

$$\begin{aligned}
& x^{-\alpha} y(\xi) \\
= & x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}; \alpha, \alpha-\gamma+1, \alpha-\beta+1, \delta; \frac{1}{x} \right) \\
= & x^{-\alpha} \left\{ {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; 1+\alpha-\beta; \eta \right) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1+\alpha-\beta} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
& \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\gamma-1} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \left. \right\} \\
& \times {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; 1+\alpha-\beta; \overleftrightarrow{w}_{1,n} \right) \left. \right\} z^n \Bigg\} \tag{.2.10}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{-\varphi \pm \sqrt{\varphi^2 - 4\left(1 + \frac{1}{a}\right) \left(\frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a} \right)}}{2\left(1 + \frac{1}{a}\right)} \\ \Pi_{n-k}^{\pm} = \frac{-(\varphi + 4\left(1 + \frac{1}{a}\right)(n-k)) \pm \sqrt{\varphi^2 - 4\left(1 + \frac{1}{a}\right) \left(\frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a} \right)}}{2\left(1 + \frac{1}{a}\right)} \end{cases}$$

$$.2.6 \quad \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replacing coefficients a, q, α, x, c_0 and λ by $1 - a, -q + \gamma\beta, -\alpha + \gamma + \delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.3.6). Multiply $\left(1 - \frac{x}{a}\right)^{-\beta}$ and the new (2.3.6) together.

$$\begin{aligned}
& \left(1 - \frac{x}{a}\right)^{-\beta} y(\xi) \\
&= \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right) \\
&= \left(1 - \frac{x}{a}\right)^{-\beta} \left\{ {}_2F_1(-q_0, q_0 + \Omega; \gamma; \eta) \right. \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)-\alpha+\gamma+\delta)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{-\alpha-\beta+\gamma+\delta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta} \left. \right\} \\
&\times {}_2F_1(-q_0, q_0 + \Omega; \gamma; \overleftrightarrow{w}_{1,n}) \left. \right\} z^n \Bigg\} \tag{.2.11}
\end{aligned}$$

where

$$\begin{cases} \xi = \frac{(1-a)x}{x-a} \\ z = -\frac{1}{1-a}\xi^2 \\ \eta = \frac{2-a}{1-a}\xi \\ \varphi = -\alpha + \beta + \gamma + (1-a)(\gamma + \delta - 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = \gamma\beta + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients a, q, α, x, c_0 and λ by $1-a, -q+\gamma\beta, -\alpha+\gamma+\delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.3.18). Multiply $(1-\frac{x}{a})^{-\beta}$ and the new (2.3.18) together.

$$\begin{aligned}
& \left(1 - \frac{x}{a}\right)^{-\beta} y(\xi) \\
&= \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1-a, -q+\gamma\beta; -\alpha+\gamma+\delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right) \\
&= \left(1 - \frac{x}{a}\right)^{-\beta} \left\{ {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \gamma; \eta\right) \right. \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)-\alpha+\gamma+\delta)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{-\alpha-\beta+\gamma+\delta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta} \Big\} \\
&\times \left. {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \gamma; \overleftrightarrow{w}_{1,n}\right) \right\} z^n \Big\} \tag{2.12}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(2-a)(-q+\gamma\beta)}}{2(2-a)} \\ \Pi_{n-k}^{\pm} = \frac{-(\varphi+4(2-a)(n-k)) \pm \sqrt{\varphi^2 - 4(2-a)(-q+\gamma\beta)}}{2(2-a)} \end{cases}$$

.2.7

$$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl \left(1-a, -q+\gamma[(\delta-1)a+\beta-\delta+1]; -\alpha+\gamma+1, \beta-\delta+1, \gamma, 2-\delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replacing coefficients $a, q, \alpha, \beta, \delta, x, c_0$ and λ by $1-a, -q+\gamma[(\delta-1)a+\beta-\delta+1], -\alpha+\gamma+1, \beta-\delta+1, 2-\delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.3.6). Multiply

$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}$ and the new (2.3.6) together.

$$\begin{aligned}
& (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} y(\xi) \\
= & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl\left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma \right. \\
& \left. , 2 - \delta; (1-a)x(x-a)^{-1}\right) \\
= & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} \left\{ {}_2F_1(-q_0, q_0 + \Omega; \gamma; \eta) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1-\alpha+\gamma)} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{-\alpha-\beta+\gamma+\delta} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \Big\} \\
& \times {}_2F_1(-q_0, q_0 + \Omega; \gamma; \overleftarrow{w}_{1,n}) \Big\} z^n \Big\} \tag{.2.13}
\end{aligned}$$

where

$$\begin{cases} \xi = \frac{(1-a)x}{x-a} \\ z = -\frac{1}{1-a} \xi^2 \\ \eta = \frac{(2-a)}{1-a} \xi \\ \varphi = -\alpha + \beta + \gamma + (1-a)(\gamma - \delta + 1) \\ \Omega = \frac{\varphi}{(2-a)} \\ q = \gamma[(\delta-1)a + \beta - \delta + 1] + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \alpha, \beta, \delta, x, c_0$ and λ by $1-a, q^* = -q + \gamma[(\delta-1)a + \beta - \delta + 1], -\alpha + \gamma + 1, \beta - \delta + 1, 2 - \delta, \frac{(1-a)x}{x-a}, 1$ and zero into (2.3.18). Multiply

$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}$ and the new (2.3.18) together.

$$\begin{aligned}
& (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} y(\xi) \\
= & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl\left(1-a, q^* = -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1 \right. \\
& \left. , \beta - \delta + 1, \gamma, 2 - \delta; (1-a)x(x-a)^{-1}\right) \\
= & (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} \left\{ {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \gamma; \eta\right) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
& \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1-\alpha+\gamma)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{-\alpha-\beta+\gamma+\delta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \Big\} \\
& \times \left. {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \gamma; \overleftrightarrow{w}_{1,n}\right) \right\} z^n \Big\} \tag{.2.14}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(2-a)(-q + \gamma[(\delta-1)a + \beta - \delta + 1])}}{2(2-a)} \\ \Pi_{n-k}^\pm = \frac{-(\varphi + 4(2-a)(n-k)) \pm \sqrt{\varphi^2 - 4(2-a)(-q + \gamma[(\delta-1)a + \beta - \delta + 1])}}{2(2-a)} \end{cases}$$

$$\textbf{.2.8} \quad x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}; \alpha, \alpha - \gamma + 1, \delta, \alpha - \beta + 1; \frac{x-1}{x} \right)$$

Polynomial of type 2

Replacing coefficients $a, q, \beta, \gamma, \delta, x, c_0$ and λ by $\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}, \alpha - \gamma + 1, \delta, \alpha - \beta + 1, \frac{x-1}{x}, 1$ and zero into (2.3.6). Multiply $x^{-\alpha}$ and the new (2.3.6) together.

$$\begin{aligned}
& x^{-\alpha} y(\xi) \\
&= x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}; \alpha, \alpha - \gamma + 1, \delta, \alpha - \beta + 1; \frac{x-1}{x} \right) \\
&= x^{-\alpha} \left\{ {}_2F_1(-q_0, q_0 + \Omega; \delta; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \right. \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\gamma-1} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \Big\} \\
&\times {}_2F_1(-q_0, q_0 + \Omega; \delta; \overleftarrow{w}_{1,n}) \Big\} z^n \Big\} \tag{.2.15}
\end{aligned}$$

where

$$\begin{cases} \xi = \frac{x-1}{x} \\ z = \frac{-a}{a-1} \xi^2 \\ \eta = \frac{2a-1}{a-1} \xi \\ \varphi = \alpha + \beta - \gamma + \frac{a-1}{a}(\alpha - \beta + \delta) \\ \Omega = \frac{a\varphi}{(2a-1)} \\ q = \alpha(\delta a + \beta - \delta) + (q_j + 2j)\{a\varphi + (2a-1)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $a, q, \beta, \gamma, \delta, x, c_0$ and λ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha-\gamma+1, \delta, \alpha-\beta+1, \frac{x-1}{x}, 1$ and zero into (2.3.18). Multiply $x^{-\alpha}$ and the new (2.3.18) together.

$$\begin{aligned}
& x^{-\alpha} y(\xi) \\
= & x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}; \alpha, \alpha-\gamma+1, \delta, \alpha-\beta+1; \frac{x-1}{x} \right) \\
= & x^{-\alpha} \left\{ {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \delta; \eta \right) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\gamma-1} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \left. \right\} \\
& \times {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \delta; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \Bigg\} \tag{.2.16}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{-a\varphi \pm a\sqrt{\varphi^2-4\frac{(2a-1)}{a}\left(\frac{-q+\alpha(\delta a+\beta-\delta)}{a}\right)}}{2(2a-1)} \\ \Pi_{n-k}^{\pm} = \frac{-a\left\{\varphi+4\frac{(2a-1)}{a}(n-k)\right\} \pm a\sqrt{\varphi^2-4\frac{(2a-1)}{a}\left(\frac{-q+\alpha(\delta a+\beta-\delta)}{a}\right)}}{2(2a-1)} \end{cases}$$

$$\mathbf{.2.9} \quad \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right)$$

Polynomial of type 2

Replacing coefficients $q, \beta, \gamma, \delta, x, c_0$ and λ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$ and zero into (2.3.6). Multiply $\left(\frac{x-a}{1-a} \right)^{-\alpha}$ and the new (2.3.6) together.

$$\begin{aligned} & \left(\frac{x-a}{1-a} \right)^{-\alpha} y(\xi) \\ = & \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right) \\ = & \left(\frac{x-a}{1-a} \right)^{-\alpha} \left\{ {}_2F_1(-q_0, q_0 + \Omega; \delta; \eta) \right. \\ & + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \right. \right. \\ & \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\ & \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega)} \\ & \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\beta-\gamma-\delta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)-\beta+\gamma+\delta} \Big\} \\ & \times {}_2F_1(-q_0, q_0 + \Omega; \delta; \overleftrightarrow{w}_{1,n}) \Big\} z^n \Big\} \end{aligned} \quad (.2.17)$$

where

$$\begin{cases} \xi = \frac{a(x-1)}{x-a} \\ z = -\frac{1}{a}\xi^2 \\ \eta = \frac{(1+a)}{a}\xi \\ \varphi = \alpha - \beta + \delta + a(\gamma + \delta - 1) \\ \Omega = \frac{\varphi}{(1+a)} \\ q = (\beta - \delta)\alpha - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Infinite series

Replacing coefficients $q, \beta, \gamma, \delta, x, c_0$ and λ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$ and zero into (2.3.18). Multiply $\left(\frac{x-a}{1-a}\right)^{-\alpha}$ and the new (2.3.18) together.

$$\begin{aligned}
& \left(\frac{x-a}{1-a}\right)^{-\alpha} y(\xi) \\
&= \left(\frac{x-a}{1-a}\right)^{-\alpha} Hl\left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a}\right) \\
&= \left(\frac{x-a}{1-a}\right)^{-\alpha} \left\{ {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \delta; \eta\right) \right. \\
&+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}}\right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}}\right) \overleftarrow{w}_{n-k,n}^{\alpha+\beta-\gamma-\delta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}}\right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)-\beta+\gamma+\delta} \Big\} \\
&\times \left. {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \delta; \overleftarrow{w}_{1,n}\right) \right\} z^n \Big\} \tag{.2.18}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)(q - (\beta - \delta)\alpha)}}{2(1+a)} \\ \Pi_{n-k}^{\pm} = \frac{-(\varphi + 4(1+a)(n-k)) \pm \sqrt{\varphi^2 - 4(1+a)(q - (\beta - \delta)\alpha)}}{2(1+a)} \end{cases}$$

Bibliography

- [1] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806.
- [2] Choun, Y.S., “The analytic solution for the power series expansion of Heun function,” <http://dx.doi.org/10.1016/j.aop.2013.06.020> Available as arXiv:1303.0830.
- [3] Choun, Y.S., “Asymptotic behavior of Heun function and its integral formalism,” arXiv:1303.0876.
- [4] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische Annalen* 33, (1889)161.
- [5] Maier, R.S., “The 192 solutions of the Heun equation,” *Math. Comp.* 33, (2007)811–843.
- [6] Suzuki, H., Takasugi, E., Umetsu, H., “Perturbations of Kerr-de Sitter Black Hole and Heun’s Equation,” *Prog. Theor. Phys.* 100, (1998)491–505.
- [7] Suzuki, H., Takasugi, E., Umetsu, H., “Analytic solution of Teukolsky Equation in Kerr-de Sitter and Kerr-Newman-de Sitter Geometries,” *Prog. Theor. Phys.* 102, (1999)253–272.
- [8] Lau, S.R., “Rapid Evaluation of Radiation Boundary Kernels for Timedomain Wave Propagation on Blackholes,” *J. Comput. Phys.* 199, (2004)376–422. Available as arXiv:gr-qc/0401001.
- [9] Birkandan, T., Hortacsu, M., Examples of Heun and Mathieu functions as solutions of wave equations in curved spaces, *J. Phys. A: Math. Theor.* 40 (2007) 1105–1116.
- [10] Ronveaux, A., “Heuns Differential Equations” Oxford University Press, (1995).
- [11] M. Hortacsu, Heun Functions and their uses in Physics, arXiv:1101.0471

- [12] Epstein, P.S., The stark effect from the point of view of Schrödinger quantum theory, *Phys. Rev.* 2 (1926) 695.
- [13] Slavyanov, S.Y. and Lay, W., *Special Functions, A Unified Theory Based on Singularities*, (Oxford University Press, 2000).
- [14] Wilson, A.H., A generalized spheroidal wave equation. and The ionized hydrogen molecule, *Proc. Roy. soc. London A* 118 (1928) 617–647.
- [15] Teukolsky, S.A., Perturbations of a Rotating Black Hole I. Fundamental Equations for Gravitational, Electromagnetic, and Neutrino-Field Perturbations, *Astroph. J.* 185 (1973) 635.
- [16] Leaver, E.W., An Analytic Representation for the Quasi-Normal Modes of Kerr Black Holes, *Proc. Roy. soc. London A* 402 (1985) 285.
- [17] Batic, D., Schmid, H. and Winklmeier, M., The generalized Heun equation in QFT in curved space-times, *J. Phys. A* 116 (2006) 517.
- [18] Batic, D. and Schmid, H., Heun equation, Teukolsky equation, and type-D metrics, *J. Math. Phys.* 48 (2007) 042502.
- [19] Batic, D. and Sandoval, M., The hypergeneralized Heun equation in QFT in curved space-times, *Central Europ. J. Phys.* 8 (2010) 490.
- [20] Takemura, K., The Heun equation and the Calogero-Moser-Sutherland system I: the Bethe Ansatz method, *Commun. Math. Phys.* 235 (2003) 467–494.
- [21] Lambe, C.G., Ward, D.R., “Some differential equations and associated integral equations,” *Q. J. Math. (Oxford)* 5 (1934) 81–97.
- [22] Erdélyi, A., “Integral equations for Heun functions,” *Q. J. Math. (Oxford)* 13 (1942), 107–112.
- [23] Sleeman, B.D., “Integral representations for solutions of Heun’s equation,” *Proc. Camb. Phil. Soc.* 65 (1969) 447–459.
- [24] Sleeman, B.D., “Non-linear integral equations for Heun functions,” *Proc. Edinburgh Math. Sci.* (2) 16 (1969) 281–299.
- [25] Arscott, F.M., “Two-parameter eigenvalue problems in differential equations,” *Proc. Lond. Math. Soc.* 14 (1964) 459–470.
- [26] Schmidt, D., Wolf, G., “A method of generating integral relations by the simultaneous separability of generalized Schrödinger equations,” *SIAM J. Math. Anal.* 10 (1979) 823–838.

Chapter 3

The generating function for Heun polynomial using reversible three-term recurrence formula

In this chapter, by applying the generating function for Jacobi polynomial using hypergeometric functions into the integral representation of Heun polynomial of type 2¹, I consider the generating function of Heun polynomial including all higher terms of B_n 's².

Nine examples of 192 local solutions of the Heun equation (Maier, 2007) are provided in the appendices. For each example, I construct the generating function for Heun polynomial of type 2 including all higher terms of B_n 's.

3.1 Introduction

In 1837, Gabriel Lamé introduced second ordinary differential equation which has four regular singular points in the method of separation of variables applied to the Laplace equation in elliptic coordinates[6]. Various authors has called this equation as 'Lamé equation' or 'ellipsoidal harmonic equation'[23].

In 1889, K. Heun worked on the second ordinary differential equation which has four regular singular points. Later, the Heun function became a general function of all

¹polynomial of type 2 is a polynomial which makes A_n term terminated in three term recursion relation of the power series in a linear differential equation.

²"higher terms of B_n 's" means at least two terms of B_n 's.

well-known special functions: Mathieu, Lamé and Coulomb spheroidal functions.

Lamé ordinary differential equation represented either in the algebraic form or in Weierstrass's form is the special case of Heun's differential equation: Lamé equation is derived from Heun equation by changing all coefficients and an independent variable. The solutions of Lamé and Heun equations are a form of power series that can be expressed as three term recurrence. In contrast, most of well-known special functions consist of two term recursion relation (Hypergeometric, Bessel, Legendre, Kummer functions, etc). Lamé and Heun functions arise from deriving the Laplace equation in general Jacobi ellipsoidal or conical coordinates.[8] Also Heun polynomial arises in the separable coordinate systems on the n-sphere.[7]

According to E. G. Kalnins and W. Miller Jr.(1990 [8]) , “Lamé [6] and Heun functions have received relatively little attention, since they are rather intractable. Unfortunately the beautiful identities appearing have received little notice, probably because the methods of proof seemed obscure.”

Because Lamé and Heun equations have three term recursion relation in power series representation: three different coefficients in its power series create the complicated mathematical calculations. In general, most problems of physical and mechanical system in nature are nonlinear. For the simplified purposes and better approximation of physical systems, mathematically we have used linear (ordinary) differential equations to linearize these nonlinear system. Traditionally, we have explained these linear systems by only using two term recurrence relation of the power series in linear ordinary differential equations until 19th century. However, since modern physics (quantum gravity, SUSY, general relativity, etc) comes out the world, it seems to require at least three or four term recurrence relations in power series expansions. Furthermore these type of problems can not be reduced to two term recurrence relations.[18]

In 1939 Svartholm showed how Heun equation can be obtained in the form of series of degenerate hypergeometric functions. [11] A few years later, His solution was developed by Erdélyi in three papers (1942, 1944), Sleeman (1969), Schmidt and Wolf (1979), Kalnins and Miller (1991).[7, 12, 13, 14, 15, 16, 18]

In Ref.[3, 4], by applying 3TRF, I show power series expansion in closed forms of Heun equation and its representation of the form of integrals for infinite series and polynomial of type 1.³ I show that a ${}_2F_1$ function recurs in each of sub-integral forms of Heun function: the first sub-integral form contains zero term of A'_n 's, the second one contains one term of A_n 's, the third one contains two terms of A_n 's, etc.

³polynomial of type 1 is a polynomial which makes B_n term terminated in three term recursion relation of the power series in a linear differential equation.

In chapter 2, by applying R3TRF, I show power series expansion in closed forms of Heun function and its representation of the form of integrals for infinite series and polynomial of type 2. I also show that a ${}_2F_1$ function recurs in each of sub-integral forms of Heun function: the first sub-integral form contains zero term of B'_n 's, the second one contains one term of B_n 's, the third one contains two terms of B_n 's, etc.

In mathematics, a generating function $F(x, t)$ is a formal (it need not converge) power series in one indeterminate t^n , whose coefficients encode information about a sequence of numbers $f_n(x)$ that is indexed by the natural numbers (in quantum mechanically, the index n corresponds to the eigenvalues denoted by natural numbers). In the case of Legendre equation, $f_n(x)$ corresponds to the Legendre polynomials. Since the expansion of $F(x, t)$ has generated the set $f_n(x)$ in t , $F(x, t)$ can be described as an analytic function, combined coefficients x and t , in closed form without index n . In general, a generating function in closed form means that it is expressed in a direct form without summation signs. For instance, the generating function for Legendre polynomials is given by

$$\sum_{n=0}^{\infty} P_n(x) t^n = \frac{1}{\sqrt{1-2xt+t^2}} \quad (3.1.1)$$

$$= 1 + xt + \frac{1}{2}(3x^2 - 1)t^2 + \frac{1}{2}(5x^3 - 3x)t^3 + \frac{1}{8}(35x^4 - 30x^2 + 3)t^4 \\ + \frac{1}{8}(63x^5 - 70x^3 + 15x)t^5 + \frac{1}{16}(231x^6 - 315x^4 + 105x^2 - 5)t^6 \dots \quad (3.1.2)$$

The power series representation (3.1.2) is not in closed form while an analytic function (3.1.1) is.

In Ref.[5], by apply a generating function for Jacobi polynomial using hypergeometric functions into the general representation in the form of integral of Lamé polynomial of type 1 in Weierstrass's form, I construct the generating function for Lamé polynomial analytically. As we see generating functions for Lamé polynomial of type 1 of the first and second kinds, it is described as an analytic function in closed forms with removed index α_j where $j, \alpha_j \in \mathbb{N}_0$. α_j is the natural numbers (eigenvalues) which makes B_n term terminated at specific values of index n in sequence c_n : the sequence c_n consists of combinations A_n and B_n terms in three term recurrence relation of the power series representation of Lamé equation.

In this chapter, by using same technique for the generating function for Lamé polynomial of type 1, I consider the generating function for Heun polynomial of type 2 including all higher terms of B_n 's by applying a generating function for Jacobi polynomial using hypergeometric functions into the general integral form of Heun polynomial of type 2. Since the generating function for Heun polynomial is derived, we might be possible to

construct orthogonal relations of Heun polynomial. In the physical point of view we might be possible to obtain the normalized constant for the wave function in modern physics, recursion relation and its expectation value of any physical quantities from the generating function for Heun polynomial. For the case of hydrogen-like atoms, the normalized wave function is derived from the generating function for associated Laguerre polynomials. And the expectation value of physical quantities such as position and momentum is constructed by applying the recursive relation of associated Laguerre polynomials.

In future papers I will derive Heun polynomial of type 3 and its generating function that has fixed integer values of α and/or β , just as it has a fixed value of q . In this paper I construct the generating function for Heun polynomial of type 2: I treat α , β , γ and δ as free variables and the accessory parameter q as a fixed value.

Heun's equation is a second-order linear ordinary differential equation of the form [1].

$$\frac{d^2 y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0 \quad (3.1.3)$$

With the condition $\epsilon = \alpha + \beta - \gamma - \delta + 1$. The parameters play different roles: $a \neq 0$ is the singularity parameter, α , β , γ , δ , ϵ are exponent parameters, q is the accessory parameter which in many physical applications appears as a spectral parameter. Also, α and β are identical to each other. The total number of free parameters is six. It has four regular singular points which are 0, 1, a and ∞ with exponents $\{0, 1 - \gamma\}$, $\{0, 1 - \delta\}$, $\{0, 1 - \epsilon\}$ and $\{\alpha, \beta\}$.

3.2 Generating function for Heun polynomial of type 2

Let's investigate the generating function for Heun polynomial of type 2 of the first and second kinds.

Lemma 3.2.1 *The generating function for Jacobi polynomial using hypergeometric functions is defined by*

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} w^{q_0} {}_2F_1(-q_0, q_0 + A; \gamma; x) \\ &= 2^{A-1} \frac{\left(1 - w + \sqrt{w^2 - 2(1-2x)w + 1}\right)^{1-\gamma} \left(1 + w + \sqrt{w^2 - 2(1-2x)w + 1}\right)^{\gamma-A}}{\sqrt{w^2 - 2(1-2x)w + 1}} \\ & \text{where } |w| < 1 \end{aligned} \quad (3.2.1)$$

Proof Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ can be written in terms of hypergeometric function using

$${}_2F_1(-n, n + \alpha + \beta + 1; \alpha + 1; x) = \frac{n!}{(\alpha + 1)_n} P_n^{(\alpha, \beta)}(1 - 2x) \quad (3.2.2)$$

And

$$P_n^{(\alpha, \beta)}(x) = \frac{\Gamma(n + \alpha + 1)}{n! \Gamma(n + \alpha + \beta + 1)} \sum_{m=0}^n \binom{n}{m} \frac{\Gamma(n + m + \alpha + \beta + 1)}{\Gamma(m + \alpha + 1)} \left(\frac{x - 1}{2} \right)^m \quad (3.2.3)$$

The generating function for Jacobi polynomials is given by

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x) w^n = 2^{\alpha+\beta} \frac{\left(1 - w + \sqrt{w^2 - 2xw + 1}\right)^{-\alpha} \left(1 + w + \sqrt{w^2 - 2xw + 1}\right)^{-\beta}}{\sqrt{w^2 - 2xw + 1}} \quad (3.2.4)$$

Replace n , α and β by q_0 , $\gamma - 1$ and $A - \gamma$ in (3.2.2), and acting the summation operator

$$\sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} w^{q_0} \text{ on the new (3.2.2)}$$

$$\sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} w^{q_0} {}_2F_1(-q_0, q_0 + A; \gamma; x) = \sum_{q_0=0}^{\infty} P_n^{(\gamma-1, A-\gamma)}(1 - 2x) w^{q_0} \quad (3.2.5)$$

Replace α , β and x by $\gamma - 1$, $A - \gamma$ and $1 - 2x$ in (3.2.4). As we take the new (3.2.4) into (3.2.5), we obtain (3.2.1). ■

Definition 3.2.2 I define that

$$\left\{ \begin{array}{l} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{if } a > b \\ s_a & \text{if } a = b \end{cases} \\ \tilde{w}_{i,j} = \begin{cases} \frac{\tilde{w}_{i+1,j} t_i u_i \{1 + (s_i + 2\tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))s_i\}}{2(1 - \tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))^2 s_i} \\ - \frac{\tilde{w}_{i+1,j} t_i u_i (1 + s_i) \sqrt{s_i^2 - 2(1 - 2\tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))s_i + 1}}{2(1 - \tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))^2 s_i} & \text{where } i < j \\ \frac{\eta t_i u_i \{1 + (s_{i,\infty} + 2\eta(1 - t_i)(1 - u_i))s_{i,\infty}\}}{2(1 - \eta(1 - t_i)(1 - u_i))^2 s_{i,\infty}} \\ - \frac{\eta t_i u_i (1 + s_{i,\infty}) \sqrt{s_{i,\infty}^2 - 2(1 - 2\eta(1 - t_i)(1 - u_i))s_{i,\infty} + 1}}{2(1 - \eta(1 - t_i)(1 - u_i))^2 s_{i,\infty}} & \text{where } i = j \end{cases} \end{array} \right. \quad (3.2.6)$$

where

$$a, b, i, j \in \mathbb{N}_0$$

And we have

$$\sum_{q_i=q_j}^{\infty} r_i^{q_i} = \frac{r_i^{q_j}}{(1-r_i)} \quad (3.2.7)$$

Theorem 3.2.3 *In chapter 2, the representation in the form of integral of Heun polynomial of type 2 is given by⁴*

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\ &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\ &\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \\ &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\ &\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\left(4(n-k) + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\ &\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)+\beta+\lambda} \left. \right\} \\ &\quad \times \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,n}^{i_0} \left. \right\} z^n \Bigg\} \end{aligned} \quad (3.2.8)$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1) \\ q = -(q_j + 2j + \lambda)\{\varphi + (1+a)(q_j + 2j + \lambda)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

⁴ $y_1(x)$ means the sub-integral form in (3.2.8) contains one term of $B'_n s$, $y_2(x)$ means the sub-integral form in (3.2.8) contains two terms of $B'_n s$, $y_3(x)$ means the sub-integral form in (3.2.8) contains three terms of $B'_n s$, etc.

and

$$\overleftrightarrow{w}_{i,j} = \begin{cases} \frac{v_i}{(v_i - 1)} \frac{\overleftrightarrow{w}_{i+1,j} t_i u_i}{1 - \overleftrightarrow{w}_{i+1,j} v_i (1 - t_i)(1 - u_i)} & \text{where } i \leq j \\ \eta & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of $B'_n s$, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (3.2.8) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (3.2.6) and (3.2.7),

Theorem 3.2.4 *The general expression of the generating function for Heun polynomial of type 2 is given by*

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y(x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \Upsilon(\lambda; s_{0,\infty}; \eta) \\ &+ \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_{1,\infty}^2 - 2(1 - 2\eta(1 - t_1)(1 - u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\ &\times \left(\frac{(1 + s_{1,\infty}) + \sqrt{s_{1,\infty}^2 - 2(1 - 2\eta(1 - t_1)(1 - u_1))s_{1,\infty} + 1}}{2} \right)^{-\left(3 + \frac{\varphi+2(1+\alpha)\lambda}{(1+\alpha)}\right)} \\ &\times \tilde{w}_{1,1}^{-(\alpha+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta+\lambda} \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) z \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1+\lambda} \int_0^1 du_n u_n^{2(n-1)+\gamma+\lambda} (s_{n,\infty}^2 - 2(1 - 2\eta(1 - t_n)(1 - u_n))s_{n,\infty} + 1)^{-\frac{1}{2}} \right. \\ &\times \left(\frac{(1 + s_{n,\infty}) + \sqrt{s_{n,\infty}^2 - 2(1 - 2\eta(1 - t_n)(1 - u_n))s_{n,\infty} + 1}}{2} \right)^{-\left(4n-1 + \frac{\varphi+2(1+\alpha)\lambda}{(1+\alpha)}\right)} \\ &\times \tilde{w}_{n,n}^{-(2(n-1)+\alpha+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\beta+\lambda} \\ &\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} (s_{n-k}^2 - 2(1 - 2\tilde{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k}))s_{n-k} + 1)^{-\frac{1}{2}} \right. \\ &\times \left(\frac{(1 + s_{n-k}) + \sqrt{s_{n-k}^2 - 2(1 - 2\tilde{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k}))s_{n-k} + 1}}{2} \right)^{-\left(4(n-k)-1 + \frac{\varphi+2(1+\alpha)\lambda}{(1+\alpha)}\right)} \\ &\times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\beta+\lambda} \left. \right\} \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) \Big\} z^n \end{aligned} \quad (3.2.9)$$

where

$$\left\{ \begin{array}{l} \Upsilon(\lambda; s_{0,\infty}; \eta) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_{0,\infty}^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right) \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \tilde{w}_{1,n}^{i_0} \right) \end{array} \right.$$

Proof Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on the integral form of Heun polynomial of type 2 $y(x)$,

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y(x) \\ &= \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} \{y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots\} \end{aligned} \quad (3.2.10)$$

According to (3.2.8), integral forms of sub-summation $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ are

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \quad (3.2.11a)$$

$$\begin{aligned} y_1(x) &= \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \frac{1}{1-\eta(1-t_1)(1-u_1)v_1} \right)^{q_1} \\ &\times (1-\eta(1-t_1)(1-u_1)v_1)^{-\left(4+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\ &\times \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\beta+\lambda} \\ &\times \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right) z \end{aligned} \quad (3.2.11b)$$

$$\begin{aligned}
y_2(x) = & \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \frac{1}{1-\eta(1-t_2)(1-u_2)v_2} \right)^{q_2} \\
& \times (1-\eta(1-t_2)(1-u_2)v_2)^{-\left(8+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{\alpha-\beta} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{2+\beta+\lambda} \\
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \frac{1}{1-\overleftarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
& \times (1-\overleftarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\left(4+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,1} \partial_{\overleftarrow{w}_{1,1}} \right) \overleftarrow{w}_{1,1}^{\alpha-\beta} \left(\overleftarrow{w}_{1,1} \partial_{\overleftarrow{w}_{1,1}} \right) \overleftarrow{w}_{1,1}^{\beta+\lambda} \\
& \times \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,2}^{i_0} \right) z^2 \tag{3.2.11c}
\end{aligned}$$

$$\begin{aligned}
y_3(x) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3-1}{v_3} \frac{1}{1-\eta(1-t_3)(1-u_3)v_3} \right)^{q_3} \\
& \times (1-\eta(1-t_3)(1-u_3)v_3)^{-\left(12+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{3,3}^{-(4+\alpha+\lambda)} \left(\overleftarrow{w}_{3,3} \partial_{\overleftarrow{w}_{3,3}} \right) \overleftarrow{w}_{3,3}^{\alpha-\beta} \left(\overleftarrow{w}_{3,3} \partial_{\overleftarrow{w}_{3,3}} \right) \overleftarrow{w}_{3,3}^{4+\beta+\lambda} \\
& \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \frac{1}{1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \right)^{q_2} \\
& \times (1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2)^{-\left(8+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{2,3}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,3} \partial_{\overleftarrow{w}_{2,3}} \right) \overleftarrow{w}_{2,3}^{\alpha-\beta} \left(\overleftarrow{w}_{2,3} \partial_{\overleftarrow{w}_{2,3}} \right) \overleftarrow{w}_{2,3}^{2+\beta+\lambda} \\
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \frac{1}{1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
& \times (1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1)^{-\left(4+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
& \times \overleftarrow{w}_{1,3}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,3} \partial_{\overleftarrow{w}_{1,3}} \right) \overleftarrow{w}_{1,3}^{\alpha-\beta} \left(\overleftarrow{w}_{1,3} \partial_{\overleftarrow{w}_{1,3}} \right) \overleftarrow{w}_{1,3}^{\beta+\lambda} \\
& \times \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,3}^{i_0} \right) z^3 \tag{3.2.11d}
\end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (3.2.11a),

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_0(x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_{0,\infty}^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right) \end{aligned} \quad (3.2.12)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (3.2.11b),

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\eta(1-t_1)(1-u_1)v_1)^{-\left(4+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\ & \times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_{1,\infty}}{1-\eta(1-t_1)(1-u_1)v_1} \right)^{q_1} \overleftarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,1} \partial \overleftarrow{w}_{1,1} \right) \overleftarrow{w}_{1,1}^{\alpha-\beta} \left(\overleftarrow{w}_{1,1} \partial \overleftarrow{w}_{1,1} \right) \overleftarrow{w}_{1,1}^{\beta+\lambda} \\ & \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,1}^{i_0} \right) z \end{aligned} \quad (3.2.13)$$

Replace q_i , q_j and r_i by q_1 , q_0 and $\frac{v_1-1}{v_1} \frac{s_{1,\infty}}{1-\eta(1-t_1)(1-u_1)v_1}$ in (3.2.7). Take the new (3.2.7) into (3.2.13).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{(1-\eta(1-t_1)(1-u_1)v_1)^{-\left(3+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)}}{-\eta(1-t_1)(1-u_1)v_1^2 + (1-s_{1,\infty})v_1 + s_{1,\infty}} \\ & \times \overleftarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,1} \partial \overleftarrow{w}_{1,1} \right) \overleftarrow{w}_{1,1}^{\alpha-\beta} \left(\overleftarrow{w}_{1,1} \partial \overleftarrow{w}_{1,1} \right) \overleftarrow{w}_{1,1}^{\beta+\lambda} \\ & \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} \left(\frac{v_1-1}{v_1} \frac{s_{0,\infty}}{1-\eta(1-t_1)(1-u_1)v_1} \right)^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,1}^{i_0} \right) z \end{aligned} \quad (3.2.14)$$

By using Cauchy's integral formula, the contour integrand has poles at

$$v_1 = \frac{1 - s_{1,\infty} - \sqrt{(1 - s_{1,\infty})^2 + 4\eta(1 - t_1)(1 - u_1)s_{1,\infty}}}{2\eta(1 - t_1)(1 - u_1)} \text{ or } \frac{1 - s_{1,\infty} + \sqrt{(1 - s_{1,\infty})^2 + 4\eta(1 - t_1)(1 - u_1)s_{1,\infty}}}{2\eta(1 - t_1)(1 - u_1)} \text{ and}$$

$$\frac{1 - s_{1,\infty} - \sqrt{(1 - s_{1,\infty})^2 + 4\eta(1 - t_1)(1 - u_1)s_{1,\infty}}}{2\eta(1 - t_1)(1 - u_1)}$$

is only inside the unit circle. As we compute the residue there in (3.2.14) we obtain

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_{1,\infty}^2 - 2(1 - 2\eta(1 - t_1)(1 - u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\ &\quad \times \left(\frac{1 + s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1 - 2\eta(1 - t_1)(1 - u_1))s_{1,\infty} + 1}}{2} \right)^{-\left(3 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\ &\quad \times \tilde{w}_{1,1}^{-(\alpha+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta+\lambda} \\ &\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) z \end{aligned} \quad (3.2.15)$$

where

$$\begin{aligned} \tilde{w}_{1,1} &= \frac{v_1}{(v_1 - 1)} \frac{\eta t_1 u_1}{1 - \eta v_1 (1 - t_1)(1 - u_1)} \Bigg|_{v_1 = \frac{1 - s_{1,\infty} - \sqrt{(1 - s_{1,\infty})^2 + 4\eta(1 - t_1)(1 - u_1)s_{1,\infty}}}{2\eta(1 - t_1)(1 - u_1)}} \\ &= \frac{\eta t_1 u_1 \{1 + (s_{1,\infty} + 2\eta(1 - t_1)(1 - u_1))s_{1,\infty}\}}{2(1 - \eta(1 - t_1)(1 - u_1))^2 s_{1,\infty}} \\ &\quad - \frac{\eta t_1 u_1 (1 + s_{1,\infty}) \sqrt{s_{1,\infty}^2 - 2(1 - 2\eta(1 - t_1)(1 - u_1))s_{1,\infty} + 1}}{2(1 - \eta(1 - t_1)(1 - u_1))^2 s_{1,\infty}} \end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (3.2.11c),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} (1-\eta(1-t_2)(1-u_2)v_2)^{-\left(8+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\times \sum_{q_2=q_1}^{\infty} \left(\frac{v_2-1}{v_2} \frac{s_{2,\infty}}{1-\eta(1-t_2)(1-u_2)v_2} \right)^{q_2} \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha-\beta} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2+\beta+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\left(4+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_1}{1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \overleftrightarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\alpha-\beta} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\beta+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right) z^2 \tag{3.2.16}
\end{aligned}$$

Replace q_i , q_j and r_i by q_2 , q_1 and $\frac{v_2-1}{v_2} \frac{s_{2,\infty}}{1-\eta(1-t_2)(1-u_2)v_2}$ in (3.2.7). Take the new

(3.2.7) into (3.2.16).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{(1-\eta(1-t_2)(1-u_2)v_2)^{-\left(7+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)}}{-\eta(1-t_2)(1-u_2)v_2^2 + (1-s_{2,\infty})v_2 + s_{2,\infty}} \\
&\times \overleftarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overrightarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{\alpha-\beta} \left(\overrightarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{2+\beta+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\overleftarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\left(4+\frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_2-1}{v_2} \frac{s_{1,\infty}}{1-\eta(1-t_2)(1-u_2)v_2} \frac{v_1-1}{v_1} \frac{1}{1-\overleftarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
&\times \overleftarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overrightarrow{w}_{1,2} \partial_{\overleftarrow{w}_{1,2}} \right) \overleftarrow{w}_{1,2}^{\alpha-\beta} \left(\overrightarrow{w}_{1,2} \partial_{\overleftarrow{w}_{1,2}} \right) \overleftarrow{w}_{1,2}^{\beta+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftarrow{w}_{1,2}^{i_0} \right) z^2 \tag{3.2.17}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$v_2 = \frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4\eta(1-t_2)(1-u_2)s_{2,\infty}}}{2\eta(1-t_2)(1-u_2)} \quad \text{or} \quad \frac{1-s_{2,\infty} + \sqrt{(1-s_{2,\infty})^2 + 4\eta(1-t_2)(1-u_2)s_{2,\infty}}}{2\eta(1-t_2)(1-u_2)}$$

and $\frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4\eta(1-t_2)(1-u_2)s_{2,\infty}}}{2\eta(1-t_2)(1-u_2)}$ is only inside the unit circle. As

we compute the residue there in (3.2.17) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-\left(7 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{2,2}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\alpha-\beta} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2+\beta+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\left(4 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \ddot{w}_{1,2}^{-(\alpha+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\alpha-\beta} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\beta+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) z^2 \tag{3.2.18}
\end{aligned}$$

where

$$\begin{aligned}
\tilde{w}_{2,2} &= \frac{v_2}{(v_2-1)} \frac{\eta t_2 u_2}{1 - \eta v_2(1-t_2)(1-u_2)} \Bigg|_{v_2 = \frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4\eta(1-t_2)(1-u_2)s_{2,\infty}}}{2\eta(1-t_2)(1-u_2)}} \\
&= \frac{\eta t_2 u_2 \{1 + (s_{2,\infty} + 2\eta(1-t_2)(1-u_2))s_{2,\infty}\}}{2(1-\eta(1-t_2)(1-u_2))^2 s_{2,\infty}} \\
&\quad - \frac{\eta t_2 u_2 (1 + s_{2,\infty}) \sqrt{s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2(1-\eta(1-t_2)(1-u_2))^2 s_{2,\infty}}
\end{aligned}$$

and

$$\ddot{w}_{1,2} = \frac{v_1}{(v_1-1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2} v_1(1-t_1)(1-u_1)}$$

Replace q_i , q_j and r_i by q_1 , q_0 and $\frac{v_1-1}{v_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1}$ in (3.2.7). Take the

new (3.2.7) into (3.2.18).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-\left(7 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{2,2}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\alpha-\beta} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2+\beta+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\left(3 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)}}{-\tilde{w}_{2,2}(1-t_1)(1-u_1)v_1^2 + (1-s_1)v_1 + s_1} \\
&\quad \times \ddot{w}_{1,2}^{-(\alpha+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\alpha-\beta} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\beta+\lambda} \tag{3.2.19} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} \left(\frac{v_1 - 1}{v_1} \frac{s_{0,1}}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) z^2
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$v_1 = \frac{1 - s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)} \quad \text{or} \quad \frac{1 - s_1 + \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)}$$

and $\frac{1 - s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)}$ is only inside the unit circle. As we

compute the residue there in (3.2.19) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2\eta(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-\left(7 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{2,2}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\alpha-\beta} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2+\beta+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-\left(3 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{1,2}^{-(\alpha+\lambda)} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\alpha-\beta} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\beta+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \tilde{w}_{1,2}^{i_0} \right) z^2
\end{aligned} \tag{3.2.20}$$

where

$$\begin{aligned}
\tilde{w}_{1,2} &= \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2} v_1 (1 - t_1)(1 - u_1)} \Bigg|_{v_1 = \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)}} \\
&= \frac{\tilde{w}_{2,2} t_1 u_1 \{1 + (s_1 + 2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1\}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1} \\
&\quad - \frac{\tilde{w}_{2,2} t_1 u_1 (1 + s_1) \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1}
\end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (3.2.11d),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_3(x) \\
&= \prod_{k=4}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} (s_{3,\infty}^2 - 2(1-2\eta(1-t_3)(1-u_3))s_{3,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{3,\infty} + \sqrt{s_{3,\infty}^2 - 2(1-2\eta(1-t_3)(1-u_3))s_{3,\infty} + 1}}{2} \right)^{-\left(11 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{3,3}^{-(4+\alpha+\lambda)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{\alpha-\beta} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{4+\beta+\lambda} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_2 + \sqrt{s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1}}{2} \right)^{-\left(7 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{2,3}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{\alpha-\beta} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{2+\beta+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-\left(3 + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
&\quad \times \tilde{w}_{1,3}^{-(\alpha+\lambda)} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\alpha-\beta} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\beta+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \tilde{w}_{1,3}^{i_0} \right) z^3
\end{aligned} \tag{3.2.21}$$

where

$$\begin{aligned}
\tilde{w}_{3,3} &= \frac{v_3}{(v_3 - 1)} \frac{\eta t_3 u_3}{1 - \eta(1 - t_3)(1 - u_3)v_3} \Bigg|_{v_3 = \frac{1 - s_{3,\infty} - \sqrt{(1 - s_{3,\infty})^2 + 4\eta(1 - t_3)(1 - u_3)s_{3,\infty}}}{2\eta(1 - t_3)(1 - u_3)}} \\
&= \frac{\eta t_3 u_3 \{1 + (s_{3,\infty} + 2\eta(1 - t_3)(1 - u_3))s_{3,\infty}\}}{2(1 - \eta(1 - t_3)(1 - u_3))^2 s_{3,\infty}} \\
&\quad - \frac{\eta t_3 u_3 (1 + s_{3,\infty}) \sqrt{s_{3,\infty}^2 - 2(1 - 2\eta(1 - t_3)(1 - u_3))s_{3,\infty} + 1}}{2(1 - \eta(1 - t_3)(1 - u_3))^2 s_{3,\infty}} \\
\\
\tilde{w}_{2,3} &= \frac{v_2}{(v_2 - 1)} \frac{\tilde{w}_{3,3} t_2 u_2}{1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2)v_2} \Bigg|_{v_2 = \frac{1 - s_2 - \sqrt{(1 - s_2)^2 + 4\tilde{w}_{3,3}(1 - t_2)(1 - u_2)s_2}}{2\tilde{w}_{3,3}(1 - t_2)(1 - u_2)}} \\
&= \frac{\tilde{w}_{3,3} t_2 u_2 \{1 + (s_2 + 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2\}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2} \\
&\quad - \frac{\tilde{w}_{3,3} t_2 u_2 (1 + s_2) \sqrt{s_2^2 - 2(1 - 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2 + 1}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2} \\
\\
\tilde{w}_{1,3} &= \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,3} t_1 u_1}{1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1)v_1} \Bigg|_{v_1 = \frac{1 - s_1 - \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,3}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,3}(1 - t_1)(1 - u_1)}} \\
&= \frac{\tilde{w}_{2,3} t_1 u_1 \{1 + (s_1 + 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1\}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1} \\
&\quad - \frac{\tilde{w}_{2,3} t_1 u_1 (1 + s_1) \sqrt{s_1^2 - 2(1 - 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1 + 1}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1}
\end{aligned}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$

terms where $m > 3$, I obtain every $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(x)$ terms.

Substitute (3.2.12), (3.2.15), (3.2.20), (3.2.21) and including all

$$\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(x) \text{ terms where } m > 3 \text{ into (3.2.10).} \quad \blacksquare$$

Remark 3.2.5 *The generating function for Heun polynomial of type 2 of the first kind*

as $q = -(q_j + 2j)\{\alpha + \beta - \delta + a(\delta + \gamma - 1) + (1 + a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} HF_{q_j}^R \left(q_j = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j \right. \\
& \left. , \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_1 = \frac{\varphi}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
& = 2^{\frac{\varphi}{(1+a)}-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^\gamma \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\alpha} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^\beta \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\beta} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\beta} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Big\} \quad (3.2.22)
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{cases}$$

and

$$\left\{ \begin{aligned} \mathbf{A}(s_{0,\infty}; \eta) &= \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{1-\gamma}}{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\Omega_1 - \gamma} \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{1-\gamma}}{\left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\Omega_1 - \gamma} \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{1-\gamma}}{\left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\Omega_1 - \gamma} \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \right.$$

Proof Replace $\mathbf{A}$, w and x by $\frac{\varphi}{(1+a)}$, $s_{0,\infty}$ and η in (3.2.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_{0,\infty}^{q_0} {}_2F_1 \left(-q_0, q_0 + \frac{\varphi}{(1+a)}; \gamma; \eta \right) \\ &= 2^{\frac{\varphi}{(1+a)} - 1} \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{1-\gamma}}{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{\varphi}{(1+a)} - \gamma} \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \end{aligned} \quad (3.2.23)$$

Replace $\mathbf{A}$, w and x by $\frac{\varphi}{(1+a)}$, s_0 and $\tilde{w}_{1,1}$ in (3.2.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1 \left(-q_0, q_0 + \frac{\varphi}{(1+a)}; \gamma; \tilde{w}_{1,1} \right) \\ &= 2^{\frac{\varphi}{(1+a)} - 1} \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{1-\gamma}}{\left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{\varphi}{(1+a)} - \gamma} \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (3.2.24)$$

Replace $\mathbf{A}$, w and x by $\frac{\varphi}{(1+a)}$, s_0 and $\tilde{w}_{1,n}$ in (3.2.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1 \left(-q_0, q_0 + \frac{\varphi}{(1+a)}; \gamma; \tilde{w}_{1,n} \right) \\ &= 2^{\frac{\varphi}{(1+a)} - 1} \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{1-\gamma}}{\left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{\varphi}{(1+a)} - \gamma} \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (3.2.25)$$

Put $c_0 = 1$, $\lambda = 0$ and $\gamma' = \gamma$ in (3.2.9). Substitute (3.2.23), (3.2.24) and (3.2.25) into the new (3.2.9). ■

Remark 3.2.6 *The generating function for Heun polynomial of type 2 of the second kind as $q = -(q_j + 2j + 1 - \gamma)\{\alpha + \beta + 1 - \gamma - (1 - a)\delta + (1 + a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} HS_{q_j}^R \left(q_j = \frac{-\{\varphi + 2(1+a)(1-\gamma)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j \right. \\
& , \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_2 = \frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \left. \right) \\
& = 2^{\frac{\varphi+2(1+a)(1/2-\gamma)}{(1+a)}} \eta^{1-\gamma} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{B}(s_{0,\infty}; \eta) \right. \\
& + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{2-\gamma} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \eta) \\
& \times \tilde{w}_{1,1}^{-(\alpha-\gamma+1)}(\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta}(\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta-\gamma+1} \mathbf{B}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\gamma} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-(2n-1+\alpha-\gamma)}(\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta}(\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\beta-\gamma} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)}(\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta}(\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\beta-\gamma} \left. \right\} \\
& \times \mathbf{B}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \left. \right\} \tag{3.2.26}
\end{aligned}$$

where

$$\left\{ \begin{array}{l} \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)} s_{1,\infty} + 1}{2} \right)^{-(3+\Omega_2)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)} s_{1,\infty} + 1} \\ \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)} s_{n,\infty} + 1}{2} \right)^{-(4n-1+\Omega_2)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)} s_{n,\infty} + 1} \\ \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{(1+s_{n-k}) + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))} s_{n-k} + 1}{2} \right)^{-(4(n-k)-1+\Omega_2)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))} s_{n-k} + 1} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{B}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{2-\gamma-\Omega_2}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{1-\gamma} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{2-\gamma-\Omega_2}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{2-\gamma-\Omega_2}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Proof Replace γ , A , w and x by $2-\gamma$, $\frac{\varphi+2(1+a)(1-\gamma)}{(1+a)}$, $s_{0,\infty}$ and η in (3.2.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_{0,\infty}^{q_0} {}_2F_1 \left(-q_0, q_0 + \frac{\varphi+2(1+a)(1-\gamma)}{(1+a)}; 2-\gamma; \eta \right) \\ &= 2^{\frac{\varphi+2(1+a)(1/2-\gamma)}{(1+a)}} \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{-\frac{\varphi-(1+a)\gamma}{(1+a)}}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{1-\gamma} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \end{aligned} \quad (3.2.27)$$

Replace γ , A , w and x by $2 - \gamma$, $\frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}$, s_0 and $\tilde{w}_{1,1}$ in (3.2.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1 \left(-q_0, q_0 + \frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}; 2-\gamma; \tilde{w}_{1,1} \right) \\ &= 2^{\frac{\varphi+2(1+a)(1/2-\gamma)}{(1+a)}} \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{-\frac{\varphi-(1+a)\gamma}{(1+a)}}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (3.2.28)$$

Replace γ , A , w and x by $2 - \gamma$, $\frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}$, s_0 and $\tilde{w}_{1,n}$ in (3.2.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1 \left(-q_0, q_0 + \frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}; 2-\gamma; \tilde{w}_{1,n} \right) \\ &= 2^{\frac{\varphi+2(1+a)(1/2-\gamma)}{(1+a)}} \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{-\frac{\varphi-(1+a)\gamma}{(1+a)}}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (3.2.29)$$

Put $c_0 = \left(\frac{1+a}{a} \right)^{1-\gamma}$, $\lambda = 1 - \gamma$ and $\gamma' = 2 - \gamma$ in (3.2.9). Substitute (3.2.27), (3.2.28) and (3.2.29) into the new (3.2.9). ■

3.3 Summary

As we see all solutions of power series expansions in Heun function by using 3TRF, denominators and numerators in all B_n terms arise with Pochhammer symbol.[3] Also the Frobenius solutions in Heun function by using R3TRF, denominators and numerators in all A_n terms arise with Pochhammer symbol: the meaning of this is that the analytic solutions of Heun's ordinary differential equations with three recursive coefficients can be described as Hypergeometric functions in a strict mathematical way.

We can express representations in closed form integrals of Heun function since we have power series expansions with Pochhammer symbols in numerators and denominators: (1) look at Remark 1 through Remark 6 in Ref.[4]. (2) look at Remark 2.3.2 through Remark 2.3.5 in chapter 2. We can transform any special functions into all other well-known special functions such as Legendre, Bessel, Laguerre, Kummer functions and etc because a ${}_2F_1$ function recurs in each of sub-integral forms of Heun function. After we replace ${}_2F_1$

functions in integral forms of Heun function to other special functions, we can rebuild the power series expansion of Heun function and its linear ordinary differential equation in a backward: understanding the connection between other special functions is important in the mathematical and physical points of views as we all know.

Since integral representation of Heun function is derived from power series expansion in closed forms, I construct generating function for Heun polynomial of type 2 by applying the generating function for Jacobi polynomial using hypergeometric functions into the general integral representation of Heun polynomial. The generating function is really helpful in order to derive orthogonal relations, recursion relations and expectation values of any physical quantities as we all recognize; i.e. the normalized wave function of hydrogen-like atoms and expectation values of its physical quantities such as position and momentum.

Appendices

.1 Generating functions for 192 Heun polynomials of type 2

In appendix of chapter 2, by applying R3TRF, I construct the fundamental power series expansions in closed forms and integral forms of Heun function (infinite series and polynomial of type 2) of nine out of the 192 local solution of Heun function in Table 2, obtained by Robert S. Maier(2007) [2].

In this appendix, replacing coefficients in the general expression of generating function for Heun polynomial of type 2, I derive generating functions for Heun polynomial of type 2 for the previous nine examples of the 192 local solution of Heun equation [2].⁵

⁵I treat α , β , γ and δ as free variables and a fixed value of q to construct the generating functions for Heun polynomial of type 2 for all nine examples of the 192 local solution of Heun function.

$$\mathbf{.1.1} \quad (1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x)$$

Replacing coefficients q , α , β and δ by $q - (\delta-1)\gamma a$, $\alpha - \delta + 1$, $\beta - \delta + 1$ and $2 - \delta$ into (3.2.22). Multiply $(1-x)^{1-\delta}$ and the new (3.2.22) together.

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} (1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x) \\
&= \frac{2^{\frac{\varphi}{(1+a)}-1}}{(1-x)^{\delta-1}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\gamma} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
&\times \tilde{w}_{1,1}^{-\alpha+\delta-1} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta-\delta+1} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \tilde{w}_{n,n}^{-(2n-1+\alpha-\delta)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\beta-\delta} \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
&\times \tilde{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\delta)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \left. \right\} \\
&\times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Big\} \tag{.1.1}
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - \delta + a(\gamma - \delta + 1) \\ \Omega_1 = \frac{\varphi}{(1+a)} \\ q = (\delta-1)\gamma a - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\gamma - 1} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\gamma - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\gamma - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

.1.2 $x^{1-\gamma}(1-x)^{1-\delta} Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x)$

Replacing coefficients q, α, β, γ and δ by $q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1)$, $\alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma$ and $2 - \delta$ into (3.2.22). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and

the new (3.2.22) together.

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} x^{1-\gamma} (1-x)^{1-\delta} Hl(a, q - (\gamma + \delta - 2)a \\
& \quad - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1); \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x) \\
& = \frac{2^{\frac{\varphi}{(1+a)}-1}}{x^{\gamma-1}(1-x)^{\delta-1}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\
& \quad + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{2-\gamma} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
& \quad \times \tilde{w}_{1,1}^{-\alpha+\gamma+\delta-2} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta-\gamma-\delta+2} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& \quad + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \quad \times \tilde{w}_{n,n}^{-(2n+\alpha-\gamma-\delta)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n+\beta-\gamma-\delta} \\
& \quad \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \quad \times \tilde{w}_{n-k,n}^{-(2(n-k)+\alpha-\gamma-\delta)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)+\beta-\gamma-\delta} \left. \right\} \\
& \quad \times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Bigg\} \tag{.1.2}
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - 2\gamma - \delta + 2 + a(3 - \gamma - \delta) \\ \Omega_1 = \frac{\varphi}{(1+a)} \\ q = (\gamma + \delta - 2)a + (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1) - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{2-\gamma-\Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{1-\gamma} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{2-\gamma-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{2-\gamma-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

.1.3 $Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x)$

Replacing coefficients a, q, γ, δ and x by $1-a, -q + \alpha\beta, \delta, \gamma$ and $1-x$ into (3.2.22).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x) \\
&= 2^{\frac{\varphi}{(2-a)}-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^\delta \overleftarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
&\times \tilde{w}_{1,1}^{-\alpha} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^\beta \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\delta} \overleftarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\beta} \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \overleftarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
&\times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\beta} \left. \right\} \\
&\times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Bigg\} \tag{.1.3}
\end{aligned}$$

where

$$\begin{cases} \xi = 1-x \\ z = \frac{-1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = \alpha + \beta - \delta + (1-a)(\delta + \gamma - 1) \\ \Omega_1 = \frac{\varphi}{(2-a)} \\ q = \alpha\beta + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\delta - \Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\delta - 1} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\delta - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\delta - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\delta - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\delta - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

.1.4 $(1-x)^{1-\delta} Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x)$

Replacing coefficients $a, q, \alpha, \beta, \gamma, \delta$ and x by $1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma$ and $1-x$ into

(3.2.22). Multiply $(1-x)^{1-\delta}$ and the new (3.2.22) together.

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} (1-x)^{1-\delta} \\
& \times Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x) \\
& = \frac{2^{\frac{\varphi}{(2-a)}-1}}{(1-x)^{\delta-1}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\
& + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{2-\delta} \overleftarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
& \times \tilde{w}_{1,1}^{-\alpha+\delta-1} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta-\delta+1} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\delta} \overleftarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-(2n-1+\alpha-\delta)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\beta-\delta} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\delta} \overleftarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\delta)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \left. \right\} \\
& \times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Bigg\} \tag{.1.4}
\end{aligned}$$

where

$$\begin{cases} \xi = 1-x \\ z = \frac{-1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = \alpha + \beta - \gamma - 2\delta + 2 + (1-a)(\gamma - \delta + 1) \\ \Omega_1 = \frac{\varphi}{(2-a)} \\ q = (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1) + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{2-\delta-\Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{1-\delta} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{2-\delta-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{1-\delta} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{2-\delta-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{1-\delta} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

$$\mathbf{.1.5} \quad x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1, \alpha - \beta + 1, \delta; \frac{1}{x} \right)$$

Replacing coefficients a, q, β, γ and x by $\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}, \alpha - \gamma + 1, \alpha - \beta + 1$ and $\frac{1}{x}$ into (3.2.22). Multiply $x^{-\alpha}$ and the new (3.2.22) together.

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1 \right. \\ & \left. , \alpha - \beta + 1, \delta; \frac{1}{x} \right) \\ &= \frac{2^{\frac{a\varphi}{(1+a)}-1}}{x^{\alpha}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\ & + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\alpha-\beta+1} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\ & \times \tilde{w}_{1,1}^{-\alpha} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\gamma-1} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\gamma+1} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ & + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-1+\alpha-\beta} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ & \times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\gamma-1} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\alpha-\gamma} \\ & \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1+\alpha-\beta} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ & \times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\gamma-1} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \left. \right\} \\ & \left. \times \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} z^n \Bigg\} \end{aligned} \quad (.1.5)$$

where

$$\begin{cases} \xi = \frac{1}{x} \\ z = -a\xi^2 \\ \eta = (1+a)\xi \\ \varphi = 2\alpha - \gamma - \delta + 1 + \frac{1}{a}(\alpha - \beta + \delta) \\ \Omega_1 = \frac{a\varphi}{(1+a)} \\ q = -\alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta] - a(q_j + 2j)\{\varphi + (1 + 1/a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\alpha-\beta+1-\Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\alpha-\beta} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\alpha-\beta+1-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\alpha-\beta} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\alpha-\beta+1-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\alpha-\beta} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

$$\mathbf{.1.6} \quad \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right)$$

Replacing coefficients a , q , α and x by $1 - a$, $-q + \gamma\beta$, $-\alpha + \gamma + \delta$ and $\frac{(1-a)x}{x-a}$ into (3.2.22). Multiply $\left(1 - \frac{x}{a}\right)^{-\beta}$ and the new (3.2.22) together.

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right) \\ &= \frac{2^{\frac{\varphi}{(2-a)}-1}}{\left(1 - \frac{x}{a}\right)^{\beta}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\ &+ \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\gamma} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\ &\times \tilde{w}_{1,1}^{\alpha-\gamma-\delta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\alpha-\beta+\gamma+\delta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ &\times \tilde{w}_{n,n}^{-(2(n-1)-\alpha+\gamma+\delta)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{-\alpha-\beta+\gamma+\delta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\beta} \\ &\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ &\times \tilde{w}_{n-k,n}^{-(2(n-k-1)-\alpha+\gamma+\delta)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{-\alpha-\beta+\gamma+\delta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\beta} \\ &\left. \left. \times \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} z^n \right\} \end{aligned} \quad (.1.6)$$

where

$$\begin{cases} \xi = \frac{(1-a)x}{x-a} \\ z = -\frac{1}{1-a} \xi^2 \\ \eta = \frac{2-a}{1-a} \xi \\ \varphi = -\alpha + \beta + \gamma + (1-a)(\gamma + \delta - 1) \\ \Omega_1 = \frac{\varphi}{(2-a)} \\ q = \gamma\beta + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\gamma - 1} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\gamma - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\gamma - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

.1.7

$$(1-x)^{1-\delta} \left(1 - \frac{x}{a} \right)^{-\beta+\delta-1} Hl \left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma, 2 - \delta; \frac{(1-a)x}{x-a} \right)$$

Replacing coefficients $a, q, \alpha, \beta, \delta$ and x by $1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1], -\alpha + \gamma + 1, \beta - \delta + 1, 2 - \delta$ and $\frac{(1-a)x}{x-a}$ into (3.2.22). Multiply $(1-x)^{1-\delta} \left(1 - \frac{x}{a} \right)^{-\beta+\delta-1}$

and the new (3.2.22) together.

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} \\
& \times Hl \left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma, 2 - \delta; \frac{(1-a)x}{x-a} \right) \\
& = \frac{2^{\frac{\varphi}{(2-a)}-1} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}}{(1-x)^{\delta-1}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\
& + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\gamma} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
& \times \tilde{w}_{1,1}^{\alpha-\gamma-1} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\alpha-\beta+\gamma+\delta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta-\delta+1} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-(2n-1-\alpha+\gamma)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{-\alpha-\beta+\gamma+\delta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\beta-\delta} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-(2(n-k)-1-\alpha+\gamma)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{-\alpha-\beta+\gamma+\delta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\beta-\delta} \left. \right\} \\
& \times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \quad (.1.7)
\end{aligned}$$

where

$$\begin{cases} \xi = \frac{(1-a)x}{x-a} \\ z = -\frac{1}{1-a} \xi^2 \\ \eta = \frac{(2-a)}{1-a} \xi \\ \varphi = -\alpha + \beta + \gamma + (1-a)(\gamma - \delta + 1) \\ \Omega_1 = \frac{\varphi}{(2-a)} \\ q = \gamma[(\delta-1)a + \beta - \delta + 1] + (q_j + 2j)\{\varphi + (2-a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\gamma - 1} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\gamma - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\gamma - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\gamma - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

$$\mathbf{.1.8} \quad x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}; \alpha, \alpha - \gamma + 1, \delta, \alpha - \beta + 1; \frac{x-1}{x} \right)$$

Replacing coefficients $a, q, \beta, \gamma, \delta$ and x by $\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}, \alpha - \gamma + 1, \delta, \alpha - \beta + 1$ and $\frac{x-1}{x}$ into (3.2.22). Multiply $x^{-\alpha}$ and the new (3.2.22) together.

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}; \alpha, \alpha - \gamma + 1, \delta, \right. \\ & \quad \left. \alpha - \beta + 1; \frac{x-1}{x} \right) \\ &= \frac{2^{\frac{a\varphi}{(2a-1)}-1}}{x^\alpha} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\ & \quad + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^\delta \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\ & \quad \times \tilde{w}_{1,1}^{-\alpha} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\gamma-1} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\gamma+1} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ & \quad + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\delta} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ & \quad \times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\gamma-1} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\alpha-\gamma} \\ & \quad \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ & \quad \times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\gamma-1} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \left. \right\} \\ & \quad \times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \end{aligned} \quad (.1.8)$$

where

$$\begin{cases} \xi = \frac{x-1}{x} \\ z = \frac{-a}{a-1} \xi^2 \\ \eta = \frac{2a-1}{a-1} \xi \\ \varphi = \alpha + \beta - \gamma + \frac{a-1}{a} (\alpha - \beta + \delta) \\ \Omega_1 = \frac{a\varphi}{(2a-1)} \\ q = \alpha(\delta a + \beta - \delta) + (q_j + 2j) \{a\varphi + (2a-1)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ \left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)} \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\delta - \Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\delta - 1} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\delta - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\delta - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\delta - \Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\delta - 1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

$$\mathbf{.1.9} \quad \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right)$$

Replacing coefficients q, β, γ, δ and x by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma$ and $\frac{a(x-1)}{x-a}$ into (3.2.22). Multiply $\left(\frac{x-a}{1-a} \right)^{-\alpha}$ and the new (3.2.22) together.

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right) \\ &= \frac{2^{\frac{\varphi}{(1+a)}-1}}{\left(\frac{x-a}{1-a} \right)^{\alpha}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\ &+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\delta} \overleftarrow{\mathbf{\Gamma}}_1(s_{1,\infty}; t_1, u_1, \eta) \\ &\times \tilde{w}_{1,1}^{-\alpha}(\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\beta-\gamma-\delta}(\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\beta+\gamma+\delta} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\delta} \overleftarrow{\mathbf{\Gamma}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ &\times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)}(\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\beta-\gamma-\delta}(\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)-\beta+\gamma+\delta} \\ &\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\delta} \overleftarrow{\mathbf{\Gamma}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ &\times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)}(\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\beta-\gamma-\delta}(\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)-\beta+\gamma+\delta} \left. \right\} \\ &\times \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Big\} \end{aligned} \quad (.1.9)$$

where

$$\begin{cases} \xi = \frac{a(x-1)}{x-a} \\ z = -\frac{1}{a} \xi^2 \\ \eta = \frac{(1+a)}{a} \xi \\ \varphi = \alpha - \beta + \delta + a(\gamma + \delta - 1) \\ \Omega_1 = \frac{\varphi}{(1+a)} \\ q = (\beta - \delta)\alpha - (q_j + 2j)\{\varphi + (1+a)(q_j + 2j)\} \text{ as } j, q_j \in \mathbb{N}_0 \end{cases}$$

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n})(1-t_{n-k})(1-u_{n-k})s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\delta-\Omega_1}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\delta-1} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\delta-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\delta-1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\delta-\Omega_1}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\delta-1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Bibliography

- [1] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische. Annalen.* **33**, 161(1889).
- [2] Maier, R.S., “The 192 solutions of the Heun equation,” *Math. Comp.* **33**, (2007)811–843.
- [3] Choun, Y.S., “The analytic solution for the power series expansion of Heun function,” <http://dx.doi.org/10.1016/j.aop.2013.06.020> Available as arXiv:1303.0830.
- [4] Choun, Y.S., “Asymptotic behavior of Heun function and its integral formalism,” arXiv:1303.0876.
- [5] Choun, Y.S., “The generating functions of Lamé equation in Weierstrass’s form,” arXiv:1303.0879.
- [6] Lamé, G., Sur les surfaces isothermes dans les corps homogènes en équilibre de température. *J. Math. Pures. Appl.*, 2, 147-188 (French)(1837)
- [7] Kalnins, E. G. and Miller, W. Jr., “Hypergeometric Expansions of Heun Polynomials,” *SIAM J. Math. Anal.* **22**(5), 1450-1459(1991).
- [8] Kalnins, E. G. and Miller, W., Jr., “Jacobi elliptic coordinates, functions of Heun and Lamé type and the Niven transform,” *Regul. Chaotic. Dyn.* **10**(4), 487-508(2005).
- [9] Bateman, H., Erdélyi, A., *Higher transcendental functions 3. Automorphic functions*, McGraw-Hill, (1955).
- [10] Arscott, F.M., *Periodic differential equations*, (Pergamon Press, 1964).
- [11] Svartholm, N., “Die Lösung der Fuchs’schen Differentialgleichung zweiter Ordnung durch Hypergeometrische Polynome,” *Math. Ann.* **116**, 413–421(1939).

- [12] Erdélyi, A., “The Fuchsian equation of second order with four singularities,” *Duke Math. J.* 9, 48–58(1942).
- [13] Erdélyi, A., “Integral equations for Heun functions,” *Q. J. Math. (Oxford)* 13, 107–112(1942).
- [14] Erdélyi, A., “Certain expansions of solutions of the Heun equation,” *Q. J. Math. (Oxford)* 15, 62–69(1944).
- [15] Sleeman, B.D., “Integral representations for solutions of Heun’s equation,” *Proc. Camb. Phil. Sci.* 65, 447–459(1969).
- [16] Sleeman, B.D., “Non-linear integral equations for Heun functions,” *Proc. Edinburgh Math. Sci. (2)* 16, 281–299(1969).
- [17] Schmidt, D., Wolf, G., “A method of generating integral relations by the simultaneous separability of generalized Schrödinger equations,” *SIAM J. Math. Anal.* 10, 823–838(1979).
- [18] M. Hortacsu, “Heun Functions and their uses in Physics,” [arXiv:1101.0471](https://arxiv.org/abs/1101.0471)

Chapter 4

Confluent Heun function using three-term recurrence formula

The Confluent Heun equation (CHE) is one of 4 confluent forms of Heun's differential equation. The Confluent Heun function (CHF) is applicable to areas such as general relativity, theory of rotating/non-rotating black hole, the gauge theories on thick brane worlds, hydrogen molecule ion in Stark effect and etc.

In this chapter I will apply three term recurrence formula (3TRF) [1] to the power series expansion in closed forms and its integral forms of the CHE for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's.¹ Indeed, I construct the generating function for Confluent Heun polynomial (CHP) which makes B_n term terminated.

4.1 Introduction

4.1.1 Non-symmetrical canonical form of Confluent Heun Equation

In Ref.[1, 2], Heun's equation is a second-order linear ordinary differential equation of the form

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0 \quad (4.1.1)$$

¹“ higher terms of A_n 's” means at least two terms of A_n 's.

With the condition $\epsilon = \alpha + \beta - \gamma - \delta + 1$. The parameters play different roles: $a \neq 0$ is the singularity parameter, $\alpha, \beta, \gamma, \delta, \epsilon$ are exponent parameters, q is the accessory parameter. Also, α and β are identical to each other. The total number of free parameters is six. It has four regular singular points which are $0, 1, a$ and ∞ with exponents $\{0, 1 - \gamma\}, \{0, 1 - \delta\}, \{0, 1 - \epsilon\}$ and $\{\alpha, \beta\}$.

Heun equation has the four kinds of confluent forms: (1) Confluent Heun (two regular and one irregular singularities), (2) Doubly Confluent Heun (two irregular singularities), (3) Biconfluent Heun (one regular and one irregular singularities)², (4) Triconfluent Heun equations (one irregular singularity). We can derive these four confluent forms from Heun equation by combining two or more regular singularities to take form an irregular singularity. Its process, converting Heun equation to other confluent forms, is similar to deriving of confluent hypergeometric equation from the hypergeometric equation. For the CHE, divide (4.1.1) by a to get

$$x(x-1) \left(\frac{x}{a} - 1 \right) \frac{d^2 y}{dx^2} + \left(\gamma(x-1) \left(\frac{x}{a} - 1 \right) + \delta x \left(\frac{x}{a} - 1 \right) + \frac{\epsilon}{a} x(x-1) \right) \frac{dy}{dx} + \left(\alpha \frac{\beta}{a} x - \frac{q}{a} \right) y = 0$$

Let $a, \beta, \epsilon, q \rightarrow \infty$ in such a way that

$$\frac{\beta}{a} \rightarrow \frac{\beta}{a} \rightarrow -\beta, \frac{q}{a} \rightarrow -q$$

the new β and q being constants, we find the non-symmetrical canonical form of the Confluent Heun Equation (CHE).[2, 21, 8]

$$\frac{d^2 y}{dx^2} + \left(\beta + \frac{\gamma}{x} + \frac{\delta}{x-1} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)} y = 0 \quad (4.1.2)$$

(4.1.2) has three singular points: two regular singular points which are 0 and 1 with exponents $\{0, 1 - \gamma\}$ and $\{0, 1 - \delta\}$, and one irregular singular point which is ∞ with an exponent α . Its solution is denoted as $H_c(\alpha, \beta, \gamma, \delta, q; x)$.³

4.1.2 Generalized spheroidal equation (GSE)

The Bôcher symmetrical form of the CHE in applications of mathematical physics is written as [9, 10]

$$\frac{d}{dz}(z^2 - 1) \frac{d}{dz} y(z) + \left(-p^2(z^2 - 1) + 2p\beta z - \lambda - \frac{m^2 + s^2 + 2msz}{(z^2 - 1)} \right) y(z) = 0 \quad (4.1.3)$$

²Biconfluent Heun equation is derived from the grand confluent hypergeometric equation by changing all coefficients $\mu = 1$ and $\varepsilon\omega = -q$. [36, 37]

³Several authors denote as coefficients $4p$ and σ instead of β and q . And they define the solution of the CHE as $H_c^{(a)}(p, \alpha, \gamma, \delta, \sigma; x)$.

It is called the generalized spheroidal equation (GSE). The GSE has three singular points: two regular singular points which are ± 1 and one irregular singular point which is ∞ . Putting $y(z) = (1 - z^2)^{-\frac{1}{2}}v(z)$ in (4.1.2), we obtain the normal symmetrical form of the CHE:

$$(z^2 - 1)\frac{d^2v}{dz^2} + \left(-p^2(z^2 - 1) + 2p\beta z - \lambda - \frac{m^2 + s^2 - 1 + 2msz}{(z^2 - 1)}\right)v(z) = 0 \quad (4.1.4)$$

It is called a normal form of the GSE. By putting $y(z) = (z - 1)^{\pm\frac{1}{2}(m+s)}(z + 1)^{\pm\frac{1}{2}(m-s)}e^{\pm pz}w(z)$ into (4.1.3), The solution of the GSE can be described in the form of non-symmetrical canonical form of the CHE. The 16 possible solutions are as follows.

-
1. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{pz}H_c(\beta+m+1, 4p, m-s+1, m+s+1, 2p(\beta+1)-m(m+1)-\lambda; \frac{1+z}{2})$
 2. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta+m+1, -4p, m-s+1, m+s+1, 2p(\beta-1)-m(m+1)-\lambda; \frac{1+z}{2})$
 3. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{pz}H_c(\beta+m+1, -4p, m+s+1, m-s+1, -2p(\beta+2m+1)-m(m+1)-\lambda; \frac{1-z}{2})$
 4. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta+m+1, 4p, m+s+1, m-s+1, -2p(\beta-2m-1)-m(m+1)-\lambda; \frac{1-z}{2})$
 5. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{pz}H_c(\beta-m+1, 4p, -m+s+1, -m-s+1, 2p(\beta+1)+m(-m+1)-\lambda; \frac{1+z}{2})$
 6. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta-m+1, -4p, -m+s+1, -m-s+1, 2p(\beta-1)+m(-m+1)-\lambda; \frac{1+z}{2})$
 7. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{pz}H_c(\beta-m+1, -4p, -m-s+1, -m+s+1, -2p(\beta-2m+1)+m(-m+1)-\lambda; \frac{1-z}{2})$
 8. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta-m+1, 4p, -m-s+1, -m+s+1, -2p(\beta-2m-1)+m(-m+1)-\lambda; \frac{1-z}{2})$
 9. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{pz}H_c(\beta+s+1, 4p, -m+s+1, m+s+1, 2p(\beta+1)-s(s+1)-\lambda; \frac{1+z}{2})$
 10. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta+s+1, -4p, -m+s+1, m+s+1, 2p(\beta-1)-s(s+1)-\lambda; \frac{1+z}{2})$
 11. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{pz}H_c(\beta+s+1, -4p, m+s+1, -m+s+1, -2p(\beta+2s+1)-s(s+1)-\lambda; \frac{1-z}{2})$
 12. $y(z) = (z-1)^{\frac{1}{2}(m+s)}(z+1)^{-\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta+s+1, 4p, m+s+1, -m+s+1, -2p(\beta-2s-1)-s(s+1)-\lambda; \frac{1-z}{2})$
 13. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{pz}H_c(\beta-s+1, 4p, m-s+1, -m-s+1, 2p(\beta+1)+s(-s+1)-\lambda; \frac{1+z}{2})$
 14. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta-s+1, -4p, m-s+1, -m-s+1, 2p(\beta-1)+s(-s+1)-\lambda; \frac{1+z}{2})$
 15. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{pz}H_c(\beta-s+1, -4p, -m-s+1, m-s+1, -2p(\beta-2s+1)+s(-s+1)-\lambda; \frac{1-z}{2})$
 16. $y(z) = (z-1)^{-\frac{1}{2}(m+s)}(z+1)^{\frac{1}{2}(m-s)}e^{-pz}H_c(-\beta-s+1, 4p, -m-s+1, m-s+1, -2p(\beta+2s+1)+s(-s+1)-\lambda; \frac{1-z}{2})$
-

Table 4.1: The 16 possible solutions of the GSE using non-symmetrical canonical form of the CHE

4.1.3 Generalized spheroidal equation in the Leaver version

The generalized spheroidal wave equation (GSWE), one of standard forms of the CHE, in the Leaver version reads [6]

$$z(z - z_0) \frac{d^2 U}{dz^2} + (B_1 + B_2 z) \frac{dU}{dz} + (B_3 - 2\eta\omega(z - z_0) + \omega^2 z(z - z_0)) U(z) = 0 \quad \text{where } \omega \neq 0 \quad (4.1.5)$$

where B_1, B_2, B_3, η and ω are constant. (4.1.5) has three singular points: two regular singular points which are 0 and z_0 with exponents $\{0, 1 + \frac{B_1}{z_0}\}$ and $\{0, 1 - B_2 - \frac{B_1}{z_0}\}$, and one irregular singular point which is ∞ . To permit the non-symmetrical canonical form of the CHE limit, the first step is given by the substitutions

$$z = z_0(1 - x), \quad U(z) = \exp(i\omega z_0(1 - x)) y(x)$$

which convert (4.1.5) into

$$\frac{d^2 y}{dx^2} + \left(-2i\omega z_0 + \frac{B_2 + \frac{B_1}{z_0}}{x} + \frac{\frac{-B_1}{z_0}}{x - 1} \right) \frac{dy}{dx} + \frac{-2i\omega z_0 \left(i\eta + \frac{B_2}{2} \right) x - \left(-B_2 - i\omega z_0 \left(B_2 + \frac{B_1}{z_0} \right) \right)}{x(x - 1)} y(x) = 0 \quad (4.1.6)$$

As we compare all coefficients in (4.1.6) with (4.1.2), we find that

$$\begin{cases} \alpha \rightarrow i\eta + \frac{B_2}{2} \\ \beta \rightarrow -2i\omega z_0 \\ \gamma \rightarrow B_2 + \frac{B_1}{z_0} \\ \delta \rightarrow -\frac{B_1}{z_0} \\ q \rightarrow -B_2 - i\omega z_0 \left(B_2 + \frac{B_1}{2} \right) \end{cases}$$

Then the solution $U(z)$ for the GSWE in the Leaver version is described by the non-symmetrical canonical form of the CHE.

$$\begin{aligned} & U(B_1, B_2, B_3; z_0, \omega, \eta; z) \\ &= \exp(i\omega z_0(1 - x)) H_c \left(i\eta + \frac{B_2}{2}, -2i\omega z_0, B_2 + \frac{B_1}{z_0}, -\frac{B_1}{z_0}, -B_2 - i\omega z_0 \left(B_2 + \frac{B_1}{2} \right); z_0(1 - x) \right) \end{aligned} \quad (4.1.7)$$

4.2 Confluent Heun function with a regular singular point at zero

In this chapter, by applying 3TRF [1], I'll construct the power series expansion in closed forms, the integral form and the generating function for non-symmetrical canonical form

of the CHE for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's: I prefer (4.1.2) as an analytic solution of the CHE rather than the GSE and its Leaver version because of more convenient mathematical calculations.

Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} \quad \text{where } \lambda = \text{indicial root} \quad (4.2.1)$$

Plug (4.2.1) into (4.1.2). Then we get a three-term recurrence relation for the coefficients c_n :

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (4.2.2)$$

where,

$$A_n = \frac{(n+\lambda)(n+\lambda-\beta+\gamma+\delta-1)-q}{(n+\lambda+1)(n+\lambda+\gamma)} \quad (4.2.3a)$$

$$B_n = \frac{\beta(n+\lambda+\alpha-1)}{(n+\lambda+1)(n+\lambda+\gamma)} \quad (4.2.3b)$$

$$c_1 = A_0 c_0 \quad (4.2.3c)$$

We have two indicial roots which are $\lambda_1 = 0$ and $\lambda_2 = 1 - \gamma$.

Now let's test for convergence of infinite series of the analytic function $y(x)$. As $n \gg 1$, (4.2.3a)–(4.2.3c) are

$$\lim_{n \gg 1} A_n = 1 \quad (4.2.4a)$$

$$\lim_{n \gg 1} B_n = \frac{\beta}{n} \quad (4.2.4b)$$

(4.2.4b) is negligible because of $n \gg 1$. Substitute (4.2.4a) into (4.2.2) with $B_n = 0$. For $n = 0, 1, 2, \dots$, it give

$$\begin{aligned} c_0 & \\ c_1 &= c_0 \\ c_2 &= c_0 \\ c_3 &= c_0 \\ &\vdots \quad \vdots \end{aligned} \quad (4.2.5)$$

When a function $y(x)$, analytic at $x = 0$, is expanded in a power series putting $c_0 = 1$ by using (4.2.5), we write

$$\lim_{n \gg 1} y(x) \approx \sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \quad \text{where } |x| < 1 \quad (4.2.6)$$

For polynomial which makes A_n terminated, (4.2.4b) is only available for the asymptotic behavior of the minimum $y(z)$.⁴ Substitute (4.2.4b) into (4.2.2) with $A_n = 0$. For $n = 0, 1, 2, \dots$, it gives

$$\begin{array}{ll}
 c_0 & c_1 \\
 c_2 = \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{3}{2})} \frac{\beta}{2} c_0 & c_3 = \frac{1}{1!} \frac{\beta}{2} c_1 \\
 c_4 = \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{5}{2})} \left(\frac{\beta}{2}\right)^2 c_0 & c_5 = \frac{1}{2!} \left(\frac{\beta}{2}\right)^2 c_1 \\
 c_6 = \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{7}{2})} \left(\frac{\beta}{2}\right)^3 c_0 & c_7 = \frac{1}{3!} \left(\frac{\beta}{2}\right)^3 c_1 \\
 \vdots & \vdots
 \end{array} \tag{4.2.7}$$

Put (4.2.7) into (4.2.1) taking $c_0 = c_1 = 1$ and $\lambda = 0$ for simplicity.

$$\lim_{n \gg 1} y(z) > 1 + e^\eta (x + \sqrt{\pi\eta} \operatorname{Erf}(\sqrt{\eta})) \quad \text{where } \eta = \frac{1}{2}\beta x^2 \tag{4.2.8}$$

On the above $\operatorname{Erf}(y)$ is an error function which is

$$\operatorname{Erf}(y) = \frac{2}{\sqrt{\pi}} \int_0^y dt e^{-t^2}$$

4.2.1 Power series

Polynomial of type 1

As I mentioned in chapter 2, there are three types of polynomials in three term recurrence relation of a linear ordinary differential equation: (1) type 1 is the polynomial which makes B_n term terminated: A_n term is not terminated, (2) type 2 is the polynomial which makes A_n term terminated: B_n term is not terminated, (3) type 3 is the polynomial which makes A_n and B_n terms terminated at the same time.⁵ In general the CHP is defined as type 3 polynomial where A_n and B_n terms terminated. The CHP comes from the CHE that has a fixed integer value of α , just as it has a fixed value of q . In three term recurrence relation, polynomial of type 3 I categorize as complete polynomial. In future papers I will derive type 3 CHP. In this chapter I construct the power series expansion, its integral forms and the generating function for the CHP of type 1: I treat β , γ , δ and q as free variables and α as a fixed value.

⁴(4.2.4a) is negligible for the minimum $y(z)$ because A_n term will be terminated at the specific eigenvalues.

⁵If A_n and B_n terms are not terminated, it turns to be infinite series.

In Ref.[1], the general expression of power series of $y(x)$ for polynomial of type 1 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\beta_0} \left(\prod_{i_1=0}^{i_0-1} B_{2i_1+1} \right) x^{2i_0+\lambda} + \sum_{i_0=0}^{\beta_0} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \sum_{i_2=i_0}^{\beta_1} \left(\prod_{i_3=i_0}^{i_2-1} B_{2i_3+2} \right) \right\} x^{2i_2+1+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\beta_0} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\beta_k} A_{2i_{2k}+k} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} B_{2i_{2k+1}+(k+1)} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\beta_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} B_{2i_{2N+1}+(N+1)} \right) \right\} \right\} x^{2i_{2N}+N+\lambda} \Big\} \quad (4.2.9)
\end{aligned}$$

In the above, $\beta_i \leq \beta_j$ only if $i \leq j$ where $i, j, \beta_i, \beta_j \in \mathbb{N}_0$.

For a polynomial, we need a condition, which is:

$$B_{2\beta_i+(i+1)} = 0 \quad \text{where } i = 0, 1, 2, \dots, \beta_i = 0, 1, 2, \dots \quad (4.2.10)$$

In the above, β_i is an eigenvalue that makes B_n term terminated at certain value of index n . (4.2.10) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (4.2.9).

In (4.2.3b) replace α by $-2\alpha_i - i - \lambda$. In (4.2.9) and (4.2.10) replace index β_i by α_i . Take the new (4.2.3a)–(4.2.3c), (4.2.10) and put them in (4.2.9). After the replacement process,

the general expression of power series of the CHE for polynomial of type 1 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(i_0 + \frac{\lambda}{2}) (i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \right. \\
&\quad \times \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \sum_{i_1=i_0}^{\alpha_1} \frac{(-\alpha_1)_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(-\alpha_1)_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \eta^{i_1} \Big\} x \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(i_0 + \frac{\lambda}{2}) (i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\alpha_k} \frac{(i_k + \frac{k}{2} + \frac{\lambda}{2}) (i_k + \frac{1}{2}(-\beta + \gamma + \delta + \lambda + k - 1)) - \frac{q}{4}}{(i_k + \frac{k}{2} + \frac{1}{2} + \frac{\lambda}{2})(i_k + \frac{k}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \right. \\
&\quad \times \frac{(-\alpha_k)_{i_k} (1 + \frac{k}{2} + \frac{\lambda}{2})_{i_{k-1}} (\frac{1}{2} + \frac{k}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_{k-1}}}{(-\alpha_k)_{i_{k-1}} (1 + \frac{k}{2} + \frac{\lambda}{2})_{i_k} (\frac{1}{2} + \frac{k}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\alpha_n} \frac{(-\alpha_n)_{i_n} (1 + \frac{n}{2} + \frac{\lambda}{2})_{i_{n-1}} (\frac{1}{2} + \frac{n}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_{n-1}}}{(-\alpha_n)_{i_{n-1}} (1 + \frac{n}{2} + \frac{\lambda}{2})_{i_n} (\frac{1}{2} + \frac{n}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_n}} \eta^{i_n} \Big\} x^n \Big\} \quad (4.2.11)
\end{aligned}$$

where

$$\begin{cases} \eta = \frac{1}{2}\beta x^2 \\ \alpha = -2\alpha_j - j - \lambda \\ \alpha_i \leq \alpha_j \text{ only if } i \leq j \text{ where } i, j, \alpha_i, \alpha_j \in \mathbb{N}_0 \end{cases}$$

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ for the second one in (4.2.11).

Remark 4.2.1 *The power series expansion of the CHE of the first kind for polynomial of*

type 1 about $x = 0$ as $\alpha = -(2\alpha_j + j)$ where $j, \alpha_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= H_c^{(a)} F_{\alpha_j} \left(\alpha = -(2\alpha_j + j), \beta, \gamma, \delta, q; \eta = \frac{1}{2}\beta x^2 \right) \\
&= \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1)_{i_0} \left(\frac{1}{2} + \frac{\gamma}{2}\right)_{i_0}} \eta^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{\alpha_0} \frac{i_0 \left(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1)\right) - \frac{q}{4}}{(i_0 + \frac{1}{2})(i_0 + \frac{\gamma}{2})} \frac{(-\alpha_0)_{i_0}}{(1)_{i_0} \left(\frac{1}{2} + \frac{\gamma}{2}\right)_{i_0}} \sum_{i_1=i_0}^{\alpha_1} \frac{(-\alpha_1)_{i_1} \left(\frac{3}{2}\right)_{i_0} \left(1 + \frac{\gamma}{2}\right)_{i_0}}{(-\alpha_1)_{i_0} \left(\frac{3}{2}\right)_{i_1} \left(1 + \frac{\gamma}{2}\right)_{i_1}} \eta^{i_1} \right\} x \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{i_0 \left(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1)\right) - \frac{q}{4}}{(i_0 + \frac{1}{2})(i_0 + \frac{\gamma}{2})} \frac{(-\alpha_0)_{i_0}}{(1)_{i_0} \left(\frac{1}{2} + \frac{\gamma}{2}\right)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\alpha_k} \frac{(i_k + \frac{k}{2}) \left(i_k + \frac{1}{2}(-\beta + \gamma + \delta + k - 1)\right) - \frac{q}{4}}{(i_k + \frac{k}{2} + \frac{1}{2})(i_k + \frac{k}{2} + \frac{\gamma}{2})} \frac{(-\alpha_k)_{i_k} \left(\frac{k}{2} + 1\right)_{i_{k-1}} \left(\frac{k}{2} + \frac{1}{2} + \frac{\gamma}{2}\right)_{i_{k-1}}}{(-\alpha_k)_{i_{k-1}} \left(\frac{k}{2} + 1\right)_{i_k} \left(\frac{k}{2} + \frac{1}{2} + \frac{\gamma}{2}\right)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\alpha_n} \frac{(-\alpha_n)_{i_n} \left(\frac{n}{2} + 1\right)_{i_{n-1}} \left(\frac{n}{2} + \frac{1}{2} + \frac{\gamma}{2}\right)_{i_{n-1}}}{(-\alpha_n)_{i_{n-1}} \left(\frac{n}{2} + 1\right)_{i_n} \left(\frac{n}{2} + \frac{1}{2} + \frac{\gamma}{2}\right)_{i_n}} \eta^{i_n} \right\} x^n \quad (4.2.12)
\end{aligned}$$

Remark 4.2.2 The power series expansion of the CHE of the second kind for polynomial of type 1 about $x = 0$ as $\alpha = -(2\alpha_j + j + 1 - \gamma)$ where $j, \alpha_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= H_c^{(a)} S_{\alpha_j} \left(\alpha = -(2\alpha_j + j + 1 - \gamma), \beta, \gamma, \delta, q; \eta = \frac{1}{2}\beta x^2 \right) \\
&= x^{1-\gamma} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{\left(\frac{3}{2} - \frac{\gamma}{2}\right)_{i_0} (1)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\alpha_0} \frac{\left(i_0 + \frac{1}{2} - \frac{\gamma}{2}\right) \left(i_0 + \frac{1}{2}(-\beta + \delta)\right) - \frac{q}{4}}{(i_0 + 1 - \frac{\gamma}{2})(i_0 + \frac{1}{2})} \frac{(-\alpha_0)_{i_0}}{\left(\frac{3}{2} - \frac{\gamma}{2}\right)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\alpha_1} \frac{(-\alpha_1)_{i_1} \left(2 - \frac{\gamma}{2}\right)_{i_0} \left(\frac{3}{2}\right)_{i_0}}{(-\alpha_1)_{i_0} \left(2 - \frac{\gamma}{2}\right)_{i_1} \left(\frac{3}{2}\right)_{i_1}} \eta^{i_1} \right\} x \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{\left(i_0 + \frac{1}{2} - \frac{\gamma}{2}\right) \left(i_0 + \frac{1}{2}(-\beta + \delta)\right) - \frac{q}{4}}{(i_0 + 1 - \frac{\gamma}{2})(i_0 + \frac{1}{2})} \frac{(-\alpha_0)_{i_0}}{\left(\frac{3}{2} - \frac{\gamma}{2}\right)_{i_0} (1)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\alpha_k} \frac{\left(i_k + \frac{k}{2} + \frac{1}{2} - \frac{\gamma}{2}\right) \left(i_k + \frac{1}{2}(-\beta + \delta + k)\right) - \frac{q}{4}}{(i_k + \frac{k}{2} + 1 - \frac{\gamma}{2})(i_k + \frac{k}{2} + \frac{1}{2})} \frac{(-\alpha_k)_{i_k} \left(\frac{k}{2} + \frac{3}{2} - \frac{\gamma}{2}\right)_{i_{k-1}} \left(\frac{k}{2} + 1\right)_{i_{k-1}}}{(-\alpha_k)_{i_{k-1}} \left(\frac{k}{2} + \frac{3}{2} - \frac{\gamma}{2}\right)_{i_k} \left(\frac{k}{2} + 1\right)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\alpha_n} \frac{(-\alpha_n)_{i_n} \left(\frac{n}{2} + \frac{3}{2} - \frac{\gamma}{2}\right)_{i_{n-1}} \left(\frac{n}{2} + 1\right)_{i_{n-1}}}{(-\alpha_n)_{i_{n-1}} \left(\frac{n}{2} + \frac{3}{2} - \frac{\gamma}{2}\right)_{i_n} \left(\frac{n}{2} + 1\right)_{i_n}} \eta^{i_n} \right\} x^n \quad (4.2.13)
\end{aligned}$$

Infinite series

In Ref.[1], the general expression of power series of $y(x)$ for infinite series is defined by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} B_{2i_1+1} \right) x^{2i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} B_{2i_3+2} \right) \right\} x^{2i_2+1+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} A_{2i_{2k}+k} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} B_{2i_{2k+1}+(k+1)} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} B_{2i_{2N+1}+(N+1)} \right) \right\} \right\} x^{2i_{2N}+N+\lambda} \Bigg\} \quad (4.2.14)
\end{aligned}$$

Substitute (4.2.3a)–(4.2.3c) into (4.2.14). The general expression of power series of the CHE for infinite series about $x = 0$ is

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^{\lambda} \left\{ \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\lambda}{2}) (i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(\frac{1}{2} + \frac{\alpha}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(\frac{1}{2} + \frac{\alpha}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \eta^{i_1} \Bigg\} x \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\lambda}{2}) (i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + \frac{k}{2} + \frac{\lambda}{2}) (i_k + \frac{1}{2}(-\beta + \gamma + \delta - 1 + k + \lambda)) - \frac{q}{4}}{(i_k + \frac{k}{2} + \frac{1}{2} + \frac{\lambda}{2})(i_k + \frac{k}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \right. \\
&\quad \times \frac{(\frac{k}{2} + \frac{\alpha}{2} + \frac{\lambda}{2})_{i_k} (\frac{k}{2} + 1 + \frac{\lambda}{2})_{i_{k-1}} (\frac{k}{2} + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_{k-1}}}{(\frac{k}{2} + \frac{\alpha}{2} + \frac{\lambda}{2})_{i_{k-1}} (\frac{k}{2} + 1 + \frac{\lambda}{2})_{i_k} (\frac{k}{2} + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_k}} \Bigg\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(\frac{n}{2} + \frac{\alpha}{2} + \frac{\lambda}{2})_{i_n} (\frac{n}{2} + 1 + \frac{\lambda}{2})_{i_{n-1}} (\frac{n}{2} + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_{n-1}}}{(\frac{n}{2} + \frac{\alpha}{2} + \frac{\lambda}{2})_{i_{n-1}} (\frac{n}{2} + 1 + \frac{\lambda}{2})_{i_n} (\frac{n}{2} + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_n}} \eta^{i_n} \Bigg\} x^n \Bigg\} \quad (4.2.15)
\end{aligned}$$

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ in (4.2.15).

Remark 4.2.3 *The power series expansion of the CHE of the first kind for infinite series about $x = 0$ using 3TRF is*

$$\begin{aligned}
y(x) &= H_c^{(a)} F \left(\alpha, \beta, \gamma, \delta, q; \eta = \frac{1}{2} \beta x^2 \right) \\
&= \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2})_{i_0}}{(1)_{i_0} (\frac{1}{2} + \frac{\gamma}{2})_{i_0}} \eta^{i_0} \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{i_0 (i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1)) - \frac{q}{4}}{(i_0 + \frac{1}{2})(i_0 + \frac{\gamma}{2})} \frac{(\frac{\alpha}{2})_{i_0}}{(1)_{i_0} (\frac{1}{2} + \frac{\gamma}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\frac{1}{2} + \frac{\alpha}{2})_{i_1} (\frac{3}{2})_{i_0} (1 + \frac{\gamma}{2})_{i_0}}{(\frac{1}{2} + \frac{\alpha}{2})_{i_0} (\frac{3}{2})_{i_1} (1 + \frac{\gamma}{2})_{i_1}} \eta^{i_1} \right\} x \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{i_0 (i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1)) - \frac{q}{4}}{(i_0 + \frac{1}{2})(i_0 + \frac{\gamma}{2})} \frac{(\frac{\alpha}{2})_{i_0}}{(1)_{i_0} (\frac{1}{2} + \frac{\gamma}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + \frac{k}{2}) (i_k + \frac{1}{2}(-\beta + \gamma + \delta - 1 + k)) - \frac{q}{4}}{(i_k + \frac{k}{2} + \frac{1}{2})(i_k + \frac{k}{2} + \frac{\gamma}{2})} \right. \\
&\quad \times \left. \frac{(\frac{k}{2} + \frac{\alpha}{2})_{i_k} (\frac{k}{2} + 1)_{i_{k-1}} (\frac{k}{2} + \frac{1}{2} + \frac{\gamma}{2})_{i_{k-1}}}{(\frac{k}{2} + \frac{\alpha}{2})_{i_{k-1}} (\frac{k}{2} + 1)_{i_k} (\frac{k}{2} + \frac{1}{2} + \frac{\gamma}{2})_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\frac{n}{2} + \frac{\alpha}{2})_{i_n} (\frac{n}{2} + 1)_{i_{n-1}} (\frac{n}{2} + \frac{1}{2} + \frac{\gamma}{2})_{i_{n-1}}}{(\frac{n}{2} + \frac{\alpha}{2})_{i_{n-1}} (\frac{n}{2} + 1)_{i_n} (\frac{n}{2} + \frac{1}{2} + \frac{\gamma}{2})_{i_n}} \eta^{i_n} \right\} x^n \tag{4.2.16}
\end{aligned}$$

Remark 4.2.4 *The power series expansion of the CHE of the second kind for infinite*

series for infinite series about $x = 0$ using 3TRF is

$$\begin{aligned}
y(x) &= H_c^{(a)} S \left(\alpha, \beta, \gamma, \delta, q; \eta = \frac{1}{2} \beta x^2 \right) \\
&= x^{1-\gamma} \left\{ \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_0}}{(\frac{3}{2} - \frac{\gamma}{2})_{i_0} (1)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} - \frac{\gamma}{2}) (i_0 + \frac{1}{2} (-\beta + \delta)) - \frac{q}{4} (\frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_0}}{(i_0 + 1 - \frac{\gamma}{2}) (i_0 + \frac{1}{2}) (\frac{3}{2} - \frac{\gamma}{2})_{i_0} (1)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(1 + \frac{\alpha}{2} - \frac{\gamma}{2})_{i_1} (2 - \frac{\gamma}{2})_{i_0} (\frac{3}{2})_{i_0}}{(1 + \frac{\alpha}{2} - \frac{\gamma}{2})_{i_0} (2 - \frac{\gamma}{2})_{i_1} (\frac{3}{2})_{i_1}} \eta^{i_1} \Big\} x \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} - \frac{\gamma}{2}) (i_0 + \frac{1}{2} (-\beta + \delta)) - \frac{q}{4} (\frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_0}}{(i_0 + 1 - \frac{\gamma}{2}) (i_0 + \frac{1}{2}) (\frac{3}{2} - \frac{\gamma}{2})_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + \frac{k}{2} + \frac{1}{2} - \frac{\gamma}{2}) (i_k + \frac{1}{2} (-\beta + \delta + k)) - \frac{q}{4}}{(i_k + \frac{k}{2} + 1 - \frac{\gamma}{2}) (i_k + \frac{k}{2} + \frac{1}{2})} \right. \\
&\quad \times \frac{(\frac{k}{2} + \frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_k} (\frac{k}{2} + \frac{3}{2} - \frac{\gamma}{2})_{i_{k-1}} (\frac{k}{2} + 1)_{i_{k-1}}}{(\frac{k}{2} + \frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_{k-1}} (\frac{k}{2} + \frac{3}{2} - \frac{\gamma}{2})_{i_k} (\frac{k}{2} + 1)_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(\frac{n}{2} + \frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_n} (\frac{n}{2} + \frac{3}{2} - \frac{\gamma}{2})_{i_{n-1}} (\frac{n}{2} + 1)_{i_{n-1}}}{(\frac{n}{2} + \frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2})_{i_{n-1}} (\frac{n}{2} + \frac{3}{2} - \frac{\gamma}{2})_{i_n} (\frac{n}{2} + 1)_{i_n}} \eta^{i_n} \Big\} x^n \Big\} \quad (4.2.17)
\end{aligned}$$

It is required that $\gamma \neq 0, -1, -2, \dots$ for the first kind of independent solution of the CHF for all cases. Because if it does not, its solution will be divergent. And it is required that $\gamma \neq 2, 3, 4, \dots$ for the second kind of independent solution of the CHF for all cases.

4.2.2 Integral formalism

The CHE could not be constructed in the definite or contour integral form of any well-known simple function because of three different coefficients in its power series expansion. The three term recurrence relation in power series of the CHE creates mathematical difficulty to be analyzed it into direct or contour integral formalism.

Instead in earlier literature the integral relations of the CHE were constructed by using Fredholm integral equations; such integral relationships express one analytic solution in terms of another analytic solution such as a confluent hypergeometric function with a

branch point at zero. [2] There are many other forms of integral relations in the CHE. [14, 15, 16, 17, 18, 19]

Now I consider the combined definite and contour integral forms of the CHF by using 3TRF. Expressing the CHF in integral forms resulting in a precise and simplified transformation of the CHF to other well-known special functions such as confluent hypergeometric function.

Polynomial of type 1

Let's investigate the integral representation for polynomial of type 1. There is a generalized hypergeometric function which is

$$\begin{aligned} I_l &= \sum_{i_l=i_{l-1}}^{\alpha_l} \frac{(-\alpha_l)_{i_l} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}}}{(-\alpha_l)_{i_{l-1}} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_l} \eta^{i_l} \\ &= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2}, j+1) B(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2}, j+1) (i_{l-1} - \alpha_l)_j}{(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})^{-1} (i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})^{-1} (1)_j j!} \eta^j \end{aligned} \quad (4.2.18)$$

By using integral form of beta function,

$$B\left(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2}, j+1\right) = \int_0^1 dt_l t_l^{i_{l-1} + \frac{l}{2} - 1 + \frac{\lambda}{2}} (1 - t_l)^j \quad (4.2.19a)$$

$$B\left(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2}, j+1\right) = \int_0^1 du_l u_l^{i_{l-1} + \frac{l}{2} - \frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2}} (1 - u_l)^j \quad (4.2.19b)$$

Substitute (4.2.19a) and (4.2.19b) into (4.2.18). And divide $(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})$ into I_l .

$$\begin{aligned} K_l &= \frac{1}{(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \sum_{i_l=i_{l-1}}^{\alpha_l} \frac{(-\alpha_l)_{i_l} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}}}{(-\alpha_l)_{i_{l-1}} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_l}} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{\frac{l}{2} - 1 + \frac{\lambda}{2}} \int_0^1 du_l u_l^{\frac{l}{2} - \frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2}} (\eta t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(i_{l-1} - \alpha_l)_j}{(1)_j j!} [\eta(1 - t_l)(1 - u_l)]^j \end{aligned}$$

Confluent hypergeometric polynomial of the first kind is defined by

$$F_{\alpha_0}(\gamma; z) = \frac{\Gamma(\alpha_0 + \gamma)}{\Gamma(\gamma)} \sum_{j=0}^{\alpha_0} \frac{(-\alpha_0)_j}{(\gamma)_j j!} z^j = \frac{\Gamma(\alpha_0 + 1)}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{zv_l}{(1-v_l)}\right)}{v_l^{\alpha_0+1} (1-v_l)^\gamma} \quad (4.2.20)$$

replaced α_0 , γ and z by $\alpha_l - i_{l-1}$, 1 and $\eta(1 - t_l)(1 - u_l)$ in (4.2.20)

$$\sum_{j=0}^{\infty} \frac{(i_{l-1} - \alpha_l)_j}{(1)_j j!} [\eta(1 - t_l)(1 - u_l)]^j = \frac{1}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{v_l}{(1-v_l)}\eta(1 - t_l)(1 - u_l)\right)}{v_l^{\alpha_l - i_{l-1} + 1}(1 - v_l)} \quad (4.2.21)$$

Substitute (4.2.21) into K_l .

$$\begin{aligned} K_l &= \frac{1}{(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \sum_{i_l=i_{l-1}}^{\alpha_l} \frac{(-\alpha_l)_{i_l}(1 + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}}(\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}}}{(-\alpha_l)_{i_{l-1}}(1 + \frac{l}{2} + \frac{\lambda}{2})_{i_l}(\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_l}} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{\frac{l}{2}-1+\frac{\lambda}{2}} \int_0^1 du_l u_l^{\frac{l}{2}-\frac{3}{2}+\frac{\gamma}{2}+\frac{\lambda}{2}} \frac{1}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{v_l}{(1-v_l)}\eta(1 - t_l)(1 - u_l)\right)}{v_l^{\alpha_l+1}(1 - v_l)} (\eta t_l u_l v_l)^{i_{l-1}} \quad (4.2.22) \end{aligned}$$

Substitute (4.2.22) into (4.2.11) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(x)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(x)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(x)$, etc.⁶

Theorem 4.2.5 *The general representation in the form of integral of the CHP of type 1 about $x = 0$ is given by*

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} y_n(x) \\ &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\ &\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2+\lambda)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-3+\gamma+\lambda)} \right. \right. \\ &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})\right)}{v_{n-k}^{\alpha_{n-k}+1}(1 - v_{n-k})} \\ &\quad \times \left(w_{n-k,n}^{-\frac{1}{2}(n-k-1+\lambda)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda+n-k-2)} - \frac{q}{4} \right) \Big\} \\ &\quad \times \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,n}^{i_0} \Big\} x^n \Big\} \quad (4.2.23) \end{aligned}$$

⁶ $y_1(x)$ means the sub-power series in (4.2.11) contains one term of $A'_n s$, $y_2(x)$ means the sub-power series in (4.2.11) contains two terms of $A'_n s$, $y_3(x)$ means the sub-power series in (4.2.11) contains three terms of $A'_n s$, etc.

where

$$w_{a,b} = \begin{cases} \eta \prod_{l=a}^b t_l u_l v_l & \text{where } a \leq b \\ \eta & \text{only if } a > b \end{cases}$$

In the above, the first sub-integral form contains one term of A'_n s, the second one contains two terms of A_n 's, the third one contains three terms of A_n 's, etc.

Proof According to (4.2.11),

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \quad (4.2.24)$$

In the above, sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of the CHP of type 1 using 3TRF about $x = 0$ are

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \quad (4.2.25a)$$

$$\begin{aligned} y_1(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\ & \times \left. \sum_{i_1=i_0}^{\alpha_1} \frac{(-\alpha_1)_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(-\alpha_1)_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \eta^{i_1} \right\} x \end{aligned} \quad (4.2.25b)$$

$$\begin{aligned} y_2(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\ & \times \sum_{i_1=i_0}^{\alpha_1} \frac{(i_1 + \frac{1}{2} + \frac{\lambda}{2})(i_1 + \frac{1}{2}(-\beta + \gamma + \delta + \lambda)) - \frac{q}{4}}{(i_1 + 1 + \frac{\lambda}{2})(i_1 + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_1)_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(-\alpha_1)_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \\ & \times \left. \sum_{i_2=i_1}^{\alpha_2} \frac{(-\alpha_2)_{i_2} (2 + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}}{(-\alpha_2)_{i_1} (2 + \frac{\lambda}{2})_{i_2} (\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_2}} \eta^{i_2} \right\} x^2 \end{aligned} \quad (4.2.25c)$$

$$\begin{aligned}
y_3(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\alpha_1} \frac{(i_1 + \frac{1}{2} + \frac{\lambda}{2})(i_1 + \frac{1}{2}(-\beta + \gamma + \delta + \lambda)) - \frac{q}{4}}{(i_1 + 1 + \frac{\lambda}{2})(i_1 + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_1)_{i_1}(\frac{3}{2} + \frac{\lambda}{2})_{i_0}(1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(-\alpha_1)_{i_0}(\frac{3}{2} + \frac{\lambda}{2})_{i_1}(1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \\
& \times \sum_{i_2=i_1}^{\alpha_2} \frac{(i_2 + 1 + \frac{\lambda}{2})(i_2 + \frac{1}{2}(-\beta + \gamma + \delta + 1 + \lambda)) - \frac{q}{4}}{(i_2 + \frac{3}{2} + \frac{\lambda}{2})(i_2 + 1 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_2)_{i_2}(2 + \frac{\lambda}{2})_{i_1}(\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}}{(-\alpha_2)_{i_1}(2 + \frac{\lambda}{2})_{i_2}(\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_2}} \\
& \left. \times \sum_{i_3=i_2}^{\alpha_3} \frac{(-\alpha_3)_{i_3}(\frac{5}{2} + \frac{\lambda}{2})_{i_2}(2 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_2}}{(-\alpha_3)_{i_2}(\frac{5}{2} + \frac{\lambda}{2})_{i_3}(2 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_3}} \eta^{i_3} \right\} x^3 \quad (4.2.25d)
\end{aligned}$$

Put $l = 1$ in (4.2.22). Take the new (4.2.22) into (4.2.25b).

$$\begin{aligned}
y_1(x) = & \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}(1-t_1)(1-u_1)\eta\right)}{v_1^{\alpha_1+1}(1-v_1)} \\
& \times \left(w_{1,1}^{-\frac{\lambda}{2}} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,1}^{i_0} \right\} x \quad (4.2.26)
\end{aligned}$$

where

$$w_{1,1} = \eta \prod_{l=1}^1 t_l u_l v_l$$

Put $l = 2$ in (4.2.22). Take the new (4.2.22) into (4.2.25c).

$$\begin{aligned}
y_2(x) = & c_0 x^\lambda \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}(1-t_2)(1-u_2)\eta\right)}{v_2^{\alpha_2+1}(1-v_2)} \\
& \times \left(w_{2,2}^{-\frac{1}{2}(1+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
& \times \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\alpha_1} \frac{(-\alpha_1)_{i_1}(\frac{3}{2} + \frac{\lambda}{2})_{i_0}(1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(-\alpha_1)_{i_0}(\frac{3}{2} + \frac{\lambda}{2})_{i_1}(1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} w_{2,2}^{i_1} \left. \right\} x^2 \quad (4.2.27)
\end{aligned}$$

where

$$w_{2,2} = \eta \prod_{l=2}^2 t_l u_l v_l$$

Put $l = 1$ and $\eta = w_{2,2}$ in (4.2.22). Take the new (4.2.22) into (4.2.27).

$$\begin{aligned}
y_2(x) = & \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}(1-t_2)(1-u_2)\eta\right)}{v_2^{\alpha_2+1}(1-v_2)} \\
& \times \left(w_{2,2}^{-\frac{1}{2}(1+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
& \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}(1-t_1)(1-u_1)w_{2,2}\right)}{v_1^{\alpha_1+1}(1-v_1)} \\
& \times \left(w_{1,2}^{-\frac{\lambda}{2}} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{1}{2}+\frac{\gamma}{2}+\frac{\lambda}{2})_{i_0}} w_{1,2}^{i_0} \right\} x^2 \tag{4.2.28}
\end{aligned}$$

where

$$w_{1,2} = \eta \prod_{l=1}^2 t_l u_l v_l$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) = & \int_0^1 dt_3 t_3^{\frac{1}{2}(1+\lambda)} \int_0^1 du_3 u_3^{\frac{1}{2}(\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_3 \frac{\exp\left(-\frac{v_3}{(1-v_3)}(1-t_3)(1-u_3)\eta\right)}{v_3^{\alpha_3+1}(1-v_3)} \\
& \times \left(w_{3,3}^{-\frac{1}{2}(2+\lambda)} (w_{3,3} \partial_{w_{3,3}}) w_{3,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{3,3} \partial_{w_{3,3}}) w_{3,3}^{\frac{1}{2}(-\beta+\gamma+\delta+1+\lambda)} - \frac{q}{4} \right) \\
& \times \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}(1-t_2)(1-u_2)w_{3,3}\right)}{v_2^{\alpha_2+1}(1-v_2)} \\
& \times \left(w_{2,3}^{-\frac{1}{2}(1+\lambda)} (w_{2,3} \partial_{w_{2,3}}) w_{2,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,3} \partial_{w_{2,3}}) w_{2,3}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
& \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}(1-t_1)(1-u_1)w_{2,3}\right)}{v_1^{\alpha_1+1}(1-v_1)} \\
& \times \left(w_{1,3}^{-\frac{\lambda}{2}} (w_{1,3} \partial_{w_{1,3}}) w_{1,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,3} \partial_{w_{1,3}}) w_{1,3}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{1}{2}+\frac{\gamma}{2}+\frac{\lambda}{2})_{i_0}} w_{1,3}^{i_0} \right\} x^3 \tag{4.2.29}
\end{aligned}$$

where

$$w_{3,3} = \eta \prod_{l=3}^3 t_l u_l v_l \quad w_{2,3} = \eta \prod_{l=2}^3 t_l u_l v_l \quad w_{1,3} = \eta \prod_{l=1}^3 t_l u_l v_l$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Substitute (4.2.25a), (4.2.26), (4.2.28), (4.2.29) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (4.2.24). ■

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ in (4.2.23).

Remark 4.2.6 *The integral representation of the CHE of the first kind for polynomial of type 1 about $x = 0$ as $\alpha = -(2\alpha_j + j)$ where $j, \alpha_j \in \mathbb{N}_0$ is*

$$\begin{aligned} y(x) &= H_c^{(a)} F_{\alpha_j} \left(\alpha = -(2\alpha_j + j), \beta, \gamma, \delta, q; \eta = \frac{1}{2}\beta x^2 \right) \\ &= {}_1F_1 \left(-\alpha_0; \frac{1}{2} + \frac{\gamma}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-3+\gamma)} \right. \right. \\ &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp \left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n} (1-t_{n-k})(1-u_{n-k}) \right)}{v_{n-k}^{\alpha_{n-k}+1} (1-v_{n-k})} \\ &\quad \times \left(w_{n-k,n}^{-\frac{1}{2}(n-k-1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(-\beta+\gamma+\delta+n-k-2)} - \frac{q}{4} \right) \Big\} \\ &\quad \times {}_1F_1 \left(-\alpha_0; \frac{1}{2} + \frac{\gamma}{2}; w_{1,n} \right) \Big\} x^n \end{aligned} \quad (4.2.30)$$

Remark 4.2.7 *The integral representation of the CHE of the second kind for polynomial*

of type 1 about $x = 0$ as $\alpha = -(2\alpha_j + j + 1 - \gamma)$ where $j, \alpha_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= H_c^{(a)} S_{\alpha_j} \left(\alpha = -(2\alpha_j + j + 1 - \gamma), \beta, \gamma, \delta, q; \eta = \frac{1}{2}\beta x^2 \right) \\
&= x^{1-\gamma} \left\{ {}_1F_1 \left(-\alpha_0; \frac{3}{2} - \frac{\gamma}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-1-\gamma)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-2)} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp \left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n} (1-t_{n-k})(1-u_{n-k}) \right)}{v_{n-k}^{\alpha_{n-k}+1} (1-v_{n-k})} \\
&\quad \times \left(w_{n-k,n}^{-\frac{1}{2}(n-k-\gamma)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(-\beta+\delta+n-k-1)} - \frac{q}{4} \right) \Bigg\} \\
&\quad \times {}_1F_1 \left(-\alpha_0; \frac{3}{2} - \frac{\gamma}{2}; w_{1,n} \right) \Bigg\} x^n \Bigg\} \quad (4.2.31)
\end{aligned}$$

In the above, ${}_1F_1(a; b; z)$ is a Kummer function of the first kind which is defined by

$$\begin{aligned}
{}_1F_1(a; b; z) &= M(a, b, z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n n!} z^n = e^z M(b-a, b, -z) \\
&= -\frac{1}{2\pi i} \frac{\Gamma(1-a)\Gamma(b)}{\Gamma(b-a)} \oint dv_j e^{zv_j} (-v_j)^{a-1} (1-v_j)^{b-a-1} \\
&= \frac{\Gamma(a)}{2\pi i} \oint dv_j e^{v_j} v_j^{-b} \left(1 - \frac{z}{v_j} \right)^{-a} \\
&= \frac{1}{2\pi i} \frac{\Gamma(1-a)\Gamma(b)}{\Gamma(b-a)} \oint dv_j e^{-\frac{z}{1-v_j}} v_j^{a-1} (1-v_j)^{-b} \quad (4.2.32)
\end{aligned}$$

Infinite series

Let's consider the integral representation of the CHE about $x = 0$ for infinite series by applying 3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned}
M_l &= \sum_{i_l=i_{l-1}}^{\infty} \frac{\left(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2}\right)_{i_l} \left(1 + \frac{l}{2} + \frac{\lambda}{2}\right)_{i_{l-1}} \left(\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2}\right)_{i_{l-1}}}{\left(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2}\right)_{i_{l-1}} \left(1 + \frac{l}{2} + \frac{\lambda}{2}\right)_{i_l} \left(\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2}\right)_l} \eta^{i_l} \\
&= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2}, j+1) B(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2}, j+1) \left(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2} + i_{l-1}\right)_j}{(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})^{-1} (i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})^{-1} (1)_j j!} \eta^j \quad (4.2.33)
\end{aligned}$$

Substitute (4.2.19a) and (4.2.19b) into (4.2.33). And divide $(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})$ into the new (4.2.33).

$$\begin{aligned} V_l &= \frac{1}{(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_l} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}}}{(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_l} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{\frac{l}{2}-1+\frac{\lambda}{2}} \int_0^1 du_l u_l^{\frac{l}{2}-\frac{3}{2}+\frac{\gamma}{2}+\frac{\lambda}{2}} (\eta t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2} + i_{l-1})_j}{(1)_j j!} [\eta(1-t_l)(1-u_l)]^j \end{aligned}$$

Replace a , b and z by $\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2} + i_{l-1}$, 1 and $\eta(1-t_j)(1-u_j)$ in (4.2.32). Take the new (4.2.32) into V_l

$$\begin{aligned} V_l &= \frac{1}{(i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})(i_{l-1} + \frac{l}{2} - \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_l} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}}}{(\frac{\alpha}{2} + \frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (1 + \frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{1}{2} + \frac{\gamma}{2} + \frac{l}{2} + \frac{\lambda}{2})_l} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{\frac{l}{2}-1+\frac{\lambda}{2}} \int_0^1 du_l u_l^{\frac{l}{2}-\frac{3}{2}+\frac{\gamma}{2}+\frac{\lambda}{2}} \frac{1}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{v_l}{(1-v_l)} \eta(1-t_l)(1-u_l)\right)}{v_l^{-\frac{1}{2}(\alpha+l+\lambda)+1} (1-v_l)} (\eta t_l u_l v_l)^{i_{l-1}} \quad (4.2.34) \end{aligned}$$

Substitute (4.2.34) into (4.2.15) where $l = 1, 2, 3, \dots$; apply V_1 into the second summation of sub-power series $y_1(x)$, apply V_2 into the third summation and V_1 into the second summation of sub-power series $y_2(x)$, apply V_3 into the forth summation, V_2 into the third summation and V_1 into the second summation of sub-power series $y_3(x)$, etc.⁷

Theorem 4.2.8 *The general expression of an integral form of the CHE for infinite*

⁷ $y_1(x)$ means the sub-power series in (4.2.15) contains one term of $A'_n s$, $y_2(x)$ means the sub-power series in (4.2.15) contains two terms of $A'_n s$, $y_3(x)$ means the sub-power series in (4.2.15) contains three terms of $A'_n s$, etc.

series about $x = 0$ using 3TRF is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2+\lambda)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-3+\gamma+\lambda)} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})\right)}{v_{n-k}^{-\frac{1}{2}(\alpha+n-k+\lambda)+1} (1-v_{n-k})} \\
&\quad \times \left(w_{n-k,n}^{-\frac{1}{2}(n-k-1+\lambda)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda+n-k-2)} - \frac{q}{4} \right) \Big\} \\
&\quad \times \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,n}^{i_0} \Big\} x^n \Big\} \tag{4.2.35}
\end{aligned}$$

In the above, the first sub-integral form contains one term of A'_n s, the second one contains two terms of A_n 's, the third one contains three terms of A_n 's, etc.

Proof In (4.2.15) sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of the CHE for infinite series using 3TRF about $x = 0$ are

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \tag{4.2.36a}$$

$$\begin{aligned}
y_1(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \eta^{i_1} \Big\} x \tag{4.2.36b}
\end{aligned}$$

$$\begin{aligned}
y_2(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + \frac{1}{2} + \frac{\lambda}{2})(i_1 + \frac{1}{2}(-\beta + \gamma + \delta + \lambda)) - \frac{q}{4}}{(i_1 + 1 + \frac{\lambda}{2})(i_1 + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \\
& \left. \times \sum_{i_2=i_1}^{\infty} \frac{(\frac{\alpha}{2} + 1 + \frac{\lambda}{2})_{i_2} (2 + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}}{(\frac{\alpha}{2} + 1 + \frac{\lambda}{2})_{i_1} (2 + \frac{\lambda}{2})_{i_2} (\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_2}} \eta^{i_2} \right\} x^2 \quad (4.2.36c)
\end{aligned}$$

$$\begin{aligned}
y_3(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(i_0 + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0}(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + \frac{1}{2} + \frac{\lambda}{2})(i_1 + \frac{1}{2}(-\beta + \gamma + \delta + \lambda)) - \frac{q}{4}}{(i_1 + 1 + \frac{\lambda}{2})(i_1 + \frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} \\
& \times \sum_{i_2=i_1}^{\infty} \frac{(i_2 + 1 + \frac{\lambda}{2})(i_1 + \frac{1}{2}(-\beta + \gamma + \delta + 1 + \lambda)) - \frac{q}{4}}{(i_2 + \frac{3}{2} + \frac{\lambda}{2})(i_2 + 1 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + 1 + \frac{\lambda}{2})_{i_2} (2 + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}}{(\frac{\alpha}{2} + 1 + \frac{\lambda}{2})_{i_1} (2 + \frac{\lambda}{2})_{i_2} (\frac{3}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_2}} \\
& \left. \times \sum_{i_3=i_2}^{\infty} \frac{(\frac{\alpha}{2} + \frac{3}{2} + \frac{\lambda}{2})_{i_3} (\frac{5}{2} + \frac{\lambda}{2})_{i_2} (2 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_2}}{(\frac{\alpha}{2} + \frac{3}{2} + \frac{\lambda}{2})_{i_2} (\frac{5}{2} + \frac{\lambda}{2})_{i_3} (2 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_3}} \eta^{i_3} \right\} x^3 \quad (4.2.36d)
\end{aligned}$$

Put $l = 1$ in (4.2.34). Take the new (4.2.34) into (4.2.36b).

$$\begin{aligned}
y_1(x) = & \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}(1-t_1)(1-u_1)\eta\right)}{v_1^{-\frac{1}{2}(\alpha+1+\lambda)+1}(1-v_1)} \\
& \times \left(w_{1,1}^{-\frac{\lambda}{2}} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,1}^{i_0} \right\} x \quad (4.2.37)
\end{aligned}$$

where

$$w_{1,1} = \eta \prod_{l=1}^1 t_l u_l v_l$$

Put $l = 2$ in (4.2.34). Take the new (4.2.34) into (4.2.36c).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}(1-t_2)(1-u_2)\eta\right)}{v_2^{-\frac{1}{2}(\alpha+2+\lambda)+1}(1-v_2)} \\
&\times \left(w_{2,2}^{-\frac{1}{2}(1+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \left\{ \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})(i_0 + \frac{1}{2}(-\beta + \gamma + \delta - 1 + \lambda)) - \frac{q}{4}}{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})(i_0 + \frac{\gamma}{2} + \frac{\lambda}{2})} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \right. \\
&\times \left. \sum_{i_1=i_0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} + \frac{\lambda}{2})_{i_0} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}}{(\frac{\alpha}{2} + \frac{1}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} + \frac{\lambda}{2})_{i_1} (1 + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_1}} w_{2,2}^{i_1} \right\} x^2 \quad (4.2.38)
\end{aligned}$$

where

$$w_{2,2} = \eta \prod_{l=2}^2 t_l u_l v_l$$

Put $l = 1$ and $\eta = w_{2,2}$ in (4.2.34). Take the new (4.2.34) into (4.2.38).

$$\begin{aligned}
y_2(x) &= \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}(1-t_2)(1-u_2)\eta\right)}{v_2^{-\frac{1}{2}(\alpha+2+\lambda)+1}(1-v_2)} \\
&\times \left(w_{2,2}^{-\frac{1}{2}(1+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}(1-t_1)(1-u_1)w_{2,2}\right)}{v_1^{-\frac{1}{2}(\alpha+1+\lambda)+1}(1-v_1)} \\
&\times \left(w_{1,2}^{-\frac{\lambda}{2}} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\frac{\alpha}{2} + \frac{\lambda}{2})_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,2}^{i_0} \right\} x^2 \quad (4.2.39)
\end{aligned}$$

where

$$w_{1,2} = \eta \prod_{l=1}^2 t_l u_l v_l$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the

integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) = & \int_0^1 dt_3 t_3^{\frac{1}{2}(1+\lambda)} \int_0^1 du_3 u_3^{\frac{1}{2}(\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_3 \frac{\exp\left(-\frac{v_3}{(1-v_3)}(1-t_3)(1-u_3)\eta\right)}{v_3^{-\frac{1}{2}(\alpha+3+\lambda)+1}(1-v_3)} \\
& \times \left(w_{3,3}^{-\frac{1}{2}(2+\lambda)} (w_{3,3} \partial_{w_{3,3}}) w_{3,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{3,3} \partial_{w_{3,3}}) w_{3,3}^{\frac{1}{2}(-\beta+\gamma+\delta+1+\lambda)} - \frac{q}{4} \right) \\
& \times \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}(1-t_2)(1-u_2)w_{3,3}\right)}{v_2^{-\frac{1}{2}(\alpha+2+\lambda)+1}(1-v_2)} \\
& \times \left(w_{2,3}^{-\frac{1}{2}(1+\lambda)} (w_{2,3} \partial_{w_{2,3}}) w_{2,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,3} \partial_{w_{2,3}}) w_{2,3}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
& \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}(1-t_1)(1-u_1)w_{2,3}\right)}{v_1^{-\frac{1}{2}(\alpha+1+\lambda)+1}(1-v_1)} \\
& \times \left(w_{1,3}^{-\frac{\lambda}{2}} (w_{1,3} \partial_{w_{1,3}}) w_{1,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,3} \partial_{w_{1,3}}) w_{1,3}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{\left(\frac{\alpha}{2} + \frac{\lambda}{2}\right)_{i_0}}{\left(1 + \frac{\lambda}{2}\right)_{i_0} \left(\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2}\right)_{i_0}} w_{1,3}^{i_0} \right\} x^3 \tag{4.2.40}
\end{aligned}$$

where

$$w_{3,3} = \eta \prod_{l=3}^3 t_l u_l v_l \quad w_{2,3} = \eta \prod_{l=2}^3 t_l u_l v_l \quad w_{1,3} = \eta \prod_{l=1}^3 t_l u_l v_l$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute (4.2.36a), (4.2.37), (4.2.39), (4.2.40) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (4.2.15), we obtain (4.2.35).⁸ ■

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ in (4.2.35).

Remark 4.2.9 *The integral representation of the CHE of the first kind for infinite series*

⁸Or replace the finite summation with an interval $[0, \alpha_0]$ by infinite summation with an interval $[0, \infty]$ in (4.2.23). Replace α_0 and α_{n-k} by $-\frac{1}{2}(\alpha + \lambda)$ and $-\frac{1}{2}(\alpha + n - k + \lambda)$ into the new (4.2.23). Its solution is also equivalent to (4.2.35)

about $x = 0$ using 3TRF is

$$\begin{aligned}
y(x) &= H_c^{(a)} F \left(\alpha, \beta, \gamma, \delta, q; \eta = \frac{1}{2} \beta x^2 \right) \\
&= {}_1F_1 \left(\frac{\alpha}{2}; \frac{1}{2} + \frac{\gamma}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-3+\gamma)} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp \left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n} (1-t_{n-k})(1-u_{n-k}) \right)}{v_{n-k}^{-\frac{1}{2}(\alpha+n-k-2)} (1-v_{n-k})} \\
&\quad \times \left(w_{n-k,n}^{-\frac{1}{2}(n-k-1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(-\beta+\gamma+\delta+n-k-2)} - \frac{q}{4} \right) \Big\} \\
&\quad \times {}_1F_1 \left(\frac{\alpha}{2}; \frac{1}{2} + \frac{\gamma}{2}; w_{1,n} \right) \Big\} x^n \tag{4.2.41}
\end{aligned}$$

Remark 4.2.10 The integral representation of the CHE of the second kind for infinite series about $x = 0$ using 3TRF is

$$\begin{aligned}
y(x) &= H_c^{(a)} S \left(\alpha, \beta, \gamma, \delta, q; \eta = \frac{1}{2} \beta x^2 \right) \\
&= x^{1-\gamma} \left\{ {}_1F_1 \left(\frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2}; \frac{3}{2} - \frac{\gamma}{2}; \eta \right) \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-1-\gamma)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-2)} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp \left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n} (1-t_{n-k})(1-u_{n-k}) \right)}{v_{n-k}^{-\frac{1}{2}(\alpha+n-k-1-\gamma)} (1-v_{n-k})} \\
&\quad \times \left(w_{n-k,n}^{-\frac{1}{2}(n-k-\gamma)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(-\beta+\delta+n-k-1)} - \frac{q}{4} \right) \Big\} \\
&\quad \times {}_1F_1 \left(\frac{\alpha}{2} + \frac{1}{2} - \frac{\gamma}{2}; \frac{3}{2} - \frac{\gamma}{2}; w_{1,n} \right) \Big\} x^n \Big\} \tag{4.2.42}
\end{aligned}$$

4.2.3 Generating function for the CHP of type 1

I consider the generating function for the CHP of type 1. Since the generating function for the CHP is derived, we might be possible to construct orthogonal relations of the

CHP. In the physical point of view we might be possible to obtain the normalized constant for the wave function in modern physics (especially black hole problems) recursion relation and its expectation value of any physical quantities from the generating function for the CHP.⁹ For the case of hydrogen-like atoms, the normalized wave function is derived from the generating function for associated Laguerre polynomial. And the expectation value of physical quantities such as position and momentum is constructed by applying the recursive relation of associated Laguerre polynomial.

Let's investigate the generating function for the first and second kinds of the CHP about $x = 0$ as $B'_n s$ term terminated at certain eigenvalue.

Definition 4.2.11 *I define that*

$$\left\{ \begin{array}{l} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{where } a > b \\ s_a & \text{only if } a = b \end{cases} \\ \tilde{w}_{a,b} = \eta s_{a,\infty} \prod_{l=a}^b t_l v_l \end{array} \right. \quad (4.2.43)$$

where

$$a, b \in \mathbb{N}_0$$

And we have

$$\sum_{\alpha_i=\alpha_j}^{\infty} s_i^{\alpha_i} = \frac{s_i^{\alpha_j}}{(1-s_i)} \quad (4.2.44)$$

Acting the summation operator $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\}$ on (4.2.23) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (4.2.43) and (4.2.44),

Theorem 4.2.12 *The general expression of the generating function for the CHP of type*

⁹In this chapter I consider the generating functions for type 1 CHP. In the next chapter and future paper, I will construct the generating functions for type 2 and 3 CHP's. For the type 1 polynomial, I treat β, γ, δ and q as free variables and α as a fixed value. For the type 2 polynomial, I treat α, β, γ and δ as free variables and q as a fixed value. For the type 3 polynomial, I treat β, γ and δ as free variables and α and q as fixed values.

1 about $x = 0$ is given by

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y(x) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \Upsilon(\gamma', \lambda; s_{0,\infty}; \eta) \\
&+ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \exp\left(-\frac{s_{1,\infty}}{(1 - s_{1,\infty})} \eta(1 - t_1)(1 - u_1)\right) \\
&\times \left(\tilde{w}_{1,1}^{-\frac{\lambda}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \Upsilon(\gamma', \lambda; s_0; \tilde{w}_{1,1}) x \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{\frac{1}{2}(n-2+\lambda)} \int_0^1 du_n u_n^{\frac{1}{2}(n-3+\gamma+\lambda)} \exp\left(-\frac{s_{n,\infty}}{(1 - s_{n,\infty})} \eta(1 - t_n)(1 - u_n)\right) \right. \\
&\times \left(\tilde{w}_{n,n}^{-\frac{1}{2}(n-1+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(-\beta+\gamma+\delta+n-2+\lambda)} - \frac{q}{4} \right) \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2+\lambda)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-3+\gamma+\lambda)} \right. \\
&\times \frac{\exp\left(-\frac{s_{n-k}}{(1-s_{n-k})} \tilde{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})\right)}{(1 - s_{n-k})} \\
&\times \left. \left(\tilde{w}_{n-k,n}^{-\frac{1}{2}(n-k-1+\lambda)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(-\beta+\gamma+\delta+n-k-2+\lambda)} - \frac{q}{4} \right) \right\} \\
&\times \Upsilon(\gamma', \lambda; s_0; \tilde{w}_{1,n}) \Big\} x^n
\end{aligned} \tag{4.2.45}$$

where

$$\begin{cases} \Upsilon(\gamma', \lambda; s_{0,\infty}; \eta) = \sum_{\alpha_0=0}^{\infty} \frac{s_{0,\infty}^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right) \\ \Upsilon(\gamma', \lambda; s_0; \tilde{w}_{1,1}) = \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \\ \Upsilon(\gamma', \lambda; s_0; \tilde{w}_{1,n}) = \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \tilde{w}_{1,n}^{i_0} \right) \end{cases}$$

Proof Acting the summation operator $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\}$ on the form of integral of the CHP of type 1 $y(x)$,

$$\begin{aligned} & \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y(x) \\ &= \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} \{y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots\} \end{aligned} \quad (4.2.46)$$

Acting the summation operator $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\}$ on (4.2.25a) by applying (4.2.44),

$$\begin{aligned} & \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_0(x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \left(c_0 x^\lambda \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right) \end{aligned} \quad (4.2.47)$$

Acting the summation operator $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\}$ on (4.2.26),

$$\begin{aligned} & \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} \eta(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\ & \times \sum_{\alpha_1=\alpha_0}^{\infty} \left(\frac{s_{1,\infty}}{v_1} \right)^{\alpha_1} \left(w_{1,1}^{-\frac{\lambda}{2}} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\ & \times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,1}^{i_0} \right) x \end{aligned} \quad (4.2.48)$$

Replace α_i , α_j and s_i by α_1 , α_0 and $\frac{s_{1,\infty}}{v_1}$ in (4.2.44). Take the new (4.2.44) into (4.2.48).

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_1(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}\eta(1-t_1)(1-u_1)\right)}{(1-v_1)(v_1-s_{1,\infty})} \\
&\times \left(w_{1,1}^{-\frac{\lambda}{2}} (w_{1,1}\partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,1}\partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{1}{\alpha_0!} \left(\frac{s_{0,\infty}}{v_1} \right)^{\alpha_0} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{1}{2}+\frac{\gamma}{2}+\frac{\lambda}{2})_{i_0}} w_{1,1}^{i_0} \right) x \tag{4.2.49}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $v_1 = 1$ or $s_{1,\infty}$ and $s_{1,\infty}$ is only inside the unit circle. As we compute the residue there in (4.2.49) we obtain

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_1(x) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \exp\left(-\frac{s_{1,\infty}}{(1-s_{1,\infty})}\eta(1-t_1)(1-u_1)\right) \\
&\times \left(\tilde{w}_{1,1}^{-\frac{\lambda}{2}} (\tilde{w}_{1,1}\partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{1,1}\partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{1}{2}+\frac{\gamma}{2}+\frac{\lambda}{2})_{i_0}} \tilde{w}_{1,1}^{i_0} \right) x \tag{4.2.50}
\end{aligned}$$

where $\tilde{w}_{1,1} = \eta s_{1,\infty} \prod_{l=1}^1 t_l u_l$

Acting the summation operator $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\}$ on (4.2.28),

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{v_2(1-v_2)} \\
&\times \sum_{\alpha_2=\alpha_1}^{\infty} \left(\frac{s_{2,\infty}}{v_2}\right)^{\alpha_2} \left(w_{2,2}^{-\frac{1}{2}(1+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}w_{2,2}(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\
&\times \sum_{\alpha_1=\alpha_0}^{\infty} \left(\frac{s_{1,\infty}}{v_1}\right)^{\alpha_1} \left(w_{1,2}^{-\frac{\lambda}{2}} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,2}^{i_0} \right) x^2 \tag{4.2.51}
\end{aligned}$$

Replace α_i , α_j and s_i by α_2 , α_1 and $\frac{s_{2,\infty}}{v_2}$ in (4.2.44). Take the new (4.2.44) into (4.2.51).

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{(1-v_2)(v_2 - s_{2,\infty})} \\
&\times \left(w_{2,2}^{-\frac{1}{2}(1+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}w_{2,2}(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\
&\times \sum_{\alpha_1=\alpha_0}^{\infty} \left(\frac{s_{1,\infty}}{v_1 v_2}\right)^{\alpha_1} \left(w_{1,2}^{-\frac{\lambda}{2}} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1 + \frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} w_{1,2}^{i_0} \right) x^2 \tag{4.2.52}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $v_2 = 1$ or $s_{2,\infty}$ and $s_{2,\infty}$ is only inside the unit circle. As we compute the residue there in (4.2.52) we obtain

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_2(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \exp\left(-\frac{s_{2,\infty}}{(1-s_{2,\infty})} \eta(1-t_2)(1-u_2)\right) \\
&\times \left(\tilde{w}_{2,2}^{-\frac{1}{2}(1+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} \tilde{w}_{2,2}(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\
&\times \sum_{\alpha_1=\alpha_0}^{\infty} \left(\frac{s_1}{v_1} \right)^{\alpha_1} \left(\ddot{w}_{1,2}^{-\frac{\lambda}{2}} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \ddot{w}_{1,2}^{i_0} \right) x^2 \quad (4.2.53)
\end{aligned}$$

$$\text{where } \tilde{w}_{2,2} = \eta s_{2,\infty} \prod_{l=2}^2 t_l u_l \text{ and } \ddot{w}_{1,2} = \eta v_1 s_{2,\infty} \prod_{l=1}^2 t_l u_l$$

Replace α_i , α_j and s_i by α_1 , α_0 and $\frac{s_1}{v_1}$ in (4.2.44). Take the new (4.2.44) into (4.2.53).

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_2(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \exp\left(-\frac{s_{2,\infty}}{(1-s_{2,\infty})} \eta(1-t_2)(1-u_2)\right) \\
&\times \left(\tilde{w}_{2,2}^{-\frac{1}{2}(1+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} \tilde{w}_{2,2}(1-t_1)(1-u_1)\right)}{(1-v_1)(v_1-s_1)} \\
&\times \left(\ddot{w}_{1,2}^{-\frac{\lambda}{2}} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{1}{\alpha_0!} \left(\frac{s_{0,1}}{v_1} \right)^{\alpha_0} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \ddot{w}_{1,2}^{i_0} \right) x^2 \quad (4.2.54)
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $v_1 = 1$ or s_1 and s_1 is only inside the unit circle. As we compute the residue there in (4.2.54) we obtain

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_2(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \exp\left(-\frac{s_{2,\infty}}{(1-s_{2,\infty})} \eta(1-t_2)(1-u_2)\right) \\
&\times \left(\tilde{w}_{2,2}^{-\frac{1}{2}(1+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{\exp\left(-\frac{s_1}{(1-s_1)} \tilde{w}_{2,2}(1-t_1)(1-u_1)\right)}{(1-s_1)} \\
&\times \left(\tilde{w}_{1,2}^{-\frac{\lambda}{2}} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^{\lambda} \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{1}{2}+\frac{\gamma}{2}+\frac{\lambda}{2})_{i_0}} \tilde{w}_{1,2}^{i_0} \right) x^2 \tag{4.2.55}
\end{aligned}$$

where $\tilde{w}_{1,2} = \eta s_{1,\infty} \prod_{l=1}^2 t_l u_l$

Acting the summation operator $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\}$ on (4.2.29),

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_3(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{\frac{1}{2}(1+\lambda)} \int_0^1 du_3 u_3^{\frac{1}{2}(\gamma+\lambda)} \exp\left(-\frac{s_{3,\infty}}{(1-s_{3,\infty})} \eta(1-t_3)(1-u_3)\right) \\
&\times \left(\tilde{w}_{3,3}^{-\frac{1}{2}(2+\lambda)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{\frac{1}{2}(-\beta+\gamma+\delta+1+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_2 t_2^{\frac{\lambda}{2}} \int_0^1 du_2 u_2^{\frac{1}{2}(-1+\gamma+\lambda)} \frac{\exp\left(-\frac{s_2}{(1-s_2)} \tilde{w}_{3,3}(1-t_2)(1-u_2)\right)}{(1-s_2)} \\
&\times \left(\tilde{w}_{2,3}^{-\frac{1}{2}(-1+\lambda)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{\frac{1}{2}(-\beta+\gamma+\delta+\lambda)} - \frac{q}{4} \right) \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1+\lambda)} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma+\lambda)} \frac{\exp\left(-\frac{s_1}{(1-s_1)} \tilde{w}_{2,3}(1-t_1)(1-u_1)\right)}{(1-s_1)} \\
&\times \left(\tilde{w}_{1,3}^{-\frac{\lambda}{2}} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\frac{1}{2}(-\beta+\gamma+\delta-1+\lambda)} - \frac{q}{4} \right) \\
&\times \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \left(c_0 x^\lambda \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{1}{2} + \frac{\gamma}{2} + \frac{\lambda}{2})_{i_0}} \tilde{w}_{1,3}^{i_0} \right) x^3 \tag{4.2.56}
\end{aligned}$$

$$\text{where } \tilde{w}_{3,3} = \eta s_{3,\infty} \prod_{l=3}^3 t_l u_l, \quad \tilde{w}_{2,3} = \eta s_{2,\infty} \prod_{l=2}^3 t_l u_l \quad \text{and} \quad \tilde{w}_{1,3} = \eta s_{1,\infty} \prod_{l=1}^3 t_l u_l$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$

terms where $m > 3$, I obtain every $\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} y_m(x)$ terms.

Substitute (4.2.47), (4.2.50), (4.2.55), (4.2.56) and including all

$$\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{(q_0)!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(x) \text{ terms where } m > 3 \text{ into (4.2.46).} \quad \blacksquare$$

Remark 4.2.13 *The generating function for the CHP of type 1 of the first kind about*

$x = 0$ as $\alpha = -(2\alpha_j + j)$ where $j, \alpha_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{1}{2} + \frac{\gamma}{2})}{\Gamma(\frac{1}{2} + \frac{\gamma}{2})} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} H_c^{(a)} F_{\alpha_j} \left(\alpha = -(2\alpha_j + j), \beta, \gamma, \delta, q; \eta = \frac{1}{2}\beta x^2 \right) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \\
&+ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{-\frac{1}{2}} \int_0^1 du_1 u_1^{\frac{1}{2}(-2+\gamma)} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
&\times \left((\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\beta+\gamma+\delta-1)} - \frac{q}{4} \right) \mathbf{A}(s_0; \tilde{w}_{1,1}) x \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{\frac{1}{2}(n-2)} \int_0^1 du_n u_n^{\frac{1}{2}(n-3+\gamma)} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \left(\tilde{w}_{n,n}^{-\frac{1}{2}(n-1)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(-\beta+\gamma+\delta+n-2)} - \frac{q}{4} \right) \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-3+\gamma)} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
&\times \left. \left(\tilde{w}_{n-k,n}^{-\frac{1}{2}(n-k-1)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(-\beta+\gamma+\delta+n-k-2)} - \frac{q}{4} \right) \right\} \\
&\times \mathbf{A}(s_0; \tilde{w}_{1,n}) \Big\} x^n \tag{4.2.57}
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \exp\left(-\frac{s_{1,\infty}}{(1-s_{1,\infty})}\eta(1-t_1)(1-u_1)\right) \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \exp\left(-\frac{s_{n,\infty}}{(1-s_{n,\infty})}\eta(1-t_n)(1-u_n)\right) \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) = \frac{\exp\left(-\frac{s_{n-k}}{(1-s_{n-k})}\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})\right)}{(1-s_{n-k})} \end{cases}$$

and

$$\begin{cases} \mathbf{A}(s_{0,\infty}; \eta) = (1 - s_{0,\infty})^{-\frac{1}{2}(1+\gamma)} \exp\left(-\frac{\eta s_{0,\infty}}{(1-s_{0,\infty})}\right) \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = (1 - s_0)^{-\frac{1}{2}(1+\gamma)} \exp\left(-\frac{\tilde{w}_{1,1}s_0}{(1-s_0)}\right) \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = (1 - s_0)^{-\frac{1}{2}(1+\gamma)} \exp\left(-\frac{\tilde{w}_{1,n}s_0}{(1-s_0)}\right) \end{cases}$$

Proof The generating function for Confluent hypergeometric (Kummer's) polynomial of the first kind is given by

$$\sum_{\alpha_0=0}^{\infty} \frac{t^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \gamma')}{\Gamma(\gamma')} {}_1F_1(-\alpha_0; \gamma'; z) = \frac{\exp\left(-\frac{zt}{(1-t)}\right)}{(1-t)^{\gamma'}} \quad (4.2.58)$$

Replace t , γ' and z by $s_{0,\infty}$, $\frac{1}{2} + \frac{\gamma}{2}$ and η in (4.2.58).

$$\sum_{\alpha_0=0}^{\infty} \frac{s_{0,\infty}^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{1}{2} + \frac{\gamma}{2})}{\Gamma(\frac{1}{2} + \frac{\gamma}{2})} {}_1F_1\left(-\alpha_0; \frac{1}{2} + \frac{\gamma}{2}; \eta\right) = (1 - s_{0,\infty})^{-\frac{1}{2}(1+\gamma)} \exp\left(-\frac{\eta s_{0,\infty}}{(1 - s_{0,\infty})}\right) \quad (4.2.59)$$

Replace t , γ' and z by s_0 , $\frac{1}{2} + \frac{\gamma}{2}$ and $\tilde{w}_{1,1}$ in (4.2.58).

$$\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{1}{2} + \frac{\gamma}{2})}{\Gamma(\frac{1}{2} + \frac{\gamma}{2})} {}_1F_1\left(-\alpha_0; \frac{1}{2} + \frac{\gamma}{2}; \tilde{w}_{1,1}\right) = (1 - s_0)^{-\frac{1}{2}(1+\gamma)} \exp\left(-\frac{\tilde{w}_{1,1}s_0}{(1 - s_0)}\right) \quad (4.2.60)$$

Replace t , γ' and z by s_0 , $\frac{1}{2} + \frac{\gamma}{2}$ and $\tilde{w}_{1,n}$ in (4.2.58).

$$\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{1}{2} + \frac{\gamma}{2})}{\Gamma(\frac{1}{2} + \frac{\gamma}{2})} {}_1F_1\left(-\alpha_0; \frac{1}{2} + \frac{\gamma}{2}; \tilde{w}_{1,n}\right) = (1 - s_0)^{-\frac{1}{2}(1+\gamma)} \exp\left(-\frac{\tilde{w}_{1,n}s_0}{(1 - s_0)}\right) \quad (4.2.61)$$

Put $c_0 = 1$, $\lambda = 0$ and $\gamma' = \frac{1}{2} + \frac{\gamma}{2}$ in (4.2.45). Substitute (4.2.59), (4.2.60) and (4.2.61) into the new (4.2.45). ■

Remark 4.2.14 The generating function for the CHP of type 1 of the second kind about

$x = 0$ as $\alpha = -(2\alpha_j + j + 1 - \gamma)$ where $j, \alpha_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{3}{2} - \frac{\gamma}{2})}{\Gamma(\frac{3}{2} - \frac{\gamma}{2})} \prod_{n=1}^{\infty} \left\{ \sum_{\alpha_n=\alpha_{n-1}}^{\infty} s_n^{\alpha_n} \right\} H_c^{(a)} S_{\alpha_j} \left(\alpha = -(2\alpha_j + j + 1 - \gamma), \beta, \gamma, \delta, q; \eta = \frac{1}{2}\beta x^2 \right) \\
&= x^{1-\gamma} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{B}(s_{0,\infty}; \eta) \right. \\
&+ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{-\frac{\gamma}{2}} \int_0^1 du_1 u_1^{-\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
&\times \left(\tilde{w}_{1,1}^{-\frac{1}{2}(1-\gamma)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\beta+\delta)} - \frac{q}{4} \right) \mathbf{B}(s_0; \tilde{w}_{1,1}) \quad x \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{\frac{1}{2}(n-1-\gamma)} \int_0^1 du_n u_n^{\frac{1}{2}(n-2)} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \left(\tilde{w}_{n,n}^{-\frac{1}{2}(n-\gamma)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(-\beta+\delta+n-1)} - \frac{q}{4} \right) \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-1-\gamma)} \int_0^1 du_{n-k} u_{n-k}^{\frac{1}{2}(n-k-2)} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
&\times \left. \left(\tilde{w}_{n-k,n}^{-\frac{1}{2}(n-k-\gamma)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(\beta-\gamma-\delta+1)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(-\beta+\delta+n-k-1)} - \frac{q}{4} \right) \right\} \\
&\times \mathbf{B}(s_0; \tilde{w}_{1,n}) \left. \right\} x^n \Bigg\} \tag{4.2.62}
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \exp\left(-\frac{s_{1,\infty}}{(1-s_{1,\infty})} \eta (1-t_1)(1-u_1)\right) \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \exp\left(-\frac{s_{n,\infty}}{(1-s_{n,\infty})} \eta (1-t_n)(1-u_n)\right) \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) = \frac{\exp\left(-\frac{s_{n-k}}{(1-s_{n-k})} \tilde{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})\right)}{(1-s_{n-k})} \end{cases}$$

and

$$\begin{cases} \mathbf{B}(s_{0,\infty}; \eta) = (1-s_{0,\infty})^{-\frac{1}{2}(3-\gamma)} \exp\left(-\frac{\eta s_{0,\infty}}{(1-s_{0,\infty})}\right) \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = (1-s_0)^{-\frac{1}{2}(3-\gamma)} \exp\left(-\frac{\tilde{w}_{1,1} s_0}{(1-s_0)}\right) \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = (1-s_0)^{-\frac{1}{2}(3-\gamma)} \exp\left(-\frac{\tilde{w}_{1,n} s_0}{(1-s_0)}\right) \end{cases}$$

Proof Replace t , γ' and z by $s_{0,\infty}$, $\frac{3}{2} - \frac{\gamma}{2}$ and η in (4.2.58).

$$\sum_{\alpha_0=0}^{\infty} \frac{s_{0,\infty}^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{3}{2} - \frac{\gamma}{2})}{\Gamma(\frac{3}{2} - \frac{\gamma}{2})} {}_1F_1 \left(-\alpha_0; \frac{3}{2} - \frac{\gamma}{2}; \eta \right) = (1 - s_{0,\infty})^{-\frac{1}{2}(3-\gamma)} \exp \left(-\frac{\eta s_{0,\infty}}{(1 - s_{0,\infty})} \right) \quad (4.2.63)$$

Replace t , γ' and z by s_0 , $\frac{3}{2} - \frac{\gamma}{2}$ and $\tilde{w}_{1,1}$ in (4.2.58).

$$\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{3}{2} - \frac{\gamma}{2})}{\Gamma(\frac{3}{2} - \frac{\gamma}{2})} {}_1F_1 \left(-\alpha_0; \frac{3}{2} - \frac{\gamma}{2}; \tilde{w}_{1,1} \right) = (1 - s_0)^{-\frac{1}{2}(3-\gamma)} \exp \left(-\frac{\tilde{w}_{1,1} s_0}{(1 - s_0)} \right) \quad (4.2.64)$$

Replace t , γ' and z by s_0 , $\frac{3}{2} - \frac{\gamma}{2}$ and $\tilde{w}_{1,n}$ in (4.2.58).

$$\sum_{\alpha_0=0}^{\infty} \frac{s_0^{\alpha_0}}{\alpha_0!} \frac{\Gamma(\alpha_0 + \frac{3}{2} - \frac{\gamma}{2})}{\Gamma(\frac{3}{2} - \frac{\gamma}{2})} {}_1F_1 \left(-\alpha_0; \frac{3}{2} - \frac{\gamma}{2}; \tilde{w}_{1,n} \right) = (1 - s_0)^{-\frac{1}{2}(3-\gamma)} \exp \left(-\frac{\tilde{w}_{1,n} s_0}{(1 - s_0)} \right) \quad (4.2.65)$$

Put $c_0 = 1$, $\lambda = 1 - \gamma$ and $\gamma' = \frac{3}{2} - \frac{\gamma}{2}$ in (4.2.45). Substitute (4.2.63), (4.2.64) and (4.2.65) into the new (4.2.45). ■

4.3 Summary

In this chapter I apply the 3TRF to (1) the power series expansion in closed forms of the CHE about regular singularity $x = 0$, (2) its integral representation for infinite series and polynomial of type 1 and (3) generating functions of the CHP of type 1.

As we see the power series expansions of the CHE for infinite series and polynomial, denominators and numerators in every B_n terms arise with Pochhammer symbol: the meaning of this is that the analytic solutions of the CHF can be described as hypergeometric functions in a strict mathematical way. I show how to construct representations in closed form integrals resulting in a precise since we have power series expansions with Pochhammer symbols in numerators and denominators. We can transform the CHF into all other well-known special functions with two recursive coefficients because a ${}_1F_1$ function recurs in each of sub-integral forms of the CHF.

I show how to analyze generating functions for the CHP of type 1 from the general representation of an integral form of the CHP. We can derive orthogonal relations, recursion relations and expectation values of physical quantities from the generating functions for the CHP: the processes in order to obtain orthogonal and recursion relations

of the CHP are similar to the case of a normalized wave function for the hydrogen-like atoms.

The second independent solution of the CHF is only valid for non-integer values of $1 - \gamma$ in the domain $|x| < 1$. Its analytic solution is defined by its behavior at infinity in the domain $|x| > 1$. The local solution of the CHF about regular singularity $x = 0$ is called angular solution of the CHE, denoted as $Hc^{(a)}(\alpha, \beta, \gamma, \delta, q; x)$. And the local solution of the CHE about the singular point at infinity is called radial solution of the CHE, denoted as $Hc^{(r)}(\alpha, \beta, \gamma, \delta, q; x)$. [2] Various local solutions of the CHE can be constructed using the combination of transformed independent parameters which resembles Kummer's 24 local solutions of the hypergeometric equation. All possible local solutions of the CHE (Regge-Wheeler and Teukolsky equations) were constructed by Fiziev.[11, 12] In general the power series expansion for the CHE about the singular point at infinity is not convergent by only asymptotic. [2, 13] Putting $z = \frac{1}{x}$ into (4.1.2),

$$z^2 \frac{d^2 y}{dz^2} + z \left((2 - \gamma) - \frac{\beta}{z} + \frac{\delta}{z - 1} \right) \frac{dy}{dz} + \frac{qz - \alpha\beta}{z(z - 1)} y = 0 \quad (4.3.1)$$

Assume that its solution is

$$y(z) = \sum_{n=0}^{\infty} c_n z^{n+\lambda} \quad \text{where } \lambda = \text{indicial root} \quad (4.3.2)$$

Plug (4.3.2) into (4.3.1). We again obtain a three-term recurrence relation for the coefficients c_n :

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (4.3.3)$$

where,

$$A_n = \frac{(n + \lambda)(n + \lambda + \beta - \gamma - \delta + 1) - q}{\beta(n + 1 - \alpha + \lambda)} \quad (4.3.4a)$$

$$B_n = \frac{-(n + \lambda - 1)(n + \lambda - \gamma)}{\beta(n + \lambda + 1 - \alpha)} \quad (4.3.4b)$$

$$c_1 = A_0 c_0 \quad (4.3.4c)$$

We only have an indicial root which is $\lambda = \alpha$. Let's test for the asymptotic behavior of the function $y(z)$. As n goes to infinity,

$$A = \lim_{n \gg 1} A_n = \frac{n}{\beta} \quad (4.3.5a)$$

$$B = \lim_{n \gg 1} B_n = -\frac{n}{\beta} \quad (4.3.5b)$$

As we see (4.3.5a) and (4.3.5b), the function $y(z)$ for infinite series is divergent. Also, for polynomial of type 1, (4.3.5b) is negligible for the minimum $y(z)$ because B_n term will be

terminated at the specific eigenvalues. (4.3.5a) is only survived. And its recurrence relation is

$$c_{n+1} = \frac{n}{\beta} c_n \quad (4.3.6)$$

The local solution for (4.3.6) is divergent. By similar reason, polynomial of type 2 also will be divergent. Therefore, the local solution of the CHE about the singular point at infinity is only valid for the type 3 polynomial where A_n and B_n terms terminated at the same time. Also we need a condition $\alpha > 0$ in order to make a function $y(z)$ convergent. If it does not, the function $y(z)$ will be divergent as x goes to infinity. For the black hole problems, x is correspond to the radius.[11, 12, 13] In the quantum mechanical points of views any physical quantities should be disappeared as the distance goes to the infinity as we all know. In the next series I will construct the Frobenius solution in closed forms of the CHE about the singular point at infinity, its integral representation and its generating functions for the type 3 polynomial. I also derive the mathematical formula of the spectral parameter q for this type polynomial.

Bibliography

- [1] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806.
- [2] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische. Annalen.* **33**, 161(1889).
- [3] Ronveaux, A., “Heuns Differential Equations” Oxford University Press, (1995).
- [4] Choun, Y.S., “Analytic solution for grand confluent hypergeometric function,” arXiv:1303.0813.
- [5] Choun, Y.S., “The integral formalism and the generating function of grand confluent hypergeometric function,” arXiv:1303.0819.
- [6] Leaver, E.W., “Solutions to a generalized spheroidal wave equation: Teukolsky’s equations in general relativity, and the two-center problem in molecular quantum mechanics,” *J. Math. Phys.* **27**, 1238–1265(1986).
- [7] Decarreau, A., Maroni, P. and Robert, A., “Sur les équations confluentes de l’équation de Heun,” *Ann. Soc. Sci. Bruxelles.* **T92(III)**, 151–189 (1978).
- [8] Decarreau, A., Dumont-Lepage, M. C., Maroni, P., Robert, A. and Ronveaux, A., “Formes canoniques des équations confluentes de l’équation de Heun,” *Ann. Soc. Sci. Bruxelles.* **T92(I-II)**, 53–78 (1978).
- [9] Wilson, A. H., “A generalised spheroidal wave equation,” *Proc. Roy. Soc. London* **A118**, 61–635 (1928).
- [10] Wilson, A. H., “The ionised hydrogen molecule,” *Proc. Roy. Soc. London* **A118**, 635–647 (1928).
- [11] Fiziev, P.P., “Classes of exact solutions to Regge-Wheeler and Teukolsky equations,” arXiv:0902.1277.

- [12] Fiziev, P.P., “Classes of exact solutions to the Teukolsky master equations,” *Class. Quantum. Grav.* **27**, 135001 (2010).
- [13] Fiziev, P.P., “Exact solutions of Teukolsky master equation with continuous spectrum,” arXiv:0903.3617.
- [14] Lambe, C.G. and Ward, D.R., “Some differential equations and associated integral equations,” *Q. J. Math. (Oxford), Ser.* **13**, 107-112 (1942).
- [15] Erdélyi, A., “Integral equations for Heun functions,” *Q.J. Math. (Oxford), Ser.* **13**, 107–112 (1942).
- [16] Sleeman, B.D., “Non-linear integral equations for Heun functions,” *Proc. Edinburgh Math. Soc.*, Vol. **16**, 281–289 (1969).
- [17] Abramov, D.I. and Komarov, I.V., “Integral equations and relations for the Coulomb spheroidal functions,” *Sov. J. Teor. Mat. Phys.* Vol. **29**, 234–243 (1976).
- [18] Schmidt, D. and Wolf, G., “A method of generating integral relations by simultaneous separability of generalized Schrödinger equation,” *SIAM J. Math. Anal.* Vol. **10**, 823–838 (1979).
- [19] Kazakov, A.Ya., “The Laplace transformation and the differential equations with polynomial coefficients,” *Sov. J. Differ. Uravn.* Vol. **22**, 354–356 (1986).

Chapter 5

Confluent Heun function using reversible three-term recurrence formula

In this chapter I apply reversible three term recurrence formula to (1) the power series expansions in closed forms, (2) its integral forms of the Confluent Heun function (CHF) for infinite series and polynomial of type 2¹ including all higher terms of B_n 's² and (3) the generating function for Confluent Heun polynomial (CHP) of type 2.

5.1 Non-symmetrical canonical form of Confluent Heun Equation

In general, there are three types of the Confluent Heun equation (CHE): (1) Generalized spheroidal equation (GSE), (2) the generalized spheroidal wave equation (GSWE) in the Leaver version, (3) Non-symmetrical canonical form of the CHE.

In 1928, A.H. Wilson worked on the wave equation for the ion of the hydrogen molecule H_2^+ with considering the protons as at rest: he assume that the mass of the electron is negligible compared with that of the protons. Neglecting the mass of electrons in his wave equation, he obtained the second ordinary differential equation in elliptic

¹polynomial of type 2 is a polynomial which makes A_n term terminated in three term recursion relation of the power series in a linear differential equation.

²“ higher terms of B_n 's” means at least two terms of B_n 's.

coordinates for the approximative wave equation. And he introduced on the solution of the more general form of the Helmholtz equation in prolate spheroidal coordinates which is written by [9, 10]

$$\frac{d}{d\xi} \left((1 - \xi^2) \frac{dX}{d\xi} \right) + \left(\lambda^2 \xi^2 - 2p\lambda\xi - \frac{n_3^2}{1 - \xi^2} + \mu' \right) X = 0$$

When $p = 0$ this equation becomes the equation giving the solution of $\nabla^2 X - \lambda^2 X = 0$ in prolate spheroidal coordinates. Therefore, this equation is called the generalized spheroidal equation (GSE). The GSE has three singular points: two regular singular points which are ± 1 and one irregular singular point which is ∞ .

In 1986, E.W. Leaver introduced the different version of the GSE which is given by [6]

$$z(z - z_0) \frac{d^2 U}{dz^2} + (B_1 + B_2 z) \frac{dU}{dz} + (B_3 - 2\eta\omega(z - z_0) + \omega^2 z(z - z_0)) U(z) = 0 \quad \text{where } \omega \neq 0$$

This equation has three singular points: two regular singular points which are 0 and z_0 with exponents $\left\{0, 1 + \frac{B_1}{z_0}\right\}$ and $\left\{0, 1 - B_2 - \frac{B_1}{z_0}\right\}$, and one irregular singular point which is ∞ . It arises in the scattering problem of non-relativistic electrons from polar molecule and ions, in wave equation on the background geometry of the Schwarzschild black hole (Teukolsky's equation) and etc.

In Ref.[1, 2], Heun equation is a second-order linear ordinary differential equation of the form

$$\frac{d^2 y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0 \quad (5.1.1)$$

It has four regular singular points which are 0, 1, a and ∞ with exponents $\{0, 1 - \gamma\}$, $\{0, 1 - \delta\}$, $\{0, 1 - \epsilon\}$ and $\{\alpha, \beta\}$.

Heun equation has the four kinds of confluent forms: (1) Confluent Heun (two regular and one irregular singularities), (2) Doubly Confluent Heun (two irregular singularities), (3) Biconfluent Heun (one regular and one irregular singularities),³ (4) Triconfluent Heun equations (one irregular singularity). We can derive these four confluent forms from Heun equation by combining two or more regular singularities to take form an irregular singularity. Its process, converting Heun equation to other confluent forms, is similar to deriving of confluent hypergeometric equation from the hypergeometric equation. For the non-symmetrical canonical form of the Confluent Heun Equation (CHE).[2, 21, 8]

$$\frac{d^2 y}{dx^2} + \left(\beta + \frac{\gamma}{x} + \frac{\delta}{x-1} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)} y = 0 \quad (5.1.2)$$

³Biconfluent Heun equation is derived from the grand confluent hypergeometric equation by changing all coefficients $\mu = 1$ and $\varepsilon\omega = -q$. [36, 37]

(5.1.2) has three singular points: two regular singular points which are 0 and 1 with exponents $\{0, 1 - \gamma\}$ and $\{0, 1 - \delta\}$, and one irregular singular point which is ∞ with an exponent α . Its solution is denoted as $H_c(\alpha, \beta, \gamma, \delta, q; x)$.⁴ We can convert the GSE and its Leaver version to the non-symmetrical canonical form of the CHE by changing coefficients, adding the new variables and combining regular singularities.

5.2 Confluent Heun function with a regular singular point at 0

In chapter 4 I showed the analytic solutions of the CHE for infinite series and polynomial of type 1⁵ including all higher terms of A_n 's by applying 3TRF [1]; (1) the power series expansion of the CHE, (2) the integral representation of the CHF, (3) its generating function for the Confluent Heun polynomial (CHP) of type 1. Expressing the CHF in integral forms resulting in a precise and simplified transformation of the CHF to a confluent hypergeometric function. Also, the orthogonal relations of the CHF can be obtained from the integral forms.

In this chapter, by applying R3TRF, I construct the power series expansion in closed forms of non-symmetrical canonical form of the CHE (for infinite series and polynomial of type 2 including all higher terms of B_n 's), its integral forms and the generating function for the CHP of type 2 analytically.

Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} \quad \text{where } \lambda = \text{indicial root} \quad (5.2.1)$$

Above λ is indicial root. Plug (5.2.1) into (5.1.2). Then we get a three-term recurrence relation for the coefficients c_n :

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (5.2.2)$$

⁴Several authors denote as coefficients $4p$ and σ instead of β and q . And they define the solution of the CHE as $H_c^{(a)}(p, \alpha, \gamma, \delta, \sigma; x)$.

⁵polynomial of type 1 is a polynomial which makes B_n term terminated in three term recursion relation of the power series in a linear differential equation.

where,

$$\begin{aligned}
 A_n &= \frac{(n + \lambda)(n + \lambda - \beta + \gamma + \delta - 1) - q}{(n + \lambda + 1)(n + \lambda + \gamma)} \\
 &= \frac{\left(n - \frac{-(\varphi+2\lambda)-\sqrt{\varphi^2+4q}}{2}\right) \left(n - \frac{-(\varphi+2\lambda)+\sqrt{\varphi^2+4q}}{2}\right)}{(n + \lambda + 1)(n + \lambda + \gamma)}
 \end{aligned} \tag{5.2.3a}$$

and

$$\varphi = -\beta + \gamma + \delta - 1$$

$$B_n = \frac{\beta(n + \lambda + \alpha - 1)}{(n + \lambda + 1)(n + \lambda + \gamma)} \tag{5.2.3b}$$

$$c_1 = A_0 c_0 \tag{5.2.3c}$$

We have two indicial roots which are $\lambda_1 = 0$ and $\lambda_2 = 1 - \gamma$.

5.2.1 Power series

Polynomial of type 2

In the previous chapter, by applying 3TRF, I construct the power series expansion, its integral forms and the generating function for the CHP of type 1: I treat β , γ , δ and q as free variables and α as a fixed value. In this chapter, by applying R3TRF, I will show how to derive the power series expansion, the integral representation and the generating function for the CHP of type 2: I treat α , β , γ and δ as free variables and q as a fixed value.

In chapter 1 the general expression of power series of $y(x)$ for polynomial of type 2 is

defined by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\alpha_0} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\alpha_k} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\alpha_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (5.2.4)
\end{aligned}$$

In the above, $\alpha_i \leq \alpha_j$ only if $i \leq j$ where $i, j, \alpha_i, \alpha_j \in \mathbb{N}_0$.

For a polynomial, we need a condition which is:

$$A_{\alpha_i+2i} = 0 \quad \text{where } i, \alpha_i = 0, 1, 2, \dots \quad (5.2.5)$$

In the above, α_i is an eigenvalue that makes A_n term terminated at certain value of n .

(5.2.5) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (5.2.4).

The case of $\sqrt{\varphi^2 + 4q} = -\{\varphi + 2\lambda + 2(q_i + 2i)\}$ In (5.2.3a)-(5.2.3c) replace $\sqrt{\varphi^2 + 4q}$ by $-\{\varphi + 2\lambda + 2(q_i + 2i)\}$. In (5.2.5) replace index α_i by q_i . Take the new (5.2.3a)-(5.2.3c), (5.2.5) and put them in (1.3.11). After the replacement process, the

general expression of power series of the CHE for polynomial of type 2 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} x^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \gamma + \lambda)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + \varphi + 4 + 2\lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(-q_1)_{i_0} (q_1 + \varphi + 4 + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} x^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \gamma + \lambda)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha + \lambda)}{(i_k + 2(k+1) + \lambda)(i_k + 2k + 1 + \gamma + \lambda)} \right. \\
&\quad \times \left. \frac{(-q_k)_{i_k} (q_k + 4k + \varphi + 2\lambda)_{i_k} (2k + 1 + \lambda)_{i_{k-1}} (2k + \gamma + \lambda)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \varphi + 2\lambda)_{i_{k-1}} (2k + 1 + \lambda)_{i_k} (2k + \gamma + \lambda)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \varphi + 2\lambda)_{i_n} (2n + 1 + \lambda)_{i_{n-1}} (2n + \gamma + \lambda)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \varphi + 2\lambda)_{i_{n-1}} (2n + 1 + \lambda)_{i_n} (2n + \gamma + \lambda)_{i_n}} x^{i_n} \right\} z^n \Bigg\} \quad (5.2.6)
\end{aligned}$$

where

$$\begin{cases} z = \beta x^2 \\ \varphi = -\beta + \gamma + \delta - 1 \\ q = (q_j + 2j + \lambda)(q_j + 2j + \lambda + \varphi) \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

The case of $\sqrt{\varphi^2 + 4q} = \varphi + 2\lambda + 2(q_i + 2i)$ In (5.2.3a)-(5.2.3c) replace $\sqrt{\varphi^2 + 4q}$ by $\varphi + 2\lambda + 2(q_i + 2i)$. In (5.2.5) replace index α_i by q_i . Take the new (5.2.3a)-(5.2.3c), (5.2.5) and put them in (5.2.4). After the replacement process, its solution is equivalent to (5.2.6).

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = \frac{1}{2}$ for the second one in (5.2.6).

Remark 5.2.1 *The power series expansion of the CHE of the first kind for polynomial of*

type 2 about $x = 0$ as $q = (q_j + 2j)(q_j + 2j - \beta + \gamma + \delta - 1)$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= H_c^{(a)} F_{q_j}^R(\alpha, \beta, \gamma, \delta, q = (q_j + 2j)(q_j + 2j + \varphi); \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&= \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} x^{i_0} \\
&\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \varphi)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + \varphi + 4)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(-q_1)_{i_0} (q_1 + \varphi + 4)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma)_{i_1}} x^{i_1} \Big\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \varphi)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \right. \\
&\quad \times \frac{(-q_k)_{i_k} (q_k + 4k + \varphi)_{i_k} (2k + 1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + \varphi)_{i_{k-1}} (2k + 1)_{i_k} (2k + \gamma)_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + \varphi)_{i_n} (2n + 1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + \varphi)_{i_{n-1}} (2n + 1)_{i_n} (2n + \gamma)_{i_n}} x^{i_n} \Big\} z^n \quad (5.2.7)
\end{aligned}$$

Remark 5.2.2 The power series expansion of the CHE of the first kind for polynomial of

type 2 about $x = 0$ as $q = (q_j + 2j + 1 - \gamma)(q_j + 2j - \beta + \delta)$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
 y(x) &= H_c^{(a)} S_{\alpha_j}^R (\alpha, \beta, \gamma, \delta, q = (q_j + 2j + 1 - \gamma)(q_j + 2j - \beta + \delta); \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
 &= x^{1-\gamma} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2(1 - \gamma))_{i_0}}{(2 - \gamma)_{i_0} (1)_{i_0}} x^{i_0} \right. \\
 &\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha + 1 - \gamma)}{(i_0 + 3 - \gamma)(i_0 + 2)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2(1 - \gamma))_{i_0}}{(2 - \gamma)_{i_0} (1)_{i_0}} \right. \\
 &\quad \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + \varphi + 2(3 - \gamma))_{i_1}}{(-q_1)_{i_0} (q_1 + \varphi + 2(3 - \gamma))_{i_0}} \frac{(4 - \gamma)_{i_0} (3)_{i_0}}{(4 - \gamma)_{i_1} (3)_{i_1}} x^{i_1} \Big\} z \\
 &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \alpha + 1 - \gamma)}{(i_0 + 3 - \gamma)(i_0 + 2)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2(1 - \gamma))_{i_0}}{(2 - \gamma)_{i_0} (1)_{i_0}} \right. \\
 &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \alpha + 1 - \gamma)}{(i_k + 2k + 3 - \gamma)(i_k + 2k + 2)} \right. \\
 &\quad \times \frac{(-q_k)_{i_k} (q_k + \varphi + 2(2k + 1 - \gamma))_{i_k}}{(-q_k)_{i_{k-1}} (q_k + \varphi + 2(2k + 1 - \gamma))_{i_{k-1}}} \frac{(2k + 2 - \gamma)_{i_{k-1}} (2k + 1)_{i_{k-1}}}{(2k + 2 - \gamma)_{i_k} (2k + 1)_{i_k}} \Big\} \\
 &\quad \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + \varphi + 2(2n + 1 - \gamma))_{i_n}}{(-q_n)_{i_{n-1}} (q_n + \varphi + 2(2n + 1 - \gamma))_{i_{n-1}}} \frac{(2n + 2 - \gamma)_{i_{n-1}} (2n + 1)_{i_{n-1}}}{(2n + 2 - \gamma)_{i_n} (2n + 1)_{i_n}} x^{i_n} \Big\} z^n \Big\} \quad (5.2.8)
 \end{aligned}$$

In chapter 4 I treat α as a fixed value and β, γ, δ, q as free variables to construct the CHP of type 1: (1) if $\alpha = -(2\alpha_j + j)$ where $j, \alpha_j \in \mathbb{N}_0$, an analytic solution of the CHE turns to be the first kind of independent solution of the CHP of type 1. (2) if $\alpha = -(2\alpha_j + j + 1 - \gamma)$, an analytic solution of the CHE turns to be the second kind of independent solution of the CHP of type 1.

In this chapter I treat q as a fixed value and $\alpha, \beta, \gamma, \delta$ as free variables to construct the CHP of type 2: (1) if $q = (q_j + 2j)(q_j + 2j - \beta + \gamma + \delta - 1)$ where $j, q_j \in \mathbb{N}_0$, an analytic solution of the CHE turns to be the first kind of independent solution of the CHP of type 2. (2) if $q = (q_j + 2j + 1 - \gamma)(q_j + 2j - \beta + \delta)$, an analytic solution of the CHE turns to be the second kind of independent solution of the CHP of type 2.

Infinite series

In chapter 1 the general expression of power series of $y(x)$ for infinite series is defined by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (5.2.9)
\end{aligned}$$

Substitute (5.2.3a)–(5.2.3c) into (5.2.9). The general expression of power series of the CHE for infinite series about $x = 0$ is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^{\lambda} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} x^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} x^{i_1} \Bigg\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+\lambda+\alpha)}{(i_k+2(k+1)+\lambda)(i_k+2k+1+\gamma+\lambda)} \right. \\
&\quad \times \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1+\lambda)_{i_{k-1}} (2k+\gamma+\lambda)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1+\lambda)_{i_k} (2k+\gamma+\lambda)_{i_k}} \Bigg\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1+\lambda)_{i_{n-1}} (2n+\gamma+\lambda)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1+\lambda)_{i_n} (2n+\gamma+\lambda)_{i_n}} x^{i_n} \Bigg\} z^n \Bigg\} \quad (5.2.10)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi+2\lambda\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_1^\pm = \frac{\varphi+4+2\lambda\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_k^\pm = \frac{\varphi+2\lambda+4k\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_n^\pm = \frac{\varphi+2\lambda+4n\pm\sqrt{\varphi^2+4q}}{2} \end{cases}$$

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ for the second one in (5.2.10).

Remark 5.2.3 *The power series expansion of the CHE of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned} y(x) &= H_c^{(a)} F^R(\alpha, \beta, \gamma, \delta, q; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\ &= \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} x^{i_0} \right. \\ &\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \alpha)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (2 + \gamma)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (2 + \gamma)_{i_1}} x^{i_1} \right\} z \\ &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \alpha)}{(i_0 + 2)(i_0 + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\gamma)_{i_0}} \right. \\ &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \alpha)}{(i_k + 2(k+1))(i_k + 2k + 1 + \gamma)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1)_{i_{k-1}} (2k + \gamma)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1)_{i_k} (2k + \gamma)_{i_k}} \right\} \\ &\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1)_{i_{n-1}} (2n + \gamma)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1)_{i_n} (2n + \gamma)_{i_n}} x^{i_n} \right\} z^n \end{aligned} \quad (5.2.11)$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_1^\pm = \frac{\varphi+4\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_k^\pm = \frac{\varphi+4k\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_n^\pm = \frac{\varphi+4n\pm\sqrt{\varphi^2+4q}}{2} \end{cases}$$

Remark 5.2.4 *The power series expansion of the CHE of the second kind for infinite*

series about $x = 0$ using R3TRF is

$$\begin{aligned}
y(x) &= H_c^{(a)} S^R(\alpha, \beta, \gamma, \delta, q; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&= x^{1-\gamma} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} x^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+1-\gamma+\alpha)}{(i_0+3-\gamma)(i_0+2)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (4-\gamma)_{i_0} (3)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (4-\gamma)_{i_1} (3)_{i_1}} x^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+1-\gamma+\alpha)}{(i_0+3-\gamma)(i_0+2)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(2-\gamma)_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+1-\gamma+\alpha)}{(i_k+2k+3-\gamma)(i_k+2k+2)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+2-\gamma)_{i_{k-1}} (2k+1)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+2-\gamma)_{i_k} (2k+1)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+2-\gamma)_{i_{n-1}} (2n+1)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+2-\gamma)_{i_n} (2n+1)_{i_n}} x^{i_n} \right\} z^n \Bigg\} \quad (5.2.12)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\varphi+2(1-\gamma)\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_1^{\pm} = \frac{\varphi+2(3-\gamma)\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_k^{\pm} = \frac{\varphi+2(2k+1-\gamma)\pm\sqrt{\varphi^2+4q}}{2} \\ \Delta_n^{\pm} = \frac{\varphi+2(2n+1-\gamma)\pm\sqrt{\varphi^2+4q}}{2} \end{cases}$$

It is required that $\gamma \neq 0, -1, -2, \dots$ for the first kind of independent solution of the CHE for all cases. Because if it does not, its solution will be divergent. And it is required that $\gamma \neq 2, 3, 4, \dots$ for the second kind of independent solution of the CHE for all cases.

The infinite series in this chapter are equivalent to the infinite series in chapter 4. In this chapter B_n is the leading term in sequence c_n of the analytic function $y(x)$. In chapter 4 A_n is the leading term in sequence c_n of the analytic function $y(x)$.

For the special case, as $|\beta| \ll 1$ in (5.2.11) and (5.2.12), we have

$$\begin{aligned}
&\lim_{|\beta| \ll 1} H_c^{(a)} F^R(\alpha, \beta, \gamma, \delta, q; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&\approx {}_2F_1 \left(\frac{\gamma + \delta - 1 - \sqrt{(\gamma + \delta - 1)^2 + 4q}}{2}, \frac{\gamma + \delta - 1 + \sqrt{(\gamma + \delta - 1)^2 + 4q}}{2}; \gamma; x \right) \quad (5.2.13a)
\end{aligned}$$

And,

$$\begin{aligned} & \lim_{|\beta| \ll 1} H_c^{(a)} S^R(\alpha, \beta, \gamma, \delta, q; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\ & \approx x^{1-\gamma} {}_2F_1\left(\frac{-\gamma + \delta + 1 - \sqrt{(\gamma + \delta - 1)^2 + 4q}}{2}, \frac{-\gamma + \delta + 1 + \sqrt{(\gamma + \delta - 1)^2 + 4q}}{2}; 2 - \gamma; x\right) \end{aligned} \quad (5.2.13b)$$

(5.2.13a) and (5.2.13b) are Gauss Hypergeometric function.

5.2.2 Integral formalism

In earlier literature the integral relations of the CHE were constructed by using Fredholm integral equations; such integral relationships express one analytic solution in terms of another analytic solution such as a confluent hypergeometric function with a branch point at zero. [2] There are many other forms of integral relations in the CHE. [14, 15, 16, 17, 18, 19]

In chapter 4 I derive the combined definite and contour integral forms of the CHF by applying 3TRF: a ${}_1F_1$ function recurs in each of sub-integral forms of the CHF. Now I consider the combined definite and contour integral representations of the CHF by using R3TRF.

Polynomial of type 2

There is a generalized hypergeometric function which is given by

$$\begin{aligned} I_l &= \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} (q_l + 4l + \varphi + 2\lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \gamma + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} (q_l + 4l + \varphi + 2\lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \gamma + \lambda)_{i_l}} x^{i_l} \\ &= x^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1} + 2l + \lambda, j + 1) B(i_{l-1} + 2l - 1 + \gamma + \lambda, j + 1)}{(i_{l-1} + 2l + \lambda)^{-1} (i_{l-1} + 2l - 1 + \gamma + \lambda)^{-1}} \\ &\quad \times \frac{(i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + \varphi + 2\lambda)_j}{(1)_j j!} x^j \end{aligned} \quad (5.2.14)$$

By using integral form of beta function,

$$B(i_{l-1} + 2l + \lambda, j + 1) = \int_0^1 dt_l t_l^{i_{l-1} + 2l - 1 + \lambda} (1 - t_l)^j \quad (5.2.15a)$$

$$B(i_{l-1} + 2l - 1 + \gamma + \lambda, j + 1) = \int_0^1 du_l u_l^{i_{l-1} + 2(l-1) + \gamma + \lambda} (1 - u_l)^j \quad (5.2.15b)$$

Substitute (5.2.15a) and (5.2.15b) into (5.2.14). And divide $(i_{l-1} + 2l + \lambda)(i_{l-1} + 2l - 1 + \gamma + \lambda)$ into I_l .

$$\begin{aligned}
K_l &= \frac{1}{(i_{l-1} + 2l + \lambda)(i_{l-1} + 2l - 1 + \gamma + \lambda)} \\
&\times \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} (q_l + 4l + \varphi + 2\lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \gamma + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} (q_l + 4l + \varphi + 2\lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \gamma + \lambda)_{i_l}} x^{i_l} \\
&= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} (xt_l u_l)^{i_{l-1}} \\
&\times \sum_{j=0}^{\infty} \frac{(i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + \varphi + 2\lambda)_j}{(1)_j j!} [x(1 - t_l)(1 - u_l)]^j
\end{aligned}$$

The integral form of Gauss hypergeometric function is defined by

$$\begin{aligned}
{}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{j=0}^{\infty} \frac{(\alpha)_j (\beta)_j}{(\gamma)_j (j!)} z^j \\
&= -\frac{1}{2\pi i} \frac{\Gamma(1-\alpha)\Gamma(\gamma)}{\Gamma(\gamma-\alpha)} \oint dv_l (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \quad (5.2.16) \\
&\text{where } \operatorname{Re}(\gamma - \alpha) > 0
\end{aligned}$$

replaced α, β, γ and z by $i_{l-1} - q_l, i_{l-1} + q_l + 4l + \varphi + 2\lambda, 1$ and $x(1 - t_l)(1 - u_l)$ in (5.2.16)

$$\begin{aligned}
&\sum_{j=0}^{\infty} \frac{(i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + \varphi + 2\lambda)_j}{(1)_j j!} [x(1 - t_l)(1 - u_l)]^j \\
&= \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \frac{1}{1 - x(1 - t_l)(1 - u_l)v_l} \right)^{q_l} (1 - x(1 - t_l)(1 - u_l)v_l)^{-(4l+\varphi+2\lambda)} \\
&\times \left(\frac{v_l}{v_l - 1} \frac{1}{1 - x(1 - t_l)(1 - u_l)v_l} \right)^{i_{l-1}} \quad (5.2.17)
\end{aligned}$$

Substitute (5.2.17) into K_l .

$$\begin{aligned}
K_l &= \frac{1}{(i_{l-1} + 2l + \lambda)(i_{l-1} + 2l - 1 + \gamma + \lambda)} \\
&\times \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} (q_l + 4l + \varphi + 2\lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \gamma + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} (q_l + 4l + \varphi + 2\lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \gamma + \lambda)_{i_l}} x^{i_l} \\
&= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \frac{1}{1 - x(1 - t_l)(1 - u_l)v_l} \right)^{q_l} \\
&\times (1 - x(1 - t_l)(1 - u_l)v_l)^{-(4l+\varphi+2\lambda)} \left(\frac{v_l}{v_l - 1} \frac{xt_l u_l}{1 - x(1 - t_l)(1 - u_l)v_l} \right)^{i_{l-1}} \quad (5.2.18)
\end{aligned}$$

Substitute (5.2.18) into (5.2.6) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(x)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(x)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(x)$, etc.⁶

Theorem 5.2.5 *The general representation in the form of integral of the CHP of type 2 is given by*

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} x^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\varphi+2\lambda)} \\
&\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)+\alpha+\lambda} \left. \right\} \\
&\quad \times \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftarrow{w}_{1,n}^{i_0} \left. \right\} z^n \Bigg\} \tag{5.2.19}
\end{aligned}$$

where

$$\overleftarrow{w}_{i,j} = \begin{cases} \frac{v_i}{(v_i - 1)} \frac{\overleftarrow{w}_{i+1,j} t_i u_i}{1 - \overleftarrow{w}_{i+1,j} v_i (1 - t_i)(1 - u_i)} & \text{where } i \leq j \\ x & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof According to (5.2.6),

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots \tag{5.2.20}$$

⁶ $y_1(x)$ means the sub-power series in (5.2.6) contains one term of B'_n 's, $y_2(x)$ means the sub-power series in (5.2.6) contains two terms of B'_n 's, $y_3(x)$ means the sub-power series in (5.2.6) contains three terms of B'_n 's, etc.

In the above, the power series expansion of sub-summation $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of the type 2 CHP using R3TRF about $x = 0$ are

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} x^{i_0} \quad (5.2.21a)$$

$$y_1(x) = c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\ \left. \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \varphi + 2\lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \varphi + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} x^{i_1} \right\} z \quad (5.2.21b)$$

$$y_2(x) = c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\ \times \sum_{i_1=i_0}^{q_1} \frac{(i_1 + 2 + \lambda + \alpha)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(-q_1)_{i_1} (q_1 + 4 + \varphi + 2\lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \varphi + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\ \left. \times \sum_{i_2=i_1}^{q_2} \frac{(-q_2)_{i_2} (q_2 + 8 + \varphi + 2\lambda)_{i_2} (5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(-q_2)_{i_1} (q_2 + 8 + \varphi + 2\lambda)_{i_1} (5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} x^{i_2} \right\} z^2 \quad (5.2.21c)$$

$$y_3(x) = c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\ \times \sum_{i_1=i_0}^{q_1} \frac{(i_1 + 2 + \lambda + \alpha)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(-q_1)_{i_1} (q_1 + 4 + \varphi + 2\lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \varphi + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\ \times \sum_{i_2=i_1}^{q_2} \frac{(i_2 + 4 + \lambda + \alpha)}{(i_2 + \lambda + 6)(i_2 + \lambda + 5 + \gamma)} \frac{(-q_2)_{i_2} (q_2 + 8 + \varphi + 2\lambda)_{i_2} (5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(-q_2)_{i_1} (q_2 + 8 + \varphi + 2\lambda)_{i_1} (5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} \\ \left. \times \sum_{i_3=i_2}^{q_3} \frac{(-q_3)_{i_3} (q_2 + 12 + \varphi + 2\lambda)_{i_3} (7 + \lambda)_{i_2} (6 + \gamma + \lambda)_{i_2}}{(-q_3)_{i_2} (q_2 + 12 + \varphi + 2\lambda)_{i_2} (7 + \lambda)_{i_3} (6 + \gamma + \lambda)_{i_3}} x^{i_3} \right\} z^3 \quad (5.2.21d)$$

Put $l = 1$ in (5.2.18). Take the new (5.2.18) into (5.2.21b).

$$y_1(x) = \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - x(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \\ \times (1 - x(1 - t_1)(1 - u_1)v_1)^{-(4+\varphi+2\lambda)} \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\lambda} \\ \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right\} z \quad (5.2.22)$$

where

$$\overleftarrow{w}_{1,1} = \frac{v_1}{v_1 - 1} \frac{xt_1 u_1}{1 - x(1 - t_1)(1 - u_1)v_1}$$

Put $l = 2$ in (5.2.18). Take the new (5.2.18) into (5.2.21c).

$$\begin{aligned} y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - x(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\ &\times (1 - x(1 - t_2)(1 - u_2)v_2)^{-(8+\varphi+2\lambda)} \overleftarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{2+\alpha+\lambda} \\ &\times \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\ &\times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + \varphi + 2\lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(-q_1)_{i_0} (q_1 + 4 + \varphi + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \overleftarrow{w}_{2,2}^{i_1} \right\} z^2 \end{aligned} \quad (5.2.23)$$

where

$$\overleftarrow{w}_{2,2} = \frac{v_2}{v_2 - 1} \frac{xt_2 u_2}{1 - x(1 - t_2)(1 - u_2)v_2}$$

Put $l = 1$ and $\eta = \overleftarrow{w}_{2,2}$ in (5.2.18). Take the new (5.2.18) into (5.2.23).

$$\begin{aligned} y_2(x) &= \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - x(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\ &\times (1 - x(1 - t_2)(1 - u_2)v_2)^{-(8+\varphi+2\lambda)} \overleftarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{2+\alpha+\lambda} \\ &\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \\ &\times (1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1)^{-(4+\varphi+2\lambda)} \overleftarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,2} \partial_{\overleftarrow{w}_{1,2}} \right) \overleftarrow{w}_{1,2}^{\alpha+\lambda} \\ &\times \left\{ c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftarrow{w}_{1,2}^{i_0} \right\} z^2 \end{aligned} \quad (5.2.24)$$

where

$$\overleftarrow{w}_{1,2} = \frac{v_1}{v_1 - 1} \frac{\overleftarrow{w}_{2,2} t_1 u_1}{1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1}$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the

integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3-1}{v_3} \frac{1}{1-x(1-t_3)(1-u_3)v_3} \right)^{q_3} \\
& \times (1-x(1-t_3)(1-u_3)v_3)^{-(12+\varphi+2\lambda)} \overleftarrow{w}_{3,3}^{-(4+\alpha+\lambda)} \left(\overleftarrow{w}_{3,3} \partial \overleftarrow{w}_{3,3} \right) \overleftarrow{w}_{3,3}^{4+\alpha+\lambda} \\
& \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \frac{1}{1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \right)^{q_2} \\
& \times (1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2)^{-(8+\varphi+2\lambda)} \overleftarrow{w}_{2,3}^{-(2+\alpha+\lambda)} \left(\overleftarrow{w}_{2,3} \partial \overleftarrow{w}_{2,3} \right) \overleftarrow{w}_{2,3}^{2+\alpha+\lambda} \\
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \frac{1}{1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
& \times (1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1)^{-(4+\varphi+2\lambda)} \overleftarrow{w}_{1,3}^{-(\alpha+\lambda)} \left(\overleftarrow{w}_{1,3} \partial \overleftarrow{w}_{1,3} \right) \overleftarrow{w}_{1,3}^{\alpha+\lambda} \\
& \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftarrow{w}_{1,3}^{i_0} \right\} z^3 \tag{5.2.25}
\end{aligned}$$

where

$$\begin{cases} \overleftarrow{w}_{3,3} = \frac{v_3}{v_3-1} \frac{xt_3u_3}{1-x(1-t_3)(1-u_3)v_3} \\ \overleftarrow{w}_{2,3} = \frac{v_2}{v_2-1} \frac{\overleftarrow{w}_{3,3}t_2u_2}{1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \\ \overleftarrow{w}_{1,3} = \frac{v_1}{v_1-1} \frac{\overleftarrow{w}_{2,3}t_1u_1}{1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Substitute (5.2.21a), (5.2.22), (5.2.24), (5.2.25) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (5.2.20).

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ for the second one in (5.2.19).

Remark 5.2.6 *The integral representation of the CHE of the first kind for polynomial of*

type 2 about $x = 0$ as $q = (q_j + 2j)(-\beta + \gamma + \delta + q_j + 2j - 1)$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= H_c^{(a)} F_{q_j}^R(\alpha, \beta, \gamma, \delta, q = (q_j + 2j)(q_j + 2j + \varphi); \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&= {}_2F_1(-q_0, q_0 + \varphi; \gamma; x) \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\varphi)} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\alpha} \Big\} \\
&\quad \times {}_2F_1(-q_0, q_0 + \varphi; \gamma; \overleftrightarrow{w}_{1,n}) \Big\} z^n \tag{5.2.26}
\end{aligned}$$

Remark 5.2.7 The integral representation of the CHE of the first kind for polynomial of type 2 about $x = 0$ as $q = (q_j + 2j + 1 - \gamma)(-\beta + \delta + q_j + 2j)$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= H_c^{(a)} S_{\alpha_j}^R(\alpha, \beta, \gamma, \delta, q = (q_j + 2j + 1 - \gamma)(q_j + 2j + 1 - \gamma + \varphi); \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&= x^{1-\gamma} \left\{ {}_2F_1(-q_0, q_0 + \varphi + 2(1 - \gamma); 2 - \gamma; x) \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\varphi+2(1-\gamma))} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \Big\} \\
&\quad \times {}_2F_1(-q_0, q_0 + \varphi + 2(1 - \gamma); 2 - \gamma; \overleftrightarrow{w}_{1,n}) \Big\} z^n \tag{5.2.27}
\end{aligned}$$

Infinite series

Let's consider the integral representation of the CHE about $x = 0$ for infinite series by applying R3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned} M_l &= \sum_{i_l=i_{l-1}}^{\infty} \frac{(\Delta_l^-)_{i_l} (\Delta_l^+)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(\Delta_l^-)_{i_{l-1}} (\Delta_l^+)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} x^{i_l} \\ &= x^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+2l+\lambda, j+1) B(i_{l-1}+2l-1+\gamma+\lambda, j+1) (\Delta_l^- + i_{l-1})_j (\Delta_l^+ + i_{l-1})_j}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-1+\gamma+\lambda)^{-1} (1)_j j!} x^j \end{aligned} \quad (5.2.28)$$

where

$$\Delta_l^{\pm} = \frac{\varphi + 2\lambda + 4l \pm \sqrt{\varphi^2 + 4q}}{2}$$

Substitute (5.2.15a) and (5.2.15b) into (5.2.28). And divide $(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)$ into the new (5.2.28).

$$\begin{aligned} V_l &= \frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\Delta_l^-)_{i_l} (\Delta_l^+)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(\Delta_l^-)_{i_{l-1}} (\Delta_l^+)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} x^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} (x t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(\Delta_l^- + i_{l-1})_j (\Delta_l^+ + i_{l-1})_j}{(1)_j j!} (x(1-t_l)(1-u_l))^j \end{aligned}$$

The hypergeometric function is defined by

$$\begin{aligned} {}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n (n!)} z^n \\ &= \frac{1}{2\pi i} \frac{\Gamma(1+\alpha-\gamma)}{\Gamma(\alpha)} \int_0^{(1+)} dv_l (-1)^\gamma (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \quad (5.2.29) \\ &\text{where } \gamma - \alpha \neq 1, 2, 3, \dots, \operatorname{Re}(\alpha) > 0 \end{aligned}$$

Replace α, β, γ and z by $\Delta_l^- + i_{l-1}, \Delta_l^+ + i_{l-1}, 1$ and $x(1-t_l)(1-u_l)$ in (5.2.29). Take the new (5.2.29) into V_l .

$$\begin{aligned} V_l &= \frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\gamma+\lambda)} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\Delta_l^-)_{i_l} (\Delta_l^+)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\gamma+\lambda)_{i_{l-1}}}{(\Delta_l^-)_{i_{l-1}} (\Delta_l^+)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\gamma+\lambda)_{i_l}} x^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2(l-1)+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l-1}{v_l} \right)^{-\Delta_l^-} \\ &\quad \times (1-x(1-t_l)(1-u_l)v_l)^{-\Delta_l^+} \left(\frac{v_l}{v_l-1} \frac{x t_l u_l}{1-x(1-t_l)(1-u_l)v_l} \right)^{i_{l-1}} \end{aligned} \quad (5.2.30)$$

Substitute (5.2.30) into (5.2.10) where $l = 1, 2, 3, \dots$; apply V_1 into the second summation of sub-power series $y_1(x)$, apply V_2 into the third summation and V_1 into the second summation of sub-power series $y_2(x)$, apply V_3 into the forth summation, V_2 into the third summation and V_1 into the second summation of sub-power series $y_3(x)$, etc.⁷

Theorem 5.2.8 *The general representation in the form of integral of the CHE for infinite series about $x = 0$ using R3TRF is given by*

$$\begin{aligned}
y(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} x^{i_0} \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\Delta_{n-k}^+} \\
& \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\alpha+\lambda} \left. \right\} \\
& \times \left. \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \right\} z^n \Bigg\} \tag{5.2.31}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi+2\lambda \pm \sqrt{\varphi^2+4q}}{2} \\ \Delta_{n-k}^\pm = \frac{\varphi+2\lambda+4(n-k) \pm \sqrt{\varphi^2+4q}}{2} \end{cases}$$

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof In (5.2.10) sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of the CHE for infinite series about $x = 0$ using R3TRF are given by

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} x^{i_0} \tag{5.2.32a}$$

⁷ $y_1(x)$ means the sub-power series in (5.2.10) contains one term of B'_n 's, $y_2(x)$ means the sub-power series in (5.2.10) contains two terms of B'_n 's, $y_3(x)$ means the sub-power series in (5.2.10) contains three terms of B'_n 's, etc.

$$\begin{aligned}
y_1(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
& \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} x^{i_1} \right\} z \quad (5.2.32b)
\end{aligned}$$

$$\begin{aligned}
y_2(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + 2 + \lambda + \alpha)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\
& \times \left. \sum_{i_2=i_1}^{\infty} \frac{(\Delta_2^-)_{i_2} (\Delta_2^+)_{i_2} (5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(\Delta_2^-)_{i_1} (\Delta_2^+)_{i_1} (5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} x^{i_2} \right\} z^2 \quad (5.2.32c)
\end{aligned}$$

$$\begin{aligned}
y_3(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \lambda + \alpha)}{(i_0 + \lambda + 2)(i_0 + \lambda + 1 + \gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + 2 + \lambda + \alpha)}{(i_1 + \lambda + 4)(i_1 + \lambda + 3 + \gamma)} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3 + \lambda)_{i_0} (2 + \gamma + \lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3 + \lambda)_{i_1} (2 + \gamma + \lambda)_{i_1}} \\
& \times \sum_{i_2=i_1}^{\infty} \frac{(i_2 + 4 + \lambda + \alpha)}{(i_2 + \lambda + 6)(i_2 + \lambda + 5 + \gamma)} \frac{(\Delta_2^-)_{i_2} (\Delta_2^+)_{i_2} (5 + \lambda)_{i_1} (4 + \gamma + \lambda)_{i_1}}{(\Delta_2^-)_{i_1} (\Delta_2^+)_{i_1} (5 + \lambda)_{i_2} (4 + \gamma + \lambda)_{i_2}} \\
& \times \left. \sum_{i_3=i_2}^{\infty} \frac{(\Delta_3^-)_{i_3} (\Delta_3^+)_{i_3} (7 + \lambda)_{i_2} (6 + \gamma + \lambda)_{i_2}}{(\Delta_3^-)_{i_2} (\Delta_3^+)_{i_2} (7 + \lambda)_{i_3} (6 + \gamma + \lambda)_{i_3}} x^{i_3} \right\} z^3 \quad (5.2.32d)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi + 2\lambda \pm \sqrt{\varphi^2 + 4q}}{2} \\ \Delta_1^\pm = \frac{\varphi + 2\lambda + 4 \pm \sqrt{\varphi^2 + 4q}}{2} \\ \Delta_2^\pm = \frac{\varphi + 2\lambda + 8 \pm \sqrt{\varphi^2 + 4q}}{2} \\ \Delta_3^\pm = \frac{\varphi + 2\lambda + 12 \pm \sqrt{\varphi^2 + 4q}}{2} \end{cases}$$

Put $l = 1$ in (5.2.30). Take the new (5.2.30) into (5.2.32b).

$$\begin{aligned}
y_1(x) &= \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \right)^{-\Delta_1^-} \\
&\quad \times (1-x(1-t_1)(1-u_1)v_1)^{-\Delta_1^+} \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\lambda} \\
&\quad \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right\} z
\end{aligned} \tag{5.2.33}$$

where

$$\overleftrightarrow{w}_{1,1} = \frac{v_1}{v_1-1} \frac{xt_1 u_1}{1-x(1-t_1)(1-u_1)v_1}$$

Put $l = 2$ in (5.2.30). Take the new (5.2.30) into (5.2.32c).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \right)^{-\Delta_2^-} \\
&\quad \times (1-x(1-t_2)(1-u_2)v_2)^{-\Delta_2^+} \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2+\alpha+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \overleftrightarrow{w}_{2,2}^{i_1} \right\} z^2
\end{aligned} \tag{5.2.34}$$

where

$$\overleftrightarrow{w}_{2,2} = \frac{v_2}{v_2-1} \frac{xt_2 u_2}{1-x(1-t_2)(1-u_2)v_2}$$

Put $l = 1$ and $\eta = \overleftrightarrow{w}_{2,2}$ in (5.2.30). Take the new (5.2.30) into (5.2.34).

$$\begin{aligned}
y_2(x) &= \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \right)^{-\Delta_2^-} \\
&\quad \times (1-x(1-t_2)(1-u_2)v_2)^{-\Delta_2^+} \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2+\alpha+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \right)^{-\Delta_1^-} \\
&\quad \times (1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\Delta_1^+} \overleftrightarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\alpha+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right\} z^2
\end{aligned} \tag{5.2.35}$$

where

$$\overleftrightarrow{w}_{1,2} = \frac{v_1}{v_1 - 1} \frac{\overleftrightarrow{w}_{2,2} t_1 u_1}{1 - \overleftrightarrow{w}_{2,2} (1 - t_1) (1 - u_1) v_1}$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned} y_3(x) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3 - 1}{v_3} \right)^{-\Delta_3^-} \\ & \times (1 - x(1 - t_3)(1 - u_3)v_3)^{-\Delta_3^+} \overleftrightarrow{w}_{3,3}^{-(4+\alpha+\lambda)} \left(\overleftrightarrow{w}_{3,3} \partial_{\overleftrightarrow{w}_{3,3}} \right) \overleftrightarrow{w}_{3,3}^{4+\alpha+\lambda} \\ & \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \right)^{-\Delta_2^-} \\ & \times (1 - \overleftrightarrow{w}_{3,3}(1 - t_2)(1 - u_2)v_2)^{-\Delta_2^+} \overleftrightarrow{w}_{2,3}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,3} \partial_{\overleftrightarrow{w}_{2,3}} \right) \overleftrightarrow{w}_{2,3}^{2+\alpha+\lambda} \\ & \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \right)^{-\Delta_1^-} \\ & \times (1 - \overleftrightarrow{w}_{2,3}(1 - t_1)(1 - u_1)v_1)^{-\Delta_1^+} \overleftrightarrow{w}_{1,3}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,3} \partial_{\overleftrightarrow{w}_{1,3}} \right) \overleftrightarrow{w}_{1,3}^{\alpha+\lambda} \\ & \times \left\{ c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,3}^{i_0} \right\} z^3 \end{aligned} \quad (5.2.36)$$

where

$$\begin{cases} \overleftrightarrow{w}_{3,3} = \frac{v_3}{v_3-1} \frac{x t_3 u_3}{1-x(1-t_3)(1-u_3)v_3} \\ \overleftrightarrow{w}_{2,3} = \frac{v_2}{v_2-1} \frac{\overleftrightarrow{w}_{3,3} t_2 u_2}{1-\overleftrightarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \\ \overleftrightarrow{w}_{1,3} = \frac{v_1}{v_1-1} \frac{\overleftrightarrow{w}_{2,3} t_1 u_1}{1-\overleftrightarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute (5.2.32a), (5.2.33), (5.2.35), (5.2.36) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (5.2.10), we obtain (5.2.31).⁸

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the CHE and $\lambda = 1 - \gamma$ for the second one in (5.2.31).

⁸Or replace the finite summation with an interval $[0, q_0]$ by infinite summation with an interval $[0, \infty]$ in (5.2.19). Replace q_0 and q_{n-k} by $\frac{-(\varphi+2\lambda)+\sqrt{\varphi^2+4q}}{2}$ and $\frac{-(\varphi+2(\lambda+2(n-k)))+\sqrt{\varphi^2+4q}}{2}$ into the new (5.2.19). Its solution is also equivalent to (5.2.31)

Remark 5.2.9 *The integral representation of the CHE of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
y(x) &= H_c^{(a)} F^R(\alpha, \beta, \gamma, \delta, q; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&= {}_2F_1(\Delta_0^-, \Delta_0^+; \gamma; x) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) v_{n-k})^{-\Delta_{n-k}^+} \\
&\quad \times \left. \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\alpha} \right\} {}_2F_1(\Delta_0^-, \Delta_0^+; \gamma; \overleftrightarrow{w}_{1,n}) \Big\} z^n \quad (5.2.37)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi \pm \sqrt{\varphi^2 + 4q}}{2} \\ \Delta_{n-k}^\pm = \frac{\varphi + 4(n-k) \pm \sqrt{\varphi^2 + 4q}}{2} \end{cases}$$

Remark 5.2.10 *The integral representation of the CHE of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
y(x) &= H_c^{(a)} S^R(\alpha, \beta, \gamma, \delta, q; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
&= x^{1-\gamma} \left\{ {}_2F_1(\Delta_0^-, \Delta_0^+; 2 - \gamma; x) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) v_{n-k})^{-\Delta_{n-k}^+} \\
&\quad \times \left. \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \right\} \\
&\quad \times \left. {}_2F_1(\Delta_0^-, \Delta_0^+; 2 - \gamma; \overleftrightarrow{w}_{1,n}) \right\} \Big\} z^n \quad (5.2.38)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi + 2(1-\gamma) \pm \sqrt{\varphi^2 + 4q}}{2} \\ \Delta_{n-k}^\pm = \frac{\varphi + 2(1-\gamma) + 4(n-k) \pm \sqrt{\varphi^2 + 4q}}{2} \end{cases}$$

5.2.3 Generating function for the CHP of type 2

In chapter 4 I construct the generating function for the CHP of type 1 by applying 3TRF. Now I consider the generating function for the CHP of type 2. Since the

generating function for the CHP is derived, we might be possible to construct orthogonal relations of the CHP.⁹

Lemma 5.2.11 *The generating function for Jacobi polynomial using hypergeometric functions is given by*

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} w^{q_0} {}_2F_1(-q_0, q_0 + A; \gamma; x) \\ &= 2^{A-1} \frac{\left(1 - w + \sqrt{w^2 - 2(1 - 2x)w + 1}\right)^{1-\gamma} \left(1 + w + \sqrt{w^2 - 2(1 - 2x)w + 1}\right)^{\gamma-A}}{\sqrt{w^2 - 2(1 - 2x)w + 1}} \end{aligned} \quad (5.2.39)$$

where $|w| < 1$

Proof The proof of this lemma is given in Lemma 3.2.1

Definition 5.2.12 *I define that*

$$\left\{ \begin{aligned} s_{a,b} &= \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{if } a > b \\ s_a & \text{if } a = b \end{cases} \\ \tilde{w}_{i,j} &= \begin{cases} \frac{\tilde{w}_{i+1,j} t_i u_i \{1 + (s_i + 2\tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))s_i\}}{2(1 - \tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))^2 s_i} \\ - \frac{\tilde{w}_{i+1,j} t_i u_i (1 + s_i) \sqrt{s_i^2 - 2(1 - 2\tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))s_i + 1}}{2(1 - \tilde{w}_{i+1,j}(1 - t_i)(1 - u_i))^2 s_i} & \text{where } i < j \\ \frac{x t_i u_i \{1 + (s_{i,\infty} + 2x(1 - t_i)(1 - u_i))s_{i,\infty}\}}{2(1 - x(1 - t_i)(1 - u_i))^2 s_{i,\infty}} \\ - \frac{x t_i u_i (1 + s_{i,\infty}) \sqrt{s_{i,\infty}^2 - 2(1 - 2x(1 - t_i)(1 - u_i))s_{i,\infty} + 1}}{2(1 - x(1 - t_i)(1 - u_i))^2 s_{i,\infty}} & \text{where } i = j \end{cases} \end{aligned} \right. \quad (5.2.40)$$

where

$$a, b, i, j \in \mathbb{N}_0$$

⁹Polynomial of type 3 is a polynomial which makes A_n and B_n terms terminated at the same time in three term recursion relation of the power series in a linear differential equation. For the type 1 polynomial, I treat β, γ, δ and q as free variables and α as a fixed value. For the type 2 polynomial, I treat α, β, γ and δ as free variables and q as a fixed value. For the type 3 polynomial, I treat β, γ and δ as free variables and α and q as fixed values. In the next series, I will construct the generating functions for the CHP of type 3.

And we have

$$\sum_{q_i=q_j}^{\infty} r_i^{q_i} = \frac{r_i^{q_j}}{(1 - r_i)} \quad (5.2.41)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (5.2.19) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (5.2.40) and (5.2.41),

Theorem 5.2.13 *The general expression of the generating function for the CHP of type*

$\mathcal{Z}$ is given by

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y(x) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \Upsilon(\lambda; s_{0,\infty}; x) \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\varphi+2\lambda)} \\
&\times \tilde{w}_{1,1}^{-(\alpha+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\lambda} \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) z \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1+\lambda} \int_0^1 du_n u_n^{2(n-1)+\gamma+\lambda} \right. \\
&\times (s_{n,\infty}^2 - 2(1-2x(1-t_n)(1-u_n))s_{n,\infty} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2x(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\varphi+2\lambda)} \\
&\times \tilde{w}_{n,n}^{-(2(n-1)+\alpha+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\alpha+\lambda} \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \\
&\times (s_{n-k}^2 - 2(1-2\tilde{w}_{n+1-k,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n+1-k,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\varphi+2\lambda)} \\
&\times \left. \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\alpha+\lambda} \right\} \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) \Big\} z^n \tag{5.2.42}
\end{aligned}$$

where

$$\begin{cases} \mathfrak{Y}(\lambda; s_{0,\infty}; x) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_{0,\infty}^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} x^{i_0} \right) \\ \mathfrak{Y}(\lambda; s_0; \tilde{w}_{1,1}) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \\ \mathfrak{Y}(\lambda; s_0; \tilde{w}_{1,n}) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \tilde{w}_{1,n}^{i_0} \right) \end{cases}$$

Proof Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on the form of integral of the type 2 CHP $y(x)$,

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y(x) \\ &= \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} \{y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots\} \end{aligned} \quad (5.2.43)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (5.2.21a),

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_0(x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_{0,\infty}^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} x^{i_0} \right) \end{aligned} \quad (5.2.44)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (5.2.22),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-x(1-t_1)(1-u_1)v_1)^{-(4+\varphi+2\lambda)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_{1,\infty}}{1-x(1-t_1)(1-u_1)v_1} \right)^{q_1} \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right) z \tag{5.2.45}
\end{aligned}$$

Replace q_i, q_j and r_i by q_1, q_0 and $\frac{v_1-1}{v_1} \frac{s_{1,\infty}}{1-x(1-t_1)(1-u_1)v_1}$ in (5.2.41). Take the new (5.2.41) into (5.2.45).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{(1-x(1-t_1)(1-u_1)v_1)^{-(3+\varphi+2\lambda)}}{-x(1-t_1)(1-u_1)v_1^2 + (1-s_{1,\infty})v_1 + s_{1,\infty}} \\
&\times \overleftrightarrow{w}_{1,1}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\lambda} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} \left(\frac{v_1-1}{v_1} \frac{s_{0,\infty}}{1-x(1-t_1)(1-u_1)v_1} \right)^{q_0} \\
&\times \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right) z \tag{5.2.46}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$v_1 = \frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)}$$

$$\text{or } \frac{1-s_{1,\infty} + \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)} \text{ and}$$

$$\frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)} \text{ is only inside the unit circle. As we}$$

compute the residue there in (5.2.46) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\varphi+2\lambda)} \\
&\times \tilde{w}_{1,1}^{-(\alpha+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\lambda} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) z
\end{aligned} \tag{5.2.47}$$

where

$$\begin{aligned}
\tilde{w}_{1,1} &= \frac{v_1}{(v_1 - 1)} \frac{xt_1 u_1}{1 - xv_1(1-t_1)(1-u_1)} \Bigg|_{v_1 = \frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)}} \\
&= \frac{xt_1 u_1 \{1 + (s_{1,\infty} + 2x(1-t_1)(1-u_1))s_{1,\infty}\}}{2(1-x(1-t_1)(1-u_1))^2 s_{1,\infty}} \\
&\quad - \frac{xt_1 u_1 (1 + s_{1,\infty}) \sqrt{s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2(1-x(1-t_1)(1-u_1))^2 s_{1,\infty}}
\end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (5.2.24),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} (1-x(1-t_2)(1-u_2)v_2)^{-(8+\varphi+2\lambda)} \\
&\times \sum_{q_2=q_1}^{\infty} \left(\frac{v_2-1}{v_2} \frac{s_{2,\infty}}{1-x(1-t_2)(1-u_2)v_2} \right)^{q_2} \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial \overleftrightarrow{w}_{2,2} \right) \overleftrightarrow{w}_{2,2}^{2+\alpha+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-(4+\varphi+2\lambda)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_1}{1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \overleftrightarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,2} \partial \overleftrightarrow{w}_{1,2} \right) \overleftrightarrow{w}_{1,2}^{\alpha+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right) z^2 \tag{5.2.48}
\end{aligned}$$

Replace q_i , q_j and r_i by q_2 , q_1 and $\frac{v_2-1}{v_2} \frac{s_{2,\infty}}{1-x(1-t_2)(1-u_2)v_2}$ in (5.2.41). Take the new (5.2.41) into (5.2.48).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{(1-x(1-t_2)(1-u_2)v_2)^{-(7+\varphi+2\lambda)}}{-x(1-t_2)(1-u_2)v_2^2 + (1-s_{2,\infty})v_2 + s_{2,\infty}} \\
&\times \overleftrightarrow{w}_{2,2}^{-(2+\alpha+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial \overleftrightarrow{w}_{2,2} \right) \overleftrightarrow{w}_{2,2}^{2+\alpha+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-(4+\varphi+2\lambda)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_2-1}{v_2} \frac{s_{1,\infty}}{1-x(1-t_2)(1-u_2)v_2} \frac{v_1-1}{v_1} \frac{1}{1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
&\times \overleftrightarrow{w}_{1,2}^{-(\alpha+\lambda)} \left(\overleftrightarrow{w}_{1,2} \partial \overleftrightarrow{w}_{1,2} \right) \overleftrightarrow{w}_{1,2}^{\alpha+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right) z^2 \tag{5.2.49}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$v_2 = \frac{1 - s_{2,\infty} - \sqrt{(1 - s_{2,\infty})^2 + 4x(1 - t_2)(1 - u_2)s_{2,\infty}}}{2x(1 - t_2)(1 - u_2)}$$

or $\frac{1 - s_{2,\infty} + \sqrt{(1 - s_{2,\infty})^2 + 4x(1 - t_2)(1 - u_2)s_{2,\infty}}}{2x(1 - t_2)(1 - u_2)}$ and $\frac{1 - s_{2,\infty} - \sqrt{(1 - s_{2,\infty})^2 + 4x(1 - t_2)(1 - u_2)s_{2,\infty}}}{2x(1 - t_2)(1 - u_2)}$ is only inside the unit circle. As we compute the residue there in (5.2.49) we obtain

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\ &= \prod_{k=3}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\ & \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+\varphi+2\lambda)} \\ & \times \tilde{w}_{2,2}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2+\alpha+\lambda} \\ & \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)v_1)^{-(4+\varphi+2\lambda)} \\ & \times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1 - 1}{v_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \ddot{w}_{1,2}^{-(\alpha+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\alpha+\lambda} \\ & \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) z^2 \end{aligned} \quad (5.2.50)$$

where

$$\begin{aligned} \tilde{w}_{2,2} &= \frac{v_2}{(v_2 - 1)} \frac{xt_2u_2}{1 - xv_2(1 - t_2)(1 - u_2)} \Bigg|_{v_2 = \frac{1 - s_{2,\infty} - \sqrt{(1 - s_{2,\infty})^2 + 4x(1 - t_2)(1 - u_2)s_{2,\infty}}}{2x(1 - t_2)(1 - u_2)}} \\ &= \frac{xt_2u_2 \{1 + (s_{2,\infty} + 2x(1 - t_2)(1 - u_2))s_{2,\infty}\}}{2(1 - x(1 - t_2)(1 - u_2))^2 s_{2,\infty}} \\ & \quad - \frac{xt_2u_2(1 + s_{2,\infty})\sqrt{s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1}}{2(1 - x(1 - t_2)(1 - u_2))^2 s_{2,\infty}} \end{aligned}$$

and

$$\ddot{w}_{1,2} = \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2} v_1 (1 - t_1)(1 - u_1)}$$

Replace q_i, q_j and r_i by q_1, q_0 and $\frac{v_1 - 1}{v_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)v_1}$ in (5.2.41). Take the new (5.2.41) into (5.2.50).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+\varphi+2\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2+\alpha+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{(1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)v_1)^{-(3+\varphi+2\lambda)}}{-\tilde{w}_{2,2}(1 - t_1)(1 - u_1)v_1^2 + (1 - s_1)v_1 + s_1} \\
&\quad \times \ddot{w}_{1,2}^{-(\alpha+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\alpha+\lambda} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} \left(\frac{v_1 - 1}{v_1} \frac{s_{0,1}}{1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)v_1} \right)^{q_0} \\
&\quad \times \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\gamma + \lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) z^2 \tag{5.2.51}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$\begin{aligned}
v_1 &= \frac{1 - s_1 - \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,2}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,2}(1 - t_1)(1 - u_1)} \\
&\text{or } \frac{1 - s_1 + \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,2}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,2}(1 - t_1)(1 - u_1)} \text{ and} \\
&\frac{1 - s_1 - \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,2}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,2}(1 - t_1)(1 - u_1)} \text{ is only inside the unit circle. As we}
\end{aligned}$$

compute the residue there in (5.2.51) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_{2,\infty}^2 - 2(1-2x(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2x(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+\varphi+2\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2+\alpha+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-(3+\varphi+2\lambda)} \\
&\quad \times \tilde{w}_{1,2}^{-(\alpha+\lambda)} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\alpha+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \tilde{w}_{1,2}^{i_0} \right) z^2
\end{aligned} \tag{5.2.52}$$

where

$$\begin{aligned}
\tilde{w}_{1,2} &= \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2} v_1 (1 - t_1)(1 - u_1)} \Bigg|_{v_1 = \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)}} \\
&= \frac{\tilde{w}_{2,2} t_1 u_1 \{1 + (s_1 + 2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1\}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1} \\
&\quad - \frac{\tilde{w}_{2,2} t_1 u_1 (1 + s_1) \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1}
\end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (5.2.25),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_3(x) \\
&= \prod_{k=4}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\gamma+\lambda} (s_{3,\infty}^2 - 2(1-2x(1-t_3)(1-u_3))s_{3,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{3,\infty} + \sqrt{s_{3,\infty}^2 - 2(1-2x(1-t_3)(1-u_3))s_{3,\infty} + 1}}{2} \right)^{-(11+\varphi+2\lambda)} \\
&\quad \times \tilde{w}_{3,3}^{-(4+\alpha+\lambda)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{4+\alpha+\lambda} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\gamma+\lambda} (s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_2 + \sqrt{s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1}}{2} \right)^{-(7+\varphi+2\lambda)} \tilde{w}_{2,3}^{-(2+\alpha+\lambda)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{2+\alpha+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\gamma+\lambda} (s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-(3+\varphi+2\lambda)} \tilde{w}_{1,3}^{-(\alpha+\lambda)} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\alpha+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 x^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + \varphi + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\gamma + \lambda)_{i_0}} \tilde{w}_{1,3}^{i_0} \right) z^3 \tag{5.2.53}
\end{aligned}$$

where

$$\begin{aligned}
\tilde{w}_{3,3} &= \frac{v_3}{(v_3 - 1)} \frac{xt_3 u_3}{1 - x(1-t_3)(1-u_3)v_3} \Big|_{v_3 = \frac{1-s_{3,\infty} - \sqrt{(1-s_{3,\infty})^2 + 4x(1-t_3)(1-u_3)s_{3,\infty}}}{2x(1-t_3)(1-u_3)}} \\
&= \frac{xt_3 u_3 \{1 + (s_{3,\infty} + 2x(1-t_3)(1-u_3))s_{3,\infty}\}}{2(1-x(1-t_3)(1-u_3))^2 s_{3,\infty}} \\
&\quad - \frac{xt_3 u_3 (1 + s_{3,\infty}) \sqrt{s_{3,\infty}^2 - 2(1-2x(1-t_3)(1-u_3))s_{3,\infty} + 1}}{2(1-x(1-t_3)(1-u_3))^2 s_{3,\infty}}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{2,3} &= \frac{v_2}{(v_2 - 1)} \frac{\tilde{w}_{3,3} t_2 u_2}{1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2)v_2} \Bigg|_{v_2 = \frac{1-s_2 - \sqrt{(1-s_2)^2 + 4\tilde{w}_{3,3}(1-t_2)(1-u_2)s_2}}{2\tilde{w}_{3,3}(1-t_2)(1-u_2)}} \\
&= \frac{\tilde{w}_{3,3} t_2 u_2 \{1 + (s_2 + 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2\}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2} \\
&\quad - \frac{\tilde{w}_{3,3} t_2 u_2 (1 + s_2) \sqrt{s_2^2 - 2(1 - 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2 + 1}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{1,3} &= \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,3} t_1 u_1}{1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1)v_1} \Bigg|_{v_1 = \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,3}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,3}(1-t_1)(1-u_1)}} \\
&= \frac{\tilde{w}_{2,3} t_1 u_1 \{1 + (s_1 + 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1\}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1} \\
&\quad - \frac{\tilde{w}_{2,3} t_1 u_1 (1 + s_1) \sqrt{s_1^2 - 2(1 - 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1 + 1}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1}
\end{aligned}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$

terms where $m \geq 4$, I obtain every $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(x)$ terms.

Substitute (5.2.44), (5.2.47), (5.2.52), (5.2.53) and including all

$$\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(x) \text{ terms where } m > 3 \text{ into (5.2.43).} \quad \blacksquare$$

Remark 5.2.14 *The generating function for the CHP of type 2 of the first kind about*

$x = 0$ as $q = (q_j + 2j)(-\beta + \gamma + \delta - 1 + q_j + 2j)$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} H_c^{(a)} F_{q_j}^R(\alpha, \beta, \gamma, \delta, q = (q_j + 2j)(q_j + 2j + \varphi) \\
& ; \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
& = 2^{\varphi-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; x) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\gamma} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, x) \right. \\
& \times \tilde{w}_{1,1}^{-\alpha}(\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, x) \right. \\
& \times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)}(\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\alpha} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)}(\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\alpha} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Bigg\} \quad (5.2.54)
\end{aligned}$$

where

$$\begin{aligned}
& \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, x) = \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-(3+\varphi)}}{\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}} \\
& \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, x) = \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-(4n-1+\varphi)}}{\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}} \\
& \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\
& = \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)-1+\varphi)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}
\end{aligned}$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; x) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{\gamma-\varphi}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{\gamma-1} \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{\gamma-1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{\gamma-1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Proof Replace $\mathbf{A}$ and w by φ and $s_{0,\infty}$ in (5.2.39).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_{0,\infty}^{q_0} {}_2F_1(-q_0, q_0 + \varphi; \gamma; x) \\ &= 2^{\varphi-1} \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{\gamma-\varphi}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{\gamma-1} \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}} \end{aligned} \quad (5.2.55)$$

Replace $\mathbf{A}$, w and x by φ , s_0 and $\tilde{w}_{1,1}$ in (5.2.39).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1(-q_0, q_0 + \varphi; \gamma; \tilde{w}_{1,1}) \\ &= 2^{\varphi-1} \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{\gamma-1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (5.2.56)$$

Replace $\mathbf{A}$, w and x by φ , s_0 and $\tilde{w}_{1,n}$ in (5.2.39).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1(-q_0, q_0 + \varphi; \gamma; \tilde{w}_{1,n}) \\ &= 2^{\varphi-1} \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{\gamma-1} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (5.2.57)$$

Put $c_0 = 1$, $\lambda = 0$ and $\gamma' = \gamma$ in (5.2.42). Substitute (5.2.55), (5.2.56) and (5.2.57) into the new (5.2.42). ■

Remark 5.2.15 *The generating function for the CHP of type 2 of the second kind about $x = 0$ as $q = (q_j + 2j + 1 - \gamma)(-\beta + \delta + q_j + 2j)$ where $j, q_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} H_c^{(a)} S_{\alpha_j}^R(\alpha, \beta, \gamma, \delta \\
& , q = (q_j + 2j + 1 - \gamma)(q_j + 2j + 1 - \gamma + \varphi); \varphi = -\beta + \gamma + \delta - 1; z = \beta x^2) \\
& = \frac{2^{\varphi+1-2\gamma}}{x^{\gamma-1}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{B}(s_{0,\infty}; x) \right. \\
& + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{2-\gamma} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, x) \\
& \times \tilde{w}_{1,1}^{-(\alpha-\gamma+1)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\gamma+1} \mathbf{B}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\gamma} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, x) \right. \\
& \times \tilde{w}_{n,n}^{-(2n-1+\alpha-\gamma)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\alpha-\gamma} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\alpha-\gamma} \left. \right\} \mathbf{B}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \quad (5.2.58)
\end{aligned}$$

where

$$\left\{ \begin{aligned}
& \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, x) = \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-(5-2\gamma+\varphi)}}{\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}} \\
& \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, x) = \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-(4n+1-2\gamma+\varphi)}}{\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}} \\
& \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\
& = \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)+1-2\gamma+\varphi)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}
\end{aligned} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{B}(s_{0,\infty}; x) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{-2+3\gamma-\varphi}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{1-\gamma} \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{-2+3\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{-2+3\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Proof Replace γ , A and w by $2 - \gamma$, $\varphi + 2(1 - \gamma)$ and $s_{0,\infty}$ in (5.2.39).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_{0,\infty}^{q_0} {}_2F_1(-q_0, q_0 + \varphi + 2(1-\gamma); 2-\gamma; x) \\ &= 2^{\varphi+1-2\gamma} \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{-2+3\gamma-\varphi}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}\right)^{1-\gamma} \sqrt{s_{0,\infty}^2 - 2(1-2x)s_{0,\infty} + 1}} \end{aligned} \quad (5.2.59)$$

Replace γ , A , w and x by $2 - \gamma$, $\varphi + 2(1 - \gamma)$, s_0 and $\tilde{w}_{1,1}$ in (5.2.39).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1(-q_0, q_0 + \varphi + 2(1-\gamma); 2-\gamma; \tilde{w}_{1,1}) \\ &= 2^{\varphi+1-2\gamma} \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{-2+3\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}\right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (5.2.60)$$

Replace γ , A , w and x by $2 - \gamma$, $\varphi + 2(1 - \gamma)$, s_0 and $\tilde{w}_{1,n}$ in (5.2.39).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} {}_2F_1(-q_0, q_0 + \varphi + 2(1-\gamma); 2-\gamma; \tilde{w}_{1,n}) \\ &= 2^{\varphi+1-2\gamma} \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{-2+3\gamma-\varphi}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}\right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (5.2.61)$$

Put $c_0 = 1$, $\lambda = 1 - \gamma$ and $\gamma' = 2 - \gamma$ in (5.2.42). Substitute (5.2.59), (5.2.60) and (5.2.61) into the new (5.2.42). $\blacksquare$

5.3 Summary

The second independent solutions of the CHF by using 3TRF and R3TRF are only available for non-integer value of γ in the domain $|x| < 1$. If γ is an integer, one of two solutions will be dependent. Its analytic solution is defined by its behavior at infinity in the domain $|x| > 1$. All possible local solutions of the CHE (Regge-Wheeler and Teukolsky equations) were constructed by Fiziev.[11, 12] The power series representation of the CHE about the singular point at infinity is not convergent by only asymptotic [2, 13]: Its solution will be constructed in the future paper for (1) the power series expansion in closed forms of the CHF about regular singularity $x = \infty$, (2) its integral representation and (3) the generating functions for type 3 CHP.

In chapter 4 I show how to obtain the power series expansion in closed forms and its integral forms of the CHE for infinite series and polynomial of type 1 including all higher terms of A_n 's by applying 3TRF. This was done by letting A_n in sequence c_n is the leading term in the analytic function $y(x)$: the sequence c_n consists of combinations A_n and B_n . For polynomial of type 1, I treat β , γ , δ and q as free variables and a fixed value of α .

In this chapter I show how to construct the power series expansion in closed forms and its integral forms of the CHE for infinite series and polynomial of type 2 including all higher terms of B_n 's by applying R3TRF. This is done by letting B_n in sequence c_n is the leading term in the analytic function $y(x)$. For polynomial of type 2, I treat α , β , γ and δ as free variables and a fixed value of q .

The power series expansion and integral forms of the CHE for infinite series about $x = 0$ in this chapter are equivalent to infinite series of the CHE in chapter 4. In this chapter B_n is the leading term in sequence c_n in the analytic function $y(x)$. In chapter 4 A_n is the leading term in sequence c_n in the analytic function $y(x)$.

In chapter 4 as we see the power series expansions of the CHE about $x = 0$ for infinite series and polynomial of type 1, the denominators and numerators in all B_n terms of $y(x)$ arise with Pochhammer symbol. Again in this chapter the denominators and numerators in all A_n terms of the CHF about $x = 0$ arise with Pochhammer symbol. Since we construct the power series expansion with Pochhammer symbols in numerators and denominators, we are able to describe integral forms of the CHF in the mathematical rigor. As we observe representations in closed form integrals of the CHF by applying 3TRF, a ${}_1F_1$ function recurs in each of sub-integral forms of the CHF. Also ${}_2F_1$ function recurs in each of sub-integral forms of the CHF as we observe integral form of the CHF by applying R3TRF.

We can transform the CHF into any special functions having two recursive coefficients because of ${}_1F_1$ and ${}_2F_1$ functions in each of sub-integral forms of the CHF. After we replace ${}_1F_1$ and ${}_2F_1$ functions in integral forms of the CHF to other special functions, we are able to rebuild the power series expansion of the CHF in a backward.

In chapter 4 I show how to analyze the generating function for the CHP of type 1 from its general integral representation. And in this chapter I construct the generating function for the CHP of type 2 from its integral form. We are able to derive orthogonal relations, recursion relations and expectation values of physical quantities from these two generating functions compared with the case of a normalized wave function for the hydrogen-like atoms.¹⁰

Mathematical structure of the generating function for the CHP of type 2 closely resembles the generating function for the Heun polynomial of type 2 in chapter 3. If $a \rightarrow 0$, $\eta \rightarrow x$, $\alpha = \beta$, $z = -\frac{1}{a}x^2 \rightarrow \beta x^2$, $\varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1) \rightarrow -\beta + \gamma + \delta - 1$, $q = -(q_j + 2j + \lambda)\{\varphi + (1 + a)(q_j + 2j + \lambda)\} \rightarrow (q_j + 2j + \lambda)(\varphi + q_j + 2j + \lambda)$, $(\tilde{w}_{1,1}\partial_{\tilde{w}_{1,1}})^2 \rightarrow \tilde{w}_{1,1}\partial_{\tilde{w}_{1,1}}$, $(\tilde{w}_{n,n}\partial_{\tilde{w}_{n,n}})^2 \rightarrow \tilde{w}_{n,n}\partial_{\tilde{w}_{n,n}}$ and $(\tilde{w}_{n-k,n}\partial_{\tilde{w}_{n-k,n}})^2 \rightarrow \tilde{w}_{n-k,n}\partial_{\tilde{w}_{n-k,n}}$ in the general expression of the generating function for the Heun polynomial of type 2, its solution is equivalent to the generating function for the CHP of type 2; compare theorem 3.2.4 in chapter 3 with theorem 5.2.12 in this chapter. Because I derive the both generating functions for the CHP and Heun polynomial of type 2 by applying the generating function for the Jacobi polynomial using ${}_2F_1$ functions into integral forms of the CHP and Heun polynomial: ${}_2F_1$ function recurs in each of sub-integral forms of them. By similar reason, mathematical structure of the generating function for the CHP of type 1 closely resembles the generating function for the GCH polynomial of type 1 [37]. The generating function for confluent hypergeometric polynomial is applied into the integral representations of the CHP and GCH polynomial: ${}_1F_1$ function recurs in each of sub-integral forms of them.

¹⁰The normalized constant for this wave function is obtained by applying the generating function for associated Laguerre polynomials into the Helmholtz equation in the spherical coordinates. It only has one eigenvalue. And the CHP of either type 1 or 2 has infinite eigenvalues because of three different coefficients in its power series.

Bibliography

- [1] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806.
- [2] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische. Annalen.* **33**, 161(1889).
- [3] Ronveaux, A., “Heuns Differential Equations” Oxford University Press, (1995).
- [4] Choun, Y.S., “Analytic solution for grand confluent hypergeometric function,” arXiv:1303.0813.
- [5] Choun, Y.S., “The integral formalism and the generating function of grand confluent hypergeometric function,” arXiv:1303.0819.
- [6] Leaver, E.W., “Solutions to a generalized spheroidal wave equation: Teukolsky’s equations in general relativity, and the two-center problem in molecular quantum mechanics,” *J. Math. Phys.* **27**, 1238–1265(1986).
- [7] Decarreau, A., Maroni, P. and Robert, A., “Sur les équations confluentes de l’équation de Heun,” *Ann. Soc. Sci. Bruxelles.* **T92(III)**, 151–189 (1978).
- [8] Decarreau, A., Dumont-Lepage, M. C., Maroni, P., Robert, A. and Ronveaux, A., “Formes canoniques des équations confluentes de l’équation de Heun,” *Ann. Soc. Sci. Bruxelles.* **T92(I-II)**, 53–78 (1978).
- [9] Wilson, A. H., “A generalised spheroidal wave equation,” *Proc. Roy. Soc. London* **A118**, 61–635 (1928).
- [10] Wilson, A. H., “The ionised hydrogen molecule,” *Proc. Roy. Soc. London* **A118**, 635–647 (1928).
- [11] Fiziev, P.P., “Classes of exact solutions to Regge-Wheeler and Teukolsky equations,” arXiv:0902.1277.

- [12] Fiziev, P.P., “Classes of exact solutions to the Teukolsky master equations,” *Class. Quantum. Grav.* **27**, 135001 (2010).
- [13] Fiziev, P.P., “Exact solutions of Teukolsky master equation with continuous spectrum,” arXiv:0903.3617.
- [14] Lambe, C.G. and Ward, D.R., “Some differential equations and associated integral equations,” *Q. J. Math. (Oxford), Ser.* **13**, 107-112 (1942).
- [15] Erdélyi, A., “Integral equations for Heun functions,” *Q.J. Math. (Oxford), Ser.* **13**, 107–112 (1942).
- [16] Sleeman, B.D., “Non-linear integral equations for Heun functions,” *Proc. Edinburgh Math. Soc.*, Vol. **16**, 281–289 (1969).
- [17] Abramov, D.I. and Komarov, I.V., “Integral equations and relations for the Coulomb spheroidal functions,” *Sov. J. Teor. Mat. Phys.* Vol. **29**, 234–243 (1976).
- [18] Schmidt, D. and Wolf, G., “A method of generating integral relations by simultaneous separability of generalized Schrödinger equation,” *SIAM J. Math. Anal.* Vol. **10**, 823–838 (1979).
- [19] Kazakov, A.Ya., “The Laplace transformation and the differential equations with polynomial coefficients,” *Sov. J. Differ. Uravn.* Vol. **22**, 354–356 (1986).

Chapter 6

Grand Confluent Hypergeometric function using reversible three-term recurrence formula

Biconfluent Heun (BCH) equation, a confluent form of Heun equation[1, 2], is the special case of Grand Confluent Hypergeometric (GCH) equation¹: this has a regular singularity at $x = 0$, and an irregular singularity at ∞ of rank 2.

In this chapter I will apply reversible three term recurrence formula (R3TRF) in chapter 1 to (1) the power series expansion in closed forms, (2) its integral forms of the GCH equation for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's² and (3) the generating function for the GCH polynomial which makes A_n term terminated.

6.1 Introduction

$$x \frac{d^2 y}{dx^2} + (\mu x^2 + \varepsilon x + \nu) \frac{dy}{dx} + (\Omega x + \varepsilon \omega) y = 0 \quad (6.1.1)$$

(6.1.1) is Grand Confluent Hypergeometric (GCH) differential equation where $\mu, \varepsilon, \nu, \Omega$ and ω are real or complex parameters [34, 36]. GCH ordinary differential equation is of

¹For the canonical form of the BCH equation[2], replace $\mu, \varepsilon, \nu, \Omega$ and ω by $-2, -\beta, 1 + \alpha, \gamma - \alpha - 2$ and $1/2(\delta/\beta + 1 + \alpha)$ in (6.1.1). For DLFM version [19] or in Ref.[20], replace μ and ω by 1 and $-q/\varepsilon$ in (6.1.1).

²“higher terms of B_n 's” means at least two terms of B_n 's.

Fuchsian types with two singular points: one regular singular point which is zero with exponents $\{0, 1 - \nu\}$, and one irregular singular point which is infinity with an exponent $\frac{\Omega}{\mu}$. In contrast, Heun equation of Fuchsian types has the four regular singularities. Heun equation has the four kind of confluent forms: (1) Confluent Heun (two regular and one irregular singularities), (2) Doubly confluent Heun (two irregular singularities), (3) Biconfluent Heun (one regular and one irregular singularities), (4) Triconfluent Heun equations (one irregular singularity).

BCH equation is derived from the GCH equation by changing all coefficients.[2] The GCH (or BCH) equation is applicable into the modern physics. [3, 4, 5, 6, 7] The BCH equation appears in the radial Schrödinger equation with those potentials such as the rotating harmonic oscillator [8], the doubly anharmonic oscillator [9, 10, 11], a three-dimensional anharmonic oscillator [12, 13, 11], Coulomb potential with a linear confining potential [11, 14] and other kinds of potentials [15, 16].

The fundamental solution of the BCH equation for infinite series and the BCH spectral polynomial about $x = 0$ in the canonical form was obtained by applying the power series expansion.[21, 22, 23, 24] For the case of the irregular singular point $x = \infty$, the three term recurrence of the power series in the BCH equation was derived.[17, 25] And they left the analytic solution of the BCH equation as solutions of recurrences because of three different coefficients in its power series expansion of the BCH equation.³ In comparison with the two term recursion relation of the power series in the linear differential equation, the analytic solution in closed forms on the three term recurrence relation of the power series is unknown currently because of its complex mathematical calculation.

As I know, there are no examples for the analytic solution of the BCH equation about $x = 0$ and $x = \infty$ in the form of a definite or contour integral containing the well-known special functions which consists of two term recursion relation in its power series of the linear differential equation such as a ${}_2F_1$ or ${}_1F_1$ function. In place of describing the integral representation of the BCH equation involving only simple functions, especially for confluent hypergeometric functions, the BCH equation is obtained by Fredholm-type integral equations; such integral relationships express one analytic solution in terms of another analytic solution.[27, 28, 29, 30, 31, 32, 33]

In Ref.[36, 37] I construct analytic solutions of GCH equation about the regular singular point at $x = 0$ by applying three term recurrence formula (3TRF). [35]; (1) the power series expansion in closed forms, (2) its integral forms of the GCH equation (for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's⁴) and (3) the generating function for GCH polynomial which makes B_n term

³For the special case, the explicit solutions of the BCH equation in the canonical form was constructed when one of coefficients $\beta = 0$. [26]

⁴“ higher terms of A_n 's” means at least two terms of A_n 's.

terminated. And I show four examples of the analytic wave functions and its eigenvalues in the radial Schrödinger equation with certain potentials: (1) Schrödinger equation with the rotating harmonic oscillator and a class of confinement potentials, (2) The spin free Hamiltonian involving only scalar potential for the $q - \bar{q}$ system, (3) The radial Schrödinger equation with Confinement potentials, (4) Two interacting electrons in a uniform magnetic field and a parabolic potential. The Frobenius solutions in closed forms and its combined definite and contour integrals of these four quantum mechanical wave functions are derived analytically.

In this chapter, by applying R3TRF in chapter 1, I construct the power series expansion in closed forms of GCH equation about the regular singular point at $x = 0$ for infinite series and polynomial which makes A_n term terminated. The integral forms of GCH equation and its generating function for the GCH polynomial which makes A_n term terminated are derived analytically. Also the Frobenius solution of GCH equation about irregular singular point at $x = \infty$ by applying 3TRF [35] is constructed analytically including its integral representation and the generating function for the GCH polynomial which makes B_n term terminated.

6.2 GCH equation about regular singular point at zero

Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} \quad (6.2.1)$$

where λ is an indicial root. Plug (6.2.1) into (6.1.1). We obtain a three-term recurrence relation for the coefficients c_n :

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (6.2.2)$$

where,

$$A_n = \frac{-\varepsilon(n + \omega + \lambda)}{(n + 1 + \lambda)(n + \nu + \lambda)} \quad (6.2.3a)$$

$$B_n = -\frac{\Omega + \mu(n - 1 + \lambda)}{(n + 1 + \lambda)(n + \nu + \lambda)} \quad (6.2.3a)$$

$$c_1 = A_0 c_0 \quad (6.2.3b)$$

We have two indicial roots which are $\lambda = 0$ and $1 - \nu$.

6.2.1 Power series

Polynomial of type 2

There are two types of power series expansion using the two term recurrence relation in a linear ordinary differential equation which are a polynomial and an infinite series. In contrast there are three types of polynomials and an infinite series in three term recurrence relation of linear ordinary differential equation: (1) polynomial which makes B_n term terminated: A_n term is not terminated, (2) polynomial which makes A_n term terminated: B_n term is not terminated, (3) polynomial which makes A_n and B_n terms terminated at the same time.⁵ In general the GCH polynomial is defined as type 3 polynomial where A_n and B_n terms terminated. For the type 3 GCH polynomial about $x = 0$, it has a fixed integer value of Ω , just as it has a fixed value of ω . In three term recurrence relation, polynomial of type 3 I categorize as complete polynomial. In the previous paper [36, 37] I construct the type 1 GCH polynomial about $x = 0$. In future papers I will derive type 3 polynomial about $x = 0$. In this chapter I construct the power series expansion, its integral forms and the generating function for the GCH polynomial of type 2 about $x = 0$: I treat μ, ε, ν and Ω as free variables and ω as a fixed value.

In chapter 1 the general expression of power series of $y(x)$ for polynomial of type 2 is defined by

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 \left\{ \sum_{i_0=0}^{\alpha_0} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
 &\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\alpha_k} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
 &\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\alpha_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Big\} \quad (6.2.4)
 \end{aligned}$$

In the above, $\alpha_i \leq \alpha_j$ only if $i \leq j$ where $i, j, \alpha_i, \alpha_j \in \mathbb{N}_0$.

For a polynomial, we need a condition which is:

$$A_{\alpha_i+2i} = 0 \quad \text{where } i, \alpha_i = 0, 1, 2, \dots \quad (6.2.5)$$

⁵If A_n and B_n terms are not terminated, it turns to be infinite series.

In the above, α_i is an eigenvalue that makes A_n term terminated at certain value of index n . (6.2.5) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (6.2.4). In (6.2.3a)-(6.2.3b) replace ω by $-(\omega_i + 2i + \lambda)$. In (6.2.5) replace index α_i by ω_i . Take the new (6.2.3a)-(6.2.3b), (6.2.5) and put them in (6.2.4). After the replacement process, the general expression of power series of the GCH equation for polynomial of type 2 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \sum_{i_1=i_0}^{\omega_1} \frac{(-\omega_1)_{i_1}(3+\lambda)_{i_0}(2+\nu+\lambda)_{i_0}}{(-\omega_1)_{i_0}(3+\lambda)_{i_1}(2+\nu+\lambda)_{i_1}} \eta^{i_1} \right\} \rho \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\omega_k} \frac{(i_k + 2k + \Omega/\mu + \lambda)}{(i_k + 2k + 2 + \lambda)(i_k + 2k + 1 + \nu + \lambda)} \frac{(-\omega_k)_{i_k}(2k+1+\lambda)_{i_{k-1}}(2k+\nu+\lambda)_{i_{k-1}}}{(-\omega_k)_{i_{k-1}}(2k+1+\lambda)_{i_k}(2k+\nu+\lambda)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\omega_n} \frac{(-\omega_n)_{i_n}(2n+1+\lambda)_{i_{n-1}}(2n+\nu+\lambda)_{i_{n-1}}}{(-\omega_n)_{i_{n-1}}(2n+1+\lambda)_{i_n}(2n+\nu+\lambda)_{i_n}} \eta^{i_n} \right\} \rho^n \Bigg\} \quad (6.2.6)
\end{aligned}$$

where

$$\begin{cases} \eta = -\varepsilon x \\ \rho = -\mu x^2 \\ \omega = -(\omega_j + 2j + \lambda) \text{ as } j, \omega_j \in \mathbb{N}_0 \\ \omega_i \leq \omega_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the GCH equation and $\lambda = 1 - \nu$ for the second one in (6.2.6).

Remark 6.2.1 *The power series expansion of the GCH equation of the first kind for*

polynomial of type 2 about $x = 0$ as $\omega = -(\omega_j + 2j)$ where $j, \omega_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= QW_{\omega_j}^R(\mu, \varepsilon, \nu, \Omega, \omega = -(\omega_j + 2j); \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1)_{i_0}(\nu)_{i_0}} \eta^{i_0} + \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu)}{(i_0 + 2)(i_0 + 1 + \nu)} \frac{(-\omega_0)_{i_0}}{(1)_{i_0}(\nu)_{i_0}} \sum_{i_1=i_0}^{\omega_1} \frac{(-\omega_1)_{i_1}(3)_{i_0}(2 + \nu)_{i_0}}{(-\omega_1)_{i_0}(3)_{i_1}(2 + \nu)_{i_1}} \eta^{i_1} \right\} \rho \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu)}{(i_0 + 2)(i_0 + 1 + \nu)} \frac{(-\omega_0)_{i_0}}{(1)_{i_0}(\nu)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\omega_k} \frac{(i_k + 2k + \Omega/\mu)}{(i_k + 2k + 2)(i_k + 2k + 1 + \nu)} \frac{(-\omega_k)_{i_k}(2k + 1)_{i_{k-1}}(2k + \nu)_{i_{k-1}}}{(-\omega_k)_{i_{k-1}}(2k + 1)_{i_k}(2k + \nu)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\omega_n} \frac{(-\omega_n)_{i_n}(2n + 1)_{i_{n-1}}(2n + \nu)_{i_{n-1}}}{(-\omega_n)_{i_{n-1}}(2n + 1)_{i_n}(2n + \nu)_{i_n}} \eta^{i_n} \right\} \rho^n
\end{aligned} \tag{6.2.7}$$

Remark 6.2.2 The power series expansion of the GCH equation of the second kind for polynomial of type 2 about $x = 0$ as $\omega = -(\omega_j + 2j + 1 - \nu)$ where $j, \omega_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= RW_{\omega_j}^R(\mu, \varepsilon, \nu, \Omega, \omega = -(\omega_j + 2j + 1 - \nu); \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= x^{1-\nu} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(2 - \nu)_{i_0}(1)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + 1 + \Omega/\mu - \nu)}{(i_0 + 3 - \nu)(i_0 + 2)} \frac{(-\omega_0)_{i_0}}{(2 - \nu)_{i_0}(1)_{i_0}} \sum_{i_1=i_0}^{\omega_1} \frac{(-\omega_1)_{i_1}(4 - \nu)_{i_0}(3)_{i_0}}{(-\omega_1)_{i_0}(4 - \nu)_{i_1}(3)_{i_1}} \eta^{i_1} \right\} \rho \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + 1 + \Omega/\mu - \nu)}{(i_0 + 3 - \nu)(i_0 + 2)} \frac{(-\omega_0)_{i_0}}{(2 - \nu)_{i_0}(1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\omega_k} \frac{(i_k + 2k + 1 + \Omega/\mu - \nu)}{(i_k + 2k + 3 - \nu)(i_k + 2k + 2)} \frac{(-\omega_k)_{i_k}(2k + 2 - \nu)_{i_{k-1}}(2k + 1)_{i_{k-1}}}{(-\omega_k)_{i_{k-1}}(2k + 2 - \nu)_{i_k}(2k + 1)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\omega_n} \frac{(-\omega_n)_{i_n}(2n + 2 - \nu)_{i_{n-1}}(2n + 1)_{i_{n-1}}}{(-\omega_n)_{i_{n-1}}(2n + 2 - \nu)_{i_n}(2n + 1)_{i_n}} \eta^{i_n} \right\} \rho^n \Big\}
\end{aligned} \tag{6.2.8}$$

In Ref.[36, 37] I treat Ω as a fixed value and $\mu, \varepsilon, \nu, \omega$ as free variables to construct the GCH polynomial of type 1: (1) if $\Omega = -\mu(2\beta_j + j)$ where $j, \beta_j \in \mathbb{N}_0$, an analytic solution of the GCH equation turns to be the first kind of independent solution of the GCH polynomial of type 1. (2) if $\Omega = -\mu(2\psi_j + j + 1 - \nu)$ where $j, \psi_j \in \mathbb{N}_0$, an analytic

solution of the GCH equation turns to be the second kind of independent solution of the GCH polynomial of type 1.

In this chapter I treat ω as a fixed value and $\mu, \varepsilon, \nu, \Omega$ as free variables to construct the GCH polynomial of type 2: (1) if $\omega = -(\omega_j + 2j)$ where $j, \omega_j \in \mathbb{N}_0$, an analytic solution of the GCH equation turns to be the first kind of independent solution of the GCH polynomial of type 2. (2) if $\omega = -(\omega_j + 2j + 1 - \nu)$, an analytic solution of the GCH equation turns to be the second kind of independent solution of the GCH polynomial of type 2.

Infinite series

In chapter 1 the general expression of power series of $y(x)$ for infinite series is defined by

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
 &\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
 &\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (6.2.9)
 \end{aligned}$$

Substitute (6.2.3a)–(6.2.3b) into (6.2.9). The general expression of power series of the GCH equation for infinite series about $x = 0$ is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\omega + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \nu + \lambda)_{i_0}}{(\omega + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \nu + \lambda)_{i_1}} \eta^{i_1} \right\} \rho \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \Omega/\mu + \lambda)}{(i_k + 2k + 2 + \lambda)(i_k + 2k + 1 + \nu + \lambda)} \right. \\
&\quad \times \left. \frac{(\omega + 2k + \lambda)_{i_k} (2k + 1 + \lambda)_{i_{k-1}} (2k + \nu + \lambda)_{i_{k-1}}}{(\omega + 2k + \lambda)_{i_{k-1}} (2k + 1 + \lambda)_{i_{k-1}} (2k + \nu + \lambda)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\omega + 2n + \lambda)_{i_n} (2n + 1 + \lambda)_{i_{n-1}} (2n + \nu + \lambda)_{i_{n-1}}}{(\omega + 2n + \lambda)_{i_{n-1}} (2n + 1 + \lambda)_{i_{n-1}} (2n + \nu + \lambda)_{i_n}} \eta^{i_n} \right\} \rho^n \Bigg\} \quad (6.2.10)
\end{aligned}$$

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the GCH equation and $\lambda = 1 - \nu$ for the second one in (6.2.10).

Remark 6.2.3 *The power series expansion of the GCH equation of the first kind for*

infinite series about $x = 0$ using $R3TRF$ is

$$\begin{aligned}
y(x) &= QW^R(\mu, \varepsilon, \nu, \Omega, \omega; \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= \sum_{i_0=0}^{\infty} \frac{(\omega)_{i_0}}{(1)_{i_0}(\nu)_{i_0}} \eta^{i_0} + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu)}{(i_0 + 2)(i_0 + 1 + \nu)} \frac{(\omega)_{i_0}}{(1)_{i_0}(\nu)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\omega + 2)_{i_1}(3)_{i_0}(2 + \nu)_{i_0}}{(\omega + 2)_{i_0}(3)_{i_1}(2 + \nu)_{i_1}} \eta^{i_1} \right\} \rho \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu)}{(i_0 + 2)(i_0 + 1 + \nu)} \frac{(\omega)_{i_0}}{(1)_{i_0}(\nu)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \Omega/\mu)}{(i_k + 2k + 2)(i_k + 2k + 1 + \nu)} \frac{(\omega + 2k)_{i_k}(2k + 1)_{i_{k-1}}(2k + \nu)_{i_{k-1}}}{(\omega + 2k)_{i_{k-1}}(2k + 1)_{i_{k-1}}(2k + \nu)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\omega + 2n)_{i_n}(2n + 1)_{i_{n-1}}(2n + \nu)_{i_{n-1}}}{(\omega + 2n)_{i_{n-1}}(2n + 1)_{i_{n-1}}(2n + \nu)_{i_n}} \eta^{i_n} \right\} \rho^n \tag{6.2.11}
\end{aligned}$$

Remark 6.2.4 The power series expansion of the GCH equation of the second kind for infinite series about $x = 0$ using $R3TRF$ is

$$\begin{aligned}
y(x) &= RW^R(\mu, \varepsilon, \nu, \Omega, \omega; \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= x^{1-\nu} \left\{ \sum_{i_0=0}^{\infty} \frac{(\omega + 1 - \nu)_{i_0}}{(2 - \nu)_{i_0}(1)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \Omega/\mu - \nu)}{(i_0 + 3 - \nu)(i_0 + 2)} \frac{(\omega + 1 - \nu)_{i_0}}{(2 - \nu)_{i_0}(1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\omega + 3 - \nu)_{i_1}(4 - \nu)_{i_0}(3)_{i_0}}{(\omega + 3 - \nu)_{i_0}(4 - \nu)_{i_1}(3)_{i_1}} \eta^{i_1} \right\} \rho \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \Omega/\mu - \nu)}{(i_0 + 3 - \nu)(i_0 + 2)} \frac{(\omega + 1 - \nu)_{i_0}}{(2 - \nu)_{i_0}(1)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + 1 + \Omega/\mu - \nu)}{(i_k + 2k + 3 - \nu)(i_k + 2k + 2)} \frac{(\omega + 2k + 1 - \nu)_{i_k}(2k + 2 - \nu)_{i_{k-1}}(2k + 1)_{i_{k-1}}}{(\omega + 2k + 1 - \nu)_{i_{k-1}}(2k + 2 - \nu)_{i_{k-1}}(2k + 1)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\omega + 2n + 1 - \nu)_{i_n}(2n + 2 - \nu)_{i_{n-1}}(2n + 1)_{i_{n-1}}}{(\omega + 2n + 1 - \nu)_{i_{n-1}}(2n + 2 - \nu)_{i_{n-1}}(2n + 1)_{i_n}} \eta^{i_n} \right\} \rho^n \left. \right\} \tag{6.2.12}
\end{aligned}$$

It is required that $\nu \neq 0, -1, -2, \dots$ for the first kind of independent solutions of the GCH equation for all cases. Because if it does not, its solution will be divergent. And it is required that $\nu \neq 2, 3, 4, \dots$ for the second kind of independent solutions of the GCH equation for all cases.

The infinite series in this chapter are equivalent to the infinite series in Ref.[36, 37]. In this chapter B_n is the leading term in sequence c_n of the analytic function $y(x)$. In Ref.[36, 37] A_n is the leading term in sequence c_n of the analytic function $y(x)$.⁶

6.2.2 Integral formalism

Polynomial of type 2

There is a generalized hypergeometric function which is

$$\begin{aligned} I_l &= \sum_{i_l=i_{l-1}}^{\omega_l} \frac{(-\omega_l)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\nu+\lambda)_{i_{l-1}}}{(-\omega_l)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\nu+\lambda)_{i_l}} \eta^{i_l} \\ &= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+2l+\lambda, j+1) B(i_{l-1}+2l-1+\nu+\lambda, j+1) (i_{l-1}-\omega_l)_j}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-1+\nu+\lambda)^{-1} (1)_j j!} \eta^j \end{aligned} \quad (6.2.13)$$

By using integral form of beta function,

$$B(i_{l-1}+2l+\lambda, j+1) = \int_0^1 dt_l t_l^{i_{l-1}+2l-1+\lambda} (1-t_l)^j \quad (6.2.14a)$$

$$B(i_{l-1}+2l-1+\nu+\lambda, j+1) = \int_0^1 du_l u_l^{i_{l-1}+2l-2+\nu+\lambda} (1-u_l)^j \quad (6.2.14b)$$

Substitute (6.2.14a) and (6.2.14b) into (6.2.13), and divide $(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\nu+\lambda)$ into I_l .

$$\begin{aligned} &\frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\nu+\lambda)} \sum_{i_l=i_{l-1}}^{\omega_l} \frac{(-\omega_l)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\nu+\lambda)_{i_{l-1}}}{(-\omega_l)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\nu+\lambda)_{i_l}} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-2+\nu+\lambda} (\eta t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(i_{l-1}-\omega_l)_j}{(1)_j j!} [\eta(1-t_l)(1-u_l)]^j \end{aligned} \quad (6.2.15)$$

The integral form of confluent hypergeometric function of the first kind is given by

$$\sum_{j=0}^{\infty} \frac{(-\alpha_0)_j}{(\gamma)_j j!} z^j = \frac{\Gamma(\alpha_0+1)\Gamma(\gamma)}{2\pi i \Gamma(\alpha_0+\gamma)} \oint dv_l \frac{\exp\left(-\frac{zv_l}{(1-v_l)}\right)}{v_l^{\alpha_0+1} (1-v_l)^\gamma} \quad (6.2.16)$$

⁶As $\frac{\Gamma(1/2+\nu/2-\Omega/(2\mu))}{\Gamma(1/2+\nu/2)}$ is multiplied on (6.2.11), the new (6.2.11) is equivalent to the first kind solution of the GCH equation for the infinite series using 3TRF.[36] Again, As $(-\mu/2)^{1/2(1-\nu)} \frac{\Gamma(1-\Omega/(2\mu))}{\Gamma(3/2-\nu/2)}$ is multiplied on (6.2.12), the new (6.2.12) corresponds to the second kind solution of the GCH equation for the infinite series using 3TRF.[36]

replace α_0 , γ and z by $\omega_l - i_{l-1}$, 1 and $\eta(1 - t_l)(1 - u_l)$ in (6.2.16).

$$\sum_{j=0}^{\infty} \frac{(i_{l-1} - \omega_l)_j}{(1)_j j!} [\eta(1 - t_l)(1 - u_l)]^j = \frac{1}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{v_l}{(1-v_l)} \eta(1 - t_l)(1 - u_l)\right)}{v_l^{\omega_l+1-i_{l-1}}(1 - v_l)} \quad (6.2.17)$$

Substitute (6.2.17) into (6.2.15).

$$\begin{aligned} K_l &= \frac{1}{(i_{l-1} + 2l + \lambda)(i_{l-1} + 2l - 1 + \nu + \lambda)} \sum_{i_l=i_{l-1}}^{\omega_l} \frac{(-\omega_l)_{i_l}(2l + 1 + \lambda)_{i_{l-1}}(2l + \nu + \lambda)_{i_{l-1}}}{(-\omega_l)_{i_{l-1}}(2l + 1 + \lambda)_{i_l}(2l + \nu + \lambda)_{i_l}} \eta^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{v_l}{(1-v_l)} \eta(1 - t_l)(1 - u_l)\right)}{v_l^{\omega_l+1}(1 - v_l)} (\eta t_l u_l v_l)^{i_{l-1}} \end{aligned} \quad (6.2.18)$$

Substitute (6.2.18) into (6.2.6) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(x)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(x)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(x)$, etc.⁷

Theorem 6.2.5 *The general representation in the form of integral of the GCH polynomial of type 2 is given by*

$$\begin{aligned} y(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \eta^{i_0} \right. \\ &\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\nu+\lambda} \right. \right. \\ &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})\right)}{v_{n-k}^{\omega_{n-k}+1}(1 - v_{n-k})} \\ &\quad \times w_{n-k,n}^{-(\Omega/\mu+2(n-k-1)+\lambda)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\Omega/\mu+2(n-k-1)+\lambda} \left. \right\} \\ &\quad \times \left. \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} w_{1,n}^{i_0} \right\} \rho^n \Big\} \end{aligned} \quad (6.2.19)$$

where

$$w_{a,b} = \begin{cases} \eta \prod_{l=a}^b t_l u_l v_l & \text{where } a \leq b \\ \eta & \text{only if } a > b \end{cases}$$

⁷ $y_1(x)$ means the sub-power series in (6.2.6) contains one term of $B'_n s$, $y_2(x)$ means the sub-power series in (6.2.6) contains two terms of $B'_n s$, $y_3(x)$ means the sub-power series in (6.2.6) contains three terms of $B'_n s$, etc.

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof In (6.2.6) the power series expansions of sub-summation $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of the GCH polynomial of type 2 are

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \quad (6.2.20)$$

where

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \eta^{i_0} \quad (6.2.21a)$$

$$\begin{aligned} y_1(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \right. \\ & \times \left. \sum_{i_1=i_0}^{\omega_1} \frac{(-\omega_1)_{i_1}(3+\lambda)_{i_0}(2+\nu+\lambda)_{i_0}}{(-\omega_1)_{i_0}(3+\lambda)_{i_1}(2+\nu+\lambda)_{i_1}} \eta^{i_1} \right\} \rho \end{aligned} \quad (6.2.21b)$$

$$\begin{aligned} y_2(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \right. \\ & \times \sum_{i_1=i_0}^{\omega_1} \frac{(i_1 + 2 + \Omega/\mu + \lambda)}{(i_1 + 4 + \lambda)(i_1 + 3 + \nu + \lambda)} \frac{(-\omega_1)_{i_1}(3+\lambda)_{i_0}(2+\nu+\lambda)_{i_0}}{(-\omega_1)_{i_0}(3+\lambda)_{i_1}(2+\nu+\lambda)_{i_1}} \\ & \times \left. \sum_{i_2=i_1}^{\omega_2} \frac{(-\omega_2)_{i_2}(5+\lambda)_{i_1}(4+\nu+\lambda)_{i_1}}{(-\omega_2)_{i_1}(5+\lambda)_{i_2}(4+\nu+\lambda)_{i_2}} \eta^{i_2} \right\} \rho^2 \end{aligned} \quad (6.2.21c)$$

$$\begin{aligned} y_3(x) = & c_0 x^\lambda \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \right. \\ & \times \sum_{i_1=i_0}^{\omega_1} \frac{(i_1 + 2 + \Omega/\mu + \lambda)}{(i_1 + 4 + \lambda)(i_1 + 3 + \nu + \lambda)} \frac{(-\omega_1)_{i_1}(3+\lambda)_{i_0}(2+\nu+\lambda)_{i_0}}{(-\omega_1)_{i_0}(3+\lambda)_{i_1}(2+\nu+\lambda)_{i_1}} \\ & \times \sum_{i_2=i_1}^{\omega_2} \frac{(i_2 + 4 + \Omega/\mu + \lambda)}{(i_2 + 6 + \lambda)(i_2 + 5 + \nu + \lambda)} \frac{(-\omega_2)_{i_2}(5+\lambda)_{i_1}(4+\nu+\lambda)_{i_1}}{(-\omega_2)_{i_1}(5+\lambda)_{i_2}(4+\nu+\lambda)_{i_2}} \\ & \times \left. \sum_{i_3=i_2}^{\omega_3} \frac{(-\omega_3)_{i_3}(7+\lambda)_{i_2}(6+\nu+\lambda)_{i_2}}{(-\omega_3)_{i_2}(7+\lambda)_{i_3}(6+\nu+\lambda)_{i_3}} \eta^{i_3} \right\} \rho^3 \end{aligned} \quad (6.2.21d)$$

Put $l = 1$ in (6.2.18). Take the new (6.2.18) into (6.2.21b).

$$\begin{aligned}
y_1(x) &= c_0 x^\lambda \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}\eta(1-t_1)(1-u_1)\right)}{v_1^{\omega_1+1}(1-v_1)} \\
&\quad \times \left\{ \sum_{i_0=0}^{\omega_0} (i_0 + \Omega/\mu + \lambda) \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} (\eta t_1 u_1 v_1)^{i_0} \right\} \rho \\
&= c_0 x^\lambda \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}\eta(1-t_1)(1-u_1)\right)}{v_1^{\omega_1+1}(1-v_1)} \\
&\quad \times w_{1,1}^{-(\Omega/\mu+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\Omega/\mu+\lambda} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,1}^{i_0} \right\} \rho \quad (6.2.22)
\end{aligned}$$

where $w_{1,1} = \eta \prod_{l=1}^1 t_l u_l v_l$

Put $l = 2$ in (6.2.18). Take the new (6.2.18) into (6.2.21c).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{v_2^{\omega_2+1}(1-v_2)} \\
&\quad \times w_{2,2}^{-(\Omega/\mu+2+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{\omega_0} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\omega_1} \frac{(-\omega_1)_{i_1}(3+\lambda)_{i_0}(2+\nu+\lambda)_{i_0}}{(-\omega_1)_{i_0}(3+\lambda)_{i_1}(2+\nu+\lambda)_{i_1}} w_{2,2}^{i_1} \right\} \rho^2 \quad (6.2.23)
\end{aligned}$$

where $w_{2,2} = \eta \prod_{l=2}^2 t_l u_l v_l$

Put $l = 1$ and $\eta = w_{2,2}$ in (6.2.18). Take the new (6.2.18) into (6.2.23).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{v_2^{\omega_2+1}(1-v_2)} \\
&\quad \times w_{2,2}^{-(\Omega/\mu+2+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}w_{2,2}(1-t_1)(1-u_1)\right)}{v_1^{\omega_1+1}(1-v_1)} \\
&\quad \times w_{1,2}^{-(\Omega/\mu+\lambda)} (w_{1,2}\partial_{w_{1,2}}) w_{2,2}^{\Omega/\mu+\lambda} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,2}^{i_0} \right\} \rho^2 \quad (6.2.24)
\end{aligned}$$

$$\text{where } w_{1,2} = \eta \prod_{l=1}^2 t_l u_l v_l$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) &= c_0 x^\lambda \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\nu+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{\exp\left(-\frac{v_3}{(1-v_3)}\eta(1-t_3)(1-u_3)\right)}{v_3^{\omega_3+1}(1-v_3)} \\
&\quad \times w_{3,3}^{-(\Omega/\mu+4+\lambda)} (w_{3,3}\partial_{w_{3,3}}) w_{3,3}^{\Omega/\mu+4+\lambda} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}w_{3,3}(1-t_2)(1-u_2)\right)}{v_2^{\omega_2+1}(1-v_2)} \\
&\quad \times w_{2,3}^{-(\Omega/\mu+2+\lambda)} (w_{2,3}\partial_{w_{2,3}}) w_{2,3}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}w_{2,3}(1-t_1)(1-u_1)\right)}{v_1^{\omega_1+1}(1-v_1)} \\
&\quad \times w_{1,3}^{-(\Omega/\mu+\lambda)} (w_{1,3}\partial_{w_{1,3}}) w_{1,3}^{\Omega/\mu+\lambda} \left\{ \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,3}^{i_0} \right\} \rho^3 \quad (6.2.25)
\end{aligned}$$

where

$$\begin{cases} w_{3,3} = \eta \prod_{l=3}^3 t_l u_l v_l \\ w_{2,3} = \eta \prod_{l=2}^3 t_l u_l v_l \\ w_{1,3} = \eta \prod_{l=1}^3 t_l u_l v_l \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute (6.2.21a), (6.2.22), (6.2.24), (6.2.25) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (6.2.20), we obtain (6.2.19). ■

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the GCH equation and $\lambda = 1 - \nu$ for the second one in (6.2.19).

Remark 6.2.6 *The integral representation of the GCH equation of the first kind for polynomial of type 2 about $x = 0$ as $\omega = -(\omega_j + 2j)$ where $j, \omega_j = 0, 1, 2, \dots$ is*

$$\begin{aligned}
y(x) &= QW_{\omega_j}^R(\mu, \varepsilon, \nu, \Omega, \omega = -(\omega_j + 2j); \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= {}_1F_1(-\omega_0; \nu; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\nu} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})\right)}{v_{n-k}^{\omega_{n-k}+1}(1-v_{n-k})} \\
&\quad \left. \left. \times w_{n-k,n}^{-(\Omega/\mu+2(n-k-1))} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\Omega/\mu+2(n-k-1)} \right\} {}_1F_1(-\omega_0; \nu; w_{1,n}) \right\} \rho^n \quad (6.2.26)
\end{aligned}$$

Remark 6.2.7 *The integral representation of the GCH equation of the second kind for polynomial of type 2 about $x = 0$ as $\omega = -(\omega_j + 2j + 1 - \nu)$ where $j, \omega_j = 0, 1, 2, \dots$ is*

$$\begin{aligned}
y(x) &= RW_{\omega_j}^R(\mu, \varepsilon, \nu, \Omega, \omega = -(\omega_j + 2j + 1 - \nu); \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= x^{1-\nu} \left\{ {}_1F_1(-\omega_0; 2 - \nu; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\nu} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})\right)}{v_{n-k}^{\omega_{n-k}+1}(1-v_{n-k})} \\
&\quad \left. \left. \times w_{n-k,n}^{-(\Omega/\mu+2(n-k)-1-\nu)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\Omega/\mu+2(n-k)-1-\nu} \right\} \right. \\
&\quad \left. \times {}_1F_1(-\omega_0; 2 - \nu; w_{1,n}) \right\} \rho^n \quad (6.2.27)
\end{aligned}$$

In the above, ${}_1F_1(a; b; z)$ is a Kummer function of the first kind which is defined by

$$\begin{aligned}
{}_1F_1(a; b; z) &= M(a, b, z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n n!} z^n = e^z M(b-a, b, -z) \\
&= -\frac{1}{2\pi i} \frac{\Gamma(1-a)\Gamma(b)}{\Gamma(b-a)} \oint dv_j e^{zv_j} (-v_j)^{a-1} (1-v_j)^{b-a-1} \\
&= \frac{\Gamma(a)}{2\pi i} \oint dv_j e^{v_j} v_j^{-b} \left(1 - \frac{z}{v_j}\right)^{-a} \\
&= \frac{1}{2\pi i} \frac{\Gamma(1-a)\Gamma(b)}{\Gamma(b-a)} \oint dv_j e^{-\frac{z}{1-v_j}} v_j^{a-1} (1-v_j)^{-b} \quad (6.2.28)
\end{aligned}$$

Infinite series

Let's consider the integral representation of the GCH equation about $x = 0$ for infinite series by applying R3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned}
M_l &= \sum_{i_l=i_{l-1}}^{\infty} \frac{(\omega+2l+\lambda)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\nu+\lambda)_{i_{l-1}}}{(\omega+2l+\lambda)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\nu+\lambda)_{i_l}} \eta^{i_l} \\
&= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+2l+\lambda, j+1) B(i_{l-1}+2l-1+\nu+\lambda, j+1) (\omega+2l+\lambda+i_{l-1})_j}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-1+\nu+\lambda)^{-1} (1)_j j!} \eta^j \quad (6.2.29)
\end{aligned}$$

Substitute (6.2.14a) and (6.2.14b) into (6.2.29), and divide $(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\nu+\lambda)$ into the new (6.2.29).

$$\begin{aligned}
&\frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\nu+\lambda)} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\omega+2l+\lambda)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\nu+\lambda)_{i_{l-1}}}{(\omega+2l+\lambda)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\nu+\lambda)_{i_l}} \eta^{i_l} \\
&= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-2+\nu+\lambda} (\eta t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(\omega+2l+\lambda+i_{l-1})_j}{(1)_j j!} (\eta(1-t_l)(1-u_l))^j \quad (6.2.30)
\end{aligned}$$

Replace a, b and z by $\omega+2l+\lambda+i_{l-1}, 1$ and $\eta(1-t_j)(1-u_j)$ in (6.2.28). Take the new (6.2.28) into (6.2.30).

$$\begin{aligned}
V_l &= \frac{1}{(i_{l-1}+2l+\lambda)(i_{l-1}+2l-1+\nu+\lambda)} \sum_{i_l=i_{l-1}}^{\infty} \frac{(\omega+2l+\lambda)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\nu+\lambda)_{i_{l-1}}}{(\omega+2l+\lambda)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\nu+\lambda)_{i_l}} \eta^{i_l} \\
&= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{\exp\left(-\frac{v_l}{(1-v_l)} \eta(1-t_l)(1-u_l)\right)}{v_l^{-(\omega+2l-1+\lambda)} (1-v_l)} (\eta t_l u_l v_l)^{i_{l-1}} \quad (6.2.31)
\end{aligned}$$

Substitute (6.2.31) into (6.2.10) where $l = 1, 2, 3, \dots$; apply V_1 into the second summation of sub-power series $y_1(x)$, apply V_2 into the third summation and V_1 into the second summation of sub-power series $y_2(x)$, apply V_3 into the forth summation, V_2 into the third summation and V_1 into the second summation of sub-power series $y_3(x)$, etc.⁸

Theorem 6.2.8 *The general representation in the form of integral of the GCH equation for infinite series about $x = 0$ using R3TRF is given by*

$$\begin{aligned}
y(x) &= \sum_{m=0}^{\infty} y_m(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\nu+\lambda} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})\right)}{v_{n-k}^{-(\omega+2(n-k)-1+\lambda)} (1-v_{n-k})} \\
&\quad \times w_{n-k,n}^{-(\Omega/\mu+2(n-k-1)+\lambda)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\Omega/\mu+2(n-k-1)+\lambda} \left. \right\} \\
&\quad \times \sum_{i_0=0}^{\infty} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} w_{1,n}^{i_0} \left. \right\} \rho^n \Bigg\} \tag{6.2.32}
\end{aligned}$$

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof In (6.2.10) sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of the CHE for infinite series using 3TRF about $x = 0$ are given by

$$y_0(x) = c_0 x^\lambda \sum_{i_0=0}^{\infty} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} \eta^{i_0} \tag{6.2.33a}$$

⁸ $y_1(x)$ means the sub-power series in (6.2.10) contains one term of B'_n 's, $y_2(x)$ means the sub-power series in (6.2.10) contains two terms of B'_n 's, $y_3(x)$ means the sub-power series in (6.2.10) contains three terms of B'_n 's, etc.

$$\begin{aligned}
y_1(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\omega + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \nu + \lambda)_{i_0}}{(\omega + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \nu + \lambda)_{i_1}} \eta^{i_1} \right\} \rho
\end{aligned} \tag{6.2.33b}$$

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + 2 + \Omega/\mu + \lambda)}{(i_1 + 4 + \lambda)(i_1 + 3 + \nu + \lambda)} \frac{(\omega + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \nu + \lambda)_{i_0}}{(\omega + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \nu + \lambda)_{i_1}} \\
&\quad \times \left. \sum_{i_2=i_1}^{\infty} \frac{(\omega + 4 + \lambda)_{i_2} (5 + \lambda)_{i_1} (4 + \nu + \lambda)_{i_1}}{(\omega + 4 + \lambda)_{i_1} (5 + \lambda)_{i_2} (4 + \nu + \lambda)_{i_2}} \eta^{i_2} \right\} \rho^2
\end{aligned} \tag{6.2.33c}$$

$$\begin{aligned}
y_3(x) &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + 2 + \Omega/\mu + \lambda)}{(i_1 + 4 + \lambda)(i_1 + 3 + \nu + \lambda)} \frac{(\omega + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \nu + \lambda)_{i_0}}{(\omega + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \nu + \lambda)_{i_1}} \\
&\quad \times \sum_{i_2=i_1}^{\infty} \frac{(i_2 + 4 + \Omega/\mu + \lambda)}{(i_2 + 6 + \lambda)(i_2 + 5 + \nu + \lambda)} \frac{(\omega + 4 + \lambda)_{i_2} (5 + \lambda)_{i_1} (4 + \nu + \lambda)_{i_1}}{(\omega + 4 + \lambda)_{i_1} (5 + \lambda)_{i_2} (4 + \nu + \lambda)_{i_2}} \\
&\quad \times \left. \sum_{i_3=i_2}^{\infty} \frac{(\omega + 6 + \lambda)_{i_3} (7 + \lambda)_{i_2} (6 + \nu + \lambda)_{i_2}}{(\omega + 6 + \lambda)_{i_2} (7 + \lambda)_{i_3} (6 + \nu + \lambda)_{i_3}} \eta^{i_3} \right\} \rho^3
\end{aligned} \tag{6.2.33d}$$

Put $l = 1$ in (6.2.31). Take the new (6.2.31) into (6.2.33b).

$$\begin{aligned}
y_1(x) &= c_0 x^\lambda \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}\eta(1-t_1)(1-u_1)\right)}{v_1^{-(\omega+1+\lambda)}(1-v_1)} \\
&\quad \times \left\{ \sum_{i_0=0}^{\infty} (i_0 + \Omega/\mu + \lambda) \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} (\eta t_1 u_1 v_1)^{i_0} \right\} \rho \\
&= c_0 x^\lambda \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}\eta(1-t_1)(1-u_1)\right)}{v_1^{-(\omega+1+\lambda)}(1-v_1)} \\
&\quad \times w_{1,1}^{-(\Omega/\mu+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\Omega/\mu+\lambda} \left\{ \sum_{i_0=0}^{\infty} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} w_{1,1}^{i_0} \right\} \rho \quad (6.2.34)
\end{aligned}$$

where $w_{1,1} = \eta \prod_{l=1}^1 t_l u_l v_l$

Put $l = 2$ in (6.2.31). Take the new (6.2.31) into (6.2.33c).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{v_2^{-(\omega+3+\lambda)}(1-v_2)} \\
&\quad \times w_{2,2}^{-(\Omega/\mu+2+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \Omega/\mu + \lambda)}{(i_0 + 2 + \lambda)(i_0 + 1 + \nu + \lambda)} \frac{(\omega + \lambda)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\omega + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} (2 + \nu + \lambda)_{i_0}}{(\omega + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} (2 + \nu + \lambda)_{i_1}} w_{2,2}^{i_1} \right\} \rho^2 \quad (6.2.35)
\end{aligned}$$

where $w_{2,2} = \eta \prod_{l=2}^2 t_l u_l v_l$

Put $l = 1$ and $\eta = w_{2,2}$ in (6.2.31). Take the new (6.2.31) into (6.2.35).

$$\begin{aligned}
y_2(x) &= c_0 x^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)} \eta(1-t_2)(1-u_2)\right)}{v_2^{-(\omega+3+\lambda)}(1-v_2)} \\
&\quad \times w_{2,2}^{-(\Omega/\mu+2+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} w_{2,2}(1-t_1)(1-u_1)\right)}{v_1^{-(\omega+1+\lambda)}(1-v_1)} \\
&\quad \times w_{1,2}^{-(\Omega/\mu+\lambda)} (w_{1,2} \partial_{w_{1,2}}) w_{2,2}^{\Omega/\mu+\lambda} \left\{ \sum_{i_0=0}^{\infty} \frac{(\omega+\lambda)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,2}^{i_0} \right\} \rho^2 \quad (6.2.36)
\end{aligned}$$

$$\text{where } w_{1,2} = \eta \prod_{l=1}^2 t_l u_l v_l$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) &= c_0 x^\lambda \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\nu+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{\exp\left(-\frac{v_3}{(1-v_3)} \eta(1-t_3)(1-u_3)\right)}{v_3^{-(\omega+5+\lambda)}(1-v_3)} \\
&\quad \times w_{3,3}^{-(\Omega/\mu+4+\lambda)} (w_{3,3} \partial_{w_{3,3}}) w_{3,3}^{\Omega/\mu+4+\lambda} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)} w_{3,3}(1-t_2)(1-u_2)\right)}{v_2^{-(\omega+3+\lambda)}(1-v_2)} \\
&\quad \times w_{2,3}^{-(\Omega/\mu+2+\lambda)} (w_{2,3} \partial_{w_{2,3}}) w_{2,3}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} w_{2,3}(1-t_1)(1-u_1)\right)}{v_1^{-(\omega+1+\lambda)}(1-v_1)} \\
&\quad \times w_{1,3}^{-(\Omega/\mu+\lambda)} (w_{1,3} \partial_{w_{1,3}}) w_{1,3}^{\Omega/\mu+\lambda} \left\{ \sum_{i_0=0}^{\infty} \frac{(\omega+\lambda)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,3}^{i_0} \right\} \rho^3 \quad (6.2.37)
\end{aligned}$$

where

$$\begin{cases} w_{3,3} = \eta \prod_{l=3}^3 t_l u_l v_l \\ w_{2,3} = \eta \prod_{l=2}^3 t_l u_l v_l \\ w_{1,3} = \eta \prod_{l=1}^3 t_l u_l v_l \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute (6.2.33a), (6.2.34), (6.2.36), (6.2.37) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (6.2.10), we obtain (6.2.32).⁹ ■

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of the GCH equation and $\lambda = 1 - \nu$ for the second one in (6.2.32).

Remark 6.2.9 *The integral representation of the GCH equation of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
 y(x) &= QW^R(\mu, \varepsilon, \nu, \Omega, \omega; \rho = -\mu x^2; \eta = -\varepsilon x) \\
 &= {}_1F_1(\omega; \nu; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\nu} \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})\right)}{v_{n-k}^{-(\omega+2(n-k)-1)}(1-v_{n-k})} \\
 &\quad \left. \left. \times w_{n-k,n}^{-(\Omega/\mu+2(n-k-1))} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\Omega/\mu+2(n-k-1)} \right\} {}_1F_1(\omega; \nu; w_{1,n}) \right\} \rho^n \quad (6.2.38)
 \end{aligned}$$

Remark 6.2.10 *The integral representation of the GCH equation of the second kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
 y(x) &= RW^R(\mu, \varepsilon, \nu, \Omega, \omega; \rho = -\mu x^2; \eta = -\varepsilon x) \\
 &= x^{1-\nu} \left\{ {}_1F_1(\omega + 1 - \nu; 2 - \nu; \eta) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\nu} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{\exp\left(-\frac{v_{n-k}}{(1-v_{n-k})} w_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})\right)}{v_{n-k}^{-(\omega+2(n-k)-\nu)}(1-v_{n-k})} \\
 &\quad \left. \left. \times w_{n-k,n}^{-(\Omega/\mu+2(n-k)-1-\nu)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\Omega/\mu+2(n-k)-1-\nu} \right\} \right. \\
 &\quad \left. \times {}_1F_1(\omega + 1 - \nu; 2 - \nu; w_{1,n}) \right\} \rho^n \quad (6.2.39)
 \end{aligned}$$

⁹Or replace the finite summation with an interval $[0, \omega_0]$ by infinite summation with an interval $[0, \infty]$ in (6.2.19). Replace ω_0 and ω_{n-j} by $-(\omega + \lambda)$ and $-(\omega + 2(n-k) + \lambda)$ into the new (6.2.19). Its solution is also equivalent to (6.2.32)

As $\frac{\Gamma(1/2 + \nu/2 - \Omega/(2\mu))}{\Gamma(1/2 + \nu/2)}$ is multiplied on (6.2.38), the new (6.2.38) is equivalent to the integral form of the first kind solution of the GCH equation for the infinite series using 3TRF.[36] Again, As $(-\mu/2)^{1/2(1-\nu)} \frac{\Gamma(1 - \Omega/(2\mu))}{\Gamma(3/2 - \nu/2)}$ is multiplied on (6.2.39), the new (6.2.39) corresponds to the integral representation of the second kind solution of the GCH equation for the infinite series using 3TRF.[36]

6.2.3 Generating function for the GCH polynomial of type 2

Now let's investigate the generating function for the type 2 GCH polynomials of the first and second kind about $x = 0$.

Definition 6.2.11 *I define that*

$$\begin{cases} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{where } a > b \\ s_a & \text{only if } a = b \end{cases} \\ \tilde{w}_{a,b} = \eta s_{a,\infty} \prod_{l=a}^b t_l v_l \end{cases} \quad (6.2.40)$$

where

$$a, b \in \mathbb{N}_0$$

And we have

$$\sum_{\omega_i=\omega_j}^{\infty} s_i^{\omega_i} = \frac{s_i^{\omega_j}}{(1-s_i)} \quad (6.2.41)$$

Acting the summation operator $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\}$ on (6.2.19) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (6.2.40) and (6.2.41),

Theorem 6.2.12 *The general expression of the generating function for the GCH*

polynomial of type 2 about $x = 0$ is given by

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y(x) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \Upsilon(\lambda; s_{0,\infty}; \eta) \\
&+ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \exp \left(-\frac{s_{1,\infty}}{(1 - s_{1,\infty})} \eta(1 - t_1)(1 - u_1) \right) \\
&\times \tilde{w}_{1,1}^{-(\Omega/\mu+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\Omega/\mu+\lambda} \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) \rho \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{k=n}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_n t_n^{2n-1+\lambda} \int_0^1 du_n u_n^{2(n-1)+\nu+\lambda} \exp \left(-\frac{s_{n,\infty}}{(1 - s_{n,\infty})} \eta(1 - t_n)(1 - u_n) \right) \right. \\
&\times \tilde{w}_{n,n}^{-(\Omega/\mu+2(n-1)+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\Omega/\mu+2(n-1)+\lambda} \\
&\times \prod_{j=1}^{n-1} \left\{ \int_0^1 dt_{n-j} t_{n-j}^{2(n-j)-1+\lambda} \int_0^1 du_{n-j} u_{n-j}^{2(n-j-1)+\nu+\lambda} \frac{\exp \left(-\frac{s_{n-j}}{(1 - s_{n-j})} \tilde{w}_{n-j+1,n}(1 - t_{n-j})(1 - u_{n-j}) \right)}{(1 - s_{n-j})} \right. \\
&\times \tilde{w}_{n-j,n}^{-(\Omega/\mu+2(n-j-1)+\lambda)} (\tilde{w}_{n-j,n} \partial_{\tilde{w}_{n-j,n}}) \tilde{w}_{n-j,n}^{\Omega/\mu+2(n-j-1)+\lambda} \left. \right\} \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) \left. \right\} \rho^n \tag{6.2.42}
\end{aligned}$$

where

$$\begin{cases} \Upsilon(\lambda; s_{0,\infty}; \eta) = \sum_{\omega_0=0}^{\infty} \frac{s_{0,\infty}^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \eta^{i_0} \right\} \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) = \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right\} \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) = \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0}(\nu + \lambda)_{i_0}} \tilde{w}_{1,n}^{i_0} \right\} \end{cases}$$

Proof Acting the summation operator $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\}$ on the

form of integral of the type 2 GCH polynomial $y(x)$,

$$\begin{aligned} & \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y(x) \\ &= \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} (y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots) \end{aligned} \quad (6.2.43)$$

Acting the summation operator $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\}$ on (6.2.21a) by using (6.2.40) and (6.2.41),

$$\begin{aligned} & \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_0(x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \sum_{\omega_0=0}^{\infty} \frac{s_{0,\infty}^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} \eta^{i_0} \right\} \end{aligned} \quad (6.2.44)$$

Acting the summation operator $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\}$ on (6.2.22) by using (6.2.40) and (6.2.41),

$$\begin{aligned} & \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} \eta(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\ &\quad \times \sum_{\omega_1=\omega_0}^{\infty} \left(\frac{s_{1,\infty}}{v_1} \right)^{\omega_1} w_{1,1}^{-(\Omega/\mu+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\Omega/\mu+\lambda} \\ &\quad \times \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1 + \lambda)_{i_0} (\nu + \lambda)_{i_0}} w_{1,1}^{i_0} \right\} \rho \end{aligned} \quad (6.2.45)$$

Replace ω_i , ω_j and s_i by ω_1 , ω_0 and $\frac{s_{1,\infty}}{v_1}$ in (6.2.41). Take the new (6.2.41) into (6.2.45).

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_1(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}\eta(1-t_1)(1-u_1)\right)}{(1-v_1)(v_1-s_{1,\infty})} \\
&\times w_{1,1}^{-(\Omega/\mu+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\Omega/\mu+\lambda} \\
&\times \sum_{\omega_0=0}^{\infty} \frac{1}{\omega_0!} \left(\frac{s_{0,\infty}}{v_1}\right)^{\omega_0} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,1}^{i_0} \right\} \rho \quad (6.2.46)
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $v_1 = 1$ or $s_{1,\infty}$, and $s_{1,\infty}$ is only inside the unit circle. As we compute the residue there in (6.2.46) we obtain

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_1(x) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \exp\left(-\frac{s_{1,\infty}}{(1-s_{1,\infty})}\eta(1-t_1)(1-u_1)\right) \\
&\times \tilde{w}_{1,1}^{-(\Omega/\mu+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\Omega/\mu+\lambda} \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right\} \rho \quad (6.2.47)
\end{aligned}$$

where

$$\tilde{w}_{1,1} = \eta s_{1,\infty} \prod_{l=1}^1 t_l u_l$$

Acting the summation operator $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\}$ on (6.2.24) by

using (6.2.40) and (6.2.41),

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{v_2(1-v_2)} \\
&\times \sum_{\omega_2=\omega_1}^{\infty} \left(\frac{s_{2,\infty}}{v_2}\right)^{\omega_2} w_{2,2}^{-(\Omega/\mu+2+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\Omega/\mu+2+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}w_{2,2}(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\
&\times \sum_{\omega_1=\omega_0}^{\infty} \left(\frac{s_1}{v_1}\right)^{\omega_1} w_{1,2}^{-(\Omega/\mu+\lambda)} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\Omega/\mu+\lambda} \\
&\times \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,2}^{i_0} \right\} \rho^2 \tag{6.2.48}
\end{aligned}$$

Replace ω_i , ω_j and s_i by ω_2 , ω_1 and $\frac{s_{2,\infty}}{v_2}$ in (6.2.41). Take the new (6.2.41) into (6.2.48).

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{\exp\left(-\frac{v_2}{(1-v_2)}\eta(1-t_2)(1-u_2)\right)}{(1-v_2)(v_2-s_{2,\infty})} \\
&\times w_{2,2}^{-(\Omega/\mu+2+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\Omega/\mu+2+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)}w_{2,2}(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \\
&\times \sum_{\omega_1=\omega_0}^{\infty} \left(\frac{s_{1,\infty}}{v_1 v_2}\right)^{\omega_1} w_{1,2}^{-(\Omega/\mu+\lambda)} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\Omega/\mu+\lambda} \\
&\times \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} w_{1,2}^{i_0} \right\} \rho^2 \tag{6.2.49}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $v_2 = 1$ or $s_{2,\infty}$, and $s_{2,\infty}$ is only inside the unit circle. As we compute the residue there in (6.2.49) we

obtain

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_2(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \exp\left(-\frac{s_{2,\infty}}{(1-s_{2,\infty})} \eta(1-t_2)(1-u_2)\right) \\
&\quad \times \tilde{w}_{2,2}^{-(\Omega/\mu+2+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} \tilde{w}_{2,2}(1-t_1)(1-u_1)\right)}{v_1(1-v_1)} \sum_{\omega_1=\omega_0}^{\infty} \left(\frac{s_1}{v_1}\right)^{\omega_1} \\
&\quad \times \ddot{w}_{1,2}^{-(\Omega/\mu+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\Omega/\mu+\lambda} \\
&\quad \times \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right\} \rho^2 \tag{6.2.50}
\end{aligned}$$

where

$$\tilde{w}_{2,2} = \eta s_{2,\infty} \prod_{l=2}^2 t_l u_l \quad \ddot{w}_{1,2} = \eta s_{2,\infty} v_1 \prod_{l=1}^2 t_l u_l$$

Replace ω_i , ω_j and s_i by ω_1 , ω_0 and $\frac{s_1}{v_1}$ in (6.2.41). Take the new (6.2.41) into (6.2.50).

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_2(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \exp\left(-\frac{s_{2,\infty}}{(1-s_{2,\infty})} \eta(1-t_2)(1-u_2)\right) \\
&\quad \times \tilde{w}_{2,2}^{-(\Omega/\mu+2+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{\exp\left(-\frac{v_1}{(1-v_1)} \tilde{w}_{2,2}(1-t_1)(1-u_1)\right)}{(1-v_1)(v_1-s_1)} \\
&\quad \times \ddot{w}_{1,2}^{-(\Omega/\mu+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\Omega/\mu+\lambda} \\
&\quad \times \sum_{\omega_0=0}^{\infty} \frac{1}{\omega_0!} \left(\frac{s_{0,1}}{v_1}\right)^{\omega_0} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right\} \rho^2 \tag{6.2.51}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $v_1 = 1$ or s_1 , and

s_1 is only inside the unit circle. As we compute the residue there in (6.2.51) we obtain

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_2(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \exp\left(-\frac{s_{2,\infty}}{(1-s_{2,\infty})} \eta(1-t_2)(1-u_2)\right) \\
&\quad \times \tilde{w}_{2,2}^{-(\Omega/\mu+2+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{\exp\left(-\frac{s_1}{(1-s_1)} \tilde{w}_{2,2}(1-t_1)(1-u_1)\right)}{(1-s_1)} \tilde{w}_{1,2}^{-(\Omega/\mu+\lambda)} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\Omega/\mu+\lambda} \\
&\quad \times \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \tilde{w}_{1,2}^{i_0} \right\} \rho^2 \tag{6.2.52}
\end{aligned}$$

where

$$\tilde{w}_{1,2} = \eta s_{1,\infty} \prod_{l=1}^2 t_l u_l$$

Acting the summation operator $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\}$ on (6.2.52) by using (6.2.40) and (6.2.41),

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_3(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{4+\nu+\lambda} \exp\left(-\frac{s_{3,\infty}}{(1-s_{3,\infty})} \eta(1-t_3)(1-u_3)\right) \\
&\quad \times \tilde{w}_{3,3}^{-(\Omega/\mu+4+\lambda)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{\Omega/\mu+4+\lambda} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{2+\nu+\lambda} \frac{\exp\left(-\frac{s_2}{(1-s_2)} \tilde{w}_{3,3}(1-t_2)(1-u_2)\right)}{(1-s_2)} \tilde{w}_{2,3}^{-(\Omega/\mu+2+\lambda)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{\Omega/\mu+2+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\nu+\lambda} \frac{\exp\left(-\frac{s_1}{(1-s_1)} \tilde{w}_{2,3}(1-t_1)(1-u_1)\right)}{(1-s_1)} \tilde{w}_{1,3}^{-(\Omega/\mu+\lambda)} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\Omega/\mu+\lambda} \\
&\quad \times \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \left\{ c_0 x^\lambda \sum_{i_0=0}^{\omega_0} \frac{(-\omega_0)_{i_0}}{(1+\lambda)_{i_0}(\nu+\lambda)_{i_0}} \tilde{w}_{1,3}^{i_0} \right\} \rho^3 \tag{6.2.53}
\end{aligned}$$

where

$$\tilde{w}_{3,3} = \eta s_{3,\infty} \prod_{l=3}^3 t_l u_l \quad \tilde{w}_{2,3} = \eta s_{2,\infty} v_1 \prod_{l=2}^3 t_l u_l \quad \tilde{w}_{1,3} = \eta s_{1,\infty} v_1 \prod_{l=1}^3 t_l u_l$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m > 3$, we obtain every $\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_m(x)$ terms.

Since we substitute (6.2.44), (6.2.47), (6.2.52), (6.2.53) and including all

$$\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma')}{\Gamma(\gamma')} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} y_m(x) \text{ terms where } m > 3 \text{ into (6.2.43), we obtain}$$

(6.2.42) ■

Remark 6.2.13 *The generating function for the GCH polynomial of type 2 of the first kind about $x = 0$ as $\omega = -(\omega_j + 2j)$ where $j, \omega_j = 0, 1, 2, \dots$ is*

$$\begin{aligned} & \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \nu)}{\Gamma(\nu)} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} QW_{\omega_j}^R(\mu, \varepsilon, \nu, \Omega, \omega = -(\omega_j + 2j); \rho = -\mu x^2; \eta = -\varepsilon x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \\ &+ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\nu} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \tilde{w}_{1,1}^{-\Omega/\mu}(\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\Omega/\mu} \mathbf{A}(s_0; \tilde{w}_{1,1}) \rho \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{k=n}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\nu} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ &\times \tilde{w}_{n,n}^{-(\Omega/\mu+2(n-1))}(\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\Omega/\mu+2(n-1)} \\ &\times \prod_{j=1}^{n-1} \left\{ \int_0^1 dt_{n-j} t_{n-j}^{2(n-j)-1} \int_0^1 du_{n-j} u_{n-j}^{2(n-j-1)+\nu} \overleftrightarrow{\mathbf{F}}_{n-j}(s_{n-j}; t_{n-j}, u_{n-j}, \tilde{w}_{n-j+1,n}) \right. \\ &\times \tilde{w}_{n-j,n}^{-(\Omega/\mu+2(n-j-1))}(\tilde{w}_{n-j,n} \partial_{\tilde{w}_{n-j,n}}) \tilde{w}_{n-j,n}^{\Omega/\mu+2(n-j-1)} \left. \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} \rho^n \end{aligned} \quad (6.2.54)$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \exp\left(-\frac{s_{1,\infty}}{(1-s_{1,\infty})}\eta(1-t_1)(1-u_1)\right) \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \exp\left(-\frac{s_{n,\infty}}{(1-s_{n,\infty})}\eta(1-t_n)(1-u_n)\right) \\ \overleftrightarrow{\mathbf{F}}_{n-j}(s_{n-j}; t_{n-j}, u_{n-j}, \tilde{w}_{n-j+1,n}) = \frac{\exp\left(-\frac{s_{n-j}}{(1-s_{n-j})}\tilde{w}_{n-j+1,n}(1-t_{n-j})(1-u_{n-j})\right)}{(1-s_{n-j})} \end{cases}$$

and

$$\begin{cases} \mathbf{A}(s_{0,\infty}; \eta) = (1-s_{0,\infty})^{-\nu} \exp\left(-\frac{\eta s_{0,\infty}}{(1-s_{0,\infty})}\right) \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = (1-s_0)^{-\nu} \exp\left(-\frac{\tilde{w}_{1,1}s_0}{(1-s_0)}\right) \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = (1-s_0)^{-\nu} \exp\left(-\frac{\tilde{w}_{1,n}s_0}{(1-s_0)}\right) \end{cases}$$

Proof The generating function for confluent Hypergeometric polynomial of the first kind is given by

$$\sum_{\omega_0=0}^{\infty} \frac{t^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \gamma)}{\Gamma(\gamma)} {}_1F_1(-\omega_0; \gamma; z) = (1-t)^{-\gamma} \exp\left(-\frac{zt}{(1-t)}\right) \quad (6.2.55)$$

Replace t , γ and z by $s_{0,\infty}$, ν and η in (6.2.55).

$$\sum_{\omega_0=0}^{\infty} \frac{s_{0,\infty}^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \nu)}{\Gamma(\nu)} {}_1F_1(-\omega_0; \nu; \eta) = (1-s_{0,\infty})^{-\nu} \exp\left(-\frac{\eta s_{0,\infty}}{(1-s_{0,\infty})}\right) \quad (6.2.56)$$

Replace t , γ and z by s_0 , ν and $\tilde{w}_{1,1}$ in (6.2.55).

$$\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \nu)}{\Gamma(\nu)} {}_1F_1(-\omega_0; \nu; \tilde{w}_{1,1}) = (1-s_0)^{-\nu} \exp\left(-\frac{\tilde{w}_{1,1}s_0}{(1-s_0)}\right) \quad (6.2.57)$$

Replace t , γ and z by s_0 , ν and $\tilde{w}_{1,n}$ in (6.2.55).

$$\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + \nu)}{\Gamma(\nu)} {}_1F_1(-\omega_0; \nu; \tilde{w}_{1,n}) = (1-s_0)^{-\nu} \exp\left(-\frac{\tilde{w}_{1,n}s_0}{(1-s_0)}\right) \quad (6.2.58)$$

Put $c_0=1$, $\lambda=0$ and $\gamma'=\nu$ in (6.2.42). Substitute (6.2.56), (6.2.57) and (6.2.58) into the new (6.2.42). ■

Remark 6.2.14 *The generating function for the GCH polynomial of type 2 of the second kind about $x = 0$ as $\omega = -(\omega_j + 2j + 1 - \nu)$ where $j, \omega_j = 0, 1, 2, \dots$ is*

$$\begin{aligned}
& \sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + 2 - \nu)}{\Gamma(2 - \nu)} \prod_{n=1}^{\infty} \left\{ \sum_{\omega_n=\omega_{n-1}}^{\infty} s_n^{\omega_n} \right\} RW_{\omega_j}^R(\mu, \varepsilon, \nu, \Omega, \omega = -(\omega_j + 2j + 1 - \nu); \rho = -\mu x^2; \eta = -\varepsilon x) \\
&= x^{1-\nu} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{B}(s_{0,\infty}; \eta) \right. \\
&+ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{2-\nu} \int_0^1 du_1 u_1 \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \tilde{w}_{1,1}^{-(\Omega/\mu+1-\nu)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\Omega/\mu+1-\nu} \mathbf{B}(s_0; \tilde{w}_{1,1}) \rho \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{k=n}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_n t_n^{2n-\nu} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \tilde{w}_{n,n}^{-(\Omega/\mu+2n-1-\nu)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\Omega/\mu+2n-1-\nu} \\
&\times \prod_{j=1}^{n-1} \left\{ \int_0^1 dt_{n-j} t_{n-j}^{2(n-j)-\nu} \int_0^1 du_{n-j} u_{n-j}^{2(n-j)-1} \overleftrightarrow{\mathbf{F}}_{n-j}(s_{n-j}; t_{n-j}, u_{n-j}, \tilde{w}_{n-j+1,n}) \right. \\
&\times \tilde{w}_{n-j,n}^{-(\Omega/\mu+2(n-j)-1-\nu)} (\tilde{w}_{n-j,n} \partial_{\tilde{w}_{n-j,n}}) \tilde{w}_{n-j,n}^{\Omega/\mu+2(n-j)-1-\nu} \left. \left. \right\} \mathbf{B}(s_0; \tilde{w}_{1,n}) \right\} \rho^n \Bigg\} \quad (6.2.59)
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \exp\left(-\frac{s_{1,\infty}}{(1 - s_{1,\infty})} \eta (1 - t_1)(1 - u_1)\right) \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \exp\left(-\frac{s_{n,\infty}}{(1 - s_{n,\infty})} \eta (1 - t_n)(1 - u_n)\right) \\ \overleftrightarrow{\mathbf{F}}_{n-j}(s_{n-j}; t_{n-j}, u_{n-j}, \tilde{w}_{n-j+1,n}) = \frac{\exp\left(-\frac{s_{n-j}}{(1 - s_{n-j})} \tilde{w}_{n-j+1,n} (1 - t_{n-j})(1 - u_{n-j})\right)}{(1 - s_{n-j})} \end{cases}$$

and

$$\begin{cases} \mathbf{B}(s_{0,\infty}; \eta) = (1 - s_{0,\infty})^{\nu-2} \exp\left(-\frac{\eta s_{0,\infty}}{(1 - s_{0,\infty})}\right) \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = (1 - s_0)^{\nu-2} \exp\left(-\frac{\tilde{w}_{1,1} s_0}{(1 - s_0)}\right) \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = (1 - s_0)^{\nu-2} \exp\left(-\frac{\tilde{w}_{1,n} s_0}{(1 - s_0)}\right) \end{cases}$$

Proof Replace t , γ and z by $s_{0,\infty}$, $2 - \nu$ and η in (6.2.55).

$$\sum_{\omega_0=0}^{\infty} \frac{s_{0,\infty}^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + 2 - \nu)}{\Gamma(2 - \nu)} {}_1F_1(-\omega_0; 2 - \nu; \eta) = (1 - s_{0,\infty})^{\nu-2} \exp\left(-\frac{\eta s_{0,\infty}}{(1 - s_{0,\infty})}\right) \quad (6.2.60)$$

Replace t , γ and z by s_0 , $2 - \nu$ and $\tilde{w}_{1,1}$ in (6.2.55).

$$\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + 2 - \nu)}{\Gamma(2 - \nu)} {}_1F_1(-\omega_0; 2 - \nu; \tilde{w}_{1,1}) = (1 - s_0)^{\nu-2} \exp\left(-\frac{\tilde{w}_{1,1}s_0}{(1 - s_0)}\right) \quad (6.2.61)$$

Replace t , γ and z by s_0 , $2 - \nu$ and $\tilde{w}_{1,n}$ in (6.2.55).

$$\sum_{\omega_0=0}^{\infty} \frac{s_0^{\omega_0}}{\omega_0!} \frac{\Gamma(\omega_0 + 2 - \nu)}{\Gamma(2 - \nu)} {}_1F_1(-\omega_0; 2 - \nu; \tilde{w}_{1,n}) = (1 - s_0)^{\nu-2} \exp\left(-\frac{\tilde{w}_{1,n}s_0}{(1 - s_0)}\right) \quad (6.2.62)$$

Put $c_0 = 1$, $\lambda = 1 - \nu$ and $\gamma' = 2 - \nu$ in (6.2.42). Substitute (6.2.60), (6.2.61) and (6.2.62) into the new (6.2.42). ■

6.3 GCH equation about irregular singular point at infinity

Let $z = \frac{1}{x}$ in (6.1.1) in order to obtain the analytic solution of the GCH equation about $x = \infty$.

$$z^4 \frac{d^2 y}{dz^2} + ((2 - \mu)z^3 - \varepsilon z^2 - \mu z) \frac{dy}{dz} + (\Omega + \varepsilon \omega z) y = 0 \quad (6.3.1)$$

Assume that its solution is

$$y(z) = \sum_{n=0}^{\infty} c_n z^{n+\lambda} \quad (6.3.2)$$

where λ is indicial root. Plug (6.3.2) into (6.3.1). We get a three-term recurrence relation for the coefficients c_n :

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (6.3.3)$$

where,

$$A_n = -\frac{\varepsilon}{\mu} \frac{(n - \omega + \lambda)}{(n + 1 - \Omega/\mu + \lambda)}$$

$$B_n = \frac{1}{\mu} \frac{(n - 1 + \lambda)(n - \nu + \lambda)}{(n + 1 - \Omega/\mu + \lambda)} \quad (6.3.4a)$$

$$c_1 = A_0 c_0 \quad (6.3.4b)$$

We have an indicial roots which is $\lambda = \Omega/\mu$.

Now let's test for convergence of the analytic function $y(z)$. As $n \gg 1$, (6.3.4a)–(6.3.4b) are

$$\lim_{n \gg 1} A_n = -\frac{\varepsilon}{\mu} \quad (6.3.5a)$$

$$\lim_{n \gg 1} B_n = \frac{n}{\mu} \quad (6.3.5b)$$

There are no analytic solutions for type 2 polynomial and infinite series. Because the $y(z)$ will be divergent as $n \gg 1$ in (6.3.5b). So there are only two types of the analytic solutions of the GCH equation about $x = \infty$ which are type 1 and type 3 polynomials. In future papers I will derive type 3 polynomial about $x = \infty$: I treat μ , ε and Ω as free variables and ν and ω as fixed values. In this chapter I construct the power series expansion, its integral forms and the generating function for the GCH polynomial of type 1 about $x = \infty$: I treat μ , ε , ω and Ω as free variables and ν as a fixed value.

For type 1 polynomial, (6.3.5a) is only available for the asymptotic behavior of the minimum $y(z)$.¹⁰ Substitute (6.3.5a) into (6.3.3) with $B_n = 0$. For $n = 0, 1, 2, \dots$, it give

$$\begin{aligned} c_0 & \\ c_1 &= -\frac{\varepsilon}{\mu} c_0 \\ c_2 &= \left(-\frac{\varepsilon}{\mu}\right)^2 c_0 \\ c_3 &= \left(-\frac{\varepsilon}{\mu}\right)^3 c_0 \\ c_4 &= \left(-\frac{\varepsilon}{\mu}\right)^4 c_0 \\ &\vdots \end{aligned} \quad (6.3.6)$$

Put (6.3.6) into (6.3.2) taking $c_0 = 1$ for simplicity and $\lambda = \Omega/\mu$.

$$\lim_{n \gg 1} y(z) > z^{\Omega/\mu} \sum_{n=0}^{\infty} \left(-\frac{\varepsilon}{\mu}\right)^n z^n = \frac{z^{\Omega/\mu}}{1 + \frac{\varepsilon}{\mu} z} \quad \text{where } \left|\frac{\varepsilon}{\mu} z\right| < 1 \quad (6.3.7)$$

Since $|\varepsilon|, |\mu| \gg 1$, (6.3.5b) is negligible relative to (6.3.5a). The approximative $y(z)$ as $|\varepsilon|, |\mu| \gg 1$ is

$$\lim_{\substack{n \gg 1 \\ |\varepsilon|, |\mu| \gg 1}} y(z) = \frac{z^{\Omega/\mu}}{1 + \frac{\varepsilon}{\mu} z} \quad (6.3.8)$$

¹⁰(6.3.5b) is negligible for the minimum $y(z)$ because B_n term will be terminated at the specific eigenvalues.

6.3.1 Power series for polynomial of type 1

In Ref.[35], the general expression of power series of $y(x)$ for polynomial of type 1 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\beta_0} \left(\prod_{i_1=0}^{i_0-1} B_{2i_1+1} \right) x^{2i_0+\lambda} + \sum_{i_0=0}^{\beta_0} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \sum_{i_2=i_0}^{\beta_1} \left(\prod_{i_3=i_0}^{i_2-1} B_{2i_3+2} \right) \right\} x^{2i_2+1+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\beta_0} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\beta_k} A_{2i_{2k}+k} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} B_{2i_{2k+1}+(k+1)} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\beta_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} B_{2i_{2N+1}+(N+1)} \right) \right\} \right\} x^{2i_{2N}+N+\lambda} \Bigg\} \quad (6.3.9)
\end{aligned}$$

In the above, $\beta_i \leq \beta_j$ only if $i \leq j$ where $i, j, \beta_i, \beta_j \in \mathbb{N}_0$.

For a polynomial, we need a condition, which is:

$$B_{2\beta_i+(i+1)} = 0 \quad \text{where } i = 0, 1, 2, \dots, \beta_i = 0, 1, 2, \dots \quad (6.3.10)$$

In the above, β_i is an eigenvalue that makes B_n term terminated at certain value of index n . (6.3.10) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (6.3.9).

In (6.3.4a) replace ν by $2\nu_i + i + 1 + \lambda$. In (6.3.9) and (6.3.10) replace an independent variable x and an index β_i by z and ν_i . Take the new (6.3.4a)–(6.3.4b), (6.3.10) and put them in (6.3.9). After the replacement process, the general expression of power series of

the GCH equation for polynomial of type 1 about $x = \infty$ is given by

$$\begin{aligned}
y(z) &= \sum_{n=0}^{\infty} y_n(z) \\
&= c_0 z^\lambda \left\{ \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{(i_0 + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \sum_{i_1=i_0}^{\nu_1} \frac{(-\nu_1)_{i_1} (\frac{1}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}}{(-\nu_1)_{i_0} (\frac{1}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_1}} \eta^{i_1} \right\} \xi \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{(i_0 + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\nu_k} \frac{(i_k + \frac{k}{2} - \frac{\omega}{2} + \frac{\lambda}{2})}{(i_k + \frac{k}{2} + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \frac{(-\nu_k)_{i_k} (\frac{k}{2} + \frac{\lambda}{2})_{i_k} (\frac{k}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_{k-1}}}{(-\nu_k)_{i_{k-1}} (\frac{k}{2} + \frac{\lambda}{2})_{i_{k-1}} (\frac{k}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_k}} \right\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\nu_n} \frac{(-\nu_n)_{i_n} (\frac{n}{2} + \frac{\lambda}{2})_{i_n} (\frac{n}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_{n-1}}}{(-\nu_n)_{i_{n-1}} (\frac{n}{2} + \frac{\lambda}{2})_{i_{n-1}} (\frac{n}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_n}} \eta^{i_n} \left. \right\} \xi^n \Bigg\} \quad (6.3.11)
\end{aligned}$$

where

$$\begin{cases} \eta = \frac{2}{\mu} z^2 \\ \xi = -\frac{\varepsilon}{\mu} z \\ \nu = 2\nu_j + j + 1 + \lambda \\ z = \frac{1}{x} \\ \nu_i \leq \nu_j \text{ only if } i \leq j \text{ where } i, j, \nu_i, \nu_j \in \mathbb{N}_0 \dots \end{cases}$$

Put $c_0 = 1$ as $\lambda = \Omega/\mu$ in (6.3.11).

Remark 6.3.1 *The power series expansion of the GCH equation of the first kind for*

polynomial of type 1 about $x = \infty$ as $\nu = 2\nu_j + j + 1 + \Omega/\mu$ where $j, \nu_j \in \mathbb{N}_0$ is

$$\begin{aligned}
 y(z) &= Q^{(i)} W_{\nu_j} \left(\mu, \varepsilon, \Omega, \omega, \nu = 2\nu_j + j + 1 + \frac{\Omega}{\mu}; z = \frac{1}{x}; \xi = -\frac{\varepsilon}{\mu} z; \eta = \frac{2}{\mu} z^2 \right) \\
 &= z^{\frac{\Omega}{\mu}} \left\{ \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} (\frac{\Omega}{2\mu})_{i_0}}{(1)_{i_0}} \eta^{i_0} \right. \\
 &\quad + \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\Omega}{2\mu})}{(i_0 + \frac{1}{2})} \frac{(-\nu_0)_{i_0} (\frac{\Omega}{2\mu})_{i_0}}{(1)_{i_0}} \sum_{i_1=i_0}^{\nu_1} \frac{(-\nu_1)_{i_1} (\frac{1}{2} + \frac{\Omega}{2\mu})_{i_1} (\frac{3}{2})_{i_0}}{(-\nu_1)_{i_0} (\frac{1}{2} + \frac{\Omega}{2\mu})_{i_0} (\frac{3}{2})_{i_1}} \eta^{i_1} \right\} \xi \\
 &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\Omega}{2\mu})}{(i_0 + \frac{1}{2})} \frac{(-\nu_0)_{i_0} (\frac{\Omega}{2\mu})_{i_0}}{(1)_{i_0}} \right. \\
 &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\nu_k} \frac{(i_k + \frac{k}{2} - \frac{\omega}{2} + \frac{\Omega}{2\mu})}{(i_k + \frac{k}{2} + \frac{1}{2})} \frac{(-\nu_k)_{i_k} (\frac{k}{2} + \frac{\Omega}{2\mu})_{i_k} (\frac{k}{2} + 1)_{i_{k-1}}}{(-\nu_k)_{i_{k-1}} (\frac{k}{2} + \frac{\Omega}{2\mu})_{i_{k-1}} (\frac{k}{2} + 1)_{i_k}} \right\} \\
 &\quad \times \left. \sum_{i_n=i_{n-1}}^{\nu_n} \frac{(-\nu_n)_{i_n} (\frac{n}{2} + \frac{\Omega}{2\mu})_{i_n} (\frac{n}{2} + 1)_{i_{n-1}}}{(-\nu_n)_{i_{n-1}} (\frac{n}{2} + \frac{\Omega}{2\mu})_{i_{n-1}} (\frac{n}{2} + 1)_{i_n}} \eta^{i_n} \right\} \xi^n \Bigg\} \quad (6.3.12)
 \end{aligned}$$

6.3.2 Integral formalism for polynomial of type 1

There is a generalized hypergeometric function which is given by

$$\begin{aligned}
 L_l &= \sum_{i_l=i_{l-1}}^{\nu_l} \frac{(-\nu_l)_{i_l} (\frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{l}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_{l-1}}}{(-\nu_l)_{i_{l-1}} (\frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{l}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_l}} \eta^{i_l} \\
 &= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B \left(i_{l-1} + \frac{l}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}, j+1 \right) (i_{l-1} - \nu_l)_j (i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})_j}{(i_{l-1} + \frac{l}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})^{-1} (1)_j} \eta^j \quad (6.3.13)
 \end{aligned}$$

By using integral form of beta function,

$$B \left(i_{l-1} + \frac{l}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}, j+1 \right) = \int_0^1 dt_l t_l^{i_{l-1} + \frac{l}{2} - 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}} (1 - t_l)^j \quad (6.3.14)$$

Substitute (6.3.14) into (6.3.13), and divide $(i_{l-1} + \frac{l}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})$ into L_l .

$$\begin{aligned}
 G_l &= \frac{1}{(i_{l-1} + \frac{l}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \sum_{i_l=i_{l-1}}^{\nu_l} \frac{(-\nu_l)_{i_l} (\frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{l}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_{l-1}}}{(-\nu_l)_{i_{l-1}} (\frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{l}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_l}} \eta^{i_l} \\
 &= \int_0^1 dt_l t_l^{\frac{l}{2} - 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}} (\eta t_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(i_{l-1} - \nu_l)_j (i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})_j}{(1)_j} (\eta(1 - t_l))^j \quad (6.3.15)
 \end{aligned}$$

Tricomi's function is defined by

$$U(a, b, z) = \frac{\Gamma(1-b)}{\Gamma(a-b+1)} M(a, b, z) + \frac{\Gamma(b-1)}{\Gamma(a)} z^{1-b} M(a-b+1, 2-b, z) \quad (6.3.16)$$

The contour integral form of (6.3.16) is given by [19]

$$U(a, b, z) = e^{-a\pi i} \frac{\Gamma(1-a)}{2\pi i} \int_{\infty}^{(0+)} dp_l e^{-zp_l} p_l^{a-1} (1+p_l)^{b-a-1} \text{ where } a \neq 1, 2, 3, \dots, \quad |\text{ph}z| < \frac{1}{2}\pi \quad (6.3.17)$$

Also (6.3.16) is written by [19]

$$U(a, b, z) = z^{-a} \sum_{j=0}^{\infty} \frac{(a)_j (a-b+1)_j}{(1)_j} (-z^{-1})^j = z^{-a} {}_2F_0(a, a-b+1; -; -z^{-1}) \quad (6.3.18)$$

Replace a, b and z by $i_{l-1} - \nu_l, -\nu_l + 1 - \frac{l}{2} - \frac{\lambda}{2}$ and $\frac{-1}{\eta(1-t_l)}$ into (6.3.18).

$$\sum_{j=0}^{\infty} \frac{(i_{l-1} - \nu_l)_j (i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})_j}{(1)_j} (\eta(1-t_l))^j = \left(\frac{-1}{\eta(1-t_l)} \right)^{i_{l-1} - \nu_l} U \left(i_{l-1} - \nu_l, -\nu_l + 1 - \frac{l}{2} - \frac{\lambda}{2}, \frac{-1}{\eta(1-t_l)} \right) \quad (6.3.19)$$

Replace a, b and z by $i_{l-1} - \nu_l, -\nu_l + 1 - \frac{l}{2} - \frac{\lambda}{2}$ and $\frac{-1}{\eta(1-t_l)}$ into (6.3.17). Take the new (6.3.17) into (6.3.19)

$$\begin{aligned} & \sum_{j=0}^{\infty} \frac{(i_{l-1} - \nu_l)_j (i_{l-1} + \frac{l}{2} + \frac{\lambda}{2})_j}{(1)_j} (\eta(1-t_l))^j \\ &= \frac{\Gamma(\nu_l - i_{l-1} + 1)}{2\pi i} \int_{\infty}^{(0+)} dp_l \exp \left(\frac{p_l}{\eta(1-t_l)} \right) p_l^{-1} (1+p_l)^{-\frac{1}{2}(l+\lambda)} \\ & \times \left(\frac{\eta(1-t_l)}{p_l} \right)^{\nu_l} \left(\frac{p_l}{\eta(1-t_l)(1+p_l)} \right)^{i_{l-1}} \end{aligned} \quad (6.3.20)$$

The definition of the Gamma function $\Gamma(z)$ is written by

$$\Gamma(z) = \int_0^{\infty} du_l e^{-u_l} u_l^{z-1} \text{ where } \text{Re}(z) > 0 \quad (6.3.21)$$

Put $z = \nu_l - i_{l-1} + 1$ in (6.3.21).

$$\Gamma(\nu_l - i_{l-1} + 1) = \int_0^{\infty} du_l e^{-u_l} u_l^{\nu_l - i_{l-1}} \quad (6.3.22)$$

Put (6.3.22) in (6.3.20). Take the new (6.3.20) into (6.3.15)

$$\begin{aligned}
G_l &= \frac{1}{(i_{l-1} + \frac{l}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \sum_{i_l=i_{l-1}}^{\nu_l} \frac{(-\nu_l)_{i_l} (\frac{l}{2} + \frac{\lambda}{2})_{i_l} (\frac{l}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_{l-1}}}{(-\nu_l)_{i_{l-1}} (\frac{l}{2} + \frac{\lambda}{2})_{i_{l-1}} (\frac{l}{2} + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_l}} \eta^{i_l} \\
&= \int_0^1 dt_l t_l^{\frac{1}{2}(l-2-\frac{\Omega}{\mu}+\lambda)} \int_0^\infty du_l e^{-u_l} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_l \exp\left(\frac{p_l}{\eta(1-t_l)}\right) p_l^{-1} (1+p_l)^{-\frac{1}{2}(l+\lambda)} \\
&\quad \times \left(\frac{\eta u_l(1-t_l)}{p_l}\right)^{\nu_l} \left(\frac{t_l p_l}{u_l(1-t_l)(1+p_l)}\right)^{i_{l-1}} \tag{6.3.23}
\end{aligned}$$

Substitute (6.3.23) into (6.3.11) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(z)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(z)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(z)$, etc.¹¹

Theorem 6.3.2 *The general representation in the form of integral of the GCH polynomial of type 1 about $x = \infty$ is given by*

$$\begin{aligned}
y(z) &= c_0 z^\lambda \left\{ \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \eta^{i_0} + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_{n-k} e^{-u_{n-k}} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \int_\infty^{(0+)} dp_{n-k} p_{n-k}^{-1} (1+p_{n-k})^{-\frac{1}{2}(n-k+\lambda)} \exp\left(\frac{p_{n-k}}{w_{n-k+1,n}(1-t_{n-k})}\right) \\
&\quad \times \left(\frac{w_{n-k+1,n} u_{n-k}(1-t_{n-k})}{p_{n-k}}\right)^{\nu_{n-k}} \\
&\quad \times w_{n-k,n}^{-\frac{1}{2}(n-k-1-\omega+\lambda)} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(n-k-1-\omega+\lambda)} \left. \right\} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} w_{1,n}^{i_0} \left. \right\} \xi^n \tag{6.3.24}
\end{aligned}$$

where

$$w_{i,j} = \begin{cases} \frac{t_i p_i}{u_i(1-t_i)(1+p_i)} & \text{where } i \leq j \\ \eta & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of A'_n 's, the second one contains two terms of A_n 's, the third one contains three terms of A_n 's, etc.

¹¹ $y_1(z)$ means the sub-power series in (6.3.11) contains one term of A'_n 's, $y_2(x)$ means the sub-power series in (6.3.11) contains two terms of A'_n 's, $y_3(x)$ means the sub-power series in (6.3.11) contains three terms of A'_n 's, etc.

Proof In (6.3.11) the power series expansions of sub-summation $y_0(z)$, $y_1(z)$, $y_2(z)$ and $y_3(z)$ of the GCH polynomial of type 1 about $x = \infty$ are

$$y(z) = \sum_{n=0}^{\infty} c_n z^{n+\lambda} = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \quad (6.3.25)$$

where

$$y_0(z) = \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \eta^{i_0} \quad (6.3.26a)$$

$$\begin{aligned} y_1(z) = & c_0 z^\lambda \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{\left(i_0 + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \right. \\ & \times \left. \sum_{i_1=i_0}^{\nu_1} \frac{(-\nu_1)_{i_1} \left(\frac{1}{2} + \frac{\lambda}{2}\right)_{i_1} \left(\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}}{(-\nu_1)_{i_0} \left(\frac{1}{2} + \frac{\lambda}{2}\right)_{i_0} \left(\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_1}} \eta^{i_1} \right\} \xi \end{aligned} \quad (6.3.26b)$$

$$\begin{aligned} y_2(z) = & c_0 z^\lambda \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{\left(i_0 + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \right. \\ & \times \sum_{i_1=i_0}^{\nu_1} \frac{(i_1 + \frac{1}{2} - \frac{\omega}{2} + \frac{\lambda}{2})}{\left(i_1 + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)} \frac{(-\nu_1)_{i_1} \left(\frac{1}{2} + \frac{\lambda}{2}\right)_{i_1} \left(\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}}{(-\nu_1)_{i_0} \left(\frac{1}{2} + \frac{\lambda}{2}\right)_{i_0} \left(\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_1}} \\ & \times \left. \sum_{i_2=i_1}^{\nu_2} \frac{(-\nu_2)_{i_2} \left(1 + \frac{\lambda}{2}\right)_{i_2} \left(2 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_1}}{(-\nu_2)_{i_1} \left(1 + \frac{\lambda}{2}\right)_{i_1} \left(2 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_2}} \eta^{i_2} \right\} \xi^2 \end{aligned} \quad (6.3.26c)$$

$$\begin{aligned}
y_3(z) = & c_0 z^\lambda \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{(i_0 + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\nu_1} \frac{(i_1 + \frac{1}{2} - \frac{\omega}{2} + \frac{\lambda}{2})}{(i_1 + 1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \frac{(-\nu_1)_{i_1} (\frac{1}{2} + \frac{\lambda}{2})_{i_1} (\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}}{(-\nu_1)_{i_0} (\frac{1}{2} + \frac{\lambda}{2})_{i_0} (\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_1}} \\
& \times \sum_{i_2=i_1}^{\nu_2} \frac{(i_2 + 1 - \frac{\omega}{2} + \frac{\lambda}{2})}{(i_2 + \frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})} \frac{(-\nu_2)_{i_2} (1 + \frac{\lambda}{2})_{i_2} (2 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_1}}{(-\nu_2)_{i_1} (1 + \frac{\lambda}{2})_{i_1} (2 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_2}} \\
& \left. \times \sum_{i_3=i_2}^{\nu_3} \frac{(-\nu_3)_{i_3} (\frac{3}{2} + \frac{\lambda}{2})_{i_3} (\frac{5}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_2}}{(-\nu_3)_{i_2} (\frac{3}{2} + \frac{\lambda}{2})_{i_2} (\frac{5}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_3}} \eta^{i_3} \right\} \xi^3 \quad (6.3.26d)
\end{aligned}$$

Put $l = 1$ in (6.3.23). Take the new (6.3.23) into (6.3.26b).

$$\begin{aligned}
y_1(z) = & c_0 z^\lambda \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_1 \exp\left(\frac{p_1}{\eta(1-t_1)}\right) p_1^{-1} (1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
& \times \left(\frac{\eta u_1(1-t_1)}{p_1}\right)^{\nu_1} \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} \left(\frac{t_1 p_1}{u_1(1-t_1)(1+p_1)}\right)^{i_0} \right\} \xi \\
= & c_0 z^\lambda \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_1 \exp\left(\frac{p_1}{\eta(1-t_1)}\right) p_1^{-1} (1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
& \times \left(\frac{\eta u_1(1-t_1)}{p_1}\right)^{\nu_1} w_{1,1}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\omega+\lambda)} \\
& \times \left\{ \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} (\frac{\lambda}{2})_{i_0}}{(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2})_{i_0}} w_{1,1}^{i_0} \right\} \xi \quad (6.3.27)
\end{aligned}$$

where $w_{1,1} = \frac{t_1 p_1}{u_1(1-t_1)(1+p_1)}$

Put $l = 2$ in (6.3.23). Take the new (6.3.23) into (6.3.26c).

$$\begin{aligned}
 y_2(z) &= c_0 z^\lambda \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 e^{-u_2} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_2 \exp\left(\frac{p_2}{\eta(1-t_2)}\right) p_2^{-1} (1+p_2)^{-\frac{1}{2}(2+\lambda)} \\
 &\quad \times \left(\frac{\eta u_2(1-t_2)}{p_2}\right)^{\nu_2} w_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
 &\quad \times \left\{ \sum_{i_0=0}^{\nu_0} \frac{(i_0 - \frac{\omega}{2} + \frac{\lambda}{2})}{\left(i_0 + \frac{1}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \sum_{i_1=i_0}^{\nu_1} \frac{(-\nu_1)_{i_1} \left(\frac{1}{2} + \frac{\lambda}{2}\right)_{i_1} \left(\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}}{(-\nu_1)_{i_0} \left(\frac{1}{2} + \frac{\lambda}{2}\right)_{i_0} \left(\frac{3}{2} - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_1}} w_{2,2}^{i_1} \right\} \xi^2 \\
 \text{where } w_{2,2} &= \frac{t_2 p_2}{u_2(1-t_2)(1+p_2)}
 \end{aligned} \tag{6.3.28}$$

Put $l = 1$ and $\eta = w_{2,2}$ in (6.3.23). Take the new (6.3.23) into (6.3.28).

$$\begin{aligned}
 y_2(z) &= c_0 z^\lambda \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 e^{-u_2} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_2 \exp\left(\frac{p_2}{\eta(1-t_2)}\right) p_2^{-1} (1+p_2)^{-\frac{1}{2}(2+\lambda)} \\
 &\quad \times \left(\frac{\eta u_2(1-t_2)}{p_2}\right)^{\nu_2} w_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (w_{2,2} \partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
 &\quad \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_1 \exp\left(\frac{p_1}{w_{2,2}(1-t_1)}\right) p_1^{-1} (1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
 &\quad \times \left(\frac{w_{2,2} u_1(1-t_1)}{p_1}\right)^{\nu_1} w_{1,2}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\omega+\lambda)} \\
 &\quad \times \left\{ \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} w_{1,2}^{i_0} \right\} \xi^2 \\
 \text{where } w_{1,2} &= \frac{t_1 p_1}{u_1(1-t_1)(1+p_1)}
 \end{aligned} \tag{6.3.29}$$

By using similar process for the previous cases of integral forms of $y_1(z)$ and $y_2(z)$, the

integral form of sub-power series expansion of $y_3(z)$ is

$$\begin{aligned}
y_3(z) = & c_0 z^\lambda \int_0^1 dt_3 t_3^{\frac{1}{2}\left(1-\frac{\Omega}{\mu}+\lambda\right)} \int_0^1 du_3 e^{-u_3} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_3 \exp\left(\frac{p_3}{\eta(1-t_3)}\right) p_3^{-1} (1+p_3)^{-\frac{1}{2}(3+\lambda)} \\
& \times \left(\frac{\eta u_3(1-t_3)}{p_3}\right)^{\nu_3} w_{3,3}^{-\frac{1}{2}(2-\omega+\lambda)} (w_{3,3} \partial_{w_{3,3}}) w_{3,3}^{\frac{1}{2}(2-\omega+\lambda)} \\
& \times \int_0^1 dt_2 t_2^{\frac{1}{2}\left(-\frac{\Omega}{\mu}+\lambda\right)} \int_0^1 du_2 e^{-u_2} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_2 \exp\left(\frac{p_2}{w_{3,3}(1-t_2)}\right) p_2^{-1} (1+p_2)^{-\frac{1}{2}(2+\lambda)} \\
& \times \left(\frac{w_{3,3} u_2(1-t_2)}{p_2}\right)^{\nu_2} w_{2,3}^{-\frac{1}{2}(1-\omega+\lambda)} (w_{2,3} \partial_{w_{2,3}}) w_{2,3}^{\frac{1}{2}(1-\omega+\lambda)} \\
& \times \int_0^1 dt_1 t_1^{\frac{1}{2}\left(-1-\frac{\Omega}{\mu}+\lambda\right)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_\infty^{(0+)} dp_1 \exp\left(\frac{p_1}{w_{2,3}(1-t_1)}\right) p_1^{-1} (1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
& \times \left(\frac{w_{2,3} u_1(1-t_1)}{p_1}\right)^{\nu_1} w_{1,3}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,3} \partial_{w_{1,3}}) w_{1,3}^{\frac{1}{2}(-\omega+\lambda)} \\
& \times \left\{ \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1-\frac{\Omega}{2\mu}+\frac{\lambda}{2}\right)_{i_0}} w_{1,3}^{i_0} \right\} \xi^3 \tag{6.3.30}
\end{aligned}$$

where

$$\begin{cases} w_{3,3} = \frac{t_3 p_3}{u_3(1-t_3)(1+p_3)} \\ w_{2,3} = \frac{t_2 p_2}{u_2(1-t_2)(1+p_2)} \\ w_{1,3} = \frac{t_1 p_1}{u_1(1-t_1)(1+p_1)} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(z)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(z)$ terms. Since we substitute (6.3.26a), (6.3.27), (6.3.29), (6.3.30) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (6.3.25), we obtain (6.3.24). ■

Remark 6.3.3 *The integral representation of the GCH equation of the first kind for*

polynomial of type 1 about $x = \infty$ as $\nu = 2\nu_j + j + 1 + \Omega/\mu$ where $j, \nu_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(z) &= Q^{(i)} W_{\nu_j} \left(\mu, \varepsilon, \Omega, \omega, \nu = 2\nu_j + j + 1 + \frac{\Omega}{\mu}; z = \frac{1}{x}; \xi = -\frac{\varepsilon}{\mu} z; \eta = \frac{2}{\mu} z^2 \right) \\
&= z^{\frac{\Omega}{\mu}} \left\{ (-\eta)^{\nu_0} U \left(-\nu_0, -\nu_0 + 1 - \frac{\Omega}{\mu}, -\eta^{-1} \right) \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{\frac{1}{2}(n-k-2)} \int_0^1 du_{n-k} e^{-u_{n-k}} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_{n-k} p_{n-k}^{-1} (1 + p_{n-k})^{-\frac{1}{2}(n-k+\frac{\Omega}{\mu})} \exp \left(\frac{p_{n-k}}{w_{n-k+1,n}(1-t_{n-k})} \right) \\
&\quad \times \left(\frac{w_{n-k+1,n} u_{n-k} (1-t_{n-k})}{p_{n-k}} \right)^{\nu_{n-k}} w_{n-k,n}^{-\frac{1}{2}(n-k-1-\omega+\frac{\Omega}{\mu})} (w_{n-k,n} \partial_{w_{n-k,n}}) w_{n-k,n}^{\frac{1}{2}(n-k-1-\omega+\frac{\Omega}{\mu})} \left. \right\} \\
&\quad \times (-w_{1,n})^{\nu_0} U \left(-\nu_0, -\nu_0 + 1 - \frac{\Omega}{\mu}, -w_{1,n}^{-1} \right) \left. \right\} \xi^n \quad (6.3.31)
\end{aligned}$$

Proof Replace a , b and z by $-\nu_0$, $-\nu_0 + 1 - \frac{\Omega}{2\mu}$ and $-\eta^{-1}$ into (6.3.18).

$$\sum_{j=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \eta^j = (-\eta)^{\nu_0} U \left(-\nu_0, -\nu_0 + 1 - \frac{\Omega}{\mu}, -\eta^{-1} \right) \quad (6.3.32)$$

Replace a , b and z by $-\nu_0$, $-\nu_0 + 1 - \frac{\Omega}{2\mu}$ and $-w_{1,n}^{-1}$ into (6.3.18).

$$\sum_{j=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} w_{1,n}^j = (-w_{1,n})^{\nu_0} U \left(-\nu_0, -\nu_0 + 1 - \frac{\Omega}{\mu}, -w_{1,n}^{-1} \right) \quad (6.3.33)$$

Put $c_0 = 1$ and $\lambda = \Omega/\mu$ in (6.3.24). Substitute (6.3.32) and (6.3.33) into the new (6.3.24). ■

6.3.3 Generating function of the GCH polynomial of type 1

Let's investigate the generating function for the GCH polynomial of type 1 of the first kind about $x = \infty$.

Definition 6.3.4 *I define that*

$$\begin{cases} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{where } a > b \\ s_a & \text{only if } a = b \end{cases} \\ \tilde{w}_{i,j} = \begin{cases} \frac{s_i t_i \tilde{w}_{i+1,j}}{1 + s_i u_i (1 - t_i) \tilde{w}_{i+1,j}} & \text{where } i < j \\ \frac{s_i t_i \eta}{1 + s_i u_i (1 - t_i) \eta} & \text{only if } i = j \end{cases} \end{cases} \quad (6.3.34)$$

where

$$a, b, i, j \in \mathbb{N}_0$$

And we have

$$\sum_{\nu_i=\nu_j}^{\infty} s_i^{\nu_i} = \frac{s_i^{\nu_j}}{(1 - s_i)} \quad (6.3.35)$$

Acting the summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\}$ on (6.3.24) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (6.3.34) and (6.3.35),

Theorem 6.3.5 *The general expression of the generating function for the GCH*

polynomial of type 1 about $x = \infty$ is given by

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y(z) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \Upsilon(\lambda; s_{0,\infty}; \eta) \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^{\infty} du_1 \exp(-(1-s_{1,\infty})u_1) (1+s_{1,\infty}u_1(1-t_1)\eta)^{-\frac{1}{2}(1+\lambda)} \\
&\times \tilde{w}_{1,1}^{-\frac{1}{2}(-\omega+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\omega+\lambda)} \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) \xi \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{k=n+1}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_n t_n^{\frac{1}{2}(n-2-\frac{\Omega}{\mu}+\lambda)} \int_0^{\infty} du_n \exp(-(1-s_{n,\infty})u_n) \right. \\
&\times (1+s_{n,\infty}u_n(1-t_n)\eta)^{-\frac{1}{2}(n+\lambda)} \tilde{w}_{n,n}^{-\frac{1}{2}(n-1-\omega+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(n-1-\omega+\lambda)} \\
&\times \prod_{j=1}^{n-1} \left\{ \int_0^1 dt_{n-j} t_{n-j}^{\frac{1}{2}(n-j-2-\frac{\Omega}{\mu}+\lambda)} \int_0^{\infty} du_{n-j} \exp(-(1-s_{n-j})u_{n-j}) \right. \\
&\times (1+s_{n-j}u_{n-j}(1-t_{n-j})\tilde{w}_{n-j+1,n})^{-\frac{1}{2}(n-j+\lambda)} \\
&\times \left. \left. \tilde{w}_{n-j,n}^{-\frac{1}{2}(n-j-1-\omega+\lambda)} (\tilde{w}_{n-j,n} \partial_{\tilde{w}_{n-j,n}}) \tilde{w}_{n-j,n}^{\frac{1}{2}(n-j-1-\omega+\lambda)} \right\} \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) \right\} \xi^n
\end{aligned} \tag{6.3.36}$$

where

$$\begin{cases} \Upsilon(\lambda; s_{0,\infty}; \eta) = \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^\lambda \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \eta^{i_0} \right\} \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) = \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^\lambda \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \tilde{w}_{1,1}^{i_0} \right\} \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) = \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^\lambda \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \tilde{w}_{1,n}^{i_0} \right\} \end{cases}$$

Proof Acting the summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\}$ on the form of integral

of the type 1 GCH polynomial $y(z)$,

$$\begin{aligned} & \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y(z) \\ &= \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} (y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots) \end{aligned} \quad (6.3.37)$$

Acting the summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\}$ on (6.3.26a) by using (6.3.34) and (6.3.35),

$$\begin{aligned} & \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_0(z) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \sum_{\nu_0=0}^{\infty} \frac{s_{0,\infty}^{\nu_0}}{\nu_0!} \left\{ c_0 z^\lambda \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \eta^{i_0} \right\} \end{aligned} \quad (6.3.38)$$

Acting the summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\}$ on (6.3.27) by using (6.3.34) and (6.3.35),

$$\begin{aligned} & \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_1(z) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_1 \exp\left(\frac{p_1}{\eta(1-t_1)}\right) \\ & \times p_1^{-1} (1+p_1)^{-\frac{1}{2}(1+\lambda)} \sum_{\nu_1=\nu_0}^{\infty} \left(\frac{s_{1,\infty} \eta u_1 (1-t_1)}{p_1} \right)^{\nu_1} w_{1,1}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\omega+\lambda)} \\ & \times \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^\lambda \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} w_{1,1}^{i_0} \right\} \xi \end{aligned} \quad (6.3.39)$$

Replace ν_i , ν_j and s_i by ν_1 , ν_0 and $\frac{s_{1,\infty} \eta u_1 (1-t_1)}{p_1}$ in (6.3.35). Take the new (6.3.35)

into (6.3.39).

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_1(z) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_1 \exp\left(\frac{p_1}{\eta(1-t_1)}\right) \\
&\times \frac{(1+p_1)^{-\frac{1}{2}(1+\lambda)}}{p_1 - s_{1,\infty}\eta u_1(1-t_1)} w_{1,1}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,1} \partial_{w_{1,1}}) w_{1,1}^{\frac{1}{2}(-\omega+\lambda)} \\
&\times \sum_{\nu_0=0}^{\infty} \left(\frac{s_{0,\infty}\eta u_1(1-t_1)}{p_1} \right)^{\nu_0} \frac{1}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} w_{1,1}^{i_0} \right\} \xi \quad (6.3.40)
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $p_1 = s_{1,\infty}\eta u_1(1-t_1)$, and $s_{1,\infty}\eta u_1(1-t_1)$ is inside the unit circle. As we compute the residue there in (6.3.40) we obtain

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_1(z) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 \exp(-(1-s_{1,\infty})u_1) (1+s_{1,\infty}u_1(1-t_1)\eta)^{-\frac{1}{2}(1+\lambda)} \\
&\times \tilde{w}_{1,1}^{-\frac{1}{2}(-\omega+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\omega+\lambda)} \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \tilde{w}_{1,1}^{i_0} \right\} \xi \quad (6.3.41)
\end{aligned}$$

where

$$\tilde{w}_{1,1} = \frac{t_1 p_1}{u_1(1-t_1)(1+p_1)} \Big|_{p_1=s_{1,\infty}\eta u_1(1-t_1)} = \frac{s_{1,\infty} t_1 \eta}{1+s_{1,\infty} u_1(1-t_1)\eta}$$

Acting the summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\}$ on (6.3.29) by using (6.3.34)

and (6.3.35),

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 e^{-u_2} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_2 \exp\left(\frac{p_2}{\eta(1-t_2)}\right) \\
&\times p_2^{-1}(1+p_2)^{-\frac{1}{2}(2+\lambda)} \sum_{\nu_2=\nu_1}^{\infty} \left(\frac{s_{2,\infty}\eta u_2(1-t_2)}{p_2}\right)^{\nu_2} w_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_1 \exp\left(\frac{p_1}{w_{2,2}(1-t_1)}\right) p_1^{-1}(1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
&\times \sum_{\nu_1=\nu_0}^{\infty} \left(\frac{s_1 w_{2,2} u_1 (1-t_1)}{p_1}\right)^{\nu_1} w_{1,2}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\omega+\lambda)} \\
&\times \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1-\frac{\Omega}{2\mu}+\frac{\lambda}{2}\right)_{i_0}} w_{1,2}^{i_0} \right\} \xi^2 \tag{6.3.42}
\end{aligned}$$

Replace ν_i , ν_j and s_i by ν_2 , ν_1 and $\frac{s_{2,\infty}\eta u_2(1-t_2)}{p_2}$ in (6.3.35). Take the new (6.3.35) into (6.3.42).

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 e^{-u_2} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_2 \exp\left(\frac{p_2}{\eta(1-t_2)}\right) \\
&\times \frac{(1+p_2)^{-\frac{1}{2}(2+\lambda)}}{p_2 - s_{2,\infty}\eta u_2(1-t_2)} w_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (w_{2,2}\partial_{w_{2,2}}) w_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_1 \exp\left(\frac{p_1}{w_{2,2}(1-t_1)}\right) p_1^{-1}(1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
&\times \sum_{\nu_1=\nu_0}^{\infty} \left(\frac{s_{1,\infty}\eta u_2(1-t_2)w_{2,2}u_1(1-t_1)}{p_1 p_2}\right)^{\nu_1} w_{1,2}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,2}\partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\omega+\lambda)} \\
&\times \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1-\frac{\Omega}{2\mu}+\frac{\lambda}{2}\right)_{i_0}} w_{1,2}^{i_0} \right\} \xi^2 \tag{6.3.43}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $p_2 = s_{2,\infty}\eta u_2(1-t_2)$, and $s_{2,\infty}\eta u_2(1-t_2)$ is inside the unit circle. As we compute the residue there in (6.3.43) we obtain

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 \exp(-(1-s_{2,\infty})u_2) (1+s_{2,\infty}u_2(1-t_2)\eta)^{-\frac{1}{2}(2+\lambda)} \\
&\times \tilde{w}_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
&\times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_1 \exp\left(\frac{p_1}{\tilde{w}_{2,2}(1-t_1)}\right) p_1^{-1} (1+p_1)^{-\frac{1}{2}(1+\lambda)} \\
&\times \sum_{\nu_1=\nu_0}^{\infty} \left(\frac{s_1 \tilde{w}_{2,2} u_1 (1-t_1)}{p_1} \right)^{\nu_1} w_{1,2}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\omega+\lambda)} \\
&\times \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1-\frac{\Omega}{2\mu}+\frac{\lambda}{2}\right)_{i_0}} w_{1,2}^{i_0} \right\} \xi^2 \tag{6.3.44}
\end{aligned}$$

where

$$\tilde{w}_{2,2} = \frac{t_2 p_2}{u_2(1-t_2)(1+p_2)} \Big|_{p_2=s_{2,\infty}\eta u_2(1-t_2)} = \frac{s_{2,\infty} t_2 \eta}{1+s_{2,\infty} u_2(1-t_2)\eta}$$

Replace ν_i , ν_j and s_i by ν_1 , ν_0 and $\frac{s_1 \tilde{w}_{2,2} u_1 (1-t_1)}{p_1}$ in (6.3.35). Take the new (6.3.35)

into (6.3.44).

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 \exp(-(1-s_{2,\infty})u_2) (1+s_{2,\infty}u_2(1-t_2)\eta)^{-\frac{1}{2}(2+\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
&\quad \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 e^{-u_1} \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp_1 \exp\left(\frac{p_1}{\tilde{w}_{2,2}(1-t_1)}\right) \frac{(1+p_1)^{-\frac{1}{2}(1+\lambda)}}{p_1 - s_1 \tilde{w}_{2,2} u_1 (1-t_1)} \\
&\quad \times w_{1,2}^{-\frac{1}{2}(-\omega+\lambda)} (w_{1,2} \partial_{w_{1,2}}) w_{1,2}^{\frac{1}{2}(-\omega+\lambda)} \\
&\quad \times \sum_{\nu_0=0}^{\infty} \left(\frac{s_{0,1} \tilde{w}_{2,2} u_1 (1-t_1)}{p_1} \right)^{\nu_0} \frac{1}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} w_{1,2}^{i_0} \right\} \xi^2 \tag{6.3.45}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at $p_1 = s_1 \tilde{w}_{2,2} u_1 (1-t_1)$, and $s_1 \tilde{w}_{2,2} u_1 (1-t_1)$ is inside the unit circle. As we compute the residue there in (6.3.45) we obtain

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 \exp(-(1-s_{2,\infty})u_2) (1+s_{2,\infty}u_2(1-t_2)\eta)^{-\frac{1}{2}(2+\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-\frac{1}{2}(1-\omega+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\frac{1}{2}(1-\omega+\lambda)} \\
&\quad \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 \exp(-(1-s_1)u_1) (1+s_1 u_1 (1-t_1) \tilde{w}_{2,2})^{-\frac{1}{2}(1+\lambda)} \\
&\quad \times \tilde{w}_{1,2}^{-\frac{1}{2}(-\omega+\lambda)} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\frac{1}{2}(-\omega+\lambda)} \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^{\lambda} \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1 - \frac{\Omega}{2\mu} + \frac{\lambda}{2}\right)_{i_0}} \tilde{w}_{1,2}^{i_0} \right\} \xi^2 \tag{6.3.46}
\end{aligned}$$

where

$$\tilde{w}_{1,2} = \frac{t_1 p_1}{u_1 (1-t_1) (1+p_1)} \Big|_{p_1=s_1 \tilde{w}_{2,2} u_1 (1-t_1)} = \frac{s_1 t_1 \tilde{w}_{2,2}}{1 + s_1 u_1 (1-t_1) \tilde{w}_{2,2}}$$

Acting the summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\}$ on (6.3.30) by using (6.3.34)

and (6.3.35),

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_3(z) \\
&= \prod_{k=4}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{\frac{1}{2}(1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_3 \exp(-(1-s_{3,\infty})u_3) (1+s_{3,\infty}u_3(1-t_3)\eta)^{-\frac{1}{2}(3+\lambda)} \\
&\quad \times \tilde{w}_{3,3}^{-\frac{1}{2}(2-\omega+\lambda)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{\frac{1}{2}(2-\omega+\lambda)} \\
&\quad \times \int_0^1 dt_2 t_2^{\frac{1}{2}(-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_2 \exp(-(1-s_2)u_2) (1+s_2u_2(1-t_2)\tilde{w}_{3,3})^{-\frac{1}{2}(2+\lambda)} \\
&\quad \times \tilde{w}_{2,3}^{-\frac{1}{2}(1-\omega+\lambda)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{\frac{1}{2}(1-\omega+\lambda)} \\
&\quad \times \int_0^1 dt_1 t_1^{\frac{1}{2}(-1-\frac{\Omega}{\mu}+\lambda)} \int_0^1 du_1 \exp(-(1-s_1)u_1) (1+s_1u_1(1-t_1)\tilde{w}_{2,3})^{-\frac{1}{2}(1+\lambda)} \\
&\quad \times \tilde{w}_{1,3}^{-\frac{1}{2}(-\omega+\lambda)} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\frac{1}{2}(-\omega+\lambda)} \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \left\{ c_0 z^\lambda \sum_{i_0=0}^{\nu_0} \frac{(-\nu_0)_{i_0} \left(\frac{\lambda}{2}\right)_{i_0}}{\left(1-\frac{\Omega}{2\mu}+\frac{\lambda}{2}\right)_{i_0}} \tilde{w}_{1,3}^{i_0} \right\} \xi^3 \quad (6.3.47)
\end{aligned}$$

where

$$\begin{cases} \tilde{w}_{3,3} = \frac{t_3 p_3}{u_3(1-t_3)(1+p_3)} \Big|_{p_3=s_{3,\infty}\eta u_3(1-t_3)} = \frac{s_{3,\infty} t_3 \eta}{1+s_{3,\infty} u_3(1-t_3)\eta} \\ \tilde{w}_{2,3} = \frac{t_2 p_2}{u_2(1-t_2)(1+p_2)} \Big|_{p_2=s_2 \tilde{w}_{3,3} u_2(1-t_2)} = \frac{s_2 t_2 \tilde{w}_{3,3}}{1+s_2 u_2(1-t_2)\tilde{w}_{3,3}} \\ \tilde{w}_{1,3} = \frac{t_1 p_1}{u_1(1-t_1)(1+p_1)} \Big|_{p_1=s_1 \tilde{w}_{2,3} u_1(1-t_1)} = \frac{s_1 t_1 \tilde{w}_{2,3}}{1+s_1 u_1(1-t_1)\tilde{w}_{2,3}} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(z)$

terms where $m > 3$, we obtain every $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_m(z)$ terms. Since we substitute (6.3.38), (6.3.41), (6.3.46), (6.3.47) and including all

$$\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} y_m(z) \text{ terms where } m > 3 \text{ into (6.3.37), we obtain (6.3.36) } \blacksquare$$

Remark 6.3.6 *The generating function for the GCH polynomial of type 1 of the first*

kind about $x = \infty$ as $\nu = 2\nu_j + j + 1 + \Omega/\mu$ where $j, \nu_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \prod_{n=1}^{\infty} \left\{ \sum_{\nu_n=\nu_{n-1}}^{\infty} s_n^{\nu_n} \right\} Q^{(i)} W_{\nu_j} \left(\mu, \varepsilon, \Omega, \omega, \nu = 2\nu_j + j + 1 + \frac{\Omega}{\mu}; z = \frac{1}{x}; \xi = -\frac{\varepsilon}{\mu} z; \eta = \frac{2}{\mu} z^2 \right) \\
&= z^{\frac{\Omega}{\mu}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) \right. \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{-\frac{1}{2}} \int_0^{\infty} du_1 \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) \\
&\times \tilde{w}_{1,1}^{-\frac{1}{2}(-\omega+\frac{\Omega}{\mu})} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(-\omega+\frac{\Omega}{\mu})} \mathbf{A}(s_0; \tilde{w}_{1,1}) \xi \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{k=n+1}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_n t_n^{\frac{1}{2}(n-2)} \int_0^{\infty} du_n \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \tilde{w}_{n,n}^{-\frac{1}{2}(n-1-\omega+\frac{\Omega}{\mu})} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(n-1-\omega+\frac{\Omega}{\mu})} \\
&\times \prod_{j=1}^{n-1} \left\{ \int_0^1 dt_{n-j} t_{n-j}^{\frac{1}{2}(n-j-2)} \int_0^{\infty} du_{n-j} \overleftrightarrow{\Gamma}_{n-j}(s_{n-j}; t_{n-j}, u_{n-j}, \tilde{w}_{n-j+1,n}) \right. \\
&\times \tilde{w}_{n-j,n}^{-\frac{1}{2}(n-j-1-\omega+\frac{\Omega}{\mu})} (\tilde{w}_{n-j,n} \partial_{\tilde{w}_{n-j,n}}) \tilde{w}_{n-j,n}^{\frac{1}{2}(n-j-1-\omega+\frac{\Omega}{\mu})} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} \xi^n \Bigg\} \quad (6.3.48)
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \exp(-(1-s_{1,\infty})u_1) (1+s_{1,\infty}u_1(1-t_1)\eta)^{-\frac{1}{2}(1+\frac{\Omega}{\mu})} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \exp(-(1-s_{n,\infty})u_n) (1+s_{n,\infty}u_n(1-t_n)\eta)^{-\frac{1}{2}(n+\frac{\Omega}{\mu})} \\ \overleftrightarrow{\Gamma}_{n-j}(s_{n-j}; t_{n-j}, u_{n-j}, \tilde{w}_{n-j+1,n}) \\ = \exp(-(1-s_{n-j})u_{n-j}) (1+s_{n-j}u_{n-j}(1-t_{n-j})\tilde{w}_{n-j+1,n})^{-\frac{1}{2}(n-j+\frac{\Omega}{\mu})} \end{cases}$$

and

$$\begin{cases} \mathbf{A}(s_{0,\infty}; \eta) = \exp(s_{0,\infty}) (1+s_{0,\infty}\eta)^{-\frac{\Omega}{2\mu}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \exp(s_0) (1+s_0\tilde{w}_{1,1})^{-\frac{\Omega}{2\mu}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \exp(s_0) (1+s_0\tilde{w}_{1,n})^{-\frac{\Omega}{2\mu}} \end{cases}$$

Proof Replace a, b, j and z by $-\nu_0, -\nu_0 + 1 - a, i_0$ and $-z^{-1}$ in (6.3.18). Acting the

summation operator $\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!}$ on the new (6.3.18)

$$\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \sum_{i_0=\nu_0}^{\infty} \frac{(-\nu_0)_{i_0} (a)_{i_0}}{(1)_{i_0}} z^{i_0} = \sum_{\nu_0=0}^{\infty} \frac{(-s_0 z)^{\nu_0}}{\nu_0!} U(-\nu_0, -\nu_0 + 1 - a, -z^{-1}) \quad (6.3.49)$$

Replace a, b, p_l and z by $-\nu_0, -\nu_0 + 1 - a, p$ and $-z^{-1}$ in (6.3.17).

$$U(-\nu_0, -\nu_0 + 1 - a, -z^{-1}) = e^{\nu_0 \pi i} \frac{\nu_0!}{2\pi i} \int_{\infty}^{(0+)} dp e^{\frac{p}{z}} p^{-\nu_0-1} (1+p)^{-a} \quad (6.3.50)$$

Put (6.3.50) in (6.3.49).

$$\begin{aligned} \sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \sum_{i_0=\nu_0}^{\infty} \frac{(-\nu_0)_{i_0} (a)_{i_0}}{(1)_{i_0}} z^{i_0} &= \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp e^{\frac{p}{z}} p^{-1} (1+p)^{-a} \sum_{\nu_0=0}^{\infty} \left(\frac{s_0 z}{p} \right)^{\nu_0} \\ &= \frac{1}{2\pi i} \int_{\infty}^{(0+)} dp e^{\frac{p}{z}} \frac{(1+p)^{-a}}{(p - s_0 z)} \\ &= \exp(s_0) (1 + s_0 z)^{-a} \end{aligned} \quad (6.3.51)$$

Replace s_0, a and z by $s_{0,\infty}, \frac{\Omega}{2\mu}$ and η in (6.3.51).

$$\sum_{\nu_0=0}^{\infty} \frac{s_{0,\infty}^{\nu_0}}{\nu_0!} \sum_{i_0=\nu_0}^{\infty} \frac{(-\nu_0)_{i_0} \left(\frac{\Omega}{2\mu} \right)_{i_0}}{(1)_{i_0}} \eta^{i_0} = \exp(s_{0,\infty}) (1 + s_{0,\infty} \eta)^{-\frac{\Omega}{2\mu}} \quad (6.3.52)$$

Replace a and z by $\frac{\Omega}{2\mu}$ and $\tilde{w}_{1,1}$ in (6.3.51).

$$\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \sum_{i_0=\nu_0}^{\infty} \frac{(-\nu_0)_{i_0} \left(\frac{\Omega}{2\mu} \right)_{i_0}}{(1)_{i_0}} \tilde{w}_{1,1}^{i_0} = \exp(s_0) (1 + s_0 \tilde{w}_{1,1})^{-\frac{\Omega}{2\mu}} \quad (6.3.53)$$

Replace a and z by $\frac{\Omega}{2\mu}$ and $\tilde{w}_{1,n}$ in (6.3.51).

$$\sum_{\nu_0=0}^{\infty} \frac{s_0^{\nu_0}}{\nu_0!} \sum_{i_0=\nu_0}^{\infty} \frac{(-\nu_0)_{i_0} \left(\frac{\Omega}{2\mu} \right)_{i_0}}{(1)_{i_0}} \tilde{w}_{1,n}^{i_0} = \exp(s_0) (1 + s_0 \tilde{w}_{1,n})^{-\frac{\Omega}{2\mu}} \quad (6.3.54)$$

Put $c_0 = 1$ and $\lambda = \frac{\Omega}{\mu}$ in (6.3.36). Substitute (6.3.52), (6.3.53) and (6.3.54) into the new (6.3.36). ■

6.4 Summary

The canonical form of the biconfluent Heun equation is defined by [2, 17]

$$x \frac{d^2 y}{dx^2} + (1 + \alpha - \beta x - 2x^2) \frac{dy}{dx} + \left((\gamma - \alpha - 2)x - \frac{1}{2}[\delta + (1 + \alpha)\beta] \right) y = 0 \quad (6.4.1)$$

in which $(\alpha, \beta, \gamma, \delta) \in \mathbb{C}^4$. This equation has two singular points which are a regular singularity at $x = 0$ and an irregular singularity at ∞ . This equation is derived from the GCH equation [34] by replacing all coefficients $\mu, \varepsilon, \nu, \Omega$ and ω by $-2, -\beta, 1 + \alpha, \gamma - \alpha - 2$ and $\frac{1}{2} \left(\frac{\delta}{\beta} + 1 + \alpha \right)$ in (6.1.1).

In my previous two papers [36, 37] I show how to derive the power series expansion in closed forms, its integral forms (each sub-integral $y_m(x)$ where $m = 0, 1, 2, \dots$ is composed of $2m$ terms of definite integrals and m terms of contour integrals) and the generating function for the GCH equation about $x = 0$ by applying 3TRF. [35]

In this chapter I show how to construct the power series expansion in closed forms and its integral forms of the GCH equation about $x = 0$ for infinite series and polynomial of type 2 by applying R3TRF. This is done by letting B_n in sequence c_n is the leading term in the analytic function $y(x)$. For polynomial of type 2, I treat ω as a fixed value and $\mu, \varepsilon, \nu, \Omega$ as free variables.

The power series expansion and the integral representation of the GCH equation for infinite series about $x = 0$ in this chapter are equivalent to infinite series of the GCH equation in Ref.[36, 37]. In this chapter B_n is the leading term in sequence c_n in the analytic function $y(x)$. In Ref.[36, 37] A_n is the leading term in sequence c_n in the analytic function $y(x)$.

In Ref.[36, 37], as we see the power series expansions of the GCH equation for infinite series and polynomial, the denominators and numerators in all B_n terms of each sub-power series expansion $y_m(x)$ where $m = 0, 1, 2, \dots$ arise with Pochhammer symbol. And in this chapter the denominators and numerators in all A_n terms of each sub-power series expansion $y_m(x)$ also arise with Pochhammer symbol. Since we construct the power series expansions with Pochhammer symbols in numerators and denominators, we are able to describe the integral representation of the GCH equation analytically. As we observe representations in closed form integrals of the GCH equation about $x = 0$ by applying either 3TRF or R3TRF, a ${}_1F_1$ function (the Kummer function of the first kind) recurs in each of sub-integral forms of the GCH equation. It means that we are able to transform the GCH (or BCH) function about $x = 0$ into any well-known special functions having two recursive coefficients in the power series of its ordinary differential equation because ${}_1F_1$ function arises in each of sub-integral forms on the GCH equation. After we

replace ${}_1F_1$ functions in its integral forms to other special functions, we can rebuild the Frobenius solution of the GCH equation about $x = 0$ in a backward.

In Ref.[37] and this chapter, I show how to derive the generating function for the type 1 and type 2 GCH polynomials from its analytic integral representation. And in this chapter I construct the generating function for the type 2 GCH polynomial from its integral representation. We are able to derive orthogonal relations, recursion relations and expectation values of the physical quantities from these two generating functions: the processes in order to obtain orthogonal and recursion relations of the GCH polynomial are similar as the case of a normalized wave function for the hydrogen-like atoms.¹²

In section 6.3 I construct the Frobenius solution of the GCH equation about $x = \infty$ for the type 1 polynomial by applying 3TRF analytically [35]. Also its integral representation and the generating function for the GCH equation are derived analytically. There are no such solutions for infinite series and the type 2 polynomial because the B_n term is divergent in (6.3.5b) since the index $n \gg 1$.¹³ In comparison with integral forms of the GCH polynomials of type 1 and 2 about $x = 0$, a Tricomi's function (Kummer's function of the second kind) recurs in each of sub-integral forms of the GCH polynomial of type 1 about $x = \infty$.

¹²For instance, in the quantum mechanical aspects, if the eigenenergy is contained in B_n term in its three different coefficients of the power series expansion, we have to apply the type 1 GCH polynomial. If the eigenenergy is included in A_n term in its three different coefficients, we should apply the type 2 GCH polynomial. If the first eigenenergy (mathematically, it is denoted by a spectral parameter) is included in A_n term and the second one is involved in B_n terms, we must apply the type 3 GCH polynomial. In my future paper I will discuss about the type 3 GCH polynomial.

¹³There are only two types of the analytic solution of the GCH equation about $x = \infty$ which are type 1 and type 3 polynomials. Its type 3 polynomial will be constructed in the future paper.

Bibliography

- [1] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische Annalen* **33**, (1889)161.
- [2] Ronveaux, A., “Heuns Differential Equations” Oxford University Press, (1995).
- [3] Slavyanov, S.Y., “Asymptotic Solutions of the One-dimensional Schrödinger Equation,” *Amer. Math. Soc. Trans. of Math. Monographs* **151**, (1996).
- [4] Ralko, A. and Truong, T.T., “Behavior of three charged particles on a plane under perpendicular magnetic field,” *J. Phys. A: Math. Theor.* **35**, 9573–9583(2002).
- [5] Kandemir, B.S., “Two interacting electrons in a uniform magnetic field and a parabolic potential: The general closed-form solution,” *J. Math. Phys.* **46**, 032110(2005).
- [6] M. Hortacsu, “Heun Functions and their uses in Physics,” arXiv:1101.0471 [math-ph].
- [7] Arriola, E.R., Zarzo, A., and Dehesa, J.S., “Spectral Properties of the Biconfluent Heun Differential Equation,” *J. Comput. Appl. Math.* **37**, 161–169(1991).
- [8] Masson, D., “The rotating harmonic oscillator eigenvalue problem,” *J. Math. Phys.* **24**, 2074–2088 and 2089–2094(1983).
- [9] Chaudhuri, R. N., “The Hill determinant: an application to a class of confinement potentials,” *J. Phys. A* **16**, 209-211(1983).
- [10] Chauduri, R.N. and Mukkerjee, B., “On the $\mu x^2 + \lambda x^4 + \eta x^6$ interaction,” *J. Phys. A: Math. Gen.* **17**, 3327–3335(1984).
- [11] Léauté, B. and Marcihacy, G., “On the Schrödinger equation of rotating harmonic, three-dimensional and double anharmonic oscillators and a class of confinement potentials in connexion with the biconfluent Heun differential equation,” *J. Phys. A: Math. Gen.* **19**, 3527–3533(1986).

- [12] Flessas, G. P. and Das, K. P., "On the three-dimensional anharmonic oscillator," Phys. Lett. **78**, no. 1, 19-21(1980).
- [13] Flessas, G. P. and Anagnostatos, G. S., "On the applicability of the Hill determinant and the analytic continued fraction method to anharmonic oscillators," J. Phys. A **15**, no. 10, L537-L542(1982).
- [14] Pons, R. and Marcihacy, G., "Contribution à l'étude de l'équation de Schrödinger avec un potentiel de Coulomb et un terme de confinement linéaire [Contribution to the study of the Schrödinger equation with a Coulomb potential and a linear confinement term]," (French) Ann. Soc. Sci. Bruxelles Sér. I **102**, 1-2 and 3-7(1988).
- [15] Léauté, B., Marcihacy, G., Pons, R. and Skinazi, J., "On the connection problem for some Schrödinger equations in relation to the biconfluent Heun differential equation," SIAM J. Math. Anal. **21**, no. 3, 793-798(1990).
- [16] Lemieux, A. and Bose, A. K., "Construction de potentiels pour lesquels l'équation de Schrödinger est soluble," (French) Ann. Inst. H. Poincaré Sect. A (N.S.) **10**, no. 3, 259-270(1969).
- [17] Maroni, P., "Sur la forme bi-confluente de l'équation de Heun," (French) C. R. Acad. Sci. Paris **264**, 503-505(1967).
- [18] NIST Digital Library of Mathematical Functions, "Confluent Forms of Heuns Equation," <http://dlmf.nist.gov/31.12>
- [19] NIST Digital Library of Mathematical Functions, "Confluent Hypergeometric Functions," <http://dlmf.nist.gov/13>
- [20] Slavyanov, S. Yu., Lay W. *Special Functions: A Unified Theory Based on Singularities*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000.
- [21] Decarreau, A., Maroni, P. and Robert, A., "Sur les équations confluentes de l'équation de Heun," (French) Ann. Soc. Sci. Bruxelles **92**, 151-189(1978).
- [22] Batola, F., "Quelques propriétés de l'équation biconfluente de l'équation de Heun," Thèse de troisième cycle, Univ. P. et M. Curie, Paris (1977).
- [23] Hautot, A., "Sur les solutions polynomiales de l'équation différentielle $zP_n'' + (az^2 + bz + c)P_n' + (d + ez + fz^2)P_n = 0$," Bull. Soc. R. Sci. Liège **40**, 13-23(1969).
- [24] Urwin, K.M., "Note on certain polynomial solutions of Heun's equation and two confluent forms," Bull. Soc. R. Sci. Liège **44**, 169-172(1975).

- [25] Meixner, J., “Die Greensche Funktion des wellenmechanischen Keplerproblems,” (German) *Mathematische Zeitschrift* **36** (1): 677-707(1933).
- [26] Exton, H., “On a non-trivial special case of the biconfluent Heun equation,” *Ann. Soc. Sci. Bruxelles* **102**, 87–95(1989).
- [27] Batola, F., “Une généralisation d’une formule d’Erdélyi-Tricomi,” *Ark. Mat.* **20**, 87–99(1982).
- [28] Batola, F., “Une généralisation d’une formule de Meixner-Tricomi,” *Can. J. Math.* **34**, 411-422(1982).
- [29] Maroni, P., “Sur quelques relations intégrales entre les solutions de l’équation biconfluente de Heun,” *Ann. Inst. H.P.* **30**, 315–332(1979).
- [30] Maroni, P., “Un problème aux limites pour l’équation représentative de la classe $(0, 1, 1_4)$,” *Publ. Lab. Anal. Num. Uni. P. et M. Curie-C.N.R.S., Paris*, no. **79004**, (1979).
- [31] Maroni, P., “Une représentation intégrale pour une solution particulière de l’équation biconfluente de Heun,” *Publ. Lab. Anal. Num. Uni. P. et M. Curie-C.N.R.S., Paris*, no. **84026**, (1984).
- [32] Chiang, Y.M. and Yu, G.F., “On complex oscillation theory, quasi-exact solvability and Fredholm integral equations,” *arXiv:1310.5507*.
- [33] S. Belmehdi, S. and Chehab, J.-P., “Integral representation of the solutions to Heun’s biconfluent equation,” *Abstract and Applied Analysis*, vol. 2004, no. **4**, 295–306(2004).
- [34] Choun, Y.S., “Approximative solution of the spin free Hamiltonian involving only scalar potential for the $q - \bar{q}$ system,” *arXiv:1302.7309*
- [35] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications.” *arXiv:1303.0806*.
- [36] Choun, Y.S., “Analytic solution for grand confluent hypergeometric function,” *arXiv:1303.0813*.
- [37] Choun, Y.S., “The integral formalism and the generating function of grand confluent hypergeometric function,” *arXiv:1303.0819*.

Chapter 7

Mathieu function using reversible three-term recurrence formula

In Ref.[21], by applying three term recurrence formula (3TRF)[20], I construct the power series expansion in closed forms and its integral forms of Mathieu equation for infinite series including all higher terms of A_n 's.¹

In this chapter I will apply reversible three term recurrence formula (R3TRF) in chapter 1 to (1) the power series expansion in closed forms, (2) its integral forms of Mathieu equation (for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's²) and (3) the generating function for Mathieu polynomial which makes A_n term terminated.

7.1 Introduction

Mathieu ordinary differential equation is of Fuchsian types with the two regular and one irregular singularities. In contrast, Heun equation of Fuchsian types has the four regular singularities. Heun equation has the four kind of confluent forms: (1) Confluent Heun (two regular and one irregular singularities), (2) Doubly confluent Heun (two irregular singularities), (3) Biconfluent Heun (one regular and one irregular singularities), (4) Triconfluent Heun equations (one irregular singularity). For DLFM version [19], Mathieu equation in algebraic forms is also derived from the Confluent Heun equation by changing

¹“ higher terms of A_n 's” means at least two terms of A_n 's.

²“ higher terms of B_n 's” means at least two terms of B_n 's.

all coefficients $\delta = \gamma = \frac{1}{2}$, $\epsilon = 0$, $\alpha = q$ and $q = \frac{\lambda+2q}{4}$.

The simplest example of three term recursion relations in the power series of a linear ordinary differential equation is the Mathieu equation, introduced by Mathieu (1868)[11], while investigating the vibration of an elliptical drumhead: it is derived from the Helmholtz equation in elliptic cylinder coordinates by using the method of separation of variables. Mathieu equation, known for the elliptic cylinder equation appears in diverse areas such as astronomy and physical problems involving Schrödinger equation for a periodic potentials [17], the parametric Resonance in the reheating process of universe [18], and wave equations in general relativity[14], etc. Mathieu function has been used in various areas in modern physics and mathematics.[1, 2, 3, 4, 5, 6, 7, 8, 9, 15, 16]

Unfortunately, even though Mathieu equation has been observed in various areas mentioned above, the Mathieu function could not be described in the form of a definite or contour integral of any elementary functions analytically. Besides the analytic solutions of its power series expansion in closed forms have not been constructed yet for almost 150 years because of its three different coefficients in the Frobenius solution. Three recursive relation in its power series creates the mathematical complex for the calculation. Instead, the analytic solutions for any linear ordinary differential equation having two term recurrence relation in its power series expansion can be derived. Mathieu function has only been described in numerical approximations (Whittaker 1914[13], Frenkel and Portugal 2001[10]). Sips 1949[12], Frenkel and Portugal 2001[10] argued that it is not possible to represent analytically the Mathieu function in a simple and handy way. All these authors treat solutions of Mathieu equation as the periodic-function with period 2π . It is not possible for arbitrary parameters λ and q in (7.1.1). They leave the fundamental solution of Mathieu equation by applying Floquet's theorem as solutions of recurrences because of three different coefficients in its power series expansion. In place of using the Bloch's theorem I will derive the analytic solution of Mathieu equation for any arbitrary λ and q by applying R3TRF in this chapter.

In Ref.[21] I construct analytic solutions of Mathieu equation about the regular singular point at $x = 0$ by applying 3TRF[20]; for the power series expansion in closed forms and its integral representation for infinite series including all higher terms of A_n 's.

In this chapter, by applying R3TRF in chapter 1, I construct the power series expansion in closed forms of Mathieu equation about the regular singular point at $x = 0$ (for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's) analytically. The integral representation of Mathieu equation and the generating function for Mathieu polynomial which makes A_n term terminated are derived in a mathematical rigor. Also Frobenius solution of the Mathieu equation about regular singular point at $x = 1$ and irregular singular one at $x = \infty$ by applying R3TRF is constructed including its integral forms and the generating function of it.

Mathieu equation is

$$\frac{d^2 y}{dz^2} + (\lambda - 2q \cos 2z) y = 0 \quad (7.1.1)$$

where λ and q are parameters. This is an equation with periodic-function coefficient.

Mathieu equation also can be described in algebraic forms putting $x = \cos^2 z$:

$$4x(1-x)\frac{d^2 y}{dx^2} + 2(1-2x)\frac{dy}{dx} + (\lambda + 2q - 4qx)y = 0 \quad (7.1.2)$$

This equation has two regular singularities: $x = 0$ and $x = 1$; the other singularity $x = \infty$ is irregular. Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\nu} \quad (7.1.3)$$

where ν is an indicial root. Plug (7.1.3) into (7.1.2).

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (7.1.4)$$

where,

$$A_n = \frac{4(n+\nu)^2 - (\lambda + 2q)}{2(n+1+\nu)(2(n+\nu)+1)} = \frac{(n+\nu-\varphi)(n+\nu+\varphi)}{(n+1+\nu)(n+\nu+\frac{1}{2})} \quad (7.1.5a)$$

and

$$\varphi = \frac{\sqrt{\lambda + 2q}}{2}$$

$$B_n = \frac{q}{(n+1+\nu)(n+\nu+\frac{1}{2})} \quad (7.1.5b)$$

$$c_1 = A_0 c_0 \quad (7.1.5c)$$

We have two indicial roots which are $\nu = 0$ and $\frac{1}{2}$. As we see (7.1.5b), there are no way to build the analytic solution of Mathieu equation for polynomial which makes B_n term terminated at certain value of n . Because the numerator of (7.1.5b) is just consist of constant q parameter.³

Now let's test for convergence of the analytic function $y(x)$. As $n \gg 1$, (7.1.5a)–(7.1.5c) are

$$\lim_{n \gg 1} A_n = 1 \quad (7.1.6a)$$

$$\lim_{n \gg 1} B_n = \frac{q}{n^2} \quad (7.1.6b)$$

³Whenever index n increase in (7.1.5b), B_n term never be terminated with a fixed constant parameter q .

As $n \gg 1$, (7.1.6b) is negligible. Put (7.1.6a) with $B_n = 0$ into (7.1.4).

$$\begin{aligned}
 c_0 & \\
 c_1 &= c_0 \\
 c_2 &= c_0 \\
 c_3 &= c_0 \\
 &\vdots \quad \vdots
 \end{aligned} \tag{7.1.7}$$

When a function $y(x)$, analytic at $x = 0$, is expanded in a power series putting $c_0 = 1$ by using (7.1.7), we write

$$\lim_{n \gg 1} y(x) \approx \sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \quad \text{where } 0 \leq x = \cos^2 z < 1 \tag{7.1.8}$$

For being convergent of $y(x)$ in (7.1.8), an independent variable $x = \cos^2 z$ should be less than 1.

The function $y(x)$ should be convergent as $x = 1$ in terms of the application of the mathematical physics. As A_n term is terminated as certain value of index n , the function $y(x)$ will be convergent even if $x = 1$. (7.1.8) is not applicable for the asymptotic behavior of $y(x)$ any more from the quantum mechanical point of view. Put (7.1.6b) into (7.1.4) with $A_n = 0$.⁴ For $n = 0, 1, 2, \dots$, it give

$$\begin{aligned}
 c_2 &= \frac{q}{1^2} c_0 & c_3 &= \frac{q}{2^2} c_1 \\
 c_4 &= \frac{q}{1^2 \cdot 3^2} c_0 & c_5 &= \frac{q}{2^2 \cdot 4^2} c_1 \\
 c_6 &= \frac{q}{1^2 \cdot 3^2 \cdot 5^2} c_0 & c_7 &= \frac{q}{2^2 \cdot 4^2 \cdot 6^2} c_1 \\
 &\vdots & &\vdots \\
 c_{2n} &= \left(\frac{\Gamma(\frac{1}{2})}{\Gamma(n+\frac{1}{2})} \right)^2 \left(\frac{q}{4} \right)^n c_0 & c_{2n+1} &= \frac{1}{(n!)^2} \left(\frac{q}{4} \right)^n c_1
 \end{aligned} \tag{7.1.9}$$

Put (7.1.9) into the power series expansion where $\sum_{n=0}^{\infty} c_n x^n$ putting $c_0 = c_1 = 1$ for simplicity.

$$\lim_{n \gg 1} y(x) > {}_1F_2 \left(1; \frac{1}{2}, \frac{1}{2}; \frac{q}{4} x^2 \right) + x I_0(\sqrt{q}x) \tag{7.1.10}$$

⁴Actually A_n can not be zero for $n \gg 1$. However, I only interesting in the minimum value of $y(x)$ for asymptotic behavior.

On the above ${}_1F_2(a; b, c; z)$ is a generalized hypergeometric function which is given by

$${}_1F_2(a; b, c; z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n (c)_n} \frac{z^n}{n!}$$

And $I_m(z)$ is a modified Bessel function which is defined by

$$I_m(z) = \sum_{n=0}^{\infty} \frac{1}{(n+m)! n!} \left(\frac{z}{2}\right)^{2n+m}$$

7.2 Mathieu equation about regular singular point at zero

7.2.1 Power series

Polynomial of type 2

There are two types of power series expansions using the two term recurrence relation in a linear ordinary differential equation which are a polynomial and an infinite series. In contrast there are three types of polynomials and an infinite series in three term recurrence relation of it: (1) polynomial which makes B_n term terminated: A_n term is not terminated, (2) polynomial which makes A_n term terminated: B_n term is not terminated, (3) polynomial which makes A_n and B_n terms terminated at the same time.⁵ In three term recurrence relation, polynomial of type 3 I categorize as complete polynomial. The solutions for the type 1 and 3 polynomials of Mathieu equation in the algebraic form does not exist because of the numerator in B_n term. There are only two type solutions of Mathieu equation which are type 2 polynomial and infinite series. For type 2 Mathieu polynomial about $x = 0$, I treat a parameter q as a free variable and λ as a fixed value.

In chapter 1 the general expression of power series of $y(x)$ for polynomial of type 2 is

⁵If A_n and B_n terms are not terminated, it turns to be infinite series.

defined by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\alpha_0} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\alpha_k} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\alpha_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (7.2.1)
\end{aligned}$$

In the above, $\alpha_i \leq \alpha_j$ only if $i \leq j$ where $i, j, \alpha_i, \alpha_j \in \mathbb{N}_0$.

For a polynomial, we need a condition which is:

$$A_{\alpha_i+2i} = 0 \quad \text{where } i, \alpha_i = 0, 1, 2, \dots \quad (7.2.2)$$

In the above, α_i is an eigenvalue that makes A_n term terminated at certain value of index n . (7.2.2) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (7.2.1).

The case of $\varphi = \frac{1}{2}\sqrt{\lambda+2q} = \lambda_i + 2i + \nu$ In (7.1.5a)-(7.1.5c) replace $\varphi = \frac{1}{2}\sqrt{\lambda+2q}$ by $\lambda_i + 2i + \nu$. In (7.2.2) replace index α_i by λ_i . Take the new (7.1.5a)-(7.1.5c), (7.2.2) and put them in (7.2.1). After the replacement process, the general expression of power series

of Mathieu equation for polynomial of type 2 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\nu \left\{ \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} x^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0 + 2 + \nu) (i_0 + \frac{3}{2} + \nu)} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1} (\lambda_1 + 4 + 2\nu)_{i_1} (3 + \nu)_{i_1} (\frac{5}{2} + \nu)_{i_1}}{(-\lambda_1)_{i_0} (\lambda_1 + 4 + 2\nu)_{i_0} (3 + \nu)_{i_1} (\frac{5}{2} + \nu)_{i_1}} x^{i_1} \Big\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0 + 2 + \nu) (i_0 + \frac{3}{2} + \nu)} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\lambda_k} \frac{1}{(i_k + 2k + 2 + \nu) (i_k + 2k + \frac{3}{2} + \nu)} \right. \\
&\quad \times \frac{(-\lambda_k)_{i_k} (\lambda_k + 4k + 2\nu)_{i_k} (2k + 1 + \nu)_{i_{k-1}} (2k + \frac{1}{2} + \nu)_{i_{k-1}}}{(-\lambda_k)_{i_{k-1}} (\lambda_k + 4k + 2\nu)_{i_{k-1}} (2k + 1 + \nu)_{i_k} (2k + \frac{1}{2} + \nu)_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\lambda_n} \frac{(-\lambda_n)_{i_n} (\lambda_n + 4n + 2\nu)_{i_n} (2n + 1 + \nu)_{i_{n-1}} (2n + \frac{1}{2} + \nu)_{i_{n-1}}}{(-\lambda_n)_{i_{n-1}} (\lambda_n + 4n + 2\nu)_{i_{n-1}} (2n + 1 + \nu)_{i_n} (2n + \frac{1}{2} + \nu)_{i_n}} x^{i_n} \Big\} \eta^n \Big\} \quad (7.2.3)
\end{aligned}$$

where

$$\begin{cases} x = \cos^2 z \\ \eta = qx^2 \\ \lambda = 2^2 (\lambda_j + 2j + \nu)^2 - 2q \text{ as } j, \lambda_j \in \mathbb{N}_0 \\ \lambda_i \leq \lambda_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

The case of $\varphi = \frac{1}{2}\sqrt{\lambda + 2q} = -(\lambda_i + 2i + \nu)$ In (7.1.5a)-(7.1.5c) replace $\frac{1}{2}\sqrt{\lambda + 2q}$ by $-(\lambda_i + 2i + \nu)$. In (7.2.2) replace index α_i by λ_i . Take the new (7.1.5a)-(7.1.5c), (7.2.2) and put them in (7.2.1). After the replacement process, its solution is equivalent to (7.2.3).

Put $c_0 = 1$ as $\nu = 0$ for the first kind of independent solutions of Mathieu equation and $\nu = 1/2$ for the second one in (7.2.3).

Remark 7.2.1 The power series expansion of Mathieu equation of the first kind for polynomial of type 2 about $x = 0$ as $\lambda = 2^2(\lambda_j + 2j)^2 - 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= MF_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j)^2 - 2q; \eta = qx^2, x = \cos^2 z) \\
&= \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} x^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2)(i_0+\frac{3}{2})} \frac{(-\lambda_0)_{i_0} (\lambda_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1} (\lambda_1+4)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-\lambda_1)_{i_0} (\lambda_1+4)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} x^{i_1} \right\} \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2)(i_0+\frac{3}{2})} \frac{(-\lambda_0)_{i_0} (\lambda_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\lambda_k} \frac{1}{(i_k+2k+2)(i_k+2k+\frac{3}{2})} \frac{(-\lambda_k)_{i_k} (\lambda_k+4k)_{i_k} (2k+1)_{i_{k-1}} (2k+\frac{1}{2})_{i_{k-1}}}{(-\lambda_k)_{i_{k-1}} (\lambda_k+4k)_{i_{k-1}} (2k+1)_{i_k} (2k+\frac{1}{2})_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\lambda_n} \frac{(-\lambda_n)_{i_n} (\lambda_n+4n)_{i_n} (2n+1)_{i_{n-1}} (2n+\frac{1}{2})_{i_{n-1}}}{(-\lambda_n)_{i_{n-1}} (\lambda_n+4n)_{i_{n-1}} (2n+1)_{i_n} (2n+\frac{1}{2})_{i_n}} x^{i_n} \right\} \eta^n \quad (7.2.4)
\end{aligned}$$

Remark 7.2.2 The power series expansion of Mathieu equation of the second kind for polynomial of type 2 about $x = 0$ as $\lambda = 2^2(\lambda_j + 2j + 1/2)^2 - 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= MS_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j + 1/2)^2 - 2q; \eta = qx^2, x = \cos^2 z) \\
&= x^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+1)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} x^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+\frac{5}{2})(i_0+2)} \frac{(-\lambda_0)_{i_0} (\lambda_0+1)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1} (\lambda_1+5)_{i_1} (\frac{7}{2})_{i_0} (3)_{i_0}}{(-\lambda_1)_{i_0} (\lambda_1+5)_{i_0} (\frac{7}{2})_{i_1} (3)_{i_1}} x^{i_1} \right\} \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+\frac{5}{2})(i_0+2)} \frac{(-\lambda_0)_{i_0} (\lambda_0+1)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\lambda_k} \frac{1}{(i_k+2k+\frac{5}{2})(i_k+2k+2)} \frac{(-\lambda_k)_{i_k} (\lambda_k+4k+1)_{i_k} (2k+\frac{3}{2})_{i_{k-1}} (2k+1)_{i_{k-1}}}{(-\lambda_k)_{i_{k-1}} (\lambda_k+4k+1)_{i_{k-1}} (2k+\frac{3}{2})_{i_k} (2k+1)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\lambda_n} \frac{(-\lambda_n)_{i_n} (\lambda_n+4n+1)_{i_n} (2n+\frac{3}{2})_{i_{n-1}} (2n+1)_{i_{n-1}}}{(-\lambda_n)_{i_{n-1}} (\lambda_n+4n+1)_{i_{n-1}} (2n+\frac{3}{2})_{i_n} (2n+1)_{i_n}} x^{i_n} \right\} \eta^n \quad (7.2.5)
\end{aligned}$$

Infinite series

In chapter 1 the general expression of power series of $y(x)$ for infinite series is

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
&\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Big\} \quad (7.2.6)
\end{aligned}$$

Substitute (7.1.5a)–(7.1.5c) into (7.2.6). The general expression of power series of Mathieu equation for infinite series about $x = 0$ is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^{\nu} \left\{ \sum_{i_0=0}^{\infty} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} \left(\frac{1}{2} + \nu\right)_{i_0}} x^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0 + 2 + \nu) \left(i_0 + \frac{3}{2} + \nu\right)} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} \left(\frac{1}{2} + \nu\right)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\nu + 2 - \varphi)_{i_1} (\nu + 2 + \varphi)_{i_1} (3 + \nu)_{i_0} \left(\frac{5}{2} + \nu\right)_{i_0}}{(\nu + 2 - \varphi)_{i_0} (\nu + 2 + \varphi)_{i_0} (3 + \nu)_{i_1} \left(\frac{5}{2} + \nu\right)_{i_1}} x^{i_1} \right\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0 + 2 + \nu) \left(i_0 + \frac{3}{2} + \nu\right)} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} \left(\frac{1}{2} + \nu\right)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{1}{(i_k + 2k + 2 + \nu) \left(i_k + 2k + \frac{3}{2} + \nu\right)} \right. \\
&\quad \times \left. \frac{(\nu + 2k - \varphi)_{i_k} (\nu + 2k + \varphi)_{i_k} (2k + 1 + \nu)_{i_{k-1}} \left(2k + \frac{1}{2} + \nu\right)_{i_{k-1}}}{(\nu + 2k - \varphi)_{i_{k-1}} (\nu + 2k + \varphi)_{i_{k-1}} (2k + 1 + \nu)_{i_k} \left(2k + \frac{1}{2} + \nu\right)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\nu + 2n - \varphi)_{i_n} (\nu + 2n + \varphi)_{i_n} (2n + 1 + \nu)_{i_{n-1}} \left(2n + \frac{1}{2} + \nu\right)_{i_{n-1}}}{(\nu + 2n - \varphi)_{i_{n-1}} (\nu + 2n + \varphi)_{i_{n-1}} (2n + 1 + \nu)_{i_n} \left(2n + \frac{1}{2} + \nu\right)_{i_n}} x^{i_n} \right\} \eta^n \Big\} \quad (7.2.7)
\end{aligned}$$

where

$$\begin{cases} x = \cos^2 z \\ \eta = qx^2 \\ \varphi = \frac{\sqrt{\lambda+2q}}{2} \end{cases}$$

Put $c_0 = 1$ as $\nu = 0$ for the first kind of independent solutions of Mathieu equation and $\nu = 1/2$ for the second one in (7.2.7).

Remark 7.2.3 *The power series expansion of Mathieu equation of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned} y(x) &= MF^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda + 2q}; \eta = qx^2, x = \cos^2 z \right) \\ &= \sum_{i_0=0}^{\infty} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} x^{i_0} \\ &+ \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2) \left(i_0 + \frac{3}{2}\right)} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(2-\varphi)_{i_1} (2+\varphi)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{(2-\varphi)_{i_0} (2+\varphi)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} x^{i_1} \right\} \eta \\ &+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2) \left(i_0 + \frac{3}{2}\right)} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \right. \\ &\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{1}{(i_k+2k+2) \left(i_k + 2k + \frac{3}{2}\right)} \frac{(2k-\varphi)_{i_k} (2k+\varphi)_{i_k} (2k+1)_{i_{k-1}} \left(2k + \frac{1}{2}\right)_{i_{k-1}}}{(2k-\varphi)_{i_{k-1}} (2k+\varphi)_{i_{k-1}} (2k+1)_{i_k} \left(2k + \frac{1}{2}\right)_{i_k}} \right\} \\ &\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(2n-\varphi)_{i_n} (2n+\varphi)_{i_n} (2n+1)_{i_{n-1}} \left(2n + \frac{1}{2}\right)_{i_{n-1}}}{(2n-\varphi)_{i_{n-1}} (2n+\varphi)_{i_{n-1}} (2n+1)_{i_n} \left(2n + \frac{1}{2}\right)_{i_n}} x^{i_n} \right\} \eta^n \end{aligned} \quad (7.2.8)$$

Remark 7.2.4 *The power series expansion of Mathieu equation of the second kind for*

infinite series about $x = 0$ using R3TRF is

$$\begin{aligned}
y(x) &= MS^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda + 2q}; \eta = qx^2, x = \cos^2 z \right) \\
&= x^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{\left(\frac{1}{2} - \varphi\right)_{i_0} \left(\frac{1}{2} + \varphi\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} x^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{1}{\left(i_0 + \frac{5}{2}\right) (i_0 + 2)} \frac{\left(\frac{1}{2} - \varphi\right)_{i_0} \left(\frac{1}{2} + \varphi\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{\left(\frac{5}{2} - \varphi\right)_{i_1} \left(\frac{5}{2} + \varphi\right)_{i_1} \left(\frac{7}{2}\right)_{i_0} (3)_{i_0}}{\left(\frac{5}{2} - \varphi\right)_{i_0} \left(\frac{5}{2} + \varphi\right)_{i_0} \left(\frac{7}{2}\right)_{i_1} (3)_{i_1}} x^{i_1} \right\} \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{1}{\left(i_0 + \frac{5}{2}\right) (i_0 + 2)} \frac{\left(\frac{1}{2} - \varphi\right)_{i_0} \left(\frac{1}{2} + \varphi\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{1}{\left(i_k + 2k + \frac{5}{2}\right) (i_k + 2k + 2)} \frac{\left(2k + \frac{1}{2} - \varphi\right)_{i_k} \left(2k + \frac{1}{2} + \varphi\right)_{i_k} \left(2k + \frac{3}{2}\right)_{i_{k-1}} (2k + 1)_{i_{k-1}}}{\left(2k + \frac{1}{2} - \varphi\right)_{i_{k-1}} \left(2k + \frac{1}{2} + \varphi\right)_{i_{k-1}} \left(2k + \frac{3}{2}\right)_{i_k} (2k + 1)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{\left(2n + \frac{1}{2} - \varphi\right)_{i_n} \left(2n + \frac{1}{2} + \varphi\right)_{i_n} \left(2n + \frac{3}{2}\right)_{i_{n-1}} (2n + 1)_{i_{n-1}}}{\left(2n + \frac{1}{2} - \varphi\right)_{i_{n-1}} \left(2n + \frac{1}{2} + \varphi\right)_{i_{n-1}} \left(2n + \frac{3}{2}\right)_{i_n} (2n + 1)_{i_n}} x^{i_n} \right\} \eta^n \Bigg\} \quad (7.2.9)
\end{aligned}$$

Two power series expansions of Mathieu equation for infinite series in this chapter and Ref.[21] are equivalent to each other. In this chapter, B_n is the leading term in sequence c_n of the analytic function $y(x)$. In Ref.[21], A_n is the leading term in sequence c_n of the analytic function $y(x)$.

For the special case, as $|q| \ll 1$ in (7.2.8) and (7.2.9), we have

$$\begin{aligned}
&\lim_{|q| \ll 1} MF^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda + 2q}; \eta = qx^2, x = \cos^2 z \right) \\
&\approx \sum_{i_0=0}^{\infty} \frac{\left(-\frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{\sqrt{\lambda}}{2}\right)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} x^{i_0} \\
&+ \eta \sum_{i_0=0}^{\infty} \frac{1}{\left(i_0 + 2\right) \left(i_0 + \frac{3}{2}\right)} \frac{\left(-\frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{\sqrt{\lambda}}{2}\right)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{\left(2 - \frac{\sqrt{\lambda}}{2}\right)_{i_1} \left(2 + \frac{\sqrt{\lambda}}{2}\right)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{\left(2 - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(2 + \frac{\sqrt{\lambda}}{2}\right)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} x^{i_1} \\
&> \cos \left(\sqrt{\lambda} \sin^{-1} (\sqrt{x}) \right) \quad (7.2.10a)
\end{aligned}$$

And,

$$\begin{aligned}
& \lim_{|q| \ll 1} MS^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda + 2q}; \eta = qx^2, x = \cos^2 z \right) \\
& \approx x^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{\left(\frac{1}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{1}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} x^{i_0} \right. \\
& \quad \left. + \eta \sum_{i_0=0}^{\infty} \frac{1}{\left(i_0 + \frac{5}{2}\right) (i_0 + 2)} \frac{\left(\frac{1}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{1}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{\left(\frac{5}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_1} \left(\frac{5}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_1} \left(\frac{7}{2}\right)_{i_0} (3)_{i_0}}{\left(\frac{5}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{5}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{7}{2}\right)_{i_1} (3)_{i_1}} x^{i_1} \right\} \\
& > \sqrt{\frac{x}{1-x}} \cos \left(\sqrt{\lambda} \sin^{-1} (\sqrt{x}) \right) \tag{7.2.10b}
\end{aligned}$$

7.2.2 Integral formalism

The Mathieu function could not be constructed in a definite or contour integral form of any well-known simple functions because of three different coefficients in its power series expansion. The three term recurrence relation in power series of Mathieu equation creates mathematical difficulty to be analyzed it into a direct or contour integrals.

In place of describing the Mathieu function into the integral representation of any simple functions which have two term recursion in its power series expansion of a linear ordinary differential equation, in earlier literature the integral equations of the Mathieu function were constructed by using trigonometric functions kernels, Bessel-function kernels and etc; such integral relationships express one analytic solution in terms of another analytic solution such as the Fourier series of the periodic Mathieu functions; one of ce, se, me, Ce and Se, Me, Ge and Ge functions.[22, 23, 24, 25] There are many other forms of integral relations in Mathieu equation. [26, 27, 28, 29]

In Ref.[21], I show integral representation (each sub-integral $y_m(x)$ where $m = 0, 1, 2, \dots$ is composed of $2m$ terms of definite integrals and m terms of contour integrals) of the Mathieu function using 3TRF: a Modified Bessel function recurs in each of sub-integral forms of the Mathieu function.

Now I consider integral forms of the Mathieu function by using R3TRF.

Polynomial of type 2

Now I consider an integral form of the Mathieu polynomial of type 2 by using R3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned} I_l &= \sum_{i_l=i_{l-1}}^{\lambda_l} \frac{(-\lambda_l)_{i_l} (\lambda_l + 4l + 2\nu)_{i_l} (2l + 1 + \nu)_{i_{l-1}} (2l + \frac{1}{2} + \nu)_{i_{l-1}}}{(-\lambda_l)_{i_{l-1}} (\lambda_l + 4l + 2\nu)_{i_{l-1}} (2l + 1 + \nu)_{i_l} (2l + \frac{1}{2} + \nu)_{i_l}} x^{i_l} \\ &= x^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1} + 2l - \frac{1}{2} + \nu, j + 1) B(i_{l-1} + 2l + \nu, j + 1) (i_{l-1} - \lambda_l)_j (i_{l-1} + \lambda_l + 4l + 2\nu)_j}{(i_{l-1} + 2l + \nu)^{-1} (i_{l-1} + 2l - \frac{1}{2} + \nu)^{-1} (1)_j j!} x^j \end{aligned} \quad (7.2.11)$$

By using integral form of beta function,

$$B(i_{l-1} + 2l + \nu, j + 1) = \int_0^1 dt_l t_l^{i_{l-1} + 2l - 1 + \nu} (1 - t_l)^j \quad (7.2.12a)$$

$$B\left(i_{l-1} + 2l - \frac{1}{2} + \nu, j + 1\right) = \int_0^1 du_l u_l^{i_{l-1} + 2l - \frac{3}{2} + \nu} (1 - u_l)^j \quad (7.2.12b)$$

Substitute (7.2.12a) and (7.2.12b) into (7.2.11). And divide $(i_{l-1} + 2l + \nu) (i_{l-1} + 2l - \frac{1}{2} + \nu)$ into the new (7.2.11).

$$\begin{aligned} K_l &= \frac{1}{(i_{l-1} + 2l + \nu) (i_{l-1} + 2l - \frac{1}{2} + \nu)} \\ &\times \sum_{i_l=i_{l-1}}^{\lambda_l} \frac{(-\lambda_l)_{i_l} (\lambda_l + 4l + 2\nu)_{i_l} (2l + 1 + \nu)_{i_{l-1}} (2l + \frac{1}{2} + \nu)_{i_{l-1}}}{(-\lambda_l)_{i_{l-1}} (\lambda_l + 4l + 2\nu)_{i_{l-1}} (2l + 1 + \nu)_{i_l} (2l + \frac{1}{2} + \nu)_{i_l}} x^{i_l} \\ &= \int_0^1 dt_l t_l^{2l - 1 + \nu} \int_0^1 du_l u_l^{2l - \frac{3}{2} + \nu} (x t_l u_l)^{i_{l-1}} \\ &\times \sum_{j=0}^{\infty} \frac{(i_{l-1} - \lambda_l)_j (i_{l-1} + \lambda_l + 4l + 2\nu)_j}{(1)_j j!} [x(1 - t_l)(1 - u_l)]^j \end{aligned}$$

The contour integral form of Gauss hypergeometric function is given by

$$\begin{aligned} {}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n (n!)} z^n \\ &= -\frac{1}{2\pi i} \frac{\Gamma(1 - \alpha) \Gamma(\gamma)}{\Gamma(\gamma - \alpha)} \oint dp_l (-p_l)^{\alpha-1} (1 - p_l)^{\gamma-\alpha-1} (1 - zp_l)^{-\beta} \quad (7.2.13) \\ &\text{where } \operatorname{Re}(\gamma - \alpha) > 0 \end{aligned}$$

replaced α, β, γ and z by $i_{l-1} - \lambda_l, i_{l-1} + \lambda_l + 4l + 2\nu, 1$ and $x(1 - t_l)(1 - u_l)$ in (7.2.13)

$$\begin{aligned}
& \sum_{j=0}^{\infty} \frac{(i_{l-1} - \lambda_l)_j (i_{l-1} + \lambda_l + 4l + 2\nu)_j}{(1)_j j!} [x(1 - t_l)(1 - u_l)]^j \\
&= \frac{1}{2\pi i} \oint dp_l \frac{1}{p_l} (1 - x(1 - t_l)(1 - u_l)p_l)^{-(4l+2\nu)} \\
&\quad \times \left(\frac{p_l - 1}{p_l} \frac{1}{1 - x(1 - t_l)(1 - u_l)p_l} \right)^{\lambda_l} \left(\frac{p_l}{p_l - 1} \frac{1}{1 - x(1 - t_l)(1 - u_l)p_l} \right)^{i_{l-1}} \quad (7.2.14)
\end{aligned}$$

Substitute (7.2.14) into K_l .

$$\begin{aligned}
K_l &= \frac{1}{(i_{l-1} + 2l + \nu)(i_{l-1} + 2l - \frac{1}{2} + \nu)} \\
&\quad \times \sum_{i_l=i_{l-1}}^{\lambda_l} \frac{(-\lambda_l)_{i_l} (\lambda_l + 4l + 2\nu)_{i_l} (2l + 1 + \nu)_{i_{l-1}} (2l + \frac{1}{2} + \nu)_{i_{l-1}}}{(-\lambda_l)_{i_{l-1}} (\lambda_l + 4l + 2\nu)_{i_{l-1}} (2l + 1 + \nu)_{i_l} (2l + \frac{1}{2} + \nu)_{i_l}} x^{i_l} \\
&= \int_0^1 dt_l t_l^{2l-1+\nu} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\nu} \frac{1}{2\pi i} \oint dp_l \frac{1}{p_l} (1 - x(1 - t_l)(1 - u_l)p_l)^{-(4l+2\nu)} \\
&\quad \times \left(\frac{p_l - 1}{p_l} \frac{1}{1 - x(1 - t_l)(1 - u_l)p_l} \right)^{\lambda_l} \left(\frac{p_l}{p_l - 1} \frac{xt_l u_l}{1 - x(1 - t_l)(1 - u_l)p_l} \right)^{i_{l-1}} \quad (7.2.15)
\end{aligned}$$

Substitute (7.2.15) into (7.2.3) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(x)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(x)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(x)$, etc.⁶

Theorem 7.2.5 *The general representation in the form of integral of the Mathieu*

⁶ $y_1(x)$ means the sub-power series in (7.2.3) contains one term of $B'_n s$, $y_2(x)$ means the sub-power series in (7.2.3) contains two terms of $B'_n s$, $y_3(x)$ means the sub-power series in (7.2.3) contains three terms of $B'_n s$, etc.

polynomial of type 2 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\nu \left\{ \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\nu + \frac{1}{2})_{i_0}} x^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\nu} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}+\nu} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) p_{n-k})^{-(4(n-k)+2\nu)} \\
&\quad \times \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) p_{n-k}} \right)^{\lambda_{n-k}} \left. \right\} \\
&\quad \times \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\nu + \frac{1}{2})_{i_0}} w_{1,n}^{i_0} \left. \right\} \eta^n \Bigg\} \tag{7.2.16}
\end{aligned}$$

where

$$w_{i,j} = \begin{cases} \frac{p_i}{(p_i - 1)} \frac{w_{i+1,j} t_i u_i}{1 - w_{i+1,j} p_i (1 - t_i)(1 - u_i)} & \text{where } i \leq j \\ x & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof According to (7.2.3),

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\nu} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \tag{7.2.17}$$

In the above, the power series expansions of sub-summation $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of Mathieu equation using R3TRF about $x = 0$ are

$$y_0(x) = c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} x^{i_0} \tag{7.2.18a}$$

$$\begin{aligned}
y_1(x) &= c_0 x^\nu \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0 + 2 + \nu) (i_0 + \frac{3}{2} + \nu)} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1} (\lambda_1 + 4 + 2\nu)_{i_1} (3 + \nu)_{i_0} (\frac{5}{2} + \nu)_{i_0}}{(-\lambda_1)_{i_0} (\lambda_1 + 4 + 2\nu)_{i_0} (3 + \nu)_{i_1} (\frac{5}{2} + \nu)_{i_1}} x^{i_1} \left. \right\} \eta \tag{7.2.18b}
\end{aligned}$$

$$\begin{aligned}
y_2(x) = & c_0 x^\nu \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2+\nu)(i_0+\frac{3}{2}+\nu)} \frac{(-\lambda_0)_{i_0}(\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\lambda_1} \frac{1}{(i_1+4+\nu)(i_1+\frac{7}{2}+\nu)} \frac{(-\lambda_1)_{i_1}(\lambda_1+4+2\nu)_{i_1}(3+\nu)_{i_0}(\frac{5}{2}+\nu)_{i_0}}{(-\lambda_1)_{i_0}(\lambda_1+4+2\nu)_{i_0}(3+\nu)_{i_1}(\frac{5}{2}+\nu)_{i_1}} \\
& \times \left. \sum_{i_2=i_1}^{\lambda_2} \frac{(-\lambda_2)_{i_2}(\lambda_2+8+2\nu)_{i_2}(5+\nu)_{i_1}(\frac{9}{2}+\nu)_{i_1}}{(-\lambda_2)_{i_1}(\lambda_2+8+2\nu)_{i_1}(5+\nu)_{i_2}(\frac{9}{2}+\nu)_{i_2}} x^{i_2} \right\} \eta^2 \quad (7.2.18c)
\end{aligned}$$

$$\begin{aligned}
y_3(x) = & c_0 x^\nu \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2+\nu)(i_0+\frac{3}{2}+\nu)} \frac{(-\lambda_0)_{i_0}(\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\lambda_1} \frac{1}{(i_1+4+\nu)(i_1+\frac{7}{2}+\nu)} \frac{(-\lambda_1)_{i_1}(\lambda_1+4+2\nu)_{i_1}(3+\nu)_{i_0}(\frac{5}{2}+\nu)_{i_0}}{(-\lambda_1)_{i_0}(\lambda_1+4+2\nu)_{i_0}(3+\nu)_{i_1}(\frac{5}{2}+\nu)_{i_1}} \\
& \times \sum_{i_2=i_1}^{\lambda_2} \frac{1}{(i_2+6+\nu)(i_2+\frac{11}{2}+\nu)} \frac{(-\lambda_2)_{i_2}(\lambda_2+8+2\nu)_{i_2}(5+\nu)_{i_1}(\frac{9}{2}+\nu)_{i_1}}{(-\lambda_2)_{i_1}(\lambda_2+8+2\nu)_{i_1}(5+\nu)_{i_2}(\frac{9}{2}+\nu)_{i_2}} \\
& \times \left. \sum_{i_3=i_2}^{\lambda_3} \frac{(-\lambda_3)_{i_3}(\lambda_3+12+2\nu)_{i_3}(7+\nu)_{i_2}(\frac{13}{2}+\nu)_{i_2}}{(-\lambda_3)_{i_2}(\lambda_3+12+2\nu)_{i_2}(7+\nu)_{i_3}(\frac{13}{2}+\nu)_{i_3}} x^{i_3} \right\} \eta^3 \quad (7.2.18d)
\end{aligned}$$

Put $l = 1$ in (7.2.15). Take the new (7.2.15) into (7.2.18b).

$$\begin{aligned}
y_1(x) = & \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-x(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\
& \times \left(\frac{p_1-1}{p_1} \frac{1}{1-x(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \left\{ c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0}(\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} w_{1,1}^{i_0} \right\} \eta \quad (7.2.19)
\end{aligned}$$

where

$$w_{1,1} = \frac{p_1}{p_1-1} \frac{xt_1u_1}{1-x(1-t_1)(1-u_1)p_1}$$

Put $l = 2$ in (7.2.15). Take the new (7.2.15) into (7.2.18c).

$$\begin{aligned}
y_2(x) &= c_0 x^\nu \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} (1-x(1-t_2)(1-u_2)p_2)^{-(8+2\nu)} \\
&\quad \times \left(\frac{p_2-1}{p_2} \frac{1}{1-x(1-t_2)(1-u_2)p_2} \right)^{\lambda_2} \\
&\quad \times \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2+\nu)(i_0+\frac{3}{2}+\nu)} \frac{(-\lambda_0)_{i_0}(\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1}(\lambda_1+4+2\nu)_{i_1}(3+\nu)_{i_0}(\frac{5}{2}+\nu)_{i_0} w_{2,2}^{i_1}}{(-\lambda_1)_{i_0}(\lambda_1+4+2\nu)_{i_0}(3+\nu)_{i_1}(\frac{5}{2}+\nu)_{i_1}} \right\} \eta^2 \quad (7.2.20)
\end{aligned}$$

where

$$w_{2,2} = \frac{p_2}{p_2-1} \frac{xt_2u_2}{1-x(1-t_2)(1-u_2)p_2}$$

Put $l = 1$ and $\eta = w_{2,2}$ in (7.2.15). Take the new (7.2.15) into (7.2.20).

$$\begin{aligned}
y_2(x) &= \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} (1-x(1-t_2)(1-u_2)p_2)^{-(8+2\nu)} \\
&\quad \times \left(\frac{p_2-1}{p_2} \frac{1}{1-x(1-t_2)(1-u_2)p_2} \right)^{\lambda_2} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-w_{2,2}(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\
&\quad \times \left(\frac{p_1-1}{p_1} \frac{1}{1-w_{2,2}(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \left\{ c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0}(\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} w_{1,2}^{i_0} \right\} \eta^2 \quad (7.2.21)
\end{aligned}$$

where

$$w_{1,2} = \frac{p_1}{p_1-1} \frac{w_{2,2}t_1u_1}{1-w_{2,2}(1-t_1)(1-u_1)p_1}$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the

integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned}
y_3(x) = & \int_0^1 dt_3 t_3^{5+\nu} \int_0^1 du_3 u_3^{\frac{9}{2}+\nu} \frac{1}{2\pi i} \oint dp_3 \frac{1}{p_3} (1-x(1-t_3)(1-u_3)p_3)^{-(12+2\nu)} \\
& \times \left(\frac{p_3-1}{p_3} \frac{1}{1-x(1-t_3)(1-u_3)p_3} \right)^{\lambda_3} \\
& \times \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} (1-w_{3,3}(1-t_2)(1-u_2)p_2)^{-(8+2\nu)} \\
& \times \left(\frac{p_2-1}{p_2} \frac{1}{1-w_{3,3}(1-t_2)(1-u_2)p_2} \right)^{\lambda_2} \\
& \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-w_{2,3}(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\
& \times \left(\frac{p_1-1}{p_1} \frac{1}{1-w_{2,3}(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \left\{ c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2}+\nu)_{i_0}} w_{1,3}^{i_0} \right\} \eta^3 \quad (7.2.22)
\end{aligned}$$

where

$$\begin{cases} w_{3,3} = \frac{p_3}{p_3-1} \frac{xt_3u_3}{1-x(1-t_3)(1-u_3)p_3} \\ w_{2,3} = \frac{p_2}{p_2-1} \frac{w_{3,3}t_2u_2}{1-w_{3,3}(1-t_2)(1-u_2)p_2} \\ w_{1,3} = \frac{p_1}{p_1-1} \frac{w_{2,3}t_1u_1}{1-w_{2,3}(1-t_1)(1-u_1)p_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Substitute (7.2.18a), (7.2.19), (7.2.21), (7.2.22) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (7.2.16).

Put $c_0 = 1$ as $\nu = 0$ for the first kind of independent solutions of Mathieu equation and $\nu = 1/2$ for the second one in (7.2.16).

Remark 7.2.6 *The integral representation of Mathieu equation of the first kind for*

polynomial of type 2 about $x = 0$ as $\lambda = 2^2(\lambda_j + 2j)^2 - 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= MF_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j)^2 - 2q; \eta = qx^2, x = \cos^2 z) \\
&= {}_2F_1\left(-\lambda_0, \lambda_0; \frac{1}{2}; x\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-4(n-k)} \\
&\quad \times \left. \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k}} \right)^{\lambda_{n-k}} \right\} \\
&\quad \times {}_2F_1\left(-\lambda_0, \lambda_0; \frac{1}{2}; w_{1,n}\right) \Big\} \eta^n
\end{aligned} \tag{7.2.23}$$

Remark 7.2.7 The integral representation of Mathieu equation of the second kind for polynomial of type 2 about $x = 0$ as $\lambda = 2^2(\lambda_j + 2j + 1/2)^2 - 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(x) &= MS_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j + 1/2)^2 - 2q; \eta = qx^2, x = \cos^2 z) \\
&= x^{\frac{1}{2}} \left\{ {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; x\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-(4(n-k)+1)} \\
&\quad \times \left. \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k}} \right)^{\lambda_{n-k}} \right\} \\
&\quad \times {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; w_{1,n}\right) \Big\} \eta^n
\end{aligned} \tag{7.2.24}$$

Infinite series

Let's consider the integral representation of Mathieu equation about $x = 0$ for infinite series by applying R3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned}
M_l &= \sum_{i_l=i_{l-1}}^{\infty} \frac{(\nu + 2l - \varphi)_{i_l} (\nu + 2l + \varphi)_{i_l} (2l + 1 + \nu)_{i_{l-1}} (2l + \frac{1}{2} + \nu)_{i_{l-1}}}{(\nu + 2l - \varphi)_{i_{l-1}} (\nu + 2l + \varphi)_{i_{l-1}} (2l + 1 + \nu)_{i_l} (2l + \frac{1}{2} + \nu)_{i_l}} x^{i_l} \\
&= x^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1} + 2l - \frac{1}{2} + \nu, j + 1) B(i_{l-1} + 2l + \nu, j + 1) (\nu + 2l - \varphi + i_{l-1})_j (\nu + 2l + \varphi + i_{l-1})_j}{(i_{l-1} + 2l + \nu)^{-1} (i_{l-1} + 2l - \frac{1}{2} + \nu)^{-1} (1)_j j!} x^j
\end{aligned} \tag{7.2.25}$$

Substitute (7.2.12a) and (7.2.12b) into (7.2.25). And divide $(i_{l-1} + 2l + \nu) (i_{l-1} + 2l - \frac{1}{2} + \nu)$ into the new (7.2.25).

$$\begin{aligned}
 V_l &= \frac{1}{(i_{l-1} + 2l + \nu) (i_{l-1} + 2l - \frac{1}{2} + \nu)} \\
 &\times \sum_{i_l=i_{l-1}}^{\infty} \frac{(\nu + 2l - \varphi)_{i_l} (\nu + 2l + \varphi)_{i_l} (2l + 1 + \nu)_{i_{l-1}} (2l + \frac{1}{2} + \nu)_{i_{l-1}}}{(\nu + 2l - \varphi)_{i_{l-1}} (\nu + 2l + \varphi)_{i_{l-1}} (2l + 1 + \nu)_{i_l} (2l + \frac{1}{2} + \nu)_{i_l}} x^{i_l} \\
 &= \int_0^1 dt_l t_l^{2l-1+\nu} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\nu} (xt_l u_l)^{i_{l-1}} \\
 &\times \sum_{j=0}^{\infty} \frac{(\nu + 2l - \varphi + i_{l-1})_j (\nu + 2l + \varphi + i_{l-1})_j}{(1)_j j!} (x(1-t_l)(1-u_l))^j
 \end{aligned}$$

The hypergeometric function is defined by

$$\begin{aligned}
 {}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n n!} z^n \\
 &= \frac{1}{2\pi i} \frac{\Gamma(1+\alpha-\gamma)}{\Gamma(\alpha)} \int_0^{(1+)} dv_l (-1)^\gamma (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \quad (7.2.26) \\
 &\text{where } \gamma - \alpha \neq 1, 2, 3, \dots, \operatorname{Re}(\alpha) > 0
 \end{aligned}$$

Replace α, β, γ and z by $\nu + 2l - \varphi + i_{l-1}, \nu + 2l + \varphi + i_{l-1}, 1$ and $x(1-t_l)(1-u_l)$ in (7.2.26). Take the new (7.2.26) into V_l .

$$\begin{aligned}
 V_l &= \frac{1}{(i_{l-1} + 2l + \nu) (i_{l-1} + 2l - \frac{1}{2} + \nu)} \\
 &\times \sum_{i_l=i_{l-1}}^{\infty} \frac{(\nu + 2l - \varphi)_{i_l} (\nu + 2l + \varphi)_{i_l} (2l + 1 + \nu)_{i_{l-1}} (2l + \frac{1}{2} + \nu)_{i_{l-1}}}{(\nu + 2l - \varphi)_{i_{l-1}} (\nu + 2l + \varphi)_{i_{l-1}} (2l + 1 + \nu)_{i_l} (2l + \frac{1}{2} + \nu)_{i_l}} x^{i_l} \\
 &= \int_0^1 dt_l t_l^{2l-1+\nu} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\nu} \frac{1}{2\pi i} \oint dp_l \frac{1}{p_l} \left(\frac{p_l - 1}{p_l} \right)^{-(\nu+2l)+\varphi} \\
 &\times (1 - x(1-t_l)(1-u_l)p_l)^{-(\nu+2l)-\varphi} \left(\frac{p_l}{p_l - 1} \frac{xt_l u_l}{1 - x(1-t_l)(1-u_l)p_l} \right)^{i_{l-1}} \quad (7.2.27)
 \end{aligned}$$

Substitute (7.2.27) into (7.2.7) where $l = 1, 2, 3, \dots$; apply V_1 into the second summation of sub-power series $y_1(x)$, apply V_2 into the third summation and V_1 into the second summation of sub-power series $y_2(x)$, apply V_3 into the forth summation, V_2 into the third summation and V_1 into the second summation of sub-power series $y_3(x)$, etc.⁷

⁷ $y_1(x)$ means the sub-power series in (7.2.7) contains one term of $B'_n s$, $y_2(x)$ means the sub-power series in (7.2.7) contains two terms of $B'_n s$, $y_3(x)$ means the sub-power series in (7.2.7) contains three terms of $B'_n s$, etc.

Theorem 7.2.8 *The general expression of an integral form of Mathieu equation for infinite series about $x = 0$ using R3TRF is given by*

$$\begin{aligned}
y(x) = & c_0 x^\nu \left\{ \sum_{i_0=0}^{\infty} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} (\nu + \frac{1}{2})_{i_0}} x^{i_0} \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\nu} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}+\nu} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} \left(\frac{p_{n-k} - 1}{p_{n-k}} \right)^{-(\nu+2(n-k))+\varphi} \\
& \times (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-(\nu+2(n-k))-\varphi} \left. \right\} \\
& \times \sum_{i_0=0}^{\infty} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} (\nu + \frac{1}{2})_{i_0}} w_{1,n}^{i_0} \left. \right\} \eta^n \Bigg\} \quad (7.2.28)
\end{aligned}$$

where

$$\varphi = \frac{1}{2} \sqrt{\lambda + 2q}$$

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof In (7.2.7) sub-power series $y_0(x)$, $y_1(x)$, $y_2(x)$ and $y_3(x)$ of Mathieu equation for infinite series about $x = 0$ using R3TRF are given by

$$y_0(x) = c_0 x^\nu \sum_{i_0=0}^{\infty} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} x^{i_0} \quad (7.2.29a)$$

$$\begin{aligned}
y_1(x) = & c_0 x^\nu \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0 + 2 + \nu) (i_0 + \frac{3}{2} + \nu)} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{(\nu + 2 - \varphi)_{i_1} (\nu + 2 + \varphi)_{i_1} (3 + \nu)_{i_0} (\frac{5}{2} + \nu)_{i_0}}{(\nu + 2 - \varphi)_{i_0} (\nu + 2 + \varphi)_{i_0} (3 + \nu)_{i_1} (\frac{5}{2} + \nu)_{i_1}} x^{i_1} \left. \right\} \eta \quad (7.2.29b)
\end{aligned}$$

$$\begin{aligned}
y_2(x) = & c_0 x^\nu \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2+\nu)(i_0+\frac{3}{2}+\nu)} \frac{(\nu-\varphi)_{i_0}(\nu+\varphi)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{1}{(i_1+4+\nu)(i_1+\frac{7}{2}+\nu)} \frac{(\nu+2-\varphi)_{i_1}(\nu+2+\varphi)_{i_1}(3+\nu)_{i_0}(\frac{5}{2}+\nu)_{i_0}}{(\nu+2-\varphi)_{i_0}(\nu+2+\varphi)_{i_0}(3+\nu)_{i_1}(\frac{5}{2}+\nu)_{i_1}} \\
& \times \left. \sum_{i_2=i_1}^{\infty} \frac{(\nu+4-\varphi)_{i_2}(\nu+4+\varphi)_{i_2}(5+\nu)_{i_1}(\frac{9}{2}+\nu)_{i_1}}{(\nu+4-\varphi)_{i_1}(\nu+4+\varphi)_{i_1}(5+\nu)_{i_2}(\frac{9}{2}+\nu)_{i_2}} x^{i_2} \right\} \eta^2 \quad (7.2.29c)
\end{aligned}$$

$$\begin{aligned}
y_3(x) = & c_0 x^\nu \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2+\nu)(i_0+\frac{3}{2}+\nu)} \frac{(\nu-\varphi)_{i_0}(\nu+\varphi)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{\infty} \frac{1}{(i_1+4+\nu)(i_1+\frac{7}{2}+\nu)} \frac{(\nu+2-\varphi)_{i_1}(\nu+2+\varphi)_{i_1}(3+\nu)_{i_0}(\frac{5}{2}+\nu)_{i_0}}{(\nu+2-\varphi)_{i_0}(\nu+2+\varphi)_{i_0}(3+\nu)_{i_1}(\frac{5}{2}+\nu)_{i_1}} \\
& \times \sum_{i_2=i_1}^{\infty} \frac{1}{(i_2+6+\nu)(i_2+\frac{11}{2}+\nu)} \frac{(\nu+4-\varphi)_{i_2}(\nu+4+\varphi)_{i_2}(5+\nu)_{i_1}(\frac{9}{2}+\nu)_{i_1}}{(\nu+4-\varphi)_{i_1}(\nu+4+\varphi)_{i_1}(5+\nu)_{i_2}(\frac{9}{2}+\nu)_{i_2}} \\
& \times \left. \sum_{i_3=i_2}^{\infty} \frac{(\nu+6-\varphi)_{i_3}(\nu+6+\varphi)_{i_3}(7+\nu)_{i_2}(\frac{13}{2}+\nu)_{i_2}}{(\nu+6-\varphi)_{i_2}(\nu+6+\varphi)_{i_2}(7+\nu)_{i_3}(\frac{13}{2}+\nu)_{i_3}} x^{i_3} \right\} \eta^3 \quad (7.2.29d)
\end{aligned}$$

Put $l = 1$ in (7.2.27). Take the new (7.2.27) into (7.2.29b).

$$\begin{aligned}
y_1(x) = & \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} \left(\frac{p_1-1}{p_1} \right)^{-(\nu+2)+\varphi} \\
& \times (1-x(1-t_1)(1-u_1)p_1)^{-(\nu+2)-\varphi} \left\{ c_0 x^\nu \sum_{i_0=0}^{\infty} \frac{(\nu-\varphi)_{i_0}(\nu+\varphi)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} w_{1,1}^{i_0} \right\} \eta \quad (7.2.30)
\end{aligned}$$

where

$$w_{1,1} = \frac{p_1}{p_1-1} \frac{xt_1u_1}{1-x(1-t_1)(1-u_1)p_1}$$

Put $l = 2$ in (7.2.27). Take the new (7.2.27) into (7.2.29c).

$$\begin{aligned}
y_2(x) = & c_0 x^\nu \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} \left(\frac{p_2-1}{p_2} \right)^{-(\nu+4)+\varphi} \\
& \times (1-x(1-t_2)(1-u_2)p_2)^{-(\nu+4)-\varphi} \\
& \times \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2+\nu)(i_0+\frac{3}{2}+\nu)} \frac{(\nu-\varphi)_{i_0}(\nu+\varphi)_{i_0}}{(1+\nu)_{i_0}(\frac{1}{2}+\nu)_{i_0}} \right. \\
& \times \left. \sum_{i_1=i_0}^{\infty} \frac{(\nu+2-\varphi)_{i_1}(\nu+2+\varphi)_{i_1}(3+\nu)_{i_0}(\frac{5}{2}+\nu)_{i_0}}{(\nu+2-\varphi)_{i_0}(\nu+2+\varphi)_{i_0}(3+\nu)_{i_1}(\frac{5}{2}+\nu)_{i_1}} w_{2,2}^{i_1} \right\} \eta^2 \quad (7.2.31)
\end{aligned}$$

where

$$w_{2,2} = \frac{p_2}{p_2 - 1} \frac{xt_2u_2}{1 - x(1 - t_2)(1 - u_2)p_2}$$

Put $l = 1$ and $\eta = w_{2,2}$ in (7.2.27). Take the new (7.2.27) into (7.2.31).

$$\begin{aligned} y_2(x) &= \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} \left(\frac{p_2 - 1}{p_2} \right)^{-(\nu+4)+\varphi} \\ &\quad \times (1 - x(1 - t_2)(1 - u_2)p_2)^{-(\nu+4)-\varphi} \\ &\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} \left(\frac{p_1 - 1}{p_1} \right)^{-(\nu+2)+\varphi} \\ &\quad \times (1 - w_{2,2}(1 - t_1)(1 - u_1)p_1)^{-(\nu+2)-\varphi} \left\{ c_0 x^\nu \sum_{i_0=0}^{\infty} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} w_{1,2}^{i_0} \right\} \eta^2 \quad (7.2.32) \end{aligned}$$

where

$$w_{1,2} = \frac{p_1}{p_1 - 1} \frac{w_{2,2}t_1u_1}{1 - w_{2,2}(1 - t_1)(1 - u_1)p_1}$$

By using similar process for the previous cases of integral forms of $y_1(x)$ and $y_2(x)$, the integral form of sub-power series expansion of $y_3(x)$ is

$$\begin{aligned} y_3(x) &= \int_0^1 dt_3 t_3^{5+\nu} \int_0^1 du_3 u_3^{\frac{9}{2}+\nu} \frac{1}{2\pi i} \oint dp_3 \frac{1}{p_3} \left(\frac{p_3 - 1}{p_3} \right)^{-(\nu+6)+\varphi} \\ &\quad \times (1 - x(1 - t_3)(1 - u_3)p_3)^{-(\nu+6)-\varphi} \\ &\quad \times \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} \left(\frac{p_2 - 1}{p_2} \right)^{-(\nu+4)+\varphi} \\ &\quad \times (1 - w_{3,3}(1 - t_2)(1 - u_2)p_2)^{-(\nu+4)-\varphi} \\ &\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} \left(\frac{p_1 - 1}{p_1} \right)^{-(\nu+2)+\varphi} \\ &\quad \times (1 - w_{2,3}(1 - t_1)(1 - u_1)p_1)^{-(\nu+2)-\varphi} \left\{ c_0 x^\nu \sum_{i_0=0}^{\infty} \frac{(\nu - \varphi)_{i_0} (\nu + \varphi)_{i_0}}{(1 + \nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} w_{1,3}^{i_0} \right\} \eta^3 \quad (7.2.33) \end{aligned}$$

where

$$\begin{cases} w_{3,3} = \frac{p_3}{p_3 - 1} \frac{xt_3u_3}{1 - x(1 - t_3)(1 - u_3)p_3} \\ w_{2,3} = \frac{p_2}{p_2 - 1} \frac{w_{3,3}t_2u_2}{1 - w_{3,3}(1 - t_2)(1 - u_2)p_2} \\ w_{1,3} = \frac{p_1}{p_1 - 1} \frac{w_{2,3}t_1u_1}{1 - w_{2,3}(1 - t_1)(1 - u_1)p_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute

(7.2.29a), (7.2.30), (7.2.32), (7.2.33) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (7.2.7), we obtain (7.2.28).⁸

Put $c_0 = 1$ as $\nu = 0$ for the first kind of independent solutions of Mathieu equation and $\nu = 1/2$ for the second one in (7.2.28).

Remark 7.2.9 *The integral representation of Mathieu equation of the first kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
 y(x) &= MF^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda + 2q}; \eta = qx^2, x = \cos^2 z \right) \\
 &= {}_2F_1 \left(-\varphi, \varphi; \frac{1}{2}; x \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} \left(\frac{p_{n-k}-1}{p_{n-k}} \right)^{-2(n-k)+\varphi} \\
 &\quad \times (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-2(n-k)-\varphi} \left. \right\} \\
 &\quad \times {}_2F_1 \left(-\varphi, \varphi; \frac{1}{2}; w_{1,n} \right) \left. \right\} \eta^n
 \end{aligned} \tag{7.2.34}$$

Remark 7.2.10 *The integral representation of Mathieu equation of the second kind for infinite series about $x = 0$ using R3TRF is*

$$\begin{aligned}
 y(x) &= MS^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda + 2q}; \eta = qx^2, x = \cos^2 z \right) \\
 &= x^{\frac{1}{2}} \left\{ {}_2F_1 \left(\frac{1}{2} - \varphi, \frac{1}{2} + \varphi; \frac{3}{2}; x \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} \left(\frac{p_{n-k}-1}{p_{n-k}} \right)^{-(2(n-k)+\frac{1}{2})+\varphi} \\
 &\quad \times (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-(2(n-k)+\frac{1}{2})-\varphi} \left. \right\} \\
 &\quad \times {}_2F_1 \left(\frac{1}{2} - \varphi, \frac{1}{2} + \varphi; \frac{3}{2}; w_{1,n} \right) \left. \right\} \eta^n
 \end{aligned} \tag{7.2.35}$$

⁸Or replace the finite summation with an interval $[0, \lambda_0]$ by infinite summation with an interval $[0, \infty]$ in (7.2.16). Replace λ_0 and λ_{n-k} by $\frac{1}{2}\sqrt{\lambda+2q} - \nu$ and $\frac{1}{2}\sqrt{\lambda+2q} - 2(n-k) - \nu$ into the new (7.2.16). Its solution is also equivalent to (7.2.28)

Two integral forms of Mathieu equation for infinite series in this chapter and Ref.[21] are equivalent to each other. In this chapter, ${}_2F_1$ function recurs in each of sub-integral forms of the Mathieu function. In Ref.[21], the Modified Bessel function recurs in each of sub-integral forms of it.

7.2.3 Generating function for the Mathieu polynomial of type 2

I consider the generating function for the Mathieu polynomial of type 2. Since its generating function is derived, we might be possible to construct orthogonal relations of it.

Lemma 7.2.11 *The generating function for Jacobi polynomial using hypergeometric functions is given by*

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\gamma)_{\lambda_0}}{\lambda_0!} w^{\lambda_0} {}_2F_1(-\lambda_0, \lambda_0 + A; \gamma; x) \\ &= 2^{A-1} \frac{\left(1 - w + \sqrt{w^2 - 2(1 - 2x)w + 1}\right)^{1-\gamma} \left(1 + w + \sqrt{w^2 - 2(1 - 2x)w + 1}\right)^{\gamma-A}}{\sqrt{w^2 - 2(1 - 2x)w + 1}} \\ & \text{where } |w| < 1 \end{aligned} \quad (7.2.36)$$

Proof Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ can be written in terms of hypergeometric function using

$${}_2F_1(-n, n + \alpha + \beta + 1; \alpha + 1; x) = \frac{n!}{(\alpha + 1)_n} P_n^{(\alpha, \beta)}(1 - 2x) \quad (7.2.37)$$

And

$$P_n^{(\alpha, \beta)}(x) = \frac{\Gamma(n + \alpha + 1)}{n! \Gamma(n + \alpha + \beta + 1)} \sum_{m=0}^n \binom{n}{m} \frac{\Gamma(n + m + \alpha + \beta + 1)}{\Gamma(m + \alpha + 1)} \left(\frac{x - 1}{2}\right)^m \quad (7.2.38)$$

The generating function for the Jacobi polynomials is written by

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x) w^n = 2^{\alpha+\beta} \frac{\left(1 - w + \sqrt{w^2 - 2xw + 1}\right)^{-\alpha} \left(1 + w + \sqrt{w^2 - 2xw + 1}\right)^{-\beta}}{\sqrt{w^2 - 2xw + 1}} \quad (7.2.39)$$

Replace n, α and β by $\lambda_0, \gamma - 1$ and $A - \gamma$ in (7.2.37), and acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma)_{\lambda_0}}{\lambda_0!} w^{\lambda_0}$ on the new (7.2.37)

$$\sum_{\lambda_0=0}^{\infty} \frac{(\gamma)_{\lambda_0}}{\lambda_0!} w^{\lambda_0} {}_2F_1(-\lambda_0, \lambda_0 + A; \gamma; x) = \sum_{\lambda_0=0}^{\infty} P_n^{(\gamma-1, A-\gamma)}(1-2x) w^{\lambda_0} \quad (7.2.40)$$

Replace α, β and x by $\gamma - 1, A - \gamma$ and $1 - 2x$ in (7.2.39). As we take the new (7.2.39) into (7.2.40), we obtain (7.2.36). ■

Definition 7.2.12 *I define that*

$$\left\{ \begin{array}{l} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{if } a > b \\ s_a & \text{if } a = b \end{cases} \\ \\ \tilde{w}_{i,j} = \begin{cases} \frac{\tilde{w}_{i+1,j} t_i u_i \{1 + (s_i + 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i\}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} \\ - \frac{\tilde{w}_{i+1,j} t_i u_i (1 + s_i) \sqrt{s_i^2 - 2(1 - 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i + 1}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} & \text{where } i < j \\ \\ \frac{x t_i u_i \{1 + (s_{i,\infty} + 2x(1-t_i)(1-u_i))s_{i,\infty}\}}{2(1 - x(1-t_i)(1-u_i))^2 s_{i,\infty}} \\ - \frac{x t_i u_i (1 + s_{i,\infty}) \sqrt{s_{i,\infty}^2 - 2(1 - 2x(1-t_i)(1-u_i))s_{i,\infty} + 1}}{2(1 - x(1-t_i)(1-u_i))^2 s_{i,\infty}} & \text{where } i = j \end{cases} \end{array} \right. \quad (7.2.41)$$

where

$$a, b, i, j \in \mathbb{N}_0$$

And we have

$$\sum_{\lambda_i=\lambda_j}^{\infty} s_i^{\lambda_i} = \frac{s_i^{\lambda_j}}{(1-s_i)} \quad (7.2.42)$$

Acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\}$ on (7.2.16) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (7.2.41) and (7.2.42),

Theorem 7.2.13 *The general expression of the generating function for the Mathieu polynomial of type 2 is given by*

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y(x) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \Upsilon(\nu; s_{0,\infty}; x) \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \overleftrightarrow{\Xi}_1(\nu; s_{1,\infty}; t_1, u_1, x) \Upsilon(\nu; s_0; \tilde{w}_{1,1}) \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1+\nu} \int_0^1 du_n u_n^{2n-\frac{3}{2}+\nu} \overleftrightarrow{\Xi}_n(\nu; s_{n,\infty}; t_n, u_n, x) \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\nu} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}+\nu} \overleftrightarrow{\Xi}_{n-k}(\nu; s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right\} \\
&\times \Upsilon(\nu; s_0; \tilde{w}_{1,n}) \left. \right\} \eta^n
\end{aligned} \tag{7.2.43}$$

where

$$\left\{ \begin{aligned}
& \overleftrightarrow{\Xi}_1(\nu; s_{1,\infty}; t_1, u_1, x) = \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-(3+2\nu)}}{\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}} \\
& \overleftrightarrow{\Xi}_n(\nu; s_{n,\infty}; t_n, u_n, x) = \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-(4n-1+2\nu)}}{\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}} \\
& \overleftrightarrow{\Xi}_{n-k}(\nu; s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\
& \quad \left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)-1+2\nu)} \\
& = \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)-1+2\nu)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}
\end{aligned} \right.$$

and

$$\begin{cases} \mathfrak{Y}(\nu; s_{0,\infty}; x) = \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_{0,\infty}^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} x^{i_0} \right) \\ \mathfrak{Y}(\nu; s_0; \tilde{w}_{1,1}) = \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \\ \mathfrak{Y}(\nu; s_0; \tilde{w}_{1,n}) = \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \tilde{w}_{1,n}^{i_0} \right) \end{cases}$$

Proof Acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\}$ on the form of integral of the Mathieu polynomial of type 2 $y(x)$,

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y(x) \\ &= \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} \{y_0(x) + y_1(x) + y_2(x) + y_3(x) + \dots\} \end{aligned} \quad (7.2.44)$$

Acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\}$ on (7.2.18a),

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_0(x) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_{0,\infty}^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} x^{i_0} \right) \end{aligned} \quad (7.2.45)$$

Acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\}$ on (7.2.19),

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-x(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\ &\times \sum_{\lambda_1=\lambda_0}^{\infty} \left(\frac{p_1-1}{p_1} \frac{s_{1,\infty}}{1-x(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2}+\nu)_{i_0}} w_{1,1}^{i_0} \right) \eta \quad (7.2.46) \end{aligned}$$

Replace λ_i , λ_j and s_i by λ_1 , λ_0 and $\frac{p_1-1}{p_1} \frac{s_{1,\infty}}{1-x(1-t_1)(1-u_1)p_1}$ in (7.2.42). Take the new (7.2.42) into (7.2.46).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_1(x) \\ &= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{(1-x(1-t_1)(1-u_1)p_1)^{-(3+2\nu)}}{-x(1-t_1)(1-u_1)p_1^2 + (1-s_{1,\infty})p_1 + s_{1,\infty}} \\ &\times \sum_{\lambda_0=0}^{\infty} \left(\frac{p_1-1}{p_1} \frac{s_{0,\infty}}{1-x(1-t_1)(1-u_1)p_1} \right)^{\lambda_0} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2}+\nu)_{i_0}} w_{1,1}^{i_0} \right) \eta \quad (7.2.47) \end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$p_1 = \frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)}$$

$$\text{or } \frac{1-s_{1,\infty} + \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)} \text{ and}$$

$$\frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)} \text{ is only inside the unit circle. As we}$$

compute the residue there in (7.2.47) we obtain

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_1(x) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} (s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+2\nu)} \\
&\quad \times \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \eta
\end{aligned} \tag{7.2.48}$$

where

$$\begin{aligned}
\tilde{w}_{1,1} &= \frac{p_1}{(p_1 - 1)} \frac{xt_1 u_1}{1 - x(1-t_1)(1-u_1)p_1} \Big|_{p_1 = \frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4x(1-t_1)(1-u_1)s_{1,\infty}}}{2x(1-t_1)(1-u_1)}} \\
&= \frac{xt_1 u_1 \{1 + (s_{1,\infty} + 2x(1-t_1)(1-u_1))s_{1,\infty}\}}{2(1-x(1-t_1)(1-u_1))^2 s_{1,\infty}} \\
&\quad - \frac{xt_1 u_1 (1 + s_{1,\infty}) \sqrt{s_{1,\infty}^2 - 2(1-2x(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2(1-x(1-t_1)(1-u_1))^2 s_{1,\infty}}
\end{aligned}$$

Acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\}$ on (7.2.21),

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{1}{p_2} (1-x(1-t_2)(1-u_2)p_2)^{-(8+2\nu)} \\
&\quad \times \sum_{\lambda_2=\lambda_1}^{\infty} \left(\frac{p_2-1}{p_2} \frac{s_{2,\infty}}{1-x(1-t_2)(1-u_2)p_2} \right)^{\lambda_2} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-w_{2,2}(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\
&\quad \times \sum_{\lambda_1=\lambda_0}^{\infty} \left(\frac{p_1-1}{p_1} \frac{s_1}{1-w_{2,2}(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} w_{1,2}^{i_0} \right) \eta^2
\end{aligned} \tag{7.2.49}$$

Replace λ_i , λ_j and s_i by λ_2 , λ_1 and $\frac{p_2-1}{p_2} \frac{s_{2,\infty}}{1-x(1-t_2)(1-u_2)p_2}$ in (7.2.42). Take the new (7.2.42) into (7.2.49).

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} \frac{1}{2\pi i} \oint dp_2 \frac{(1-x(1-t_2)(1-u_2)p_2)^{-(7+2\nu)}}{-x(1-t_2)(1-u_2)p_2^2 + (1-s_{2,\infty})p_2 + s_{2,\infty}} \\
&\times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-w_{2,2}(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\
&\times \sum_{\lambda_1=\lambda_0}^{\infty} \left(\frac{p_2-1}{p_2} \frac{s_{1,\infty}}{1-x(1-t_2)(1-u_2)p_2} \frac{p_1-1}{p_1} \frac{1}{1-w_{2,2}(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \\
&\times \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+2\nu)_{i_0} w_{1,2}^{i_0}}{(1+\nu)_{i_0} (\frac{1}{2}+\nu)_{i_0}} \right) \eta^2 \tag{7.2.50}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$p_2 = \frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4x(1-t_2)(1-u_2)s_{2,\infty}}}{2x(1-t_2)(1-u_2)}$$

$$\text{or } \frac{1-s_{2,\infty} + \sqrt{(1-s_{2,\infty})^2 + 4x(1-t_2)(1-u_2)s_{2,\infty}}}{2x(1-t_2)(1-u_2)} \text{ and}$$

$$\frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4x(1-t_2)(1-u_2)s_{2,\infty}}}{2x(1-t_2)(1-u_2)}$$

is only inside the unit circle. As we compute the residue there in (7.2.50) we obtain

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} (s_{2,\infty}^2 - 2(1-2x(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1+s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2x(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+2\nu)} \\
&\times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{1}{p_1} (1-\tilde{w}_{2,2}(1-t_1)(1-u_1)p_1)^{-(4+2\nu)} \\
&\times \sum_{\lambda_1=\lambda_0}^{\infty} \left(\frac{p_1-1}{p_1} \frac{s_1}{1-\tilde{w}_{2,2}(1-t_1)(1-u_1)p_1} \right)^{\lambda_1} \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+2\nu)_{i_0} \tilde{w}_{1,2}^{i_0}}{(1+\nu)_{i_0} (\frac{1}{2}+\nu)_{i_0}} \right) \eta^2 \tag{7.2.51}
\end{aligned}$$

where

$$\begin{aligned}\tilde{w}_{2,2} &= \frac{p_2}{(p_2 - 1)} \frac{xt_2u_2}{1 - x(1 - t_2)(1 - u_2)p_2} \Bigg|_{p_2 = \frac{1 - s_{2,\infty} - \sqrt{(1 - s_{2,\infty})^2 + 4x(1 - t_2)(1 - u_2)s_{2,\infty}}}{2x(1 - t_2)(1 - u_2)}} \\ &= \frac{xt_2u_2 \{1 + (s_{2,\infty} + 2x(1 - t_2)(1 - u_2))s_{2,\infty}\}}{2(1 - x(1 - t_2)(1 - u_2))^2 s_{2,\infty}} \\ &\quad - \frac{xt_2u_2(1 + s_{2,\infty})\sqrt{s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1}}{2(1 - x(1 - t_2)(1 - u_2))^2 s_{2,\infty}}\end{aligned}$$

and

$$\ddot{w}_{1,2} = \frac{p_1}{(p_1 - 1)} \frac{\tilde{w}_{2,2}t_1u_1}{1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)p_1}$$

Replace λ_i , λ_j and s_i by λ_1 , λ_0 and $\frac{p_1 - 1}{p_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)p_1}$ in (7.2.42). Take the new (7.2.42) into (7.2.51).

$$\begin{aligned}& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_2(x) \\ &= \prod_{k=3}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} (s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\ &\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1 - 2x(1 - t_2)(1 - u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+2\nu)} \\ &\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} \frac{1}{2\pi i} \oint dp_1 \frac{(1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)p_1)^{-(3+2\nu)}}{-\tilde{w}_{2,2}(1 - t_1)(1 - u_1)p_1^2 + (1 - s_1)p_1 + s_1} \\ &\quad \times \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} \left(\frac{p_1 - 1}{p_1} \frac{s_{0,1}}{1 - \tilde{w}_{2,2}(1 - t_1)(1 - u_1)p_1} \right)^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1 + \nu)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) \eta^2 \quad (7.2.52)\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$\begin{aligned}p_1 &= \frac{1 - s_1 - \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,2}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,2}(1 - t_1)(1 - u_1)} \\ \text{or } &\frac{1 - s_1 + \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,2}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,2}(1 - t_1)(1 - u_1)} \text{ and} \\ &\frac{1 - s_1 - \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,2}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,2}(1 - t_1)(1 - u_1)} \text{ is only inside the unit circle. As we}\end{aligned}$$

compute the residue there in (7.2.52) we obtain

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_2(x) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} (s_{2,\infty}^2 - 2(1-2x(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2x(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+2\nu)} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} (s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-(3+2\nu)} \\
&\quad \times \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \tilde{w}_{1,2}^{i_0} \right) \eta^2 \tag{7.2.53}
\end{aligned}$$

where

$$\begin{aligned}
\tilde{w}_{1,2} &= \frac{p_1}{(p_1 - 1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)p_1} \Bigg|_{p_1 = \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)}} \\
&= \frac{\tilde{w}_{2,2} t_1 u_1 \left\{ 1 + (s_1 + 2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 - (1+s_1)\sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1} \right\}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1}
\end{aligned}$$

Acting the summation operator $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\}$ on (7.2.22),

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_3(x) \\
&= \prod_{k=4}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{5+\nu} \int_0^1 du_3 u_3^{\frac{9}{2}+\nu} (s_{3,\infty}^2 - 2(1-2x(1-t_3)(1-u_3))s_{3,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{3,\infty} + \sqrt{s_{3,\infty}^2 - 2(1-2x(1-t_3)(1-u_3))s_{3,\infty} + 1}}{2} \right)^{-(11+2\nu)} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\nu} \int_0^1 du_2 u_2^{\frac{5}{2}+\nu} (s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_2 + \sqrt{s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1}}{2} \right)^{-(7+2\nu)} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\nu} \int_0^1 du_1 u_1^{\frac{1}{2}+\nu} (s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-(3+2\nu)} \\
&\quad \times \sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \left(c_0 x^\nu \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0 + 2\nu)_{i_0} \tilde{w}_{1,3}^{i_0}}{(1+\nu)_{i_0} (\frac{1}{2} + \nu)_{i_0}} \right) \eta^3 \tag{7.2.54}
\end{aligned}$$

where

$$\begin{aligned}
\tilde{w}_{3,3} &= \frac{p_3}{(p_3 - 1)} \frac{xt_3u_3}{1 - x(1-t_3)(1-u_3)p_3} \Bigg|_{p_3 = \frac{1-s_{3,\infty} - \sqrt{(1-s_{3,\infty})^2 + 4x(1-t_3)(1-u_3)s_{3,\infty}}}{2x(1-t_3)(1-u_3)}} \\
&= \frac{xt_3u_3 \{1 + (s_{3,\infty} + 2x(1-t_3)(1-u_3))s_{3,\infty}\}}{2(1-x(1-t_3)(1-u_3))^2 s_{3,\infty}} \\
&\quad - \frac{xt_3u_3(1+s_{3,\infty})\sqrt{s_{3,\infty}^2 - 2(1-2x(1-t_3)(1-u_3))s_{3,\infty} + 1}}{2(1-x(1-t_3)(1-u_3))^2 s_{3,\infty}}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{2,3} &= \frac{p_2}{(p_2 - 1)} \frac{\tilde{w}_{3,3} t_2 u_2}{1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2)} \Bigg|_{p_2 = \frac{1-s_2 - \sqrt{(1-s_2)^2 + 4\tilde{w}_{3,3}(1-t_2)(1-u_2)s_2}}{2\tilde{w}_{3,3}(1-t_2)(1-u_2)}} \\
&= \frac{\tilde{w}_{3,3} t_2 u_2 \left\{ 1 + (s_2 + 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2 - (1 + s_2)\sqrt{s_2^2 - 2(1 - 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2 + 1} \right\}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2} \\
\tilde{w}_{1,3} &= \frac{p_1}{(p_1 - 1)} \frac{\tilde{w}_{2,3} t_1 u_1}{1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1)} \Bigg|_{p_1 = \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,3}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,3}(1-t_1)(1-u_1)}} \\
&= \frac{\tilde{w}_{2,3} t_1 u_1 \left\{ 1 + (s_1 + 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1 - (1 + s_1)\sqrt{s_1^2 - 2(1 - 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1 + 1} \right\}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1}
\end{aligned}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$

terms where $m \geq 4$, I obtain every $\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_m(x)$ terms.

Substitute (7.2.45), (7.2.48), (7.2.53), (7.2.54) and including all

$$\sum_{\lambda_0=0}^{\infty} \frac{(\gamma')_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} y_m(x) \text{ terms where } m > 3 \text{ into (7.2.44).} \quad \blacksquare$$

Remark 7.2.14 The generating function for the Mathieu polynomial of type 2 of the first kind about $x = 0$ as $\lambda = 2^2(\lambda_j + 2j)^2 - 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
&\sum_{\lambda_0=0}^{\infty} \frac{(\frac{1}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} MF_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j)^2 - 2q; \eta = qx^2, x = \cos^2 z) \\
&= \frac{1}{2} \left\{ \prod_{l=1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \mathbf{A}(s_{0,\infty}; x) \right. \\
&+ \prod_{l=2}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, x) \mathbf{A}(s_0; \tilde{w}_{1,1}) \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, x) \right. \\
&\times \left. \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} \eta^n \Bigg\}
\end{aligned} \tag{7.2.55}$$

where

$$\left\{ \begin{array}{l} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, x) = \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-3}}{\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, x) = \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-(4n-1)}}{\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) = \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)-1)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; x) = \frac{\left(1-s_{0,\infty}+\sqrt{s_{0,\infty}^2-2(1-2x)s_{0,\infty}+1}\right)^{\frac{1}{2}} \left(1+s_{0,\infty}+\sqrt{s_{0,\infty}^2-2(1-2x)s_{0,\infty}+1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2-2(1-2x)s_{0,\infty}+1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1-s_0+\sqrt{s_0^2-2(1-2\tilde{w}_{1,1})s_0+1}\right)^{\frac{1}{2}} \left(1+s_0+\sqrt{s_0^2-2(1-2\tilde{w}_{1,1})s_0+1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2-2(1-2\tilde{w}_{1,1})s_0+1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1-s_0+\sqrt{s_0^2-2(1-2\tilde{w}_{1,n})s_0+1}\right)^{\frac{1}{2}} \left(1+s_0+\sqrt{s_0^2-2(1-2\tilde{w}_{1,n})s_0+1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2-2(1-2\tilde{w}_{1,n})s_0+1}} \end{array} \right.$$

Proof Replace A , w and γ by 0 , $s_{0,\infty}$ and $\frac{1}{2}$ in (7.2.36).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{1}{2})_{\lambda_0}}{\lambda_0!} s_{0,\infty}^{\lambda_0} {}_2F_1\left(-\lambda_0, \lambda_0; \frac{1}{2}; x\right) \\ &= \frac{\left(1-s_{0,\infty}+\sqrt{s_{0,\infty}^2-2(1-2x)s_{0,\infty}+1}\right)^{\frac{1}{2}} \left(1+s_{0,\infty}+\sqrt{s_{0,\infty}^2-2(1-2x)s_{0,\infty}+1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2-2(1-2x)s_{0,\infty}+1}} \end{aligned} \quad (7.2.56)$$

Replace A , w , γ , x by 0 , s_0 , $\frac{1}{2}$ and $\tilde{w}_{1,1}$ in (7.2.36).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{1}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} {}_2F_1\left(-\lambda_0, \lambda_0; \frac{1}{2}; \tilde{w}_{1,1}\right) \\ &= \frac{\left(1-s_0+\sqrt{s_0^2-2(1-2\tilde{w}_{1,1})s_0+1}\right)^{\frac{1}{2}} \left(1+s_0+\sqrt{s_0^2-2(1-2\tilde{w}_{1,1})s_0+1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2-2(1-2\tilde{w}_{1,1})s_0+1}} \end{aligned} \quad (7.2.57)$$

Replace A, w, γ, x by $0, s_0, \frac{1}{2}$ and $\tilde{w}_{1,n}$ in (7.2.36).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{1}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} {}_2F_1 \left(-\lambda_0, \lambda_0; \frac{1}{2}; \tilde{w}_{1,n} \right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1} \right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1} \right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (7.2.58)$$

Put $c_0 = 1$, $\lambda = 0$ and $\gamma' = \frac{1}{2}$ in (7.2.43). Substitute (7.2.56), (7.2.57) and (7.2.58) into the new (7.2.43). ■

Remark 7.2.15 *The generating function for the Mathieu polynomial of type 2 of the second kind about $x = 0$ as $\lambda = 2^2(\lambda_j + 2j + 1/2)^2 - 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is*

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{3}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} MS_{\lambda_j}^R \left(q, \lambda = 2^2(\lambda_j + 2j + 1/2)^2 - 2q; \eta = qx^2, x = \cos^2 z \right) \\ &= x^{\frac{1}{2}} \left\{ \prod_{l=1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \mathbf{B}(s_{0,\infty}; x) + \prod_{l=2}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_1 t_1^{\frac{3}{2}} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, x) \mathbf{B}(s_0; \tilde{w}_{1,1}) \eta \right. \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\frac{1}{2}} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, x) \right. \\ &\times \left. \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right\} \mathbf{B}(s_0; \tilde{w}_{1,n}) \right\} \eta^n \Big\} \end{aligned} \quad (7.2.59)$$

where

$$\begin{cases} \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, x) = \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-4}}{\sqrt{s_{1,\infty}^2-2(1-2x(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, x) = \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-4n}}{\sqrt{s_{n,\infty}^2-2(1-2x(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) = \frac{\left(\frac{(1+s_{n-k})+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-4(n-k)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{cases}$$

and

$$\left\{ \begin{array}{l} \mathbf{B}(s_{0,\infty}; x) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2x)s_{0,\infty} + 1}\right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2x)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2x)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Proof Replace A , w and γ by 1, $s_{0,\infty}$ and $\frac{3}{2}$ in (7.2.36).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{3}{2})_{\lambda_0}}{\lambda_0!} s_{0,\infty}^{\lambda_0} {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; x\right) \\ &= \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2x)s_{0,\infty} + 1}\right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2x)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2x)s_{0,\infty} + 1}} \end{aligned} \quad (7.2.60)$$

Replace A , w , γ , x by 1, s_0 , $\frac{3}{2}$ and $\tilde{w}_{1,1}$ in (7.2.36).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{3}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; \tilde{w}_{1,1}\right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (7.2.61)$$

Replace A , w , γ , x by 1, s_0 , $\frac{3}{2}$ and $\tilde{w}_{1,n}$ in (7.2.36).

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{3}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; \tilde{w}_{1,n}\right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (7.2.62)$$

Put $c_0 = 1$, $\lambda = 1/2$ and $\gamma' = 3/2$ in (7.2.43). Substitute (7.2.60), (7.2.61) and (7.2.62) into the new (7.2.43). $\blacksquare$

7.3 Mathieu equation about regular singular point at one

Let $\zeta = 1 - x$ in (7.1.2) to obtain analytic solutions of the Mathieu equation about $x = 1$.

$$4\zeta(1 - \zeta)\frac{d^2y}{d\zeta^2} + 2(1 - 2\zeta)\frac{dy}{d\zeta} + (\lambda - 2q + 4q\zeta)y = 0 \quad (7.3.1)$$

As we compare (7.3.1) with (7.1.2), an independent variable and a coefficient on the above are correspondent to the following way.

$$\begin{aligned} x &\longrightarrow \zeta = 1 - x \\ q &\longrightarrow -q \end{aligned} \quad (7.3.2)$$

For the asymptotic behavior of the function $y(x)$ about $x = 1$ for infinite series, replace x by $\zeta = 1 - x$ in (7.1.8).

$$\lim_{n \gg 1} y(\zeta) \approx \frac{1}{x} \quad \text{where } x = \cos^2 z$$

For polynomial of type 2, replace x and q by $\zeta = 1 - x$ and $-q$ in (7.1.10).

$$\lim_{n \gg 1} y(\zeta) > {}_1F_2 \left(1; \frac{1}{2}, \frac{1}{2}; -\frac{q}{4}\zeta^2 \right) + \zeta I_0(\sqrt{-q}\zeta)$$

As we put (7.3.2) into analytic solutions of Mathieu equation about $x = 0$ using 3TRF[21] and R3TRF, we obtain (1) the Frobenius solution in closed form of Mathieu equation about $x = 1$, (2) its integral form, (3) the generating function for the Mathieu polynomial of type 2. The local solutions are as follow.

7.3.1 Mathieu equation about $x = 1$ using 3TRF

Power series for infinite series

Remark 7.3.1 *The power series expansion of Mathieu equation of the first kind for infinite series about $x = 1$ using 3TRF is*

$$\begin{aligned}
 y(\zeta) &= M^{(o)}F\left(q, \lambda; \zeta = 1 - x, \eta = -\frac{1}{4}q\zeta^2, x = \cos^2 z\right) \\
 &= \sum_{i_0=0}^{\infty} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} \eta^{i_0} + \left\{ \sum_{i_0=0}^{\infty} \frac{i_0^2 - \frac{1}{4^2}(\lambda - 2q)}{(i_0 + \frac{1}{2})(i_0 + \frac{1}{4})} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\frac{3}{2})_{i_0}(\frac{5}{4})_{i_0}}{(\frac{3}{2})_{i_1}(\frac{5}{4})_{i_1}} \eta^{i_1} \right\} \zeta \\
 &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{i_0^2 - \frac{1}{4^2}(\lambda - 2q)}{(i_0 + \frac{1}{2})(i_0 + \frac{1}{4})} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} \right. \\
 &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + \frac{k}{2})^2 - \frac{1}{4^2}(\lambda - 2q)}{(i_k + \frac{k}{2} + \frac{1}{2})(i_k + \frac{k}{2} + \frac{1}{4})} \frac{(1 + \frac{k}{2})_{i_{k-1}}(\frac{3}{4} + \frac{k}{2})_{i_{k-1}}}{(1 + \frac{k}{2})_{i_k}(\frac{3}{4} + \frac{k}{2})_{i_k}} \right\} \\
 &\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(1 + \frac{n}{2})_{i_{n-1}}(\frac{3}{4} + \frac{n}{2})_{i_{n-1}}}{(1 + \frac{n}{2})_{i_n}(\frac{3}{4} + \frac{n}{2})_{i_n}} \eta^{i_n} \right\} \zeta^n \tag{7.3.3}
 \end{aligned}$$

Remark 7.3.2 *The power series expansion of Mathieu equation of the second kind for infinite series about $x = 1$ using 3TRF is*

$$\begin{aligned}
 y(\zeta) &= M^{(o)}S\left(q, \lambda; \zeta = 1 - x, \eta = -\frac{1}{4}q\zeta^2, x = \cos^2 z\right) \\
 &= \zeta^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}} \eta^{i_0} + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{4})^2 - \frac{1}{4^2}(\lambda - 2q)}{(i_0 + \frac{3}{4})(i_0 + \frac{1}{2})} \frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\frac{7}{4})_{i_0}(\frac{3}{2})_{i_0}}{(\frac{7}{4})_{i_1}(\frac{3}{2})_{i_1}} \eta^{i_1} \right\} \zeta \right. \\
 &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{4})^2 - \frac{1}{4^2}(\lambda - 2q)}{(i_0 + \frac{3}{4})(i_0 + \frac{1}{2})} \frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}} \right. \\
 &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + \frac{1}{4} + \frac{k}{2})^2 - \frac{1}{4^2}(\lambda - 2q)}{(i_k + \frac{3}{4} + \frac{k}{2})(i_k + \frac{1}{2} + \frac{k}{2})} \frac{(\frac{5}{4} + \frac{k}{2})_{i_{k-1}}(1 + \frac{k}{2})_{i_{k-1}}}{(\frac{5}{4} + \frac{k}{2})_{i_k}(1 + \frac{k}{2})_{i_k}} \right\} \\
 &\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\frac{5}{4} + \frac{n}{2})_{i_{n-1}}(1 + \frac{n}{2})_{i_{n-1}}}{(\frac{5}{4} + \frac{n}{2})_{i_n}(1 + \frac{n}{2})_{i_n}} \eta^{i_n} \right\} \zeta^n \left. \right\} \tag{7.3.4}
 \end{aligned}$$

Integral formalism for infinite series

Remark 7.3.3 *The integral representation of Mathieu equation of the first kind for infinite series about $x = 1$ using 3TRF is*

$$\begin{aligned}
y(\zeta) &= M^{(o)} F \left(q, \lambda; \zeta = 1 - x, \eta = -\frac{1}{4} q \zeta^2, x = \cos^2 z \right) \\
&= \Gamma(3/4) \left\{ \eta^{\frac{1}{8}} I_{-\frac{1}{4}}(2\sqrt{\eta}) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{-\frac{5}{4} + \frac{1}{2}(n-k)} \int_0^1 du_{n-k} u_{n-k}^{-1 + \frac{1}{2}(n-k)} \right. \right. \right. \\
&\quad \times I_0 \left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})} \right) \\
&\quad \times \left. \left. \left. \left(w_{n-k,n}^{-\frac{1}{2}(n-1-k)} (w_{n-k,n} \partial_{w_{n-k,n}})^2 w_{n-k,n}^{\frac{1}{2}(n-1-k)} - \frac{1}{4^2} (\lambda - 2q) \right) \right\} w_{1,n}^{\frac{1}{8}} I_{-\frac{1}{4}}(2\sqrt{w_{1,n}}) \right\} \zeta^n \right\} \quad (7.3.5)
\end{aligned}$$

Remark 7.3.4 *The integral representation of Mathieu equation of the second kind for infinite series about $x = 1$ using 3TRF is*

$$\begin{aligned}
y(\zeta) &= M^{(o)} S \left(q, \lambda; \zeta = 1 - x, \eta = -\frac{1}{4} q \zeta^2, x = \cos^2 z \right) \\
&= \Gamma(5/4) \zeta^{\frac{1}{2}} \left\{ \eta^{-\frac{1}{8}} I_{\frac{1}{4}}(2\sqrt{\eta}) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{-1 + \frac{1}{2}(n-k)} \int_0^1 du_{n-k} u_{n-k}^{-\frac{3}{4} + \frac{1}{2}(n-k)} \right. \right. \right. \\
&\quad \times I_0 \left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})} \right) \\
&\quad \times \left. \left. \left. \left(w_{n-k,n}^{-\frac{1}{2}(n-k-\frac{1}{2})} (w_{n-k,n} \partial_{w_{n-k,n}})^2 w_{n-k,n}^{\frac{1}{2}(n-k-\frac{1}{2})} - \frac{1}{4^2} (\lambda - 2q) \right) \right\} w_{1,n}^{-\frac{1}{8}} I_{\frac{1}{4}}(2\sqrt{w_{1,n}}) \right\} \zeta^n \right\} \quad (7.3.6)
\end{aligned}$$

In (7.3.5) and (7.3.6),

$$w_{a,b} = \begin{cases} \eta \prod_{l=a}^b t_l u_l & \text{where } a \leq b \\ \eta & \text{only if } a > b \end{cases}$$

7.3.2 Mathieu equation about $x = 1$ using R3TRF**Power series****Polynomial of type 2**

Remark 7.3.5 The power series expansion of Mathieu equation of the first kind for polynomial of type 2 about $x = 1$ as $\lambda = 2^2(\lambda_j + 2j)^2 + 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(\zeta) &= M^{(o)} F_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j)^2 + 2q; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z) \\
&= \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \zeta^{i_0} \\
&\quad + \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2)(i_0+\frac{3}{2})} \frac{(-\lambda_0)_{i_0} (\lambda_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1} (\lambda_1+4)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-\lambda_1)_{i_0} (\lambda_1+4)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \zeta^{i_1} \right\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+2)(i_0+\frac{3}{2})} \frac{(-\lambda_0)_{i_0} (\lambda_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\lambda_k} \frac{1}{(i_k+2k+2)(i_k+2k+\frac{3}{2})} \frac{(-\lambda_k)_{i_k} (\lambda_k+4k)_{i_k} (2k+1)_{i_{k-1}} (2k+\frac{1}{2})_{i_{k-1}}}{(-\lambda_k)_{i_{k-1}} (\lambda_k+4k)_{i_{k-1}} (2k+1)_{i_k} (2k+\frac{1}{2})_{i_k}} \right\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\lambda_n} \frac{(-\lambda_n)_{i_n} (\lambda_n+4n)_{i_n} (2n+1)_{i_{n-1}} (2n+\frac{1}{2})_{i_{n-1}}}{(-\lambda_n)_{i_{n-1}} (\lambda_n+4n)_{i_{n-1}} (2n+1)_{i_n} (2n+\frac{1}{2})_{i_n}} \zeta^{i_n} \left. \right\} \eta^n \tag{7.3.7}
\end{aligned}$$

Remark 7.3.6 The power series expansion of Mathieu equation of the second kind for polynomial of type 2 about $x = 1$ as $\lambda = 2^2(\lambda_j + 2j + 1/2)^2 + 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(\zeta) &= M^{(o)} S_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j + 1/2)^2 + 2q; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z) \\
&= \zeta^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{(-\lambda_0)_{i_0} (\lambda_0+1)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \zeta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+\frac{5}{2})(i_0+2)} \frac{(-\lambda_0)_{i_0} (\lambda_0+1)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\lambda_1} \frac{(-\lambda_1)_{i_1} (\lambda_1+5)_{i_1} (\frac{7}{2})_{i_0} (3)_{i_0}}{(-\lambda_1)_{i_0} (\lambda_1+5)_{i_0} (\frac{7}{2})_{i_1} (3)_{i_1}} \zeta^{i_1} \right\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\lambda_0} \frac{1}{(i_0+\frac{5}{2})(i_0+2)} \frac{(-\lambda_0)_{i_0} (\lambda_0+1)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\lambda_k} \frac{1}{(i_k+2k+\frac{5}{2})(i_k+2k+2)} \frac{(-\lambda_k)_{i_k} (\lambda_k+4k+1)_{i_k} (2k+\frac{3}{2})_{i_{k-1}} (2k+1)_{i_{k-1}}}{(-\lambda_k)_{i_{k-1}} (\lambda_k+4k+1)_{i_{k-1}} (2k+\frac{3}{2})_{i_k} (2k+1)_{i_k}} \right\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\lambda_n} \frac{(-\lambda_n)_{i_n} (\lambda_n+4n+1)_{i_n} (2n+\frac{3}{2})_{i_{n-1}} (2n+1)_{i_{n-1}}}{(-\lambda_n)_{i_{n-1}} (\lambda_n+4n+1)_{i_{n-1}} (2n+\frac{3}{2})_{i_n} (2n+1)_{i_n}} \zeta^{i_n} \left. \right\} \eta^n \left. \right\} \tag{7.3.8}
\end{aligned}$$

Infinite series

Remark 7.3.7 *The power series expansion of Mathieu equation of the first kind for infinite series about $x = 1$ using R3TRF is*

$$\begin{aligned}
y(\zeta) &= M^{(o)} F^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda - 2q}; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z \right) \\
&= \sum_{i_0=0}^{\infty} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \zeta^{i_0} + \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2) (i_0+\frac{3}{2})} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(2-\varphi)_{i_1} (2+\varphi)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(2-\varphi)_{i_0} (2+\varphi)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \zeta^{i_1} \right\} \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2) (i_0+\frac{3}{2})} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{1}{(i_k+2k+2) (i_k+2k+\frac{3}{2})} \frac{(2k-\varphi)_{i_k} (2k+\varphi)_{i_k} (2k+1)_{i_{k-1}} (2k+\frac{1}{2})_{i_{k-1}}}{(2k-\varphi)_{i_{k-1}} (2k+\varphi)_{i_{k-1}} (2k+1)_{i_k} (2k+\frac{1}{2})_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(2n-\varphi)_{i_n} (2n+\varphi)_{i_n} (2n+1)_{i_{n-1}} (2n+\frac{1}{2})_{i_{n-1}}}{(2n-\varphi)_{i_{n-1}} (2n+\varphi)_{i_{n-1}} (2n+1)_{i_n} (2n+\frac{1}{2})_{i_n}} \zeta^{i_n} \right\} \eta^n \quad (7.3.9)
\end{aligned}$$

Remark 7.3.8 *The power series expansion of Mathieu equation of the second kind for infinite series about $x = 1$ using R3TRF is*

$$\begin{aligned}
y(\zeta) &= M^{(o)} S^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda - 2q}; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z \right) \\
&= \zeta^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{(\frac{1}{2}-\varphi)_{i_0} (\frac{1}{2}+\varphi)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \zeta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+\frac{5}{2}) (i_0+2)} \frac{(\frac{1}{2}-\varphi)_{i_0} (\frac{1}{2}+\varphi)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\frac{5}{2}-\varphi)_{i_1} (\frac{5}{2}+\varphi)_{i_1} (\frac{7}{2})_{i_0} (3)_{i_0}}{(\frac{5}{2}-\varphi)_{i_0} (\frac{5}{2}+\varphi)_{i_0} (\frac{7}{2})_{i_1} (3)_{i_1}} \zeta^{i_1} \right\} \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{1}{(i_0+\frac{5}{2}) (i_0+2)} \frac{(\frac{1}{2}-\varphi)_{i_0} (\frac{1}{2}+\varphi)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{1}{(i_k+2k+\frac{5}{2}) (i_k+2k+2)} \frac{(2k+\frac{1}{2}-\varphi)_{i_k} (2k+\frac{1}{2}+\varphi)_{i_k} (2k+\frac{3}{2})_{i_{k-1}} (2k+1)_{i_{k-1}}}{(2k+\frac{1}{2}-\varphi)_{i_{k-1}} (2k+\frac{1}{2}+\varphi)_{i_{k-1}} (2k+\frac{3}{2})_{i_k} (2k+1)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(2n+\frac{1}{2}-\varphi)_{i_n} (2n+\frac{1}{2}+\varphi)_{i_n} (2n+\frac{3}{2})_{i_{n-1}} (2n+1)_{i_{n-1}}}{(2n+\frac{1}{2}-\varphi)_{i_{n-1}} (2n+\frac{1}{2}+\varphi)_{i_{n-1}} (2n+\frac{3}{2})_{i_n} (2n+1)_{i_n}} \zeta^{i_n} \right\} \eta^n \quad (7.3.10)
\end{aligned}$$

The case of small values of q As $|q| \ll 1$ in (7.3.9) and (7.3.10), we have

$$\begin{aligned}
& \lim_{|q| \ll 1} M^{(o)} F^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda - 2q}; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z \right) \\
& \approx \sum_{i_0=0}^{\infty} \frac{\left(-\frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{\sqrt{\lambda}}{2}\right)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \zeta^{i_0} \\
& + \eta \sum_{i_0=0}^{\infty} \frac{1}{(i_0+2) \left(i_0 + \frac{3}{2}\right)} \frac{\left(-\frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{\sqrt{\lambda}}{2}\right)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{\left(2 - \frac{\sqrt{\lambda}}{2}\right)_{i_1} \left(2 + \frac{\sqrt{\lambda}}{2}\right)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{\left(2 - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(2 + \frac{\sqrt{\lambda}}{2}\right)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} \zeta^{i_1} \\
& > \cos \left(\sqrt{\lambda} \sin^{-1} \left(\sqrt{\zeta} \right) \right) \tag{7.3.11a}
\end{aligned}$$

And,

$$\begin{aligned}
& \lim_{|q| \ll 1} M^{(o)} S^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda - 2q}; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z \right) \\
& \approx \zeta^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{\left(\frac{1}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{1}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \zeta^{i_0} \right. \\
& + \eta \sum_{i_0=0}^{\infty} \frac{1}{\left(i_0 + \frac{5}{2}\right) (i_0+2)} \frac{\left(\frac{1}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{1}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{\left(\frac{5}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_1} \left(\frac{5}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_1} \left(\frac{7}{2}\right)_{i_0} (3)_{i_0}}{\left(\frac{5}{2} - \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{5}{2} + \frac{\sqrt{\lambda}}{2}\right)_{i_0} \left(\frac{7}{2}\right)_{i_1} (3)_{i_1}} \zeta^{i_1} \left. \right\} \\
& > \sqrt{\frac{\zeta}{1-\zeta}} \cos \left(\sqrt{\lambda} \sin^{-1} \left(\sqrt{\zeta} \right) \right) \tag{7.3.11b}
\end{aligned}$$

Integral formalism

Polynomial of type 2

Remark 7.3.9 *The integral representation of Mathieu equation of the first kind for*

polynomial of type 2 about $x = 1$ as $\lambda = 2^2(\lambda_j + 2j)^2 + 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(\zeta) &= M^{(o)} F_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j)^2 + 2q; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z) \\
&= {}_2F_1\left(-\lambda_0, \lambda_0; \frac{1}{2}; \zeta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-4(n-k)} \\
&\quad \times \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k}} \right)^{\lambda_{n-k}} \left. \right\} \\
&\quad \times {}_2F_1\left(-\lambda_0, \lambda_0; \frac{1}{2}; w_{1,n}\right) \Big\} \eta^n \tag{7.3.12}
\end{aligned}$$

Remark 7.3.10 The integral representation of Mathieu equation of the second kind for polynomial of type 2 about $x = 1$ as $\lambda = 2^2(\lambda_j + 2j + 1/2)^2 + 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(\zeta) &= M^{(o)} S_{\lambda_j}^R(q, \lambda = 2^2(\lambda_j + 2j + 1/2)^2 + 2q; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z) \\
&= \zeta^{\frac{1}{2}} \left\{ {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; \zeta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k})^{-(4(n-k)+1)} \\
&\quad \times \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})p_{n-k}} \right)^{\lambda_{n-k}} \left. \right\} \\
&\quad \times {}_2F_1\left(-\lambda_0, \lambda_0 + 1; \frac{3}{2}; w_{1,n}\right) \Big\} \eta^n \tag{7.3.13}
\end{aligned}$$

Infinite series

Remark 7.3.11 The integral representation of Mathieu equation of the first kind for

infinite series about $x = 1$ using R3TRF is

$$\begin{aligned}
y(\zeta) &= M^{(o)} F^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda - 2q}; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z \right) \\
&= {}_2F_1 \left(-\varphi, \varphi; \frac{1}{2}; \zeta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) p_{n-k})^{-4(n-k)} \\
&\quad \times \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) p_{n-k}} \right)^{-2(n-k)+\varphi} \left. \right\} \\
&\quad \times {}_2F_1 \left(-\varphi, \varphi; \frac{1}{2}; w_{1,n} \right) \Big\} \eta^n
\end{aligned} \tag{7.3.14}$$

Remark 7.3.12 The integral representation of Mathieu equation of the second kind for infinite series about $x = 1$ using R3TRF is

$$\begin{aligned}
y(\zeta) &= M^{(o)} S^R \left(q, \lambda, \varphi = \frac{1}{2} \sqrt{\lambda - 2q}; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z \right) \\
&= \zeta^{\frac{1}{2}} \left\{ {}_2F_1 \left(\frac{1}{2} - \varphi, \frac{1}{2} + \varphi; \frac{3}{2}; \zeta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dp_{n-k} \frac{1}{p_{n-k}} (1 - w_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) p_{n-k})^{-(4(n-k)+1)} \\
&\quad \times \left(\frac{p_{n-k} - 1}{p_{n-k}} \frac{1}{1 - w_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) p_{n-k}} \right)^{-2(n-k)-\frac{1}{2}+\varphi} \left. \right\} \\
&\quad \times {}_2F_1 \left(\frac{1}{2} - \varphi, \frac{1}{2} + \varphi; \frac{3}{2}; w_{1,n} \right) \Big\} \eta^n
\end{aligned} \tag{7.3.15}$$

In (7.3.12)–(7.3.15),

$$w_{i,j} = \begin{cases} \frac{p_i}{(p_i - 1)} \frac{w_{i+1,j} t_i u_i}{1 - w_{i+1,j} p_i (1 - t_i)(1 - u_i)} & \text{where } i \leq j \\ \zeta & \text{only if } i > j \end{cases}$$

Generating function for the Mathieu polynomial of type 2

Remark 7.3.13 *The generating function for the Mathieu polynomial of type 2 of the first kind about $x = 1$ as $\lambda = 2^2(\lambda_j + 2j)^2 + 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
& \sum_{\lambda_0=0}^{\infty} \frac{(\frac{1}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} M^{(o)} F_{\lambda_j}^R (q, \lambda = 2^2(\lambda_j + 2j)^2 + 2q; \zeta = 1 - x, \eta = -q\zeta^2, x = \cos^2 z) \\
&= \frac{1}{2} \left\{ \prod_{l=1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \mathbf{A}(s_{0,\infty}; \zeta) \right. \\
&+ \prod_{l=2}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \zeta) \mathbf{A}(s_0; \tilde{w}_{1,1}) \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \zeta) \right. \\
&\times \left. \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} \eta^n \Bigg\}
\end{aligned} \tag{7.3.16}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \zeta) &= \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\zeta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-3}}{\sqrt{s_{1,\infty}^2 - 2(1-2\zeta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \zeta) &= \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\zeta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\zeta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{aligned} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \zeta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\zeta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\zeta)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\zeta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Remark 7.3.14 The generating function for the Mathieu polynomial of type 2 of the second kind about $x = 1$ as $\lambda = 2^2(\lambda_j + 2j + 1/2)^2 + 2q$ where $j, \lambda_j \in \mathbb{N}_0$ is

$$\begin{aligned} & \sum_{\lambda_0=0}^{\infty} \frac{(\frac{3}{2})_{\lambda_0}}{\lambda_0!} s_0^{\lambda_0} \prod_{n=1}^{\infty} \left\{ \sum_{\lambda_n=\lambda_{n-1}}^{\infty} s_n^{\lambda_n} \right\} M^{(o)} S_{\lambda_j}^R \left(q, \lambda = 2^2(\lambda_j + 2j + 1/2)^2 + 2q; \zeta = 1 - x \right. \\ & \left. , \eta = -q\zeta^2, x = \cos^2 z \right) \\ &= \zeta^{\frac{1}{2}} \left\{ \prod_{l=1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \mathbf{B}(s_{0,\infty}; \zeta) \right. \\ &+ \prod_{l=2}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_1 t_1^{\frac{3}{2}} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \zeta) \mathbf{B}(s_0; \tilde{w}_{1,1}) \eta \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\frac{1}{2}} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \zeta) \right. \\ &\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right\} \\ &\left. \times \mathbf{B}(s_0; \tilde{w}_{1,n}) \right\} \eta^n \Bigg\} \end{aligned} \quad (7.3.17)$$

where

$$\left\{ \begin{array}{l} \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \zeta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\zeta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-4}}{\sqrt{s_{1,\infty}^2 - 2(1-2\zeta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \zeta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\zeta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-4n}}{\sqrt{s_{n,\infty}^2 - 2(1-2\zeta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) = \frac{\left(\frac{(1+s_{n-k}) + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-4(n-k)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{B}(s_{0,\infty}; \zeta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\zeta)s_{0,\infty} + 1} \right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\zeta)s_{0,\infty} + 1} \right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1-2\zeta)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

In (7.3.16) and (7.3.17),

$$\left\{ \begin{array}{l} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{if } a > b \\ s_a & \text{if } a = b \end{cases} \\ \tilde{w}_{i,j} = \begin{cases} \frac{\tilde{w}_{i+1,j} t_i u_i \{1 + (s_i + 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i\}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} \\ - \frac{\tilde{w}_{i+1,j} t_i u_i (1 + s_i) \sqrt{s_i^2 - 2(1-2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i + 1}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} & \text{where } i < j \\ \frac{\zeta t_i u_i \{1 + (s_{i,\infty} + 2\zeta(1-t_i)(1-u_i))s_{i,\infty}\}}{2(1 - \zeta(1-t_i)(1-u_i))^2 s_{i,\infty}} \\ - \frac{\zeta t_i u_i (1 + s_{i,\infty}) \sqrt{s_{i,\infty}^2 - 2(1-2\zeta(1-t_i)(1-u_i))s_{i,\infty} + 1}}{2(1 - \zeta(1-t_i)(1-u_i))^2 s_{i,\infty}} & \text{where } i = j \end{cases} \end{array} \right.$$

where

$$a, b, i, j \in \mathbb{N}_0$$

7.4 Mathieu equation about irregular singular point at infinity

7.4.1 Mathieu equation about $x = \infty$

Let $\varsigma = \frac{1}{x}$ in (7.1.2) to obtain the analytic solution of Mathieu equation about $x = \infty$.

$$4\varsigma^3(\varsigma - 1)\frac{d^2y}{d\varsigma^2} + 2\varsigma^2(3\varsigma - 2)\frac{dy}{d\varsigma} + ((\lambda + 2q)\varsigma - 4q)y = 0 \quad (7.4.1)$$

Assume that its solution is

$$y(\varsigma) = \sum_{n=0}^{\infty} c_n \varsigma^{n+\nu} \quad (7.4.2)$$

Plug (7.4.2) into (7.4.1).

$$c_{n+1} = B_n c_{n-1} \quad ; n \geq 1 \quad (7.4.3)$$

where,

$$B_n = \frac{\left(n + \frac{1}{2}\sqrt{\lambda} - 1\right) \left(n + \frac{1}{2}\sqrt{\lambda} - \frac{1}{2}\right)}{n \left(n + \sqrt{\lambda}\right)} \quad (7.4.4a)$$

with

$$q = 0 \quad \nu = \frac{1}{2}\sqrt{\lambda} \quad (7.4.4b)$$

The Frobenius solution of (7.4.4a) with (7.4.4b) is

$$y(\varsigma) = M^{(i,1)}F \left(q = 0, \lambda; \varsigma = \frac{1}{x}, x = \cos^2 z \right) = \varsigma^{\frac{1}{2}\sqrt{\lambda}} {}_2F_1 \left(\frac{1}{2}\sqrt{\lambda}, \frac{1}{2}\sqrt{\lambda} + \frac{1}{2}; \sqrt{\lambda} + 1; \varsigma \right) \quad (7.4.5)$$

The power series of (7.4.4a) for polynomial does not exist. If $\sqrt{\lambda}$ is equal to $-2m$ or $-2m - 1$ where $m = 0, 1, 2, \dots$, the function $y(\varsigma)$ does not convergent because Pochhammer symbol $(\sqrt{\lambda} + 1)_n$ on the denominator will be zero at certain value of m . For infinite series, the series in $y(\varsigma)$ does not have circle $|\varsigma| = |\cos^{-2}z| < 1$ as its circle of convergence.

Lemma 7.4.1 *The hypergeometric functions is defined by*

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!} \quad \text{where } c \neq 0, -1, -2, \dots$$

converges for $|z| < 1$. On the boundary $|z| = 1$ of the region of convergence, a sufficient condition for absolute convergence of the series is $R(c - a - b) > 0$. [30]

By using this lemma, the analytic solution of (7.4.5) is given by

$$\begin{aligned} y(\varsigma) &= M^{(i,1)} F \left(q = 0, \lambda; \varsigma = \frac{1}{x} = 1, x = \cos^2 z \right) \\ &= {}_2F_1 \left(\frac{1}{2} \sqrt{\lambda}, \frac{1}{2} \sqrt{\lambda} + \frac{1}{2}; \sqrt{\lambda} + 1; 1 \right) = 2^{\sqrt{\lambda}} \quad \text{where } z = 0, \pm\pi, \pm 2\pi, \dots \end{aligned} \quad (7.4.6)$$

7.4.2 Mathieu equation about $\zeta = \infty$

Let $\phi = \frac{1}{\zeta}$ in (7.3.1) to obtain the analytic solution of Mathieu equation about $\zeta = \infty$.

$$4\phi^3(\phi - 1) \frac{d^2 y}{d\phi^2} + 2\phi^2(3\phi - 2) \frac{dy}{d\phi} + ((\lambda + 2q)\phi - 4q) y = 0 \quad (7.4.7)$$

As ς is replaced by ϕ in (7.4.1), its equation is correspondent to (7.4.7). The series of Mathieu equation about $\zeta = \infty$ is given by

$$\begin{aligned} y(\varsigma) &= M^{(i,2)} F \left(q = 0, \lambda; \phi = \frac{1}{\zeta} = 1, \zeta = 1 - x, x = \cos^2 z \right) \\ &= {}_2F_1 \left(\frac{1}{2} \sqrt{\lambda}, \frac{1}{2} \sqrt{\lambda} + \frac{1}{2}; \sqrt{\lambda} + 1; 1 \right) = 2^{\sqrt{\lambda}} \quad \text{where } z = \pm \frac{1}{2}\pi, \pm \frac{3}{2}\pi, \pm \frac{5}{2}\pi, \dots \end{aligned} \quad (7.4.8)$$

7.5 Summary

In Ref.[21] I show how to obtain the power series expansion in closed forms and its integral form of Mathieu equation for infinite series including all higher terms of A_n 's by applying 3TRF. This was done by letting A_n in sequence c_n is the leading term in the analytic function $y(x)$: the sequence c_n consists of combinations A_n and B_n . The series for the type 1 and 3 polynomials does not exist by using either 3TRF or R3TRF. Because there are no ways to terminate B_n term with a fixed parameter q of numerator in B_n whenever index n increases in (7.1.5b).

In this chapter I show how to construct the power series expansion in closed forms and its integral form of Mathieu equation for infinite series and polynomial of type 2 including all higher terms of B_n 's by applying R3TRF. This is done by letting B_n in sequence c_n is the leading term in the analytic function $y(x)$. For the type 2 polynomial, I treat q as a free variable and a fixed value of λ .

The power series expansion and its integral form of Mathieu equation for infinite series about $x = 0$ in this chapter are equivalent to infinite series of Mathieu equation in Ref.[21]. In this chapter B_n is the leading term in sequence c_n in the analytic function $y(x)$. In Ref.[21] A_n is the leading term in sequence c_n in the analytic function $y(x)$.

In Ref.[21] and this chapter, as we see the power series expansion of Mathieu equation about $x = 0$ for either polynomial or infinite series, the denominators and numerators in all A_n or B_n terms of $y(x)$ arise with Pochhammer symbol. Since we construct the power series expansion with Pochhammer symbols in numerators and denominators, we are able to describe integral representation of Mathieu equation analytically. As we observe representations in closed form integrals of Mathieu equation by applying 3TRF, a modified Bessel function recurs in each of sub-integral forms of the $y(x)$. And ${}_2F_1$ function recurs in each of sub-integral forms of $y(x)$ as we observe integral forms of Mathieu equation by applying R3TRF (each sub-integral $y_m(x)$ where $m = 0, 1, 2, \dots$ is composed of $2m$ terms of definite integrals and m terms of contour integrals). We are able to transform the Mathieu function into any well-known special functions having two recurrence relation in its power series of a linear differential equation because of modified Bessel and ${}_2F_1$ functions in each of sub-integral forms of the Mathieu function. After we replace modified Bessel and ${}_2F_1$ functions in its integral forms to other special functions (such as Bessel function, Kummer function, hypergeometric function, Laguerre function and etc), we are able to rebuild the power series expansion of Mathieu equation in a backward.

Indeed, the generating function for the Mathieu polynomial of type 2 is constructed analytically from its integral representation by applying the generating function for the Jacobi polynomial using hypergeometric functions. Also it might be possible to obtain orthogonal relations, recursion relations and expectation values of physical quantities from the generating function for the Mathieu polynomial of type 2: the processes in order to obtain orthogonal and recursion relations of the Mathieu polynomial are similar as the case of a normalized wave function for the hydrogen-like atoms.

Mathematical structure of the Frobenius solution, its integral representation and the generating function for Mathieu equation closely resembles the case of Confluent Heun equation using 3TRF and R3TRF. As $\beta \rightarrow 0$, $\delta = \gamma = \frac{1}{2}$, $\alpha = \frac{q}{0}$, $q \rightarrow \frac{\lambda+2q}{4}$ and $\lambda \rightarrow \nu$ in the power series expansion and its integral forms of the Confluent Heun equation using R3TRF in chapter 5, its analytic solution are correspondent to the Frobenius solution

and its integral representation of Mathieu equation in this chapter; compare remarks 5.2.1–5.2.9 in chapter 5 with remarks 7.2.1–7.2.9 in this chapter. By similar process, the analytic solution of the Confluent Heun equation using 3TRF for infinite series by changing all coefficients in chapter 4 is equivalent to the power series and its integral form of Mathieu equation in Ref.[21]; compare remarks 4.2.3, 4.2.4 and 4.2.8 and 4.2.9 in chapter 4 with remarks 1–4 in Ref.[21].

As $\tilde{w}_{1,1}^{-(\alpha+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\lambda}, \tilde{w}_{n,n}^{-(2(n-1)+\alpha+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\alpha+\lambda},$

$\tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\alpha+\lambda} \rightarrow 1$ with $\beta \rightarrow 0, \delta = \gamma = \frac{1}{2}, \alpha = \frac{q}{0},$

$q \rightarrow \frac{\lambda+2q}{4}$ and $\lambda \rightarrow \nu$ in the general expression of the generating function for the type 2 Confluent Heun polynomial using R3TRF in the chapter 5, its solution is equivalent to the generating function for the type 2 Mathieu polynomial in this chapter; compare theorem 5.2.12 in chapter 5 with theorem 7.2.12 in this chapter. Because, by applying the generating function for the Jacobi polynomial using ${}_2F_1$ functions into both integral forms of the Confluent Heun and Mathieu polynomials of type 2 (${}_2F_1$ function recurs in each of sub-integral forms of them), I derive generating functions for these polynomials of type 2.

Bibliography

- [1] McLachlan, N. W., *Theory and application of Mathieu functions*, Clarendon Press, Oxford, England, 1947.
- [2] Gutierrez-Vega, J. C., Rodriguez-Dagnino, R. M., Meneses-Nava, M. A. and Chavez-Cerda, S., "Mathieu functions, a visual approach," *Amer. J. Phys.* **71**(3), 233-242(2003).
- [3] Daymond, S. D., "The principal frequencies of vibrating systems with elliptic boundaries," *Quart. J. Mech. Appl. Math.* **8**(3), 361-372(1955).
- [4] Troesch, B. A. and Troesch, H. R., "Eigenfrequencies of an elliptic membrane," *Math. Comp.* **27**(124), 755-765(1973).
- [5] Alhargan, F. and Judah, S., "A general mode theory for the elliptic disk microstrip antenna," *IEEE Trans. Antennas and Propagation* **43**(6), 560-568(1995).
- [6] Shen, L., "The elliptical microstrip antenna with circular polarization," *IEEE Trans. Antennas and Propagation* **29**(1), 90-94(1981).
- [7] Bhattacharyya, A. and Shafai, L., "Theoretical and experimental investigation of the elliptical annular ring antenna," *IEEE Trans. Antennas and Propagation* **36**(11), 1526-1530(1988).
- [8] Ragheb, H. A., Shafai, L. and Hamid, M., "Plane wave scattering by a conducting elliptic cylinder coated by a nonconfocal dielectric," *IEEE Trans. Antennas and Propagation* **39**(2), 218-223(1991).
- [9] Aliev, A.N., Hortacsu, M., Kalayci, J., and Nutku, Y., "Gravitational instantons from minimal surfaces," *Class. Quantum Grav.* **16**, 631(1999).
- [10] Frenkel, D. and Protugal, R., "Algebraic Methods to Compute Mathieu Functions," *J. Phys. A: Math. Gen.* **34**, 3541(2001).

- [11] Mathieu, E., “Mémoire sur Le Mouvement Vibratoire d’une Membrane de forme Elliptique,” J. Math. Pure Appl. **13**, 137(1868).
- [12] Sips, R., “Représentation asymptotique des fonctions de Mathieu et des fonctions d’onde sphéroïdales,” Trans. Amer. Math. Soc. **66**, 93(1949).
- [13] Whittaker, E.T., “On the General Solution of Mathieu’s Equation,” Proc. Edinburgh Math. Soc. **32**, 75(1914).
- [14] Birkandan, T., Hortacsu, M., “Examples of Heun and Mathieu functions as solutions of wave equations in curved spaces,” J. Phys. A: Math. Theor. **40**, 1105–1116(2007).
- [15] Birkandan, T., Hortacsu, M., “Dirac equation in the background of the Nutku helicoid metric,” J. Phys. A: Math. Theor. **48**, 092301(2007).
- [16] Newman, E.T. and Penrose, R., “An Approach to Gravitational Radiation by a Method of Spin Coefficients,” J. Math. Phys. **3**, 566-768(1962).
- [17] Connor, J. N. L., Uzer, T., Marcus, R. A. and Smith, A. D., “Eigenvalues of the Schrödinger equation for a periodic potential with nonperiodic boundary conditions: A uniform semiclassical analysis,” J. Chem. Phys. **80**(10), 5095–6006(1984).
- [18] Zlatev, I., Huey, G. and Steinhardt, P.J., “Parametric Resonance in an Expanding Universe,” Phys. Rev. D **57**, 2152–2157(1998).
- [19] NIST Digital Library of Mathematical Functions, “Confluent Forms of Heuns Equation,” <http://dlmf.nist.gov/31.12>
- [20] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications.” arXiv:1303.0806.
- [21] Choun, Y.S., “The power series expansion of Mathieu function and its integral formalism,” arXiv:1303.0820.
- [22] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I., *Integrals and Series: More Special Functions*, Vol. 3, Gordon and Breach Science Publishers, New York, (1990).
- [23] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G., *Higher Transcendental Functions. Vol. III*, McGraw-Hill Book Company, Inc., New York-Toronto-London, (1955).
- [24] Gradshteyn, I. S. and Ryzhik, I. M., *Table of Integrals, Series, and Products*, 6th edition, Academic Press Inc., San Diego, CA., (2000).
- [25] Volkmer, H., “Integralgleichungen für periodische Lösungen Hill’scher Differentialgleichungen,” (German) Analysis **3** (1-4), 189-203(1983).

- [26] Meixner, J., Schäfer, F. W. and Wolf, G. *Mathieu Functions and Spheroidal Functions and Their Mathematical Foundations: Further Studies*, Lecture Notes in Mathematics, Vol. 837, Springer-Verlag, Berlin-New York, (1980).
- [27] Sips, R., “Quelques intégrales définies discontinues contenant des fonctions de Mathieu,” (French) Acad. Roy. Belg. Bull. Cl. Sci. (5) **56** (5), 475-491 (1970).
- [28] Arscott, F. M. *Periodic Differential Equations: An introduction to Mathieu, Lamé, and allied functions*, The Macmillan company, New York, (1964)
- [29] Wang, Z.X. and Guo, D. R., *Special Functions*, World Scientific Pub., (1989).
- [30] Rainville, E.D., *Special Functions*, The Macmillan Company (N.Y.); 1st edition, 45-47 (1960).

Chapter 8

Lame function in the algebraic form using reversible three-term recurrence formula

In Ref.[12] I apply three term recurrence formula (3TRF)[9] to the power series expansion in closed forms of Lamé equation in the algebraic form (for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n 's¹) and its integral form. I show how to transform power series expansion in Lamé equation to its integral representation for infinite series and polynomial in mathematical rigour.

In this chapter I will apply reversible three term recurrence formula (R3TRF) in chapter 1 to (1) the Frobenius solution in closed forms of Lamé equation, (2) its integral form (for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n 's²), (3) the generating function for the Lamé polynomial which makes A_n term terminated.

8.1 Introduction

The sphere is a geometrical perfect shape, the set of points which are all equidistant from its center (a fixed point) in three-dimensional space. In contrast, an ellipsoid is a imperfect one, a surface whose plane sections are all ellipses or circles; the set of points

¹“ higher terms of A_n 's” means at least two terms of A_n 's.

²“ higher terms of B_n 's” means at least two terms of B_n 's.

are not same distance from the center of the ellipsoid any more. As we all recognize, the nature is nonlinear and imperfect geometrically. For the purpose of simplification, we usually linearize those system in order to take a step to the future with a good numerical approximation. Actually, many geometrical spherical objects (earth, sun, black hole, etc) are not perfectly sphere in nature. The shape of those objects are closely better interpreted by an ellipsoid because of their rotations by themselves. For example, the ellipsoidal harmonics are represented in calculations of gravitational potential[8]. However spherical harmonic is preferred over the more mathematically complex ellipsoid harmonics (the coefficients in the power series expansion of Lamé equation have a relation between three different coefficients).

In 1837, Gabriel Lamé introduced a second ordinary differential equation which has four regular singular points in the method of separation of variables applied to the Laplace equation in elliptic coordinates[2]. Various authors has called this equation as ‘Lamé equation’ or ‘ellipsoidal harmonic equation’[17]. Lamé equation is applicable to diverse areas such as boundary value problems in ellipsoidal geometry, chaotic Hamiltonian systems, the theory of Bose-Einstein condensates, etc [3, 4, 5].

Until the present day, there are no analytic solutions in closed forms of the Lamé function[6, 7, 17]. Using Frobenius method to obtain an analytic solution of Lamé equation represented either in the algebraic form or in Weierstrass’s form, a solution automatically comes out three term recurrence relation[6, 7]. They left the analytic solution of Lamé equation as solutions of recurrences because of three different coefficients in its power series expansion. In comparison with two term recursion relation of power series in any linear ordinary differential equations, an analytic solution in closed forms on three term recurrence relation of power series is unknown currently because of its complex mathematical calculation.

In Ref.[12] I construct analytic solutions of Lamé equation in the algebraic form about the regular singular point at $x = a$ by applying 3TRF. [9]: the power series expansion in closed forms of Lamé equation and its integral form for infinite series and polynomial which makes B_n term terminated including all higher terms of A_n ’s. One interesting observation resulting from the calculations is the fact that a ${}_2F_1$ function recurs in each of sub-integral forms: the first sub-integral form contains zero term of A_n ’s, the second one contains one term of A_n ’s, the third one contains two terms of A_n ’s, etc.

In this chapter, by applying R3TRF in chapter 1, I construct the power series expansion in closed forms of Lamé equation in the algebraic form about a regular singular point at $x = a$ for infinite series and polynomial which makes A_n term terminated including all higher terms of B_n ’s. Indeed, an integral form of Lamé equation and the generating function of Lamé polynomial which makes A_n term terminated will be derived analytically.

Lame equation is a second-order linear ordinary differential equation of the algebraic form [2, 13, 15]

$$\frac{d^2 y}{dx^2} + \frac{1}{2} \left(\frac{1}{x-a} + \frac{1}{x-b} + \frac{1}{x-c} \right) \frac{dy}{dx} + \frac{-\alpha(\alpha+1)x+q}{4(x-a)(x-b)(x-c)} y = 0 \quad (8.1.1)$$

Lame equation has four regular singular points: a, b, c and ∞ ; the exponents at the first three are all 0 and $\frac{1}{2}$, and those at infinity are $-\frac{1}{2}\alpha$ and $\frac{1}{2}(\alpha+1)$. [14, 16] (8.1.1) is a special case of Heun equation. Heun equation is a second-order linear ordinary differential equation of the form [1].

$$\frac{d^2 y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} y = 0 \quad (8.1.2)$$

With the condition $\epsilon = \alpha + \beta - \gamma - \delta + 1$. The parameters play different roles: $a \neq 0$ is the singularity parameter, $\alpha, \beta, \gamma, \delta, \epsilon$ are exponent parameters, q is the accessory parameter which in many physical applications appears as a spectral parameter. Also, α and β are identical to each other. The total number of free parameters is six. It has four regular singular points which are 0, 1, a and ∞ with exponents $\{0, 1-\gamma\}$, $\{0, 1-\delta\}$, $\{0, 1-\epsilon\}$ and $\{\alpha, \beta\}$.

Let $z = x - a$ in (8.1.1).

$$\frac{d^2 y}{dz^2} + \frac{1}{2} \left(\frac{1}{z} + \frac{1}{z-(b-a)} + \frac{1}{z-(c-a)} \right) \frac{dy}{dz} + \frac{-\frac{1}{4}\alpha(\alpha+1)z - (-q + \frac{1}{4}\alpha(\alpha+1)a)}{z(z-(b-a))(z-(c-a))} y = 0 \quad (8.1.3)$$

As we compare (8.1.2) with (8.1.3), all coefficients on the above are correspondent to the following way.

$$\begin{aligned} \gamma, \delta, \epsilon &\longleftrightarrow \frac{1}{2} \\ 1 &\longleftrightarrow b-a \\ a &\longleftrightarrow c-a \\ \alpha &\longleftrightarrow \frac{1}{2}(\alpha+1) \\ \beta &\longleftrightarrow -\frac{1}{2}\alpha \\ q &\longleftrightarrow -q + \frac{1}{4}\alpha(\alpha+1)a \\ x &\longleftrightarrow z \end{aligned} \quad (8.1.4)$$

Assume that the solution of (8.1.3) is

$$y(z) = \sum_{n=0}^{\infty} c_n z^{n+\lambda} \quad (8.1.5)$$

where λ is an indicial root. Plug (8.1.5) into (8.1.3).

$$c_{n+1} = A_n c_n + B_n c_{n-1} \quad ; n \geq 1 \quad (8.1.6)$$

where,

$$\begin{aligned} A_n &= \frac{\frac{1}{4}(\alpha(\alpha+1)a - q) - (2a - b - c)(n + \lambda)^2}{(a - b)(a - c)(n + 1 + \lambda)(n + \frac{1}{2} + \lambda)} \\ &= -\frac{(2a - b - c)}{(a - b)(a - c)} \frac{(n - \varphi + \lambda)(n + \varphi + \lambda)}{(n + 1 + \lambda)(n + \frac{1}{2} + \lambda)} \end{aligned} \quad (8.1.7a)$$

and

$$\varphi = \sqrt{\frac{\alpha(\alpha+1)a - q}{4(2a - b - c)}}$$

$$\begin{aligned} B_n &= \frac{[\alpha - (1 - 2(n + \lambda))][\alpha - 2(n - 1 + \lambda)]}{2^2(a - b)(a - c)(n + 1 + \lambda)(n + \frac{1}{2} + \lambda)} \\ &= \frac{-1}{(a - b)(a - c)} \frac{(n + \frac{\alpha}{2} - \frac{1}{2} + \lambda)(n - \frac{\alpha}{2} - 1 + \lambda)}{(n + 1 + \lambda)(n + \frac{1}{2} + \lambda)} \end{aligned} \quad (8.1.7b)$$

$$c_1 = A_0 c_0 \quad (8.1.7c)$$

We have two indicial roots which are $\lambda = 0$ and $\frac{1}{2}$.

8.2 Power series

8.2.1 Polynomial of type 2

There are three types of polynomials in three-term recurrence relation of a linear ordinary differential equation: (1) polynomial which makes B_n term terminated: A_n term is not terminated, (2) polynomial which makes A_n term terminated: B_n term is not terminated, (3) polynomial which makes A_n and B_n terms terminated at the same time.³ In general Lamé (spectral) polynomial represented either in the algebraic form or in Weierstrass's form is defined as type 3 polynomial where A_n and B_n terms terminated. Lamé polynomial comes from a Lamé equation that has a fixed integer value of α , just as it has a fixed value of q . In three-term recurrence formula, polynomial of type 3 I categorize as complete polynomial. In future papers I will derive type 3 Lamé

³If A_n and B_n terms are not terminated, it turns to be infinite series.

polynomial. In Ref.[12] I derive the Frobenius solution and its integral form of the Lamé polynomial of type 1: I treat q as a free variable and α as a fixed value. In this chapter I construct the power series expansion and integral representation for the Lamé polynomial of type 2: I treat α as a free variable and q as a fixed value.

In chapter 1 the general expression of power series of $y(x)$ for polynomial of type 2 is defined by

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 \left\{ \sum_{i_0=0}^{\alpha_0} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\alpha_1} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
 &\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\alpha_k} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
 &\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\alpha_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (8.2.1)
 \end{aligned}$$

In the above, $\alpha_i \leq \alpha_j$ only if $i \leq j$ where $i, j, \alpha_i, \alpha_j \in \mathbb{N}_0$.

For a polynomial, we need a condition which is:

$$A_{\alpha_i+2i} = 0 \quad \text{where } i, \alpha_i = 0, 1, 2, \dots \quad (8.2.2)$$

In the above, α_i is an eigenvalue that makes A_n term terminated at certain value of index n . (8.2.2) makes each $y_i(x)$ where $i = 0, 1, 2, \dots$ as the polynomial in (8.2.1).

The case of $\varphi = -(q_i + 2i + \lambda)$ where $i, q_i = 0, 1, 2, \dots$

In (8.2.2) replace index α_i by q_i . In (8.1.7a)-(8.1.7c) replace φ by $-(q_i + 2i + \lambda)$. Take the new (8.1.7a)-(8.1.7c), (8.2.2) and put them in (8.2.1) with replacing variable x by z . After the replacement process, the general expression of power series of Lamé equation in

the algebraic form for polynomial of type 2 is given by

$$\begin{aligned}
y(z) &= \sum_{n=0}^{\infty} c_n z^{n+\lambda} = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \\
&= c_0 z^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \rho^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + 2\lambda)_{i_1} (3 + \lambda)_{i_1} (\frac{5}{2} + \lambda)_{i_1}}{(-q_1)_{i_0} (q_1 + 4 + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (\frac{5}{2} + \lambda)_{i_1}} \rho^{i_1} \Big\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_k + 2k - \frac{\alpha}{2} + \lambda)}{(i_k + 2k + 2 + \lambda) (i_k + 2k + \frac{3}{2} + \lambda)} \right. \\
&\quad \times \frac{(-q_k)_{i_k} (q_k + 4k + 2\lambda)_{i_k} (2k + 1 + \lambda)_{i_{k-1}} (2k + \frac{1}{2} + \lambda)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + 2\lambda)_{i_{k-1}} (2k + 1 + \lambda)_{i_k} (2k + \frac{1}{2} + \lambda)_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + 2\lambda)_{i_n} (2n + 1 + \lambda)_{i_{n-1}} (2n + \frac{1}{2} + \lambda)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + 2\lambda)_{i_{n-1}} (2n + 1 + \lambda)_{i_n} (2n + \frac{1}{2} + \lambda)_{i_n}} \rho^{i_n} \Big\} \eta^n \Big\} \quad (8.2.3)
\end{aligned}$$

where

$$\begin{cases} z = x - a \\ \rho = -\frac{2a-b-c}{(a-b)(a-c)} z \\ \eta = -\frac{1}{(a-b)(a-c)} z^2 \\ \varphi = -\beta + \gamma + \delta - 1 \\ q = \alpha(\alpha + 1)a - 4(2a - b - c)(q_j + 2j + \lambda)^2 \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

The case of $\varphi = q_i + 2i + \lambda$

In (8.2.2) replace index α_i by q_i . In (8.1.7a)-(8.1.7c) replace φ by $q_i + 2i + \lambda$. Take the new (8.1.7a)-(8.1.7c), (8.2.2) and put them in (8.2.1) with replacing variable x by z . After the replacement process, its solution is equivalent to (8.2.3).

Put $c_0 = 1$ as $\lambda = 0$ for the first independent solution of Lamé equation and $\lambda = \frac{1}{2}$ for the second one in (8.2.3).

Remark 8.2.1 *The power series expansion of Lamé equation in the algebraic form of the first kind for polynomial of type 2 about $x = a$ as $q = \alpha(\alpha + 1)a - 4(2a - b - c)(q_j + 2j)^2$ where $j, q_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
 y(z) &= LF_{q_j}^R(a, b, c, \alpha, q = \alpha(\alpha + 1)a - 4(2a - b - c)(q_j + 2j)^2; z = x - a \\
 &\quad, \rho = -\frac{2a - b - c}{(a - b)(a - c)}z, \eta = \frac{-z^2}{(a - b)(a - c)}) \\
 &= \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \rho^{i_0} \\
 &+ \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-q_0)_{i_0} (q_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-q_1)_{i_0} (q_1 + 4)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \rho^{i_1} \right\} \eta \\
 &+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-q_0)_{i_0} (q_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
 &\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \frac{\alpha}{2} + \frac{1}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(-q_k)_{i_k} (q_k + 4k)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
 &\times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \rho^{i_n} \right\} \eta^n \quad (8.2.4)
 \end{aligned}$$

Remark 8.2.2 *The power series expansion of Lamé equation in the algebraic form of the second kind for polynomial of type 2 about $x = a$ as*

$q = \alpha(\alpha + 1)a - 4(2a - b - c) \left(q_j + 2j + \frac{1}{2}\right)^2$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(z) &= LS_{q_j}^R \left(a, b, c, \alpha, q = \alpha(\alpha + 1)a - 4(2a - b - c) \left(q_j + 2j + \frac{1}{2}\right)^2 ; z = x - a \right. \\
&\quad \left. , \rho = -\frac{2a - b - c}{(a - b)(a - c)}z, \eta = \frac{-z^2}{(a - b)(a - c)} \right) \\
&= z^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 1)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \rho^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + 1) (i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2}) (i_0 + 2)} \frac{(-q_0)_{i_0} (q_0 + 1)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 5)_{i_1}}{(-q_1)_{i_0} (q_1 + 5)_{i_0}} \frac{\left(\frac{7}{2}\right)_{i_0} (3)_{i_0}}{\left(\frac{7}{2}\right)_{i_1} (3)_{i_1}} \rho^{i_1} \right\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + 1) (i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2}) (i_0 + 2)} \frac{(-q_0)_{i_0} (q_0 + 1)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k + 2k + \frac{\alpha}{2} + 1) (i_k + 2k - \frac{\alpha}{2} + \frac{1}{2})}{(i_k + 2k + \frac{5}{2}) (i_k + 2k + 2)} \right. \\
&\quad \times \frac{(-q_k)_{i_k} (q_k + 4k + 1)_{i_k} (2k + \frac{3}{2})_{i_{k-1}} (2k + 1)_{i_{k-1}}}{(-q_k)_{i_{k-1}} (q_k + 4k + 1)_{i_{k-1}} (2k + \frac{3}{2})_{i_k} (2k + 1)_{i_k}} \left. \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} (q_n + 4n + 1)_{i_n} (2n + \frac{3}{2})_{i_{n-1}} (2n + 1)_{i_{n-1}}}{(-q_n)_{i_{n-1}} (q_n + 4n + 1)_{i_{n-1}} (2n + \frac{3}{2})_{i_n} (2n + 1)_{i_n}} \rho^{i_n} \right\} \eta^n \Bigg\} \quad (8.2.5)
\end{aligned}$$

In Ref.[12] I treat α as a fixed value and q as a free variable to construct the Lamé polynomial of type 1: (1) if $\alpha = 2(2\alpha_j + j)$ or $-2(2\alpha_j + j) - 1$ where $j, \alpha_j \in \mathbb{N}_0$, an analytic solution of Lamé equation turns to be the first kind of independent solution of the Lamé polynomial of type 1. (2) if $\alpha = 2(2\alpha_j + j) + 1$ or $-2(2\alpha_j + j) + 1$, an analytic solution of Lamé equation turns to be the second kind of independent solution of the Lamé polynomial of type 1.

In this chapter I treat q as a fixed value and α as a free variable to construct the Lamé polynomial of type 2: (1) if $q = \alpha(\alpha + 1)a - 4(2a - b - c)(q_j + 2j)^2$ where $j, q_j \in \mathbb{N}_0$, an analytic solution of Lamé equation turns to be the first kind of independent solution of the Lamé polynomial of type 2. (2) if $q = \alpha(\alpha + 1)a - 4(2a - b - c) \left(q_j + 2j + \frac{1}{2}\right)^2$, an analytic solution of Lamé equation turns to be the second kind of independent solution of the Lamé polynomial of type 2.

8.2.2 Infinite series

In chapter 1 the general expression of power series of $y(x)$ for infinite series is defined by

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 \left\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} A_{i_1} \right) x^{i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} A_{i_3+2} \right) \right\} x^{i_2+2+\lambda} \right. \\
 &\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ B_{i_0+1} \prod_{i_1=0}^{i_0-1} A_{i_1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} B_{i_{2k}+2k+1} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} A_{i_{2k+1}+2k} \right) \right. \right. \\
 &\quad \times \left. \left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} A_{i_{2N+1}+2N} \right) \right\} \right\} x^{i_{2N}+2N+\lambda} \Bigg\} \quad (8.2.6)
 \end{aligned}$$

Substitute (8.1.7a)–(8.1.7c) into (8.2.6) with replacing variable x by z . The general expression of power series of Lamé equation in the algebraic form for infinite series about

$x = a$ is given by

$$\begin{aligned}
y(z) &= \sum_{n=0}^{\infty} y_n(z) = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \\
&= c_0 z^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \rho^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(-\varphi + 2 + \lambda)_{i_1} (\varphi + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} \left(\frac{5}{2} + \lambda\right)_{i_0}}{(-\varphi + 2 + \lambda)_{i_0} (\varphi + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} \left(\frac{5}{2} + \lambda\right)_{i_1}} \rho^{i_1} \Big\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_k + 2k - \frac{\alpha}{2} + \lambda)}{(i_k + 2k + 2 + \lambda) (i_k + 2k + \frac{3}{2} + \lambda)} \right. \\
&\quad \times \frac{(-\varphi + 2k + \lambda)_{i_k} (\varphi + 2k + \lambda)_{i_k} (2k + 1 + \lambda)_{i_{k-1}} (2k + \frac{1}{2} + \lambda)_{i_{k-1}}}{(-\varphi + 2k + \lambda)_{i_{k-1}} (\varphi + 2k + \lambda)_{i_{k-1}} (2k + 1 + \lambda)_{i_k} (2k + \frac{1}{2} + \lambda)_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(-\varphi + 2n + \lambda)_{i_n} (\varphi + 2n + \lambda)_{i_n} (2n + 1 + \lambda)_{i_{n-1}} (2n + \frac{1}{2} + \lambda)_{i_{n-1}}}{(-\varphi + 2n + \lambda)_{i_{n-1}} (\varphi + 2n + \lambda)_{i_{n-1}} (2n + 1 + \lambda)_{i_n} (2n + \frac{1}{2} + \lambda)_{i_n}} \rho^{i_n} \Big\} \eta^n \Big\} \quad (8.2.7)
\end{aligned}$$

Put $c_0 = 1$ as $\lambda = 0$ for the first kind of independent solutions of Lamé equation and $\lambda = \frac{1}{2}$ for the second one in (8.2.7).

Remark 8.2.3 *The power series expansion of Lamé equation in the algebraic form of the*

first kind for infinite series about $x = a$ using R3TRF is

$$\begin{aligned}
y(z) &= LFR \left(a, b, c, \alpha, q, \varphi = \sqrt{\frac{\alpha(\alpha+1)a-q}{4(2a-b-c)}}; z = x - a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z, \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
&= \sum_{i_0=0}^{\infty} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \rho^{i_0} \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(-\varphi+2)_{i_1} (\varphi+2)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-\varphi+2)_{i_0} (\varphi+2)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \rho^{i_1} \right\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-\varphi)_{i_0} (\varphi)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{\alpha}{2} + \frac{1}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(-\varphi+2k)_{i_k} (\varphi+2k)_{i_k} (2k+1)_{i_{k-1}} (2k+\frac{1}{2})_{i_{k-1}}}{(-\varphi+2k)_{i_{k-1}} (\varphi+2k)_{i_{k-1}} (2k+1)_{i_k} (2k+\frac{1}{2})_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(-\varphi+2n)_{i_n} (\varphi+2n)_{i_n} (2n+1)_{i_{n-1}} (2n+\frac{1}{2})_{i_{n-1}}}{(-\varphi+2n)_{i_{n-1}} (\varphi+2n)_{i_{n-1}} (2n+1)_{i_n} (2n+\frac{1}{2})_{i_n}} \rho^{i_n} \right\} \eta^n \tag{8.2.8}
\end{aligned}$$

Remark 8.2.4 The power series expansion of Lamé equation in the algebraic form of the

second kind for infinite series about $x = a$ using R3TRF is

$$\begin{aligned}
y(z) &= LS^R \left(a, b, c, \alpha, q, \varphi = \sqrt{\frac{\alpha(\alpha+1)a-q}{4(2a-b-c)}}; z = x - a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z, \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
&= z^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{(-\varphi + \frac{1}{2})_{i_0} (\varphi + \frac{1}{2})_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \rho^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + 1) (i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2}) (i_0 + 2)} \frac{(-\varphi + \frac{1}{2})_{i_0} (\varphi + \frac{1}{2})_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{\infty} \frac{(-\varphi + \frac{5}{2})_{i_1} (\varphi + \frac{5}{2})_{i_1} (\frac{7}{2})_{i_0} (3)_{i_0}}{(-\varphi + \frac{5}{2})_{i_0} (\varphi + \frac{5}{2})_{i_0} (\frac{7}{2})_{i_1} (3)_{i_1}} \rho^{i_1} \Big\} \eta \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + 1) (i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2}) (i_0 + 2)} \frac{(-\varphi + \frac{1}{2})_{i_0} (\varphi + \frac{1}{2})_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{\alpha}{2} + 1) (i_k + 2k - \frac{\alpha}{2} + \frac{1}{2})}{(i_k + 2k + \frac{5}{2}) (i_k + 2k + 2)} \right. \\
&\quad \times \frac{(-\varphi + 2k + \frac{1}{2})_{i_k} (\varphi + 2k + \frac{1}{2})_{i_k} (2k + \frac{3}{2})_{i_{k-1}} (2k + 1)_{i_{k-1}}}{(-\varphi + 2k + \frac{1}{2})_{i_{k-1}} (\varphi + 2k + \frac{1}{2})_{i_{k-1}} (2k + \frac{3}{2})_{i_k} (2k + 1)_{i_k}} \Big\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(-\varphi + 2n + \frac{1}{2})_{i_n} (\varphi + 2n + \frac{1}{2})_{i_n} (2n + \frac{3}{2})_{i_{n-1}} (2n + 1)_{i_{n-1}}}{(-\varphi + 2n + \frac{1}{2})_{i_{n-1}} (\varphi + 2n + \frac{1}{2})_{i_{n-1}} (2n + \frac{3}{2})_{i_n} (2n + 1)_{i_n}} \rho^{i_n} \right\} \eta^n \Big\} \quad (8.2.9)
\end{aligned}$$

The infinite series in this chapter are equivalent to the infinite series in Ref.[12]. In this chapter B_n is the leading term in sequence c_n of the analytic function $y(z)$. In Ref.[12] A_n is the leading term in sequence c_n of the analytic function $y(z)$.

8.3 Integral formalism

Lame equation could not be built in a definite or contour integral form of any well-known simple functions such as Gauss hypergeometric, Kummer functions and etc because of three different coefficients in its power series. The three term recurrence relation in the power series expansion of Lamé equation brings mathematical difficulty to be derived into a direct or contour integral.

In place of analyzing Lamé equation into its integral representation of any simple functions, in earlier literature the integral equations of Lamé equation were constructed

by using simple kernels involving Legendre functions of the Jacobian elliptic function or ellipsoidal harmonic functions: such integral relationships express one analytic solution in terms of another analytic solution involving Jacobian elliptic functions. There are many other forms of integral relations in Lamé equation.[17, 18, 19, 20, 21, 22, 23, 24, 25]

In Ref.[12], I show the integral form (each sub-integral $y_m(x)$ where $m = 0, 1, 2, \dots$ is composed of $2m$ terms of definite integrals and m terms of contour integrals) of Lamé equation using 3TRF: a ${}_2F_1$ function recurs in each of sub-integral forms of Lamé equation.

8.3.1 Polynomial of type 2

Now I consider the combined definite and contour integral representation of Lamé equation by using R3TRF. There is a generalized hypergeometric function which is given by

$$\begin{aligned} I_l &= \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} (q_l + 4l + 2\lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \frac{1}{2} + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} (q_l + 4l + 2\lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \frac{1}{2} + \lambda)_{i_l}} \rho^{i_l} \\ &= \rho^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1} + 2l + \lambda, j + 1) B(i_{l-1} + 2l - \frac{1}{2} + \lambda, j + 1) (i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + 2\lambda)_j}{(i_{l-1} + 2l + \lambda)^{-1} (i_{l-1} + 2l - \frac{1}{2} + \lambda)^{-1} (1)_j j!} \rho^j \end{aligned} \quad (8.3.1)$$

By using integral form of beta function,

$$B(i_{l-1} + 2l + \lambda, j + 1) = \int_0^1 dt_l t_l^{i_{l-1} + 2l - 1 + \lambda} (1 - t_l)^j \quad (8.3.2a)$$

$$B(i_{l-1} + 2l - \frac{1}{2} + \lambda, j + 1) = \int_0^1 du_l u_l^{i_{l-1} + 2l - \frac{3}{2} + \lambda} (1 - u_l)^j \quad (8.3.2b)$$

Substitute (8.3.2a) and (8.3.2b) into (8.3.1). And divide $(i_{l-1} + 2l + \lambda) (i_{l-1} + 2l - \frac{1}{2} + \lambda)$ into I_l .

$$\begin{aligned} K_l &= \frac{1}{(i_{l-1} + 2l + \lambda) (i_{l-1} + 2l - \frac{1}{2} + \lambda)} \\ &\times \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} (q_l + 4l + 2\lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \frac{1}{2} + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} (q_l + 4l + 2\lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \frac{1}{2} + \lambda)_{i_l}} \rho^{i_l} \\ &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\lambda} (\rho t_l u_l)^{i_{l-1}} \sum_{j=0}^{\infty} \frac{(i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + 2\lambda)_j}{(1)_j j!} (\rho(1 - t_l)(1 - u_l))^j \end{aligned}$$

The integral form of Gauss hypergeometric function is written by

$$\begin{aligned}
 {}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n (n!)} z^n \\
 &= -\frac{1}{2\pi i} \frac{\Gamma(1-\alpha)\Gamma(\gamma)}{\Gamma(\gamma-\alpha)} \oint dv_l (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \quad (8.3.3) \\
 &\text{where } \operatorname{Re}(\gamma-\alpha) > 0
 \end{aligned}$$

replaced α, β, γ and z by $i_{l-1} - q_l, i_{l-1} + q_l + 4l + 2\lambda, 1$ and $\rho(1-t_l)(1-u_l)$ in (8.3.3)

$$\begin{aligned}
 &\sum_{j=0}^{\infty} \frac{(i_{l-1} - q_l)_j (i_{l-1} + q_l + 4l + 2\lambda)_j}{(1)_j j!} (\rho(1-t_l)(1-u_l))^j \\
 &= \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \frac{1}{1 - \rho(1-t_l)(1-u_l)v_l} \right)^{q_l} (1 - \rho(1-t_l)(1-u_l)v_l)^{-(4l+2\lambda)} \\
 &\quad \times \left(\frac{v_l}{v_l - 1} \frac{1}{1 - \rho(1-t_l)(1-u_l)v_l} \right)^{i_{l-1}} \quad (8.3.4)
 \end{aligned}$$

Substitute (8.3.4) into K_l .

$$\begin{aligned}
 K_l &= \frac{1}{(i_{l-1} + 2l + \lambda) (i_{l-1} + 2l - \frac{1}{2} + \lambda)} \\
 &\quad \times \sum_{i_l=i_{l-1}}^{q_l} \frac{(-q_l)_{i_l} (q_l + 4l + 2\lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \frac{1}{2} + \lambda)_{i_{l-1}}}{(-q_l)_{i_{l-1}} (q_l + 4l + 2\lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \frac{1}{2} + \lambda)_{i_l}} \rho^{i_l} \\
 &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \frac{1}{1 - \rho(1-t_l)(1-u_l)v_l} \right)^{q_l} \\
 &\quad \times (1 - \rho(1-t_l)(1-u_l)v_l)^{-(4l+2\lambda)} \left(\frac{v_l}{v_l - 1} \frac{\rho t_l u_l}{1 - \rho(1-t_l)(1-u_l)v_l} \right)^{i_{l-1}} \quad (8.3.5)
 \end{aligned}$$

Substitute (8.3.5) into (8.2.3) where $l = 1, 2, 3, \dots$; apply K_1 into the second summation of sub-power series $y_1(z)$, apply K_2 into the third summation and K_1 into the second summation of sub-power series $y_2(z)$, apply K_3 into the forth summation, K_2 into the third summation and K_1 into the second summation of sub-power series $y_3(z)$, etc.⁴

Theorem 8.3.1 *The general representation in the form of integral of the Lamé*

⁴ $y_1(z)$ means the sub-power series in (8.2.3) contains one term of $B'_n s$, $y_2(z)$ means the sub-power series in (8.2.3) contains two terms of $B'_n s$, $y_3(z)$ means the sub-power series in (8.2.3) contains three terms of $B'_n s$, etc.

polynomial of type 2 in the algebraic form is given by

$$\begin{aligned}
y(z) &= \sum_{n=0}^{\infty} y_n(z) = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \\
&= c_0 z^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \rho^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}+\lambda} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
&\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+2\lambda)} \\
&\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-2-\frac{\alpha}{2}+\lambda} \Big\} \\
&\quad \times \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \Big\} \eta^n \Big\} \quad (8.3.6)
\end{aligned}$$

where

$$\overleftrightarrow{w}_{i,j} = \begin{cases} \frac{v_i}{(v_i - 1)} \frac{\overleftrightarrow{w}_{i+1,j} t_i u_i}{1 - \overleftrightarrow{w}_{i+1,j}(1-t_i)(1-u_i)v_i} & \text{where } i \leq j \\ \rho & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of B'_n s, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof According to (8.2.3),

$$y(z) = \sum_{n=0}^{\infty} c_n z^{n+\lambda} = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \quad (8.3.7)$$

In the above, sub-power series $y_0(z)$, $y_1(z)$, $y_2(z)$ and $y_3(z)$ of the Lamé polynomial using R3TRF about $x = a$ are given by

$$y_0(z) = c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \rho^{i_0} \quad (8.3.8a)$$

$$y_1(z) = c_0 z^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda)(i_0 - \frac{\alpha}{2} + \lambda)(-q_0)_{i_0}(q_0 + 2\lambda)_{i_0}}{(i_0 + 2 + \lambda)(i_0 + \frac{3}{2} + \lambda)(1 + \lambda)_{i_0}(\frac{1}{2} + \lambda)_{i_0}} \right. \\ \left. \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1}(q_1 + 4 + 2\lambda)_{i_1}(3 + \lambda)_{i_0}(\frac{5}{2} + \lambda)_{i_0}}{(-q_1)_{i_0}(q_1 + 4 + 2\lambda)_{i_0}(3 + \lambda)_{i_1}(\frac{5}{2} + \lambda)_{i_1}} \rho^{i_1} \right\} \eta \quad (8.3.8b)$$

$$y_2(z) = c_0 z^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda)(i_0 - \frac{\alpha}{2} + \lambda)(-q_0)_{i_0}(q_0 + 2\lambda)_{i_0}}{(i_0 + 2 + \lambda)(i_0 + \frac{3}{2} + \lambda)(1 + \lambda)_{i_0}(\frac{1}{2} + \lambda)_{i_0}} \right. \\ \times \sum_{i_1=i_0}^{q_1} \frac{(i_1 + \frac{\alpha}{2} + \frac{5}{2} + \lambda)(i_1 - \frac{\alpha}{2} + 2 + \lambda)(-q_1)_{i_1}(q_1 + 4 + 2\lambda)_{i_1}(3 + \lambda)_{i_0}(\frac{5}{2} + \lambda)_{i_0}}{(i_1 + 4 + \lambda)(i_1 + \frac{7}{2} + \lambda)(-q_1)_{i_0}(q_1 + 4 + 2\lambda)_{i_0}(3 + \lambda)_{i_1}(\frac{5}{2} + \lambda)_{i_1}} \\ \times \sum_{i_2=i_1}^{q_2} \frac{(-q_2)_{i_2}(q_2 + 8 + 2\lambda)_{i_2}(5 + \lambda)_{i_1}(\frac{9}{2} + \lambda)_{i_1}}{(-q_2)_{i_1}(q_2 + 8 + 2\lambda)_{i_1}(5 + \lambda)_{i_2}(\frac{9}{2} + \lambda)_{i_2}} \rho^{i_2} \left. \right\} \eta^2 \quad (8.3.8c)$$

$$y_3(z) = c_0 z^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda)(i_0 - \frac{\alpha}{2} + \lambda)(-q_0)_{i_0}(q_0 + 2\lambda)_{i_0}}{(i_0 + 2 + \lambda)(i_0 + \frac{3}{2} + \lambda)(1 + \lambda)_{i_0}(\frac{1}{2} + \lambda)_{i_0}} \right. \\ \times \sum_{i_1=i_0}^{q_1} \frac{(i_1 + \frac{\alpha}{2} + \frac{5}{2} + \lambda)(i_1 - \frac{\alpha}{2} + 2 + \lambda)(-q_1)_{i_1}(q_1 + 4 + 2\lambda)_{i_1}(3 + \lambda)_{i_0}(\frac{5}{2} + \lambda)_{i_0}}{(i_1 + 4 + \lambda)(i_1 + \frac{7}{2} + \lambda)(-q_1)_{i_0}(q_1 + 4 + 2\lambda)_{i_0}(3 + \lambda)_{i_1}(\frac{5}{2} + \lambda)_{i_1}} \\ \times \sum_{i_2=i_1}^{q_2} \frac{(i_2 + \frac{\alpha}{2} + \frac{9}{2} + \lambda)(i_2 - \frac{\alpha}{2} + 4 + \lambda)(-q_2)_{i_2}(q_2 + 8 + 2\lambda)_{i_2}(5 + \lambda)_{i_1}(\frac{9}{2} + \lambda)_{i_1}}{(i_2 + 6 + \lambda)(i_2 + \frac{11}{2} + \lambda)(-q_2)_{i_1}(q_2 + 8 + 2\lambda)_{i_1}(5 + \lambda)_{i_2}(\frac{9}{2} + \lambda)_{i_2}} \\ \times \sum_{i_3=i_2}^{q_3} \frac{(-q_3)_{i_3}(q_3 + 12 + 2\lambda)_{i_3}(7 + \lambda)_{i_2}(\frac{13}{2} + \lambda)_{i_2}}{(-q_3)_{i_2}(q_3 + 12 + 2\lambda)_{i_2}(7 + \lambda)_{i_3}(\frac{13}{2} + \lambda)_{i_3}} \rho^{i_3} \left. \right\} \eta^3 \quad (8.3.8d)$$

Put $l = 1$ in (8.3.5). Take the new (8.3.5) into (8.3.8b).

$$y_1(z) = \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - \rho(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \\ \times (1 - \rho(1 - t_1)(1 - u_1)v_1)^{-(4+2\lambda)} \overleftrightarrow{w}_{1,1}^{-(\frac{1}{2} + \frac{\alpha}{2} + \lambda)} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{\alpha + \frac{1}{2}} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{-\frac{\alpha}{2} + \lambda} \\ \times \left\{ c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0}(q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0}(\frac{1}{2} + \lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right\} \eta \quad (8.3.9)$$

where

$$\overleftrightarrow{w}_{1,1} = \frac{v_1}{v_1 - 1} \frac{\rho t_1 u_1}{1 - \rho(1 - t_1)(1 - u_1)v_1}$$

Put $l = 2$ in (8.3.5). Take the new (8.3.5) into (8.3.8c).

$$\begin{aligned}
y_2(z) = & c_0 z^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - \rho(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\
& \times (1 - \rho(1 - t_2)(1 - u_2)v_2)^{-(8+2\lambda)} \overleftarrow{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
& \times \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda) (-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda) (1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \right. \\
& \times \left. \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} (q_1 + 4 + 2\lambda)_{i_1} (3 + \lambda)_{i_0} (\frac{5}{2} + \lambda)_{i_0} \overleftarrow{w}_{2,2}^{i_1}}{(-q_1)_{i_0} (q_1 + 4 + 2\lambda)_{i_0} (3 + \lambda)_{i_1} (\frac{5}{2} + \lambda)_{i_1}} \right\} \eta^2 \quad (8.3.10)
\end{aligned}$$

where

$$\overleftarrow{w}_{2,2} = \frac{v_2}{v_2 - 1} \frac{\rho t_2 u_2}{1 - \rho(1 - t_2)(1 - u_2)v_2}$$

Put $l = 1$ and $\rho = \overleftarrow{w}_{2,2}$ in (8.3.5). Take the new (8.3.5) into (8.3.10).

$$\begin{aligned}
y_2(z) = & c_0 z^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \frac{1}{1 - \rho(1 - t_2)(1 - u_2)v_2} \right)^{q_2} \\
& \times (1 - \rho(1 - t_2)(1 - u_2)v_2)^{-(8+2\lambda)} \overleftarrow{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{2,2} \partial_{\overleftarrow{w}_{2,2}} \right) \overleftarrow{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \frac{1}{1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1} \right)^{q_1} \\
& \times (1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1)^{-(4+2\lambda)} \overleftarrow{w}_{1,2}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftarrow{w}_{1,2} \partial_{\overleftarrow{w}_{1,2}} \right) \overleftarrow{w}_{1,2}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{1,2} \partial_{\overleftarrow{w}_{1,2}} \right) \overleftarrow{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \\
& \times \left\{ c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0} \overleftarrow{w}_{1,2}^{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \right\} \eta^2 \quad (8.3.11)
\end{aligned}$$

where

$$\overleftarrow{w}_{1,2} = \frac{v_1}{v_1 - 1} \frac{\overleftarrow{w}_{2,2} t_1 u_1}{1 - \overleftarrow{w}_{2,2}(1 - t_1)(1 - u_1)v_1}$$

By using similar process for the previous cases of integral forms of $y_1(z)$ and $y_2(z)$, the

integral form of sub-power series expansion of $y_3(z)$ is given by

$$\begin{aligned}
y_3(z) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{\frac{9}{2}+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3-1}{v_3} \frac{1}{1-\rho(1-t_3)(1-u_3)v_3} \right)^{q_3} \\
& \times (1-\rho(1-t_3)(1-u_3)v_3)^{-(12+2\lambda)} \overleftarrow{w}_{3,3}^{-\left(\frac{9}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftarrow{w}_{3,3} \partial_{\overleftarrow{w}_{3,3}} \right) \overleftarrow{w}_{3,3}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{3,3} \partial_{\overleftarrow{w}_{3,3}} \right) \overleftarrow{w}_{3,3}^{4-\frac{\alpha}{2}+\lambda} \\
& \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \frac{1}{1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \right)^{q_2} \\
& \times (1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2)^{-(8+2\lambda)} \overleftarrow{w}_{2,3}^{-\left(\frac{5}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftarrow{w}_{2,3} \partial_{\overleftarrow{w}_{2,3}} \right) \overleftarrow{w}_{2,3}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{2,3} \partial_{\overleftarrow{w}_{2,3}} \right) \overleftarrow{w}_{2,3}^{2-\frac{\alpha}{2}+\lambda} \\
& \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \frac{1}{1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
& \times (1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1)^{-(4+2\lambda)} \overleftarrow{w}_{1,3}^{-\left(\frac{1}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftarrow{w}_{1,3} \partial_{\overleftarrow{w}_{1,3}} \right) \overleftarrow{w}_{1,3}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{1,3} \partial_{\overleftarrow{w}_{1,3}} \right) \overleftarrow{w}_{1,3}^{-\frac{\alpha}{2}+\lambda} \\
& \times \left\{ c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} \left(\frac{1}{2}+\lambda\right)_{i_0}} \overleftarrow{w}_{1,3}^{i_0} \right\} \eta^3 \tag{8.3.12}
\end{aligned}$$

where

$$\begin{cases} \overleftarrow{w}_{3,3} = \frac{v_3}{v_3-1} \frac{\rho t_3 u_3}{1-\rho(1-t_3)(1-u_3)v_3} \\ \overleftarrow{w}_{2,3} = \frac{v_2}{v_2-1} \frac{\overleftarrow{w}_{3,3} t_2 u_2}{1-\overleftarrow{w}_{3,3}(1-t_2)(1-u_2)v_2} \\ \overleftarrow{w}_{1,3} = \frac{v_1}{v_1-1} \frac{\overleftarrow{w}_{2,3} t_1 u_1}{1-\overleftarrow{w}_{2,3}(1-t_1)(1-u_1)v_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute (8.3.8a), (8.3.9), (8.3.11), (8.3.12) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (8.3.7), we obtain (8.3.6).

Put $c_0=1$ as $\lambda=0$ for the first kind of independent solutions of Lamé equation and $\lambda=\frac{1}{2}$ for the second one in (8.3.6).

Remark 8.3.2 The integral representation of Lamé equation in the algebraic form of the first kind for polynomial of type 2 about $x=a$ as $q=\alpha(\alpha+1)a-4(2a-b-c)(q_j+2j)^2$

where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
 y(z) &= LF_{q_j}^R \left(a, b, c, \alpha, q = \alpha(\alpha+1)a - 4(2a-b-c)(q_j+2j)^2; z = x-a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z \right. \\
 &\quad \left. , \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
 &= {}_2F_1 \left(-q_0, q_0; \frac{1}{2}; \rho \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
 &\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-4(n-k)} \\
 &\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-2-\frac{\alpha}{2}} \left. \right\} \\
 &\quad \times {}_2F_1 \left(-q_0, q_0; \frac{1}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} \eta^n \tag{8.3.13}
 \end{aligned}$$

Remark 8.3.3 The integral representation of Lamé equation in the algebraic form of the second kind for polynomial of type 2 about $x = a$ as $q = \alpha(\alpha+1)a - 4(2a-b-c)(q_j+2j+\frac{1}{2})^2$ where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
 y(z) &= LS_{q_j}^R \left(a, b, c, \alpha, q = \alpha(\alpha+1)a - 4(2a-b-c) \left(q_j + 2j + \frac{1}{2} \right)^2; z = x-a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z \right. \\
 &\quad \left. , \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
 &= z^{\frac{1}{2}} \left\{ {}_2F_1 \left(-q_0, q_0 + 1; \frac{3}{2}; \rho \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
 &\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+1)} \\
 &\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \left. \right\} \\
 &\quad \times {}_2F_1 \left(-q_0, q_0 + 1; \frac{3}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} \eta^n \tag{8.3.14}
 \end{aligned}$$

8.3.2 Infinite series

Let's consider the integral representation of Lamé equation about $x = 0$ for infinite series by applying R3TRF. There is a generalized hypergeometric function which is written by

$$\begin{aligned}
 M_l &= \sum_{i_l=i_{l-1}}^{\infty} \frac{(-\varphi+2l+\lambda)_{i_l} (\varphi+2l+\lambda)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\frac{1}{2}+\lambda)_{i_{l-1}}}{(-\varphi+2l+\lambda)_{i_{l-1}} (\varphi+2l+\lambda)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\frac{1}{2}+\lambda)_{i_l}} \rho^{i_l} \\
 &= \rho^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+2l+\lambda, j+1) B(i_{l-1}+2l-\frac{1}{2}+\lambda, j+1)}{(i_{l-1}+2l+\lambda)^{-1} (i_{l-1}+2l-\frac{1}{2}+\lambda)^{-1}} \\
 &\quad \times \frac{(-\varphi+2l+\lambda+i_{l-1})_j (\varphi+2l+\lambda+i_{l-1})_j}{(1)_j j!} \rho^j
 \end{aligned} \tag{8.3.15}$$

Substitute (8.3.2a) and (8.3.2b) into (8.3.15). And divide $(i_{l-1}+2l+\lambda) (i_{l-1}+2l-\frac{1}{2}+\lambda)$ into the new (8.3.15).

$$\begin{aligned}
 V_l &= \frac{1}{(i_{l-1}+2l+\lambda) (i_{l-1}+2l-\frac{1}{2}+\lambda)} \\
 &\quad \times \sum_{i_l=i_{l-1}}^{\infty} \frac{(-\varphi+2l+\lambda)_{i_l} (\varphi+2l+\lambda)_{i_l} (2l+1+\lambda)_{i_{l-1}} (2l+\frac{1}{2}+\lambda)_{i_{l-1}}}{(-\varphi+2l+\lambda)_{i_{l-1}} (\varphi+2l+\lambda)_{i_{l-1}} (2l+1+\lambda)_{i_l} (2l+\frac{1}{2}+\lambda)_{i_l}} \rho^{i_l} \\
 &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\lambda} (\rho t_l u_l)^{i_{l-1}} \\
 &\quad \times \sum_{j=0}^{\infty} \frac{(-\varphi+2l+\lambda+i_{l-1})_j (\varphi+2l+\lambda+i_{l-1})_j}{(1)_j j!} (\rho(1-t_l)(1-u_l))^j
 \end{aligned}$$

The hypergeometric function is defined by

$$\begin{aligned}
 {}_2F_1(\alpha, \beta; \gamma; z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n (n!)} z^n \\
 &= \frac{1}{2\pi i} \frac{\Gamma(1+\alpha-\gamma)}{\Gamma(\alpha)} \int_0^{(1+)} dv_l (-1)^\gamma (-v_l)^{\alpha-1} (1-v_l)^{\gamma-\alpha-1} (1-zv_l)^{-\beta} \tag{8.3.16}
 \end{aligned}$$

where $\gamma - \alpha \neq 1, 2, 3, \dots$, $\text{Re}(\alpha) > 0$

Replace α, β, γ and z by $-\varphi + 2l + \lambda + i_{l-1}$, $\varphi + 2l + \lambda + i_{l-1}$, 1 and $\rho(1 - t_l)(1 - u_l)$ in (8.3.16). Take the new (8.3.16) into V_l .

$$\begin{aligned}
 V_l &= \frac{1}{(i_{l-1} + 2l + \lambda)(i_{l-1} + 2l - \frac{1}{2} + \lambda)} \\
 &\times \sum_{i_l=i_{l-1}}^{\infty} \frac{(-\varphi + 2l + \lambda)_{i_l} (\varphi + 2l + \lambda)_{i_l} (2l + 1 + \lambda)_{i_{l-1}} (2l + \frac{1}{2} + \lambda)_{i_{l-1}}}{(-\varphi + 2l + \lambda)_{i_{l-1}} (\varphi + 2l + \lambda)_{i_{l-1}} (2l + 1 + \lambda)_{i_l} (2l + \frac{1}{2} + \lambda)_{i_l}} \rho^{i_l} \\
 &= \int_0^1 dt_l t_l^{2l-1+\lambda} \int_0^1 du_l u_l^{2l-\frac{3}{2}+\lambda} \frac{1}{2\pi i} \oint dv_l \frac{1}{v_l} \left(\frac{v_l - 1}{v_l} \right)^{\varphi-(2l+\lambda)} \\
 &\times (1 - \rho(1 - t_l)(1 - u_l)v_l)^{-\varphi-(2l+\lambda)} \left(\frac{v_l}{v_l - 1} \frac{\rho t_l u_l}{1 - \rho(1 - t_l)(1 - u_l)v_l} \right)^{i_{l-1}} \quad (8.3.17)
 \end{aligned}$$

Substitute (8.3.17) into (8.2.7) where $l = 1, 2, 3, \dots$; apply V_1 into the second summation of sub-power series $y_1(z)$, apply K_2 into the third summation and V_1 into the second summation of sub-power series $y_2(z)$, apply V_3 into the forth summation, V_2 into the third summation and V_1 into the second summation of sub-power series $y_3(z)$, etc.⁵

Theorem 8.3.4 *The general expression of an integral form of Lamé equation in the algebraic form for infinite series about $x = 0$ using R3TRF is given by*

$$\begin{aligned}
 y(z) &= \sum_{n=0}^{\infty} y_n(z) = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \dots \\
 &= c_0 z^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \rho^{i_0} \right. \\
 &+ \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}+\lambda} \right. \right. \\
 &\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \right)^{\varphi-2(n-k)-\lambda} \\
 &\times (1 - \overleftrightarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-\varphi-2(n-k)-\lambda} \\
 &\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-2-\frac{\alpha}{2}+\lambda} \Big\} \\
 &\times \left. \sum_{i_0=0}^{\infty} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \right\} \eta^n \Big\} \quad (8.3.18)
 \end{aligned}$$

⁵ $y_1(z)$ means the sub-power series in (8.2.7) contains one term of $B'_n s$, $y_2(z)$ means the sub-power series in (8.2.7) contains two terms of $B'_n s$, $y_3(z)$ means the sub-power series in (8.2.7) contains three terms of $B'_n s$, etc.

In the above, the first sub-integral form contains one term of B'_n 's, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Proof In (8.2.7) sub-power series $y_0(z)$, $y_1(z)$, $y_2(z)$ and $y_3(z)$ of the Lamé equation for infinite series about $x = a$ using R3TRF are given by

$$y_0(z) = c_0 z^\lambda \sum_{i_0=0}^{\infty} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \rho^{i_0} \quad (8.3.19a)$$

$$y_1(z) = c_0 z^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \right. \\ \left. \times \sum_{i_1=i_0}^{\infty} \frac{(-\varphi + 2 + \lambda)_{i_1} (\varphi + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} \left(\frac{5}{2} + \lambda\right)_{i_0}}{(-\varphi + 2 + \lambda)_{i_0} (\varphi + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} \left(\frac{5}{2} + \lambda\right)_{i_1}} \rho^{i_1} \right\} \eta \quad (8.3.19b)$$

$$y_2(z) = c_0 z^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \right. \\ \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + \frac{\alpha}{2} + \frac{5}{2} + \lambda) (i_1 - \frac{\alpha}{2} + 2 + \lambda)}{(i_1 + 4 + \lambda) (i_1 + \frac{7}{2} + \lambda)} \frac{(-\varphi + 2 + \lambda)_{i_1} (\varphi + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} \left(\frac{5}{2} + \lambda\right)_{i_0}}{(-\varphi + 2 + \lambda)_{i_0} (\varphi + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} \left(\frac{5}{2} + \lambda\right)_{i_1}} \\ \left. \times \sum_{i_2=i_1}^{\infty} \frac{(-\varphi + 4 + \lambda)_{i_2} (\varphi + 4 + \lambda)_{i_2} (5 + \lambda)_{i_1} \left(\frac{9}{2} + \lambda\right)_{i_1}}{(-\varphi + 4 + \lambda)_{i_1} (\varphi + 4 + \lambda)_{i_1} (5 + \lambda)_{i_2} \left(\frac{9}{2} + \lambda\right)_{i_2}} \rho^{i_2} \right\} \eta^2 \quad (8.3.19c)$$

$$y_3(z) = c_0 z^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} \left(\frac{1}{2} + \lambda\right)_{i_0}} \right. \\ \times \sum_{i_1=i_0}^{\infty} \frac{(i_1 + \frac{\alpha}{2} + \frac{5}{2} + \lambda) (i_1 - \frac{\alpha}{2} + 2 + \lambda)}{(i_1 + 4 + \lambda) (i_1 + \frac{7}{2} + \lambda)} \frac{(-\varphi + 2 + \lambda)_{i_1} (\varphi + 2 + \lambda)_{i_1} (3 + \lambda)_{i_0} \left(\frac{5}{2} + \lambda\right)_{i_0}}{(-\varphi + 2 + \lambda)_{i_0} (\varphi + 2 + \lambda)_{i_0} (3 + \lambda)_{i_1} \left(\frac{5}{2} + \lambda\right)_{i_1}} \\ \times \sum_{i_2=i_1}^{\infty} \frac{(i_2 + \frac{\alpha}{2} + \frac{9}{2} + \lambda) (i_2 - \frac{\alpha}{2} + 4 + \lambda)}{(i_2 + 6 + \lambda) (i_2 + \frac{11}{2} + \lambda)} \frac{(-\varphi + 4 + \lambda)_{i_2} (\varphi + 4 + \lambda)_{i_2} (5 + \lambda)_{i_1} \left(\frac{9}{2} + \lambda\right)_{i_1}}{(-\varphi + 4 + \lambda)_{i_1} (\varphi + 4 + \lambda)_{i_1} (5 + \lambda)_{i_2} \left(\frac{9}{2} + \lambda\right)_{i_2}} \\ \left. \times \sum_{i_3=i_2}^{\infty} \frac{(-\varphi + 6 + \lambda)_{i_3} (\varphi + 6 + \lambda)_{i_3} (7 + \lambda)_{i_2} \left(\frac{13}{2} + \lambda\right)_{i_2}}{(-\varphi + 6 + \lambda)_{i_2} (\varphi + 6 + \lambda)_{i_2} (7 + \lambda)_{i_3} \left(\frac{13}{2} + \lambda\right)_{i_3}} \rho^{i_3} \right\} \eta^3 \quad (8.3.19d)$$

Put $l = 1$ in (8.3.17). Take the new (8.3.17) into (8.3.19b).

$$\begin{aligned}
y_1(z) &= \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \right)^{\varphi-(2+\lambda)} \\
&\quad \times (1 - \rho(1-t_1)(1-u_1)v_1)^{-\varphi-(2+\lambda)} \overleftrightarrow{w}_{1,1}^{-\left(\frac{1}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{1,1} \partial_{\overleftrightarrow{w}_{1,1}} \right) \overleftrightarrow{w}_{1,1}^{-\frac{\alpha}{2}+\lambda} \\
&\quad \times \left\{ c_0 z^\lambda \sum_{i_0=0}^{\infty} \frac{(-\varphi+\lambda)_{i_0} (\varphi+\lambda)_{i_0}}{(1+\lambda)_{i_0} \left(\frac{1}{2}+\lambda\right)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right\} \eta
\end{aligned} \tag{8.3.20}$$

where

$$\overleftrightarrow{w}_{1,1} = \frac{v_1}{v_1-1} \frac{\rho t_1 u_1}{1 - \rho(1-t_1)(1-u_1)v_1}$$

Put $l = 2$ in (8.3.17). Take the new (8.3.17) into (8.3.19c).

$$\begin{aligned}
y_2(z) &= c_0 z^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \right)^{\varphi-(4+\lambda)} \\
&\quad \times (1 - \rho(1-t_2)(1-u_2)v_2)^{-\varphi-(4+\lambda)} \overleftrightarrow{w}_{2,2}^{-\left(\frac{5}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\quad \times \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2} + \lambda) (i_0 - \frac{\alpha}{2} + \lambda)}{(i_0 + 2 + \lambda) (i_0 + \frac{3}{2} + \lambda)} \frac{(-\varphi+\lambda)_{i_0} (\varphi+\lambda)_{i_0}}{(1+\lambda)_{i_0} \left(\frac{1}{2}+\lambda\right)_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\infty} \frac{(-\varphi+2+\lambda)_{i_1} (\varphi+2+\lambda)_{i_1} (3+\lambda)_{i_0} \left(\frac{5}{2}+\lambda\right)_{i_0}}{(-\varphi+2+\lambda)_{i_0} (\varphi+2+\lambda)_{i_0} (3+\lambda)_{i_1} \left(\frac{5}{2}+\lambda\right)_{i_1}} \overleftrightarrow{w}_{2,2}^{i_1} \right\} \eta^2
\end{aligned} \tag{8.3.21}$$

where

$$\overleftrightarrow{w}_{2,2} = \frac{v_2}{v_2-1} \frac{\rho t_2 u_2}{1 - \rho(1-t_2)(1-u_2)v_2}$$

Put $l = 1$ and $\rho = \overleftrightarrow{w}_{2,2}$ in (8.3.17). Take the new (8.3.17) into (8.3.21).

$$\begin{aligned}
y_2(z) &= c_0 z^\lambda \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2-1}{v_2} \right)^{\varphi-(4+\lambda)} \\
&\quad \times (1 - \rho(1-t_2)(1-u_2)v_2)^{-\varphi-(4+\lambda)} \overleftrightarrow{w}_{2,2}^{-\left(\frac{5}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1-1}{v_1} \right)^{\varphi-(2+\lambda)} \\
&\quad \times (1 - \overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-\varphi-(2+\lambda)} \overleftrightarrow{w}_{1,2}^{-\left(\frac{1}{2}+\frac{\alpha}{2}+\lambda\right)} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \\
&\quad \times \left\{ c_0 z^\lambda \sum_{i_0=0}^{\infty} \frac{(-\varphi+\lambda)_{i_0} (\varphi+\lambda)_{i_0}}{(1+\lambda)_{i_0} \left(\frac{1}{2}+\lambda\right)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right\} \eta^2
\end{aligned} \tag{8.3.22}$$

where

$$\overleftarrow{w}_{1,2} = \frac{v_1}{v_1 - 1} \frac{\overleftarrow{w}_{2,2} t_1 u_1}{1 - \overleftarrow{w}_{2,2} (1 - t_1) (1 - u_1) v_1}$$

By using similar process for the previous cases of integral forms of $y_1(z)$ and $y_2(z)$, the integral form of sub-power series expansion of $y_3(z)$ is given by

$$\begin{aligned} y_3(z) = & \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{\frac{9}{2}+\lambda} \frac{1}{2\pi i} \oint dv_3 \frac{1}{v_3} \left(\frac{v_3 - 1}{v_3} \right)^{\varphi - (6+\lambda)} \\ & \times (1 - \rho(1 - t_3)(1 - u_3)v_3)^{-\varphi - (6+\lambda)} \\ & \times \overleftarrow{w}_{3,3}^{-(\frac{9}{2} + \frac{\alpha}{2} + \lambda)} \left(\overleftarrow{w}_{3,3} \partial_{\overleftarrow{w}_{3,3}} \right) \overleftarrow{w}_{3,3}^{\alpha + \frac{1}{2}} \left(\overleftarrow{w}_{3,3} \partial_{\overleftarrow{w}_{3,3}} \right) \overleftarrow{w}_{3,3}^{4 - \frac{\alpha}{2} + \lambda} \\ & \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} \left(\frac{v_2 - 1}{v_2} \right)^{\varphi - (4+\lambda)} \\ & \times (1 - \overleftarrow{w}_{3,3}(1 - t_2)(1 - u_2)v_2)^{-\varphi - (4+\lambda)} \\ & \times \overleftarrow{w}_{2,3}^{-(\frac{5}{2} + \frac{\alpha}{2} + \lambda)} \left(\overleftarrow{w}_{2,3} \partial_{\overleftarrow{w}_{2,3}} \right) \overleftarrow{w}_{2,3}^{\alpha + \frac{1}{2}} \left(\overleftarrow{w}_{2,3} \partial_{\overleftarrow{w}_{2,3}} \right) \overleftarrow{w}_{2,3}^{2 - \frac{\alpha}{2} + \lambda} \\ & \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} \left(\frac{v_1 - 1}{v_1} \right)^{\varphi - (2+\lambda)} \\ & \times (1 - \overleftarrow{w}_{2,3}(1 - t_1)(1 - u_1)v_1)^{-\varphi - (2+\lambda)} \\ & \times \overleftarrow{w}_{1,3}^{-(\frac{1}{2} + \frac{\alpha}{2} + \lambda)} \left(\overleftarrow{w}_{1,3} \partial_{\overleftarrow{w}_{1,3}} \right) \overleftarrow{w}_{1,3}^{\alpha + \frac{1}{2}} \left(\overleftarrow{w}_{1,3} \partial_{\overleftarrow{w}_{1,3}} \right) \overleftarrow{w}_{1,3}^{-\frac{\alpha}{2} + \lambda} \\ & \times \left\{ c_0 z^\lambda \sum_{i_0=0}^{\infty} \frac{(-\varphi + \lambda)_{i_0} (\varphi + \lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \overleftarrow{w}_{1,3}^{i_0} \right\} \eta^3 \end{aligned} \quad (8.3.23)$$

where

$$\begin{cases} \overleftarrow{w}_{3,3} = \frac{v_3}{v_3 - 1} \frac{\rho t_3 u_3}{1 - \rho(1 - t_3)(1 - u_3)v_3} \\ \overleftarrow{w}_{2,3} = \frac{v_2}{v_2 - 1} \frac{\overleftarrow{w}_{3,3} t_2 u_2}{1 - \overleftarrow{w}_{3,3}(1 - t_2)(1 - u_2)v_2} \\ \overleftarrow{w}_{1,3} = \frac{v_1}{v_1 - 1} \frac{\overleftarrow{w}_{2,3} t_1 u_1}{1 - \overleftarrow{w}_{2,3}(1 - t_1)(1 - u_1)v_1} \end{cases}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(x)$ terms where $m \geq 4$, we obtain every integral forms of $y_m(x)$ terms. Since we substitute (8.3.19a), (8.3.20), (8.3.22), (8.3.23) and including all integral forms of $y_m(x)$ terms where $m \geq 4$ into (8.2.7), we obtain (8.3.18).⁶

Put $c_0 = 1$ as $\lambda = 0$ for the first independent solution of Lamé equation and $\lambda = \frac{1}{2}$ for the second one in (8.3.18).

⁶Or replace the finite summation with an interval $[0, q_0]$ by infinite summation with an interval $[0, \infty]$ in (8.3.6). Replace q_0 and q_{n-k} by $\varphi - \lambda$ and $\varphi - 2(n - k) - \lambda$ into the new (8.3.6). Its solution is also equivalent to (8.3.18)

Remark 8.3.5 *The integral representation of Lamé equation in the algebraic form of the first kind for infinite series about $x = a$ using R3TRF is*

$$\begin{aligned}
y(z) &= LFR \left(a, b, c, \alpha, q, \varphi = \sqrt{\frac{\alpha(\alpha+1)a-q}{4(2a-b-c)}}; z = x - a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z, \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
&= {}_2F_1 \left(-\varphi, \varphi; \frac{1}{2}; \rho \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\varphi-2(n-k)} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\varphi-2(n-k)} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-2-\frac{\alpha}{2}} \Big\} \\
&\times {}_2F_1 \left(-\varphi, \varphi; \frac{1}{2}; \overleftrightarrow{w}_{1,n} \right) \Big\} \eta^n
\end{aligned} \tag{8.3.24}$$

Remark 8.3.6 *The integral representation of Lamé equation in the algebraic form of the second kind for infinite series about $x = a$ using R3TRF is*

$$\begin{aligned}
y(z) &= LSR \left(a, b, c, \alpha, q, \varphi = \sqrt{\frac{\alpha(\alpha+1)a-q}{4(2a-b-c)}}; z = x - a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z, \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
&= z^{\frac{1}{2}} \left\{ {}_2F_1 \left(-\varphi + \frac{1}{2}, \varphi + \frac{1}{2}; \frac{3}{2}; \rho \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\varphi-2(n-k)-\frac{1}{2}} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\varphi-2(n-k)-\frac{1}{2}} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}+\frac{\alpha}{2}} \Big\} \\
&\times {}_2F_1 \left(-\varphi + \frac{1}{2}, \varphi + \frac{1}{2}; \frac{3}{2}; \overleftrightarrow{w}_{1,n} \right) \Big\} \eta^n
\end{aligned} \tag{8.3.25}$$

8.4 Generating function for the Lamé polynomial of type 2

I consider the generating function for the Lamé polynomial of type 2 in the algebraic form by applying R3TRF. Since the generating function for the Lamé polynomial is derived, we might be possible to construct orthogonal relations of it.⁷

⁷In the next series, I will construct the generating function for the type 3 Lamé polynomial. For the type 3 polynomial, I treat α and q as fixed values.

Lemma 8.4.1 *The generating function for the Jacobi polynomial using hypergeometric functions is given by*

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} w^{q_0} {}_2F_1(-q_0, q_0 + A; \gamma; x) \\ &= 2^{A-1} \frac{\left(1 - w + \sqrt{w^2 - 2(1-2x)w + 1}\right)^{1-\gamma} \left(1 + w + \sqrt{w^2 - 2(1-2x)w + 1}\right)^{\gamma-A}}{\sqrt{w^2 - 2(1-2x)w + 1}} \\ & \text{where } |w| < 1 \end{aligned} \quad (8.4.1)$$

Proof The proof of this lemma is given in Lemma 3.2.1

Definition 8.4.2 *I define that*

$$\left\{ \begin{aligned} s_{a,b} &= \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{if } a > b \\ s_a & \text{if } a = b \end{cases} \\ \tilde{w}_{i,j} &= \begin{cases} \frac{\tilde{w}_{i+1,j} t_i u_i \{1 + (s_i + 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i\}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} \\ - \frac{\tilde{w}_{i+1,j} t_i u_i (1 + s_i) \sqrt{s_i^2 - 2(1 - 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i + 1}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} & \text{where } i < j \\ \frac{\rho t_i u_i \{1 + (s_{i,\infty} + 2\rho(1-t_i)(1-u_i))s_{i,\infty}\}}{2(1 - \rho(1-t_i)(1-u_i))^2 s_{i,\infty}} \\ - \frac{\rho t_i u_i (1 + s_{i,\infty}) \sqrt{s_{i,\infty}^2 - 2(1 - 2\rho(1-t_i)(1-u_i))s_{i,\infty} + 1}}{2(1 - \rho(1-t_i)(1-u_i))^2 s_{i,\infty}} & \text{where } i = j \end{cases} \end{aligned} \right. \quad (8.4.2)$$

where

$$a, b, i, j \in \mathbb{N}_0$$

And we have

$$\sum_{q_i=q_j}^{\infty} r_i^{q_i} = \frac{r_i^{q_j}}{(1 - r_i)} \quad (8.4.3)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (8.3.6) where $|s_i| < 1$ as $i = 0, 1, 2, \dots$ by using (8.4.2) and (8.4.3),

Theorem 8.4.3 *The general expression of the generating function for the Lamé polynomial of type 2 in the algebraic form is given by*

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y(z) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \Upsilon(\lambda; s_{0,\infty}; \rho) \\
&+ \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} (s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+2\lambda)} \\
&\times \tilde{w}_{1,1}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{\alpha}{2}+\lambda} \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) \eta \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1+\lambda} \int_0^1 du_n u_n^{2n-\frac{3}{2}+\lambda} \right. \\
&\times (s_{n,\infty}^2 - 2(1-2\rho(1-t_n)(1-u_n))s_{n,\infty} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\rho(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+2\lambda)} \\
&\times \tilde{w}_{n,n}^{-(2n-\frac{3}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)-\frac{\alpha}{2}+\lambda} \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}+\lambda} \right. \\
&\times (s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1)^{-\frac{1}{2}} \\
&\times \left(\frac{1 + s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+2\lambda)} \\
&\times \tilde{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)-\frac{\alpha}{2}+\lambda} \left. \right\} \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) \left. \right\} \eta^n
\end{aligned} \tag{8.4.4}$$

where

$$\begin{cases} \Upsilon(\lambda; s_{0,\infty}; \rho) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_{0,\infty}^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \rho^{i_0} \right) \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,1}) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \\ \Upsilon(\lambda; s_0; \tilde{w}_{1,n}) = \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \tilde{w}_{1,n}^{i_0} \right) \end{cases}$$

Proof Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on the form of integrals of Lamé equation of type 2 $y(z)$,

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y(z) \\ &= \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} \{y_0(z) + y_1(z) + y_2(z) + y_3(z) + \dots\} \end{aligned} \quad (8.4.5)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (8.3.8a),

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_0(z) \\ &= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_{0,\infty}^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1 + \lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \rho^{i_0} \right) \end{aligned} \quad (8.4.6)$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (8.3.9),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(z) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\rho(1-t_1)(1-u_1)v_1)^{-(4+2\lambda)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_{1,\infty}}{1-\rho(1-t_1)(1-u_1)v_1} \right)^{q_1} \overleftrightarrow{w}_{1,1}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{-\frac{\alpha}{2}+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2}+\lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right) \eta \quad (8.4.7)
\end{aligned}$$

Replace q_i , q_j and r_i by q_1 , q_0 and $\frac{v_1-1}{v_1} \frac{s_{1,\infty}}{1-\rho(1-t_1)(1-u_1)v_1}$ in (8.4.3). Take the new (8.4.3) into (8.4.7).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(z) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{(1-\rho(1-t_1)(1-u_1)v_1)^{-(3+2\lambda)}}{-\rho(1-t_1)(1-u_1)v_1^2 + (1-s_{1,\infty})v_1 + s_{1,\infty}} \\
&\times \overleftrightarrow{w}_{1,1}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{1,1} \partial \overleftrightarrow{w}_{1,1} \right) \overleftrightarrow{w}_{1,1}^{-\frac{\alpha}{2}+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} \left(\frac{v_1-1}{v_1} \frac{s_{0,\infty}}{1-\rho(1-t_1)(1-u_1)v_1} \right)^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2}+\lambda)_{i_0}} \overleftrightarrow{w}_{1,1}^{i_0} \right) \eta \quad (8.4.8)
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$v_1 = \frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4\rho(1-t_1)(1-u_1)s_{1,\infty}}}{2\rho(1-t_1)(1-u_1)}$$

$$\text{or } \frac{1-s_{1,\infty} + \sqrt{(1-s_{1,\infty})^2 + 4\rho(1-t_1)(1-u_1)s_{1,\infty}}}{2\rho(1-t_1)(1-u_1)} \text{ and}$$

$$\frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4\rho(1-t_1)(1-u_1)s_{1,\infty}}}{2\rho(1-t_1)(1-u_1)} \text{ is only inside the unit circle. As we}$$

compute the residue there in (8.4.8) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_1(z) \\
&= \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} (s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+2\lambda)} \\
&\quad \times \tilde{w}_{1,1}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{\alpha}{2}+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \tilde{w}_{1,1}^{i_0} \right) \eta
\end{aligned} \tag{8.4.9}$$

where

$$\begin{aligned}
\tilde{w}_{1,1} &= \frac{v_1}{(v_1 - 1)} \frac{\rho t_1 u_1}{1 - \rho(1-t_1)(1-u_1)v_1} \Bigg|_{v_1 = \frac{1-s_{1,\infty} - \sqrt{(1-s_{1,\infty})^2 + 4\rho(1-t_1)(1-u_1)s_{1,\infty}}}{2\rho(1-t_1)(1-u_1)}} \\
&= \frac{\rho t_1 u_1 \{1 + (s_{1,\infty} + 2\rho(1-t_1)(1-u_1))s_{1,\infty}\}}{2(1 - \rho(1-t_1)(1-u_1))^2 s_{1,\infty}} \\
&\quad - \frac{\rho t_1 u_1 (1 + s_{1,\infty}) \sqrt{s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2(1 - \rho(1-t_1)(1-u_1))^2 s_{1,\infty}}
\end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (8.3.11),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{1}{v_2} (1-\rho(1-t_2)(1-u_2)v_2)^{-(8+2\lambda)} \\
&\times \sum_{q_2=q_1}^{\infty} \left(\frac{v_2-1}{v_2} \frac{s_{2,\infty}}{1-\rho(1-t_2)(1-u_2)v_2} \right)^{q_2} \overleftarrow{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftarrow{w}_{2,2} \partial \overleftarrow{w}_{2,2} \right) \overleftarrow{w}_{2,2}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{2,2} \partial \overleftarrow{w}_{2,2} \right) \overleftarrow{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1-\overleftarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-(4+2\lambda)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_1}{1-\overleftarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
&\times \overleftarrow{w}_{1,2}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftarrow{w}_{1,2} \partial \overleftarrow{w}_{1,2} \right) \overleftarrow{w}_{1,2}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{1,2} \partial \overleftarrow{w}_{1,2} \right) \overleftarrow{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2}+\lambda)_{i_0}} \overleftarrow{w}_{1,2}^{i_0} \right) \eta^2 \tag{8.4.10}
\end{aligned}$$

Replace q_i , q_j and r_i by q_2 , q_1 and $\frac{v_2-1}{v_2} \frac{s_{2,\infty}}{1-\rho(1-t_2)(1-u_2)v_2}$ in (8.4.3). Take the new

(8.4.3) into (8.4.10).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} \frac{1}{2\pi i} \oint dv_2 \frac{(1-\rho(1-t_2)(1-u_2)v_2)^{-(7+2\lambda)}}{-\rho(1-t_2)(1-u_2)v_2^2 + (1-s_{2,\infty})v_2 + s_{2,\infty}} \\
&\times \overleftrightarrow{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{2,2} \partial_{\overleftrightarrow{w}_{2,2}} \right) \overleftrightarrow{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1 - \overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-(4+2\lambda)} \\
&\times \sum_{q_1=q_0}^{\infty} \left(\frac{v_2-1}{v_2} \frac{s_{1,\infty}}{1-\rho(1-t_2)(1-u_2)v_2} \frac{v_1-1}{v_1} \frac{1}{1-\overleftrightarrow{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \\
&\times \overleftrightarrow{w}_{1,2}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{1,2} \partial_{\overleftrightarrow{w}_{1,2}} \right) \overleftrightarrow{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \\
&\times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2}+\lambda)_{i_0}} \overleftrightarrow{w}_{1,2}^{i_0} \right) \eta^2
\end{aligned} \tag{8.4.11}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$\begin{aligned}
v_2 &= \frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4\rho(1-t_2)(1-u_2)s_{2,\infty}}}{2\rho(1-t_2)(1-u_2)} \\
&\text{or } \frac{1-s_{2,\infty} + \sqrt{(1-s_{2,\infty})^2 + 4\rho(1-t_2)(1-u_2)s_{2,\infty}}}{2\rho(1-t_2)(1-u_2)} \text{ and} \\
&\frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4\rho(1-t_2)(1-u_2)s_{2,\infty}}}{2\rho(1-t_2)(1-u_2)} \text{ is only inside the unit circle. As we}
\end{aligned}$$

compute the residue there in (8.4.11) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} (s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+2\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\alpha+\frac{1}{2}} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{1}{v_1} (1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-(4+2\lambda)} \\
&\quad \times \sum_{q_1=q_0}^{\infty} \left(\frac{v_1-1}{v_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_1} \ddot{w}_{1,2}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\alpha+\frac{1}{2}} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2}+\lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) \eta^2 \tag{8.4.12}
\end{aligned}$$

where

$$\begin{aligned}
\tilde{w}_{2,2} &= \frac{v_2}{(v_2-1)} \frac{\rho t_2 u_2}{1 - \rho(1-t_2)(1-u_2)v_2} \Big|_{v_2 = \frac{1-s_{2,\infty} - \sqrt{(1-s_{2,\infty})^2 + 4\rho(1-t_2)(1-u_2)s_{2,\infty}}}{2\rho(1-t_2)(1-u_2)}} \\
&= \frac{\rho t_2 u_2 \{1 + (s_{2,\infty} + 2\rho(1-t_2)(1-u_2))s_{2,\infty}\}}{2(1 - \rho(1-t_2)(1-u_2))^2 s_{2,\infty}} \\
&\quad - \frac{\rho t_2 u_2 (1 + s_{2,\infty}) \sqrt{s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2(1 - \rho(1-t_2)(1-u_2))^2 s_{2,\infty}}
\end{aligned}$$

and

$$\ddot{w}_{1,2} = \frac{v_1}{(v_1-1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1}$$

Replace q_i , q_j and r_i by q_1 , q_0 and $\frac{v_1-1}{v_1} \frac{s_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1}$ in (8.4.3). Take the

new (8.4.3) into (8.4.12).

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} (s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+2\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\alpha+\frac{1}{2}} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} \frac{1}{2\pi i} \oint dv_1 \frac{(1-\tilde{w}_{2,2}(1-t_1)(1-u_1)v_1)^{-(3+2\lambda)}}{-\tilde{w}_{2,2}(1-t_1)(1-u_1)v_1^2 + (1-s_1)v_1 + s_1} \\
&\quad \times \ddot{w}_{1,2}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{\alpha+\frac{1}{2}} (\ddot{w}_{1,2} \partial_{\ddot{w}_{1,2}}) \ddot{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} \left(\frac{v_1-1}{v_1} \frac{s_{0,1}}{1-\tilde{w}_{2,2}(1-t_1)(1-u_1)v_1} \right)^{q_0} \\
&\quad \times \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0+2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2}+\lambda)_{i_0}} \ddot{w}_{1,2}^{i_0} \right) \eta^2 \tag{8.4.13}
\end{aligned}$$

By using Cauchy's integral formula, the contour integrand has poles at

$$\begin{aligned}
v_1 &= \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)} \\
&\text{or } \frac{1-s_1 + \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)} \text{ and} \\
&\frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)} \text{ is only inside the unit circle. As we}
\end{aligned}$$

compute the residue there in (8.4.13) we obtain

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_2(z) \\
&= \prod_{k=3}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} (s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{2,\infty} + \sqrt{s_{2,\infty}^2 - 2(1-2\rho(1-t_2)(1-u_2))s_{2,\infty} + 1}}{2} \right)^{-(7+2\lambda)} \\
&\quad \times \tilde{w}_{2,2}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{\alpha+\frac{1}{2}} (\tilde{w}_{2,2} \partial_{\tilde{w}_{2,2}}) \tilde{w}_{2,2}^{2-\frac{\alpha}{2}+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} (s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-(3+2\lambda)} \\
&\quad \times \tilde{w}_{1,2}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,2} \partial_{\tilde{w}_{1,2}}) \tilde{w}_{1,2}^{-\frac{\alpha}{2}+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \tilde{w}_{1,2}^{i_0} \right) \eta^2 \tag{8.4.14}
\end{aligned}$$

where

$$\begin{aligned}
\tilde{w}_{1,2} &= \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,2} t_1 u_1}{1 - \tilde{w}_{2,2}(1-t_1)(1-u_1)v_1} \Big|_{v_1 = \frac{1-s_1 - \sqrt{(1-s_1)^2 + 4\tilde{w}_{2,2}(1-t_1)(1-u_1)s_1}}{2\tilde{w}_{2,2}(1-t_1)(1-u_1)}} \\
&= \frac{\tilde{w}_{2,2} t_1 u_1 \{1 + (s_1 + 2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1\}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1} \\
&\quad - \frac{\tilde{w}_{2,2} t_1 u_1 (1 + s_1) \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,2}(1-t_1)(1-u_1))s_1 + 1}}{2(1 - \tilde{w}_{2,2}(1-t_1)(1-u_1))^2 s_1}
\end{aligned}$$

Acting the summation operator $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\}$ on (8.3.12),

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_3(z) \\
&= \prod_{k=4}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_3 t_3^{5+\lambda} \int_0^1 du_3 u_3^{\frac{9}{2}+\lambda} (s_{3,\infty}^2 - 2(1-2\rho(1-t_3)(1-u_3))s_{3,\infty} + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_{3,\infty} + \sqrt{s_{3,\infty}^2 - 2(1-2\rho(1-t_3)(1-u_3))s_{3,\infty} + 1}}{2} \right)^{-(11+2\lambda)} \\
&\quad \times \tilde{w}_{3,3}^{-(\frac{9}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{\alpha+\frac{1}{2}} (\tilde{w}_{3,3} \partial_{\tilde{w}_{3,3}}) \tilde{w}_{3,3}^{4-\frac{\alpha}{2}+\lambda} \\
&\quad \times \int_0^1 dt_2 t_2^{3+\lambda} \int_0^1 du_2 u_2^{\frac{5}{2}+\lambda} (s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_2 + \sqrt{s_2^2 - 2(1-2\tilde{w}_{3,3}(1-t_2)(1-u_2))s_2 + 1}}{2} \right)^{-(7+2\lambda)} \\
&\quad \times \tilde{w}_{2,3}^{-(\frac{5}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{\alpha+\frac{1}{2}} (\tilde{w}_{2,3} \partial_{\tilde{w}_{2,3}}) \tilde{w}_{2,3}^{2-\frac{\alpha}{2}+\lambda} \\
&\quad \times \int_0^1 dt_1 t_1^{1+\lambda} \int_0^1 du_1 u_1^{\frac{1}{2}+\lambda} (s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1)^{-\frac{1}{2}} \\
&\quad \times \left(\frac{1 + s_1 + \sqrt{s_1^2 - 2(1-2\tilde{w}_{2,3}(1-t_1)(1-u_1))s_1 + 1}}{2} \right)^{-(3+2\lambda)} \\
&\quad \times \tilde{w}_{1,3}^{-(\frac{1}{2}+\frac{\alpha}{2}+\lambda)} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,3} \partial_{\tilde{w}_{1,3}}) \tilde{w}_{1,3}^{-\frac{\alpha}{2}+\lambda} \\
&\quad \times \sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \left(c_0 z^\lambda \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} (q_0 + 2\lambda)_{i_0}}{(1+\lambda)_{i_0} (\frac{1}{2} + \lambda)_{i_0}} \tilde{w}_{1,3}^{i_0} \right) \eta^3
\end{aligned} \tag{8.4.15}$$

where

$$\begin{aligned}
\tilde{w}_{3,3} &= \frac{v_3}{(v_3 - 1)} \frac{\rho t_3 u_3}{1 - \rho(1 - t_3)(1 - u_3)v_3} \Bigg|_{v_3 = \frac{1 - s_{3,\infty} - \sqrt{(1 - s_{3,\infty})^2 + 4\rho(1 - t_3)(1 - u_3)s_{3,\infty}}}{2\rho(1 - t_3)(1 - u_3)}} \\
&= \frac{\rho t_3 u_3 \{1 + (s_{3,\infty} + 2\rho(1 - t_3)(1 - u_3))s_{3,\infty}\}}{2(1 - \rho(1 - t_3)(1 - u_3))^2 s_{3,\infty}} \\
&\quad - \frac{\rho t_3 u_3 (1 + s_{3,\infty}) \sqrt{s_{3,\infty}^2 - 2(1 - 2\rho(1 - t_3)(1 - u_3))s_{3,\infty} + 1}}{2(1 - \rho(1 - t_3)(1 - u_3))^2 s_{3,\infty}} \\
\tilde{w}_{2,3} &= \frac{v_2}{(v_2 - 1)} \frac{\tilde{w}_{3,3} t_2 u_2}{1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2)v_2} \Bigg|_{v_2 = \frac{1 - s_2 - \sqrt{(1 - s_2)^2 + 4\tilde{w}_{3,3}(1 - t_2)(1 - u_2)s_2}}{2\tilde{w}_{3,3}(1 - t_2)(1 - u_2)}} \\
&= \frac{\tilde{w}_{3,3} t_2 u_2 \{1 + (s_2 + 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2\}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2} \\
&\quad - \frac{\tilde{w}_{3,3} t_2 u_2 (1 + s_2) \sqrt{s_2^2 - 2(1 - 2\tilde{w}_{3,3}(1 - t_2)(1 - u_2))s_2 + 1}}{2(1 - \tilde{w}_{3,3}(1 - t_2)(1 - u_2))^2 s_2} \\
\tilde{w}_{1,3} &= \frac{v_1}{(v_1 - 1)} \frac{\tilde{w}_{2,3} t_1 u_1}{1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1)v_1} \Bigg|_{v_1 = \frac{1 - s_1 - \sqrt{(1 - s_1)^2 + 4\tilde{w}_{2,3}(1 - t_1)(1 - u_1)s_1}}{2\tilde{w}_{2,3}(1 - t_1)(1 - u_1)}} \\
&= \frac{\tilde{w}_{2,3} t_1 u_1 \{1 + (s_1 + 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1\}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1} \\
&\quad - \frac{\tilde{w}_{2,3} t_1 u_1 (1 + s_1) \sqrt{s_1^2 - 2(1 - 2\tilde{w}_{2,3}(1 - t_1)(1 - u_1))s_1 + 1}}{2(1 - \tilde{w}_{2,3}(1 - t_1)(1 - u_1))^2 s_1}
\end{aligned}$$

By repeating this process for all higher terms of integral forms of sub-summation $y_m(z)$

terms where $m \geq 4$, I obtain every $\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(z)$ terms.

Substitute (8.4.6), (8.4.9), (8.4.14), (8.4.15) and including all

$$\sum_{q_0=0}^{\infty} \frac{(\gamma')_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} y_m(z) \text{ terms where } m > 3 \text{ into (8.4.5).} \quad \blacksquare$$

Remark 8.4.4 The generating function for the Lamé polynomial of type 2 in the algebraic form of the first kind about $x = a$ as $q = \alpha(\alpha + 1)a - 4(2a - b - c)(q_j + 2j)^2$

where $j, q_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\frac{1}{2})_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} L F_{q_j}^R \left(a, b, c, \alpha, q = \alpha(\alpha+1)a - 4(2a-b-c)(q_j+2j)^2 \right. \\
& \left. ; z = x-a, \rho = -\frac{2a-b-c}{(a-b)(a-c)}z, \eta = \frac{-z^2}{(a-b)(a-c)} \right) \\
& = 2^{-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \rho) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \rho) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(1+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{\alpha}{2}} \mathbf{A}(s_0; \tilde{w}_{1,1}) \eta \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \rho) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4(n-1)-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k-1)-\alpha)} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} \eta^n \left. \right\} \quad (8.4.16)
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \rho) &= \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-3}}{\sqrt{s_{1,\infty}^2-2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \rho) &= \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2\rho(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-(4n-1)}}{\sqrt{s_{n,\infty}^2-2(1-2\rho(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)-1)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{aligned} \right.$$

and

$$\left\{ \begin{aligned} \mathbf{A}(s_{0,\infty}; \rho) &= \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \right.$$

Proof Replace w , γ , A and x by $s_{0,\infty}$, $\frac{1}{2}$, 0 and ρ in (8.4.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\frac{1}{2})_{q_0}}{q_0!} s_{0,\infty}^{q_0} {}_2F_1\left(-q_0, q_0; \frac{1}{2}; \rho\right) \\ &= \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{2\sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}} \end{aligned} \quad (8.4.17)$$

Replace $s_{0,\infty}$ and ρ by s_0 and $\tilde{w}_{1,1}$ in (8.4.17).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\frac{1}{2})_{q_0}}{q_0!} s_0^{q_0} {}_2F_1\left(-q_0, q_0; \frac{1}{2}; \tilde{w}_{1,1}\right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{2\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (8.4.18)$$

Replace $\tilde{w}_{1,1}$ by $\tilde{w}_{1,n}$ in (8.4.18).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\frac{1}{2})_{q_0}}{q_0!} s_0^{q_0} {}_2F_1\left(-q_0, q_0; \frac{1}{2}; \tilde{w}_{1,n}\right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{2\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (8.4.19)$$

Put $c_0 = 1$, $\lambda = 0$ and $\gamma' = \frac{1}{2}$ in (8.4.4). Substitute (8.4.17), (8.4.18) and (8.4.19) into the new (8.4.4). ■

Remark 8.4.5 The generating function for the Lamé polynomial of type 2 in the algebraic form of the second kind about $x = a$ as

$q = \alpha(\alpha + 1)a - 4(2a - b - c) \left(q_j + 2j + \frac{1}{2}\right)^2$ where $j, \alpha_j \in \mathbb{N}_0$ is

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{\left(\frac{3}{2}\right)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} LS_{q_j}^R \left(a, b, c, \alpha, q = \alpha(\alpha + 1)a - 4(2a - b - c) \left(q_j + 2j + \frac{1}{2}\right)^2 \right. \\
& \left. ; z = x - a, \rho = -\frac{2a - b - c}{(a - b)(a - c)}z, \eta = \frac{-z^2}{(a - b)(a - c)} \right) \\
& = z^{\frac{1}{2}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{B}(s_{0,\infty}; \rho) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{3}{2}} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \rho) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(2+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(1-\alpha)} \mathbf{B}(s_0; \tilde{w}_{1,1}) \eta \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\frac{1}{2}} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \rho) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-2+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-3-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \left. \left. \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-2+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-3-\alpha)} \right\} \mathbf{B}(s_0; \tilde{w}_{1,n}) \right\} \eta^n \Bigg\} \quad (8.4.20)
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \rho) &= \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-4}}{\sqrt{s_{1,\infty}^2 - 2(1-2\rho(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \rho) &= \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\rho(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-4n}}{\sqrt{s_{n,\infty}^2 - 2(1-2\rho(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-4(n-k)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{aligned} \right.$$

and

$$\left\{ \begin{aligned} \mathbf{B}(s_{0,\infty}; \rho) &= \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \right.$$

Proof Replace w , γ , A and x by $s_{0,\infty}$, $\frac{3}{2}$, 1 and ρ in (8.4.1).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\frac{3}{2})_{q_0}}{q_0!} s_{0,\infty}^{q_0} {}_2F_1\left(-q_0, q_0 + 1; \frac{3}{2}; \rho\right) \\ &= \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\rho)s_{0,\infty} + 1}} \end{aligned} \quad (8.4.21)$$

Replace $s_{0,\infty}$ and ρ by s_0 and $\tilde{w}_{1,1}$ in (8.4.21).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\frac{3}{2})_{q_0}}{q_0!} s_0^{q_0} {}_2F_1\left(-q_0, q_0 + 1; \frac{3}{2}; \tilde{w}_{1,1}\right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \end{aligned} \quad (8.4.22)$$

Replace $\tilde{w}_{1,1}$ by $\tilde{w}_{1,n}$ in (8.4.22).

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(\frac{3}{2})_{q_0}}{q_0!} s_0^{q_0} {}_2F_1\left(-q_0, q_0 + 1; \frac{3}{2}; \tilde{w}_{1,n}\right) \\ &= \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{aligned} \quad (8.4.23)$$

Put $c_0 = 1$, $\lambda = \frac{1}{2}$ and $\gamma' = \frac{3}{2}$ in (8.4.4). Substitute (8.4.21), (8.4.22) and (8.4.23) into the new (8.4.4). $\blacksquare$

8.5 Summary

Lame equation represented either in the algebraic form or in Weierstrass's form has a relation between three different coefficients in the power series of its equation as we see (8.1.6)–(8.1.7c). Lame equation could not be described in the form of a definite or contour integral of any elementary functions because of its three term recurrence relation in its power series. Three different coefficients in its recursion relation create complex mathematical calculations in order to obtain analytic solutions of it in closed forms.

In Ref.[12] I show how to obtain the power series expansion in closed forms and its integral form of Lame equation in the algebraic form (for infinite series and polynomial of type 1 including all higher terms of A_n 's) by applying 3TRF. This was done by letting A_n in sequence c_n is the leading term in the analytic function $y(z)$: the sequence c_n consists of combinations A_n and B_n . For polynomial of type 1, I treat q as a free variable and a fixed value of α .

In this chapter I show how to construct the power series expansion and its integral form of Lame equation in the algebraic form for infinite series and polynomial of type 2 including all higher terms of B_n 's by applying R3TRF. This is done by letting B_n in sequence c_n is the leading term in a analytic function $y(z)$. For polynomial of type 2, I treat α as a free variable and a fixed value of q .

The Frobenius solution and its integral form of Lame equation (ellipsoidal harmonics equation) for infinite series about $x = a$ in this chapter are equivalent to infinite series of Lame equation in Ref.[12]. In this chapter B_n is the leading term in sequence c_n in an analytic function $y(z)$. In Ref.[12] A_n is the leading term in sequence c_n in the analytic function $y(z)$.

In Ref.[12] and this chapter, as we see the power series expansions of Lame equation about $x = a$ for either polynomial or infinite series, the denominators and numerators in all A_n or B_n terms of $y(z)$ arise with Pochhammer symbol. Since we construct the power series expansions with Pochhammer symbols in numerators and denominators, we are able to describe integral representations of Lame equation analytically. As we observe representations in closed form integrals of Lame equation by applying either 3TRF or R3TRF, a ${}_2F_1$ function recurs in each of sub-integral forms of the $y(z)$ (each sub-integral $y_m(z)$ where $m = 0, 1, 2, \dots$ is composed of $2m$ terms of definite integrals and m terms of contour integrals). We are able to transform the Lame function into any well-known special functions having two recurrence relation in its power series of any linear differential equations because of a ${}_2F_1$ function in each of sub-integral forms of Lame function. After we replace ${}_2F_1$ function in its integral forms to other special functions (such as Bessel function, Kummer function, hypergeometric function, Laguerre function

and etc), we are able to rebuild the Frobenius solution of Lamé equation in a backward.

In this chapter I show how to construct the generating function for the type 2 Lamé polynomial from its integral representation. We are able to derive orthogonal relations, recursion relations and expectation values of physical quantities from the generating function: the processes in order to obtain orthogonal and recursion relations of the Lamé polynomials are similar as the case of a normalized wave function for the hydrogen-like atoms.

Mathematical structure of the power series expansions, integral representations and generating functions for Lamé equation closely resembles the case of Heun equation using 3TRF and R3TRF. (1) If $\gamma, \delta, \epsilon \rightarrow \frac{1}{2}$, $1 \rightarrow b - a$, $a \rightarrow c - a$, $\alpha \rightarrow \frac{1}{2}(\alpha + 1)$, $\beta \rightarrow -\frac{1}{2}\alpha$, $q \rightarrow -q + \frac{1}{4}\alpha(\alpha + 1)a$ and $x \rightarrow z$ in the power series expansions and its integral forms of Heun equation using 3TRF in Ref.[10, 11], its analytic solutions are correspondent to the Frobenius solutions and integral representations of Lamé equation in the algebraic form in Ref.[12]; compare (10), (11), (18) and (19) in Ref.[10] with remarks 1–4 in Ref.[12]. And make a comparison between remarks 1, 2, 5 and 6 in Ref.[11] and remarks 5–8 in Ref.[12]. (2) By similar reason, putting (8.1.4) into the power series, integral representations and generating functions of Heun equation using R3TRF in chapters 2 and 3, its analytic solutions are equivalent to the Frobenius solutions, integral forms and generating functions of Lamé equation using R3TRF in this chapter; compare remarks 2.2.1–2.2.4, 2.3.2–2.3.4 in chapter 2 and remarks 3.2.5–3.2.6 in chapter 3 with remarks 8.2.1–8.2.4, 8.3.2–8.3.5, 8.4.4–8.4.5 in this chapter.

Bibliography

- [1] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische Annalen* **33**, (1889)161.
- [2] Lamé, G., “Sur les surfaces isothermes dans les corps homogènes en équilibre de température,” (French) *J. Math. Pures. Appl.* **2**, 147-188(1837).
- [3] Brack, M., Mehta, M. and Tanaka, K., “Occurrence of periodic Lamé functions at bifurcations in chaotic Hamiltonian systems,” *J. Phys. A*, **34**(40) 8199-8220(2001).
- [4] Bronski, J. C. , Carr, L. D., Deconinck, B., Kutz, J. N. and Promislow, K., “Stability of repulsive Bose-Einstein condensates in a periodic potential,” *Phys. Rev. A*, **3**(63), 1-11(2001).
- [5] Zun-Tao, F., Shi-Kuo, L., Shi-Da, L. and Qiang1, Z., “Lamé Function and Its Application to Some Nonlinear Evolution Equations,” (Beijing, China) *Commun. Theor. Phys.* **40**, 53-56(2003).
- [6] Hobson, E.W., *The theory of spherical and ellipsoidal harmonics*. Cambridge Univ. Press, (1931).
- [7] Whittaker, E.T. and Watson, G.N., *A course of modern analysis*. Cambridge Univ. Press, (1952).
- [8] Romain, G. and Jean-Pierre, B., “Ellipsoidal Harmonic expansions of the gravitational potential: Theory and application,” *Celest. Mech. Dyn. Astron.*, **79**(4), 235–275(2001).
- [9] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806.
- [10] Choun, Y.S., “The analytic solution for the power series expansion of Heun function,” <http://dx.doi.org/10.1016/j.aop.2013.06.020> Available as arXiv:1303.0830.

- [11] Choun, Y.S., "Asymptotic behavior of Heun function and its integral formalism," arXiv:1303.0876.
- [12] Choun, Y.S., "Lame equation in the algebraic form," arXiv:1303.0873.
- [13] Stieltjes, T. J., "Sur certain polynomes que verifient une equation diferentielle lineaire du second ordre et sur la teorie des fonctions de Lamé," *Acta Math.* **6**, 321–326(1885).
- [14] Moon, P. and Spencer, D. E., *Field Theory for Engineers*. New York: Van Nostrand, (1961).
- [15] Zwillinger, D., *Handbook of Differential Equations*. 3rd ed. Boston, MA: Academic Press, (1997).
- [16] Bôcher, M., "Über die Reihenentwickelungen der Potentialtheorie," Leipzig: B. G. Teubner (1894).
- [17] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G., *Higher Transcendental Functions. Vol. III* McGraw-Hill Book Company, Inc., New York-Toronto-London, (1955).
- [18] Arscott, F. M., "Integral equations and relations for Lamé functions," *Quart. J. Math. Oxford Ser. (2)* **15**, 103-115(1964).
- [19] Shail, R., "On integral representations for Lamé and other special functions," *SIAM J. Math. Anal.* **11** (4), 702-723(1980).
- [20] Sleeman, B. D., "Integral equations and relations for Lamé functions and ellipsoidal wave functions," *Proc. Cambridge Philos. Soc.* **64**, 113-126(1968).
- [21] Volkmer, H., "Integral relations for Lamé functions," *SIAM J. Math. Anal.* **13** (6), 978-987(1982).
- [22] Volkmer, H., "Integralgleichungen fr periodische Lösungen Hillscher Differentialgleichungen," (German) *Analysis* **3** (1-4), 189-203(1983).
- [23] Volkmer, H., "Integral representations for products of Lamé functions by use of fundamental solutions," *SIAM J. Math. Anal.* **15** (3), 559-569(1984).
- [24] Wang, Z.X. and Guo, D. R., *Special Functions*, World Scientific Pub., 602–606(1989).
- [25] Whittaker, E.T. and Watson, G.N., "A Course of Modern Analysis (Cambridge Mathematical Library)," Cambridge University Press, 554–575(1996).

Chapter 9

Lame function in Weierstrass's form using reversible three-term recurrence formula

Lame ordinary differential equation in Weierstrass's form and Heun equation are of Fuchsian types with the four regular singularities. Lame equation in Weierstrass's form is derived from Heun equation by changing all coefficients $\gamma = \delta = \epsilon = \frac{1}{2}$, $a = \rho^{-2}$, $\alpha = \frac{1}{2}(\alpha + 1)$, $\beta = -\frac{1}{2}\alpha$, $q = -\frac{1}{4}h\rho^{-2}$ and an independent variable $x = sn^2(z, \rho)$. [1, 2]

In Ref.[10, 11] I construct the power series expansion in closed forms, its integral form and the generating function of Lame equation in Weierstrass's form by applying three term recurrence formula (3TRF)[6]; analytic solutions of Lame equation is derived for infinite series and polynomial of type 1¹ including all higher terms of A_n 's.²

In this chapter I will apply reversible three term recurrence formula (R3TRF) in chapter 1 to (1) the Frobenius solution in closed forms, (2) its integral form of Lame equation in Weierstrass's form for infinite series and polynomial of type 2³ including all higher terms of B_n 's⁴, (3) the generating function for the Lame polynomial of type 2.

Nine examples of 192 local solutions of the Heun equation (Maier, 2007) are provided in

¹polynomial of type 1 is a polynomial which makes B_n term terminated in three term recursion relation of the power series in a linear differential equation.

²“ higher terms of A_n 's” means at least two terms of A_n 's.

³polynomial of type 2 is a polynomial which makes A_n term terminated in three term recursion relation of the power series in a linear differential equation.

⁴“ higher terms of B_n 's” means at least two terms of B_n 's.

the appendix. For each example, I show how to convert local solutions of Heun equation by applying R3TRF to analytic solutions of Lamé equation in Weierstrass's form.

9.1 Introduction

In general, there are two types of Lamé equation which are described in the algebraic form and in Weierstrass's form. For computational practice, Jacobian (Weierstrass's) form is more convenient. For the general mathematical properties of analytic solutions of Lamé equation, the algebraic form is better; especially for the asymptotic expansion of Lamé equation, the algebraic form is much better and its asymptotic solutions are more various because of three different singularity parameters. In contrast, the Jacobian form only has two singularity parameters which are 1 and the modulus of the elliptic function $sn z$. Various authors have worked on the boundary value problems in ellipsoidal geometry with Lamé equation in Weierstrass's form rather than the algebraic form of it.

Many great mathematicians/physicists attempted to describe Lamé equation in a definite or contour integral form of any well-known simple functions such as Gauss hypergeometric, Kummer functions, and etc. Due to its mathematical complexity there are no analytic solutions in closed forms of the Lamé function [16, 4, 5]. Because Lamé equation represented either in the algebraic form or in Weierstrass's form is a form of the power series that is expressed as three term recurrence relation [4, 5]. In contrast, most of well-known special functions consist of two term recursion relation (Hypergeometric, Bessel, Legendre, Kummer functions, etc). They just left analytic solutions of Lamé equation as solutions of recurrences because of three different coefficients in its power series expansion. Three or more terms recursion relation in the power series of any linear ordinary differential equations creates the complex mathematical calculation.

In Ref.[9] I construct the power series expansion in closed forms and its integral representation of Lamé equation in the algebraic form by applying 3TRF [6]: analytic solutions of Lamé equation are derived for infinite series and polynomial of type 1 including all higher terms of A_n 's. One interesting observation resulting from calculations is the fact that a ${}_2F_1$ function recurs in each of sub-integral forms: the first sub-integral form contains zero term of A_n 's, the second one contains one term of A_n 's, the third one contains two terms of A_n 's, etc. Also asymptotic expansions of the Lamé function for infinite series are derived analytically including for special cases as $|2a - b - c| \gg 1$ and $2a - b - c \approx 0$.

In Ref.[10], by changing all coefficients in the general expression of the power series and its integral representation of Lamé equation in the algebraic form, I obtain the Frobenius solution and its integral form of Lamé equation in Weierstrass's form for infinite series

and polynomial of type 1. In Ref.[11] I construct the generating function for the Lamé polynomial of type 1 in Weierstrass's form from the general expression of an integral representation of Lamé equation in Weierstrass's form by applying the generating function for the Jacobi polynomial using hypergeometric functions.

In chapter 8, by applying R3TRF, I derive (1) the power series expansion, (2) its integral form of Lamé equation in the algebraic form for infinite series and polynomial of type 2 including all higher terms of B_n 's and (3) the generating function for the Lamé polynomial of type 2.

In this chapter, by changing all coefficients in the general expression of power series, its integral representation and the generating function of Heun equation in chapter 2 and 3, I consider (1) the power series expansion in closed forms of Lamé equation in Weierstrass's form (for infinite series and polynomial of type 2 including all higher terms of B_n 's), (2) its integral representation and (3) the generating function for the Lamé polynomial of type 2 analytically.⁵ Especially we are able to obtain a mathematical construction for an orthogonal relation of Lamé equation in Weierstrass's form for polynomial of type 1 or 2 from the generating function for the Lamé polynomial. After that, we might be possible to build recurrence relations of the Lamé polynomial, a normalized constant of any wave functions for Laplace equation in ellipsoidal coordinates, an expectation value of its physical quantities and etc. For an example, a normalized wave function of hydrogen-like atoms is derived from the generating function for associated Laguerre polynomials in explicit mathematical calculations. Also the expectation value of physical quantities such as position and momentum is constructed by applying the recursive relation of associated Laguerre polynomials.

I show how to transform nine examples of 192 local solutions of Heun equation [26] to analytic solutions of Lamé equation in Weierstrass's form in the appendix. For each example, I convert local solutions of Heun equation by applying R3TRF to analytic solutions of Lamé equation in Weierstrass's form; the Frobenius solution in closed form of Lamé equation, its integral representation and the generating function of it are derived from changing all coefficients and an independent variable in Heun equation.

Lamé equation in Weierstrass's form is given by

$$\frac{d^2 y}{dz^2} = \{\alpha(\alpha + 1)\rho^2 sn^2(z, \rho) - h\}y(z) \quad (9.1.1)$$

where ρ , α and h are real parameters such that $0 < \rho < 1$ and $\alpha \geq -\frac{1}{2}$. If we take $sn^2(z, \rho) = \xi$ as an independent variable, Lamé equation becomes

$$\frac{d^2 y}{d\xi^2} + \frac{1}{2} \left(\frac{1}{\xi} + \frac{1}{\xi - 1} + \frac{1}{\xi - \rho^{-2}} \right) \frac{dy}{d\xi} + \frac{-\alpha(\alpha + 1)\xi + h\rho^{-2}}{4\xi(\xi - 1)(\xi - \rho^{-2})} y(\xi) = 0 \quad (9.1.2)$$

⁵Or we can obtain analytic solutions of Lamé equation in Weierstrass's form by applying R3TRF directly.

This is an equation of Fuchsian type with the four regular singularities: $\xi = 0, 1, \rho^{-2}, \infty$. The first three, namely $0, 1, \rho^{-2}$, have the property that the corresponding exponents are $0, \frac{1}{2}$ which is the same as the case of Lamé equation in the algebraic form.

Lamé equation is a second-order linear ordinary differential equation of the algebraic form [3, 12, 14]

$$\frac{d^2y}{dx^2} + \frac{1}{2} \left(\frac{1}{x-a} + \frac{1}{x-b} + \frac{1}{x-c} \right) \frac{dy}{dx} + \frac{-\alpha(\alpha+1)x+q}{4(x-a)(x-b)(x-c)}y = 0 \quad (9.1.3)$$

Lamé equation has four regular singular points: a, b, c and ∞ ; the exponents at the first three are all 0 and $\frac{1}{2}$, and those at infinity are $-\frac{1}{2}\alpha$ and $\frac{1}{2}(\alpha+1)$. [13, 15]

As we compare (9.1.2) with (9.1.3), all coefficients on the above are correspondent to the following way.

$$\begin{aligned} a &\longrightarrow 0 \\ b &\longrightarrow 1 \\ c &\longrightarrow \rho^{-2} \\ q &\longrightarrow h\rho^{-2} \\ x &\longrightarrow \xi = sn^2(z, \rho) \end{aligned} \quad (9.1.4)$$

Also (9.1.2) is a special case of Heun's equation. Heun equation is a second-order linear ordinary differential equation of the form [1].

$$\frac{d^2y}{dx^2} + \left(\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right) \frac{dy}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)}y = 0 \quad (9.1.5)$$

With the condition $\epsilon = \alpha + \beta - \gamma - \delta + 1$. The parameters play different roles: $a \neq 0$ is the singularity parameter, $\alpha, \beta, \gamma, \delta, \epsilon$ are exponent parameters, q is the accessory parameter which in many physical applications appears as a spectral parameter. Also, α and β are identical to each other. The total number of free parameters is six. It has four regular singular points which are $0, 1, a$ and ∞ with exponents $\{0, 1 - \gamma\}$, $\{0, 1 - \delta\}$, $\{0, 1 - \epsilon\}$ and $\{\alpha, \beta\}$.

As we compare (9.1.2) with (9.1.5), all coefficients on the above are correspondent to the

following way.

$$\begin{aligned}
 \gamma, \delta, \epsilon &\longleftrightarrow \frac{1}{2} \\
 a &\longleftrightarrow \rho^{-2} \\
 \alpha &\longleftrightarrow \frac{1}{2}(\alpha + 1) \\
 \beta &\longleftrightarrow -\frac{1}{2}\alpha \\
 q &\longleftrightarrow -\frac{1}{4}h\rho^{-2} \\
 x &\longleftrightarrow \xi = sn^2(z, \rho)
 \end{aligned} \tag{9.1.6}$$

9.2 Power series

9.2.1 Polynomial of type 2

In general Lamé (spectral) polynomial represented either in the algebraic form or in Weierstrass's form is defined as polynomial of type 3.⁶ Lamé polynomial comes from a Lamé equation that has a fixed integer value of α , just as it has a fixed value of h . In three-term recurrence formula, polynomial of type 3 I categorize as complete polynomial. In future papers I will derive the type 3 Lamé polynomial. In Ref.[10, 11] I derive the Frobenius solution, its integral form and the generating function for the Lamé polynomial of type 1: I treat h as a free variable and α as a fixed value. In this chapter I construct the power series expansion and its integral representation of the Lamé polynomial of type 2: I treat α as a free variable and h as a fixed value. Also the generating function for the lamé polynomial of type 2 is constructed analytically.

In chapter 2 the general expression of power series of Heun equation about $x = 0$ for

⁶polynomial of type 3 is a polynomial which makes A_n and B_n terms terminated at the same time in three term recursion relation of the power series in a linear differential equation. If A_n and B_n terms are not terminated, it turns to be infinite series.

polynomial of type 2 is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0+\lambda+\alpha)(i_0+\lambda+\beta)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \sum_{i_1=i_0}^{q_1} \frac{(-q_1)_{i_1} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_1}}{(-q_1)_{i_0} \left(q_1 + 4 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}} \frac{(3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \eta^{i_1} \Big\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{q_0} \frac{(i_0+\lambda+\alpha)(i_0+\lambda+\beta)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{q_k} \frac{(i_k+2k+\lambda+\alpha)(i_k+2k+\lambda+\beta)}{(i_k+2(k+1)+\lambda)(i_k+2k+1+\gamma+\lambda)} \right. \\
&\quad \times \frac{(-q_k)_{i_k} \left(q_k + 4k + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_k}}{(-q_k)_{i_{k-1}} \left(q_k + 4k + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_{k-1}}} \frac{(2k+1+\lambda)_{i_{k-1}} (2k+\gamma+\lambda)_{i_{k-1}}}{(2k+1+\lambda)_{i_k} (2k+\gamma+\lambda)_{i_k}} \Big\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{q_n} \frac{(-q_n)_{i_n} \left(q_n + 4n + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_n}}{(-q_n)_{i_{n-1}} \left(q_n + 4n + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_{n-1}}} \frac{(2n+1+\lambda)_{i_{n-1}} (2n+\gamma+\lambda)_{i_{n-1}}}{(2n+1+\lambda)_{i_n} (2n+\gamma+\lambda)_{i_n}} \eta^{i_n} \Big\} z^n \Big\} \quad (9.2.1)
\end{aligned}$$

where

$$\begin{cases} z = -\frac{1}{a}x^2 \\ \eta = \frac{(1+a)}{a}x \\ \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1) \\ q = -(q_j + 2j + \lambda)\{\varphi + (1+a)(q_j + 2j + \lambda)\} \text{ as } j, q_j \in \mathbb{N}_0 \\ q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0 \end{cases}$$

In the above, We have two indicial roots which are $\lambda = 0$ and $1 - \gamma$.

Put (9.1.6) in (9.2.1) with replacing q_i by h_i where $h_i \in \mathbb{N}_0$. Take $c_0 = 1$ as $\lambda = 0$ for the first independent solution of Lamé equation and $\lambda = \frac{1}{2}$ for the second one into the new (9.2.1).

Remark 9.2.1 *The power series expansion of Lamé equation in Weierstrass's form of the first kind for polynomial of type 2 about $\xi = 0$ as $h = 4(1 + \rho^2)(h_j + 2j)^2$ where $j, h_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
 y(\xi) &= LF_{h_j}^R(\rho, \alpha, h = 4(1 + \rho^2)(h_j + 2j)^2; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\
 &= \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \\
 &\quad + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-h_1)_{i_0} (h_1 + 4)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
 &\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
 &\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{\alpha}{2} + \frac{1}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(-h_k)_{i_k} (h_k + 4k)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
 &\quad \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \tag{9.2.2}
 \end{aligned}$$

Remark 9.2.2 *The power series expansion of Lamé equation in Weierstrass's form of the second kind for polynomial of type 2 about $\xi = 0$ as $h = 4(1 + \rho^2)(h_j + 2j + \frac{1}{2})^2$*

where $j, h_j \in \mathbb{N}_0$ is

$$\begin{aligned}
y(\xi) &= LS_{h_j}^R \left(\rho, \alpha, h = 4(1 + \rho^2) \left(h_j + 2j + \frac{1}{2} \right)^2 ; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2 \xi^2 \right) \\
&= \xi^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + 1)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{\alpha}{2} + 1) (i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2}) (i_0 + 2)} \frac{(-h_0)_{i_0} (h_0 + 1)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 5)_{i_1}}{(-h_1)_{i_0} (h_1 + 5)_{i_0}} \frac{\left(\frac{7}{2}\right)_{i_0} (3)_{i_0}}{\left(\frac{7}{2}\right)_{i_1} (3)_{i_1}} \eta^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{\alpha}{2} + 1) (i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2}) (i_0 + 2)} \frac{(-h_0)_{i_0} (h_0 + 1)_{i_0}}{\left(\frac{3}{2}\right)_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{\alpha}{2} + 1) (i_k + 2k - \frac{\alpha}{2} + \frac{1}{2})}{(i_k + 2k + \frac{5}{2}) (i_k + 2k + 2)} \right. \\
&\quad \times \frac{(-h_k)_{i_k} (h_k + 4k + 1)_{i_k} (2k + \frac{3}{2})_{i_{k-1}} (2k + 1)_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k + 1)_{i_{k-1}} (2k + \frac{3}{2})_{i_k} (2k + 1)_{i_k}} \left. \right\} \\
&\quad \times \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n + 1)_{i_n} (2n + \frac{3}{2})_{i_{n-1}} (2n + 1)_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n + 1)_{i_{n-1}} (2n + \frac{3}{2})_{i_n} (2n + 1)_{i_n}} \eta^{i_n} \left. \right\} z^n \Bigg\} \quad (9.2.3)
\end{aligned}$$

In Ref.[10, 11] I treat α as a fixed value and h as a free variable to construct Lamé polynomial of type 1 in Weierstrass's: (1) if $\alpha = 2(2\alpha_j + j)$ or $-2(2\alpha_j + j) - 1$ where $j, \alpha_j \in \mathbb{N}_0$, an analytic solution of Lamé equation turns to be the first kind of independent solutions of the Lamé polynomial of type 1. (2) if $\alpha = 2(2\alpha_j + j) + 1$ or $-2(2\alpha_j + j + 1)$, an analytic solution of Lamé equation turns to be the second kind of independent solutions of the Lamé polynomial of type 1.⁷

In this chapter I treat h as a fixed value and α as a free variable to construct the Lamé polynomial of type 2 in Weierstrass's form : (1) if $h = 4(1 + \rho^2)(h_j + 2j)^2$ where $j, h_j \in \mathbb{N}_0$, an analytic solution of Lamé equation turns to be the first kind of independent solutions of the Lamé polynomial of type 2. (2) if $h = 4(1 + \rho^2) \left(h_j + 2j + \frac{1}{2} \right)^2$, an analytic solution of the Lamé equation turns to be the second kind of independent solutions of the Lamé polynomial of type 2.

⁷If we take $\alpha \geq -\frac{1}{2}$, $\alpha = -2(2\alpha_j + j) - 1$ and $-2(2\alpha_j + j + 1)$ are not available any more. I consider α as arbitrary.

9.2.2 Infinite series

In chapter 2 the general expression of power series of Heun equation for infinite series about $x = 0$ by applying R3TRF is given by

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)(i_0+\lambda+\beta)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3+\lambda)_{i_0} (2+\gamma+\lambda)_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3+\lambda)_{i_1} (2+\gamma+\lambda)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0+\lambda+\alpha)(i_0+\lambda+\beta)}{(i_0+\lambda+2)(i_0+\lambda+1+\gamma)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+2k+\lambda+\alpha)(i_k+2k+\lambda+\beta)}{(i_k+2(k+1)+\lambda)(i_k+2k+1+\gamma+\lambda)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1+\lambda)_{i_{k-1}} (2k+\gamma+\lambda)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1+\lambda)_{i_k} (2k+\gamma+\lambda)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1+\lambda)_{i_{n-1}} (2n+\gamma+\lambda)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1+\lambda)_{i_n} (2n+\gamma+\lambda)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (9.2.4)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\{\varphi+2(1+a)\lambda\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_1^\pm = \frac{\{\varphi+2(1+a)(\lambda+2)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_k^\pm = \frac{\{\varphi+2(1+a)(\lambda+2k)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_n^\pm = \frac{\{\varphi+2(1+a)(\lambda+2n)\} \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \end{cases}$$

Put (9.1.6) in (9.2.4). Take $c_0 = 1$ as $\lambda = 0$ for the first independent solution of Lamé equation and $\lambda = \frac{1}{2}$ for the second one into the new (9.2.4).

Remark 9.2.3 *The power series expansion of Lamé equation in Weierstrass's form of*

the first kind for infinite series about $\xi = 0$ using R3TRF is given by

$$\begin{aligned}
y(\xi) &= LF^R(\rho, \alpha, h; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\
&= \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + \frac{1}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{\alpha}{2} + \frac{1}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \tag{9.2.5}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_1^{\pm} = 2 \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_k^{\pm} = 2k \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_n^{\pm} = 2n \pm \sqrt{\frac{h}{4(1+\rho^2)}} \end{cases}$$

Remark 9.2.4 The power series expansion of Lamé equation in Weierstrass's form of

the second kind for infinite series about $\xi = 0$ using R3TRF is given by

$$\begin{aligned}
y(\xi) &= LS^R(\rho, \alpha, h; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\
&= \xi^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + 1)(i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2})(i_0 + 2)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0}} \frac{(\frac{7}{2})_{i_0} (3)_{i_0}}{(\frac{7}{2})_{i_1} (3)_{i_1}} \eta^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{\alpha}{2} + 1)(i_0 - \frac{\alpha}{2} + \frac{1}{2})}{(i_0 + \frac{5}{2})(i_0 + 2)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(\frac{3}{2})_{i_0} (1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{\alpha}{2} + 1)(i_k + 2k - \frac{\alpha}{2} + \frac{1}{2})}{(i_k + 2k + \frac{5}{2})(i_k + 2k + 2)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}}} \frac{(2k + \frac{3}{2})_{i_{k-1}} (2k + 1)_{i_{k-1}}}{(2k + \frac{3}{2})_{i_k} (2k + 1)_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}}} \frac{(2n + \frac{3}{2})_{i_{n-1}} (2n + 1)_{i_{n-1}}}{(2n + \frac{3}{2})_{i_n} (2n + 1)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (9.2.6)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{1}{2} \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_1^{\pm} = \frac{5}{2} \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_k^{\pm} = 2k + \frac{1}{2} \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_n^{\pm} = 2n + \frac{1}{2} \pm \sqrt{\frac{h}{4(1+\rho^2)}} \end{cases}$$

The infinite series in this chapter are equivalent to the infinite series in Ref.[10]. In this chapter B_n is the leading term in sequence c_n of the analytic function $y(\xi)$. In Ref.[10] A_n is the leading term in sequence c_n of the analytic function $y(\xi)$.

9.3 Integral formalism

Lame equation could not be built in a definite or contour integral form of any well-known simple functions such as Gauss hypergeometric, Kummer functions and etc. According to Erdelyi (1940[17]), “there is no corresponding representation of simple integral formalisms of the solutions in ordinary linear differential equations with four regular singularities; Heun equation, Lamé equation and Mathieu equation. It appears that the theory of integral equations connected with periodic solutions of Lamé equation is not as complete as the corresponding theory of integral representations of, say, Legendre

functions.” Because Heun, Lamé and Mathieu equations are composed of three different coefficients in the power series expansion. The three term recurrence relation in their power series creates complicated mathematical calculations to be constructed into its direct or contour integrals.

Instead of analyzing Lamé equation into its integral representation of any simple functions, in earlier literature the integral equations of Lamé equation were constructed by using simple kernels involving Legendre functions of Jacobian elliptic function or ellipsoidal harmonic functions: such integral relationships express one analytic solution in terms of another analytic solution involving Jacobian elliptic functions. There are many other forms of integral relations in the Lamé equation.[16, 18, 19, 20, 21, 22, 23, 24, 25]

In Ref.[10], I show integral forms (each sub-integral $y_m(\xi)$ where $m = 0, 1, 2, \dots$ is composed of $2m$ terms of definite integrals and m terms of contour integrals) of Lamé equation in Weierstrass's form by changing all coefficients and an independent variable in the general expression of the integral form of Lamé equation in the algebraic form in Ref.[9]: a ${}_2F_1$ function recurs in each of sub-integral forms of Lamé equation. Now I consider integral representations of Lamé equation by changing all coefficients and an independent variable in the general expression of Heun equation in chapter 2.

9.3.1 Polynomial of type 2

In chapter 2 the general representation in the form of integral of Heun polynomial of type 2 about $x = 0$ is given by

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} \right. \\
 &\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \\
 &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{q_{n-k}} \\
 &\quad \times (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\left(4(n-k) + \frac{\varphi+2(1+a)\lambda}{(1+a)}\right)} \\
 &\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta+\lambda} \left. \right\} \\
 &\quad \times \sum_{i_0=0}^{q_0} \frac{(-q_0)_{i_0} \left(q_0 + \frac{\varphi+2(1+a)\lambda}{(1+a)} \right)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \left. \right\} z^n \Bigg\} \tag{9.3.1}
 \end{aligned}$$

where

$$\begin{cases}
 z = -\frac{1}{a}x^2 \\
 \eta = \frac{(1+a)}{a}x \\
 \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1) \\
 q = -(q_j + 2j + \lambda)\{\varphi + (1+a)(q_j + 2j + \lambda)\} \text{ as } j, q_j \in \mathbb{N}_0 \\
 q_i \leq q_j \text{ only if } i \leq j \text{ where } i, j \in \mathbb{N}_0
 \end{cases}$$

and

$$\overleftrightarrow{w}_{i,j} = \begin{cases} \frac{v_i}{(v_i-1)} \frac{\overleftrightarrow{w}_{i+1,j} t_i u_i}{1 - \overleftrightarrow{w}_{i+1,j} v_i (1-t_i)(1-u_i)} & \text{where } i \leq j \\ \eta & \text{only if } i > j \end{cases}$$

In the above, the first sub-integral form contains one term of B'_n s, the second one contains two terms of B_n 's, the third one contains three terms of B_n 's, etc.

Put (9.1.6) in (9.3.1) with replacing q_i by h_i where $h_i \in \mathbb{N}_0$. Take $c_0 = 1$ as $\lambda = 0$ for the first independent solution of Lamé equation and $\lambda = \frac{1}{2}$ for the second one into the new (9.3.1).

Remark 9.3.1 *The integral representation of Lamé equation in Weierstrass's form of the first kind for polynomial of type 2 about $\xi = 0$ as $h = 4(1 + \rho^2)(h_j + 2j)^2$ where $j, h_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
y(\xi) &= LF_{h_j}^R(\rho, \alpha, h = 4(1 + \rho^2)(h_j + 2j)^2; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\
&= {}_2F_1\left(-h_0, h_0; \frac{1}{2}; \eta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
&\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-4(n-k)} \\
&\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-2-\frac{\alpha}{2}} \Big\} \\
&\quad \times {}_2F_1\left(-h_0, h_0; \frac{1}{2}; \overleftarrow{w}_{1,n}\right) \Big\} z^n \tag{9.3.2}
\end{aligned}$$

Remark 9.3.2 *The integral representation of Lamé equation in Weierstrass's form of the second kind for polynomial of type 2 about $\xi = 0$ as $h = 4(1 + \rho^2)(h_j + 2j + \frac{1}{2})^2$ where $j, h_j \in \mathbb{N}_0$ is*

$$\begin{aligned}
y(\xi) &= LS_{h_j}^R\left(\rho, \alpha, h = 4(1 + \rho^2)\left(h_j + 2j + \frac{1}{2}\right)^2; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2\right) \\
&= \xi^{\frac{1}{2}} \left\{ {}_2F_1\left(-h_0, h_0 + 1; \frac{3}{2}; \eta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
&\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+1)} \\
&\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \Big\} \\
&\quad \times {}_2F_1\left(-h_0, h_0 + 1; \frac{3}{2}; \overleftarrow{w}_{1,n}\right) \Big\} z^n \tag{9.3.3}
\end{aligned}$$

9.3.2 Infinite series

In chapter 2 the general representation in the form of integral of Heun equation for infinite series about $x = 0$ using R3TRF is given by

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
 &= c_0 x^\lambda \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \eta^{i_0} + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1+\lambda} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma+\lambda} \right. \right. \right. \\
 &\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\Delta_{n-k}^+} \\
 &\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k-1)+\alpha+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha-\beta} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)+\beta+\lambda} \Big\} \\
 &\times \left. \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1+\lambda)_{i_0} (\gamma+\lambda)_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \right\} z^n \Big\} \tag{9.3.4}
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \frac{\varphi+2(1+a)\lambda \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \\ \Delta_{n-k}^\pm = \frac{\varphi+2(1+a)(\lambda+2(n-k)) \pm \sqrt{\varphi^2-4(1+a)q}}{2(1+a)} \end{cases}$$

Put (9.1.6) in (9.3.4). Take $c_0 = 1$ as $\lambda = 0$ for the first independent solution of Lamé equation and $\lambda = \frac{1}{2}$ for the second one into the new (9.3.4).

Remark 9.3.3 *The integral representation of Lamé equation in Weierstrass's form of the first kind for infinite series about $\xi = 0$ using R3TRF is*

$$\begin{aligned}
 y(\xi) &= LF^R(\rho, \alpha, h; \xi = sn^2(z, \rho), \eta = (1+\rho^2)\xi, z = -\rho^2\xi^2) \\
 &= {}_2F_1\left(\Delta_0^-, \Delta_0^+; \frac{1}{2}; \eta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
 &\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1-t_{n-k})(1-u_{n-k})v_{n-k})^{-\Delta_{n-k}^+} \\
 &\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-2-\frac{\alpha}{2}} \Big\} \\
 &\times \left. {}_2F_1\left(\Delta_0^-, \Delta_0^+; \frac{1}{2}; \overleftrightarrow{w}_{1,n}\right) \right\} z^n \tag{9.3.5}
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^\pm = \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_{n-k}^\pm = 2(n-k) \pm \sqrt{\frac{h}{4(1+\rho^2)}} \end{cases}$$

Remark 9.3.4 The integral representation of Lamé equation in Weierstrass's form of the second kind for infinite series about $\xi = 0$ using R3TRF is

$$\begin{aligned} y(\xi) &= LS^R(\rho, \alpha, h; \xi = sn^2(z, \rho), \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\ &= \xi^{\frac{1}{2}} \left\{ {}_2F_1\left(\Delta_0^-, \Delta_0^+; \frac{3}{2}; \eta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \right. \right. \right. \\ &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{-\Delta_{n-k}^-} (1 - \overleftrightarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-\Delta_{n-k}^+} \\ &\quad \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \left. \right\} \\ &\quad \times {}_2F_1\left(\Delta_0^-, \Delta_0^+; \frac{3}{2}; \overleftrightarrow{w}_{1,n}\right) \left. \right\} z^n \end{aligned} \quad (9.3.6)$$

where

$$\begin{cases} \Delta_0^\pm = \frac{1}{2} \pm \sqrt{\frac{h}{4(1+\rho^2)}} \\ \Delta_{n-k}^\pm = 2(n-k) + \frac{1}{2} \pm \sqrt{\frac{h}{4(1+\rho^2)}} \end{cases}$$

9.4 Generating function for the Lamé polynomial of type 2

Definition 9.4.1 *I define that*

$$\left\{ \begin{array}{l} s_{a,b} = \begin{cases} s_a \cdot s_{a+1} \cdot s_{a+2} \cdots s_{b-2} \cdot s_{b-1} \cdot s_b & \text{if } a > b \\ s_a & \text{if } a = b \end{cases} \\ \\ \tilde{w}_{i,j} = \begin{cases} \frac{\tilde{w}_{i+1,j} t_i u_i \{1 + (s_i + 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i\}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} \\ - \frac{\tilde{w}_{i+1,j} t_i u_i (1 + s_i) \sqrt{s_i^2 - 2(1 - 2\tilde{w}_{i+1,j}(1-t_i)(1-u_i))s_i + 1}}{2(1 - \tilde{w}_{i+1,j}(1-t_i)(1-u_i))^2 s_i} & \text{where } i < j \\ \\ \frac{\eta t_i u_i \{1 + (s_{i,\infty} + 2\eta(1-t_i)(1-u_i))s_{i,\infty}\}}{2(1 - \eta(1-t_i)(1-u_i))^2 s_{i,\infty}} \\ - \frac{\eta t_i u_i (1 + s_{i,\infty}) \sqrt{s_{i,\infty}^2 - 2(1 - 2\eta(1-t_i)(1-u_i))s_{i,\infty} + 1}}{2(1 - \eta(1-t_i)(1-u_i))^2 s_{i,\infty}} & \text{where } i = j \end{cases} \end{array} \right.$$

where

$$a, b, i, j \in \mathbb{N}_0$$

In chapter 3 the generating function for the Heun polynomial of type 2 of the first kind about $x = 0$ as $q = -(q_j + 2j)\{\alpha + \beta - \delta + a(\delta + \gamma - 1) + (1 + a)(q_j + 2j)\}$ where

$j, q_j \in \mathbb{N}_0$ is given by

$$\begin{aligned}
& \sum_{q_0=0}^{\infty} \frac{(\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} H_{F_{q_j}}^R \left(q_j = \frac{-\varphi \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j \right. \\
& \left. , \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_1 = \frac{\varphi}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\
& = 2^{\frac{\varphi}{(1+a)}-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^\gamma \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\alpha} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^\beta \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2(n-1)+\gamma} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-(2(n-1)+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2(n-1)+\beta} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k-1)+\gamma} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \left. \left. \times \tilde{w}_{n-k,n}^{-(2(n-k-1)+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k-1)+\beta} \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} z^n \Bigg\} \quad (9.4.1)
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-(3+\Omega_1)}}{\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-(4n-1+\Omega_1)}}{\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-(4(n-k)-1+\Omega_1)}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{cases}$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{1-\gamma} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\gamma - \Omega_1}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{1-\gamma} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\gamma - \Omega_1}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{1-\gamma} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\gamma - \Omega_1}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

In chapter 3 the generating function for the Heun polynomial of type 2 of the second kind about $x = 0$ as $q = -(q_j + 2j + 1 - \gamma)\{\alpha + \beta + 1 - \gamma - (1 - a)\delta + (1 + a)(q_j + 2j)\}$ where $j, q_j \in \mathbb{N}_0$ with replacing $c_0 = \left(\frac{1+a}{a}\right)^{1-\gamma}$ by 1 is

$$\begin{aligned} & \sum_{q_0=0}^{\infty} \frac{(2-\gamma)_{q_0}}{q_0!} s_0^{q_0} \prod_{n=1}^{\infty} \left\{ \sum_{q_n=q_{n-1}}^{\infty} s_n^{q_n} \right\} HS_{q_j}^R \left(q_j = \frac{-\{\varphi + 2(1+a)(1-\gamma)\} \pm \sqrt{\varphi^2 - 4(1+a)q}}{2(1+a)} - 2j \right. \\ & \left. , \varphi = \alpha + \beta - \delta + a(\delta + \gamma - 1), \Omega_2 = \frac{\varphi + 2(1+a)(1-\gamma)}{(1+a)}; \eta = \frac{(1+a)}{a}x; z = -\frac{1}{a}x^2 \right) \\ & = 2^{\frac{\varphi + 2(1+a)(1/2-\gamma)}{(1+a)}} x^{1-\gamma} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{B}(s_{0,\infty}; \eta) \right. \\ & + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1^{2-\gamma} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \eta) \\ & \times \tilde{w}_{1,1}^{-(\alpha-\gamma+1)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha-\beta} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\beta-\gamma+1} \mathbf{B}(s_0; \tilde{w}_{1,1}) z \\ & + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\gamma} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ & \times \tilde{w}_{n,n}^{-(2n-1+\alpha-\gamma)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha-\beta} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{2n-1+\beta-\gamma} \\ & \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\gamma} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ & \left. \left. \times \tilde{w}_{n-k,n}^{-(2(n-k)-1+\alpha-\gamma)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha-\beta} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{2(n-k)-1+\beta-\gamma} \right\} \mathbf{B}(s_0; \tilde{w}_{1,n}) \right\} z^n \left. \right\} \end{aligned} \quad (9.4.2)$$

where

$$\left\{ \begin{array}{l} \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)}s_{1,\infty} + 1}{2} \right)^{-(3+\Omega_2)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)}s_{1,\infty} + 1} \\ \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)}s_{n,\infty} + 1}{2} \right)^{-(4n-1+\Omega_2)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)}s_{n,\infty} + 1} \\ \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{(1+s_{n-k}) + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))}s_{n-k} + 1}{2} \right)^{-(4(n-k)-1+\Omega_2)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))}s_{n-k} + 1} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{B}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{2-\gamma-\Omega_2}}{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{1-\gamma} \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{2-\gamma-\Omega_2}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{2-\gamma-\Omega_2}}{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{1-\gamma} \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Put (9.1.6) in (9.4.1) and (9.4.2) with replacing q_i by h_i where $h_i \in \mathbb{N}_0$ for the first and second independent solutions of Lamé equation.

Remark 9.4.2 The generating function for the Lamé polynomial of type 2 in Weierstrass's form of the first kind about $\xi = 0$ as $h = 4(1 + \rho^2)(h_j + 2j)^2$ where

$j, h_j \in \mathbb{N}_0$ is given by

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} L F_{h_j}^R(\rho, \alpha, h = 4(1 + \rho^2)(h_j + 2j)^2; \xi = sn^2(z, \rho) \\
& , \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\
& = 2^{-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(1+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{\alpha}{2}} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4(n-1)-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k-1)-\alpha)} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Big\} \quad (9.4.3)
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) &= \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-3}}{\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) &= \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-4n+1}}{\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-4(n-k)+1}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{aligned} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

Remark 9.4.3 The generating function for the Lamé polynomial of type 2 in Weierstrass's form of the second kind about $\xi = 0$ as $h = 4(1 + \rho^2)(h_j + 2j + \frac{1}{2})^2$ where $j, h_j \in \mathbb{N}_0$ is given by

$$\begin{aligned} & \sum_{h_0=0}^{\infty} \frac{(\frac{3}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} L_{h_j}^{SR} \left(\rho, \alpha, h = 4(1 + \rho^2) \left(h_j + 2j + \frac{1}{2} \right)^2; \xi = sn^2(z, \rho) \right. \\ & , \eta = (1 + \rho^2)\xi, z = -\rho^2\xi^2) \\ & = \xi^{\frac{1}{2}} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{B}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1^{\frac{3}{2}} \int_0^1 du_1 u_1 \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\ & \times \tilde{w}_{1,1}^{-\frac{1}{2}(2+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(1-\alpha)} \mathbf{B}(s_0; \tilde{w}_{1,1}) z \\ & + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-\frac{1}{2}} \int_0^1 du_n u_n^{2n-1} \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ & \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-2+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-3-\alpha)} \\ & \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-\frac{1}{2}} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-1} \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ & \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-2+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-3-\alpha)} \left. \right\} \mathbf{B}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Big\} \end{aligned} \quad (9.4.4)$$

where

$$\left\{ \begin{array}{l} \overleftrightarrow{\Psi}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-4}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Psi}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-4n}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Psi}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) = \frac{\left(\frac{(1+s_{n-k}) + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-4(n-k)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \mathbf{B}(s_{0,\infty}; \eta) = \frac{\left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{-\frac{1}{2}} \left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{-\frac{1}{2}} \left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{B}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{-\frac{1}{2}} \left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

9.5 Summary

Lame equation represented either in the algebraic form or in Weierstrass's form is the special case of Heun's differential equation. By changing all coefficients including an independent variable in Heun equation, we obtain the Lamé differential equation (ellipsoidal harmonics equation).

In Ref.[9] I construct the power series expansion of Lamé equation in the algebraic form (for infinite series and polynomial of type 1) and its integral representation by applying 3TRF. In Ref.[10] I derive the Frobenius solution and its integral form of Lamé equation in Weierstrass's form for infinite series and polynomial of type 1 by changing all coefficients and a variable of Lamé equation in the algebraic form in Ref.[9]. In Ref.[11] I obtain the generating function for the Lamé polynomial of type 1 in Weierstrass's form

by applying the generating function for the Jacobi polynomial using hypergeometric functions into the general representation in closed form integrals of Lamé equation in Ref.[10].

In chapter 8 I build a mathematical structure of the power series expansion of Lamé equation in the algebraic form (for infinite series and polynomial of type 2), its integral form and the generating function for the type 2 Lamé polynomial by applying R3TRF. In this chapter I construct analytic solutions of Lamé equation in Weierstrass's form for infinite series and polynomial of type 2 by changing all coefficients and a variable in the power series of Heun equation and its integral form in chapter 2. Also generating functions for the type 2 Lamé polynomial in Weierstrass's form of the first and second kinds is obtained by changing all coefficients and a variable of generating functions for the type 2 Heun polynomial of the first and second kinds in chapter 3.

The Frobenius solutions and its integral forms of Lamé equation in Weierstrass's form for infinite series about $\xi = 0$ in this chapter are equivalent to the infinite series of Lamé equation in Ref.[10]. In this chapter B_n is the leading term in sequence c_n in an analytic function $y(\xi)$. In Ref.[10] A_n is the leading term in sequence c_n in an analytic function $y(\xi)$.

As we see the power series expansions of Heun, Confluent Heun, GCH, Mathieu, Lamé equations for either polynomial or infinite series in series "Special functions and three term recurrence formula (3TRF)"⁸ and every chapters in this series, the denominators and numerators in all A_n or B_n terms in these analytic solutions arise with Pochhammer symbol. Since we construct the power series expansions with Pochhammer symbols in numerators and denominators, we are able to describe integral representations of all these equations analytically. The mathematical technique to obtain integral forms of these equations is also available for any linear ordinary differential equations having three different coefficients in its power series. As we observe representations in closed form integrals of these equations by applying either 3TRF or R3TRF, a ${}_1F_1$ or ${}_2F_1$ function recurs in each of sub-integral forms of these equations (every integral forms of all these equations are composed of the definite and contour integrals). We might be possible to convert these equations into all other well-known special functions such as Legendre, Bessel, Laguerre, Kummer functions and etc because a ${}_1F_1$ or ${}_2F_1$ function recurs in each of sub-integral forms of these equations. We can rebuild other confluent forms of these equations after we replace ${}_1F_1$ or ${}_2F_1$ function in integral forms of them to other special functions. After that, we can reconstruct the power series expansion of confluent forms of these equations in a backward from those remodeled integral forms. Indeed, generating functions of all these equations are constructed analytically by applying the generating function for the Jacobi polynomial using hypergeometric functions or for confluent

⁸It is available as arXiv.

hypergeometric (Kummer's) polynomial of the first kind into the general expression of the integral representation of these equations. We can build orthogonal relations, recursion relations and expectation values of any physical quantities from these generating functions; i.e. the normalized wave function of hydrogen-like atoms and an expectation value of measurable parameters in quantum physics are constructed by applying the generating function for associated Laguerre polynomials.

Appendices

Appendix. 9 local solutions of Lamé equation in Weierstrass's form by applying R3TRF out of 192 local solutions of Heun equation

A machine-generated list of 192 (isomorphic to the Coxeter group of the Coxeter diagram D_4) local solutions of the Heun equation was obtained by Robert S. Maier(2007) [26]. In appendix of Ref.[8], I apply 3TRF to the power series expansions of Heun equation for infinite series and polynomial of type 1 and its integral forms of nine out of the 192 local solution of Heun function in Table 2 [26].

In appendix of chapters 2 and 3, by applying R3TRF, I construct the fundamental power series expansions in closed forms and integral forms of Heun equation for infinite series and polynomial of type 2 of the previous nine out of the 192 local solution of Heun function in Table 2 including generating functions for the Heun polynomial of type 2.

In appendix of Ref.[10], 9 local solutions of Lamé equation in Weierstrass's form by applying 3TRF out of 192 local solutions of Heun equation are constructed: the power series expansions of Lamé equation (for infinite series and polynomial of type 1) and its integral forms are derived analytically.

In this appendix, by changing all coefficients and independent variables of the previous nine examples of 192 local solutions of Heun function into the first kind of independent solutions of Heun equation by applying R3TRF in chapters 2 and 3, I construct 9 local solutions of Lamé equation in Weierstrass's form for Frobenius solutions in closed form (for infinite series and polynomial of type 2), its integral representations and generating functions for the type 2 polynomial.

Infinite series by applying either 3TRF or R3TRF are equivalent to each other. For infinite series by applying 3TRF, A_n is the leading term in sequence c_n in all 9 local solutions of Lamé equation in appendix of Ref.[10]. For infinite series by applying R3TRF, B_n is the leading term in sequence c_n in all 9 local solutions of it in this appendix.⁹

⁹In appendix of Ref.[10], I treat h as a free variable and a fixed value of α to construct polynomials of type 1 for all 9 local solutions of Lamé equation. In this appendix, I treat α as a free variable and a fixed value of h to construct polynomials of type 2 for all 9 local solutions of it. And an independent variable $sn^2(z, \rho)$ is denoted by ξ .

.1 Power series

.1.1 $(1-x)^{1-\delta} Hl(a, q - (\delta-1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x)$

Polynomial of type 2

Replace coefficients $q, \alpha, \beta, \delta, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\delta-1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $(1-x)^{1-\delta}$ and (9.2.1) together. Put (9.1.6) into the new (9.2.1).¹⁰

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}} y(\xi) \\
&= (1-\xi)^{\frac{1}{2}} Hl \left(\rho^{-2}, -\frac{1}{4}(h-1)\rho^{-2} = -\rho^{-2}(h_j+2j)[1+(1+\rho^2)(h_j+2j)]; \frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}; \frac{1}{2}, \frac{3}{2}; \xi \right) \\
&= (1-\xi)^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0+1+\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{3}{2})} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1+4+\Omega_\rho)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-h_1)_{i_0} (h_1+4+\Omega_\rho)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0+1+\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{3}{2})} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k+2k+1+\frac{\alpha}{2})(i_k+2k+\frac{1}{2}-\frac{\alpha}{2})}{(i_k+2k+2)(i_k+2k+\frac{3}{2})} \right. \\
&\times \frac{(-h_k)_{i_k} (h_k+4k+\Omega_\rho)_{i_k} (2k+1)_{i_{k-1}} (2k+\frac{1}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k+4k+\Omega_\rho)_{i_{k-1}} (2k+1)_{i_k} (2k+\frac{1}{2})_{i_k}} \left. \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n+4n+\Omega_\rho)_{i_n} (2n+1)_{i_{n-1}} (2n+\frac{1}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n+4n+\Omega_\rho)_{i_{n-1}} (2n+1)_{i_n} (2n+\frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.1)
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{1}{1+\rho^2} \\ h = 4(h_j+2j)[1+(1+\rho^2)(h_j+2j)] + 1 \end{cases}$$

¹⁰For all 9 local solutions of Lamé equation for polynomial of type 2 in this appendix, $h_i \leq h_j$ only if $i \leq j$ where $i, j, h_i, h_j \in \mathbb{N}_0$.

Infinite series

Replace coefficients $q, \alpha, \beta, \delta, c_0$ and λ by $q - (\delta - 1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$ and zero into (9.2.4). Multiply $(1 - x)^{1-\delta}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
 & (1 - \xi)^{\frac{1}{2}} y(\xi) \\
 = & (1 - \xi)^{\frac{1}{2}} Hl \left(\rho^{-2}, -\frac{1}{4}(h-1)\rho^{-2}; \frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}; \frac{1}{2}, \frac{3}{2}; \xi \right) \\
 = & (1 - \xi)^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \right. \\
 & + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \frac{\alpha}{2}) (i_0 + \frac{1}{2} - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
 & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + 1 + \frac{\alpha}{2}) (i_0 + \frac{1}{2} - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
 & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + 1 + \frac{\alpha}{2}) (i_k + 2k + \frac{1}{2} - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
 & \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.2)
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{1 \pm \sqrt{1 + (1 + \rho^2)(h-1)}}{2(1 + \rho^2)} \\ \Delta_1^{\pm} = 2 + \frac{1 \pm \sqrt{1 + (1 + \rho^2)(h-1)}}{2(1 + \rho^2)} \\ \Delta_k^{\pm} = 2k + \frac{1 \pm \sqrt{1 + (1 + \rho^2)(h-1)}}{2(1 + \rho^2)} \\ \Delta_n^{\pm} = 2n + \frac{1 \pm \sqrt{1 + (1 + \rho^2)(h-1)}}{2(1 + \rho^2)} \end{cases}$$

On (.1.1) and (.1.2),

$$\begin{cases} \eta = (1 + \rho^2)\xi \\ z = -\rho^2\xi^2 \end{cases}$$

.1.2

$$x^{1-\gamma}(1-x)^{1-\delta}Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x)$$

Polynomial of type 2

Replace coefficients $q, \alpha, \beta, \gamma, \delta, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and (9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
& \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}y(\xi) \\
&= \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}Hl\left(\rho^{-2}, -\frac{1}{4}((h-4)\rho^{-2}-1) = -\rho^{-2}(h_j+2j)[2+\rho^2+(1+\rho^2)(h_j+2j)]\right. \\
&\quad \left.; \frac{\alpha}{2} + \frac{3}{2}, -\frac{\alpha}{2} + 1; \frac{3}{2}, \frac{3}{2}; \xi\right) \\
&= \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{3}{2}\right)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{3}{2} + \frac{\alpha}{2}) (i_0 + 1 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{5}{2})} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{3}{2}\right)_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4 + \Omega_\rho)_{i_1} (3)_{i_0} \left(\frac{7}{2}\right)_{i_0}}{(-h_1)_{i_0} (h_1 + 4 + \Omega_\rho)_{i_0} (3)_{i_1} \left(\frac{7}{2}\right)_{i_1}} \eta^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{3}{2} + \frac{\alpha}{2}) (i_0 + 1 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{5}{2})} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{3}{2}\right)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{3}{2} + \frac{\alpha}{2}) (i_k + 2k + 1 - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{5}{2})} \right. \\
&\quad \times \left. \frac{(-h_k)_{i_k} (h_k + 4k + \Omega_\rho)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{3}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k + \Omega_\rho)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{3}{2})_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n + \Omega_\rho)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{3}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n + \Omega_\rho)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{3}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.3)
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{2+\rho^2}{1+\rho^2} \\ h = 4 + \rho^2 + 4(h_j + 2j)[2 + \rho^2 + (1 + \rho^2)(h_j + 2j)] \end{cases}$$

Infinite series

Replace coefficients $q, \alpha, \beta, \gamma, \delta, c_0$ and λ by $q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1)$, $\alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$ and zero into (9.2.4). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
 & \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}y(\xi) \\
 = & \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}Hl\left(\rho^{-2}, -\frac{1}{4}((h-4)\rho^{-2}-1); \frac{\alpha}{2} + \frac{3}{2}, -\frac{\alpha}{2} + 1; \frac{3}{2}, \frac{3}{2}; \xi\right) \\
 = & \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}\left\{\sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{3}{2})_{i_0}} \eta^{i_0} \right. \\
 & + \left\{\sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{3}{2} + \frac{\alpha}{2})(i_0 + 1 - \frac{\alpha}{2})}{(i_0 + 2)(i_0 + \frac{5}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{3}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\frac{7}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\frac{7}{2})_{i_1}} \eta^{i_1}\right\} z \\
 & + \sum_{n=2}^{\infty} \left\{\sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{3}{2} + \frac{\alpha}{2})(i_0 + 1 - \frac{\alpha}{2})}{(i_0 + 2)(i_0 + \frac{5}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{3}{2})_{i_0}} \right. \\
 & \times \prod_{k=1}^{n-1} \left\{\sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{3}{2} + \frac{\alpha}{2})(i_k + 2k + 1 - \frac{\alpha}{2})}{(i_k + 2k + 2)(i_k + 2k + \frac{5}{2})} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{3}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{3}{2})_{i_k}} \right\} \\
 & \times \left.\sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{3}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{3}{2})_{i_n}} \eta^{i_n}\right\} z^n \Bigg\} \quad (.1.4)
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{2+\rho^2 \pm \sqrt{(h-1)\rho^2+h}}{2(1+\rho^2)} \\ \Delta_1^{\pm} = 2 + \frac{2+\rho^2 \pm \sqrt{(h-1)\rho^2+h}}{2(1+\rho^2)} \\ \Delta_k^{\pm} = 2k + \frac{2+\rho^2 \pm \sqrt{(h-1)\rho^2+h}}{2(1+\rho^2)} \\ \Delta_n^{\pm} = 2n + \frac{2+\rho^2 \pm \sqrt{(h-1)\rho^2+h}}{2(1+\rho^2)} \end{cases}$$

On (.1.3) and (.1.4),

$$\begin{cases} \eta = (1 + \rho^2)\xi \\ z = -\rho^2\xi^2 \end{cases}$$

$$\mathbf{.1.3} \quad Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x)$$

Polynomial of type 2

Replace coefficients $a, q, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1-a, -q + \alpha\beta, \delta, \gamma, 1-x, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
y(\varsigma) &= Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha(\alpha+1)) = -(2 - \rho^{-2})(h_j + 2j)^2; \frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; 1 - \xi \right) \\
&= \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \eta^{i_0} \\
&+ \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{(-h_1)_{i_0} (h_1 + 4)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(-h_k)_{i_k} (h_k + 4k)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \quad (1.5)
\end{aligned}$$

where

$$h = \rho^2 (\alpha(\alpha+1) - 4(2 - \rho^{-2})(h_j + 2j)^2)$$

Infinite series

Replace coefficients $a, q, \gamma, \delta, x, c_0$ and λ by $1 - a, -q + \alpha\beta, \delta, \gamma, 1 - x, 1$ and zero into (9.2.4). Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
 y(\varsigma) &= Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha(\alpha + 1)); \frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; 1 - \xi \right) \\
 &= \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \\
 &+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
 &+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
 &\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
 &\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \quad (.1.6)
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \pm \frac{1}{2} \sqrt{\frac{h\rho^{-2} + \alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Delta_1^{\pm} = 2 \pm \frac{1}{2} \sqrt{\frac{h\rho^{-2} + \alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Delta_k^{\pm} = 2k \pm \frac{1}{2} \sqrt{\frac{h\rho^{-2} + \alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Delta_n^{\pm} = 2n \pm \frac{1}{2} \sqrt{\frac{h\rho^{-2} + \alpha(\alpha+1)}{2-\rho^{-2}}} \end{cases}$$

On (.1.5) and (.1.6),

$$\begin{cases} \varsigma = 1 - \xi \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}} \varsigma \\ z = \frac{-1}{1-\rho^{-2}} \varsigma^2 \end{cases}$$

$$\mathbf{.1.4} \quad (1-x)^{1-\delta} Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x)$$

Polynomial of type 2

Replace coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma, 1-x, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $(1-x)^{1-\delta}$ and (9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}} y(\varsigma) \\
&= (1-\xi)^{\frac{1}{2}} Hl \left(1-\rho^{-2}, -\frac{1}{4}((1+h)\rho^{-2} + (\alpha-1)(\alpha+2)) = -(2-\rho^{-2})(h_j+2j)(h_j+2j+1) \right. \\
&\quad \left. ; \frac{\alpha}{2}+1, -\frac{\alpha}{2}+\frac{1}{2}, \frac{3}{2}, \frac{1}{2}; 1-\xi \right) \\
&= (1-\xi)^{\frac{1}{2}} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0+1)_{i_0}}{(1)_{i_0} \left(\frac{3}{2}\right)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0+1+\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{5}{2})} \frac{(-h_0)_{i_0} (h_0+1)_{i_0}}{(1)_{i_0} \left(\frac{3}{2}\right)_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1+5)_{i_1} (3)_{i_0} \left(\frac{7}{2}\right)_{i_0}}{(-h_1)_{i_0} (h_1+5)_{i_0} (3)_{i_1} \left(\frac{7}{2}\right)_{i_1}} \eta^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0+1+\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{5}{2})} \frac{(-h_0)_{i_0} (h_0+1)_{i_0}}{(1)_{i_0} \left(\frac{3}{2}\right)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k+2k+1+\frac{\alpha}{2})(i_k+2k+\frac{1}{2}-\frac{\alpha}{2})}{(i_k+2k+2)(i_k+2k+\frac{5}{2})} \right. \\
&\quad \times \left. \frac{(-h_k)_{i_k} (h_k+4k+1)_{i_k} (2k+1)_{i_{k-1}} (2k+\frac{3}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k+4k+1)_{i_{k-1}} (2k+1)_{i_k} (2k+\frac{3}{2})_{i_k}} \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n+4n+1)_{i_n} (2n+1)_{i_{n-1}} (2n+\frac{3}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n+4n+1)_{i_{n-1}} (2n+1)_{i_n} (2n+\frac{3}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.7)
\end{aligned}$$

where

$$h = 4(2\rho^2 - 1)(h_j + 2j)(h_j + 2j + 1) - \rho^2(\alpha - 1)(\alpha + 2) - 1$$

Infinite series

Replace coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0$ and λ by $1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma, 1-x, 1$ and zero

into (9.2.4). Multiply $(1-x)^{1-\delta}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}}y(\varsigma) \\
&= (1-\xi)^{\frac{1}{2}}Hl\left(1-\rho^{-2}, -\frac{1}{4}\left((1+h)\rho^{-2}+(\alpha-1)(\alpha+2)\right); \frac{\alpha}{2}+1, -\frac{\alpha}{2}+\frac{1}{2}, \frac{3}{2}, \frac{1}{2}; 1-\xi\right) \\
&= (1-\xi)^{\frac{1}{2}}\left\{\sum_{i_0=0}^{\infty}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}\left(\frac{3}{2}\right)_{i_0}}\eta^{i_0}\right. \\
&+ \left\{\sum_{i_0=0}^{\infty}\frac{(i_0+1+\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{5}{2})}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}\left(\frac{3}{2}\right)_{i_0}}\sum_{i_1=i_0}^{\infty}\frac{(\Delta_1^-)_{i_1}(\Delta_1^+)_{i_1}(3)_{i_0}\left(\frac{7}{2}\right)_{i_0}}{(\Delta_1^-)_{i_0}(\Delta_1^+)_{i_0}(3)_{i_1}\left(\frac{7}{2}\right)_{i_1}}\eta^{i_1}\right\}z \\
&+ \sum_{n=2}^{\infty}\left\{\sum_{i_0=0}^{\infty}\frac{(i_0+1+\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{5}{2})}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}\left(\frac{3}{2}\right)_{i_0}}\right. \\
&\times \prod_{k=1}^{n-1}\left\{\sum_{i_k=i_{k-1}}^{\infty}\frac{(i_k+2k+1+\frac{\alpha}{2})(i_k+2k+\frac{1}{2}-\frac{\alpha}{2})}{(i_k+2k+2)(i_k+2k+\frac{5}{2})}\frac{(\Delta_k^-)_{i_k}(\Delta_k^+)_{i_k}(2k+1)_{i_{k-1}}(2k+\frac{3}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}}(\Delta_k^+)_{i_{k-1}}(2k+1)_{i_k}(2k+\frac{3}{2})_{i_k}}\right\} \\
&\times \left.\sum_{i_n=i_{n-1}}^{\infty}\frac{(\Delta_n^-)_{i_n}(\Delta_n^+)_{i_n}(2n+1)_{i_{n-1}}(2n+\frac{3}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}}(\Delta_n^+)_{i_{n-1}}(2n+1)_{i_n}(2n+\frac{3}{2})_{i_n}}\eta^{i_n}\right\}z^n\Bigg\} \quad (.1.8)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{1}{2} \pm \frac{1}{2}\sqrt{\frac{h\rho^{-2}+\alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Delta_1^{\pm} = \frac{5}{2} \pm \frac{1}{2}\sqrt{\frac{h\rho^{-2}+\alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Delta_k^{\pm} = 2k + \frac{1}{2} \pm \frac{1}{2}\sqrt{\frac{h\rho^{-2}+\alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Delta_n^{\pm} = 2n + \frac{1}{2} \pm \frac{1}{2}\sqrt{\frac{h\rho^{-2}+\alpha(\alpha+1)}{2-\rho^{-2}}} \end{cases}$$

On (.1.7) and (.1.8),

$$\begin{cases} \varsigma = 1-\xi \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = \frac{-1}{1-\rho^{-2}}\varsigma^2 \end{cases}$$

$$.1.5 \quad x^{-\alpha}Hl\left(\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}; \alpha, \alpha-\gamma+1, \alpha-\beta+1, \delta; \frac{1}{x}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \beta, \gamma, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}, \alpha-\gamma+1, \alpha-\beta+1, \frac{1}{x}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $x^{-\alpha}$ and

(9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
& \xi^{\frac{1}{2}(\alpha+1)} y(\zeta) \\
= & \xi^{\frac{1}{2}(\alpha+1)} Hl \left(\rho^2, -\frac{1}{4} (h - (1 + \rho^2)(\alpha + 1)^2) = -(1 + \rho^2)(h_j + 2j)(h_j + 2j + 1 + \alpha) \right. \\
& \left. ; \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \alpha + \frac{3}{2}, \frac{1}{2}; \xi^{-1} \right) \\
= & \xi^{\frac{1}{2}(\alpha+1)} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + 1 + \alpha)_{i_0}}{(1)_{i_0} (\alpha + \frac{3}{2})_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2}) (-h_0)_{i_0} (h_0 + 1 + \alpha)_{i_0}}{(i_0 + 2) (i_0 + \frac{5}{2} + \alpha) (1)_{i_0} (\alpha + \frac{3}{2})_{i_0}} \right. \\
& \times \left. \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 5 + \alpha)_{i_1} (3)_{i_0} (\alpha + \frac{7}{2})_{i_0}}{(-h_1)_{i_0} (h_1 + 5 + \alpha)_{i_0} (3)_{i_1} (\alpha + \frac{7}{2})_{i_1}} \eta^{i_1} \right\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2}) (-h_0)_{i_0} (h_0 + 1 + \alpha)_{i_0}}{(i_0 + 2) (i_0 + \frac{5}{2} + \alpha) (1)_{i_0} (\alpha + \frac{3}{2})_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2}) (i_k + 2k + 1 - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{5}{2} + \alpha)} \right. \\
& \times \left. \frac{(-h_k)_{i_k} (h_k + 4k + 1 + \alpha)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{3}{2} + \alpha)_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k + 1 + \alpha)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{3}{2} + \alpha)_{i_k}} \right\} \\
& \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n + 1 + \alpha)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{3}{2} + \alpha)_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n + 1 + \alpha)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{3}{2} + \alpha)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.9)
\end{aligned}$$

where

$$h = (1 + \rho^2) (2(h_j + 2j) + 1 + \alpha)^2$$

Infinite series

Replace coefficients $a, q, \beta, \gamma, x, c_0$ and λ by $\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}, \alpha - \gamma + 1, \alpha - \beta + 1, \frac{1}{x}, 1$ and zero into (9.2.4). Multiply $x^{-\alpha}$ and (9.2.4) together. Put (9.1.6) into

the new (9.2.4).

$$\begin{aligned}
& \xi^{\frac{1}{2}(\alpha+1)} y(\varsigma) \\
&= \xi^{\frac{1}{2}(\alpha+1)} Hl \left(\rho^2, -\frac{1}{4} (h - (1 + \rho^2)(\alpha + 1)^2); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \alpha + \frac{3}{2}, \frac{1}{2}; \xi^{-1} \right) \\
&= \xi^{\frac{1}{2}(\alpha+1)} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\alpha + \frac{3}{2})_{i_0}} \eta^{i_0} \right. \\
&+ \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{5}{2} + \alpha)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\alpha + \frac{3}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\alpha + \frac{7}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\alpha + \frac{7}{2})_{i_1}} \eta^{i_1} \right\} z \\
&+ \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{5}{2} + \alpha)} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\alpha + \frac{3}{2})_{i_0}} \right. \\
&\times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2}) (i_k + 2k + 1 + \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{5}{2} + \alpha)} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{3}{2} + \alpha)_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{3}{2} + \alpha)_{i_k}} \right\} \\
&\times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{3}{2} + \alpha)_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{3}{2} + \alpha)_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.10)
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{1}{2}(\alpha + 1) \pm \frac{1}{2}\sqrt{\frac{h}{1+\rho^2}} \\ \Delta_1^{\pm} = \frac{1}{2}(\alpha + 5) \pm \frac{1}{2}\sqrt{\frac{h}{1+\rho^2}} \\ \Delta_k^{\pm} = \frac{1}{2}(\alpha + 4k + 1) \pm \frac{1}{2}\sqrt{\frac{h}{1+\rho^2}} \\ \Delta_n^{\pm} = \frac{1}{2}(\alpha + 4n + 1) \pm \frac{1}{2}\sqrt{\frac{h}{1+\rho^2}} \end{cases}$$

On (.1.9) and (.1.10),

$$\begin{cases} \varsigma = \xi^{-1} \\ \eta = (1 + \rho^{-2})\varsigma \\ z = -\rho^{-2}\varsigma^2 \end{cases}$$

$$.1.6 \quad \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \alpha, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1 - a, -q + \gamma\beta, -\alpha + \gamma + \delta, \frac{(1-a)x}{x-a}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $(1 - \frac{x}{a})^{-\beta}$ and

(9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
& (1 - \rho^2 \xi)^{\frac{\alpha}{2}} y(\varsigma) \\
= & (1 - \rho^2 \xi)^{\frac{\alpha}{2}} Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha) = (h_j + 2j) (\alpha - (2 - \rho^{-2})(h_j + 2j)) \right. \\
& \left. ; -\frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; \frac{(1 - \rho^{-2})\xi}{\xi - \rho^{-2}} \right) \\
= & (1 - \rho^2 \xi)^{\frac{\alpha}{2}} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 - \Omega_\rho)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} - \frac{\alpha}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0 - \Omega_\rho)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4 - \Omega_\rho)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(-h_1)_{i_0} (h_1 + 4 - \Omega_\rho)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} - \frac{\alpha}{2}) (i_0 - \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0 - \Omega_\rho)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{1}{2} - \frac{\alpha}{2}) (i_k + 2k - \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \frac{(-h_k)_{i_k} (h_k + 4k - \Omega_\rho)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k - \Omega_\rho)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
& \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n - \Omega_\rho)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n - \Omega_\rho)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.11)
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{\alpha}{2 - \rho^{-2}} \\ h = 4\rho^2 \left(\alpha (h_j + 2j + \frac{1}{4}) - (2 - \rho^{-2})(h_j + 2j)^2 \right) \end{cases}$$

Infinite series

Replace coefficients a, q, α, x, c_0 and λ by $1-a, -q+\gamma\beta, -\alpha+\gamma+\delta, \frac{(1-a)x}{x-a}, 1$ and zero into (9.2.4). Multiply $(1-\frac{x}{a})^{-\beta}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
& (1-\rho^2\xi)^{\frac{\alpha}{2}}y(\varsigma) \\
&= (1-\rho^2\xi)^{\frac{\alpha}{2}}Hl\left(1-\rho^{-2}, \frac{1}{4}(h\rho^{-2}-\alpha); -\frac{\alpha}{2}+\frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}}\right) \\
&= (1-\rho^2\xi)^{\frac{\alpha}{2}}\left\{\sum_{i_0=0}^{\infty}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(\frac{1}{2})_{i_0}}\eta^{i_0}\right. \\
&+ \left\{\sum_{i_0=0}^{\infty}\frac{(i_0+\frac{1}{2}-\frac{\alpha}{2})(i_0-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{3}{2})}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(\frac{1}{2})_{i_0}}\sum_{i_1=i_0}^{\infty}\frac{(\Delta_1^-)_{i_1}(\Delta_1^+)_{i_1}(3)_{i_0}(\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0}(\Delta_1^+)_{i_0}(3)_{i_1}(\frac{5}{2})_{i_1}}\eta^{i_1}\right\}z \\
&+ \sum_{n=2}^{\infty}\left\{\sum_{i_0=0}^{\infty}\frac{(i_0+\frac{1}{2}-\frac{\alpha}{2})(i_0-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{3}{2})}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(\frac{1}{2})_{i_0}}\right. \\
&\times \prod_{k=1}^{n-1}\left\{\sum_{i_k=i_{k-1}}^{\infty}\frac{(i_k+2k+\frac{1}{2}-\frac{\alpha}{2})(i_k+2k-\frac{\alpha}{2})}{(i_k+2k+2)(i_k+2k+\frac{3}{2})}\frac{(\Delta_k^-)_{i_k}(\Delta_k^+)_{i_k}(2k+1)_{i_{k-1}}(2k+\frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}}(\Delta_k^+)_{i_{k-1}}(2k+1)_{i_k}(2k+\frac{1}{2})_{i_k}}\right\} \\
&\times \left.\sum_{i_n=i_{n-1}}^{\infty}\frac{(\Delta_n^-)_{i_n}(\Delta_n^+)_{i_n}(2n+1)_{i_{n-1}}(2n+\frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}}(\Delta_n^+)_{i_{n-1}}(2n+1)_{i_n}(2n+\frac{1}{2})_{i_n}}\eta^{i_n}\right\}z^n\Bigg\} \tag{.1.12}
\end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{-\alpha \pm \sqrt{\alpha^2 - (2-\rho^{-2})(h\rho^{-2}-\alpha)}}{2(2-\rho^{-2})} \\ \Delta_1^{\pm} = 2 + \frac{-\alpha \pm \sqrt{\alpha^2 - (2-\rho^{-2})(h\rho^{-2}-\alpha)}}{2(2-\rho^{-2})} \\ \Delta_k^{\pm} = 2k + \frac{-\alpha \pm \sqrt{\alpha^2 - (2-\rho^{-2})(h\rho^{-2}-\alpha)}}{2(2-\rho^{-2})} \\ \Delta_n^{\pm} = 2n + \frac{-\alpha \pm \sqrt{\alpha^2 - (2-\rho^{-2})(h\rho^{-2}-\alpha)}}{2(2-\rho^{-2})} \end{cases}$$

On (.1.11) and (.1.12),

$$\begin{cases} \varsigma = \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}} \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = -\frac{1}{1-\rho^{-2}}\varsigma^2 \end{cases}$$

.1.7

$$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1} Hl \left(1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma, 2 - \delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \alpha, \beta, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1-a, -q + \gamma[(\delta-1)a + \beta - \delta + 1], -\alpha + \gamma + 1, \beta - \delta + 1, 2 - \delta, \frac{(1-a)x}{x-a}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}$ and (9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}} (1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)} y(\varsigma) \\
= & (1-\xi)^{\frac{1}{2}} (1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)} Hl \left(1-\rho^{-2}, \frac{1}{4}((h-1)\rho^{-2} + 1 - \alpha)\right. \\
& \left. = (h_j + 2j)(\alpha + 1 - (2 - \rho^{-2})(h_j + 2j + 1)); -\frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}, \frac{1}{2}, \frac{3}{2}; \frac{(1-\rho^{-2})\xi}{\xi - \rho^{-2}}\right) \\
= & (1-\xi)^{\frac{1}{2}} (1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \eta^{i_0} \right. \\
& + \left\{ \sum_{i_0=0}^{h_0} \frac{\left(i_0 + 1 - \frac{\alpha}{2}\right) \left(i_0 + \frac{1}{2} - \frac{\alpha}{2}\right) (-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(i_0 + 2) \left(i_0 + \frac{3}{2}\right) (1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \right. \\
& \times \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4 + \Omega_\rho)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{(-h_1)_{i_0} (h_1 + 4 + \Omega_\rho)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} \eta^{i_1} \Big\} z \\
& + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{\left(i_0 + 1 - \frac{\alpha}{2}\right) \left(i_0 + \frac{1}{2} - \frac{\alpha}{2}\right) (-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(i_0 + 2) \left(i_0 + \frac{3}{2}\right) (1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \right. \\
& \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{\left(i_k + 2k + 1 - \frac{\alpha}{2}\right) \left(i_k + 2k + \frac{1}{2} - \frac{\alpha}{2}\right)}{(i_k + 2k + 2) \left(i_k + 2k + \frac{3}{2}\right)} \right. \\
& \times \frac{(-h_k)_{i_k} (h_k + 4k + \Omega_\rho)_{i_k} (2k + 1)_{i_{k-1}} \left(2k + \frac{1}{2}\right)_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k + \Omega_\rho)_{i_{k-1}} (2k + 1)_{i_k} \left(2k + \frac{1}{2}\right)_{i_k}} \Big\} \\
& \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n + \Omega_\rho)_{i_n} (2n + 1)_{i_{n-1}} \left(2n + \frac{1}{2}\right)_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n + \Omega_\rho)_{i_{n-1}} (2n + 1)_{i_n} \left(2n + \frac{1}{2}\right)_{i_n}} \eta^{i_n} \right\} z^n \Big\} \quad (.1.13)
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{-\alpha+1-\rho^{-2}}{2-\rho^{-2}} \\ h = 4(h_j + 2j) ((\alpha + 1)\rho^2 + (1 - 2\rho^2)(h_j + 2j + 1)) + (\alpha - 1)\rho^2 + 1 \end{cases}$$

Infinite series

Replace coefficients $a, q, \alpha, \beta, \delta, x, c_0$ and λ by $1-a, -q+\gamma[(\delta-1)a+\beta-\delta+1], -\alpha+\gamma+1, \beta-\delta+1, 2-\delta, \frac{(1-a)x}{x-a}, 1$ and zero into (9.2.4). Multiply $(1-x)^{1-\delta} (1-\frac{x}{a})^{-\beta+\delta-1}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
 & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}y(\varsigma) \\
 = & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}Hl\left(1-\rho^{-2}, \frac{1}{4}\left((h-1)\rho^{-2}+1-\alpha\right); -\frac{\alpha}{2}+1, -\frac{\alpha}{2}+\frac{1}{2}, \frac{1}{2}, \frac{3}{2}; \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}}\right) \\
 = & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}\left\{\sum_{i_0=0}^{\infty}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(\frac{1}{2})_{i_0}}\eta^{i_0}\right. \\
 & + \left\{\sum_{i_0=0}^{\infty}\frac{(i_0+1-\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{3}{2})}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(\frac{1}{2})_{i_0}}\sum_{i_1=i_0}^{\infty}\frac{(\Delta_1^-)_{i_1}(\Delta_1^+)_{i_1}(3)_{i_0}(\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0}(\Delta_1^+)_{i_0}(3)_{i_1}(\frac{5}{2})_{i_1}}\eta^{i_1}\right\}z \\
 & + \sum_{n=2}^{\infty}\left\{\sum_{i_0=0}^{\infty}\frac{(i_0+1-\frac{\alpha}{2})(i_0+\frac{1}{2}-\frac{\alpha}{2})}{(i_0+2)(i_0+\frac{3}{2})}\frac{(\Delta_0^-)_{i_0}(\Delta_0^+)_{i_0}}{(1)_{i_0}(\frac{1}{2})_{i_0}}\right. \\
 & \times \prod_{k=1}^{n-1}\left\{\sum_{i_k=i_{k-1}}^{\infty}\frac{(i_k+2k+1-\frac{\alpha}{2})(i_k+2k+\frac{1}{2}-\frac{\alpha}{2})}{(i_k+2k+2)(i_k+2k+\frac{3}{2})}\frac{(\Delta_k^-)_{i_k}(\Delta_k^+)_{i_k}(2k+1)_{i_{k-1}}(2k+\frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}}(\Delta_k^+)_{i_{k-1}}(2k+1)_{i_k}(2k+\frac{1}{2})_{i_k}}\right\} \\
 & \times \left.\sum_{i_n=i_{n-1}}^{\infty}\frac{(\Delta_n^-)_{i_n}(\Delta_n^+)_{i_n}(2n+1)_{i_{n-1}}(2n+\frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}}(\Delta_n^+)_{i_{n-1}}(2n+1)_{i_n}(2n+\frac{1}{2})_{i_n}}\eta^{i_n}\right\}z^n\Bigg\} \tag{.1.14}
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{-\alpha+1-\rho^{-2}\pm\sqrt{(-\alpha+1-\rho^{-2})^2-(2-\rho^{-2})((h-1)\rho^{-2}+1-\alpha)}}{2(2-\rho^{-2})} \\ \Delta_1^{\pm} = 2 + \frac{-\alpha+1-\rho^{-2}\pm\sqrt{(-\alpha+1-\rho^{-2})^2-(2-\rho^{-2})((h-1)\rho^{-2}+1-\alpha)}}{2(2-\rho^{-2})} \\ \Delta_k^{\pm} = 2k + \frac{-\alpha+1-\rho^{-2}\pm\sqrt{(-\alpha+1-\rho^{-2})^2-(2-\rho^{-2})((h-1)\rho^{-2}+1-\alpha)}}{2(2-\rho^{-2})} \\ \Delta_n^{\pm} = 2n + \frac{-\alpha+1-\rho^{-2}\pm\sqrt{(-\alpha+1-\rho^{-2})^2-(2-\rho^{-2})((h-1)\rho^{-2}+1-\alpha)}}{2(2-\rho^{-2})} \end{cases}$$

On (.1.13) and (.1.14),

$$\begin{cases} \varsigma = \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}} \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = -\frac{1}{1-\rho^{-2}}\varsigma^2 \end{cases}$$

$$\mathbf{.1.8} \quad x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}; \alpha, \alpha - \gamma + 1, \delta, \alpha - \beta + 1; \frac{x-1}{x} \right)$$

Polynomial of type 2

Replace coefficients $a, q, \beta, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}, \alpha - \gamma + 1, \delta, \alpha - \beta + 1, \frac{x-1}{x}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $x^{-\alpha}$ and (9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned} & \xi^{-\frac{1}{2}(\alpha+1)} y(\zeta) \\ = & \xi^{-\frac{1}{2}(\alpha+1)} Hl \left(1 - \rho^2, \frac{1}{4} [h + (\alpha + 1) (1 - \rho^2(\alpha + 1))] \right) \\ & = -(h_j + 2j) ((1 - \rho^2)(\alpha + 1) + (2 - \rho^2)(h_j + 2j)); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \alpha + \frac{3}{2}; \frac{\xi - 1}{\xi} \Big) \\ = & \xi^{-\frac{1}{2}(\alpha+1)} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \eta^{i_0} \right. \\ & + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4 + \Omega_\rho)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{(-h_1)_{i_0} (h_1 + 4 + \Omega_\rho)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} \eta^{i_1} \right\} z \\ & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2) (i_0 + \frac{3}{2})} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \right. \\ & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2}) (i_k + 2k + 1 + \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \right. \\ & \times \frac{(-h_k)_{i_k} (h_k + 4k + \Omega_\rho)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k + \Omega_\rho)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \Big\} \\ & \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n + \Omega_\rho)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n + \Omega_\rho)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Big\} \end{aligned} \quad (.1.15)$$

where

$$\begin{cases} \Omega_\rho = \frac{(\alpha+1)(1-\rho^2)}{2-\rho^2} \\ h = -4(h_j + 2j) ((1 - \rho^2)(\alpha + 1) + (2 - \rho^2)(h_j + 2j)) - (\alpha + 1) (1 - \rho^2(\alpha + 1)) \end{cases}$$

Infinite series

Replace coefficients $a, q, \beta, \gamma, \delta, x, c_0$ and λ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha-\gamma+1, \delta, \alpha-\beta+1, \frac{x-1}{x}, 1$ and zero into (9.2.4). Multiply $x^{-\alpha}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
 & \xi^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
 = & \xi^{-\frac{1}{2}(\alpha+1)} Hl \left(1 - \rho^2, \frac{1}{4} [h + (\alpha+1)(1 - \rho^2(\alpha+1))] ; \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \alpha + \frac{3}{2}; \frac{\xi-1}{\xi} \right) \\
 = & \xi^{-\frac{1}{2}(\alpha+1)} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \right. \\
 & + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2})(i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2)(i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
 & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2})(i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2)(i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
 & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2})(i_k + 2k + 1 + \frac{\alpha}{2})}{(i_k + 2k + 2)(i_k + 2k + \frac{3}{2})} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k+1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k+1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
 & \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n+1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n+1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.16)
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{(1-\rho^2)(\alpha+1) \pm \sqrt{(\alpha+1)^2 - (2-\rho^{-2})(h+1+\alpha)}}{2(2-\rho^2)} \\ \Delta_1^{\pm} = 2 + \frac{(1-\rho^2)(\alpha+1) \pm \sqrt{(\alpha+1)^2 - (2-\rho^{-2})(h+1+\alpha)}}{2(2-\rho^2)} \\ \Delta_k^{\pm} = 2k + \frac{(1-\rho^2)(\alpha+1) \pm \sqrt{(\alpha+1)^2 - (2-\rho^{-2})(h+1+\alpha)}}{2(2-\rho^2)} \\ \Delta_n^{\pm} = 2n + \frac{(1-\rho^2)(\alpha+1) \pm \sqrt{(\alpha+1)^2 - (2-\rho^{-2})(h+1+\alpha)}}{2(2-\rho^2)} \end{cases}$$

On (.1.15) and (.1.16),

$$\begin{cases} \varsigma = 1 - \xi^{-1} \\ \eta = \frac{2-\rho^2}{1-\rho^2} \varsigma \\ z = -\frac{1}{1-\rho^2} \varsigma^2 \end{cases}$$

$$.1.9 \quad \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right)$$

Polynomial of type 2

Replace coefficients $q, \beta, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.2.1). Multiply $\left(\frac{x-a}{1-a} \right)^{-\alpha}$ and (9.2.1) together. Put (9.1.6) into the new (9.2.1).

$$\begin{aligned}
& \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}} \right)^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
&= \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}} \right)^{-\frac{1}{2}(\alpha+1)} Hl \left(\rho^{-2}, -\frac{1}{4} (h\rho^{-2} - (\alpha+1)^2) = -(h_j + 2j) ((1 + \rho^{-2})(h_j + 2j) + \alpha + 1) \right. \\
&\quad \left. ; \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \frac{1}{2}; \frac{\xi - 1}{\rho^2(\xi - \rho^{-2})} \right) \\
&= \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}} \right)^{-\frac{1}{2}(\alpha+1)} \left\{ \sum_{i_0=0}^{h_0} \frac{(-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \eta^{i_0} \right. \\
&\quad + \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2}) (-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(i_0 + 2) (i_0 + \frac{3}{2}) (1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \sum_{i_1=i_0}^{h_1} \frac{(-h_1)_{i_1} (h_1 + 4 + \Omega_\rho)_{i_1} (3)_{i_0} \left(\frac{5}{2}\right)_{i_0}}{(-h_1)_{i_0} (h_1 + 4 + \Omega_\rho)_{i_0} (3)_{i_1} \left(\frac{5}{2}\right)_{i_1}} \eta^{i_1} \right\} z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{h_0} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2}) (i_0 + 1 + \frac{\alpha}{2}) (-h_0)_{i_0} (h_0 + \Omega_\rho)_{i_0}}{(i_0 + 2) (i_0 + \frac{3}{2}) (1)_{i_0} \left(\frac{1}{2}\right)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{h_k} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2}) (i_k + 2k + 1 + \frac{\alpha}{2})}{(i_k + 2k + 2) (i_k + 2k + \frac{3}{2})} \right. \\
&\quad \times \frac{(-h_k)_{i_k} (h_k + 4k + \Omega_\rho)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(-h_k)_{i_{k-1}} (h_k + 4k + \Omega_\rho)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \left. \right\} \\
&\quad \times \left. \sum_{i_n=i_{n-1}}^{h_n} \frac{(-h_n)_{i_n} (h_n + 4n + \Omega_\rho)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(-h_n)_{i_{n-1}} (h_n + 4n + \Omega_\rho)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (.1.17)
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{\alpha+1}{1+\rho^{-2}} \\ h = 4(h_j + 2j) ((1 + \rho^2)(h_j + 2j) + \rho^2(\alpha + 1)) + \rho^2(\alpha + 1)^2 \end{cases}$$

Infinite series

Replace coefficients $q, \beta, \gamma, \delta, x, c_0$ and λ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$ and zero into (9.2.4). Multiply $\left(\frac{x-a}{1-a}\right)^{-\alpha}$ and (9.2.4) together. Put (9.1.6) into the new (9.2.4).

$$\begin{aligned}
 & \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}}\right)^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
 = & \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}}\right)^{-\frac{1}{2}(\alpha+1)} Hl\left(\rho^{-2}, -\frac{1}{4}(h\rho^{-2} - (\alpha+1)^2); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \frac{1}{2}; \frac{\xi - 1}{\rho^2(\xi - \rho^{-2})}\right) \\
 = & \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}}\right)^{-\frac{1}{2}(\alpha+1)} \left\{ \sum_{i_0=0}^{\infty} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \eta^{i_0} \right. \\
 & + \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2})(i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2)(i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \frac{(\Delta_1^-)_{i_1} (\Delta_1^+)_{i_1} (3)_{i_0} (\frac{5}{2})_{i_0}}{(\Delta_1^-)_{i_0} (\Delta_1^+)_{i_0} (3)_{i_1} (\frac{5}{2})_{i_1}} \eta^{i_1} \right\} z \\
 & + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(i_0 + \frac{1}{2} + \frac{\alpha}{2})(i_0 + 1 + \frac{\alpha}{2})}{(i_0 + 2)(i_0 + \frac{3}{2})} \frac{(\Delta_0^-)_{i_0} (\Delta_0^+)_{i_0}}{(1)_{i_0} (\frac{1}{2})_{i_0}} \right. \\
 & \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k + 2k + \frac{1}{2} + \frac{\alpha}{2})(i_k + 2k + 1 + \frac{\alpha}{2})}{(i_k + 2k + 2)(i_k + 2k + \frac{3}{2})} \frac{(\Delta_k^-)_{i_k} (\Delta_k^+)_{i_k} (2k + 1)_{i_{k-1}} (2k + \frac{1}{2})_{i_{k-1}}}{(\Delta_k^-)_{i_{k-1}} (\Delta_k^+)_{i_{k-1}} (2k + 1)_{i_k} (2k + \frac{1}{2})_{i_k}} \right\} \\
 & \times \left. \sum_{i_n=i_{n-1}}^{\infty} \frac{(\Delta_n^-)_{i_n} (\Delta_n^+)_{i_n} (2n + 1)_{i_{n-1}} (2n + \frac{1}{2})_{i_{n-1}}}{(\Delta_n^-)_{i_{n-1}} (\Delta_n^+)_{i_{n-1}} (2n + 1)_{i_n} (2n + \frac{1}{2})_{i_n}} \eta^{i_n} \right\} z^n \Bigg\} \quad (1.18)
 \end{aligned}$$

where

$$\begin{cases} \Delta_0^{\pm} = \frac{\rho^2(\alpha+1) \pm \sqrt{h(1+\rho^2) - \rho^2(\alpha+1)^2}}{2(1+\rho^2)} \\ \Delta_1^{\pm} = 2 + \frac{\rho^2(\alpha+1) \pm \sqrt{h(1+\rho^2) - \rho^2(\alpha+1)^2}}{2(1+\rho^2)} \\ \Delta_k^{\pm} = 2k + \frac{\rho^2(\alpha+1) \pm \sqrt{h(1+\rho^2) - \rho^2(\alpha+1)^2}}{2(1+\rho^2)} \\ \Delta_n^{\pm} = 2n + \frac{\rho^2(\alpha+1) \pm \sqrt{h(1+\rho^2) - \rho^2(\alpha+1)^2}}{2(1+\rho^2)} \end{cases}$$

On (1.17) and (1.18),

$$\begin{cases} \varsigma = \frac{\xi-1}{\rho^2(\xi-\rho^{-2})} \\ \eta = (1 + \rho^2)\varsigma \\ z = -\rho^2\varsigma^2 \end{cases}$$

.2 Integral representation

$$.2.1 \quad (1-x)^{1-\delta} Hl(a, q - (\delta - 1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x)$$

Polynomial of type 2

Replace coefficients $q, \alpha, \beta, \delta, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\delta - 1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $(1-x)^{1-\delta}$ and (9.3.1) together. Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}} y(\xi) \\
= & (1-\xi)^{\frac{1}{2}} Hl \left(\rho^{-2}, -\frac{1}{4}(h-1)\rho^{-2} = -\rho^{-2}(h_j+2j)[1+(1+\rho^2)(h_j+2j)]; \frac{\alpha}{2}+1, -\frac{\alpha}{2}+\frac{1}{2}; \frac{1}{2}, \frac{3}{2}; \xi \right) \\
= & (1-\xi)^{\frac{1}{2}} \left\{ {}_2F_1 \left(-h_0, h_0 + \Omega_\rho; \frac{1}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega_\rho)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \left. \right\} \\
& \times {}_2F_1 \left(-h_0, h_0 + \Omega_\rho; \frac{1}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \Bigg\} \tag{.2.1}
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{1}{1+\rho^2} \\ h = 4(h_j+2j)[1+(1+\rho^2)(h_j+2j)] + 1 \end{cases}$$

Infinite series

Replace coefficients $q, \alpha, \beta, \delta, c_0$ and λ by $q - (\delta - 1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, 1$ and zero into (9.3.4). Multiply $(1-x)^{1-\delta}$ and (9.3.4) together. Put (9.1.6) into the new

(9.3.4).

$$\begin{aligned}
& (1 - \xi)^{\frac{1}{2}} y(\xi) \\
&= (1 - \xi)^{\frac{1}{2}} Hl \left(\rho^{-2}, -\frac{1}{4}(h-1)\rho^{-2}; \frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}; \frac{1}{2}, \frac{3}{2}; \xi \right) \\
&= (1 - \xi)^{\frac{1}{2}} \left\{ {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n} (1 - t_{n-k})(1 - u_{n-k}) v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \left. \right\} \\
&\times {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \quad (.2.2)
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{-1 \pm \sqrt{1+(1+\rho^2)(h-1)}}{2(1+\rho^2)} \\ \Pi_{n-k}^{\pm} = -2(n-k) + \frac{-1 \pm \sqrt{1+(1+\rho^2)(h-1)}}{2(1+\rho^2)} \end{cases}$$

On (.2.1) and (.2.2),

$$\begin{cases} \eta = (1 + \rho^2)\xi \\ z = -\rho^2 \xi^2 \end{cases}$$

.2.2

$$x^{1-\gamma}(1-x)^{1-\delta} Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x)$$

Polynomial of type 2

Replace coefficients $q, \alpha, \beta, \gamma, \delta, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by

$q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1,$
zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$ and (9.3.1) together.

Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
& \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}y(\xi) \\
= & \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}Hl\left(\rho^{-2}, -\frac{1}{4}((h-4)\rho^{-2}-1) = -\rho^{-2}(h_j+2j)[2+\rho^2+(1+\rho^2)(h_j+2j)]\right. \\
& \left.; \frac{\alpha}{2} + \frac{3}{2}, -\frac{\alpha}{2} + 1; \frac{3}{2}, \frac{3}{2}; \xi\right) \\
= & \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}\left\{{}_2F_1\left(-h_0, h_0 + \Omega_\rho; \frac{3}{2}; \eta\right) + \sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}}\right.\right.\right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \frac{1}{1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}}\right)^{h_{n-k}} \\
& \times (1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega_\rho)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{1}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}}\right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}}\right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1-\frac{\alpha}{2}} \Big\} \\
& \left. \times {}_2F_1\left(-h_0, h_0 + \Omega_\rho; \frac{3}{2}; \overleftarrow{w}_{1,n}\right)\right\} z^n \Big\} \tag{.2.3}
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{2+\rho^2}{1+\rho^2} \\ h = 4 + \rho^2 + 4(h_j + 2j)[2 + \rho^2 + (1 + \rho^2)(h_j + 2j)] \end{cases}$$

Infinite series

Replace coefficients $q, \alpha, \beta, \gamma, \delta, c_0$ and λ by $q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1)$, $\alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta, 1$ and zero into (9.3.4). Multiply $x^{1-\gamma}(1-x)^{1-\delta}$

and (9.3.4) together. Put (9.1.6) into the new (9.3.4).

$$\begin{aligned}
& \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}y(\xi) \\
&= \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}Hl\left(\rho^{-2}, -\frac{1}{4}((h-4)\rho^{-2}-1); \frac{\alpha}{2} + \frac{3}{2}, -\frac{\alpha}{2} + 1; \frac{3}{2}, \frac{3}{2}; \xi\right) \\
&= \xi^{\frac{1}{2}}(1-\xi)^{\frac{1}{2}}\left\{{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{3}{2}; \eta\right) + \sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}}\right.\right.\right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}}\right)^{\Pi_{n-k}^+} (1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{1}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}}\right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}}\right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1-\frac{\alpha}{2}} \Big\} \\
&\times \left.{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{3}{2}; \overleftarrow{w}_{1,n}\right)\right\} z^n \Big\} \tag{.2.4}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{-(2+\rho^2)\pm\sqrt{\rho^2(h-1)+h}}{2(1+\rho^2)} \\ \Pi_{n-k}^{\pm} = -2(n-k) + \frac{-(2+\rho^2)\pm\sqrt{\rho^2(h-1)+h}}{2(1+\rho^2)} \end{cases}$$

On (.2.3) and (.2.4),

$$\begin{cases} \eta = (1+\rho^2)\xi \\ z = -\rho^2\xi^2 \end{cases}$$

$$\mathbf{.2.3} \quad Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x)$$

Polynomial of type 2

Replace coefficients $a, q, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1-a, -q + \alpha\beta, \delta, \gamma, 1-x, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
y(\varsigma) &= Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha(\alpha+1)) = -(2 - \rho^{-2})(h_j + 2j)^2; \frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; 1 - \xi \right) \\
&= {}_2F_1 \left(-h_0, h_0; \frac{1}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
&\quad \times (1 - \overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-4(n-k)} \\
&\quad \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)-\frac{\alpha}{2}} \Big\} \\
&\quad \times {}_2F_1 \left(-h_0, h_0; \frac{1}{2}; \overleftarrow{w}_{1,n} \right) \Big\} z^n \tag{.2.5}
\end{aligned}$$

where

$$h = \rho^2 (\alpha(\alpha+1) - 4(2 - \rho^{-2})(h_j + 2j)^2)$$

Infinite series

Replace coefficients $a, q, \gamma, \delta, x, c_0$ and λ by $1 - a, -q + \alpha\beta, \delta, \gamma, 1 - x, 1$ and zero into (9.3.4). Put (9.1.6) into the new (9.3.4).

$$\begin{aligned}
y(\varsigma) &= Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha(\alpha + 1)); \frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; 1 - \xi \right) \\
&= {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n} (1 - t_{n-k}) (1 - u_{n-k}) v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k-1)-\frac{\alpha}{2}} \Big\} \\
&\times {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \overleftrightarrow{w}_{1,n} \right) \Big\} z^n \Big\} \tag{.2.6}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \pm \frac{1}{2} \sqrt{\frac{h\rho^{-2} + \alpha(\alpha+1)}{1-\rho^{-2}}} \\ \Pi_{n-k}^\pm = -2(n-k) \pm \frac{1}{2} \sqrt{\frac{h\rho^{-2} + \alpha(\alpha+1)}{1-\rho^{-2}}} \end{cases}$$

On (.2.5) and (.2.6),

$$\begin{cases} \varsigma = 1 - \xi \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}} \varsigma \\ z = \frac{-1}{1-\rho^{-2}} \varsigma^2 \end{cases}$$

$$\textbf{.2.4} \quad (1-x)^{1-\delta} Hl(1-a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1); \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma; 1-x)$$

Polynomial of type 2

Replace coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1 - a, -q + (\delta-1)\gamma a + (\alpha-\delta+1)(\beta-\delta+1), \alpha-\delta+1, \beta-\delta+1, 2-\delta, \gamma, 1-x, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $(1-x)^{1-\delta}$ and (9.3.1) together. Put (9.1.6) into

the new (9.3.1).

$$\begin{aligned}
& (1 - \xi)^{\frac{1}{2}} y(\varsigma) \\
= & (1 - \xi)^{\frac{1}{2}} Hl \left(1 - \rho^{-2}, -\frac{1}{4} ((1 + h)\rho^{-2} + (\alpha - 1)(\alpha + 2)) = -(2 - \rho^{-2})(h_j + 2j)(h_j + 2j + 1) \right. \\
& \left. ; \frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}, \frac{3}{2}, \frac{1}{2}; 1 - \xi \right) \\
= & (1 - \xi)^{\frac{1}{2}} \left\{ {}_2F_1 \left(-h_0, h_0 + 1; \frac{3}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}} \right. \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-4(n-k)-1} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \left. \right\} \\
& \times {}_2F_1 \left(-h_0, h_0 + 1; \frac{3}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \quad (.2.7)
\end{aligned}$$

where

$$h = 4(2\rho^2 - 1)(h_j + 2j)(h_j + 2j + 1) - \rho^2(\alpha - 1)(\alpha + 2) - 1$$

Infinite series

Replace coefficients $a, q, \alpha, \beta, \gamma, \delta, x, c_0$ and λ by $1 - a, -q + (\delta - 1)\gamma a + (\alpha - \delta + 1)(\beta - \delta + 1), \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, \gamma, 1 - x, 1$ and zero

into (9.3.4). Multiply $(1-x)^{1-\delta}$ and (9.3.4) together. Put (9.1.6) into the new (9.3.4).

$$\begin{aligned}
 & (1-\xi)^{\frac{1}{2}}y(\varsigma) \\
 = & (1-\xi)^{\frac{1}{2}}Hl\left(1-\rho^{-2}, -\frac{1}{4}\left((1+h)\rho^{-2}+(\alpha-1)(\alpha+2)\right); \frac{\alpha}{2}+1, -\frac{\alpha}{2}+\frac{1}{2}, \frac{3}{2}, \frac{1}{2}; 1-\xi\right) \\
 = & (1-\xi)^{\frac{1}{2}}\left\{{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{3}{2}; \eta\right) + \sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}}\right.\right.\right. \\
 \times & \left.\left.\frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}}\right)^{\Pi_{n-k}^+} (1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-}\right.\right. \\
 \times & \left.\left.\overleftarrow{w}_{n-k,n}^{-(2(n-k)-1+\frac{\alpha}{2})}\left(\overleftarrow{w}_{n-k,n}\partial\overleftarrow{w}_{n-k,n}\right)\overleftarrow{w}_{n-k,n}^{\alpha+\frac{1}{2}}\left(\overleftarrow{w}_{n-k,n}\partial\overleftarrow{w}_{n-k,n}\right)\overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}}\right\}\right\} \\
 \times & \left.{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{3}{2}; \overleftarrow{w}_{1,n}\right)\right\} z^n \quad (.2.8)
 \end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = -\frac{1}{2} \pm \frac{1}{2}\sqrt{\frac{h\rho^{-2}+\alpha(\alpha+1)}{2-\rho^{-2}}} \\ \Pi_{n-k}^{\pm} = -2(n-k) - \frac{1}{2} \pm \frac{1}{2}\sqrt{\frac{h\rho^{-2}+\alpha(\alpha+1)}{2-\rho^{-2}}} \end{cases}$$

On (.2.7) and (.2.8),

$$\begin{cases} \varsigma = 1-\xi \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = \frac{-1}{1-\rho^{-2}}\varsigma^2 \end{cases}$$

$$.2.5 \quad x^{-\alpha}Hl\left(\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}; \alpha, \alpha-\gamma+1, \alpha-\beta+1, \delta; \frac{1}{x}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \beta, \gamma, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}, \alpha-\gamma+1, \alpha-\beta+1, \frac{1}{x}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $x^{-\alpha}$ and

(9.3.1) together. Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
& \xi^{\frac{1}{2}(\alpha+1)} y(\varsigma) \\
= & \xi^{\frac{1}{2}(\alpha+1)} Hl \left(\rho^2, -\frac{1}{4} (h - (1 + \rho^2)(\alpha + 1)^2) = -(1 + \rho^2)(h_j + 2j)(h_j + 2j + 1 + \alpha) \right. \\
& \left. ; \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \alpha + \frac{3}{2}, \frac{1}{2}; \xi^{-1} \right) \\
= & \xi^{\frac{1}{2}(\alpha+1)} \left\{ {}_2F_1 \left(-h_0, h_0 + \alpha + 1; \alpha + \frac{3}{2}; \eta \right) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}+\alpha} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-(4(n-k)+1+\alpha)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{-\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial_{\overleftarrow{w}_{n-k,n}} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\frac{\alpha}{2}} \left. \right\} \\
& \times {}_2F_1 \left(-h_0, h_0 + \alpha + 1; \alpha + \frac{3}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \Big\} \tag{.2.9}
\end{aligned}$$

where

$$h = (1 + \rho^2) (2(h_j + 2j) + 1 + \alpha)^2$$

Infinite series

Replace coefficients $a, q, \beta, \gamma, x, c_0$ and λ by $\frac{1}{a}, \frac{q+\alpha[(\alpha-\gamma-\delta+1)a-\beta+\delta]}{a}, \alpha - \gamma + 1, \alpha - \beta + 1, \frac{1}{x}, 1$ and zero into (9.3.4). Multiply $x^{-\alpha}$ and (9.3.4) together. Put (9.1.6) into

the new (9.3.4).

$$\begin{aligned}
& \xi^{\frac{1}{2}(\alpha+1)} y(\varsigma) \\
&= \xi^{\frac{1}{2}(\alpha+1)} Hl \left(\rho^2, -\frac{1}{4} (h - (1 + \rho^2)(\alpha + 1)^2); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \alpha + \frac{3}{2}, \frac{1}{2}; \xi^{-1} \right) \\
&= \xi^{\frac{1}{2}(\alpha+1)} \left\{ {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \alpha + \frac{3}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}+\alpha} \right. \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n} (1 - t_{n-k}) (1 - u_{n-k}) v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{-\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\frac{\alpha}{2}} \left. \right\} \\
&\times {}_2F_1 \left(-\Pi_0^+, -\Pi_0^-; \alpha + \frac{3}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \quad (.2.10)
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = -\frac{1}{2}(\alpha + 1) \pm \frac{1}{2}\sqrt{\frac{h}{1+\rho^2}} \\ \Pi_{n-k}^\pm = -2(n-k) - \frac{1}{2}(\alpha + 1) \pm \frac{1}{2}\sqrt{\frac{h}{1+\rho^2}} \end{cases}$$

On (.2.9) and (.2.10),

$$\begin{cases} \varsigma = \xi^{-1} \\ \eta = (1 + \rho^{-2})\varsigma \\ z = -\rho^{-2}\varsigma^2 \end{cases}$$

$$.2.6 \quad \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \alpha, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1 - a, -q + \gamma\beta, -\alpha + \gamma + \delta, \frac{(1-a)x}{x-a}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $(1 - \frac{x}{a})^{-\beta}$ and

(9.3.1) together. Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
& (1 - \rho^2 \xi)^{\frac{\alpha}{2}} y(\varsigma) \\
= & (1 - \rho^2 \xi)^{\frac{\alpha}{2}} Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha) = (h_j + 2j) (\alpha - (2 - \rho^{-2})(h_j + 2j)) \right. \\
& \left. ; -\frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; \frac{(1 - \rho^{-2})\xi}{\xi - \rho^{-2}} \right) \\
= & (1 - \rho^2 \xi)^{\frac{\alpha}{2}} \left\{ {}_2F_1 \left(-h_0, h_0 - \Omega_\rho; \frac{1}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-(4(n-k) - \Omega_\rho)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k) - \frac{3}{2} - \frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k-1) - \frac{\alpha}{2}} \left. \right\} \\
& \times {}_2F_1 \left(-h_0, h_0 - \Omega_\rho; \frac{1}{2}; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \quad (.2.11)
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{\alpha}{2 - \rho^{-2}} \\ h = 4\rho^2 \left(\alpha (h_j + 2j + \frac{1}{4}) - (2 - \rho^{-2})(h_j + 2j)^2 \right) \end{cases}$$

Infinite series

Replace coefficients a, q, α, x, c_0 and λ by $1-a, -q+\gamma\beta, -\alpha+\gamma+\delta, \frac{(1-a)x}{x-a}, 1$ and zero into (9.3.4). Multiply $(1-\frac{x}{a})^{-\beta}$ and (9.3.4) together. Put (9.1.6) into the new (9.3.4).

$$\begin{aligned}
& (1-\rho^2\xi)^{\frac{\alpha}{2}}y(\varsigma) \\
&= (1-\rho^2\xi)^{\frac{\alpha}{2}}Hl\left(1-\rho^{-2}, \frac{1}{4}(h\rho^{-2}-\alpha); -\frac{\alpha}{2}+\frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}}\right) \\
&= (1-\rho^2\xi)^{\frac{\alpha}{2}}\left\{{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \eta\right) + \sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}}\right.\right.\right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}}\right)^{\Pi_{n-k}^+} (1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}-\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n}\right) \overleftarrow{w}_{n-k,n}^{\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n}\right) \overleftarrow{w}_{n-k,n}^{2(n-k-1)-\frac{\alpha}{2}} \Big\} \\
&\times {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \overleftarrow{w}_{1,n}\right)\Big\} z^n \Big\} \tag{.2.12}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^{\pm} = \frac{\alpha \pm \sqrt{\alpha^2 - (2-\rho^{-2})(h\rho^{-2}-\alpha)}}{2(2-\rho^{-2})} \\ \Pi_{n-k}^{\pm} = -2(n-k) + \frac{\alpha \pm \sqrt{\alpha^2 - (2-\rho^{-2})(h\rho^{-2}-\alpha)}}{2(2-\rho^{-2})} \end{cases}$$

On (.2.11) and (.2.12),

$$\begin{cases} \varsigma = \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}} \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = -\frac{1}{1-\rho^{-2}}\varsigma^2 \end{cases}$$

.2.7

$$(1-x)^{1-\delta} \left(1-\frac{x}{a}\right)^{-\beta+\delta-1} Hl\left(1-a, -q+\gamma[(\delta-1)a+\beta-\delta+1]; -\alpha+\gamma+1, \beta-\delta+1, \gamma, 2-\delta; \frac{(1-a)x}{x-a}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \alpha, \beta, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1-a, -q+\gamma[(\delta-1)a+\beta-\delta+1], -\alpha+\gamma+1, \beta-\delta+1, 2-\delta, \frac{(1-a)x}{x-a}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $(1-x)^{1-\delta} (1-\frac{x}{a})^{-\beta+\delta-1}$ and (9.3.1) together. Put (9.1.6)

into the new (9.3.1).

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}y(\varsigma) \\
= & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}Hl\left(1-\rho^{-2},\frac{1}{4}\left((h-1)\rho^{-2}+1-\alpha\right)\right. \\
& = (h_j+2j)\left(\alpha+1-(2-\rho^{-2})(h_j+2j+1)\right);-\frac{\alpha}{2}+1,-\frac{\alpha}{2}+\frac{1}{2},\frac{1}{2},\frac{3}{2};\frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}}\Big) \\
= & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}\left\{{}_2F_1\left(-h_0,h_0+\Omega_\rho;\frac{1}{2};\eta\right)\right. \\
& +\sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1dt_{n-k}t_{n-k}^{2(n-k)-1}\int_0^1du_{n-k}u_{n-k}^{2(n-k)-\frac{3}{2}}\right.\right. \\
& \times\frac{1}{2\pi i}\oint dv_{n-k}\frac{1}{v_{n-k}}\left(\frac{v_{n-k}-1}{v_{n-k}}\frac{1}{1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k}}\right)^{h_{n-k}} \\
& \times(1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{-(4(n-k)+\Omega_\rho)} \\
& \times\overleftarrow{w}_{n-k,n}^{-(2(n-k)-1-\frac{\alpha}{2})}\left(\overleftarrow{w}_{n-k,n}\partial_{\overleftarrow{w}_{n-k,n}}\right)\overleftarrow{w}_{n-k,n}^{\frac{1}{2}}\left(\overleftarrow{w}_{n-k,n}\partial_{\overleftarrow{w}_{n-k,n}}\right)\overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}}\Big\} \\
& \times{}_2F_1\left(-h_0,h_0+\Omega_\rho;\frac{1}{2};\overleftarrow{w}_{1,n}\right)\Big\}z^n\Big\} \tag{.2.13}
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{-\alpha+1-\rho^{-2}}{2-\rho^{-2}} \\ h = 4(h_j+2j)\left((\alpha+1)\rho^2+(1-2\rho^2)(h_j+2j+1)\right)+(\alpha-1)\rho^2+1 \end{cases}$$

Infinite series

Replace coefficients $a, q, \alpha, \beta, \delta, x, c_0$ and λ by $1-a, -q+\gamma[(\delta-1)a+\beta-\delta+1], -\alpha+\gamma+1, \beta-\delta+1, 2-\delta, \frac{(1-a)x}{x-a}, 1$ and zero into (9.3.4). Multiply

$(1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{-\beta+\delta-1}$ and (9.3.4) together. Put (9.1.6) into the new (9.3.4).

$$\begin{aligned}
& (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}y(\varsigma) \\
= & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}Hl\left(1-\rho^{-2}, \frac{1}{4}\left((h-1)\rho^{-2}+1-\alpha\right); -\frac{\alpha}{2}+1, -\frac{\alpha}{2}+\frac{1}{2}, \frac{1}{2}, \frac{3}{2}; \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}}\right) \\
= & (1-\xi)^{\frac{1}{2}}(1-\rho^2\xi)^{\frac{1}{2}(\alpha-1)}\left\{{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \eta\right) + \sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}}\right.\right.\right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}}\right)^{\Pi_{n-k}^+} (1-\overleftarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-1-\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n}\right) \overleftarrow{w}_{n-k,n}^{\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n}\right) \overleftarrow{w}_{n-k,n}^{2(n-k)-\frac{3}{2}-\frac{\alpha}{2}} \Big\} \\
& \times \left.{}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \overleftarrow{w}_{1,n}\right)\right\} z^n \Big\} \tag{.2.14}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{\alpha-1+\rho^{-2} \pm \sqrt{(\alpha-1+\rho^{-2})^2 + (2-\rho^{-2})((1-h)\rho^{-2}+\alpha-1)}}{2(2-\rho^{-2})} \\ \Pi_{n-k}^\pm = -2(n-k) + \frac{\alpha-1+\rho^{-2} \pm \sqrt{(\alpha-1+\rho^{-2})^2 + (2-\rho^{-2})((1-h)\rho^{-2}+\alpha-1)}}{2(2-\rho^{-2})} \end{cases}$$

On (.2.13) and (.2.14),

$$\begin{cases} \varsigma = \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}} \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = -\frac{1}{1-\rho^{-2}}\varsigma^2 \end{cases}$$

$$\textbf{.2.8} \quad x^{-\alpha} Hl\left(\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}; \alpha, \alpha-\gamma+1, \delta, \alpha-\beta+1; \frac{x-1}{x}\right)$$

Polynomial of type 2

Replace coefficients $a, q, \beta, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha-\gamma+1, \delta, \alpha-\beta+1, \frac{x-1}{x}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $x^{-\alpha}$ and

(9.3.1) together. Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
& \xi^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
= & \xi^{-\frac{1}{2}(\alpha+1)} Hl \left(1 - \rho^2, \frac{1}{4} [h + (\alpha + 1) (1 - \rho^2(\alpha + 1))] \right) \\
& = -(h_j + 2j) ((1 - \rho^2)(\alpha + 1) + (2 - \rho^2)(h_j + 2j)); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \alpha + \frac{3}{2}; \frac{\xi - 1}{\xi} \Big) \\
= & \xi^{-\frac{1}{2}(\alpha+1)} \left\{ {}_2F_1 \left(-h_0, h_0 + \Omega_\rho; \frac{1}{2}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
& \times (1 - \overleftarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-(4(n-k) + \Omega_\rho)} \\
& \times \overleftarrow{w}_{n-k,n}^{-(2(n-k) - \frac{3}{2} + \frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{-\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k) - 1 + \frac{\alpha}{2}} \Big\} \\
& \times {}_2F_1 \left(-h_0, h_0 + \Omega_\rho; \frac{1}{2}; \overleftarrow{w}_{1,n} \right) \Big\} z^n \Big\} \tag{.2.15}
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{(\alpha+1)(1-\rho^2)}{2-\rho^2} \\ h = -4(h_j + 2j) ((1 - \rho^2)(\alpha + 1) + (2 - \rho^2)(h_j + 2j)) - (\alpha + 1) (1 - \rho^2(\alpha + 1)) \end{cases}$$

Infinite series

Replace coefficients $a, q, \beta, \gamma, \delta, x, c_0$ and λ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha - \gamma + 1, \delta, \alpha - \beta + 1, \frac{x-1}{x}, 1$ and zero into (9.3.4). Multiply $x^{-\alpha}$ and (9.3.4) together. Put (9.1.6)

into the new (9.3.4).

$$\begin{aligned}
& \xi^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
&= \xi^{-\frac{1}{2}(\alpha+1)} Hl \left(1 - \rho^2, \frac{1}{4} [h + (\alpha + 1) (1 - \rho^2(\alpha + 1))] ; \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \alpha + \frac{3}{2}, \frac{\xi - 1}{\xi} \right) \\
&= \xi^{-\frac{1}{2}(\alpha+1)} \left\{ {}_2F_1 \left(-\Pi_0^+, -\Pi_0^- ; \frac{1}{2} ; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \right)^{\Pi_{n-k}^+} (1 - \overleftarrow{w}_{n-k+1,n} (1 - t_{n-k}) (1 - u_{n-k}) v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{-\frac{1}{2}} \left(\overleftarrow{w}_{n-k,n} \partial \overleftarrow{w}_{n-k,n} \right) \overleftarrow{w}_{n-k,n}^{2(n-k)-1+\frac{\alpha}{2}} \left. \right\} \\
&\times {}_2F_1 \left(-\Pi_0^+, -\Pi_0^- ; \frac{1}{2} ; \overleftarrow{w}_{1,n} \right) \left. \right\} z^n \quad (.2.16)
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-(1-\rho^2)(\alpha+1) \pm \sqrt{(\alpha+1)^2 - (2-\rho^2)(h+\alpha+1)}}{2(2-\rho^2)} \\ \Pi_{n-k}^\pm = -2(n-k) + \frac{-(1-\rho^2)(\alpha+1) \pm \sqrt{(\alpha+1)^2 - (2-\rho^2)(h+\alpha+1)}}{2(2-\rho^2)} \end{cases}$$

On (.2.15) and (.2.16),

$$\begin{cases} \varsigma = 1 - \xi^{-1} \\ \eta = \frac{2-\rho^2}{1-\rho^2} \varsigma \\ z = -\frac{1}{1-\rho^2} \varsigma^2 \end{cases}$$

$$.2.9 \quad \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha ; \alpha, -\beta + \gamma + \delta, \delta, \gamma ; \frac{a(x-1)}{x-a} \right)$$

Polynomial of type 2

Replace coefficients $q, \beta, \gamma, \delta, x, c_0, \lambda$ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$, zero and h_j where $h_j \in \mathbb{N}_0$ into (9.3.1). Multiply $\left(\frac{x-a}{1-a} \right)^{-\alpha}$

and (9.3.1) together. Put (9.1.6) into the new (9.3.1).

$$\begin{aligned}
& \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}} \right)^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
= & \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}} \right)^{-\frac{1}{2}(\alpha+1)} Hl \left(\rho^{-2}, -\frac{1}{4} (h\rho^{-2} - (\alpha+1)^2) = -(h_j + 2j) ((1 + \rho^{-2})(h_j + 2j) + \alpha + 1) \right. \\
& \left. ; \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \frac{1}{2}; \frac{\xi - 1}{\rho^2(\xi - \rho^{-2})} \right) \\
= & \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}} \right)^{-\frac{1}{2}(\alpha+1)} \left\{ {}_2F_1 \left(-h_0, h_0 + \Omega_\rho; \frac{1}{2}; \eta \right) \right. \\
& + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \\
& \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k} - 1}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k}} \right)^{h_{n-k}} \\
& \times (1 - \overleftrightarrow{w}_{n-k+1,n}(1 - t_{n-k})(1 - u_{n-k})v_{n-k})^{-(4(n-k) + \Omega_\rho)} \\
& \times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k) - \frac{3}{2} + \frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right) \overleftrightarrow{w}_{n-k,n}^{2(n-k) - 1 + \frac{\alpha}{2}} \left. \right\} \\
& \times {}_2F_1 \left(-h_0, h_0 + \Omega_\rho; \frac{1}{2}; \overleftrightarrow{w}_{1,n} \right) \left. \right\} z^n \Bigg\} \tag{.2.17}
\end{aligned}$$

where

$$\begin{cases} \Omega_\rho = \frac{\alpha+1}{1+\rho^{-2}} \\ h = 4(h_j + 2j) ((1 + \rho^2)(h_j + 2j) + \rho^2(\alpha + 1)) + \rho^2(\alpha + 1)^2 \end{cases}$$

Infinite series

Replace coefficients $q, \beta, \gamma, \delta, x, c_0$ and λ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}, 1$ and zero into (9.3.4). Multiply $\left(\frac{x-a}{1-a}\right)^{-\alpha}$ and (9.3.4) together. Put (9.1.6) into the new (9.3.4).

$$\begin{aligned}
& \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}}\right)^{-\frac{1}{2}(\alpha+1)} y(\varsigma) \\
&= \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}}\right)^{-\frac{1}{2}(\alpha+1)} Hl\left(\rho^{-2}, -\frac{1}{4}(h\rho^{-2} - (\alpha+1)^2); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \frac{1}{2}; \frac{\xi - 1}{\rho^2(\xi - \rho^{-2})}\right) \\
&= \left(\frac{\xi - \rho^{-2}}{1 - \rho^{-2}}\right)^{-\frac{1}{2}(\alpha+1)} \left\{ {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \eta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \right. \right. \right. \\
&\times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(\frac{v_{n-k}-1}{v_{n-k}}\right)^{\Pi_{n-k}^+} (1 - \overleftrightarrow{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k})v_{n-k})^{\Pi_{n-k}^-} \\
&\times \overleftrightarrow{w}_{n-k,n}^{-(2(n-k)-\frac{3}{2}+\frac{\alpha}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}}\right) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}}\right) \overleftrightarrow{w}_{n-k,n}^{2(n-k)-1+\frac{\alpha}{2}} \Big\} \\
&\times \left. {}_2F_1\left(-\Pi_0^+, -\Pi_0^-; \frac{1}{2}; \overleftrightarrow{w}_{1,n}\right) \right\} z^n \Big\} \tag{.2.18}
\end{aligned}$$

where

$$\begin{cases} \Pi_0^\pm = \frac{-\rho^2(\alpha+1) \pm \sqrt{(1+\rho^2)h - \rho^2(\alpha+1)^2}}{2(1+\rho^2)} \\ \Pi_{n-k}^\pm = -2(n-k) + \frac{-\rho^2(\alpha+1) \pm \sqrt{(1+\rho^2)h - \rho^2(\alpha+1)^2}}{2(1+\rho^2)} \end{cases}$$

On (.2.17) and (.2.18),

$$\begin{cases} \varsigma = \frac{\xi-1}{\rho^2(\xi-\rho^{-2})} \\ \eta = (1+\rho^2)\varsigma \\ z = -\rho^2\varsigma^2 \end{cases}$$

.3 Generating function for the type 2 polynomial

.3.1 $(1-x)^{1-\delta} Hl(a, q - (\delta - 1)\gamma a; \alpha - \delta + 1, \beta - \delta + 1, \gamma, 2 - \delta; x)$

Replace coefficients q, α, β, δ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\delta - 1)\gamma a, \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta$ and h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(\rho^{-2}, -\frac{1}{4}(h-1)\rho^{-2} = -\rho^{-2}(h_j+2j)[1+(1+\rho^2)(h_j+2j)] \right. \\
& \left. ; \frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}; \frac{1}{2}, \frac{3}{2}; \xi \right) \\
& = 2^{-\Omega_\rho} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(2+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(1-\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-2+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-3-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \left. \left. \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-2+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-3-\alpha)} \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} z^n \Bigg\} \quad (.3.1)
\end{aligned}$$

where

$$\left\{ \begin{array}{l} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_\rho)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta)(1-t_1)(1-u_1)s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_\rho)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta)(1-t_n)(1-u_n)s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_\rho)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{array} \right.$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1} \right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_{0,\infty}^2 - 2(1-2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1} \right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1-2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1} \right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1-2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \eta = (1 + \rho^2)\xi \\ z = -\rho^2\xi^2 \\ \Omega_\rho = \frac{1}{1+\rho^2} \\ h = 4(h_j + 2j)[1 + (1 + \rho^2)(h_j + 2j)] + 1 \end{array} \right.$$

.3.2

$$x^{1-\gamma}(1-x)^{1-\delta}Hl(a, q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x)$$

Replace coefficients $q, \alpha, \beta, \gamma, \delta$ and q_j where $j, q_j \in \mathbb{N}_0$ by

$q - (\gamma + \delta - 2)a - (\gamma - 1)(\alpha + \beta - \gamma - \delta + 1), \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta$ and

h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{3}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(\rho^{-2}, -\frac{1}{4} ((h-4)\rho^{-2} - 1) \right. \\
& = -\rho^{-2}(h_j + 2j)[2 + \rho^2 + (1 + \rho^2)(h_j + 2j)]; \frac{\alpha}{2} + \frac{3}{2}, -\frac{\alpha}{2} + 1; \frac{3}{2}, \frac{3}{2}; \xi \Big) \\
& = 2^{\Omega_\rho} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{3}{2}} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(3+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(2-\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{1}{2}} \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-1+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-2-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}} \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-1+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-2-\alpha)} \Big\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \Big\} z^n \Big\} \quad (.3.2)
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) &= \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-4-\Omega_\rho}}{\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) &= \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-4n-\Omega_\rho}}{\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-4(n-k)-\Omega_\rho}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{aligned} \right.$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \Omega_\rho = \frac{1}{1+\rho^2} \\ h = 4 + \rho^2 + 4(h_j + 2j)[2 + \rho^2 + (1 + \rho^2)(h_j + 2j)] \\ \eta = (1 + \rho^2)\xi \\ z = -\rho^2\xi^2 \end{array} \right.$$

.3.3 $Hl(1-a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1-x)$

Replace coefficients a, q, γ, δ, x and q_j where $j, q_j \in \mathbb{N}_0$ by $1-a, -q + \alpha\beta, \delta, \gamma, 1-x$ and h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha(\alpha+1)) = -(2 - \rho^{-2})(h_j + 2j)^2 \right. \\
& \left. ; \frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; 1 - \xi \right) \\
& = 2^{-1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1-s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1-s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(1+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{\alpha}{2}} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1-s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4(n-1)-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \left. \left. \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k-1)-\alpha)} \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} z^n \Big\} \quad (.3.3)
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) &= \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-3}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) &= \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-4n+1}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-4(n-k)+1}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{aligned} \right.$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} h = \rho^2 (\alpha(\alpha + 1) - 4(2 - \rho^{-2})(h_j + 2j)^2) \\ \varsigma = 1 - \xi \\ \eta = \frac{2 - \rho^{-2}}{1 - \rho^{-2}} \varsigma \\ z = \frac{-1}{1 - \rho^{-2}} \varsigma^2 \end{array} \right.$$

$$\mathbf{.3.4} \quad (1 - x)^{1-\delta} Hl(1 - a, -q + (\delta - 1)\gamma a + (\alpha - \delta + 1)(\beta - \delta + 1); \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, \gamma; 1 - x)$$

Replace coefficients $a, q, \alpha, \beta, \gamma, \delta, x$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1 - a, -q + (\delta - 1)\gamma a + (\alpha - \delta + 1)(\beta - \delta + 1), \alpha - \delta + 1, \beta - \delta + 1, 2 - \delta, \gamma, 1 - x$ and h_j where

$h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{3}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(1 - \rho^{-2}, -\frac{1}{4} ((1+h)\rho^{-2} + (\alpha-1)(\alpha+2)) \right) \\
&= -(2 - \rho^{-2})(h_j + 2j)(h_j + 2j + 1); \frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}, \frac{3}{2}, \frac{1}{2}; 1 - \xi) \\
&= \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \\
&\times \tilde{w}_{1,1}^{-\frac{1}{2}(2+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\alpha+\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(1-\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
&+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-2+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-3-\alpha)} \\
&\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \quad (.3.4) \\
&\times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-2+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\alpha+\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-3-\alpha)} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) &= \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-4}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) &= \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-4n}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-4(n-k)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{aligned} \right.$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{-\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} h = 4(2\rho^2 - 1)(h_j + 2j)(h_j + 2j + 1) - \rho^2(\alpha - 1)(\alpha + 2) - 1 \\ \varsigma = 1 - \xi \\ \eta = \frac{2 - \rho^{-2}}{1 - \rho^{-2}} \varsigma \\ z = \frac{-1}{1 - \rho^{-2}} \varsigma^2 \end{array} \right.$$

$$\mathbf{.3.5} \quad x^{-\alpha} Hl \left(\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}; \alpha, \alpha - \gamma + 1, \alpha - \beta + 1, \delta; \frac{1}{x} \right)$$

Replace coefficients a, q, β, γ, x and q_j where $j, q_j \in \mathbb{N}_0$ by $\frac{1}{a}, \frac{q + \alpha[(\alpha - \gamma - \delta + 1)a - \beta + \delta]}{a}, \alpha - \gamma + 1, \alpha - \beta + 1, \frac{1}{x}$ and h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned} & \sum_{h_0=0}^{\infty} \frac{(\alpha + \frac{3}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(\rho^2, -\frac{1}{4} (h - (1 + \rho^2)(\alpha + 1)^2) \right) \\ &= -(1 + \rho^2)(h_j + 2j)(h_j + 2j + 1 + \alpha); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \alpha + \frac{3}{2}, \frac{1}{2}; \xi^{-1} \Big) \\ &= 2^\alpha \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\alpha + \frac{3}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\ & \quad \times \tilde{w}_{1,1}^{-\frac{1}{2}(1+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(2+\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ & \quad + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{1}{2}+\alpha} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ & \quad \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{-\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-2+\alpha)} \\ & \quad \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{1}{2}+\alpha} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ & \quad \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{-\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-2+\alpha)} \Big\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \Big\} z^n \Big\} \end{aligned} \quad (.3.5)$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) &= \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(4+\alpha)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) &= \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n+\alpha)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)+\alpha)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{aligned} \right.$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{-(\alpha + \frac{1}{2})} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-(\alpha + \frac{1}{2})} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-(\alpha + \frac{1}{2})} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} h = (1 + \rho^2)(2(h_j + 2j) + 1 + \alpha)^2 \\ \varsigma = \xi^{-1} \\ \eta = (1 + \rho^{-2})\varsigma \\ z = -\rho^{-2}\varsigma^2 \end{array} \right.$$

$$\mathbf{.3.6} \quad \left(1 - \frac{x}{a}\right)^{-\beta} Hl \left(1 - a, -q + \gamma\beta; -\alpha + \gamma + \delta, \beta, \gamma, \delta; \frac{(1-a)x}{x-a}\right)$$

Replace coefficients a, q, α, x and q_j where $j, q_j \in \mathbb{N}_0$ by $1 - a, -q + \gamma\beta, -\alpha + \gamma + \delta, \frac{(1-a)x}{x-a}$ and h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned} & \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(1 - \rho^{-2}, \frac{1}{4} (h\rho^{-2} - \alpha) = (h_j + 2j) (\alpha - (2 - \rho^{-2})(h_j + 2j)) \right. \\ & \left. ; -\frac{\alpha}{2} + \frac{1}{2}, -\frac{\alpha}{2}, \frac{1}{2}, \frac{1}{2}; \frac{(1 - \rho^{-2})\xi}{\xi - \rho^{-2}} \right) \\ & = 2^{-(\Omega_\rho+1)} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\ & \times \tilde{w}_{1,1}^{-\frac{1}{2}(1-\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{\alpha}{2}} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ & + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ & \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3-\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4(n-1)-\alpha)} \\ & \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ & \left. \left. \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3-\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k-1)-\alpha)} \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \right\} z^n \end{aligned} \quad (.3.6)$$

where

$$\begin{cases} \overleftrightarrow{\Gamma}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3-\Omega_\rho)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1-\Omega_\rho)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\Gamma}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1-\Omega_\rho)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{cases}$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2} + \Omega_\rho}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2} + \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2} + \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \Omega_\rho = \frac{\alpha}{2 - \rho^{-2}} \\ h = 4\rho^2 \left(\alpha \left(h_j + 2j + \frac{1}{4} \right) - (2 - \rho^{-2})(h_j + 2j)^2 \right) \\ \varsigma = \frac{(1 - \rho^{-2})\xi}{\xi - \rho^{-2}} \\ \eta = \frac{2 - \rho^{-2}}{1 - \rho^{-2}} \varsigma \\ z = -\frac{1}{1 - \rho^{-2}} \varsigma^2 \end{array} \right.$$

3.7

$$(1 - x)^{1 - \delta} \left(1 - \frac{x}{a}\right)^{-\beta + \delta - 1} Hl \left(1 - a, -q + \gamma[(\delta - 1)a + \beta - \delta + 1]; -\alpha + \gamma + 1, \beta - \delta + 1, \gamma, 2 - \delta; \frac{(1 - a)x}{x - a}\right)$$

Replace coefficients $a, q, \alpha, \beta, \delta, x$ and q_j where $j, q_j \in \mathbb{N}_0$ by $1 - a, -q + \gamma[(\delta - 1)a + \beta - \delta + 1], -\alpha + \gamma + 1, \beta - \delta + 1, 2 - \delta, \frac{(1 - a)x}{x - a}$ and h_j where $h_j \in \mathbb{N}_0$

into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(1 - \rho^{-2}, \frac{1}{4} ((h-1)\rho^{-2} + 1 - \alpha) \right. \\
& = (h_j + 2j) \left(\alpha + 1 - (2 - \rho^{-2})(h_j + 2j + 1) \right); -\frac{\alpha}{2} + 1, -\frac{\alpha}{2} + \frac{1}{2}, \frac{1}{2}, \frac{3}{2}; \frac{(1 - \rho^{-2})\xi}{\xi - \rho^{-2}} \Big) \\
& = 2^{-\Omega_\rho} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
& \times \tilde{w}_{1,1}^{-\frac{1}{2}(2-\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(1-\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
& + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
& \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-2-\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-3-\alpha)} \\
& \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
& \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-2-\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-3-\alpha)} \Big\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \Big\} z^n \Big\} \quad (.3.7)
\end{aligned}$$

where

$$\left\{ \begin{aligned} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) &= \frac{\left(\frac{1+s_{1,\infty}+\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}}{2} \right)^{-4+\Omega_\rho}}{\sqrt{s_{1,\infty}^2-2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) &= \frac{\left(\frac{1+s_{n,\infty}+\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}}{2} \right)^{-4n+\Omega_\rho}}{\sqrt{s_{n,\infty}^2-2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty}+1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) &= \frac{\left(\frac{1+s_{n-k}+\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}}{2} \right)^{-4(n-k)+\Omega_\rho}}{\sqrt{s_{n-k}^2-2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k}+1}} \end{aligned} \right.$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{-\frac{1}{2} + \Omega_\rho}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{-\frac{1}{2} + \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{-\frac{1}{2} + \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \Omega_\rho = \frac{\alpha+1}{2-\rho^{-2}} \\ h = 4(h_j + 2j) \left((\alpha + 1)\rho^2 + (1 - 2\rho^2)(h_j + 2j + 1) \right) + (\alpha - 1)\rho^2 + 1 \\ \varsigma = \frac{(1-\rho^{-2})\xi}{\xi-\rho^{-2}} \\ \eta = \frac{2-\rho^{-2}}{1-\rho^{-2}}\varsigma \\ z = -\frac{1}{1-\rho^{-2}}\varsigma^2 \end{array} \right.$$

$$\mathbf{.3.8} \quad x^{-\alpha} Hl \left(\frac{a-1}{a}, \frac{-q + \alpha(\delta a + \beta - \delta)}{a}; \alpha, \alpha - \gamma + 1, \delta, \alpha - \beta + 1; \frac{x-1}{x} \right)$$

Polynomial of type 2

Replace coefficients $a, q, \beta, \gamma, \delta, x$ and q_j where $j, q_j \in \mathbb{N}_0$ by $\frac{a-1}{a}, \frac{-q+\alpha(\delta a+\beta-\delta)}{a}, \alpha-\gamma+1, \delta, \alpha-\beta+1, \frac{x-1}{x}$ and h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new

(9.4.1).

$$\begin{aligned}
& \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(1 - \rho^2, \frac{1}{4} [h + (\alpha + 1) (1 - \rho^2(\alpha + 1))] \right) \\
&= -(h_j + 2j) \left((1 - \rho^2)(\alpha + 1) + (2 - \rho^2)(h_j + 2j) \right); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \alpha + \frac{3}{2}; \frac{\xi - 1}{\xi} \Big) \\
&= 2^{\Omega_\rho - 1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\
&\quad \times \tilde{w}_{1,1}^{-\frac{1}{2}(1+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(2+\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\
&\quad + \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\
&\quad \times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{-\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-2+\alpha)} \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\
&\quad \times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{-\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-2+\alpha)} \Big\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \Big\} z^n \Big\} \quad (.3.8)
\end{aligned}$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_\rho)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_\rho)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_\rho)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{cases}$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \Omega_\rho = \frac{(\alpha+1)(1-\rho^2)}{2-\rho^2} \\ h = -4(h_j + 2j) \left((1 - \rho^2)(\alpha + 1) + (2 - \rho^2)(h_j + 2j) \right) - (\alpha + 1) (1 - \rho^2(\alpha + 1)) \\ \varsigma = 1 - \xi^{-1} \\ \eta = \frac{2-\rho^2}{1-\rho^2} \varsigma \\ z = -\frac{1}{1-\rho^2} \varsigma^2 \end{array} \right.$$

$$\mathbf{.3.9} \quad \left(\frac{x-a}{1-a} \right)^{-\alpha} Hl \left(a, q - (\beta - \delta)\alpha; \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a} \right)$$

Replace coefficients $q, \beta, \gamma, \delta, x$ and q_j where $j, q_j \in \mathbb{N}_0$ by $q - (\beta - \delta)\alpha, -\beta + \gamma + \delta, \delta, \gamma, \frac{a(x-1)}{x-a}$ and h_j where $h_j \in \mathbb{N}_0$ into (9.4.1). Put (9.1.6) into the new (9.4.1).

$$\begin{aligned} & \sum_{h_0=0}^{\infty} \frac{(\frac{1}{2})_{h_0}}{h_0!} s_0^{h_0} \prod_{n=1}^{\infty} \left\{ \sum_{h_n=h_{n-1}}^{\infty} s_n^{h_n} \right\} Hl \left(\rho^{-2}, -\frac{1}{4} (h\rho^{-2} - (\alpha+1)^2) \right. \\ &= -(h_j + 2j) \left((1 + \rho^{-2})(h_j + 2j) + \alpha + 1 \right); \frac{\alpha}{2} + \frac{1}{2}, \frac{\alpha}{2} + 1, \frac{1}{2}, \frac{1}{2}; \frac{\xi - 1}{\rho^2(\xi - \rho^{-2})} \left. \right) \\ &= 2^{\Omega_\rho - 1} \left\{ \prod_{k=1}^{\infty} \frac{1}{(1 - s_{k,\infty})} \mathbf{A}(s_{0,\infty}; \eta) + \prod_{k=2}^{\infty} \frac{1}{(1 - s_{k,\infty})} \int_0^1 dt_1 t_1 \int_0^1 du_1 u_1^{\frac{1}{2}} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) \right. \\ &\times \tilde{w}_{1,1}^{-\frac{1}{2}(1+\alpha)} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{-\frac{1}{2}} (\tilde{w}_{1,1} \partial_{\tilde{w}_{1,1}}) \tilde{w}_{1,1}^{\frac{1}{2}(2+\alpha)} \mathbf{A}(s_0; \tilde{w}_{1,1}) z \\ &+ \sum_{n=2}^{\infty} \left\{ \prod_{l=n+1}^{\infty} \frac{1}{(1 - s_{l,\infty})} \int_0^1 dt_n t_n^{2n-1} \int_0^1 du_n u_n^{2n-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) \right. \\ &\times \tilde{w}_{n,n}^{-\frac{1}{2}(4n-3+\alpha)} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{-\frac{1}{2}} (\tilde{w}_{n,n} \partial_{\tilde{w}_{n,n}}) \tilde{w}_{n,n}^{\frac{1}{2}(4n-2+\alpha)} \\ &\times \prod_{k=1}^{n-1} \left\{ \int_0^1 dt_{n-k} t_{n-k}^{2(n-k)-1} \int_0^1 du_{n-k} u_{n-k}^{2(n-k)-\frac{3}{2}} \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \right. \\ &\times \tilde{w}_{n-k,n}^{-\frac{1}{2}(4(n-k)-3+\alpha)} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{-\frac{1}{2}} (\tilde{w}_{n-k,n} \partial_{\tilde{w}_{n-k,n}}) \tilde{w}_{n-k,n}^{\frac{1}{2}(4(n-k)-2+\alpha)} \left. \right\} \mathbf{A}(s_0; \tilde{w}_{1,n}) \left. \right\} z^n \Bigg\} \end{aligned} \quad (3.9)$$

where

$$\begin{cases} \overleftrightarrow{\mathbf{F}}_1(s_{1,\infty}; t_1, u_1, \eta) = \frac{\left(\frac{1+s_{1,\infty} + \sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}}{2} \right)^{-(3+\Omega_\rho)}}{\sqrt{s_{1,\infty}^2 - 2(1-2\eta(1-t_1)(1-u_1))s_{1,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_n(s_{n,\infty}; t_n, u_n, \eta) = \frac{\left(\frac{1+s_{n,\infty} + \sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}}{2} \right)^{-(4n-1+\Omega_\rho)}}{\sqrt{s_{n,\infty}^2 - 2(1-2\eta(1-t_n)(1-u_n))s_{n,\infty} + 1}} \\ \overleftrightarrow{\mathbf{F}}_{n-k}(s_{n-k}; t_{n-k}, u_{n-k}, \tilde{w}_{n-k+1,n}) \\ = \frac{\left(\frac{1+s_{n-k} + \sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}}{2} \right)^{-(4(n-k)-1+\Omega_\rho)}}{\sqrt{s_{n-k}^2 - 2(1-2\tilde{w}_{n-k+1,n}(1-t_{n-k})(1-u_{n-k}))s_{n-k} + 1}} \end{cases}$$

$$\left\{ \begin{array}{l} \mathbf{A}(s_{0,\infty}; \eta) = \frac{\left(1 - s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2}} \left(1 + s_{0,\infty} + \sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_{0,\infty}^2 - 2(1 - 2\eta)s_{0,\infty} + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,1}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,1})s_0 + 1}} \\ \mathbf{A}(s_0; \tilde{w}_{1,n}) = \frac{\left(1 - s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2}} \left(1 + s_0 + \sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}\right)^{\frac{1}{2} - \Omega_\rho}}{\sqrt{s_0^2 - 2(1 - 2\tilde{w}_{1,n})s_0 + 1}} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \Omega_\rho = \frac{\alpha+1}{1+\rho^{-2}} \\ h = 4(h_j + 2j) \left((1 + \rho^2)(h_j + 2j) + \rho^2(\alpha + 1) \right) + \rho^2(\alpha + 1)^2 \\ \varsigma = \frac{\xi-1}{\rho^2(\xi-\rho^{-2})} \\ \eta = (1 + \rho^2)\varsigma \\ z = -\rho^2\varsigma^2 \end{array} \right.$$

Bibliography

- [1] Heun, K., “Zur Theorie der Riemann’schen Functionen zweiter Ordnung mit vier Verzweigungspunkten,” *Mathematische Annalen* **33**, (1889)161.
- [2] Ronveaux, A., *Heuns Differential Equations*, (Oxford University Press, 1995).
- [3] Lamé, G., “Sur les surfaces isothermes dans les corps homogenes en equilibre de temperature,” (French) *J. Math. Pures. Appl.* **2**, 147-188(1837).
- [4] Hobson, E.W., *The theory of spherical and ellipsoidal harmonics*. Cambridge Univ. Press, (1931).
- [5] Whittaker, E.T. and Watson, G.N., *A course of modern analysis*. Cambridge Univ. Press, (1952).
- [6] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806.
- [7] Choun, Y.S., “The analytic solution for the power series expansion of Heun function,” <http://dx.doi.org/10.1016/j.aop.2013.06.020> Available as arXiv:1303.0830.
- [8] Choun, Y.S., “Asymptotic behavior of Heun function and its integral formalism,” arXiv:1303.0876.
- [9] Choun, Y.S., “Lamé equation in the algebraic form,” arXiv:1303.0873.
- [10] Choun, Y.S., “Power series and integral forms of Lamé equation in Weierstrass’s form,” arXiv:1303.0878.
- [11] Choun, Y.S., “The generating functions of Lamé equation in Weierstrass’s form,” arXiv:1303.0879.

- [12] Stieltjes, T. J., “Sur certain polynomes que verifient une equation diferentielle lineaire du second ordre et sur la teorie des fonctions de Lamé,” *Acta Math.* **6**, 321–326(1885).
- [13] Moon, P. and Spencer, D. E., *Field Theory for Engineers*. New York: Van Nostrand, (1961).
- [14] Zwillinger, D., *Handbook of Differential Equations*. 3rd ed. Boston, MA: Academic Press, (1997).
- [15] Bôcher, M., “Über die Reihenentwickelungen der Potentialtheorie,” Leipzig: B. G. Teubner (1894).
- [16] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G., *Higher Transcendental Functions. Vol. III* McGraw-Hill Book Company, Inc., New York-Toronto-London, (1955).
- [17] Erdelyi, A., *Integral equations for Lamé functions*. Proceedings of the Edinburgh Mathematical Society (Series 2) Vol.7 Issue 01 (1940), 3–15.
- [18] Arscott, F. M., “Integral equations and relations for Lamé functions,” *Quart. J. Math. Oxford Ser. (2)* **15**, 103-115(1964).
- [19] Shail, R., “On integral representations for Lamé and other special functions,” *SIAM J. Math. Anal.* **11** (4), 702-723(1980).
- [20] Sleeman, B. D., “Integral equations and relations for Lamé functions and ellipsoidal wave functions,” *Proc. Cambridge Philos. Soc.* **64**, 113-126(1968).
- [21] Volkmer, H., “Integral relations for Lamé functions,” *SIAM J. Math. Anal.* **13** (6), 978-987(1982).
- [22] Volkmer, H., “Integralgleichungen fr periodische Lösungen Hillscher Differentialgleichungen,” (German) *Analysis* **3** (1-4), 189-203(1983).
- [23] Volkmer, H., “Integral representations for products of Lamé functions by use of fundamental solutions,” *SIAM J. Math. Anal.* **15** (3), 559-569(1984).
- [24] Wang, Z.X. and Guo, D. R., *Special Functions*, World Scientific Pub., 602–606(1989).
- [25] Whittaker, E.T. and Watson, G.N., “A Course of Modern Analysis (Cambridge Mathematical Library),” Cambridge University Press, 554–575(1996).
- [26] Maier, R.S., “The 192 solutions of the Heun equation,” *Math. Comp.* **33**, (2007)811–843.