

# A conjecture on the primitive degree of Tensors <sup>\*</sup>

Pingzhi Yuan<sup>†</sup>, Zilong He, Lihua You<sup>‡</sup>

School of Mathematical Sciences, South China Normal University,  
Guangzhou, 510631, P.R. China

## Abstract

In this paper, we prove: Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Then its primitive degree  $\gamma(\mathbb{A}) \leq (n-1)^2 + 1$ , and the upper bound is sharp. This confirms a conjecture of Shao [7].

*AMS classification:* 05C50; 15A69

*Keywords:* tensor; product; primitive tensor; primitive degree.

## 1 Introduction

In [1] and [2], Chang et al investigated the properties of the spectra of nonnegative tensors. They defined the irreducibility of tensors, and the primitivity of nonnegative tensors, and extended many important properties of primitive matrices to primitive tensors. Recently, as an application of the general tensor product defined by Shao [7], Shao presented a simple characterization of the primitive tensors in terms of the zero pattern of the powers of  $\mathbb{A}$ . He also proposed the following conjecture on the primitive degree.

**Conjecture 1.1.** *When  $m$  is fixed, then there exists some polynomial  $f(n)$  on  $n$  such that  $\gamma(\mathbb{A}) \leq f(n)$  for all nonnegative primitive tensors of order  $m$  and dimension  $n$ .*

In this paper, we confirm the conjecture by proving the following theorem.

**Theorem 1.2.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Then its primitive degree  $\gamma(\mathbb{A}) \leq (n-1)^2 + 1$ , and the upper bound is sharp.*

---

<sup>\*</sup>P. Yuan's research is supported by the NSF of China (Grant No. 11271142) and the Guangdong Provincial Natural Science Foundation(Grant No. S2012010009942), L. You's research is supported by the Zhujiang Technology New Star Foundation of Guangzhou (Grant No. 2011J2200090) and Program on International Cooperation and Innovation, Department of Education, Guangdong Province (Grant No.2012gjhz0007).

<sup>†</sup>*Corresponding author:* yuanpz@scnu.edu.cn.

<sup>‡</sup>*Email address:* ylhua@scnu.edu.cn.

## 2 Preliminaries

An order  $m$  dimension  $n$  tensor  $\mathbb{A} = (a_{i_1 i_2 \dots i_m})_{1 \leq i_j \leq n \ (j=1, \dots, m)}$  over the complex field  $\mathbb{C}$  is a multidimensional array with all entries  $a_{i_1 i_2 \dots i_m} \in \mathbb{C}$  ( $i_1, \dots, i_m \in [n] = \{1, \dots, n\}$ ). The majorization matrix  $M(\mathbb{A})$  of the tensor  $\mathbb{A}$  is defined as  $(M(\mathbb{A}))_{ij} = a_{ij \dots j}$ , ( $i, j \in [n]$ ) by Pearson [4].

Let  $\mathbb{A}$  (and  $\mathbb{B}$ ) be an order  $m \geq 2$  (and  $k \geq 1$ ), dimension  $n$  tensor, respectively. Recently, Shao [7] defined a general product  $\mathbb{A}\mathbb{B}$  to be the following tensor  $\mathbb{D}$  of order  $(m-1)(k-1)+1$  and dimension  $n$ :

$$d_{i\alpha_1 \dots \alpha_{m-1}} = \sum_{i_2, \dots, i_m=1}^n a_{i i_2 \dots i_m} b_{i_2 \alpha_1} \dots b_{i_m \alpha_{m-1}} \quad (i \in [n], \alpha_1, \dots, \alpha_{m-1} \in [n]^{k-1}).$$

The tensor product possesses a very useful property: the associative law ([7], Theorem 1.1). With the general product, Shao [7] proved some results on the primitivity and primitive degree of nonnegative tensor. The following result will be used in Definition 2.3.

**Proposition 2.1.** ([7], Proposition 4.1) *Let  $\mathbb{A}$  be an order  $m$  and dimension  $n$  nonnegative tensor. Then the following three conditions are equivalent:*

- (1). *For any  $i, j \in [n]$ ,  $a_{ij \dots j} > 0$  holds.*
- (2). *For any  $j \in [n]$ ,  $\mathbb{A}e_j > 0$  holds (where  $e_j$  is the  $j^{\text{th}}$  column of the identity matrix  $I_n$ ).*
- (3). *For any nonnegative nonzero vector  $x \in \mathbb{R}^n$ ,  $\mathbb{A}x > 0$  holds.*

**Definition 2.2.** ([4], Definition 3.1) *A nonnegative tensor  $\mathbb{A}$  is called essentially positive, if it satisfies (3) of Proposition 2.1.*

By Proposition 2.1, the following Definition 2.3 is equivalent to Definition 2.2.

**Definition 2.3.** ([7], Definition 4.1) *A nonnegative tensor  $\mathbb{A}$  is called essentially positive, if it satisfies one of the three conditions in Proposition 2.1.*

In [2] and [4], Chang et al and Pearson define the primitive tensors as follows.

**Definition 2.4.** ([2, 5]) *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$ ,  $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$  a vector and  $x^{[r]} = (x_1^r, x_2^r, \dots, x_n^r)^T$ . Define the map  $T_{\mathbb{A}}$  from  $\mathbb{R}^n$  to  $\mathbb{R}^n$  as:  $T_{\mathbb{A}}(x) = (\mathbb{A}x)^{[\frac{1}{m-1}]}$ . If there exists some positive integer  $r$  such that  $T_{\mathbb{A}}^r(x) > 0$  for all nonnegative nonzero vectors  $x \in \mathbb{R}^n$ , then  $\mathbb{A}$  is called primitive and the smallest such integer  $r$  is called the primitive degree of  $\mathbb{A}$ , denoted by  $\gamma(\mathbb{A})$ .*

In [7], Shao show the following results and define the primitive degree by using the properties of tensor product and the zero patterns.

**Proposition 2.5.** ([7], Theorem 4.1) *A nonnegative tensor  $\mathbb{A}$  is primitive if and only if there exists some positive integer  $r$  such that  $\mathbb{A}^r$  is essentially positive. Furthermore, the smallest such  $r$  is the primitive degree of  $\mathbb{A}$ .*

**Remark 2.6.** *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$ . Then  $\mathbb{A}$  is primitive if and only if there exists some positive integer  $r$  such that  $M(\mathbb{A}^r) > 0$ .*

Now we prove the following necessary conditions for a tensor to be primitive firstly.

**Proposition 2.7.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ ,  $M(\mathbb{A})$  the majorization matrix of  $\mathbb{A}$ . Then we have:*

- (i). *For each  $j \in [n]$ , there exists an integer  $i \in [n] \setminus \{j\}$  such that  $(M(\mathbb{A}))_{ij} > 0$ .*
- (ii). *There exist some  $j \in [n]$  and integers  $u, v$  with  $1 \leq u < v \leq n$  such that  $(M(\mathbb{A}))_{uj} > 0$  and  $(M(\mathbb{A}))_{vj} > 0$ .*

*Proof.* We prove the results via contradiction.

(i) To obtain a contradiction, we suppose that there exists some integer  $j \in [n]$  such that  $(M(\mathbb{A}))_{ij} = 0$  for every  $i \in [n] \setminus \{j\}$ . By definitions of the tensor product and the majorization matrix, for any  $u \in [n] \setminus \{j\}$ , we have  $(M(\mathbb{A}))_{uj} = 0$  and

$$\begin{aligned} (M(\mathbb{A}^2))_{uj} &= \sum_{j_2, \dots, j_m=1}^n a_{uj_2 \dots j_m} a_{j_2 j \dots j} \cdots a_{j_m j \dots j} \\ &= \sum_{j_2, \dots, j_m=1}^n a_{uj_2 \dots j_m} (M(\mathbb{A}))_{j_2 j} \cdots (M(\mathbb{A}))_{j_m j} \\ &= (M(\mathbb{A}))_{uj} (M(\mathbb{A}))_{jj}^{m-1} = 0. \end{aligned}$$

Since

$$(M(\mathbb{A}^{r+1}))_{uj} = \sum_{j_2, \dots, j_m=1}^n a_{uj_2 \dots j_m} (M(\mathbb{A}^r))_{j_2 j} \cdots (M(\mathbb{A}^r))_{j_m j},$$

hence, by induction on  $k$ , we conclude that

$$(M(\mathbb{A}^k))_{uj} = 0$$

holds for any positive integer  $k$  and any  $u \in [n] \setminus \{j\}$ . This contradicts that  $(M(\mathbb{A}^{\gamma(\mathbb{A})}))_{uj} > 0$ , where  $\gamma(\mathbb{A})$  is the primitive degree of  $\mathbb{A}$ . (i) is proved.

(ii) Suppose, to derive a contradiction, that there is at most one nonzero element in each of the following  $n$  sets

$$\{(M(\mathbb{A}))_{uj}, u \in [n]\}, \quad j \in [n].$$

Now we will show that for any positive integer  $t$ , there is at most one nonzero element in each of the following  $n$  sets

$$\{(M(\mathbb{A}^t))_{uj}, u \in [n]\}, \quad j \in [n].$$

We prove the above assertion by induction on  $t$ . Clearly,  $t = 1$  is obvious. Assume that the assertion holds for  $t = k \geq 1$ , that is, there is at most one nonzero element in each of the following  $n$  sets

$$\{(M(\mathbb{A}^k))_{uj}, u \in [n]\}, \quad j \in [n],$$

say, for any  $j \in [n]$  and any  $u \neq u_j$ ,  $(M(\mathbb{A}^k))_{uj} = 0$ . Note that for any  $v \in [n]$ ,

$$(M(\mathbb{A}^{k+1}))_{vj} = \sum_{j_2, \dots, j_m=1}^n a_{vj_2 \dots j_m} (M(\mathbb{A}^k))_{j_2 j} \cdots (M(\mathbb{A}^k))_{j_m j} = a_{vu_j \dots u_j} ((M(\mathbb{A}^k))_{u_j j})^{m-1},$$

by the assumption, there is at most one  $v \in [n]$  such that  $a_{vu_j \dots u_j} = (M(\mathbb{A}))_{vu_j} > 0$ , therefore we have prove the assertion. It follows that  $\mathbb{A}$  is not a primitive tensor, this contradiction proves (ii).  $\square$

Follows from the proof of Proposition 2.7, we know that

$$(M(\mathbb{A}^{k+1}))_{uj} = \sum_{j_2, \dots, j_m=1}^n a_{uj_2 \dots j_m} (M(\mathbb{A}^k))_{j_2 j} \cdots (M(\mathbb{A}^k))_{j_m j}, \quad (2.1)$$

is useful and used repeatedly.

By Equation (2.1), it is easy to prove the following assertions.

**Corollary 2.8.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Then we have*

- (i). *For each  $u \in [n]$ , there is at least one index  $j_2 \cdots j_m \in [n]^{m-1}$  such that  $a_{uj_2 \dots j_m} > 0$ .*
- (ii). *Let  $k \in [n]$  be fixed. Suppose that  $T$  is a positive integer such that  $(M(\mathbb{A}^T))_{uk} > 0$  for all  $u \in [n]$ , then for any  $t \geq T$ , we have  $(M(\mathbb{A}^t))_{uk} > 0$  for all  $u \in [n]$ .*

**Proposition 2.9.** *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$ , and let  $a$  be a positive integer. Then  $\mathbb{A}$  is primitive if and only if  $\mathbb{A}^a$  is primitive.*

Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ , for any  $j = j_1 \in [n]$ , by Proposition 2.7, there exists a sequence  $j_1, j_2, \dots, j_{s+1}$  such that  $j_k \in [n], j_k \neq j_{k+1}, 1 \leq k \leq s$  and  $(M(\mathbb{A}))_{j_{k+1} j_k} > 0$ .

**Definition 2.10.** *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$ , if  $j_k \in [n]$  for  $1 \leq k \leq t$  and  $(M(\mathbb{A}))_{j_{k+1} j_k} > 0$  for  $1 \leq k \leq t-1$ , we say that  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_t$  is a walk of length  $t-1$  of  $M(\mathbb{A})$ ; if  $j_i \neq j_k$  for any  $i, k \in [t]$  with  $i \neq k$ , then we say the walk  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_t$  is a path of  $M(\mathbb{A})$ ; if  $j_i \neq j_k$  for any  $i, k \in [t-1]$  with  $i \neq k$  but  $j_1 = j_t$ , then we say the walk  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_t$  is a cycle of  $M(\mathbb{A})$ .*

**Lemma 2.11.** *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$ . Suppose that  $t > 1$  and  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_t$  is a walk of  $M(\mathbb{A})$ , then  $(M(\mathbb{A}^{t-1}))_{j_t j_1} > 0$ .*

*Proof.* We prove the assertion by induction on  $t$ . Clearly,  $t = 2$  is obvious. Assume that the result holds for  $t = k \geq 2$ , that is  $(M(\mathbb{A}^{k-1}))_{j_k j_1} > 0$ . Since

$$\begin{aligned} (M(\mathbb{A}^k))_{j_{k+1} j_1} &= \sum_{i_2, \dots, i_m=1}^n a_{j_{k+1} i_2 \dots i_m} (M(\mathbb{A}^{k-1}))_{i_2 j_1} \cdots (M(\mathbb{A}^{k-1}))_{i_m j_1} \\ &> a_{j_{k+1} j_k \dots j_k} ((M(\mathbb{A}^{k-1}))_{j_k j_1})^{m-1} \\ &= (M(\mathbb{A}))_{j_{k+1} j_k} ((M(\mathbb{A}^{k-1}))_{j_k j_1})^{m-1} > 0, \end{aligned}$$

by assumption, thus the assertion holds for any integer  $t > 1$ . We are done.  $\square$

**Lemma 2.12.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Then there exist an integer  $j \in [n]$  and an integer  $l \in [n-1]$  such that  $(M(\mathbb{A}^l))_{jj} > 0$ .*

*Proof.* By Proposition 2.7, we may assume that  $j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_{n+1}$  is a walk of length  $n$  of  $M(\mathbb{A})$ . Since  $j_k \in [n], 1 \leq k \leq n+1$ , there exist at least two integers  $u$  and  $v$  such that  $1 \leq u < v \leq n+1$  and  $j_u = j_v$ . It follows from Lemma 2.11 that  $(M(\mathbb{A}^{v-u}))_{j_u j_u} > 0$ .

**Case 1:** There exist  $u, v$  such that  $(u, v) \neq (1, n+1)$ .

Then  $v - u \leq n - 1$  and we are done by taking  $j = j_u$  and  $l = v - u$ .

**Case 2:**  $(u, v) = (1, n+1)$ .

Then  $j_1, j_2, \dots, j_n$  is a permutation of  $1, 2, \dots, n$  and  $j_{n+1} = j_1$ . By (ii) of Proposition 2.7, there exist an integer  $i \in [n]$  and two integers  $p \neq q$  such that  $1 \leq p, q \leq n$ ,  $(M(\mathbb{A}))_{pi} > 0$

and  $(M(\mathbb{A}))_{qi} > 0$ . Take  $i = j_t$  for some  $t \in [n]$ , and assume that  $p = j_s \neq j_{t+1}$ . Thus  $s \in [n]$ , and  $t \leq n - 1$  when  $s = 1$ .

**Subcase 2.1:**  $p \in \{j_1, \dots, j_{t-1}\}$ .

Then  $1 \leq s \leq t - 1$  and

$$p = j_s \rightarrow \dots \rightarrow j_{t-1} \rightarrow j_t \rightarrow p = j_s$$

is a cycle of  $M(\mathbb{A})$  with length  $t + 1 - s \leq n - 1$ . We take  $j = p$  and  $l = t + 1 - s$ .

**Subcase 2.2:**  $p = j_t$ .

Then  $s = t$  and  $j_t \rightarrow p (= j_s)$  is a cycle of  $M(\mathbb{A})$  with length 1. We take  $j = j_t$  and  $l = 1$ .

**Subcase 2.3:**  $p \in \{j_{t+2}, \dots, j_n\}$ .

Then  $t + 2 \leq s \leq n$  and

$$j_1 \rightarrow j_2 \rightarrow \dots \rightarrow j_t \rightarrow j_s (= p) \rightarrow \dots \rightarrow j_n \rightarrow j_1$$

is a cycle of  $M(\mathbb{A})$  and  $t + n - s + 1 = n - (s - t - 1) \leq n - 1$ . In this case we take  $j = j_1$  and  $l = n + t + 1 - s$ . This proves the lemma.  $\square$

**Remark 2.13.** *By the proof of Lemma 2.12, if  $\mathbb{A}$  is a nonnegative primitive tensor with order  $m$  and dimension  $n$ , then there exist an integer  $t$  with  $1 \leq t \leq n - 1$  and some integers  $j_1, j_2, \dots, j_t \in [n]$  such that  $j_1 \rightarrow j_2 \rightarrow \dots \rightarrow j_t \rightarrow j_1$  is a cycle of length  $t$  of  $M(\mathbb{A})$ , and for any  $k \in [t]$ ,  $(M(\mathbb{A}^t))_{j_k j_k} > 0$ .*

Note that by (2.1), (ii) of Corollary 2.8 also holds when  $\mathbb{A}$  is a nonnegative tensor with order  $m$  and dimension  $n$ . Therefore it makes sense to consider the primitive degree of some column of a tensor.

**Definition 2.14.** *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$ . For a fixed integer  $j \in [n]$ , if there exists a positive integer  $T$  such that*

$$(M(\mathbb{A}^T))_{uj} > 0, \text{ for all } u, 1 \leq u \leq n,$$

*then  $\mathbb{A}$  is called  $j$ -primitive and the smallest such integer  $T$  is called the  $j$ -primitive degree of  $\mathbb{A}$ , denoted by  $\gamma_j(\mathbb{A})$ .*

By Corollary 2.8 and the above definition, we have the following result.

**Proposition 2.15.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Then*

$$\gamma(\mathbb{A}) = \max_{1 \leq j \leq n} \{\gamma_j(\mathbb{A})\}.$$

### 3 Proof of the main results

In this section, we will prove Theorem 1.2. We first prove the following special case of the theorem.

**Theorem 3.1.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Suppose that there is an integer  $j \in [n]$  with  $(M(\mathbb{A}))_{jj} > 0$ , then  $\gamma_j(\mathbb{A}) \leq n - 1$ .*

*Proof.* Put

$$S_k = \{u \in [n], (M(\mathbb{A}^k))_{uj} > 0, k = 1, 2, \dots\}$$

Then  $j \in S_1$  and there exists an integer  $v \in [n] \setminus \{j\}$  such that  $v \in S_1$  by (i) of Proposition 2.7. Thus  $|S_1| \geq 2$ .

Since

$$(M(\mathbb{A}^{k+1}))_{uj} = \sum_{i_2, \dots, i_s=1}^n (\mathbb{A}^k)_{ui_2 \dots i_s} (M(\mathbb{A}))_{i_2 j} \cdots (M(\mathbb{A}))_{i_s j} \geq (M(\mathbb{A}^k))_{uj} (M(\mathbb{A}))_{jj}^{s-1},$$

we see that if  $u \in S_k$ , then  $u \in S_{k+1}$ . Therefore by induction on  $k$ , we can obtain that  $S_1 \subseteq S_2 \subseteq \cdots \subseteq S_l \subseteq S_{l+1} \subseteq \cdots$ .

Note that  $S_k \subseteq [n]$  for any  $k = 1, 2, \dots$ , and  $S_1 \subseteq S_2 \subseteq \cdots \subseteq S_l \subseteq S_{l+1} \subseteq \cdots$ , hence the sequence  $S_1, S_2, \dots, S_l, \dots$  eventually terminates. Let  $l$  be the smallest positive integer such that  $S_l = S_{l+1}$ , by (2.1) for  $k = l, l+1$ , we have

$$(M(\mathbb{A}^{l+1}))_{uj} = \sum_{j_2, \dots, j_m=1}^n a_{uj_2 \dots j_m} (M(\mathbb{A}^l))_{j_2 j} \cdots (M(\mathbb{A}^l))_{j_m j}$$

and

$$(M(\mathbb{A}^{l+2}))_{uj} = \sum_{j_2, \dots, j_m=1}^n a_{uj_2 \dots j_m} (M(\mathbb{A}^{l+1}))_{j_2 j} \cdots (M(\mathbb{A}^{l+1}))_{j_m j}.$$

It follows that  $S_{l+1} = S_{l+2} = \cdots = S_k$  for all  $k \geq l$ . Since  $\mathbb{A}$  is a nonnegative primitive tensor, hence  $S_l = S_{\gamma(\mathbb{A})} = [n]$ .

Now by the above arguments and the definition of  $l$ , we have

$$S_1 \subsetneq S_2 \subsetneq \cdots \subsetneq S_{l-1} \subsetneq S_l,$$

and thus  $2 \leq |S_1| < |S_2| < \cdots < |S_l| = n$ . It follows that  $l \leq n-1$  and  $\gamma_j(\mathbb{A}) \leq n-1$ .  $\square$

We also need the following lemmas.

**Lemma 3.2.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$ . Let  $H = \{i_1, i_2, \dots, i_s\}$  be the set of all elements  $i \in [n]$  such that  $i = k_1 \rightarrow k_2 \rightarrow \cdots \rightarrow k_t \rightarrow k_1 = i$  is a cycle with length  $t$  of  $M(\mathbb{A})$  for some  $t$  where  $1 \leq t \leq n-1$ . Suppose there exists a positive integer  $j$  with  $j \in [n] \setminus H$ , then  $1 \leq s \leq n-1$  and there exist positive integers  $i$  and  $l$  such that  $i \in H$ ,  $1 \leq l \leq n-s$  and  $(M(\mathbb{A}^l))_{ij} > 0$ .*

*Proof.* It is obvious that  $s \leq n-1$ , and  $s \geq 1$  by Lemma 2.12. By Proposition 2.7, we have the following walk of length  $n-s$  of  $M(\mathbb{A})$  starting with  $j$ :

$$j = j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_{n-s+1}.$$

If there exists an integer  $w, 1 \leq w \leq n-s+1$  such that  $j_w \in H$ , then  $(M(\mathbb{A}^{w-1}))_{j_w j} > 0$  by Lemma 2.11, and we are done. Otherwise, we have

$$\{j_1, j_2, \dots, j_{n-s+1}\} \cap H = \emptyset.$$

Since  $n-s+1 + |H| = n+1$ , so there exist two positive integers  $u$  and  $v$  such that  $1 \leq u < v \leq n-s+1$ ,  $j_u = j_v$ , and  $j_u \rightarrow j_{u+1} \rightarrow \cdots \rightarrow j_{v-1} \rightarrow j_v = j_u$  is a cycle with length  $v-u \leq n-s \leq n-1$  of  $M(\mathbb{A})$ . It follows that  $j_u \in H$ , a contradiction. This proves the lemma.  $\square$

Let  $Z(\mathbb{A})$  be the tensor obtained by replacing all the nonzero entries of  $\mathbb{A}$  by one. Then  $Z(\mathbb{A})$  is called the zero-nonzero pattern (or simply the zero pattern) of  $\mathbb{A}$ . Let  $a$  be a complex number, we define  $Z(a) = 1$  if  $a \neq 0$  and  $Z(a) = 0$  if  $a = 0$ .

**Lemma 3.3.** *Let  $\mathbb{A}$  be a nonnegative tensor with order  $m$  and dimension  $n$  such that  $a_{i_2 \dots i_m} = 0$  if  $i_2 \dots i_m \neq i_2 \dots i_2$  for any  $i \in [n]$ . Then for any positive  $r$ ,  $Z(M(\mathbb{A}^r)) = Z((M(\mathbb{A}))^r)$ .*

*Proof.* We show the result by induction on  $r$ . Clearly,  $r = 1$  is obvious.

Assume that the assertion holds for  $r = k \geq 1$ , then for any  $i, j \in [n]$ ,

$$\begin{aligned} Z[(M(\mathbb{A}^{k+1}))_{ij}] &= Z\left[\sum_{i_2, \dots, i_m=1}^n a_{ii_2 \dots i_m} (M(\mathbb{A}^k))_{i_2 j} \cdots (M(\mathbb{A}^k))_{i_m j}\right] \\ &= Z\left[\sum_{i_2=1}^n (M(\mathbb{A}))_{ii_2} ((M(\mathbb{A}^k))_{i_2 j})^{m-1}\right] \\ &= Z\left[\sum_{i_2=1}^n (M(\mathbb{A}))_{ii_2} (M(\mathbb{A}^k))_{i_2 j}\right] \\ &= Z\left[\sum_{i_2=1}^n Z[(M(\mathbb{A}))_{ii_2}] Z[(M(\mathbb{A}^k))_{i_2 j}]\right] \\ &= Z\left[\sum_{i_2=1}^n Z[(M(\mathbb{A}))_{ii_2}] Z[((M(\mathbb{A}))^k)_{i_2 j}]\right] \\ &= Z\left[\sum_{i_2=1}^n Z[(M(\mathbb{A}))_{ii_2} ((M(\mathbb{A}))^k)_{i_2 j}]\right] \\ &= Z[(M(\mathbb{A}))^{k+1}]_{ij} \end{aligned}$$

Thus  $Z(M(\mathbb{A}^{k+1})) = Z((M(\mathbb{A}))^{k+1})$ . □

**Corollary 3.4.** *Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$  such that  $a_{i_2 \dots i_m} = 0$  if  $i_2 \dots i_m \neq i_2 \dots i_2$  for any  $i \in [n]$ . If  $M(\mathbb{A})$  is primitive, then  $\gamma(\mathbb{A}) = \gamma(M(\mathbb{A}))$ .*

**Proof of Theorem 1.2:** Let  $H = \{i_1, i_2, \dots, i_s\}$  be the set of all elements  $i \in [n]$  such that  $i = k_1 \rightarrow k_2 \rightarrow \cdots \rightarrow k_t \rightarrow k_1 = i$  is a cycle with length  $t$  of  $M(\mathbb{A})$  and  $1 \leq t \leq n-1$ .

**Case 1:**  $s = n$ .

Then for any  $j \in [n]$ , there exists an integer, say,  $t_j$ , such that there exists a cycle  $j = k_1 \rightarrow k_2 \rightarrow \cdots \rightarrow k_{t_j} \rightarrow k_1 = j$  with length  $t_j$  where  $t_j \in [n-1]$ , so  $(M(\mathbb{A}^{t_j}))_{jj} > 0$ . Note that  $\mathbb{A}^{t_j}$  is primitive by Proposition 2.9 and  $\mathbb{A}$  is primitive, we have  $\gamma_j(\mathbb{A}^{t_j}) \leq n-1$  by Theorem 3.1. Hence

$$\gamma_j(\mathbb{A}) \leq t_j \gamma_j(\mathbb{A}^{t_j}) \leq (n-1)^2.$$

Thus  $\gamma(\mathbb{A}) \leq \max_{j \in [n]} \{\gamma_j(\mathbb{A})\} \leq (n-1)^2$  by Proposition 2.15.

**Case 2:**  $1 \leq s \leq n-1$ .

Then there exists at least an integer  $j \in [n] \setminus H$ .

**Subcase 2.1:**  $j \in H$ .

Similar to Case 1, for any  $j \in H$ , there exists an integer, say,  $t_j$ , such that there exists a cycle  $j = k_1 \rightarrow k_2 \rightarrow \cdots \rightarrow k_{t_j} \rightarrow k_1 = j$  with length  $t_j$  where  $t_j \leq s$ , so  $(M(\mathbb{A}^{t_j}))_{jj} > 0$ . Note that  $\mathbb{A}^{t_j}$  is primitive by Proposition 2.9 and  $\mathbb{A}$  is primitive, we have  $\gamma_j(\mathbb{A}^{t_j}) \leq n-1$  by Theorem 3.1. Hence

$$\gamma_j(\mathbb{A}) \leq t_j \gamma_j(\mathbb{A}^{t_j}) \leq t_j(n-1) \leq s(n-1).$$

Let  $T = \max_{j \in H} \{\gamma_j(\mathbb{A})\}$ , then  $T \leq s(n-1)$  and  $(M(\mathbb{A}^T))_{uj} > 0$  for any  $j \in H$  and all  $u \in [n]$ .

**Subcase 2.2:**  $j \in [n] \setminus H$ .

By Lemma 3.2, there exist positive integers  $i$  and  $l$  such that  $i \in H$ ,  $1 \leq l \leq n - s$  and  $(M(\mathbb{A}^l))_{ij} > 0$ . For any  $u \in [n]$ , since

$$(M(\mathbb{A}^{T+l}))_{uj} = \sum_{j_2, \dots, j_s=1}^n (\mathbb{A}^T)_{uj_2 \dots j_s} (M(\mathbb{A}^l))_{j_2 j} \cdots (M(\mathbb{A}^l))_{j_s j} \geq (M(\mathbb{A}^T))_{ui} ((M(\mathbb{A}^l))_{ij})^{s-1} > 0,$$

then  $\gamma_j(\mathbb{A}) \leq T + l \leq s(n - 1) + n - s \leq (n - 1)^2 + 1$ .

Thus combing Subcase 2.1 and Subcase 2.2, we have  $\gamma(\mathbb{A}) \leq \max_{j \in [n]} \{\gamma_j(\mathbb{A})\} \leq (n - 1)^2 + 1$  by Proposition 2.15.

Combing the above arguments,  $\gamma(\mathbb{A}) \leq \max\{(n - 1)^2, (n - 1)^2 + 1\} = (n - 1)^2 + 1$ .

Let  $M_1 = \begin{pmatrix} 0 & 0 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & \ddots & 0 & 0 \\ 0 & 0 & \cdots & 1 & 0 \end{pmatrix}$ . It is well known that  $M_1$  is primitive, and  $\gamma(M_1) =$

$(n - 1)^2 + 1$ .

Let  $\mathbb{A}$  be a nonnegative primitive tensor with order  $m$  and dimension  $n$  such that  $a_{i_1 i_2 \dots i_m} = 0$  if  $i_2 \cdots i_m \neq i_2 \cdots i_2$  for any  $i \in [n]$ , and  $M(\mathbb{A}) = M_1$ . Then by Corollary 3.4, we have  $\gamma(\mathbb{A}) = (n - 1)^2 + 1$ .  $\square$

**Remark 3.5.** *It is well known  $\gamma(\mathbb{A}) \leq (n - 1)^2 + 1$  for  $m = 2$  ( $\mathbb{A}$  is a matrix). It implies that the upper bound on the primitive degree of primitive tensors and the upper bound on the primitive index of primitive matrices are coincident.*

## References

- [1] K. C. Chang, K. Pearson, and T. Zhang, Perron-Frobenius theorem for nonnegative tensors, *Commun. Math. Sci.* 6(2008), 507-520.
- [2] K. C. Chang, K. Pearson, and T. Zhang, Primitivity, the convergence of the NQZ method, and the largest eigenvalue tensors, *SIAM J. Matrix Anal. Appl.* 32(2011), 806-819.
- [3] L.H. Lim, Singular values and eigenvalues of tensors, a rational approach, in *proceedings 1st IEEE international workshop on computational advances of adaptive processing (2005)*, 129-132.
- [4] K. Pearson, Essentially positive tensors, *Int. J. Algebra*, 4(2010),421-427.
- [5] K. Pearson, Primitive tensors and convergence of an iterative process for the eigenvalues of a primitive tensor, *arXiv: 1004-2423v1*, 2010.
- [6] L. Qi, Eigenvalues of a real supersymmetric tensor, *Symbolic Comput.* 40(2005), 1302-1324.
- [7] J.Y Shao, A general product of tensors with applications, *Linear Algebra and its Appl.* 439(2013), 2350-2366.