

## Local derivations on Rings containing a von Neumann algebra and a question of Kadison.

Don Hadwin, Jiankui Li, Qihui Li, and Xiujuan Ma

ABSTRACT. We prove that if  $\mathcal{M}$  is a von Neumann algebra whose abelian summand is discrete, then every local derivation on the algebra of all measurable operators affiliated with  $\mathcal{M}$  is a derivation. This answers a question of Richard Kadison.

At a conference held in Texas A & M University of 2012, Richard Kadison gave a talk about his joint work with Zhe Liu [10, 11] in which they proved that the only derivation that maps the algebra  $S(\mathcal{M})$  of closed densely defined operators affiliated with a factor von Neumann algebra  $\mathcal{M}$  of type  $\text{II}_1$  into that von Neumann algebra is zero. Kadison asked whether every local derivation on  $S(\mathcal{M})$  is a derivation.

In this note we prove a general ring-theoretic result which implies that for all von Neumann algebras  $\mathcal{M}$  whose abelian summand is discrete, every local derivation on  $S(\mathcal{M})$  of all measurable operators is a derivation.

If  $\mathcal{R}$  is a ring (resp. algebra) and  $\delta : \mathcal{R} \rightarrow \mathcal{R}$  is an additive (resp. linear) mapping, we say that  $\delta$  is a *derivation* if, for all  $a, b \in \mathcal{R}$ , we have

$$\delta(ab) = \delta(a)b + a\delta(b).$$

We say that an additive mapping  $\delta$  is a *local derivation* if, for every  $x \in \mathcal{R}$  there is a derivation  $\rho_x$  on  $\mathcal{R}$  such that

$$\delta(x) = \rho_x(x).$$

To prove our main result, we need two lemmas. The first is a result of [3, Corollary 4.5]. Suppose  $\mathcal{R}$  is a ring with identity 1 and  $n \geq 2$  is an integer. We say that a subset  $\{E_{ij} : 1 \leq i, j \leq n\} \subset \mathcal{R}$  is a *system of  $n \times n$  matrix units for  $\mathcal{R}$*  if and only if  $\sum_{i=1}^n E_{ii} = 1$  and, for  $1 \leq i, j, s, t \leq n$ ,  $E_{ij}E_{st} = 0$  if  $j \neq s$  and  $E_{ij}E_{st} = E_{it}$  if  $j = s$ .

In [3, p.11], Bresar shows that for any unital ring  $\mathcal{B}$ , the ring  $M_n(\mathcal{B})$  is generated by the set of all idempotents in  $\mathcal{B}$ , where  $2 \leq n$ .

The following Lemma 1 is a special case of [3, Corollary 4.5].

LEMMA 1. *If  $\mathcal{R}$  is a ring with identity 1 that possesses a system of  $n \times n$  matrix units for some  $n \geq 2$ , then every local derivation on  $\mathcal{R}$  is a derivation.*

---

2000 *Mathematics Subject Classification*. Primary 46L57, 13N15; Secondary 16W25.

*Key words and phrases*. local derivation, derivation, affiliated algebra, von Neumann algebra.

Note that, in general, derivations need not leave ideals invariant, e.g., differentiation on the polynomials. However, if  $p$  a central idempotent, then every derivation (hence, every local derivation)  $\delta$  leaves  $p\mathcal{R}$  and  $(1-p)\mathcal{R}$  invariant, since

$$\delta(pa) = \delta((pa)p) = pa\delta(p) + \delta(pa)p.$$

Moreover,  $p\mathcal{R}$  is isomorphic to  $\mathcal{R}/(1-p)\mathcal{R}$ . This yields the following corollary to the preceding lemma. A family  $\mathcal{P}$  of central idempotents for a ring  $\mathcal{R}$  is *separating* if and only if, for each nonzero  $x \in \mathcal{R}$ , there is a  $p \in \mathcal{P}$  such that  $px \neq 0$ .

LEMMA 2. *Suppose  $\mathcal{R}$  is a ring with identity and  $\mathcal{P}$  is a separating family of central idempotents such that, for each  $p \in \mathcal{P}$ , every local derivation on  $p\mathcal{R}$  is a derivation. Then every local derivation on  $\mathcal{R}$  is a derivation.*

PROOF. Suppose  $\delta : \mathcal{R} \rightarrow \mathcal{R}$  is a local derivation,  $a \in \mathcal{R}$  and  $p \in \mathcal{P}$ . Then there is a derivation  $\rho$  on  $\mathcal{R}$  such that

$$\delta(pa) = \rho(pa) = \rho(p(pa)) = p\rho(pa) + \rho(p)pa = p[\rho(pa) + \rho(p)a] = pgr(pa) = p\delta(pa).$$

It follows that  $\delta(p\mathcal{R}) \subset p\mathcal{R}$  and that  $\delta|_{p\mathcal{R}}$  is a local derivation. Hence  $\delta|_{p\mathcal{R}}$  is a derivation. Thus, for every  $a, b \in \mathcal{R}$  and every  $p \in \mathcal{P}$ , we have

$$\begin{aligned} p\delta(ab) &= \delta(pab) = \delta((pa)(pb)) = \\ &= pa\delta(pb) + \delta(pa)pb = p[a\delta(b) + \delta(a)b]. \end{aligned}$$

Hence

$$p[\delta(ab) - [a\delta(b) + \delta(a)b]] = 0.$$

Since  $\mathcal{P}$  is separating, we see that  $\delta$  is a derivation on  $\mathcal{R}$ . □

An abelian von Neumann algebra  $\mathcal{M}$  is discrete if it is generated by its minimal nonzero projections; equivalently, if the identity operator in  $\mathcal{M}$  is the sum of the minimal projections in  $\mathcal{M}$ . Since  $\mathcal{M}$  is abelian, it follows that  $Q\mathcal{M} = \mathbb{C}Q$  for every minimal projection  $Q$  in  $\mathcal{M}$ . Every von Neumann algebra on a Hilbert space  $H$  is the direct sum of algebras  $\mathcal{M}_n$  with  $1 \leq n < \infty$  (the finite type  $I_n$  summands) and a von Neumann algebra  $\mathcal{M}_\infty$  which is the direct sum of algebras of type  $I_\infty$ ,  $II$ , and  $III$  (not all summands need be present). Call the corresponding central projections  $P_n$  with  $1 \leq n \leq \infty$ .

THEOREM 1. *Suppose  $\mathcal{M}$  is a von Neumann algebra on a Hilbert space  $H$ . Then*

- (1) *If  $\mathcal{M}_1 = 0$  and  $\mathcal{R}$  is a ring containing  $\mathcal{M}$  with the same identity as  $\mathcal{M}$  such that  $\mathcal{P} = \{P_n : 2 \leq n \leq \infty\}$  is a separating family of central idempotents for  $\mathcal{R}$ , then every local derivation on  $\mathcal{R}$  is a derivation.*
- (2) *If  $\mathcal{M}_1$  is discrete, then every local derivation on the algebra of closed densely defined operators affiliated with  $\mathcal{M}$  is a derivation.*

PROOF. (1).

By [9, Theorem 6.6.5], we know that  $P_n\mathcal{M}$  contains an  $n \times n$  system of matrix units for  $2 \leq n < \infty$ , and it follows from [9, Lemma 6.5.6] that  $P_\infty\mathcal{M}$  contains a  $2 \times 2$  system of matrix units. Since  $P_n\mathcal{M} \subset P_n\mathcal{R}$  is a unital embedding and  $\mathcal{P}$  is separating, it follows from the two lemmas above that every local derivation on  $\mathcal{R}$  is a derivation.

(2). Let  $\mathcal{R}$  be the algebra of all measurable operators affiliated with  $\mathcal{M}$ . However, since  $\mathcal{M}_1$  is discrete,  $P_1$  is the orthogonal sum of a family  $\{Q_\lambda : \lambda \in \Lambda\}$  of minimal projections and that, for each  $\lambda \in \Lambda$ ,  $Q_\lambda \mathcal{M} = \mathbb{C}Q_\lambda$ , which means that  $Q_\lambda \mathcal{M}' = B(Q_\lambda H)Q_\lambda$ , which, in turn, implies that  $Q_\lambda \mathcal{R} = \mathbb{C}Q_\lambda$ , so every local derivation on  $Q_\lambda \mathcal{R}$  is a derivation. Since the elements of  $\mathcal{R}$  are densely defined operators on  $H$ , and  $\sum_{\lambda \in \Lambda} Q_\lambda + \sum_{2 \leq n \leq \infty} P_n = 1$ , it follows that  $\{Q_\lambda : \lambda \in \Lambda\} \cup \{P_n : 2 \leq n \leq \infty\}$  is a separating family of central idempotents for  $\mathcal{R}$ . Arguing as in the proof of (1) we can apply the lemmas to see that every local derivation on  $\mathcal{R}$  is a derivation.  $\square$

In [1, Theorem 3.8], the authors give necessary and sufficient conditions on a commutative von Neumann algebra  $\mathcal{M}$  for the existence of local derivations which are not derivations on the algebra  $S(\mathcal{R})$  of measurable operators affiliated with  $\mathcal{M}$ .

For a von Neumann algebra  $\mathcal{M}$ , we can define the set  $S(\mathcal{M})$  of all measurable operators affiliated with  $\mathcal{M}$  and the set  $LS(\mathcal{M})$  of all local measurable operators affiliated with  $\mathcal{M}$ .

In [12], Muratov and Chilin show that  $LS(\mathcal{M})$  is a unital  $*$ -algebra when equipped with algebraic operations of the strong addition, multiplication, and taking the adjoint of an operator and  $S(\mathcal{S})$  is a unital  $*$ -subalgebra of  $LS(\mathcal{M})$ .

Suppose that  $\mathcal{M}$  is a von Neumann algebra with a faithful normal semi-finite trace  $\tau$ . Let  $S(\mathcal{M}, \tau)$  denote the algebra of all  $\tau$ -measurable operators affiliated with  $\mathcal{M}$ . It is clear that  $\mathcal{M} \subseteq S(\mathcal{M}) \subseteq LS(\mathcal{M})$  ( see for details [1, 2, 13]).

In Theorem 1, if we choose that  $\mathcal{R}$  is  $\mathcal{M}$ ,  $S(\mathcal{M})$  or  $LS(\mathcal{M})$ , we can obtain [1, Theorem 2.5 and Proposition 2.7].

By Theorem 1 and [1, Theorem 3.8], we can completely answer [15, Conjecture 48].

### Acknowledgement

The first author is partially supported by a travel grant from the Simons Foundation and the second author is supported by NSF of China.

### References

- [1] S. Albeverio, Sh. A. Ayupov, K. K. Kudaybergenov and B. O. Nurjanov, *Local derivations on algebras of measurable operators*, Commun. in Contemp. Math. **13(4)** (2011), 643–657.
- [2] Sh. A. Ayupov and K. K. Kudaybergenov, *Derivations on algebras of measurable operators*, Infin. Dimens. Ana. Quantum Probab. Relat. Top. **13(2)** (2010), 305–337.
- [3] M. Bresar, *Characterizing homomorphisms, derivations and multipliers in rings with idempotents*, Proc. Roy. Soc. Edinburgh Sect.A **137** (2007), 9–21.
- [4] D. Hadwin and J. Li, *Local derivations and local automorphisms*, J. Math. Anal. Appl. **290** (2003), 702–714.
- [5] D. Hadwin and J. Li, *Local derivations and local automorphisms on some algebras*, J. Oper. Theory **60** (2008), 29–44.
- [6] B. Johnson, *Local derivations on  $C^*$ -algebras are derivations*, Trans. Amer. Math. Soc. **353** (2001), 313–325.
- [7] B. Johnson, *Symmetric amenability and the nonexistence of Lie and Jordan derivations*, Math. Proc. Cambridge Philos. Soc. **120** (1996), 455–473.
- [8] R. Kadison, *Local derivations*, J. Algebra **130** (1990), 494–509.
- [9] R. Kadison and J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras II*, Graduate Studies in Math. 16.
- [10] R. Kadison and Zhe Liu, *Derivations of Murray-von Neumann algebras*, Math. Scand. To appear.

- [11] R. Kadison and Zhe Liu, *A note on derivations of Murray-von Neumann algebras*, To appear.
- [12] M. A. Muratov and V. I. Chilin, *Algebras of Measurable and Local Measurable Operators*, (Ukrainian Academy of Sciences, 2007).
- [13] M. A. Muratov and V. I. Chilin, *\*-algebras of unbounded operators affiliated with a von Neumann algebra*, J. Math. Sci. **140** (2007), 445–451.
- [14] I. E. Segal, *A non-commutative extension of abstract integration*, Ann. Math. **57** (1953), 401–457.
- [15] <http://www.math.tamu.edu/~kerr/workshop/problems 2012.pdf>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEW HAMPSHIRE, DURHAM, NH 03824, USA  
*E-mail address:* don@unh.edu  
*URL:* <http://euclid.unh.edu/~don>

DEPARTMENT OF MATHEMATICS, EAST CHINA UNIVERSITY OF SCIENCE AND TECHNOLOGY,  
SHANGHAI 200237, CHINA  
*E-mail address:* jiankuili@yahoo.com

DEPARTMENT OF MATHEMATICS, EAST CHINA UNIVERSITY OF SCIENCE AND TECHNOLOGY,  
SHANGHAI 200237, CHINA  
*E-mail address:* lqh991978@gmail.com

DEPARTMENT OF MATHEMATICS, HEBEI UNIVERSITY OF TECHNOLOGY, TIANJING, 300130,  
CHINA  
*E-mail address:* mxjsusan@hebut.edu.cn