

On enumeration of a class of toroidal graphs

DIPENDU MAITY¹, ASHISH KUMAR UPADHYAY²

^{1,2}*Department of Mathematics, Indian Institute of Technology Patna, Bihta, 801 103, India.*

¹*dipendumaity@gmail.com, ²upadhyay@iitp.ac.in.*

April 21, 2017

Abstract

We present enumerations of a class of toroidal graphs which give rise to semi-equivelar maps. There are eleven different types of semi-equivelar maps on the torus. These are of the types $\{3^6\}$, $\{4^4\}$, $\{6^3\}$, $\{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3^4, 6\}$, $\{4, 8^2\}$, $\{3, 12^2\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$. We know the classification of the maps of types $\{3^6\}$, $\{4^4\}$, $\{6^3\}$ on the torus. In this article, we attempt to classify maps of types $\{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3^4, 6\}$, $\{4, 8^2\}$, $\{3, 12^2\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$ on the torus.

AMS classification : 52B70, 05C30, 05C38

Keywords : Toroidal Graphs, Semi-Equivelar Maps, Cycles

1 Introduction

A map M is an embedding of a graph G on a surface S such that the closure of components of $S \setminus G$, called the faces of M , are closed 2-cells, that is, each homeomorphic to 2-disk. A map M is said to be a polyhedral map (see Brehm and Schulte [3]) if the intersection of any two distinct faces is either empty, a common vertex, or a common edge. A a -cycle C_a is a finite connected 2-regular graph with a vertices, and the *face sequence* of a vertex v in a map is a finite sequence (a^p, b^q, \dots, m^r) of powers of positive integers $a, b, \dots, m \geq 3$ and $p, q, \dots, r \geq 1$ in *cyclic* order such that through the vertex v , p number of C_a (C_a denote the a -cycle), q number of C_b , \dots , r number of C_m are incident. A map K is said to be *semi-equivelar* if face sequence of each vertex is same, see [8]. Two maps of fixed type, on the torus, are called *isomorphic* if there exists a *homeomorphism* of the torus which sends vertices to vertices, edges to edges, faces to faces and preserves incidents. That is, if we consider two polyhedral complexes K_1 and K_2 then an isomorphism to be a map $f : K_1 \rightarrow K_2$ such that $f|_{V(K_1)} : V(K_1) \rightarrow V(K_2)$ is a bijection and $f(\sigma)$ is a cell in K_2 if and only if σ is a cell in K_1 . There are eleven types $\{3^6\}$, $\{4^4\}$, $\{6^3\}$, $\{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3^4, 6\}$, $\{4, 8^2\}$, $\{3, 12^2\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$ of semi-equivelar maps on the torus. In this article, we are interested to classify some of them up to isomorphism. It completes the classification of semi-equivelar maps on the torus. In this context, Altshuler [1] has shown construction and enumeration of maps of types $\{3^6\}$ and $\{6^3\}$ on the torus. Kurth [5] has given an enumeration of semi-equivelar maps of types $\{3^6\}$, $\{4^4\}$, $\{6^3\}$ on the torus. Negami [6] has studied uniqueness and faithfulness of embedding of a class of toroidal graphs. Brehm and Kühnel [2] have presented a classification of semi-equivelar maps of types $\{3^6\}$, $\{4^4\}$, $\{6^3\}$ on the torus. Tiwari and Upadhyay [7] have classified semi-equivelar maps of types $\{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3^4, 6\}$, $\{4, 8^2\}$, $\{3, 12^2\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$ on torus with up to

twenty vertices. In this article, we devise a way of enumerating all semi-equivelar maps of types $\{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3^4, 6\}$, $\{4, 8^2\}$, $\{3, 12^2\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$ on the torus and explicitly determine the maps with a small number of vertices. Therefore, we have the following theorem.

Theorem 1.1. *The semi-equivelar maps with n vertices of types $\{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3, 12^2\}$, $\{3^4, 6\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$, $\{4, 8^2\}$ can be classified up to isomorphism on the torus. In Table 1, 2, 3, 4, 5, 6, 7, 8, we have given non-isomorphic objects for few vertices.*

More precisely, let $X = \{3^3, 4^2\}$, $\{3^2, 4, 3, 4\}$, $\{3, 6, 3, 6\}$, $\{3, 12^2\}$, $\{3^4, 6\}$, $\{4, 6, 12\}$, $\{3, 4, 6, 4\}$ or $\{4, 8^2\}$ be a semi-equivelar type on the torus. We present an algorithmic approach of calculating different maps for the type X on different number of vertices in the subsequent sections.

2 Definitions

We now define some operations on graphs. Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two subgraphs of the same graph $G := (V, E)$. Then *union* $G_1 \cup G_2$ is a graph $G_3 = (V_3, E_3)$ where $V_3 = V_1 \cup V_2$ and $E_3 = E_1 \cup E_2$. Similarly, the *intersection* $G_1 \cap G_2$ is a graph $G_4 = (V_4, E_4)$ where $V_4 = V_1 \cap V_2$ and $E_4 = E_1 \cap E_2$. For more on graph theory see [4].

We denote a cycle $u_1-u_2-\dots-u_k-u_1$ by $C(u_1, u_2, \dots, u_k)$ and a path $w_1-w_2-\dots-w_x$ by $P(w_1, w_2, \dots, w_x)$. Let $Q_1 = P(u_1, \dots, u_k)$ be a path. We call a path $Q_2 = P(v_1, \dots, v_r)$ to be a path extended from Q_1 if $V(Q_1) \subset V(Q_2)$, $E(Q_1) \subset E(Q_2)$, i.e., Q_1 is a subpath of Q_2 . We say that the Q_2 is an *extended path* of the path Q_1 .

We say that a cycle is *contractible* if it bounds a 2-disk. If the cycle does not bound any 2-disk on the torus then we say that the cycle is *non-contractible*.

3 Examples

Example 2.1 : Let M be a semi-equivelar map of type $\{3^3, 4^2\}$ with n vertices on the torus. The map M has a $T(r, s, k)$ representation (defined later in Section 4) for some $r, s, k \in \mathbb{N} \cup \{0\}$. Let the number of vertices $n = 14$. We get by Lemma 4.8, $n = rs = 14$ where $2 \mid s$. Hence, $s = 2$, $r = 7$ and $k = 2, 3, 4$ by Lemma 4.8. So, $T(r, s, k) = T(7, 2, 2)$, $T(7, 2, 3)$ and $T(7, 2, 4)$, see Figure 1, 2, 3 respectively. In $T(7, 2, 2)$, $C_{1,1} = C(u_1, u_2, \dots, u_7)$ is a cycle of type A_1 (see definition of type A_1 in Section 4), $C_{1,2} = C(u_1, u_8, u_3, u_{10}, u_5, u_{12}, u_7, u_{14}, u_2, u_9, u_4, u_{11}, u_6, u_{13})$ and $C_{1,3} = C(u_1, u_8, u_4, u_{11}, u_7, u_{14}, u_3, u_{10}, u_6, u_{13}, u_2, u_9, u_5, u_{12})$ are two cycles of type A_2 (see definition of type A_2 in Section 4) and $C_{1,4} = C(u_3, u_{10}, u_5, u_4)$ is a cycle of type A_4 (see definition of type A_4 in Section 4). In $T(7, 2, 4)$, $C_{2,1} = C(v_1, v_2, \dots, v_7)$ is of type A_1 , $C_{2,2} = C(v_1, v_8, v_5, v_{12}, v_2, v_9, v_6, v_{13}, v_3, v_{10}, v_7, v_{14}, v_4, v_{11})$ and $C_{2,3} = C(v_1, v_8, v_6, v_{13}, v_4, v_{11}, v_2, v_9, v_7, v_{14}, v_5, v_{12}, v_3, v_{10})$ are of type A_2 , and $C_{2,4} = C(v_5, v_{12}, v_3, v_4)$ is of type A_4 . In $T(7, 2, 3)$, $C_{3,1} = C(w_1, w_2, \dots, w_7)$ is of type A_1 , $C_{3,2} = C(w_1, w_8, w_4, w_{11}, w_7, w_{14}, w_3, w_{10}, w_6, w_{13}, w_2, w_9, w_5, w_{12})$ and $C_{3,3} = C(w_1, w_8, w_5, w_{12}, w_2, w_9, w_6, w_{13}, w_3, w_{10}, w_7, w_{14}, w_4, w_{11})$ are of type A_2 and $C_{3,4} = C(w_4, w_5, w_6, w_7, w_{11})$ is of type A_4 .

In Section 4, by Lemma 4.7, the cycles of type A_1 have same length and the cycles of type A_2 have at most two different lengths in M . So, $O_3 \not\cong O_1$ since $\text{length}(C_{3,4}) \neq \text{length}(C_{1,4})$ and $O_3 \not\cong O_2$ since $\text{length}(C_{3,4}) \neq \text{length}(C_{2,4})$. Thus, $O_3 \not\cong O_i$ for $i = 1, 2$. Now,

$\text{length}(C_{1,1}) = \text{length}(C_{2,1}), \{\text{length}(C_{1,2}), \text{length}(C_{1,3})\} = \{\text{length}(C_{2,2}), \text{length}(C_{2,3})\}$ and $\text{length}(C_{1,4}) = \text{length}(C_{2,4})$. We cut $T(7, 2, 4)$ along the path $P(v_5, v_{12}, v_3)$ and identify along the path $P(v_1, v_8, v_5)$. This gives a presentation of $T(7, 2, 4)$ in Figure 4. The Figure 4 has $T(7, 2, 2)$ representation. So, $O_1 \cong O_2$, (see the proof of Lemma 4.9 for isomorphism map between O_1 and O_2). Thus, $O_1 \cong O_2$ and $O_3 \not\cong O_1, O_2$. Therefore, there are two semi-equivelar maps of type $\{3^3, 4^2\}$ with 14 vertices on the torus up to isomorphism.




Figure 1 : $T(7, 2, 2) : O_1$




Figure 2 : $T(7, 2, 4) : O_2$




Figure 3 : $T(7, 2, 3) : O_3$




Figure 4 : $T(7, 2, 2) : O_2$




Figure 6 : $T(4, 4, 0) : O_6$




Figure 7 : $T(4, 4, 2) : O_7$




Figure 8 : $T(8, 2, 4) : O_4$




Figure 9 : $T(8, 2, 4) : O_5$




Figure 10 : $T(8, 2, 6)$




Figure 5 : $T(6, 4, 2)$




Figure 11 : $T(7, 4, 3)$




Figure 12 : R

Example 2.2 : Let M be a semi-equivelar map of type $\{3^2, 4, 3, 4\}$ with 16 vertices on the torus. Similarly as above, by Lemma 5.5, there are three representations of M , namely, $T(8, 2, 4)$, $T(4, 4, 0)$, and $T(4, 4, 2)$, see Figure 9, 6, 7 respectively. In $T(8, 2, 4)$, $C_{1,1} = C(u_1, u_2, \dots, u_8)$ and $C_{1,2} = C(u_1, u_9, u_5, u_{13})$ are two cycles of type B_1 (see definition of type B_1 in Section 5). In $T(4, 4, 0)$, $C_{2,1} = C(v_1, v_2, v_3, v_4)$ and $C_{2,2} = C(v_1, v_5, v_9, v_{13})$ are two cycles of type B_1 . In $T(4, 4, 2)$, $C_{3,1} = C(w_1, w_2, w_3, w_4)$ and $C_{3,2} = C(w_1, w_5, w_9, w_{13}, w_3, w_7, w_{11}, w_{15})$ are two cycles of type B_1 . In Section 5, the cycles of type B_1 have at most two different lengths. So, $O_5 \not\cong O_6$ since $\{\text{length}(C_{1,1}), \text{length}(C_{1,2})\} \neq \{\text{length}(C_{2,1}), \text{length}(C_{2,2})\}$. Now, $\{\text{length}(C_{1,1}), \text{length}(C_{1,2})\} = \{\text{length}(C_{3,1}), \text{length}(C_{3,2})\}$. We identify boundaries of O_7 and cut along the cycle $C_{3,2} = C(w_1, w_5, w_9, w_{13}, w_3, w_7, w_{11}, w_{15})$ and next along $C_{3,1}$. Thus, we get a $T(8, 2, 4)$ representation in Figure 8. So, $O_5 \cong O_7$ (see the proof of Lemma 5.6 for isomorphism map). Therefore, we have two maps of type $\{3^2, 4, 3, 4\}$ with 16 vertices on the torus up to isomorphism.

4 Maps of type $\{3^3, 4^2\}$

Let M be a map of type $\{3^3, 4^2\}$ on the torus. Through each vertex in M there are three distinct types of paths as follows.

Definition 4.1. Let $P_1 := P(\dots, u_{i-1}, u_i, u_{i+1}, \dots)$ be a path in edge graph of M . We say P_1 of type A_1 if all the triangles incident with an inner (degree two in P_1) vertex u_i lie on one side and all quadrangles incident with u_i lie on the other side of the subpath $P' = P(u_{i-1}, u_i, u_{i+1})$ (as in Figure 13) at u_i . Since the link of vertex u_i is a cycle and the path P' is a cord of cycle $lk(u_i)$, so, the path P' divides the region into two parts. If u_t is a boundary vertex (degree one in P_1) of P_1 then there is an extended path of P_1 where u_t is an inner vertex.

Observe that the link of a vertex in M contains some vertices which are adjacent to the vertex. So, to identify the non adjacent vertices in the link we use bold letters. That is, if a description of link, say $lk(w)$ contains any bold letter \mathbf{a} then it indicates non adjacent vertex to w . For example, in $lk(u_i)$ vertices \mathbf{a} and \mathbf{c} are non adjacent to u_i . In this article, we consider permutation of vertices in $lk(u_i)$ of a vertex u_i counter clockwise locally at u_i .




Figure 13 : $lk(u_i)$




Figure 14 : Cylinder




Figure 16




Figure 15 : $T(r, 4, k)$

Definition 4.2. Let $P_2 := P(\dots, v_{i-1}, v_i, v_{i+1}, \dots)$ be a path in edge graph of M for which v_i, v_{i+1} are two consecutive inner vertices of P_2 or an extended path of P_2 . We say P_2 of type A_2 if $lk(v_i) = C(\mathbf{a}, v_{i-1}, \mathbf{b}, c, v_{i+1}, d, e)$ implies $lk(v_{i+1}) = C(\mathbf{a}_0, v_{i+2}, \mathbf{b}_0, d, v_i, c, p)$ and $lk(v_i) = C(\mathbf{x}, v_{i+1}, \mathbf{z}, l, v_{i-1}, k, m)$ implies $lk(v_{i+1}) = C(\mathbf{l}, v_i, \mathbf{m}, x, v_{i+2}, g, z)$. At least one of the former two conditions must occur for each vertex.

Definition 4.3. Let $P_3 := P(\cdots, w_{i-1}, w_i, w_{i+1}, \cdots)$ be a path in edge graph of M for which w_i, w_{i+1} are two inner vertices of P_3 or an extended path of P_3 . We say P_3 of type A_3 if $lk(w_i) = C(\mathbf{a}, w_{i-1}, \mathbf{b}, c, d, w_{i+1}, e)$ implies $lk(w_{i+1}) = C(\mathbf{a}_1, w_{i+2}, \mathbf{b}_1, p, e, w_i, d)$ and $lk(w_i) = C(\mathbf{a}_2, w_{i+1}, \mathbf{b}_2, p, e, w_{i-1}, d)$ implies $lk(w_{i+1}) = C(\mathbf{p}, w_i, \mathbf{d}, a_2, z_1, w_{i+2}, b_2)$.

Let Q be a maximal path (path of maximal length) of type A_t for a fixed $t \in \{1, 2, 3\}$. We show that there is an edge e in M such that $Q \cup e$ is a cycle of type A_t . So,

Lemma 4.1. If $P(u_1, \cdots, u_r)$ is a maximal path of type A_1, A_2 or A_3 in M then there is an edge $u_r u_1$ in M such that $C(u_1, u_2, \cdots, u_r)$ is a cycle.

Proof. Let $Q = P(u_1, \cdots, u_r)$ be of type A_1 and $lk(u_r) = C(\mathbf{x}, y, \mathbf{z}, w, v, u, u_{r-1})$.

If $w = u_1$ then $Q = C(u_1, u_2, \cdots, u_r)$ is a cycle. If $w \neq u_1$. Then, either $w = u_i$ for some $2 \leq i \leq r$ or $w \neq u_i$ for all $2 \leq i \leq r$. Suppose $w = u_i$ for some $2 \leq i \leq r$. Observe that $L = P(u_{i-1}, u_i, u_{i+1}) \subset Q$ and $L' = P(u_r, w, x)$ are two paths of type A_1 through u_i . By Definition 4.1, through each vertex in M we have only one path of this particular type A_1 . So, $L = L'$. This implies that $u_r = u_{i-1}$ or $u_r = u_{i+1}$. This is a contraction since, by assumption, Q is a path and $u_i \neq u_j$ for all $i \neq j, 1 \leq i, j \leq r$. Therefore, $w \neq u_i$ for all $2 \leq i \leq r$. So, if $w \neq u_i$ for all $1 \leq i \leq r$ then by the Definition 4.1, u_r is an inner vertex in the extended path of Q . Thus, we get a path namely Q_1 which is extended from Q . Hence, $\text{length}(Q) < \text{length}(Q_1)$. This is a contradiction as Q is maximal. Therefore, $w = u_1$ and $Q \cup u_r u_1 = C(u_1, u_2, \cdots, u_r)$ is a cycle.

Let $W = P(u_1, u_2, \cdots, u_r)$ be of type A_2 . We follow similar argument as in *Theorem 1* [1]. Let $lk(u_{r-1}) = (\mathbf{v}, u_r, \mathbf{z}, p, q, u_{r-2}, x)$ and $lk(u_r) = (\mathbf{p}, u_{r-1}, \mathbf{x}, v, e, u_{r+1}, z)$.

If $u_{r+1} = u_1$ then $C(u_1, u_2, \cdots, u_r)$ is a cycle. If $u_{r+1} \neq u_1$. Then, either $u_{r+1} = u_i$ for some $2 \leq i \leq r$ or $u_{r+1} \neq u_i$ for all $2 \leq i \leq r$. Suppose $u_{r+1} = u_i$ for some $2 \leq i \leq r$. Then, $u_{r+1} = u_i$ defines a cycle $R = C(u_i, u_{i+1}, \cdots, u_r)$. Now by assumption, W is a path. That is, $u_i \neq u_j$ for all $1 \leq i, j \leq r$ and $i \neq j$. By Definition 4.2, through each vertex in M we have exactly two paths of type A_2 . Hence, we have either $lk(u_i) = C(\mathbf{a}, u_{i-1}, \mathbf{b}, c, u_{i+1}, d, e)$ or $lk(u_i) = C(\mathbf{a}, u_{i-1}, \mathbf{b}, c, z, u_{i+1}, e)$. If $lk(u_i) = C(\mathbf{a}, u_{i-1}, \mathbf{b}, c, z, u_{i+1}, e)$ then $u_{i+1} = u_r$. But, $u_i \neq u_j$ for all $1 \leq i, j \leq r$ and $i \neq j$. Hence, $lk(u_i) = C(\mathbf{a}, u_{i-1}, \mathbf{b}, c, u_{i+1}, d, e)$. Thus, from the cycles $lk(u_r)$ and $lk(u_i)$, $z = u_{i+1}$, $p = u_{i+2}$, $d = u_r$ and $u_i u_{i+1} u_r$ is a triangle (see Figure 16). Consider cycle R and faces incident to it. So, these faces $u_{r-3} w_i w_{r-8}$, $u_{r-3} w_{r-8} u_{r-4}$, $[u_{r-4}, w_{r-8}, w_{r-9}, u_{r-5}]$, $u_{r-5} w_{r-9} w_{r-10}$, $u_{r-5} w_{r-10} u_{r-6}$, \cdots , $w_{i+3} u_i u_{i+6}$, $w_{i+3} u_{i+6} w_{i+2}$, $[w_{i+2}, u_{i+6}, u_{i+5}, w_{i+1}]$, $w_{i+1} u_{i+5} u_{i+4}$, $w_{i+1} u_{i+4} w_i$ define a new cycle $R' = C(w_i, w_{i+1}, \cdots, w_{r-9}, w_{r-8})$ (see Figure 16). Observe that, R' is a cycle of same type as R since the faces $[u_{r-2}, u_{r-3}, w_i, w_{i-1}]$ and $[u_{i+3}, u_{i+4}, w_i, w_{i-1}]$ have a common edge $w_{i-1} w_i$, and $\text{length}(R') < \text{length}(R)$. Similarly, we consider cycle R' and repeat the process as above. Thus, we get a sequence of cycles of same type as R . But, in this sequence, the length of cycles is gradually decreasing. So, after finite number of steps cycle of type as R may not exist since the map is finite. Therefore, $u_{r+1} \neq u_i$ for all $2 \leq i \leq r$. By Definition 4.2, $lk(u_r) = (\mathbf{z}, u_{r-1}, \mathbf{x}, v, u_{r+1}, w)$ implies $lk(u_{r+1}) = C(\mathbf{y}, u_r, \mathbf{x}, a, w, b, c)$ for some vertices b, w . Hence, we define a new path $L := P(u_1, \cdots, u_r) \cup P(u_r, u_{r+1})$ which is of type A_2 . So, we have a path Q with $\text{length}(Q) > \text{length}(P)$. This gives a contradiction as P is maximal. Therefore, $u_{r+1} = u_1$, that is, $C(u_1, u_2, \cdots, u_r)$ is cycle of type A_2 .

We use similar argument as above for maximal path of type A_3 . Similarly, we get an edge which defines a cycle of type A_3 . This completes the proof. \square

So, every maximal path of type A_1, A_2 or A_3 is a cycle. In this article, we use the

terminology cycle in place of maximal path since it is a cycle. Let C_1 and C_2 be two cycles of type A_t for a fixed $t \in \{1, 2, 3\}$. Then,

Lemma 4.2. (a) If C_1, C_2 are two cycles of same type A_1 such that $C_1 \cap C_2 \neq \emptyset$ then $C_1 = C_2$. (b) If C'_1, C'_2 are two cycles of same type A_t for a fixed $t \in \{2, 3\}$ such that $E(C'_1) \cap E(C'_2) \neq \emptyset$ then $C'_1 = C'_2$.

Proof. Let $C_1 := C(u_{1,1}, u_{1,2}, \dots, u_{1,r})$ and $C_2 := C(u_{2,1}, u_{2,2}, \dots, u_{2,s})$ be two cycles of type A_1 . If $C_1 \cap C_2 \neq \emptyset$ then $V(C_1 \cap C_2) \neq \emptyset$. Let $w \in V(C_1 \cap C_2)$. The cycles C_1 and C_2 are both well defined at the common vertex w . Let $lk(w) = C(w_1, w_2, w_3, w_4, w_5, w_6, w_7)$. By Definition 4.1, $w_4, w_7 \in V(C_1 \cap C_2)$. So, $P(w_4, w, w_7)$ is part of C_t for $t \in \{1, 2\}$. Let $w = u_{1,t_1} = u_{2,t_2}$. Then $w_4 = u_{1,t_1-1} = u_{2,t_2-1}$ and $w_7 = u_{1,t_1+1} = u_{2,t_2+1}$ for some $t_1 \in \{1, \dots, r\}$ and $t_2 \in \{1, \dots, s\}$. We can argue for w_4 and w_7 as we did for w to get two vertices, $u_{1,t_1-2} = u_{2,t_2-2}$ and $u_{1,t_1+2} = u_{2,t_2+2}$. This process stops after finite number of steps as r and s both are finite. Let $r < s$. Then $u_{1,1} = u_{2,I+1}, u_{1,2} = u_{2,I+2}, \dots, u_{1,r} = u_{2,I+r}$ and $u_{1,1} = u_{2,I+r+1}$ for some $I \in \{1, \dots, s\}$. Hence $u_{1,1} = u_{2,I+1} = u_{2,I+r+1}$. This implies that $I+1 = I+r+1$ and the cycle C_2 contains a cycle of length r . This gives $r = s$. Hence $C_1 = C_2$.

Let C'_1, C'_2 be two cycles of same type A_2 and $E(C'_1) \cap E(C'_2) \neq \emptyset$. Let $uv \in E(C'_1) \cap E(C'_2)$. We proceed with the vertex u of edge uv and in a similar way as we did for the cycles of type A_1 . (This argument we have also used in Lemma 5.2.) Thus, we get $C'_1 = C'_2$. Similarly we argue for the case of cycles of type A_3 to show that $C'_1 = C'_2$. This completes the proof. \square

Now, we show that the cycle of type A_t for each $t \in \{1, 2, 3\}$ is non-contractible.

Lemma 4.3. If a cycle C is of type A_t for some $t \in \{1, 2, 3\}$ in M then C is non-contractible.

Proof. Let C be a cycle of type A_t for a fixed $t \in \{1, 2, 3\}$ in M . We claim that the cycle C is non-contractible. Let the cycle C be of type A_1 . Suppose, C is contractible. Let D_C be a 2-disk bounded by the cycle C . Let f_0, f_1 and f_2 denote the number of vertices, edges and faces of D_C respectively. Let there be n internal vertices and m boundary vertices. So, $f_0 = n + m$, $f_1 = (5n + 3m)/2$ and $f_2 = n + (n + m)/2$ if quadrangles are incident with C , and $f_0 = n + m$, $f_1 = (5n + 4m)/2$ and $f_2 = 3n/2 + m$ if triangles are incident with C in D_C . In both the cases, $f_0 - f_1 + f_2 = 0$. This is not possible since the Euler characteristic of the 2-disk D_C is 1. Therefore, C is non-contractible.

We argue with the similar argument for the cycles of types A_2 and A_3 . Suppose a cycle W of type A_2 is contractible. Let D_W be a 2-disk which is bounded by the cycle W . Similarly as above, calculate f_0, f_1, f_2 and we get $f_0 - f_1 + f_2 = 0$, a contradiction.

We can argue similar way for cycle of type A_3 . This completes the proof. \square

Let C be a cycle of type A_t for a fixed $t \in \{1, 2, 3\}$ in M . Let S be a set of faces which are incident at u for all $u \in V(C)$. The geometric carrier $|S|$ is a cylinder since C is non-contractible. Let $S_C := |S|$. Observe that a cylinder (or an annulus) in M is a sub-complex of M with two boundary cycles. If the boundary cycles of a cylinder are same, it is a torus and we say that the cylinder has identical boundary components. Clearly, the S_C is a cylinder and has two boundary cycles. Let $\partial S_C = \{C_1, C_2\}$. Then, $\text{length}(C) = \text{length}(C_1) = \text{length}(C_2)$ by Lemma 4.4.

Lemma 4.4. *If C is a cycle of type A_i for a fixed $i \in \{1, 2, 3\}$ such that S_C is a cylinder and $\partial S_C = \{C_1, C_2\}$ then $\text{length}(C) = \text{length}(C_1) = \text{length}(C_2)$.*

Proof. Let C be a cycle of type A_1 . Let F_1, F_2, \dots, F_r be a sequence of faces in order which are incident with C and lie on one side of C . These faces F_1, F_2, \dots, F_r are also incident with C_t and lie on one side of C_t for a fixed $t \in \{1, 2\}$. Without loss of generality we assume that $C_t = C_1$. Let $\widehat{F}_1, \widehat{F}_2, \dots, \widehat{F}_r$ denote the sequence of faces incident with C that lie on the other side of C . Let \widehat{F}_i be a \widehat{d}_i -gon. Then, we get $T_1 := \{\widehat{d}_1, \widehat{d}_2, \dots, \widehat{d}_r\}$ the sequence of face-types corresponding to the sequence $\widehat{F}_1, \widehat{F}_2, \dots, \widehat{F}_r$. Again, let W_1, W_2, \dots, W_r denote the sequence of faces incident with C_1 that lie on the other side of C_1 . Similarly, let $T_2 := \{d_1, d_2, \dots, d_r\}$ for d_i -gon W_i , $i = 1, 2, \dots, r$. Since F_1, F_2, \dots, F_r is a sequence of faces that lie on one side of both C and C_1 , so, there exists a j such that $\widehat{d}_1 = d_j, \widehat{d}_2 = d_{j+1}, \dots, \widehat{d}_{k-j+1} = d_k, \widehat{d}_{k-j+2} = d_1, \dots, \widehat{d}_k = d_{j-1}$. So, the cycle C_1 is of type A_1 . Similarly we argue as above for C_2 and we get that the cycle C_2 is of type A_1 . For example in Figure 15, cycle $C = C(x_1, \dots, x_r)$, $\partial S_C = \{C_1(w_1, \dots, w_r), C_2(z_1, \dots, z_r)\}$, and C, C_1 and C_2 are cycles of same type A_1 .

Similarly we repeat above argument for the other two types A_j for $j \in \{2, 3\}$ and we get the similar results. So, the boundary cycles of S_C for a cycle C of type A_i , $i \in \{1, 2, 3\}$ are also of type A_i .

Suppose $\text{length}(C) \neq \text{length}(C_1) \neq \text{length}(C_2)$. Let $C := C(u_1, u_2, \dots, u_r)$, $C_1 := C(v_1, v_2, \dots, v_s)$ and $C_2 := C(w_1, w_2, \dots, w_l)$ denote three cycles of type A_1 . The link $lk(u_1)$ contains the vertices v_1 and w_1 . Let $P(v_1, u_1, w_1)$ be a path of type either A_2 or A_3 through u_1 . Without loss of generality, we assume $r < s$ and $r \neq l$ since $r \neq s \neq l$. Now, the path $P(v_1, u_1, w_1)$ is a shortest path between v_1 and w_1 via u_1 . So it follows that the path $P(v_i, u_i, w_i)$ is also a shortest path of type either A_2 or A_3 between v_i and w_i via u_i . Since, by assumption $r < s$, so, the link $lk(u_r)$ contains the vertices v_r, v_{r+j} for $j(> 0)$ and w_r . This gives that the link of u_r is different from the link of u_{r-1} . This is a contradiction as M is a semi-equivelar map. Therefore, $r = s = l$, that is, $\text{length}(C) = \text{length}(C_1) = \text{length}(C_2)$.

Similarly we argue for the cycles of type A_j for $j \in \{2, 3\}$. This completes the proof. \square

Let C_1 and C_2 be two cycles of same type in a semi-equivelar map M on the torus. We denote S_{C_1, C_2} a cylinder if the boundary components are C_1 and C_2 . We say that the cycle C_1 is *homologous* to C_2 if S_{C_1, C_2} exists. In particular, if $C_1 = C_2$ then consider $S_{C_1, C_2} = C_1 (= C_2)$, and hence, C_1 is homologous to C_2 .

So by above lemma, the cycle C and the boundary cycles of S_C are homologous. Let C_1, C_2, \dots, C_m be a list of cycles which are homologous to C in M . Then, the cycles have same length. That is,

Lemma 4.5. *If C_1, C_2, \dots, C_m are homologous cycles of type A_t for a fixed $t \in \{1, 2, 3\}$ then $\text{length}(C_i) = \text{length}(C_j)$ for $1 \leq i, j \leq m$.*

Proof. Let C_i be a cycle of type A_1 . Then, we have a cylinder S_{C_i} . Let $C_{t_1}, C_{t_2}, \dots, C_{t_m}$ denote a sequence of cycles $\{C_1, C_2, \dots, C_m\}$ such that $\partial S_{C_{t_j}} = \{C_{t_{j-1}}, C_{t_{j+1}}\}$ for $2 \leq j \leq (m-1)$. So, by Lemma 4.4, $\text{length}(C_{t_1}) = \text{length}(C_{t_j})$ for $j \in \{1, 2, \dots, m\}$. Thus, $\text{length}(C_i) = \text{length}(C_j)$ for $1 \leq i, j \leq m$. Similarly we argue for the cycles of type A_j for $j \in \{2, 3\}$. This completes the proof. \square

A (r, s, k)-representation in M : Let $v \in V(M)$. By Lemma 4.1, there are three cycles of types A_1, A_2, A_2 through v . Let L_1, L_2, L_3 be three cycles through the vertex v

where the cycle L_i is of type A_i , $i = 1, 2, 3$. Let $L_1 := C(a_1, a_2, \dots, a_r)$. We cut map M along the cycle L_1 . We get a cylinder which is bounded by identical cycle L_1 . We denote it by N_1 . We call that a cycle is a *horizontal cycle* if the cycle is L_1 or homologous to L_1 . Similarly, we say that a cycle is a *vertical cycle* if the cycle is L_i or homologous to L_i for $i \in \{2, 3\}$. Observe that the horizontal and vertical cycles are non-contractible by Lemma 4.3. Again, we say that a path is a *vertical path* if the path is part of a vertical cycle. We consider N_1 and make another cut in N_1 starting through the vertex v along a path $Q \subset L_3$ until reaching L_1 again for the first time where the starting adjacent face to the horizontal cycle L_1 . (For example in Figure 15, $v = v_1$.) Assume that along $P := P(w_1(= a_1), w_2, \dots, w_m) \subset L_3$ we took the second cut in N_1 . Thus, we get a planar representation which is denoted by N_2 .

Claim 1. The representation N_2 is connected.

Observe that the N_1 is connected as L_1 is a non-contractible cycle. Suppose N_2 is disconnected. This implies that there exists a 2-disk namely $D_{P_1 \cup Q_1}$ which is bounded by a cycle $P_1 \cup Q_1 = P(u_j, \dots, u_i) \cup P(a_t, \dots, a_s)$ where $P_1 \subset L_3$, $Q_1 \subset L_1$, $u_i = a_t$ and $u_j = a_s$. We consider faces which are incident with P_1 and Q_1 in $D_{P_1 \cup Q_1}$. (In this article, \square represents quadrangular face and \triangle represents triangular face.) Observe that if the quadrangular faces are incident with Q_1 and $\square_i, \triangle_{i,1}, \triangle_{i,2}, \square_{i+1}, \triangle_{i+1,1}, \triangle_{i+1,2}, \dots, \triangle_{j-1,1}, \triangle_{j-1,2}, \square_j$ are incident with P_1 in $D_{P_1 \cup Q_1}$ then as in Lemma 4.3, we calculate number of vertices f_0 , edges f_1 and faces f_2 of $D_{P_1 \cup Q_1}$. So, we get $f_0 - f_1 + f_2 = 0$. Similarly, if the triangular faces are incident with Q_1 then also we calculate f_0, f_1 and f_2 of $D_{P_1 \cup Q_1}$. Similarly, we get $f_0 - f_1 + f_2 = 0$. Which is a contradiction in both the case as $D_{P_1 \cup Q_1}$ is 2-disk. So, N_2 is connected.

Observe that N_2 is planer and bounded by L_1 and Q . Let s denote the number of cycles which are homologous to L_1 along P in N_2 . (For example in Figure 15, we took second cut along the path $P(v_1, w_1, x_1, z_1, v_{k+1})$ which is part of L_3 and $s = 4$.) Now, in N_2 , $\text{length}(L_1) = r$ and number of horizontal cycles along P is s . So, we denote N_2 by (r, s) -representation.

Observe that N_2 is bounded by cycle L_1 , path Q , cycle L'_1 and path Q' where $L_1 = L'_1$ and $Q = Q'$. We say that the cycle L_1 is a *lower (base) horizontal cycle* and L'_1 is an *upper horizontal cycle* in (r, s) -representation. Without loss, we may assume that the incident faces of L_1 are quadrangles. (For example in Figure 15, $C(v_1, \dots, v_r)$ is a lower horizontal cycle and $C(v_{k+1}, \dots, v_k)$ is an upper horizontal cycle.) So, we get identification of vertical sides of N_2 in the natural manner but the identification of the horizontal sides needs some shifting so that a vertex in the lower(base) side is identified with a vertex in the upper side. Let $L'_1 = C(a_{k+1}(= w_m), \dots, a_k)$. Then, a_{k+1} is the starting vertex in L'_1 . In (r, s) -representation, let $k =: \text{length}(P(a_1, \dots, a_{k+1}))$ if $P(a_1, \dots, a_{k+1})$ is part of L_1 . Thus, we get a new (r, s, k) -representation of the (r, s) -representation. We call boundaries of (r, s, k) -representation are the cycles and paths along which we took the cuts to construct (r, s, k) -representation of M . (For example in Figure 15, the vertex v_{k+1} is the starting vertex of the upper horizontal cycle $C(v_{k+1}, v_{k+2}, \dots, v_k)$ and $k = \text{length}(P(v_1, v_2, \dots, v_{k+1}))$.) \square

By the above construction, we see that every map of type $\{3^3, 4^2\}$ has a (r, s, k) -representation. We use $T(r, s, k)$ to represent (r, s, k) -representation. Therefore, we have the following lemma.

Lemma 4.6. *The map of type $\{3^3, 4^2\}$ on the torus has a $T(r, s, k)$ representation.*

Let $T(r, s, k)$ be a representation of M . It has two identical upper and lower horizon-

tal cycles of type A_1 , namely, $C_{lh} := C(u_1, u_2, \dots, u_r)$ and $C_{uh} := C(u_{k+1}, u_{k+2}, \dots, u_k)$ in $T(r, s, k)$ respectively. (For example in Figure 11, $C_{lh} = C(v_1, v_2, \dots, v_7)$ and $C_{uh} = C(v_4, v_5, \dots, v_3)$.) We define a new cycle in $T(r, s, k)$ using C_{lh} and C_{uh} . The vertex $u_{k+1} \in V(C_{lh})$ is the starting vertex of C_{uh} in $T(r, s, k)$. In $T(r, s, k)$, we define two paths $Q_2 = P(u_{k+1}, \dots, u_{k_1})$ of type A_2 and $Q_3 = P(u_{k+1}, \dots, u_{k_2})$ of type A_3 through u_{k+1} in $T(r, s, k)$ where $u_{k_1}, u_{k_2} \in V(C_{uh})$. Clearly, the paths Q_2 and Q_3 are not homologous to horizontal cycles, that is, these are not part of cycles of type A_1 . We define two edge disjoint paths $Q'_2 = P(u_{k_1}, \dots, u_{k+1})$ and $Q'_3 = P(u_{k_2}, \dots, u_{k+1})$ in C_{uh} where $Q'_2 \cup Q'_3 := P(u_{k_1}, \dots, u_{k+1}, \dots, u_{k_2}) \subset C_{uh}$ is a path in $T(r, s, k)$. Let $C_{4,1} := Q'_2 \cup Q_2 := C(u_{k+1}, \dots, u_{k_1}, \dots, u_{k+1})$ and $C_{4,2} := Q'_3 \cup Q_3 := C(u_{k+1}, \dots, u_{k_2}, \dots, u_{k+1})$. So, we define a new cycle C_4 using lengths of $C_{4,1}$ and $C_{4,2}$ as follows:

$$C_4 := \begin{cases} C_{4,1}, & \text{if } \text{length}(C_{4,1}) \leq \text{length}(C_{4,2}), \\ C_{4,2}, & \text{if } \text{length}(C_{4,1}) > \text{length}(C_{4,2}). \end{cases} \quad (1)$$

It follows from the definition of C_4 that $\text{length}(C_4) := \min\{\text{length}(Q'_2) + \text{length}(Q_2), \text{length}(Q'_3) + \text{length}(Q_3)\} = \min\{k + s, (r - \frac{s}{2} - k)(\text{mod } r) + s\}$. We say that the cycle C_4 is of type A_4 in $T(r, s, k)$. (In this section, we use $(r + \frac{s}{2} - k)$ in place of $(r + \frac{s}{2} - k)(\text{mod } r)$.) (For example in Figure 11, $C_{4,1} = Q'_2 \cup Q_2 = P(v_4, v_3, v_2) \cup P(v_2, z_5, x_5, w_4, v_4)$ and $C_{4,2} = Q'_3 \cup Q_3 = P(v_4, v_5, v_6, v_7) \cup P(v_7, z_4, x_4, w_4, v_4)$.) So, we have the cycles of four types A_1, A_2, A_3 and A_4 in $T(r, s, k)$.

We show that the cycles of type A_1 have same length and the cycles of type A_2 have at most two different lengths in M . So,

Lemma 4.7. *In M , the cycles of type A_1 have unique length and the cycles of type A_2 have at most two different lengths.*

Proof. Let C_1 be a cycle of type A_1 in M . By the preceding argument of this section, the geometric carrier S_{C_1} of the faces which are incident with C_1 is a cylinder and $\partial S_{C_1} := \{C_2, C_0\}$ where the cycle C_2 is homologous to C_1 and $\text{length}(C_1) = \text{length}(C_2)$. Similarly as above, the S_{C_2} is a cylinder and which is bounded by two homologous cycles C_1 and another, say C_3 of type A_1 . Again, we consider the cycle C_3 and continue with above process. In this process, let C_i denote a cycle at i^{th} step such that $\partial S_{C_i} = \{C_{i-1}, C_{i+1}\}$ and $\text{length}(C_{i-1}) = \text{length}(C_i) = \text{length}(C_{i+1})$. Since M consists of finite number of vertices, it follows that this process stops after, say $t + 1$ number of steps when the cycle C_0 appears in this process. Thus, we get C_1, C_2, \dots, C_t cycles of type A_1 which are homologous to C_1 and cover all the vertices of M as the vertices of S_{C_i} are the vertices of $C_{i-1} \cup C_i \cup C_{i+1}$ for $1 \leq i \leq t$. It is clear from the definition that there is only one cycle of type A_1 through each vertex in M . Therefore, the cycles C_1, C_2, \dots, C_t are the only cycles of type A_1 in M . Since these cycles are homologous to each other, it follows that $\text{length}(C_1) = \text{length}(C_i)$ for $i \in \{1, \dots, t\}$ by Lemma 4.5. This implies that the cycles of type A_1 have unique length in M .

Let L_1, L_2, L_3 be three cycles through a vertex of types A_1, A_2, A_3 respectively in M . We repeat above process and consider L_2 in place of C_1 . Similarly, here we get a sequence of cycles, namely, $R_1 (= L_2), R_2, \dots, R_k$ of type A_2 which are homologous to each other. Since R_i and R_j are homologous to each other for $1 \leq i, j \leq k$, it follows that $l_1 = \text{length}(L_2) = \text{length}(R_i)$ for $1 \leq i \leq k$ by Lemma 4.5. Similarly, again we consider the cycle L_3 and repeat above argument. Let $l_2 = \text{length}(L_3)$. Since the cycles L_2 and L_3 are mirror image of each other, it follows that they define same type of cycles. So, the map M contains the

cycles of type A_2 of lengths l_1 and l_2 . Therefore, the cycles of type A_2 have at most two different lengths in M . This completes the proof. \square

We define admissible relations among r, s, k of $T(r, s, k)$ such that $T(r, s, k)$ represents a map after identifying their boundaries.

Lemma 4.8. *The maps of type $\{3^3, 4^2\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 2$ even, (ii) $rs \geq 10$, (iii) $2 \leq k \leq r - 3$ if $s = 2$ & $0 \leq k \leq r - 1$ if $s \geq 4$.*

Proof. Let $T(r, s, k)$ be a representation of M . In $T(r, s, k)$, the s denote the number of horizontal cycles of type A_1 . By the preceding argument of this section and Lemma 4.7, the cycles of type A_1 are homologous to each other, cover all the vertex of M and have length r . So, the number of vertex n of $M =$ length of the cycle of type $A_1 \times$ the number of cycles of type $A_1 = rs$.

Let C be a cycle of type A_1 . By the definition of A_1 , the triangles incident with C lie on one side and 4-gons lie on the other side of C . If $s = 1$ then $T(r, 1, k)$ contains one horizontal cycle namely C . Since the incident faces of C are either triangles or 4-gons, it implies that the faces of M are either only 3-gons or 4-gons. This is a contradiction as M consists of both types of faces. So, $s \geq 2$ for all r . If s is not an even integer and C is the base horizontal cycle in $T(r, s, k)$ then the incident faces of a vertex in C are all 3-gons or 4-gons after identification of the boundaries of $T(r, s, k)$. This is a contradiction as both 3- and 4-gons are incident at each vertex of M . So, s is even.

If $s = 2$ and $r < 5$ then the representation $T(4, 2, k)$ has two horizontal cycles. If $C(u_1, u_2, u_3, u_4)$ and $C(u_5, u_6, u_7, u_8)$ are two horizontal cycles in $T(4, 2, k)$ then the link $lk(u_6)$ is not a cycle. So, $r \neq 4$. Similarly one can see that $r \neq 1, 2, 3$. Thus, $r \geq 5$. If $s \geq 4$ and $r < 3$ then one can see as above that some vertex has link which is not a cycle. So, by combining above all cases, $r \geq 3$ and $rs \geq 10$.

If $s = 2$ and $k \in \{1, \dots, r - 1\} \setminus \{2, \dots, r - 3\}$ then we proceed with as above and we get some vertex whose link is not a cycle. Thus, $s = 2$ implies $k \in \{2, \dots, r - 3\}$. Similarly we repeat the above argument for $s \geq 4$ and we get that $k \in \{1, \dots, r - 1\}$ if $s \geq 4$. This completes the proof. \square

Let M_1 and M_2 be two maps of type $\{3^3, 4^2\}$ with same number of vertices on the torus and $T_i := T(r_i, s_i, k_i)$, $i \in \{1, 2\}$ denote M_i . If $a_{i,1}$ = length of the cycle of type A_1 , $a_{i,2}$ = length of the cycle of type A_2 , $a_{i,3}$ = length of the cycle of type A_3 and $a_{i,4}$ = length of the cycle of type A_4 in T_i then we say that $T(r_i, s_i, k_i)$ has cycle-type $(a_{i,1}, a_{i,2}, a_{i,3}, a_{i,4})$ if $a_{i,2} \leq a_{i,3}$ or $(a_{i,1}, a_{i,3}, a_{i,2}, a_{i,4})$ if $a_{i,3} < a_{i,2}$. Now, we show the following isomorphism lemma.

Lemma 4.9. *The map $M_1 \cong M_2$ if and only if they have same cycle-type.*

Proof. We first assume that the maps M_1 and M_2 have same cycle-type. This gives that $a_{1,1} = a_{2,1}$, $\{a_{1,2}, a_{1,3}\} = \{a_{2,2}, a_{2,3}\}$ and $a_{1,4} = a_{2,4}$. The maps M_i for $i \in \{1, 2\}$ have a $T_i = T(r_i, s_i, k_i)$ representation.

Claim. $T_1 \cong T_2$.

The T_1 has s_1 number of horizontal cycles of type A_1 , namely, $C(1, 0) := C(u_{0,0}, u_{0,1}, \dots, u_{0,r_1-1})$, $C(1, 1) := C(u_{1,0}, u_{1,1}, \dots, u_{1,r_1-1})$, \dots , $C(1, s_1 - 1) := C(u_{s_1-1,0}, u_{s_1-1,1}, \dots, u_{s_1-1,r_1-1})$ in order. Similarly, the T_2 has s_2 number of horizontal cycles of type A_1 , namely, $C(2, 0) := C(v_{0,0}, v_{0,1}, \dots, v_{0,r_2-1})$, $C(2, 1) := C(v_{1,0}, v_{1,1}, \dots, v_{1,r_2-1}, v_{1,0})$, \dots , $C(2, s_2 - 1) := C(v_{s_2-1,0}, v_{s_2-1,1}, \dots, v_{s_2-1,r_2-1})$ in order. Now we have the following cases.

Case 1 : If $(r_1, s_1, k_1) = (r_2, s_2, k_2)$ then $r_1 = r_2, s_1 = s_2, k_1 = k_2$. We define a map $f_1 : V(T(r_1, s_1, k_1)) \rightarrow V(T(r_2, s_2, k_2))$ such that $f_1(u_{t,i}) = v_{t,i}$ for $0 \leq t \leq s-1$ and $0 \leq i \leq r-1$. Observe that $lk(u_{t,i}) = C(\mathbf{u}_{t-1,i-1}, u_{t-1,i}, \mathbf{u}_{t-1,i+1}, u_{t,i+1}, u_{t+1,i+1}, u_{t+1,i}, u_{t,i-1})$ is the link of the vertex $u_{t,i}$ in $T(r_i, s_i, k_i)$. By f_1 , $f_1(lk(u_{t,i})) = C(f_1(\mathbf{u}_{t-1,i-1}), f_1(u_{t-1,i}), f_1(\mathbf{u}_{t-1,i+1}), f_1(u_{t,i+1}), f_1(u_{t+1,i+1}), f_1(u_{t+1,i}), f_1(u_{t,i-1})) = C(\mathbf{v}_{t-1,i-1}, v_{t-1,i}, \mathbf{v}_{t-1,i+1}, v_{t,i+1}, v_{t+1,i+1}, v_{t+1,i}, v_{t,i-1})$. So, $f_1(lk(u_{t,i})) = lk(v_{t,i})$. This implies that the map f_1 sends vertices to vertices, edges to edges, faces to faces and also, preserves incidents. Therefore, the map f_1 defines an isomorphism map between $T(r_1, s_1, k_1)$ and $T(r_2, s_2, k_2)$. Thus, $T_1 \cong T_2$ by f_1 .

Case 2 : If $r_1 \neq r_2$ then $\text{length}(C_{1,1}) \neq \text{length}(C_{2,1})$. This implies that $a_{1,1} \neq a_{2,1}$, a contradiction since $a_{1,1} = a_{2,1}$. So, $r_1 = r_2$.

Case 3 : If $s_1 \neq s_2$ then $n_1 = r_1 s_1 \neq r_1 s_2 = n_2$ as $r_1 = r_2$ by Case 2. This is a contradiction since $n_1 = n_2$. So, $s_1 = s_2$.

Case 4 : Suppose $k_1 \neq k_2$. By assumption, $a_{1,4} = a_{2,4}$, $\text{length}(C_{1,4}) = \text{length}(C_{2,4})$. This implies that $\min\{k_1 + s_1, r_1 + \frac{s_1}{2} - k_1\} = \min\{k_2 + s_2, r_2 + \frac{s_2}{2} - k_2\}$. It follows that $k_1 + s_1 \neq k_2 + s_2$ since $k_1 \neq k_2$ and $s_1 = s_2$. This gives that $k_1 + s_1 = r_2 + \frac{s_2}{2} - k_2 = r_1 + \frac{s_1}{2} - k_2$ as $r_1 = r_2$ and $s_1 = s_2$. That is, $k_2 = r_1 - k_1 + \frac{s_1}{2} - s_1 = r_1 - k_1 - \frac{s_1}{2}$. In this case, identify T_2 along vertical identical boundary $P(v_{0,0}, v_{1,0}, \dots, v_{s_2-1,0}, v_{0,k_1})$ of T_2 and then cut along the path $Q = P(v_{0,0}, v_{1,0}, v_{2,1}, \dots, v_{s_2-1, \frac{s_2}{2}-1}, v_{0, \frac{s_2}{2}+k_2})$ of type A_2 through vertex $v_{0,0}$. We get a new (r, s, k) -representation of M_2 and we denote it by R . This process defines the map $f_2 : V(T(r_2, s_2, k_2)) \rightarrow V(R)$ such that $f_2(v_{t,i}) = v_{t, (r_2-i+\lfloor \frac{i}{2} \rfloor) \pmod{r_2}}$ for $0 \leq t \leq s_2-1$ and $0 \leq i \leq r_2-1$. In R , the base horizontal cycle is $C'(2,0) := C(v_{0,0}, v_{0,r_2-1}, \dots, v_{0,1})$, upper horizontal cycle is $C(v_{0, k_2+\frac{s_2}{2}}, v_{0, k_2+\frac{s_2}{2}-1}, \dots, v_{0, k_2+\frac{s_2}{2}+1})$ and the length of the path $P(v_{0,0}, v_{0, r_2-1}, \dots, v_{0, k_2+\frac{s_2}{2}})$ in $C'(2,0)$ is $r_2 - \frac{s_2}{2} - k_2$. In this process, we are not changing the both length of the horizontal cycles and number of horizontal cycles which are homologous to the cycle $C'(2,0)$. So, we get $R = T(r_2, s_2, r_2 - k_2 - \frac{s_2}{2})$. Now $r_2 - k_2 - \frac{s_2}{2} = r_2 - (r_1 - k_1 - \frac{s_1}{2}) - \frac{s_2}{2} = k_1$ since $r_1 = r_2, s_1 = s_2$ and $k_2 = r_1 - k_1 - \frac{s_1}{2}$. Thus, by f_1 , $T(r_2, s_2, r_2 - k_2 - \frac{s_2}{2}) \cong T(r_1, s_1, k_1)$. So, $T_1 \cong T_2$. So, by Cases 1, 2, 3, 4, claim follows.

Therefore, by f_1 , $M_1 \cong M_2$.

Conversely, let $M_1 \cong M_2$. Then, there is an isomorphism map $f : V(M_1) \rightarrow V(M_2)$. Let $C_{1,j}$ be cycle of type A_j for $j = 1, 2, 3, 4$ in M_1 . By f , consider $C_{2,j} := f(C_{1,j})$ for $j = 1, 2, 3, 4$. So, $\text{length}(C_{1,j}) = \text{length}(f(C_{1,j})) = \text{length}(C_{2,j})$ for $j = 1, 2, 3, 4$ since f is an isomorphism map. Hence, M_1 and M_2 have same cycle-type. \square

Thus, we state the following corollary.

Corollary 4.1. $T(r_1, s_1, k_1) \not\cong T(r_2, s_2, k_2) \forall r_1 \neq r_2, T(r_1, s_1, k_1) \not\cong T(r_2, s_2, k_2) \forall s_1 \neq s_2, T(r_1, s_1, k_1) \not\cong T(r_1, s_1, k_2)$ if $s_1 = 2$ and $k_2 \in \{2, 3, \dots, r_1 - 3\} \setminus \{k_1, r_1 - k_1 - 1\}$, $T(r_1, s_1, k_1) \not\cong T(r_1, s_1, k_2)$ if $s_1 \geq 4$ and $k_2 \in \{0, 1, \dots, r_1 - 1\} \setminus \{k_1, r_1 - k_1 - \frac{s_1}{2}\}$, $T(r_1, s_1, k_1) \cong T(r_1, s_1, r_1 - k_1 - 1)$ if $s_1 = 2$ and $r_1 \geq 5$, and $T(r_1, s_1, k_1) \cong T(r_1, s_1, r_1 - \frac{s_1}{2} - k_1)$ if $s_1 \geq 4$ and $r_1 \geq 3$.

Proof. If $r_1 \neq r_2$ then it follows that $a_{1,1} \neq a_{2,1}$. This implies that $T(r_1, s_1, k_1) \not\cong T(r_2, s_2, k_2)$ by Lemma 4.9. So, $T(r_1, s_1, k_1) \not\cong T(r_2, s_2, k_2) \forall r_1 \neq r_2$. Again, $s_1 \neq s_2$ implies $r_1 \neq r_2$ since $r_1 s_1 = r_2 s_2$. This implies that $T(r_1, s_1, k_1) \not\cong T(r_2, s_2, k_2) \forall s_1 \neq s_2$. If $k_1 \neq k_2, r_1 = r_2$ and $s_1 = s_2$ then by the argument in the proof of Lemma 4.9, $T(r_1, s_1, k_1) \cong T(r_1, s_1, k_2)$ if and only if $k_2 = r_1 - k_1 - \frac{s_1}{2}$. So, $T(r_1, s_1, k_1) \cong T(r_1, s_1, k_2)$ if $s_1 = 2$ and $k_2 \neq r_1 - k_1 - 1$. Thus, $T(r_1, s_1, k_1) \not\cong T(r_1, s_1, k_2)$ if $s_1 = 2$ and $k_2 \in$

Table 1: Maps of type $\{3^3, 4^2\}$

n	Equivalence classes	Length of cycles	$i(n)$
10	T(5, 2, 2)	(5, {10, 10}, 4)	1(10)
12	T(6, 2, 2), T(6, 2, 3) T(3, 4, 0), T(3, 4, 1) T(3, 4, 2)	(6, {6, 4}, 4) (3, {4, 12}, 4) (3, {12, 12}, 6)	3(12)
14	T(7, 2, 2), T(7, 2, 4) T(7, 2, 3)	(7, {14, 14}, 4) (7, {14, 14}, 5)	2(14)
16	T(8, 2, 2), T(8, 2, 5) T(8, 2, 3), T(8, 2, 4) T(4, 4, 0), T(4, 4, 2) T(4, 4, 1) T(4, 4, 3)	(8, {8, 16}, 4) (8, {16, 4}, 5) (4, {4, 8}, 4) (4, {16, 16}, 5) (4, {16, 16}, 7)	5(16)
18	T(9, 2, 2), T(9, 2, 6) T(9, 2, 3), T(9, 2, 5) T(9, 2, 4) T(3, 6, 0) T(3, 6, 1), T(3, 6, 2)	(9, {18, 6}, 4) (9, {6, 18}, 5) (9, {18, 18}, 6) (3, {6, 6}, 6) (3, {18, 18}, 7)	5(18)
20	T(10, 2, 2), T(10, 2, 7) T(10, 2, 3), T(10, 2, 6) T(10, 2, 4), T(10, 2, 5) T(5, 4, 0), T(5, 4, 3) T(5, 4, 1), T(5, 4, 2) T(5, 4, 4)	(10, {10, 20}, 4) (10, {20, 10}, 5) (10, {10, 4}, 6) (5, {4, 20}, 4) (5, {20, 20}, 5) (5, {20, 20}, 8)	6(20)
22	T(11, 2, 2), T(11, 2, 8) T(11, 2, 3), T(11, 2, 7) T(11, 2, 4), T(11, 2, 6) T(11, 2, 5)	(11, {22, 22}, 4) (11, {22, 22}, 5) (11, {22, 22}, 6) (11, {22, 22}, 7)	4(22)

$\{2, 3, \dots, r_1 - 3\} \setminus \{k_1, r_1 - k_1 - 1\}$, and $T(r_1, 2, k_1) \cong T(r_1, 2, r_1 - k_1 - 1)$ if $r_1 \geq 5$ (by Lemma 4.8). Again, $T(r_1, s_1, k_1) \cong T(r_1, s_1, k_2)$ if $s_1 \geq 4$ and $k_2 \neq r_1 - \frac{s_1}{2} - k_1$. So, $T(r_1, s_1, k_1) \not\cong T(r_1, s_1, k_2)$ if $s_1 \geq 4$ and $k_2 \in \{0, 1, \dots, r_1 - 1\} \setminus \{k_1, r_1 - k_1 - \frac{s_1}{2}\}$, and $T(r_1, s_1, k_1) \cong T(r_1, s_1, r_1 - \frac{s_1}{2} - k_1)$ if $s_1 \geq 4$ and $r_1 \geq 3$ (by Lemma 4.8). This completes the proof. \square

We calculate all possible $T(r, s, k)$ representations on n vertices by Lemma 4.8. Then, we calculate lengths of the cycles of type A_i for $i \in \{1, 2, 3, 4\}$. Next, we classify all $T(r, s, k)$ representation by Lemma 4.9 up to isomorphism. So, by the Lemmas 4.8, 4.9, maps of type $\{3^3, 4^2\}$ can be classified up to isomorphism. We repeat this same argument in the Sections 5, 6, 7, 8, 9, 10, 11. We have done the above calculations for the vertices $n \leq 22$. We have listed the obtained objects in the form of their (r, s, k) -representation in Table 1. In Table 1, we have used n to denote the number of vertices of a map. We put $T(r_1, s_1, k_1)$ and $T(r_2, s_2, k_2)$ in a single equivalence class if $T(r_1, s_1, k_1)$ and $T(r_2, s_2, k_2)$ are isomorphic. We have used $(a_1, \{a_2, a_3\}, a_4)$ to denote a permutation of lengths of cycles where $a_j = \text{length}(C_{1,j})$ for $j \in \{1, 2, 3, 4\}$ and $\{a_2, a_3\}$ denotes a set of lengths of the cycles $C_{1,2}$ and $C_{1,3}$ of type A_2 . We have also used $i(n)$ where i denote the number of non-isomorphic objects of type $\{3^3, 4^2\}$ on n vertices up to isomorphism. The above notations are also used in Tables 2, 3, 4, 5, 6, 7, 8.

5 Maps of type $\{3^2, 4, 3, 4\}$

Let M be a map of type $\{3^2, 4, 3, 4\}$ on the torus. Through each vertex in M there is a path as follows.

Definition 5.1. *Let $P(\dots, u_{i-1}, u_i, u_{i+1}, \dots)$ be a path in edge graph of M . We say the path P of type B_1 if $lk(u_i) = C(\mathbf{a}, u_{i+1}, \mathbf{b}, c, \mathbf{d}, u_{i-1}, e)$ implies $lk(u_{i-1}) = C(\mathbf{f}, g, e, u_i, \mathbf{c}, d, u_{i-2})$ and $lk(u_{i+1}) = C(e, a, k, u_{i+2}, \mathbf{l}, b, u_i)$, and $lk(u_i) = C(e, h, k, u_{i+1}, \mathbf{l}, b, u_{i-1})$ implies $lk(u_{i-1}) = C(\mathbf{h}, u_i, b, c, \mathbf{d}, u_{i-2}, e)$ and $lk(u_{i+1}) = C(\mathbf{s}, u_{i+2}, t, \mathbf{l}, \mathbf{b}, u_i, k)$.*

In Figure 17, $lk(u_i) = C(\mathbf{a}, b, c, u_{i+1}, \mathbf{f}, e, u_{i-1})$ and path $P(u_{i-1}, u_i, u_{i+1})$ is part of a path of type B_1 . Let P be a maximal path of type B_1 . Then, by the next Lemma 5.1, it defines a cycle.




Figure 18 : Cylinder




Figure 17 : $lk(u_i)$




Figure 20




Figure 19 : $T(r, 4, 2k)$

Lemma 5.1. *If P is a maximal path of type B_1 in M then there exists an edge e such that $P \cup e$ is a cycle.*

Proof. Let $P(u_1, u_2, \dots, u_r)$ be a maximal path of type B_1 and $lk(u_r) = C(u_{r-1}, \mathbf{a}, b, c, d, \mathbf{f}, e)$. If $d = u_1$ then $C(u_1, u_2, \dots, u_r)$ is a cycle. Suppose $d \neq u_1$ and $d = u_i$ for some $2 \leq i \leq r$. Then, it defines a cycle $L = C(u_i, u_{i+1}, \dots, u_r)$. By the similar argument as in Lemma 4.1 and by Definition 5.1, either $f = u_{i+1}$, $d = u_i$, $c = u_{i-1}$ or $c = u_{i+1}$, $d = u_i$, $f = u_{i-1}$. In both the cases, by considering faces incident with the cycle, we get a new cycle $C(w_i, w_{i+1}, \dots, w_{r-2})$ (see Figure 20) of same type as L with lesser length. By induction, it is impossible similarly as in Lemma 4.1. Therefore, $d \neq u_i$ for $2 \leq i \leq r$. So, we get a path Q which is extended from P with $\text{length}(P) < \text{length}(Q)$. This is a contradiction as P is maximal. Therefore, $d = u_1$ and the path P defines the cycle $C(u_1, u_2, \dots, u_r)$. So, every maximal path of type B_1 is a cycle. \square

In Figure 19, the path $P(v_1, v_2, \dots, v_r)$ is of type B_1 and the cycle $C(u_1, u_2, \dots, u_r)$ is of type B_1 . Let C_1 and C_2 be two cycles of type B_1 . We claim that

Lemma 5.2. *If C_1 and C_2 are two cycles of type B_1 and $E(C_1) \cap E(C_2) \neq \emptyset$ then $C_1 = C_2$.*

Proof. Let $C_1 := C(u_{1,1}, u_{1,2}, \dots, u_{1,r})$ and $C_2 := C(u_{2,1}, u_{2,2}, \dots, u_{2,s})$ and $E(C_1) \cap E(C_2) \neq \emptyset$. Then, there is an edge $e \in E(C_1 \cap C_2)$. Let $e = yx$. The cycles C_1, C_2 are both well defined at the vertices y and x . Let $lk(x) = C(\mathbf{a}, b, c, w, \mathbf{d}, e, y)$. By Definition 5.1, $w \in V(C_1 \cap C_2)$. So, the path $P(y, x, w)$ is part of both C_1 and C_2 . This implies that $y = u_{1,t_1-1} = u_{2,t_2-1}$, $x = u_{1,t_1} = u_{2,t_2}$ and $w = u_{1,t_1+1} = u_{2,t_2+1}$ for some $t_1 \in \{1, \dots, r\}$ and $t_2 \in \{1, \dots, s\}$. Again, we argue similarly for the edge xw as we did for the edge e . We continue with above process. This process stops after, say r number of steps. Let $t_2 > t_1$ and $t_2 - t_1 = m$ for some m . Then, by this process, we get $u_{1,1} = u_{2,m+1}$, $u_{1,2} = u_{2,m+2}, \dots, u_{1,r} = u_{2,m+r}$ and $u_{1,1} = u_{2,m+r+1}$. This implies that $m + 1 = m + r + 1$ and $r = s$ as $u_{1,m+r+1} = u_{2,m+1}$ and C_2 is a cycle. Hence, $C_1 = C_2$. Again, let $lk(x) = C(\mathbf{a}, b, c, w, \mathbf{d}, y, e)$. Then, by Definition 5.1, $b \in V(C_1 \cap C_2)$. Similarly

again we repeat above argument and we get $C_1 = C_2$. Therefore, by combining above two cases, $E(C_1) \cap E(C_2) \neq \emptyset$ implies $C_1 = C_2$. This completes the proof. \square

Let C be a cycle of type B_1 . Similarly we argue as in Lemma 4.3 for the cycles of type B_1 and so, the cycle C is non-contractible. Let $S := \{F \in F(M) \mid V(C) \cap V(F) \neq \emptyset\}$. The cylinder $S_C = |S|$ has two boundary cycles which are either disjoint or identical by Lemma 5.3.

Lemma 5.3. *Let C be a cycle of type B_1 , $\partial S_C = \{C_1, C_2\}$. If $C_1 \cap C_2 \neq \emptyset$ then $C_1 = C_2$.*

Proof. The cycle C is of type B_1 and $\partial S_C = \{C_1, C_2\}$. We argue similarly as in Lemma 4.1 for the cycle C and the cylinder S_C . So, the cycles C , C_1 and C_2 are of same type B_1 . Let $C_1 \cap C_2 \neq \emptyset$ and $u \in V(C_1 \cap C_2)$. Suppose $C_1 \cap C_2$ does not contain any edge which is incident at u . By Definition 5.1, the number of incident edges that lie on one side of the cycle C_i is two and on the other side is one at each vertex of C_i . Let d_i denote the number of incident edges which are incident at u and does not belong to $E(C_i)$. Hence, $d_1 + d_2 = 3$. The vertex $u \in V(C_1)$ and $u \in V(C_2)$. Since $C_1 \cap C_2$ does not contain any edge at u , so, the cycles C_1 and C_2 both contain two different edges which are incident at the vertex u . This implies that $\text{degree}(u) \geq (d_1 + d_2 + 4) = 7$. This is a contradiction as the degree of the vertex u is five. Therefore, $C_1 \cap C_2$ contains an edge at the vertex u . This implies that $C_1 = C_2$ by Lemma 5.2. Again, if $C_1 \cap C_2$ contains an edge then by Lemma 5.2, $C_1 = C_2$. Therefore, boundary cycles of a cylinder are either identical or disjoint. \square

We show that the cycles of type B_1 have at most two different lengths in the next Lemma 5.4.

Lemma 5.4. *In M , the cycles of type B_1 have at most two different lengths.*

Proof. We proceed with as in the case of Lemma 5.3. There are two cycles of type B_1 through each vertex of M (by the definition of cycle of type B_1). Let $u \in V(M)$. Let C_1 and C'_1 denote two cycles through a vertex u . Consider cylinder S_{C_1} which is defined by cycle C_1 . Let $\partial S_{C_1} = \{C_2, C_0\}$. The cycles C_1 , C_2 , and C_0 are homologous to each other and $\text{length}(C_1) = \text{length}(C_2) = \text{length}(C_0)$ by the similar argument of Lemma 4.5. Again, we proceed with above argument for the cycle C_2 in place of C_1 and continue. In this process, let C_i denote a cycle at i^{th} step where $\partial S_{C_i} = \{C_{i+1}, C_{i-1}\}$ and $\text{length}(C_{i-1}) = \text{length}(C_i) = \text{length}(C_{i+1})$. Let after $k + 1$ number of steps the process stops when the cycle C_1 appears. Thus, the cycles C_i , C_j are homologous for every $1 \leq i, j \leq k$ where $\cup_{i=1}^k V(C_i) = V(M)$ and $l_1 = \text{length}(C_1) = \text{length}(C_i)$ for all $1 \leq i \leq k$. Again, we proceed with above process for C'_1 in place of C_1 . Similarly, we get a sequence of homologous cycles, namely, $C'_1, C'_2, \dots, C'_{k_1}$ such that $\cup_{i=1}^{k_1} V(C'_i) = V(M)$ and $l_2 = \text{length}(C'_i)$ for all $1 \leq i \leq k_1$. So, M contains cycles of type B_1 at most two different lengths l_1 and l_2 . This completes the proof of lemma. \square

Similarly as in Section 4, observe that every map of type $\{3^2, 4, 3, 4\}$ on the torus has a $T(r, s, k)$ representation for some r, s, k . We define admissible relations among r, s, k of $T(r, s, k)$ such that $T(r, s, k)$ represent a map after identifying their boundaries in next lemma. We omit the proof of next lemma as its argument similarly as in Lemma 4.8.

Lemma 5.5. *The maps of type $\{3^2, 4, 3, 4\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 2$ even, (ii) $2 \mid r$, (iii) $rs \geq 16$, (iv) $k \in \{2t + 4 : 0 \leq t \leq \frac{r-8}{2}\}$ if $s = 2$ & $k \in \{2t : 0 \leq t < \frac{r}{2}\}$ if $s \geq 4$.*

Let $T_i = T(r_i, s_i, k_i)$, $i = \{1, 2\}$ denote M_i of type $\{3^2, 4, 3, 4\}$ on the torus with n_i vertices and $n_1 = n_2$. Let $C_{i,1}$ and $C_{i,2}$ be two non-homologous cycles of type B_1 in M_i for $i = 1, 2$ and $a_{i,j} = \text{length}(C_{i,j})$. Then,

Lemma 5.6. *The map $M_1 \cong M_2$ if and only if $(a_{1,1}, a_{1,2}) = (a_{2,t_1}, a_{2,t_2})$ for $t_1 \neq t_2 \in \{1, 2\}$.*

Proof. We first assume that $(a_{1,1}, a_{1,2}) = (a_{2,t_1}, a_{2,t_2})$ where $t_1, t_2 \in \{1, 2\}$ and $t_1 \neq t_2$. This implies that $\{a_{1,1}, a_{1,2}\} = \{a_{2,1}, a_{2,2}\}$.

Claim. $T_1 \cong T_2$.

From the definition of (r_i, s_i, k_i) -representation, the T_i has s_i number of horizontal cycles of type B_1 , namely, $C(1, 0) := C(u_{0,0}, u_{0,1}, \dots, u_{0,r_1-1})$, $C(1, 1) := C(u_{1,0}, u_{1,1}, \dots, u_{1,r_1-1})$, \dots , $C(1, s_1 - 1) := C(u_{s_1-1,0}, u_{s_1-1,1}, \dots, u_{s_1-1,r_1-1})$ in T_1 and $C(2, 0) := C(v_{0,0}, v_{0,1}, \dots, v_{0,r_2-1})$, $C(2, 1) := C(v_{1,0}, v_{1,1}, \dots, v_{1,r_2-1}, v_{1,0})$, \dots , $C(2, s_2 - 1) := C(v_{s_2-1,0}, v_{s_2-1,1}, \dots, v_{s_2-1,r_2-1})$ in T_2 .

Case 1 : If $(r_1, s_1, k_1) = (r_2, s_2, k_2)$ i.e. $r_1 = r_2, s_1 = s_2, k_1 = k_2$ then similarly as in the proof of the Lemma 4.9, we define an isomorphism map $f_1 : V(T(r_1, s_1, k_1)) \rightarrow V(T(r_2, s_2, k_2))$ such that $f(u_{t,i}) = v_{t,i}$ for $0 \leq t \leq s_1 - 1$ and $0 \leq i \leq r_1 - 1$. So, $T_1 \cong T_2$ by f_1 .

Case 2 : Suppose $r_1 = r_2, s_1 = s_2, k_1 \neq k_2$. Since $r_1 = r_2$, it implies that the vertical cycles in T_1 and T_2 have same length.

We define a cycle in T_1 as in equation (1) of Section 4. Let C_{lh} denote the base horizontal cycle and C_{uh} denote the upper horizontal cycle in T_2 . Let Q be a path through u_{k_1+1} of type B_1 and not homologous to C_{lh} . Again, let Q' and Q'' denote two edge disjoint paths in C_{uh} such that $C_{uh} = Q' \cup Q''$. Hence as in equation (1), we define a new cycle $C_3(1)$ using the above paths in T_1 . Similarly, there is a cycle $C_3(2)$ as $C_3(1)$ in T_2 . Since $\{a_{1,1}, a_{1,2}\} = \{a_{2,1}, a_{2,2}\}$ and $r_1 = r_2$, it follows that $\text{length}(C_3(1)) = \text{length}(C_3(2))$.

Since $\text{length}(C_3(1)) = \text{length}(C_3(2))$, it implies that $\min\{s_1 + k_1, r_1 + s_1 - k_1\} = \min\{s_2 + k_2, r_2 + s_2 - k_2\}$. It follows that $r_1 + s_1 - k_1 = s_2 + k_2$ since $k_1 \neq k_2$. So, $k_2 = r_1 - k_1$ as $s_1 = s_2$. We proceed similarly as in the Lemma 4.9. In this process, identify $T(r_2, s_2, k_2)$ along vertical boundary and cut along a path $Q := P(v_{0,i}, v_{1,i}, \dots, v_{s_2-1,i}, v_{0,i+k_2})$ for some odd $0 \leq i \leq r_1 - 1$. Thus, we get a new representation, say R of M_2 with a map $f_2 : V(T(r_2, s_2, k_2)) \rightarrow V(R)$ such that $f_2(v_{t,i'}) = v_{t,(i+r_2-i') \pmod{r_2}}$ for $0 \leq t \leq s_2 - 1$ and $0 \leq i' \leq r_2 - 1$. Clearly, f_2 maps cycle $C(2, t) := C(v_{t,0}, v_{t,1}, \dots, v_{t,r_2-1})$ to cycle $C'(2, t) := C(v_{t,i}, v_{t,i-1}, \dots, v_{t,r_2-1}, v_{t,0}, v_{t,1}, \dots, v_{t,i+1})$. Since the path $Q_1 := P(v_{0,i}, v_{0,i-1}, \dots, v_{0,i+k_2}) \subset C'(2, 0) := C(v_{0,i}, v_{0,i-1}, \dots, v_{0,i+1})$ and $\text{length}(Q_1) = i + r_2 - k_2 - i = r_2 - k_2$, it follows that R has s_2 number of horizontal cycles of length r_2 and the cycle of type B_2 has length $r_2 - k_2$ as $\text{length}(Q_1) = r_2 - k_2$. In R , the faces incident with the base horizontal cycle $C' = C(v_{0,i}, v_{0,i-1}, \dots, v_{0,r_2-1}, v_{0,0}, v_{0,1}, \dots, v_{0,i+1})$ at $v_{0,i-1}$ have two possibilities as follows. If one triangle and one 4-gon incident at $v_{0,i-1}$ in C' then R has a (r, s, k) -representation. If two triangles and one 4-gon incident at $v_{0,i-1}$ then R does not follow the definition of (r, s, k) -representation. (For an example, the Figure 12 is an example of R which does not follow the definition of (r, s, k) -representation since the number of incident triangles at w_2 is two.) In this case, if $C'(2, 0), C'(2, 1), \dots, C'(2, s_2 - 1)$ denote a sequence of horizontal cycles in R then we identify R along $C(v_{0,i}, v_{0,i-1}, \dots, v_{0,i+1})$ and cut along $C(v_{1,i}, v_{1,i-1}, \dots, v_{1,i+1})$. Thus, we get a new representation of M_2 , say R' where $C'(2, 1) := C(v_{1,i}, v_{1,i-1}, \dots, v_{1,i+1})$ denote the base horizontal cycle. In this process, $C'(2, 1) \rightarrow C'(2, 0), C'(2, 0) \rightarrow C'(2, 1), C'(2, s_2 - 1) \rightarrow C'(2, 2), \dots, C'(2, 2) \rightarrow C'(2, s_2 - 1)$. This process defines a map $f_3 : R \rightarrow R'$ such that $f_3(C'(2, t)) = C'(2, 1 - t \pmod{s_2})$ for

Table 2: Maps of type $\{3^2, 4, 3, 4\}$

n	Equivalence classes	Length of cycles	$i(n)$
16	T(8, 2, 4), T(4, 4, 2) T(4, 4, 0)	(8, 4) (4, 4)	2(16)
20	T(10, 2, 4), T(10, 2, 6)	(10, 10)	1(20)
24	T(12, 2, 4), T(12, 2, 8), T(6, 4, 2), T(6, 4, 4) T(12, 2, 6), T(4, 6, 2) T(6, 4, 0), T(4, 6, 0)	(12, 6) (4, 12) (4, 6)	3(24)
28	T(14, 2, 4), T(14, 2, 10) T(14, 2, 6), T(14, 2, 8)	(14, 14)	1(28)
32	T(16, 2, 4), T(16, 2, 12), T(8, 4, 2), T(8, 4, 6) T(16, 2, 6), T(16, 2, 10) T(16, 2, 8), T(4, 8, 2) T(8, 4, 4) T(8, 4, 0), T(4, 8, 0)	(16, 8) (16, 16) (4, 16) (8, 8) (4, 8)	5(32)

$0 \leq t \leq s_2 - 1$. In R' , $C'(2, 1), C'(2, 0), C'(2, s_2 - 1), \dots, C'(2, 2)$ denote the sequence of horizontal cycles in R' . (For an example of R' , the Figure 5 is a $T(6, 4, 2)$ representation which is defined from R in Figure 12. In Figure 12, we cut R along the cycle $C(v_1, v_2, \dots, v_6)$ and identify along $C(w_1, w_2, \dots, w_6)$. Hence, we get a representation $T(6, 4, 2)$ in Figure 5.) In the above process, we are redefining R to a desire representation R' . In this process, the length of the horizontal cycles of type B_1 are remain unchanged as we are only changing the order of the horizontal cycles. So, R' has a well defined $T(r_2, s_2, r_2 - k_2)$ representation. Thus, $T(r_2, s_2, r_2 - k_2) = T(r_1, s_1, k_1)$ since $r_1 = r_2, s_1 = s_2, k_2 = r_1 - k_1$. So, M_2 has a $T(r_1, s_1, k_1)$ representation. Therefore, by $f_1, T_1 \cong T_2$.

Case 3 : If $r_1 \neq r_2$, it implies that $a_{1,1} \neq a_{2,1}$. So by assumption $\{a_{1,1}, a_{1,2}\} = \{a_{2,1}, a_{2,2}\}$, we get that $a_{1,1} = a_{2,2}$. In this case, we identify boundaries of $T(r_2, s_2, k_2)$ and cut along the hole cycle $C(2, 2)$ in place of $C(2, 1)$. Then, take another cut along $C(2, 1)$ until we reaching $C(2, 2)$ again for the first time. Hence, we get $r_1 = \text{length}(C(1, 1)) = \text{length}(C(2, 2)) = r_2$. Thus, $r_1 s_1 = r_2 s_2$ implies that $s_1 = s_2$. Since $r_1 = r_2$ and $s_1 = s_2$, it implies that we are in Case 2. So similarly as in Case 2, we define maps f_1, f_2 and f_3 . Thus, by f_1, f_2 and $f_3, T_1 \cong T_2$.

Case 4 : If $s_1 \neq s_2$ then it implies that $n_1 = r_1 s_1 \neq r_2 s_2 = n_2$ if $r_1 = r_2$. Which is a contradiction as $n_1 = n_2$. If $r_1 \neq r_2$ then we are in Case 3. So, by combining above two Cases 2 and 3, we get an isomorphism map f_1 if $s_1 \neq s_2$. Again, let $k_1 \neq k_2$. Here, we have the following cases. If $r_1 \neq r_2$ then we are in Case 3. Similarly, if $s_1 \neq s_2$ then we are in Case 4. If $r_1 = r_2, s_1 = s_2$ and $k_1 \neq k_2$ then we are in Case 2. Thus, $(a_{1,1}, a_{1,2}) = (a_{2,t_1}, a_{2,t_2})$ where $t_1, t_2 \in \{1, 2\}$ defines $T_1 \cong T_2$. So, by Cases 1, 2, 3, 4, the claim follows.

So, by $f_1, M_1 \cong M_2$.

Conversely, let $M_1 \cong M_2$. Similarly as in Lemma 4.9, let $f : V(M_1) \rightarrow V(M_2)$ such that $C_{2,j} := f(C_{1,j})$ for $j \in \{1, 2\}$. So, $\{a_{1,1}, a_{1,2}\} = \{a_{2,1}, a_{2,2}\}$. Thus, it implies that $(a_{1,1}, a_{1,2}) = (a_{2,t_1}, a_{2,t_2})$ where $t_1 \neq t_2 \in \{1, 2\}$. \square

As in Section 4, by Lemmas 5.5, 5.6, the maps of type $\{3^2, 4, 3, 4\}$ can be classified up to isomorphism. We have done calculation for the vertices ≤ 32 and listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 2.

6 Maps of type $\{3, 6, 3, 6\}$

Let M be a semi-equivelar map of type $\{3, 6, 3, 6\}$ on the torus. We define path in M as follows. Through each vertex in M there are two paths of type X_1 as shown in Figure 20.

Definition 6.1. Let $Q_1 := P(\dots, u_{i-1}, u_i, u_{i+1}, \dots)$ be a path in edge graph of M . Let $A(v)$ denote a set of incident edges through v in M . We say the path Q_1 of type X_1 if $A(u_i) \setminus E(Q_1)$ is a set of two edges where one edge lie on one side and remaining one lie on the other side of $P(u_{i-1}, u_i, u_{i+1})$.




Figure 20 : $lk(u_i)$

For example in Figure 20, $P(u_{i-1}, u_i, u_{i+1})$ and $P(b, u_i, g)$ are two paths through u_i and both are part of paths of type X_1 . Let $P(u_1, \dots, u_r)$ be a maximal path of type X_1 in M . Now, consider vertex u_r , $lk(u_r)$ and argue similarly as in Lemma 4.1. So, we get an edge $u_1 u_r$ such that $P \cup \{u_1 u_r\}$ is a cycle of type X_1 . So, there are two cycles of type X_1 through a vertex. Let $L_1(v)$, $L_2(v)$ be two cycles of type X_1 through a vertex v . We proceed with similar argument as in Section 4 and we get a connected $T(r, s, k)$ representation of M . In this process, we cut M along the cycles $L_1(v)$ and $L_2(v)$ where we take second cut along the cycle L_2 and the starting adjacent face to the base horizontal cycle L_1 is a 3-gon. Thus, every map M has a $T(r, s, k)$ representation. The Figure 10 is an example of $T(8, 2, 6)$ representation of a map with 24 vertices on the torus.

Now, we show that map of type $\{3, 6, 3, 6\}$ contains three cycles of type X_1 up to homologous.

Lemma 6.1. *The map M contains at most three cycles of type X_1 of different lengths.*

Proof. Let $\Delta(u, v, w)$ be a 3-gon in M . The $\Delta(u, v, w)$ has three edges $e_1 = uv$, $e_2 = vw$ and $e_3 = uw$. By the definition of cycle of type X_1 , M contains at least three cycles, say C_1 , C_2 and C_3 where C_i contains edge e_i for $i \in \{1, 2, 3\}$ and C_i does not contain e_j for $j \neq i$. Since C_i does not contain e_j for $j \neq i$, so, cycles are not identical. Again since, cycles are not identical and $V(C_i) \cap C(C_j)$ is a vertex of Δ for $i \neq j$, so, cycles are not homologous. (In Figure 10, $v_1 v_2 v_{1,2,16,9}$ denote a face and the cycles which are of type X_1 and contains the edges $v_1 v_2$, $v_1 v_{1,2,16,9}$ and $v_{1,2,16,9} v_2$ are namely, $L_1 = C(v_1, v_2, \dots, v_8)$, $L_2 = C(v_1, v_{1,2,16,9}, v_9, v_{9,10,6,7}, \dots, v_{11,12,8,1})$ and $L_3 = C(v_2, v_{1,2,16,9}, v_{16}, v_{15,16,4,5}, \dots, v_{13,14,2,3})$ respectively. The cycles L_1 , L_2 and L_3 in Figure 10 are not homologous to each other.) Let Δ_1 be an 3-gon in $T(r, s, k)$ and $\Delta \neq \Delta_1$. Observe that there is a cycle of type X_1 through an edge Δ_1 and homologous C_i for some i . That is, there is a cylinder which is bounded by two cycles of type X_1 and containing edges of Δ, Δ_1 . This is true for any 3-gon in $T(r, s, k)$. We proceed as in the case of Lemma 4.5 and thus, the homologous cycles of type X_1 have same length. So, there are three different cycles C_1, C_2, C_3 of type X_1 up to homologous in M . So, the map M contains at most three cycles of type X_1 of different lengths. (In Figure 10, consider face $v_{11} v_{12} v_{11,12,8,1}$ and cycles of type X_1 which are containing the edges $v_{11} v_{12}$, $v_{11} v_{11,12,8,1}$ and $v_{12} v_{11,12,8,1}$ are namely, $L'_1 = C(v_9, v_{10}, \dots, v_{16})$, $L'_2 = C(v_{11}, v_{11,12,8,1}, v_1, v_{1,2,16,9}, \dots, v_{3,4,10,11})$ and $L'_3 = C(v_{12}, v_{11,12,8,1}, v_8, v_{7,8,14,15}, \dots, v_{5,6,12,13})$ respectively. The cycles which are containing the edges $v_1 v_2$ and $v_{11} v_{12}$ are L_1 and L'_1 respectively, and the cycles L_1 and L'_1 are

homologous. The cycles which are containing $v_1v_{1,2,16,9}$ and $v_{11}v_{11,12,8,1}$ are L_2 and L'_2 respectively and $C_2 = C'_2$. Also, the cycles which are containing $v_2v_{1,2,16,9}$ and $v_{12}v_{11,12,8,1}$ are L_3 and L'_3 respectively and $L_3 = L'_3$. \square

We define admissible relations among r, s, k of $T(r, s, k)$ in the next lemma. In this lemma we omit some cases as thier similar cases are discussed in the previous sections.

Lemma 6.2. *The maps of type $\{3, 6, 3, 6\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 1$, (ii) $2 \mid r$, (iii) number of vertices of $T(r, s, k) = \frac{3}{2}rs \geq 21$, (iv) $r \geq 14$ if $s = 1$, (v) $r \geq 8$ if $s = 2$, (vi) $r \geq 6$ if $s \geq 3$, (vii) $k \in \{2t + 6 : 0 \leq t \leq \frac{r-10}{2}\} \setminus \{2(\frac{r-10}{4}) + 6\}$ if $s = 1$, $k \in \{2t + 6 : 0 \leq t \leq \frac{r-8}{2}\}$ if $s = 2$ & $k \in \{2t : 0 \leq t < \frac{r}{2}\}$ if $s \geq 3$.*

Proof. Let $C_0(u_{0,0}, u_{0,1}, \dots, u_{0,r-1})$, $C_1(u_{1,0}, u_{1,1}, \dots, u_{1,r-1}), \dots, C_{s-1}(u_{s-1,0}, u_{s-1,1}, \dots, u_{s-1,r-1})$ be horizontal cycles of type X_1 in $T(r, s, k)$. By the definition of $T(r, s, k)$, $T(r, s, k)$ contains s number of horizontal cycles of type X_1 . Observe that, the number of adjacent vertices which are lie on one side of a horizontal cycle and not belong to any horizontal cycles is $\frac{r}{2}$. So, the total number of vertices in $T(r, s, k)$ is $(r + \frac{r}{2})s$. This implies that $n = (r + \frac{r}{2})s = \frac{3}{2}rs$.

By Euler's formula, the number of 6-gons in $T(r, s, k)$ is $2n/6$ and it is an integer. This implies that $6 \mid 2n$. So, $3 \mid \frac{3}{2}sr$ as $n = \frac{3}{2}rs$. Thus, $2 \mid r$ if $s = 1$. Again, if $s \geq 2$ & $2 \nmid r$, it gives that the link $lk(u_1)$ is not type $\{3, 6, 3, 6\}$. Which is a contradiction. So, $2 \mid r \forall s \geq 1$.

Let $s = 1$. If $r < 14$ then $r \in \{2, 4, 6, 8, 10, 12\}$. If $r = 2, 4, 6, 8, 10, 12$ then there is a vertex in $T(r, s, k)$ whose link is not a cycle. So, $r \geq 14$ if $s = 1$. Similarly as above, we get that $r \geq 8$ if $s = 2$ and $r \geq 6$ if $s \geq 3$. So, $\frac{3}{2}rs \geq 21$.

If $s = 1$ and $k \in \{t : 0 \leq t \leq r-1\} \setminus (\{2t+6 : 0 \leq t \leq \frac{r-10}{2}\} \setminus \{2(\frac{r-10}{4})+6\})$ then similarly as above we get some vertex whose link is not a cycle. Similarly, we repeat same argument as above for other two cases. So, $k \in \{2t+6 : 0 \leq t \leq \frac{r-8}{2}\}$ if $s = 2$ and $k \in \{2t : 0 \leq t \leq \frac{r}{2}\}$ if $s \geq 3$. \square

Let M_i , $i = 1, 2$ be maps of type $\{3, 6, 3, 6\}$ on n_i number of vertices and $n_1 = n_2$. Let $C_{i,j}$, $j = 1, 2, 3$ denote cycles which are of type X_1 and non-homologous in $T_i = T(r_i, s_i, k_i)$. Let $a_{i,j} = \text{length}(C_{i,j})$ for $i = 1, 2$ and $j = 1, 2, 3$. Then,

Lemma 6.3. *The map $M_1 \cong M_2$ if and only if $(a_{1,1}, a_{1,2}, a_{1,3}) = (a_{2,t_1}, a_{2,t_2}, a_{2,t_3})$ for $t_1 \neq t_2 \neq t_3 \in \{1, 2, 3\}$.*

Proof. We first assume that $(a_{1,1}, a_{1,2}, a_{1,3}) = (a_{2,t_1}, a_{2,t_2}, a_{2,t_3})$ where $t_i \in \{1, 2, 3\}$ and $t_i \neq t_j$. This implies that $\{a_{1,1}, a_{1,2}, a_{1,3}\} = \{a_{2,1}, a_{2,2}, a_{2,3}\}$.

Claim : $T_1 \cong T_2$.

Case 1 : If $(r_1, s_1, k_1) = (r_2, s_2, k_2)$ then $T(r_1, s_1, k_1) = T(r_2, s_2, k_2) = T(r, s, k)$. Let $C(1, 0) := C(u_{0,0}, u_{0,1}, \dots, u_{0,r-1})$, $C(1, 1) := C(u_{1,0}, u_{1,1}, \dots, u_{1,r-1})$, \dots , $C(1, s) := C(u_{s,0}, u_{s,1}, \dots, u_{s,r-1})$ denote sequence of horizontal cycles of type X_1 in T_1 . Again, let $G_1(t, t+1) := \{w_{t,0}, w_{t,1}, \dots, w_{t, \frac{r-2}{2}}\}$ which is a set of vertices where $w_{t,i}$ is adjacent with both $u_{t,2i}$ and $u_{t+1,2i}$ in T_1 and not belong to both $C(1, t)$ and $C(1, t+1)$ for $0 \leq t \leq s$. (For example in Figure 10, $G_1(0, 1) = \{v_{1,2,16,9}, v_{3,4,10,11}, v_{5,6,12,13}, v_{7,8,14,15}\}$ where $x_{0,0} = v_{1,2,16,9}, x_{0,1} = v_{3,4,10,11}, x_{0,2} = v_{5,6,12,13}, x_{0,3} = v_{7,8,14,15}$.) Similarly, let $C(2, 0) := C(v_{0,0}, v_{0,1}, \dots, v_{0,r-1})$, $C(2, 1) := C(v_{1,0}, v_{1,1}, \dots, v_{1,r-1})$, \dots , $C(2, s) := C(v_{s,0}, v_{s,1}, \dots, v_{s,r-1})$ denote sequence of horizontal cycles of type X_1 in T_2 and define $G_2(t, t+1) :=$

Table 3: Maps of type $\{3, 6, 3, 6\}$

n	Equivalence classes	Length of cycles	$i(n)$
21	$T(14, 1, 6), T(14, 1, 10)$	$(14, 14, 14)$	$1(21)$
24	$T(16, 1, 6), T(16, 1, 12)$ $T(8, 2, 6)$	$(16, 16, 8)$	$2(24)$
27	$T(16, 1, 8), T(16, 1, 10)$ $T(18, 1, 6), T(18, 1, 8)$ $T(18, 1, 12), T(18, 1, 14)$ $T(6, 3, 2), T(6, 3, 4)$ $T(6, 3, 0)$	$(16, 16, 4)$ $(18, 18, 6)$ $(6, 6, 6)$	$2(27)$
30	$T(20, 1, 6), T(20, 1, 8)$ $T(20, 1, 14), T(20, 1, 16)$ $T(10, 2, 2), T(10, 2, 6)$ $T(10, 2, 8)$ $T(20, 1, 10), T(20, 1, 12)$ $T(10, 2, 0), T(10, 2, 4)$	$(20, 20, 10)$ $(20, 4, 10)$	$2(30)$

$\{x_{t,0}, x_{t,1}, \dots, x_{t, \frac{r-2}{2}}\}$ which is a set of vertices where the vertex $x_{t,i}$ is adjacent with both $v_{t,2i}$ and $v_{t+1,2i}$ in T_2 for $0 \leq t \leq s$. Now we define an isomorphism map $f : V(T(r_1, s_1, k_1)) \rightarrow V(T(r_2, s_2, k_2))$ such that $f(u_{t,i}) = v_{t,i} \forall 0 \leq i \leq r-1, 0 \leq t \leq s-1$ and $f(w_{t,i}) = x_{t,i}$ for the vertices of $G_1(t, t+1)$ and $G_2(t, t+1)$ for all $0 \leq t \leq s-1$. By f , the link $lk(u_{t,i})$ maps to the link of $lk(v_{t,i})$ and $lk(w_{t,i})$ maps to $lk(x_{t,i})$. So, by f , $T_1 \cong T_2$.

Case 2 : Let $(r_1, s_1, k_1) \neq (r_2, s_2, k_2)$. If $r_1 \neq r_2$, we identify boundaries of $T(r_2, s_2, k_2)$ and cut M_2 along cycle of length r_1 and followed by, make another cut along a cycle of type X_1 to get a (r, s, k) -representation. Thus we get a new $T(r'_2, s'_2, k'_2)$ representation of M_2 . It implies that $r_1 = r'_2$ and $s_1 = s'_2$ as $n_1 = \frac{3}{2}r_1s_1 = \frac{3}{2}r'_2s'_2 = n_2$. By this process, we get a new representation $T(r_1, s_1, k'_3)$ of M_2 . If $k_1 = k'_3$ then $M_1 \cong M_2$ by f in Case 1. If $k_1 \neq k'_3$ similarly as in Lemma 5.6, we make a cut along a path which is homologous to boundary path and identify along the boundary path. Thus, we get an another representation $T(r_1, s_1, k''_3)$ of M_2 and $k_1 = k''_3$. So, the M_2 has a $T(r_1, s_1, k_1)$ representation since $k_1 = k''_3$. Therefore, there exists f and $T_1 \cong T_2$ by f . This completes the Claim.

So, by f , $M_1 \cong M_2$.

Conversely, let $M_1 \cong M_2$. We proceed as in the converse part of Lemma 5.6 and we get $\{a_{1,1}, a_{1,2}, a_{1,3}\} = \{a_{2,1}, a_{2,2}, a_{2,3}\}$. That is, $(a_{1,1}, a_{1,2}, a_{1,3}) = (a_{2,t_1}, a_{2,t_2}, a_{2,t_3})$ for $t_1 \neq t_2 \neq t_3 \in \{1, 2, 3\}$. \square

As in Section 4, by Lemmas 6.2, 6.3, the maps of type $\{3, 6, 3, 6\}$ can be classified up to isomorphism on different number of vertices. We have done the calculation for the vertices ≤ 30 . We have listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 3.

7 Maps of type $\{3, 12^2\}$

Let M be a semi-equivelar map of type $\{3, 12^2\}$ on the torus. We define a fixed type of path G_1 in the edge graph of M as shown in Figure 21. Let $Q(i) := P(u_i, u_{i+1}, u_{i+2}, u_{i+3}, u_{i+4})$ be a path in M where $lk(u_i) = C(u_{i-1}, \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \mathbf{f}, \mathbf{g}, \mathbf{g}', \mathbf{u}_{i+2}, \mathbf{u}_{i+1}, \mathbf{u}, \mathbf{t}, \mathbf{v}, \mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{d}', \mathbf{u}_{i-3}, \mathbf{u}_{i-2})$, $lk(u_{i+1}) = C(u_i, \mathbf{u}_{i-1}, \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \mathbf{f}, \mathbf{g}, \mathbf{g}', \mathbf{u}_{i+2}, \mathbf{u}_{i+3}, \mathbf{u}_{i+4}, \mathbf{d}', \mathbf{o}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{t}, \mathbf{u})$, $lk(u_{i+2}) = C(u_{i+1}, \mathbf{u}_i, \mathbf{u}_{i-1}, \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \mathbf{f}, \mathbf{g}, \mathbf{g}', \mathbf{u}_{i+3}, \mathbf{u}_{i+4}, \mathbf{d}', \mathbf{o}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{t}, \mathbf{u})$, $lk(u_{i+3}) = C(u_{i+2}, \mathbf{g}', \mathbf{g}, \mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{u}_{i+6}, \mathbf{u}_{i+5}, \mathbf{u}_{i+4}, \mathbf{d}', \mathbf{o}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{u}_{i+1})$, and $lk(u_{i+4}) = C(u_{i+3}, \mathbf{g}', \mathbf{g}, \mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{u}_{i+6}, \mathbf{u}_{i+5}, \mathbf{d}', \mathbf{o}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{u}_{i+1}, \mathbf{u}_{i+2})$.

Definition 7.1. Let $R_1 := P(\dots, v_{i-1}, v_i, v_{i+1}, \dots)$ be a path in edge graph of M . We say R_1 of type G_1 if $L_1 := P(v_t, v_{t+1}, v_{t+2}, v_{t+3}, v_{t+4})$ is a subpath of R_1 or is a part in

an extended path of R_1 then either $L_1 \mapsto Q(i)$ by $v_j \mapsto u_j$, $L_1 \mapsto Q(i+1)$ by $v_j \mapsto u_{j+1}$, $L_1 \mapsto Q(i+2)$ by $v_j \mapsto u_{j+2}$ or $L_1 \mapsto Q(i+3)$ by $v_j \mapsto u_{j+3}$ for $j \in \{t, t+1, t+2, t+3, t+4\}$.

Definition 7.2. Let $R_2 := P(\dots, x_{i-1}, x_i, x_{i+1}, \dots)$ be a path in edge graph of M . We say R_2 of type G'_1 if $L_2 := P(x_t, x_{t+1}, x_{t+2}, x_{t+3}, x_{t+4})$ is a subpath of R_2 or is a part in the extended path of R_2 then either $L_2 \mapsto Q(i)$ by $x_j \mapsto u_{2t+4-j}$, $L_2 \mapsto Q(i+1)$ by $x_j \mapsto u_{2t+4-j}$, $L_2 \mapsto Q(i+2)$ by $x_j \mapsto u_{2t+4-j}$ or $L_1 \mapsto Q(i+3)$ by $v_j \mapsto u_{2t+4-j}$ for $j \in \{t, t+1, t+2, t+3, t+4\}$.




Figure 21 : Cycle of type G_1

We argue with the similar argument of Lemma 4.1 for the path of types G_1, G'_1 and so, every maximal path of types G_1, G'_1 is a cycle and non-contractible (by the similar argument of Lemma 4.3). Observe that the cycles of type G_1 and G'_1 are mirror image of each other. So, these types define same type of cycle. (Similar argument of this is given details in Section 8 for the type $\{3^4, 6\}$.) Clearly, there are two cycles of type G_1 through each vertex of M . Let uvw be a 3-gon in M . Let $L_1(u, uw)$, $L_2(w, wv)$ and $L_3(v, vu)$ denote three cycles through u , w , and v respectively where $L_1(u, uw)$ contains the edge uw , $L_2(w, wv)$ contains the edge wv and $L_3(v, vu)$ contains the edge vu . We repeat the similar argument of Section 4 and define a $T(r, s, k)$ representation of the map M for some r, s, k . In the process, we take first cut along $L_1(u, uw)$, and then, second cut along $L_2(w, wv)$ where the starting adjacent face to horizontal base cycle $L_1(u, uw)$ is a 12-gon. Let $\text{length}(L_1(u, uw)) = r$, s denote the number of homologous cycles of $L_1(u, uw)$ of type G_1 , and k denote the distance of the starting vertex of upper horizontal cycle from the starting vertex w in $L_1(u, uw)$. By this process, we get a $T(r, s, k)$ representation of M . Now, we proceed with process of Section 6 for the map of type $\{3, 12^2\}$. We show that a map of type $\{3, 12^2\}$ contains at most three non-homologous cycles of type G_1 of different lengths.

Lemma 7.1. *The map M contains at most three cycles of type G_1 of different lengths.*

Proof. We proceed as in the case of Lemma 6.1 for the map of type $\{3, 12^2\}$. So, consider map of type $\{3, 12^2\}$ in place of $\{3, 6, 3, 6\}$, cycle of type G_1 in place of X_1 and 3-gon in the proof of Lemma 6.1. Thus, we get three non-homologous cycles of type G_1 of different lengths. \square

We define admissible relations among r, s, k of $T(r, s, k)$ in M .

Lemma 7.2. *The maps of type $\{3, 12^2\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 1$, (ii) $4 \mid r$, (iii) number of vertices of $T(r, s, k) = \frac{3}{2}rs \geq 36$, (iv) $r \geq 24$ if $s = 1$, (v) $r \geq 16$ if $s = 2$, (vi) $r \geq 12$ if $s \geq 3$, (vii) $k \in \{4t+9 : 0 \leq t \leq \frac{r-20}{4}\} \setminus \{4(\frac{r}{8}-3)+9\}$ if $s = 1$, $k \in \{4t+5 : 0 \leq t \leq \frac{r-16}{4}\}$ if $s = 2$ & $k \in \{4t+1 : 0 \leq t \leq \frac{r-4}{4}\}$ if $s \geq 3$.*

Proof. We proceed as in the case of proof of the Lemma 6.2. We consider map of type $\{3, 12^2\}$ in place of $\{3, 6, 3, 6\}$ and different values of r, s, k . Thus, we get all the cases. \square

Table 4: Maps of type $\{3, 12^2\}$

n	Equivalence classes	Length of cycles	$i(n)$
36	$T(24, 1, 13)$	$(24, 12, 8)$	1(36)
42	$T(28, 1, 9), T(28, 1, 13)$ $T(28, 1, 17)$	$(28, 28, 28)$	1(42)
48	$T(32, 1, 9), T(32, 1, 21)$ $T(16, 2, 5)$ $T(32, 1, 17)$	$(32, 32, 16)$ $(32, 32, 8)$	2(48)

Let $T_i = T(r_i, s_i, k_i)$ be representations of M_i , $i = 1, 2$ on same number of vertices. Similarly as in Section 6, let $b_{i,j} = \text{length}(L_{i,j})$, $j = 1, 2, 3$. Similarly as in Section 6, we have

Lemma 7.3. *The map $M_1 \cong M_2$ if and only if $(b_{1,1}, b_{1,2}, b_{1,3}) = (b_{2,t_1}, b_{2,t_2}, b_{2,t_3})$ for $t_1 \neq t_2 \neq t_3 \in \{1, 2, 3\}$.*

As in Section 4, by Lemmas 7.2, 7.3, maps of type $\{3, 12^2\}$ can be classified up to isomorphism on different number of vertices. We have done the calculation for the vertices ≤ 48 . We have listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 4.

8 Maps of type $\{3^4, 6\}$

Let M be a semi-equivelar map of type $\{3^4, 6\}$ on the torus. Let $Q(i) := P(w_i, w_{i+1}, w_{i+2}, w_{i+3})$ be a path in M , where $lk(w_i) = C(w_{i-1}, x_2, w_{i+1}, w_{i+2}, x_3, x_4, x_5, x_6)$, $lk(w_{i+1}) = C(w_i, x_2, x_7, x_8, w_{i+2}, x_3, x_4, x_5)$, $lk(w_{i+2}) = C(w_{i+1}, x_8, x_9, w_{i+3}, x_3, x_4, x_5, w_i)$, $lk(w_{i+3}) = C(w_{i+2}, x_9, w_{i+4}, w_{i+5}, x_{10}, x_{11}, x_{12}, x_3)$. We define two fixed types of paths Y_1 and Y_1' in the edge graph of M .

Definition 8.1. *Let $R_1 := P(\dots, v_{i-1}, v_i, v_{i+1}, \dots)$ be a path in edge graph of M . We say R_1 of type Y_1 if $L_1 := P(u_t, u_{t+1}, u_{t+2}, u_{t+3})$ is a sub-path of R_1 or in the extended path of R_1 then $L_1 \mapsto Q(i)$ by $u_j \mapsto w_j$, $L_1 \mapsto Q(i+1)$ by $u_j \mapsto w_{j+1}$ or $L_1 \mapsto Q(i+2)$ by $u_j \mapsto w_{j+2}$ for $j \in \{t, t+1, t+2, t+3\}$.*

Definition 8.2. *Let $R_2 := P(\dots, x_{i-1}, x_i, x_{i+1}, \dots)$ be a path in edge graph of M . We say R_2 of type Y_1' if $L_2 := P(x_t, x_{t+1}, x_{t+2}, x_{t+3})$ is a sub-path of R_2 or in the extended path of R_2 then $L_2 \mapsto Q(i)$ by $x_j \mapsto w_{2t+3-j}$, $L_2 \mapsto Q(i+1)$ by $x_j \mapsto w_{2t+3-j}$ or $L_2 \mapsto Q(i+2)$ by $x_j \mapsto w_{2t+3-j}$ for $j \in \{t, t+1, t+2, t+3\}$.*

Let P be a maximal path of type Y_1 or Y_1' . By the similar argument of Lemma 4.1, path P defines a cycle of type Y_1 or Y_1' , that is, there is an edge e in M such that $P \cup \{e\}$ is a cycle of type Y_1 or Y_1' . We show that the cycles of types Y_1 and Y_1' define same type of cycle. Let $C_1 := C(u_1, u_2, \dots, u_r)$ of type Y_1 and $C_2(v_1, v_2, \dots, v_r)$ of type Y_1' be two cycles of same length r . Let $P_1 := P(u_{i-1}, u_i, u_{i+1})$ be a sub path of C_1 where adjacent 6-gon lies on one side and all 3-gons lie on the other side of P_1 at the vertex u_i . Similarly, let $P_2 := P(v_{j-1}, v_j, v_{j+1})$ be a sub path of C_2 where adjacent 6-gon lies on one side and all 3-gons lie on the other side of P_2 at the vertex v_j . Define a map $f : V(C_1) \rightarrow V(C_2)$ by $f(u_i) = v_j$, $f(u_{i+1}) = v_{j-1}$, $f(u_{i+2}) = v_{j-2}, \dots$, $f(u_{i-1}) = v_{j+1}$. Let $P(u_t, \dots, u_k)$ be a sub path of C_1 . Then, $P(u_t, \dots, u_k)$ and $P(f(u_t), \dots, f(u_k))$ divide the link of the vertices u_i and $f(u_i)$ for $t \leq i \leq k$ into the same ratio. This is true for every sub path of C_1 . Therefore, cycles C_1 and C_2 are of type Y_1 , and hence, $Y_1 = Y_1'$.

Let M be a map and C be a cycle of type Y_1 in M . By the similar argument of Lemma 4.3, the cycle C is non-contractible. As in Section 4, we get that the cycles of type Y_1 which are homologous to C have same length by the similar argument of Lemma 4.5. There are three cycles of type Y_1 through each vertex of M . Let $v \in V(M)$ and $L_1(v)$, $L_2(v)$, $L_3(v)$ be three cycles of type Y_1 through the vertex v . We repeat similar construction of (r, s, k) -representation of a map as in Section 4 for M . In this process, we take first cut along $L_1(v)$ and second cut along $L_2(v)$ where the starting adjacent face to horizontal base cycle L_1 is a 3-gon. This gives a $T(r, s, k)$ representation of the map M . Thus, $T(r, s, k)$ representation exists for every M .

Now, we show that map M of type $\{3^4, 6\}$ contains at most three non-homologous cycles of type Y_1 of different lengths in Lemma 8.1.

Lemma 8.1. *The map M contains at most three non-homologous cycles of type Y_1 of different lengths.*

Proof. Let $v \in V(M)$ and $T(r, s, k)$ denote a (r, s, k) -representation of M . We have three cycles, namely, C_1 , C_2 and C_3 through v in M of type Y_1 . Cycles C_1 , C_2 and C_3 are not identical as C_i divides link $lk(v)$ into different ratio. Also, cycles are not disjoint as $v \in V(C_i) \cap V(C_j)$ for $i \neq j$ and $i, j \in \{1, 2, 3\}$. So, C_1 , C_2 and C_3 are not homologous to each other. Let $w \in V(M)$ and $v \neq w$. Consider cycles of type Y_1 at w in $T(r, s, k)$ and denoted by C'_1 , C'_2 , C'_3 . Now, by the definition of cycle of type Y_1 and considering cylinder, C_i and C'_j are homologous for some $i, j \in \{1, 2, 3\}$ as we have seen the same idea in Lemma 6.1. This is hold for any vertex of M . Therefore, M contains at most three non-homologous cycles of type Y_1 . We proceed as in the case of proof of Lemma 4.5 to show that the homologous cycles of type Y_1 have same length. Thus, the map M contains at most three non-homologous cycles of type Y_1 of different lengths. \square

We define admissible relations among r , s , k of $T(r, s, k)$ such that representation $T(r, s, k)$ gives a map of type $\{3^4, 6\}$ after identifying their boundaries.

Lemma 8.2. *The maps of type $\{3^4, 6\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 2$ even, (ii) $3 \mid r$, (iii) number of vertices of $T(r, s, k) = rs \geq 18$, (iv) $r \geq 9$ if $s = 2$, (v) $r \geq 6$ if $s \geq 4$, (vi) $k \in \{3t+5 : 0 \leq t \leq \frac{r-9}{3}\}$ if $s = 2$ & $k \in \{2+3t : 0 \leq t \leq \frac{r-3}{3}\}$ if $s \geq 4$.*

Proof. Let $T(r, s, k)$ be a representation of M . It has s number of disjoint horizontal cycles of type Y_1 of length r by the definition of (r, s, k) -representation. These cycles cover all the vertices of M . So, $n = rs$. By Euler's formula, $n - 5n/2 + 4n/3 + n/6 = 0$. So, the number of 6-gons in M is $n/6$ and which is an integer. This implies that $6 \mid n$. That is, $6 \mid rs$ as $n = rs$. So, $3 \mid r$ for $s = 2$. Let $s \geq 3$. If s is an odd integer then $T(r, s, k)$ contains odd number of horizontal cycles of type Y_1 . Consider a vertex v of base horizontal cycle which is belongs to only triangles which is a contradiction. So, $2 \mid s$. Similarly, for $3 \mid r$, we get a vertex whose link does not follow the type $\{3^4, 6\}$. So, $n = rs$ where $6 \mid n$, $2 \mid s$ and $3 \mid r$.

For $r \geq 9$, we proceed with similar argument as it is done in the proof of Lemma 4.8. We also proceed as in the case of the proof of Lemma 4.8 to show $r \geq 6$ and $3 \mid r$ and remaining all other cases. This completes the proof. \square

Let M_1 and M_2 be two maps of type $\{3^4, 6\}$ on same number of vertices. Similarly as in Sections 4, 5, let $a_{i,j} = \text{length}(C_{i,j})$ where $C_{i,k}$ for $i = 1, 2, 3$ denote three non-homologous cycles of type Y_1 in $T(r_i, s_i, k_i)$ of M_i . Then,

Table 5: Maps of type $\{3^4, 6\}$

n	Equivalence classes	Length of cycles	$i(n)$
18	$T(9, 2, 5)$	$(9, 9, 9)$	1(18)
24	$T(12, 2, 5), T(12, 2, 8)$	$(12, 6, 12)$	2(24)
	$T(6, 4, 2)$		
	$T(6, 4, 5)$	$(6, 6, 6)$	
30	$T(15, 2, 5), T(15, 2, 8)$	$(15, 15, 15)$	1(30)
	$T(15, 2, 11)$		
36	$T(18, 2, 5), T(18, 2, 14)$	$(18, 9, 18)$	2(36)
	$T(9, 4, 2)$		
	$T(18, 2, 8), T(18, 2, 11)$	$(18, 6, 9)$	
	$T(9, 4, 5), T(9, 4, 8)$		
	$T(6, 6, 2), T(6, 6, 5)$		
42	$T(21, 2, 5), T(21, 2, 8)$	$(21, 21, 21)$	1(42)
	$T(21, 2, 11), T(21, 2, 14)$		
	$T(21, 2, 17)$		

Lemma 8.3. *The map $M_1 \cong M_2$ if and only if $(a_{1,1}, a_{1,2}, a_{1,3}) = (a_{2,t_1}, a_{2,t_2}, a_{2,t_3})$ for $t_1 \neq t_2 \neq t_3 \in \{1, 2, 3\}$.*

Proof. Let $T(r_i, s_i, k_i)$ be a representation of M_i . If $r = r_1, s = s_1, k = k_1$, then, consider horizontal cycles in $T(r_i, s_i, k_i)$ of type Y_1 . Proceed with similar argument as in the Lemma 4.9. So, we get a map which defines isomorphism between $T(r_1, s_1, k_1)$ and $T(r_2, s_2, k_2)$. So, $M_1 \cong M_2$. Again, if $(r, s, k) \neq (r_1, s_1, k_1)$ then proceed with similar argument as in the proof of the Lemma 5.6. Converse part of the lemma follows from similar argument of the converse part of Lemma 4.9. This completes the proof. \square

As in Section 4, by Lemmas 8.2, 8.3, the maps of type $\{3^4, 6\}$ can be classified up to isomorphism on different number of vertices. We have done calculation for the vertices ≤ 42 . We have listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 5.

9 Maps of type $\{4, 6, 12\}$

Let M be a semi-equivelar map of type $\{4, 6, 12\}$ on the torus. We define a fixed type of path H_1 in the edge graph of M . Let $Q(i) := P(u_i, u_{i+1}, u_{i+2}, u_{i+3}, u_{i+4}, u_{i+5}, u_{i+6})$ be a path in M where $lk(u_i) = C(u_{i-1}, \mathbf{b}, \mathbf{c}, \mathbf{u}_{i+2}, u_{i+1}\mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{v}, \mathbf{u}_{i-5}, \mathbf{u}_{i-4}, \mathbf{u}_{i-3}, \mathbf{u}_{i-2})$, $lk(u_{i+1}) = C(u_i, \mathbf{u}_{i-1}\mathbf{b}, \mathbf{c}, \mathbf{u}_{i+2}, \mathbf{u}_{i+3}, \mathbf{u}_{i+4}, \mathbf{u}_{i+5}, \mathbf{u}_{i+6}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{o}, \mathbf{p}, \mathbf{q})$, $lk(u_{i+2}) = C(u_{i+1}, \mathbf{u}_i, \mathbf{u}_{i-1}\mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{u}_{i+3}, \mathbf{u}_{i+4}, \mathbf{u}_{i+5}, \mathbf{u}_{i+6}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{o}, \mathbf{p})$, $lk(u_{i+3}) = C(u_{i+2}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \mathbf{f}, \mathbf{g}, u_{i+4}\mathbf{u}_{i+5}\mathbf{u}_{i+6}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{o}, \mathbf{p}, \mathbf{u}_{i+1})$, $lk(u_{i+4}) = C(u_{i+3}, \mathbf{d}, \mathbf{e}, \mathbf{f}, \mathbf{g}, \mathbf{h}, u_{i+5}\mathbf{u}_{i+6}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{o}, \mathbf{p}, \mathbf{u}_{i+1}, \mathbf{u}_{i+2})$, $lk(u_{i+5}) = C(u_{i+4}, \mathbf{g}, \mathbf{h}, \mathbf{i}, \mathbf{u}_{i+8}, \mathbf{u}_{i+7}, u_{i+6}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{o}, \mathbf{p}, \mathbf{u}_{i+1}, \mathbf{u}_{i+2}, \mathbf{u}_{i+3})$, and $lk(u_{i+6}) = C(u_{i+5}, \mathbf{h}, \mathbf{i}, \mathbf{u}_{i+8}, u_{i+7}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{n}, \mathbf{o}, \mathbf{p}, \mathbf{u}_{i+1}, \mathbf{u}_{i+2}, \mathbf{u}_{i+3}, \mathbf{u}_{i+4})$.

Definition 9.1. *Let $P_1 := P(\dots, v_{i-1}, v_i, v_{i+1}, \dots)$ be a path in edge graph of M . We say P_1 of type H_1 if $L_1 := P(v_t, v_{t+1}, v_{t+2}, v_{t+3}, v_{t+4}, v_{t+5}, v_{t+6})$ is a subpath of P_1 or lies in the extended path of P_1 then either $L_1 \mapsto Q(i)$ by $v_j \mapsto u_j$, $L_1 \mapsto Q(i+1)$ by $v_j \mapsto u_{j+1}$, $L_1 \mapsto Q(i+2)$ by $v_j \mapsto u_{j+2}$, $L_1 \mapsto Q(i+3)$ by $v_j \mapsto u_{j+3}$, $L_1 \mapsto Q(i+4)$ by $v_j \mapsto u_{j+4}$ or $L_1 \mapsto Q(i+5)$ by $v_j \mapsto u_{j+5}$ for $j \in \{t, t+1, t+2, t+3, t+4, t+5, t+6\}$.*

Definition 9.2. *Let $P_2 := P(\dots, x_{i-1}, x_i, x_{i+1}, \dots)$ be a path in edge graph of M . We say P_2 of type H'_1 if $L_2 := P(x_t, x_{t+1}, x_{t+2}, x_{t+3}, x_{t+4}, x_{t+5}, x_{t+6})$ is a subpath of P_2 or lies in the extended path of P_2 then either $L_2 \mapsto Q(i)$ by $x_j \mapsto u_{2t+6-j}$, $L_2 \mapsto Q(i+1)$ by $x_j \mapsto u_{2t+6-j}$, $L_2 \mapsto Q(i+2)$ by $x_j \mapsto u_{2t+6-j}$, $L_2 \mapsto Q(i+3)$ by $v_j \mapsto u_{2t+6-j}$, $L_2 \mapsto Q(i+4)$ by $v_j \mapsto u_{2t+6-j}$ or $L_2 \mapsto Q(i+5)$ by $v_j \mapsto u_{2t+6-j}$ for $j \in \{t, t+1, t+2, t+3, t+4, t+5, t+6\}$.*

Let P be a maximal path of type H_1 or H'_1 in M . By the similar argument of Lemma 4.1, the map M contains an edge e which defines a cycle $P \cup \{e\}$ of type H_1 or H'_1 . The cycle $C := P \cup \{e\}$ is a non-contractible cycle (by the similar argument of Lemma 4.3). Observe that the cycles of types H_1 and H'_1 are mirror image of each other. So, it follows that they define same type of cycles as in Section 8. So, we consider only type H_1 . Let C_1, C_2, \dots, C_m be a sequence of homologous cycles of type H_1 in M . We argue with the similar argument of Lemma 4.5 and we get that $\text{length}(C_i) = \text{length}(C_j)$ $1 \leq i, j \leq m$. By Definition 9.1, there are three cycles of type H_1 through each vertex of M . Let $v \in V(M)$ and $L_1(v), L_2(v), L_3(v)$ denote three cycles through v . Define a $T(r, s, k)$ representation of M by the similar construction which is given in Section 4. In this process, we cut M along L_1 and then, take second cut along the cycle L_3 where the starting adjacent face to base horizontal cycle L_1 is a 6-gon. So, every map M has a $T(r, s, k)$ representation. In Lemma 9.1, we show that map of type $\{4, 6, 12\}$ contains at most three non-homologous cycles of type H_1 of different lengths.

Lemma 9.1. *The map M contains at most three non-homologous cycles of type H_1 of different lengths.*

Proof. We proceed as in the case of proof of the Lemma 8.1. In this process, consider map $\{4, 6, 12\}$ in place of $\{3^4, 6\}$ and cycle of type H_1 in place of Y_1 . Let $u \in V(M)$ and $T(r, s, k)$ denote a (r, s, k) -representation of M . Let L_1, L_2 and L_3 denote three cycles of type H_1 through u in M . They are not identical as L_i divides link $lk(u)$ into different ratio. Also, cycles are not disjoint as $u \in V(L_i) \cap V(L_j)$ for $i \neq j$. Therefore, cycles are not homologous to each other. Again, let $v \in V(M)$, $u \neq v$ and consider cycles of type H_1 at v in $T(r, s, k)$. Let L'_1, L'_2, L'_3 denote three cycles through v of type H_1 . Then, by the definition of cycle of type H_1 and considering cylinder, L_i and L'_j are homologous for some $i, j \in \{1, 2, 3\}$. This is hold for any vertex v of M . Thus, M contains at most three non-homologous cycles of type H_1 . We proceed as in the case of proof of Lemma 4.5 to show that the homologous cycles of type H_1 have same length. This completes the proof. \square

We define admissible relations among r, s, k of $T(r, s, k)$ such that representation $T(r, s, k)$ gives a map of type $\{4, 6, 12\}$ after identifying their boundaries.

Lemma 9.2. *The maps of type $\{4, 6, 12\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 2$ even, (ii) $6 \mid r$, (iii) number of vertices of $T(r, s, k) = rs \geq 36$, (iv) $r \geq 18$ if $s = 2$, (v) $r \geq 12$ if $s \geq 4$, (vi) $k \in \{6t + 9 : 0 \leq t \leq \frac{r-18}{6}\}$ if $s = 2$ & $k \in \{6t + 3 : 0 \leq t \leq \frac{r-6}{6}\}$ if $s \geq 4$.*

Proof. We proceed as in the case of proof of the Lemma 8.2. We proof this lemma by considering link of some vertices in $T(r, s, k)$. We consider map of type $\{4, 6, 12\}$ in place of type $\{3^4, 6\}$ and different values of r, s and k in the proof of Lemma 8.2. Thus, we get all possible ranges of r, s and k of $T(r, s, k)$. This completes the proof. \square

Let M_1 and M_2 be two maps of type $\{4, 6, 12\}$ on same number of vertices. Let $T(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i) -representation of M_i . Similarly as in Sections 4, 5, by Lemma 9.1, let $b_{i,j} = \text{length}(L_{i,j})$ where $L_{i,j}$, $j = 1, 2, 3$ denote non-homologous cycles of type H_1 in $T(r_i, s_i, k_i)$. Then

Lemma 9.3. *The map $M_1 \cong M_2$ if and only if $(b_{1,1}, b_{1,2}, b_{1,3}) = (b_{2,t_1}, b_{2,t_2}, b_{2,t_3})$ for $t_1 \neq t_2 \neq t_3 \in \{1, 2, 3\}$.*

Table 6: Maps of type $\{4, 6, 12\}$

n	Equivalence classes	Length of cycles	$i(n)$
36	$T(18, 2, 9)$	(18, 18, 18)	1(36)
48	$T(24, 2, 9), T(12, 4, 9)$	(24, 12, 24)	2(48)
	$T(24, 2, 15)$	(12, 12, 12)	
60	$T(12, 4, 3)$	(30, 30, 30)	1(60)
	$T(30, 2, 9), T(30, 2, 15)$		
	$T(30, 2, 21)$		

Proof. We proceed as in the case of the proof of Lemma 8.3. Let $r = r_1, s = s_1, k = k_1$. Consider horizontal cycles in $T(r_i, s_i, k_i)$ of type H_1 . We proceed with similar argument as in Lemma 4.9. So, we get $M_1 \cong M_2$. Again, if $(r, s, k) \neq (r_1, s_1, k_1)$ then we proceed as in the case of proof of Lemmas 5.6 and 6.3. Converse part follows from similar argument of the converse part of Lemma 4.9. This completes the proof. \square

As in Section 4, by Lemmas 9.2 and 9.3, the maps of type $\{4, 6, 12\}$ can be classified on different number of vertices. We have done calculation for the vertices ≤ 60 . We have listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 6.

10 Maps of type $\{3, 4, 6, 4\}$

Let M be a semi-equivelar map of type $\{3, 4, 6, 4\}$ on the torus. We define path W_1 in the edge graph of M on the torus. Let $Q(i) := P(u_i, u_{i+1}, u_{i+2}, u_{i+3})$ be a path in M , where $lk(u_i) = C(u_{i-1}, \mathbf{a}, b, u_{i+1}, \mathbf{i}, j, \mathbf{k}, \mathbf{l}, u_{i-2})$, $lk(u_{i+1}) = C(u_i, b, \mathbf{c}, u_{i+2}, u_{i+3}, \mathbf{g}, \mathbf{h}, i, \mathbf{j})$, $lk(u_{i+2}) = C(u_{i+1}, \mathbf{b}, c, d, \mathbf{e}, u_{i+3}, \mathbf{g}, \mathbf{h}, \mathbf{i})$ and $lk(u_{i+3}) = C(u_{i+2}, \mathbf{d}, e, u_{i+4}, \mathbf{f}, g, \mathbf{h}, \mathbf{i}, u_{i+1})$

Definition 10.1. Let $P_1 := P(\dots, v_{i-1}, v_i, v_{i+1}, \dots)$ be a path in edge graph of M . We say P_1 of type W_1 if $L_1 := P(v_t, v_{t+1}, v_{t+2}, v_{t+3})$ is a subpath of P_1 or in a path containing P_1 . In this case, either $L_1 \mapsto Q(i)$ by $v_j \mapsto u_j$, $L_1 \mapsto Q(i+1)$ by $v_j \mapsto u_{j+1}$ or $L_1 \mapsto Q(i+2)$ by $v_j \mapsto u_{j+2}$ for $j \in \{t, t+1, t+2, t+3\}$.

Definition 10.2. Let $P_2 := P(\dots, x_{i-1}, x_i, x_{i+1}, \dots)$ be a path in edge graph of M . We say P_2 of type W'_1 if $L_2 := P(x_t, x_{t+1}, x_{t+2}, x_{t+3})$ is a subpath of P_2 or in a path containing P_2 . In this case, either $L_2 \mapsto Q(i)$ by $x_j \mapsto u_{2t+3-j}$, $L_2 \mapsto Q(i+1)$ by $x_j \mapsto u_{2t+3-j}$ or $L_2 \mapsto Q(i+2)$ by $x_j \mapsto u_{2t+3-j}$ for $j \in \{t, t+1, t+2, t+3\}$.

We consider only cycle of type W_1 as W_1 and W'_1 define same type of cycle (by the similar argument as in Section 8). Repeat the similar argument which is given in Section 9 and define a $T(r, s, k)$ representation. In this process, we consider path of type W_1 in place of H_1 . Now, by Definition 10.1, there are three cycles through each vertex of M . Let $v \in V(M)$ and L_1, L_2 and L_3 be three cycles through v . We cut along L_1 and then, take second cut along L_3 where the starting adjacent face to base horizontal cycle L_1 is a 4-gon. So, every map has this $T(r, s, k)$ representation. In Lemma 10.1, we show that map of type $\{3, 4, 6, 4\}$ contains at most three non-homologous cycles of type W_1 of different lengths.

Lemma 10.1. *The map M contains at most three non-homologous cycles of type W_1 of different lengths.*

Proof. As above, proceed with similar argument of Lemma 8.1. Consider map of type $\{3, 4, 6, 4\}$ in place of $\{3^4, 6\}$ and cycle of type W_1 in place of Y_1 . Let $w_1 \neq w_2$ be two vertices of M . Let J_1, J_2, J_3 denote three cycles through w_1 and J'_1, J'_2, J'_3 denote three

Table 7: Maps of type $\{3, 4, 6, 4\}$

n	Equivalence classes	Length of cycles	$i(n)$
18	$T(9, 2, 4)$	(9, 9, 9)	1(18)
24	$T(12, 2, 4), T(12, 2, 7), T(6, 4, 4)$ $T(6, 4, 1)$	(12, 6, 12) (6, 6, 6)	2(24)
30	$T(15, 2, 4), T(15, 2, 7), T(15, 2, 10)$	(15, 15, 15)	1(30)
36	$T(18, 2, 4), T(18, 2, 13), T(9, 4, 7)$ $T(18, 2, 7), T(18, 2, 10), T(9, 4, 1)$ $T(9, 4, 4), T(6, 6, 4), T(6, 6, 1)$	(18, 9, 18) (18, 9, 6)	2(36)
42	$T(21, 2, 4), T(21, 2, 7), T(21, 2, 10)$ $T(21, 2, 13), T(21, 2, 16)$	(21, 21, 21)	1(42)
48	$T(24, 2, 4), T(24, 2, 7), T(24, 2, 16)$ $T(24, 2, 19), T(12, 4, 4), T(12, 4, 10)$ $T(24, 2, 10), T(24, 2, 13), T(6, 8, 4)$ $T(12, 4, 1), T(12, 4, 7), T(6, 8, 1)$	(24, 24, 12) (24, 24, 6) (12, 12, 6)	3(48)
54	$T(27, 2, 4), T(27, 2, 13), T(27, 2, 22)$ $T(27, 2, 7), T(27, 2, 10), T(27, 2, 16)$ $T(27, 2, 19), T(9, 6, 4), T(9, 6, 7)$ $T(9, 6, 1)$	(27, 27, 27) (27, 27, 9)	3(54)
		(9, 9, 9)	

cycles through w_2 . Then, by the definition of cycle of type W_1 , we get a cylinder which is bounded by J_i and J'_j . That is, J_i and J'_j are homologous for some $i, j \in \{1, 2, 3\}$. This is hold for arbitrary vertex of M . We proceed as in the case of proof of Lemma 4.5 to show that the homologous cycles of type W_1 have same length. Thus, the map M contains at most three non-homologous cycles of type W_1 of different lengths. This completes the proof. \square

We define admissible relations among r, s, k of $T(r, s, k)$ such that representation $T(r, s, k)$ gives a map of type $\{3, 4, 6, 4\}$ after identifying their boundaries.

Lemma 10.2. *The maps of type $\{3, 4, 6, 4\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $s \geq 2$ even, (ii) $3 \mid r$, (iii) number of vertices of $T(r, s, k) = rs \geq 18$, (iv) $r \geq 9$ if $s = 2$, (v) $r \geq 6$ if $s \geq 4$, (vi) $k \in \{3t + 4 : 0 \leq t \leq \frac{r-9}{3}\}$ if $s = 2$ & $k \in \{3t + 1 : 0 \leq t \leq \frac{r-3}{3}\}$ if $s \geq 4$.*

Proof. We follow similar argument of the proof of the Lemma 8.2. We proof this lemma by considering link of some vertices in $T(r, s, k)$ and by showing that link of those vertices are not cycle if we consider the values of r, s and k outside the given range in Lemma 10.2. So, consider map of type $\{3, 4, 6, 4\}$ in place of type $\{3^4, 6\}$ and different ranges of r, s and k in the proof of Lemma 8.2. Thus, we get all the cases of this lemma. This completes the proof. \square

Let M_i for $i = 1, 2$ be maps of type $\{3, 4, 6, 4\}$ on same number of vertices and $T_i = T(r_i, s_i, k_i)$ be (r_i, s_i, k_i) -representations of M_i . By Lemma 10.1, there are at most three non-homologous cycles of different lengths in T_i . So, let $c_{i,j} = \text{length}(N_{i,j})$ where $N_{i,j}$, $j = 1, 2, 3$ denote non-homologous cycles of type W_1 in T_i . Then,

Lemma 10.3. *The map $M_1 \cong M_2$ if and only if $(c_{1,1}, c_{1,2}, c_{1,3}) = (c_{2,t_1}, c_{2,t_2}, c_{2,t_3})$ for $t_1 \neq t_2 \neq t_3 \in \{1, 2, 3\}$.*

Proof. We proceed as in the case of proof of Lemma 8.3. Let $r = r_1, s = s_1, k = k_1$. We consider horizontal cycles of $T(r_i, s_i, k_i)$ of type W_1 . Proceed with similar argument as in the Lemma 4.9. Thus, we get $M_1 \cong M_2$. Again, if $(r, s, k) \neq (r_1, s_1, k_1)$ then we proceed as in the case of the proof of Lemma 5.6 and Lemma 6.3. Converse part follows from similar argument of the converse part of Lemma 4.9. This completes the proof. \square

As in Section 4, by Lemmas 10.2 and 10.3, the maps of type $\{3, 4, 6, 4\}$ can be classified on different number of vertices. We have done calculation for the vertices ≤ 54 . We have listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 7.

11 Maps of type $\{4, 8^2\}$

Let M be a semi-equivelar map of type $\{4, 8^2\}$ on the torus. We define a fixed type of path Z_1 in M . Let $Q(i) := P(u_i, u_{i+1}, u_{i+2}, u_{i+3}, u_{i+4})$ be a path in M , where $lk(u_i) = C(u_{i-1}, \mathbf{f}, \mathbf{g}, \mathbf{h}, \mathbf{i}, \mathbf{u}_{i+2}, u_{i+1}, \mathbf{s}, \mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{u}_{i-3}, \mathbf{u}_{i-2})$, $lk(u_{i+1}) = C(u_i, \mathbf{u}_{i-1}, \mathbf{f}, \mathbf{g}, \mathbf{h}, \mathbf{i}, u_{i+2}, \mathbf{u}_{i+3}, \mathbf{u}_{i+4}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{a})$, $lk(u_{i+2}) = C(u_{i+1}, \mathbf{u}_i, \mathbf{u}_{i-1}, \mathbf{f}, \mathbf{g}, \mathbf{h}, \mathbf{i}, \mathbf{j}, u_{i+3}, \mathbf{u}_{i+4}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s})$, $lk(u_{i+3}) = C(u_{i+2}, \mathbf{i}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{u}_{i+6}, \mathbf{u}_{i+5}, u_{i+4}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{u}_{i+1})$, $lk(u_{i+4}) = C(u_{i+3}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}, \mathbf{u}_{i+6}, u_{i+5}, \mathbf{o}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}, \mathbf{u}_{i+1}, \mathbf{u}_{i+2})$.

Definition 11.1. Let $R_1 := P(\dots, v_{i-1}, v_i, v_{i+1}, \dots)$ be a path in edge graph of M . We say R_1 of type Z_1 if $L_1 := P(v_t, v_{t+1}, v_{t+2}, v_{t+3}, v_{t+4})$ is a subpath of R_1 or in the extended path of R_1 then either $L_1 \mapsto Q(i)$ by $v_j \mapsto u_j$, $L_1 \mapsto Q(i+1)$ by $v_j \mapsto u_{j+1}$, $L_1 \mapsto Q(i+2)$ by $v_j \mapsto u_{j+2}$ or $L_1 \mapsto Q(i+3)$ by $v_j \mapsto u_{j+3}$ for $j \in \{t, t+1, t+2, t+3, t+4\}$.

Definition 11.2. Let $R_2 := P(\dots, x_{i-1}, x_i, x_{i+1}, \dots)$ be a path in edge graph of M . We say R_2 of type Z'_1 if $L_2 := P(x_t, x_{t+1}, x_{t+2}, x_{t+3}, x_{t+4})$ is a subpath of R_2 or in the extended path of R_2 then either $L_2 \mapsto Q(i)$ by $x_j \mapsto u_{2t+4-j}$, $L_2 \mapsto Q(i+1)$ by $x_j \mapsto u_{2t+4-j}$, $L_2 \mapsto Q(i+2)$ by $x_j \mapsto u_{2t+4-j}$ or $L_2 \mapsto Q(i+3)$ by $x_j \mapsto u_{2t+4-j}$ for $j \in \{t, t+1, t+2, t+3, t+4\}$.

As in Section 8, we consider path of type Z_1 as Z_1 and Z'_1 define same type of path (by the similar argument as in Section 8). Let P be a maximal path of type Z_1 . We argue similar argument of Lemma 4.1 for P . So, we get an edge e in M such that $P \cup e$ is a cycle of type Z_1 . Therefore, every maximal path of type Z_1 is a cycle. Let $C = P \cup e$. The cycle C is of type Z_1 and non-contractible (by the similar argument of Lemma 4.3). Let C_1, C_2, \dots, C_t be a sequence of homologous cycles of type Z_1 . Then, we proceed with the similar argument of Lemma 4.5. Thus, $\text{length}(C_i) = \text{length}(C_j)$ for $1 \leq i, j \leq t$. By Definition 11.1, there are two cycles of type Z_1 through each vertex of M . Let $v \in V(M)$ and $L_1(v), L_2(v)$ be two cycles through v . We repeat the similar construction of (r, s, k) -representation of Section 4 for M . So, we get a $T(r, s, k)$ representation of M . In this processes, we take first cut along L_1 and then, second cut along L_2 where the starting adjacent face to the base horizontal cycle L_1 is a 4-gon. By this construction, every map of type $\{4, 8^2\}$ on the torus has a $T(r, s, k)$ representation.

We have two cycles of type Z_1 through each vertex of M . Therefore, by Lemma 11.1, map M contains at most two non-homologous cycles of type Z_1 of different lengths.

Lemma 11.1. *The map M contains at most two non-homologous cycles of type Z_1 of different lengths.*

Proof. Let v be a vertex in M . By the definition of cycle of type Z_1 , we have two cycles namely, C_1 and C_2 through v . We proceed as in the case of proof of Lemma 8.1. So, we get that the map M contains at most two non-homologous cycles of type Z_1 of different lengths. \square

We claim in the Lemma 11.2 that map of type $\{4, 8^2\}$ does not contain cycle of type Z_1 which has length four. We use this result to classify the maps of type $\{4, 8^2\}$ on the torus.

Table 8: Maps of type $\{4, 8^2\}$

n	Equivalence classes	Length of cycles	$i(n)$
20	$T(20, 1, 6), T(20, 1, 14)$	(20, 20)	1(20)
24	$T(24, 1, 6), T(24, 1, 18)$	(24, 8)	2(24)
	$T(8, 3, 2), T(8, 3, 6)$		
	$T(24, 1, 14), T(24, 1, 10)$	(24, 24)	

Lemma 11.2. *The representation $T(r, s, k)$ does not contain cycle of type Z_1 of length four.*

Proof. Suppose, $T(r, s, k)$ has a cycle, say C of length four which is of type Z_1 . Let C_{F_8} denote a boundary cycle of a 8-gon F_8 . By the definition of cycle of type Z_1 , if $C \cap C_{F_8} \neq \emptyset$ then $C \cap C_{F_8}$ is a path of length three. So, let $C \cap C_{F_8} = P(u_1, u_2, u_3, u_4)$. If C is a cycle of length four then, $u_1 = u_4$ which is a contradiction as u_1, u_4 are in C_{F_8} and C_{F_8} is a cycle without chord. Again, by the definition of cycle of type Z_1 , C must intersect a 8-gon. Thus, $\text{length}(C) > 4$. So, a map M of type $\{4, 8^2\}$ does not contain a cycle C of type Z_1 of length four. This completes the proof. \square

We define admissible relations among r, s, k of $T(r, s, k)$.

Lemma 11.3. *The maps of type $\{4, 8^2\}$ of the form $T(r, s, k)$ exist if and only if the following holds : (i) $4 \mid r$ and $s \geq 1$, (ii) number of vertices of $T(r, s, k) = rs \geq 20$, (iii) $r \geq 20$ if $s = 1$, (iv) $r \geq 16$ if $s = 2$, (v) $r \geq 8$ if $s \geq 3$, (vi) $k \in \{4t + 6 : 0 \leq t \leq \frac{r-12}{4}\}$ if $s = 1$, $k \in \{4t + 7 : 0 \leq t \leq \frac{r-16}{4}\}$ if $s = 2$ & $k \in \{4t - 1(\text{mod } r) : 0 \leq t \leq \frac{r-4}{4}\}$ if $s \geq 3$.*

Proof. We proceed as in the case of proof of Lemma 8.2 and use Lemma 11.2 to prove this lemma. Considering link of some vertices in $T(r, s, k)$, map of type $\{4, 8^2\}$ in place of type $\{3^4, 6\}$, and different values of r, s and k in the proof of Lemma 8.2. So, we get all possible ranges of r, s and k of $T(r, s, k)$. This completes the proof. \square

Let M_i for $i = 1, 2$ be maps of type $\{4, 8^2\}$ on same number of vertices and $T_i = T(r_i, s_i, k_i)$ represent of M_i . By Lemma 11.1, there are at most two non-homologous cycles of different lengths in T_i . So, let $a_{i,j} = \text{length}(C_{i,j})$ where $C_{i,j}$, $j = 1, 2$ denote non-homologous cycles of type Z_1 in T_i . Then,

Lemma 11.4. *The map $M_1 \cong M_2$ if and only if $(a_{1,1}, a_{1,2}) = (a_{2,t_1}, a_{2,t_2})$ for $t_1 \neq t_2 \in \{1, 2\}$.*

Proof. We proceed as in the case of proof of Lemma 8.3. Let $r = r_1, s = s_1, k = k_1$. Consider horizontal cycles in $T(r_i, s_i, k_i)$ of type Z_1 . Proceed with similar argument as in Lemma 4.9. So, we get $M_1 \cong M_2$. Again, if $(r, s, k) \neq (r_1, s_1, k_1)$ then we proceed as in the case of the proof of Lemmas 5.6 and 6.3. Converse part follows from the similar argument of the converse part of Lemma 4.9. \square

As in Section 4, by Lemmas 11.3, 11.4, the maps of type $\{4, 8^2\}$ can be classified up to isomorphism on different number of vertices. We have done calculation for the vertices ≤ 24 . We have listed the obtained objects in the form of their $T(r, s, k)$ representation in Table 8.

12 Semi-equivelar maps

Proof of Theorem 1.1. The proof of the Theorem 1.1 follows from the Sections 4, 5, 6, 7, 8, 9, 10, 11. Let M be a map on n vertices of type $\{3^3, 4^2\}$ on the torus. We consider all admissible $T(r, s, k)$ representations of M by Lemma 4.8. We calculate length of the cycles of types A_1, A_2, A_3 and A_4 . We classify them by Lemma 4.9. In Table 1, we have classified upto 22 vertices. Similarly, we consider maps of types $\{3^2, 4, 3, 4\}, \{3, 6, 3, 6\}, \{3, 12^2\}, \{3^4, 6\}, \{4, 6, 12\}, \{3, 4, 6, 4\}, \{4, 8^2\}$ on the torus. That is, we consider cycles of type B_1 in maps of type $\{3^2, 4, 3, 4\}$ and classify by Lemma 5.6. Table 2 contains the classified maps upto 32 vertices. Consider cycles of type X_1 in maps of type $\{3, 6, 3, 6\}$ and classify by Lemma 6.3. Table 3 contains the maps upto 30 vertices. Consider cycles of type G_1 in maps of type $\{3, 12^2\}$ and classify by Lemma 7.3. Table 4 contains the maps upto 48 vertices. Consider cycles of type Y_1 in maps of type $\{3^4, 6\}$ and classify by Lemma 8.3. Table 5 contains the maps upto 42 vertices. Consider cycles of type H_1 in maps of type $\{4, 6, 12\}$ and classify by Lemma 9.3. Table 6 contains the maps upto 60 vertices. Consider cycles of type W_1 in maps of type $\{3, 4, 6, 4\}$ and classify by Lemma 10.3. Table 7 contains the classified maps upto 54 vertices. Consider cycles of type Z_1 in maps of type $\{4, 8^2\}$ and classify by Lemma 11.4. Table 8 contains the maps upto 24 vertices. This completes the proof. \square

13 Acknowledgement

The authors are grateful to the anonymous referee whose comments led to a substantial improvement in the paper. Work of second author is partially supported by SERB, DST grant No. SR/S4/MS:717/10.

References

- [1] A. Altshuler, Construction and enumeration of regular maps on the torus, *Discrete Math.* 4 (1973) 201-217.
- [2] U. Brehm and W. Kühnel, Equivelar maps on the torus, *European J. Combin.* 29 (2008) 1843-1861.
- [3] U. Brehm and E. Schulte, Polyhedral maps, *Handbook of Discrete and Computational Geometry* (eds. J. E. Goodman and J. O' Rourke), CRC Press NY, 1997, 345-358.
- [4] J. A. Bondy and U. S. R. Murthy, *Graph Theory*, GTM 244, Springer, 2008.
- [5] W. Kurth, Enumeration of Platonic maps on the torus, *Discrete Math.* 61 (1986) 71-83.
- [6] S. Negami, Uniqueness and faithfulness of embedding of toroidal graphs, *Discrete Math.* 44 (1983) 161-180.
- [7] A. K. Tiwari and A. K. Upadhyay, An Enumeration of Semi-Equivelar Maps on Torus and Klein bottle, *Math. Slovaca* (to appear).
- [8] A. K. Upadhyay, A. K. Tiwari and D. Maity, Semi-equivelar maps, *Beitr. Algebra Geom.* 55 (2014) 229-242.