A new optical field state as an output of diffusion channel when the input being number state

Hong-Yi Fan^{1†}[†] Sen-Yue Lou¹, Xiao-Yin Pan¹ and Li-Yun Hu^{2‡}

¹Department of Physics, Ningbo University, Ningbo 315211, P. R. China

²Department of physics, Jiangxi Normal University, Nanchang, 330022

June 30, 2021

Abstract

We theoretically propose a new optical field state

 $\rho_{new} = \lambda \left(1 - \lambda\right)^{l} : L_{k}\left(\frac{-\lambda^{2}a^{\dagger}a}{1 - \lambda}\right) e^{-\lambda a^{\dagger}a}:$

(here :: denotes normal ordeing symbol) which is named Laguerre-polynomial-weighted chaotic field. We show that such state can be implemented, i.e., when a number state enters into a diffusion channel, the output state is just this kind of states. We solve the master equation describing the diffusion process by using the summation method within ordered product of operators and the entangled state representation. The solution manifestly shows how a pure state evolves into a mixed state. The physical difference between the diffusion and the amplitude damping is pointed out.

1 Introduction

In quantum optics theory there are some typical states, e.g., number state, coherent state, and squeezed state, these are pure states; there are also some mixed states, the typical one is the chaotic state described by

$$\rho_c = (1 - e^{-\lambda}) \exp\left(-\lambda a^{\dagger} a\right), \tag{1}$$

where a and a^{\dagger} are photon annihilation and creation operators, obeying $[a, a^{\dagger}] = 1$, $tr\rho_c = 1$. The normally orderd form of ρ_c is $\rho_c = (1 - e^{-\lambda})$: exp $[(e^{-\lambda} - 1) a^{\dagger}a]$:, where the symbol : : denotes normal ordering symbol. In this work we shall report that there exists another important mixed state which appears in normally ordered form

$$\rho_{new} = \lambda \left(1 - \lambda\right)^l : L_l\left(\frac{-\lambda^2 a^{\dagger} a}{1 - \lambda}\right) e^{-\lambda a^{\dagger} a} : .$$
⁽²⁾

Here L_l is the *l*-th Laguerre polynomial, $tr\rho_{new} = 1$ (see Appendix 1). We show that this mixed state will appear experimently as it represents the output state of a diffusion process with the input state being a pure number state.

^{*}This work was supported by the National Natural Science Foundation of China (Grant Nos. 11175113 and 11264018), and the Young Talents Foundation of Jiangxi Normal University.

[†]Correspondence author

[‡]Correspondence author

When a pure state evolves into a mixed state the quantum decoherence happens. Decoherence is an important topic in quantum information processing. In nature, systems we concerned usually are surrounded by thermo reservoir, so some dissipative process or diffusion process naturally happen. An interesting question thus arises: when an input state for a diffusion channel is a number state $|l\rangle \langle l|, (|l\rangle = \frac{a^{\dagger l}}{\sqrt{l!}} |0\rangle)$, then how does it evolve with time? What kind of optical field will the output state be? The master equation describing the diffusion process is [1, 2]

$$\frac{d}{dt}\rho = -\kappa \left(a^{\dagger}a\rho - a\rho a^{\dagger} - a^{\dagger}\rho a + \rho a^{\dagger}a\right).$$
(3)

We shall first obtain $\rho(t)$ by deriving its infinite operator-sum form

$$\rho\left(t\right) = \sum_{i,j} M_{i,j} \rho_0 M_{i,j}^{\dagger},\tag{4}$$

where $M_{i,j}$ in general is named Kraus operator [3], whose concrete from will be derived for this diffusion problem, and then we examine how $\rho_0 = |l\rangle \langle l|$ evolves through the relation (4). We will employ the thermo entangled state representation and the technique of integration within an ordered product (IWOP) of operators [4-5] to realize our goal.

2 Solution of Eq. (3) obtained by entangled state representation and IWOP technique

We begin with introducing the thermo entangled state [6]

$$\left|\eta\right\rangle = \exp\left[-\frac{1}{2}\left|\eta\right|^{2} + \eta a^{\dagger} - \eta^{*}\tilde{a}^{\dagger} + a^{\dagger}\tilde{a}^{\dagger}\right]\left|0\tilde{0}\right\rangle,\tag{5}$$

where \tilde{a}^{\dagger} is a fictitious mode accompanying the real mode a^{\dagger} , $[\tilde{a}, \tilde{a}^{\dagger}] = 1$. $|\eta\rangle$ obeys the eigenvector equations

$$(a - \tilde{a}^{\dagger}) |\eta\rangle = \eta |\eta\rangle, \ (a^{\dagger} - \tilde{a}) |\eta\rangle = \eta^* |\eta\rangle, \tag{6}$$

$$\langle \eta | \left(a^{\dagger} - \tilde{a} \right) = \eta^* \langle \eta |, \quad \langle \eta | \left(a - \tilde{a}^{\dagger} \right) = \eta \langle \eta |.$$

$$\tag{7}$$

Using the normal ordering form of vacuum projector $|0\tilde{0}\rangle \langle 0\tilde{0}| =: e^{-a^{\dagger}a - \tilde{a}^{\dagger}\tilde{a}}:$, and the IWOP technique we can show the orthonormal and completeness relation

$$\langle \eta' | \eta \rangle = \pi \delta \left(\eta' - \eta \right) \delta \left(\eta'^* - \eta^* \right), \tag{8}$$

$$1 = \int \frac{d^2 \eta}{\pi} |\eta\rangle \langle \eta|$$

$$= \int \frac{d^2 \eta}{\pi} : \exp\left[-|\eta|^2 + \eta a^{\dagger} - \eta^* \tilde{a}^{\dagger} + \eta^* a - \eta \tilde{a} + a^{\dagger} \tilde{a}^{\dagger} + a \tilde{a} - a^{\dagger} a - \tilde{a}^{\dagger} \tilde{a}\right] := 1.$$
(9)

Let

$$\left|\eta = 0\right\rangle = e^{a^{\dagger}\tilde{a}^{\dagger}} \left|0\tilde{0}\right\rangle \equiv \left|I\right\rangle,\tag{10}$$

we have

$$a |I\rangle = \tilde{a}^{\dagger} |I\rangle, \ a^{\dagger} |I\rangle = \tilde{a} |I\rangle, \ (a^{\dagger}a)^{n} |I\rangle = (\tilde{a}^{\dagger}\tilde{a})^{n} |I\rangle.$$

$$(11)$$

Operating the two-sides of (3) on $|I\rangle$, noting that the real field ρ is independent of the fictitious mode, $[\rho, \tilde{a}] = 0$, $[\rho, \tilde{a}^{\dagger}] = 0$, and using (12) we have

$$\frac{d}{dt}\rho\left|I\right\rangle = -\kappa\left(a^{\dagger}a\rho - a\tilde{a}\rho - a^{\dagger}\tilde{a}^{\dagger}\rho + \tilde{a}\tilde{a}^{\dagger}\rho\right)\left|I\right\rangle.$$
(12)

Letting $\rho \left| I \right\rangle \equiv \left| \rho \right\rangle$, we see

$$\frac{d}{dt}\left|\rho\right\rangle = -\kappa\left(a^{\dagger} - \tilde{a}\right)\left(a - \tilde{a}^{\dagger}\right)\left|\rho\right\rangle,\tag{13}$$

its formal solution is

$$|\rho\rangle = \exp\left[-\kappa t \left(a^{\dagger} - \tilde{a}\right) \left(a - \tilde{a}^{\dagger}\right)\right] |\rho_{0}\rangle, \qquad (14)$$

where $|\rho_0\rangle = \rho_0 |I\rangle$. Projecting this equation onto the entanged state representation $\langle \eta |$ and using the eigenvalue equation (7) we have

$$\langle \eta | \rho \rangle = \langle \eta | \exp \left[-\kappa t \left(a^{\dagger} - \tilde{a} \right) \left(a - \tilde{a}^{\dagger} \right) \right] | \rho_0 \rangle = e^{-\kappa t |\eta|^2} \langle \eta | \rho_0 \rangle.$$
⁽¹⁵⁾

Multiplying the two-sides of (15) by $\int \frac{d^2\eta}{\pi} |\eta\rangle$ and using the completeness relation (9) as well as the IWOP technique we obtain

$$\begin{aligned} |\rho\rangle &= \int \frac{d^2\eta}{\pi} e^{-\kappa t |\eta|^2} |\eta\rangle \langle \eta |\rho_0\rangle \\ &= \int \frac{d^2\eta}{\pi} : e^{-(\kappa t+1)|\eta|^2 + \eta a^{\dagger} - \eta^* \tilde{a}^{\dagger} + \eta^* a - \eta \tilde{a} + a^{\dagger} \tilde{a}^{\dagger} + a \tilde{a} - a^{\dagger} a - \tilde{a}^{\dagger} \tilde{a}} : |\rho_0\rangle \end{aligned} \tag{16} \\ &= \frac{1}{1+\kappa t} : \exp\left[\frac{\kappa t}{1+\kappa t} \left(a^{\dagger} \tilde{a}^{\dagger} + a \tilde{a} - a^{\dagger} a - \tilde{a}^{\dagger} \tilde{a}\right)\right] : |\rho_0\rangle \\ &= \frac{1}{1+\kappa t} e^{\frac{\kappa t}{1+\kappa t} a^{\dagger} \tilde{a}^{\dagger}} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger} a + \tilde{a}^{\dagger} \tilde{a}} e^{\frac{\kappa t}{1+\kappa t} a \tilde{a}} \rho_0 |I\rangle , \end{aligned}$$

where we have noticed

$$: \exp\left[\frac{-\kappa t}{1+\kappa t} \left(a^{\dagger}a+\tilde{a}^{\dagger}\tilde{a}\right)\right] := \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger}a+\tilde{a}^{\dagger}\tilde{a}}$$
(17)

Using $[\tilde{a}, \rho_0] = 0$, $\tilde{a} \left| I \right\rangle = a^{\dagger} \left| I \right\rangle$ we have

$$e^{\frac{\kappa t}{1+\kappa t}a\tilde{a}}\rho_{0}\left|I\right\rangle = \sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{\kappa t}{1+\kappa t}a\right)^{n}\rho_{0}\tilde{a}^{n}\left|I\right\rangle$$
$$= \sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{\kappa t}{1+\kappa t}\right)^{n}a^{n}\rho_{0}a^{\dagger n}\left|I\right\rangle, \tag{18}$$

After substituting (18) into (16) and then using the property that $\tilde{a}^{\dagger}\tilde{a}$ is commutable with all real field operators and $f(a^{\dagger}a)|I\rangle = f(\tilde{a}^{\dagger}\tilde{a})|I\rangle$, we can put Eq.(16) into the following form

$$\begin{aligned} |\rho\rangle &= \frac{1}{1+\kappa t} e^{\frac{\kappa t}{1+\kappa t}a^{\dagger}\tilde{a}^{\dagger}} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger}a+\tilde{a}^{\dagger}\tilde{a}} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{\kappa t}{1+\kappa t}\right)^{n} a^{n} \rho_{0} a^{\dagger n} |I\rangle \\ &= \frac{1}{1+\kappa t} e^{\frac{\kappa t}{1+\kappa t}a^{\dagger}\tilde{a}^{\dagger}} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger}a} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{\kappa t}{1+\kappa t}\right)^{n} a^{n} \rho_{0} a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{\tilde{a}^{\dagger}\tilde{a}} |I\rangle \\ &= \frac{1}{1+\kappa t} \sum_{m=0}^{\infty} \frac{1}{m!} \left(\frac{\kappa t}{1+\kappa t}\right)^{m} a^{\dagger m} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger}a} \\ &\times \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{\kappa t}{1+\kappa t}\right)^{n} a^{n} \rho_{0} a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger}a} \tilde{a}^{\dagger m} |I\rangle. \end{aligned}$$
(19)

Finally, using $\tilde{a}^{\dagger m} \left| I \right\rangle = a^m \left| I \right\rangle$ we obtain

$$\rho(t) |I\rangle = \sum_{m,n=0}^{\infty} \frac{(\kappa t)^{m+n}}{m!n! (\kappa t+1)^{m+n+1}} a^{\dagger m} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger} a} \\
\times a^{n} \rho_{0} a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger} a} a^{m} |I\rangle.$$
(20)

It then follows the infinite sum form

$$\rho(t) = \sum_{m,n=0}^{\infty} \frac{(\kappa t)^{m+n}}{m!n! (\kappa t+1)^{m+n+1}} a^{\dagger m} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger a}} a^n \rho_0 a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger a}} a^m \qquad (21)$$

$$\equiv \sum_{m,n=0}^{\infty} M_{m,n} \rho_0 M_{m,n}^{\dagger}$$

where

$$M_{m,n} = \sqrt{\frac{1}{m!n!} \frac{(\kappa t)^{m+n}}{(\kappa t+1)^{m+n+1}}} a^{\dagger m} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger} a} a^{n},$$
(22)

satisfying $\sum_{m,n=0}^{\infty} M_{m,n}^{\dagger} M_{m,n} = 1$, which is trace conservative (see Appendix 2). Thus we have employed the entangled state representation to analytically derive the infinitive sum form of $\rho(t)$.

3 Diffusion of a number state

We now consider the case that a number state undergoes the diffusion channel, i.e., let ρ_0 in Eq. (21) be $|l\rangle \langle l|$ and we begin with considering the part of summation over n in Eq. (21)

$$\Im = \sum_{n=0}^{l} \frac{(\kappa t)^{n}}{n! (\kappa t+1)^{n}} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger} a} a^{n} |l\rangle \langle l| a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger} a} \\ = \sum_{n=0}^{l} \frac{1}{n!} \frac{[\kappa t (\kappa t+1)]^{n}}{(\kappa t+1)^{2l}} \frac{l!}{[(l-n)!]^{2}} (a^{\dagger})^{l-n} |0\rangle \langle 0| a^{l-n}.$$
(23)

Using the definition of the two-variable Hermite polynomials

$$H_{m,n}(x,y) = \sum_{l=0}^{\min(m,n)} \frac{m!n!(-1)^l}{l!(m-l)!(n-l)!} x^{m-l} y^{n-l},$$
(24)

and $\left|0\right\rangle \left\langle 0\right|=:e^{-a^{\dagger}a}\colon,$ we see

$$\Im = \frac{1}{l!} \left(\frac{-\kappa t}{\kappa t + 1} \right)^{l} : H_{l,l} \left(\frac{ia^{\dagger}}{\sqrt{\kappa t (\kappa t + 1)}}, \frac{ia}{\sqrt{\kappa t (\kappa t + 1)}} \right) e^{-a^{\dagger}a} : ,$$
(25)

then inserting (25) into (21) and using the summation method within ordered product of operators yields

$$\rho(t) = \frac{1}{l!} \left(\frac{-\kappa t}{\kappa t+1}\right)^l \sum_{m=0}^{\infty} \frac{1}{m!} \frac{(\kappa t)^m}{(\kappa t+1)^{m+1}} \\
\times : a^{\dagger m} a^m H_{l,l} \left[\frac{ia^{\dagger}}{\sqrt{\kappa t (\kappa t+1)}}, \frac{ia}{\sqrt{\kappa t (\kappa t+1)}}\right] e^{-a^{\dagger}a} : \\
= \frac{(-\kappa t)^l}{l! (\kappa t+1)^{l+1}} : e^{\frac{-1}{\kappa t+1}a^{\dagger}a} H_{l,l} \left[\frac{ia^{\dagger}}{\sqrt{\kappa t (\kappa t+1)}}, \frac{ia}{\sqrt{\kappa t (\kappa t+1)}}\right] : .$$
(26)

Using the definition of Laguerre-polynomial

$$L_l(x) = \sum {\binom{l}{l-k} \frac{(-x)^k}{k!}}$$
(27)

and

$$L_{l}(xy) = \frac{(-1)^{l}}{l!} H_{l,l}(x,y), \qquad (28)$$

then (26) becomes

$$\rho(t) = \frac{(\kappa t)^l}{(\kappa t+1)^{l+1}} \colon L_l\left(\frac{-a^{\dagger}a}{\kappa t(\kappa t+1)}\right) e^{\frac{-1}{\kappa t+1}a^{\dagger}a} \colon .$$
(29)

Noting $e^{\frac{-1}{\kappa t+1}a^{\dagger}a}$ represents a chaotic photon field, so $\rho(t)$ is a Laguerre-polynomial-weighted chaotic field. Thus we see $|l\rangle \langle l|$ evolves into the mixed state (29), so this diffusion process manifestly embodies quantum decoherence.

As Eq. (29) is just in the type of Eq. (2), so we can confirm the state described by Eq. (2) indeed exists as an quantum optical field.

Before we check $Tr\rho(t) = 1$ for Eq. (29), let us present an integration formula

$$\int \frac{d^2 \alpha}{\pi} e^{\lambda |\alpha|^2} |\alpha|^{2k} = \left(\frac{\partial}{\partial \lambda}\right)^k \int \frac{d^2 \alpha}{\pi} e^{\lambda |\alpha|^2} = k! \left(\frac{-1}{\lambda}\right)^{k+1},\tag{30}$$

then we introduce the completeness relation of coherent state

$$\int \frac{d^2 \alpha}{\pi} \left| \alpha \right\rangle \left\langle \alpha \right| = 1 \tag{31}$$

here $|\alpha\rangle = \exp\left[\alpha a^{\dagger} - |\alpha|^2/2\right]|0\rangle$. Due to

$$a |\alpha\rangle = \alpha |\alpha\rangle, \langle \alpha| : f(a^{\dagger}, a) : |\alpha\rangle = f(\alpha^*, \alpha), \qquad (32)$$

we see

$$\langle \alpha | : L_l \left(\frac{-a^{\dagger}a}{\kappa t (\kappa t + 1)} \right) e^{\frac{-1}{\kappa t + 1}a^{\dagger}a} : |\alpha\rangle$$

$$= e^{\frac{-1}{\kappa t + 1}|\alpha|^2} \sum_{k=0}^l \frac{l!(-1)^k}{k! \left[(l-k)!\right]^2} \left(\frac{-|\alpha|^2}{\kappa t (\kappa t + 1)} \right)^{l-k}.$$
(33)

Substituting (33) into $tr\rho(t) = \int \frac{d^{2}\alpha}{\pi} \langle \alpha | \rho(t) | \alpha \rangle$ we should calculate

$$tr\rho(t) = \int \frac{d^2\alpha}{\pi} \langle \alpha | \rho(t) | \alpha \rangle$$

$$= \int \frac{d^2\alpha}{\pi} \langle \alpha | \frac{(\kappa t)^l}{(\kappa t+1)^{l+1}} : e^{\frac{-1}{\kappa t+1}a^{\dagger}a} L_l\left(\frac{-a^{\dagger}a}{\kappa t(\kappa t+1)}\right) : |\alpha\rangle$$

$$= \frac{(\kappa t)^l}{(\kappa t+1)^{l+1}} \sum_{k=0}^l \frac{l!}{k! \left[(l-k)!\right]^2} \int \frac{d^2\alpha}{\pi} e^{\frac{-1}{\kappa t+1}|\alpha|^2} \left(\frac{|\alpha|^2}{\kappa t(\kappa t+1)}\right)^{l-k}.$$
 (34)

By setting $\frac{\alpha}{\sqrt{\kappa t(\kappa t+1)}} = \alpha'$ we reform the integration as

$$\int \frac{d^2 \alpha}{\pi} e^{\frac{-1}{\kappa t+1}|\alpha|^2} \left(\frac{|\alpha|^2}{\kappa t (\kappa t+1)}\right)^{l-k}$$

$$= \kappa t (\kappa t+1) \int \frac{d^2 \alpha'}{\pi} e^{-\kappa t |\alpha'|^2} (|\alpha'|^2)^{l-k}$$

$$= \kappa t (\kappa t+1) (l-k)! \left(\frac{1}{\kappa t}\right)^{l-k+1}.$$
(35)

Substituting (35) into (34) we see

$$tr\rho(t) = \frac{(\kappa t)^{l+1}}{(\kappa t+1)^l} \sum_{k=0}^l \frac{l!}{k!(l-k)!} \left(\frac{1}{\kappa t}\right)^{l-k+1} = 1$$
(36)

so it is trace conservative.

4 The photon number in the mixed state

Then we evaluate the photon number for Eq. (29)

$$Tr\left[\rho\left(t\right)a^{\dagger}a\right] = Tr\left[\rho\left(t\right)aa^{\dagger}\right] - 1$$
$$= \frac{\left(\kappa t\right)^{l}}{\left(\kappa t+1\right)^{l+1}}Tr\left[:a^{\dagger}ae^{\frac{-1}{\kappa t+1}a^{\dagger}a}L_{l}\left(\frac{-a^{\dagger}a}{\kappa t\left(\kappa t+1\right)}\right):\right] - 1.$$
(37)

By using the coherent state representation we have

$$Tr\left[:a^{\dagger}ae^{\frac{-1}{\kappa t+1}a^{\dagger}a}L_{l}\left(\frac{-a^{\dagger}a}{\kappa t(\kappa t+1)}\right):\right]$$

$$= \int \frac{d^{2}\alpha}{\pi}e^{\frac{-1}{\kappa t+1}|\alpha|^{2}}|\alpha|^{2}L_{l}\left(\frac{-|\alpha|^{2}}{\kappa t(\kappa t+1)}\right)$$

$$= [\kappa t(\kappa t+1)]^{2}\int \frac{d^{2}\alpha'}{\pi}e^{-\kappa t|\alpha'|^{2}}L_{l}(-|\alpha'|^{2})|\alpha'|^{2}$$

$$= [\kappa t(\kappa t+1)]^{2}\sum_{k=0}^{l}\frac{l!}{k!k!(l-k)!}\int \frac{d^{2}\alpha'}{\pi}e^{-\kappa t|\alpha'|^{2}}(|\alpha'|^{2})^{k+1}$$

$$= (\kappa t+1)^{2}\sum_{k=0}^{l}\frac{l!}{k!(l-k)!}(k+1)\left(\frac{1}{\kappa t}\right)^{k},$$
(38)

where

$$\sum_{k=0}^{l} \frac{l!}{k!(l-k)!} k\left(\frac{1}{\kappa t}\right)^{k} = \frac{l}{\kappa t} \sum_{k=1}^{l} \frac{(l-1)!}{(k-1)!(l-k)!} \left(\frac{1}{\kappa t}\right)^{k-1} = \frac{l}{\kappa t} \left(\frac{\kappa t+1}{\kappa t}\right)^{l-1},$$
(39)

and

$$\sum_{k=0}^{l} \frac{l!}{k!(l-k)!} \left(\frac{1}{\kappa t}\right)^{k} = \left(\frac{\kappa t+1}{\kappa t}\right)^{l}.$$
(40)

Substituting (39)-(40) into (38) we obtain

$$Tr\left[:a^{\dagger}ae^{\frac{-1}{\kappa t+1}a^{\dagger}a}L_{l}\left(\frac{-a^{\dagger}a}{\kappa t(\kappa t+1)}\right):\right]$$
$$= (\kappa t+1)^{2}\left(\frac{\kappa t+1}{\kappa t}\right)^{l-1}\frac{l+\kappa t+1}{\kappa t}.$$
(41)

Then substituting (41) into (37) we see

$$Tr\left[\rho\left(t\right)a^{\dagger}a\right] = l + \kappa t. \tag{42}$$

which tells that the photon number $l \rightarrow l + \kappa t$.

At the end of this work we point out that a diffusion process is quite different from the process in the amplitude dissipative channel (ADC) described by the following master equation [7]

$$\frac{d}{dt}\rho' = \gamma \left(2a\rho'a^{\dagger} - a^{\dagger}a\rho' - \rho'a^{\dagger}a\right)$$
(43)

where γ is the rate of dissipation. The solution to Eq. (43) is [8]

$$\rho' = \sum_{m=0}^{\infty} \frac{\left(1 - e^{-2\gamma t}\right)^n}{n!} e^{-\gamma t a^{\dagger} a} a^n \rho'_0 a^{\dagger n} e^{-\gamma t a^{\dagger} a}.$$
(44)

In ADC an initial pure number state $|l\rangle \langle l|$ will evolve into a binomial state as shown in [9]

$$\sum_{m=0}^{l} {l \choose l-m} e^{-2\gamma m t} \left(1 - e^{-2\gamma t}\right)^{l-m} |m\rangle \langle m| \equiv \rho_b'$$
(45)

with photon number decaying $tr(a^{\dagger}a\rho'_b) = le^{-2\gamma t}$. Comparing the diffusion master equation (3) with the dissipation Eq. (43) we realize that the term $a^{\dagger}\rho a$ may be responsible for diffusion.

In summary, we theoretically propose a new optical field state

$$\rho_{new} = \lambda \left(1 - \lambda\right)^l : L_k \left(\frac{-\lambda^2 a^{\dagger} a}{1 - \lambda}\right) e^{-\lambda a^{\dagger} a} :$$
(46)

which is named Laguerre-polynomial-weighted chaotic field. We show that such state can be implemented, i.e., when a number state enters into a diffusion channel, the output state is just this kind of states. We solve the master equation describing the diffusion process by using the summation method within ordered product of operators and the entangled state representation. The solution manifestly shows how a pure state evolves into a mixed state. The physical difference between the diffusion and the amplitude damping is pointed out.

Acknowledgements: This work was supported by the National Natural Science Foundation of China (Grant Nos. 11175113 and 11264018), and the Natural Science Foundation of Jiangxi Province of China (No 20132BAB212006).

Appendix 1

For ρ_{new} in Eq. (2) we prove $tr\rho_{new} = 1$. In fact, using the coherent state representation we have

$$tr\rho_{new} = \int \frac{d^2\alpha}{\pi} \langle \alpha | \lambda (1-\lambda)^l : L_l \left(\frac{-\lambda^2 a^{\dagger} a}{1-\lambda}\right) e^{-\lambda a^{\dagger} a} : |\alpha\rangle$$
$$= \lambda (1-\lambda)^l \int \frac{d^2\alpha}{\pi} e^{-\lambda |\alpha|^2} L_l \left(\frac{-\lambda^2 |\alpha|^2}{1-\lambda}\right)$$
$$= \lambda (1-\lambda)^l \int_0^\infty d\left(\frac{\lambda-1}{\lambda^2} x\right) e^{-\frac{\lambda-1}{\lambda} x} L_l (x) = 1,$$
(A1)

where we have used

$$\int_{0}^{\infty} e^{-bx} L_{l}(x) = (b-1)^{l} b^{-l-1}.$$
(A2)

Appendix 2

For $M_{m,n}$ in Eq. (22) we now prove $\sum_{m,n=0}^{\infty} M_{m,n}^{\dagger} M_{m,n} = 1$. Because

$$:e^{xaa^{\dagger}}:=\frac{1}{1-x}e^{a^{\dagger}a\ln\frac{1}{1-x}},$$
 (A3)

where : : denotes anti-normal ordering, we have

$$\sum_{m,n=0}^{\infty} \frac{1}{m!} \frac{(\kappa t)^m}{(\kappa t+1)^m} a^m a^{\dagger m}$$

= $\vdots \exp\left[\frac{\kappa t}{\kappa t+1} a a^{\dagger}\right] \vdots$
= $(\kappa t+1) e^{a^{\dagger a} \ln(\kappa t+1)}.$ (A4)

Substituting it into the sum representation of $\rho(t)$ yields

$$\sum_{m,n=0}^{\infty} M_{m,n}^{\dagger} M_{m,n} = \sum_{m,n=0}^{\infty} \frac{1}{m!n!} \frac{(\kappa t)^{m+n}}{(\kappa t+1)^{m+n+1}} a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger a}} a^{m} a^{\dagger m} \left(\frac{1}{1+\kappa t}\right)^{a^{\dagger a}} a^{n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{(\kappa t)^{n}}{(\kappa t+1)^{n}} a^{\dagger n} \left(\frac{1}{1+\kappa t}\right)^{2a^{\dagger a}} e^{a^{\dagger a} \ln(\kappa t+1)} a^{n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{(\kappa t)^{n}}{(\kappa t+1)^{n}} a^{\dagger n} e^{a^{\dagger a} \ln(\kappa t+1)} e^{2a^{\dagger a} \ln \frac{1}{1+\kappa t}} a^{n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{(\kappa t)^{n}}{(\kappa t+1)^{n}} a^{\dagger n} e^{a^{\dagger a} \ln (\kappa t+1)+2\ln \frac{1}{1+\kappa t}} a^{n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{(\kappa t)^{n}}{(\kappa t+1)^{n}} a^{\dagger n} e^{a^{\dagger a} \ln \frac{1}{1+\kappa t}} a^{n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{(\kappa t)^{n}}{(\kappa t+1)^{n}} a^{\dagger n} e^{a^{\dagger a} \ln \frac{1}{1+\kappa t}} a^{n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{(\kappa t)^{n}}{(\kappa t+1)^{n}} a^{\dagger n} = e^{a^{\dagger a} (\frac{1}{1+\kappa t}-1)} : a^{n}$$

$$= : e^{a^{\dagger a} \frac{\kappa t}{1+\kappa t}} e^{a^{\dagger a} \ln \frac{-\kappa t}{1+\kappa t}} : = 1.$$
(A5)

References

- Carmichael H J 1999 Statistical Methods in Ouantum Optics I, Master Equation and Fokker-Planck equations, Springer-Verlag, Berlin
- [2] Orszag M 2000 Quantum Optics, Springer-Verlag, Berlin.
- [3] Preskill J, 1998 Lecture Notes for Physics 229: Quantum Information and Computation, California Institution of Technology.
- [4] Fan H Y, Lu H L, Fan Y 2006 Ann. Phys. 321, 480.
- [5] Fan H Y 2003 J. Opt. B: Quantum Semiclass. Opt. 5 R147
- [6] Fan H Y, Klauder J R. 1994 Phys. Rev. A, 49 704
- [7] Gardiner C and Zoller P 2000 Quantum Noise (Springer Berlin).
- [8] Fan H Y and Hu L Y 2009 Chin. Phys. B 18 1061
- [9] FAN Hong-Yi, REN Gang, Chin. Phys. Lett.. 2010, 27 (5): 050302