Recent progresses on the pseudospin symmetry in single particle resonant states

Bing-Nan Lu¹, En-Guang Zhao^{1,2} and Shan-Gui Zhou^{1,2}

¹ State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

 2 Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, China

E-mail: sgzhou@itp.ac.cn (Shan-Gui Zhou)

Abstract. The pseudospin symmetry (PSS) is a relativistic dynamical symmetry directly connected with the small component of the nucleon Dirac wave function. Much effort has been made to study this symmetry in bound states. Recently, a rigorous justification of the PSS in single particle resonant states was achieved by examining the asymptotic behaviors of the radial Dirac wave functions: The PSS in single particle resonant states in nuclei is conserved exactly when the attractive scalar and repulsive vector potentials have the same magnitude but opposite sign. Several issues related to the exact conservation and breaking mechanism of the PSS in single particle resonances were investigated by employing spherical square well potentials in which the PSS breaking part can be well isolated in the Jost function. A threshold effect in the energy splitting and an anomaly in the width splitting of pseudospin partners were found when the depth of the square well potential varies from zero to a finite value.

1. Introduction

In 1969, the pseudospin symmetry (PSS) was found in atomic nuclei: Doublets of single particle levels with quantum numbers $(n_r, l, j = l + 1/2)$ and $(n_r - 1, l + 2, j = l + 3/2)$ in the same major shell are nearly degenerate [1, 2]. Since the PSS was observed, much effort has been devoted to investigate its origin [3, 4, 5, 6, 7]. Finally, it was revealed that the PSS in nuclei is a relativistic symmetry which is exactly conserved when the scalar potential S(r) and the vector potential V(r) have the same size but opposite sign, i.e., $\Sigma(r) \equiv S(r) + V(r) = 0$ [8], or, more generally, when $d\Sigma(r)/dr = 0$ [9, 10]. The PSS is usually viewed as a dynamical symmetry [11, 12] because, in either limit, $\Sigma(r) = 0$ or $d\Sigma(r)/dr = 0$, there are no bound states in realistic nuclei, thus in realistic nuclei the PSS is always broken.

The SS and PSS have been investigated extensively. The readers are referred to Ref. [13] for a review and to Refs. [14, 15] for overviews of recent progresses. Here we simply mention that the study of the PSS has been extended along several lines:

- (i) From single particle states in the Fermi sea to those in the Dirac sea, i.e., the spin symmetry in anti-nucleon spectra [16, 17, 18, 19].
- (ii) From normal nuclei to hypernuclei [20, 21, 22].
- (iii) From spherical systems to deformed ones [23, 24, 25, 26].
- (iv) From bound states to resonant states; next we will focus on this extension.

There has been a great interest in the exploration of continuum and resonant states in atomic nuclei [27, 28, 29]. The study of symmetries, e.g., the PSS, in resonant states is very interesting. There have been some numerical investigations of the PSS in single particle resonances [30, 31, 32, 33, 34] and the SS in single particle resonant states [35]. Recently, we gave a rigorous justification of the PSS in single particle resonant states and investigated several open problems related to the exact conservation and breaking mechanism of the PSS in single particle resonances [36, 37, 15]. In this contribution, we will present some results concerning the PSS in single particle resonant states.

2. Justification of the PSS in resonant states

In atomic nuclei, the equation of motion for nucleons, the Dirac equation, reads,

$$[\boldsymbol{\alpha} \cdot \boldsymbol{p} + \beta(M + S(\boldsymbol{r})) + V(\boldsymbol{r})] \psi(\boldsymbol{r}) = \epsilon \psi(\boldsymbol{r}), \qquad (1)$$

where $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ are Dirac matrices, M is the nucleon mass, and $S(\boldsymbol{r})$ and $V(\boldsymbol{r})$ the scalar and vector potentials.

For a spherical nucleus, the upper and lower components of the Dirac spinor are $iF_{n\kappa}(r)/r \cdot Y_{jm}^{l}(\theta,\phi)$ and $-G_{\tilde{n}\kappa}(r)/r \cdot Y_{jm}^{\tilde{l}}(\theta,\phi)$, respectively. One can derive a Schrödinger-like equation for the lower component,

$$\left[\frac{d^2}{dr^2} - \frac{1}{M_-(r)}\frac{d\Sigma(r)}{dr}\frac{d}{dr} - \frac{\tilde{l}(\tilde{l}+1)}{r^2} + \frac{1}{M_-(r)}\frac{\kappa}{r}\frac{d\Sigma(r)}{dr} - M_+(r)M_-(r)\right]G(r) = 0, \quad (2)$$

where $M_+(r) = M + \epsilon - \Delta(r)$ and $M_-(r) = M - \epsilon + \Sigma(r)$.

For the continuum in the Fermi sea, $\epsilon \ge M$; the regular solution for Eq. (2) vanishes at the origin and at large r,

$$G(r) = \frac{i}{2} \left[\mathcal{J}_{\kappa}^{G}(p) h_{\tilde{l}}^{-}(pr) - \mathcal{J}_{\kappa}^{G}(p)^{*} h_{\tilde{l}}^{+}(pr) \right], \quad r \to \infty,$$
(3)

where $h_{\tilde{l}}^{\pm}(pr)$ are the Ricatti-Hankel functions. $\mathcal{J}_{\kappa}(p)$ is the Jost function which is an analytic function of p. The zeros of $\mathcal{J}_{\kappa}^{G}(p)$ on the lower p plane and near the real axis correspond to resonant states. The resonance energy E and width Γ are determined by the relation $E - i\Gamma/2 = \epsilon - M$.

In the PSS limit, Eq. (2) is reduced as

$$\left[\frac{d^2}{dr^2} - \frac{\tilde{l}(\tilde{l}+1)}{r^2} + (\epsilon - M)M_+(r)\right]G(r) = 0.$$
(4)

For pseudospin doublets with κ and $\kappa' = -\kappa + 1$, the small components satisfy the same equation because they have the same pseudo orbital angular momentum \tilde{l} . For continuum states, $G_{\kappa}(\epsilon, r) = G_{\kappa'}(\epsilon, r)$ is true for any energy ϵ thus we have $\mathcal{J}_{\kappa'}^G(p) = \mathcal{J}_{\kappa}^G(p)$. Thus the zeros are the same for $\mathcal{J}_{\kappa'}^G(p)$ and $\mathcal{J}_{\kappa}^G(p)$ and the PSS in single particle resonant states in nuclei is exactly conserved [36].

3. Numerical Results and Discussions

Spherical square well potentials for $\Sigma(r)$ and $\Delta(r)$ read,

$$\Sigma(r) = \begin{cases} C, & r < R, \\ 0, & r \ge R, \end{cases} \qquad \Delta(r) = \begin{cases} D, & r < R, \\ 0, & r \ge R, \end{cases}$$
(5)



Figure 1. (Color online) The zeros of the Jost function \mathcal{J}_{κ}^{G} on the complex energy plane in square-well potentials (5) with C = 0 (solid symbols) and C = -66 MeV (half-filled symbols) for PS partners $h_{11/2}$ (diamond) and $j_{13/2}$ (square).



Figure 2. (Color online). The energy (the left panel) and width (the right panel) splitting between PSS partners with the pseudo orbital angular momentum $\tilde{l} = 2$, $p_{3/2}$ and $f_{5/2}$, as a function of the potential depth C. Taken from Ref. [15].

where C and D are depths and R is the width. The Jost function $\mathcal{J}^{G}_{\kappa}(p)$ is derived as [36],

$$\mathcal{J}^{G}_{\kappa}(p) = -\frac{p^{\tilde{l}}}{2ik^{\tilde{l}+1}} \left\{ j_{\tilde{l}}(kR)ph_{\tilde{l}}^{+\prime}(pR) - kj_{\tilde{l}}^{\prime}(kR)h_{\tilde{l}}^{+}(pR) - b\left[kj_{\tilde{l}}^{\prime}(kR) - \frac{\kappa}{R}j_{\tilde{l}}(kR)\right]h_{\tilde{l}}^{+}(pR)\right\}, \quad (6)$$

with $b = C/(\epsilon - M - C)$ and $k = \sqrt{(\epsilon - C - M)(\epsilon - D + M)}$. By examining the zeros of the Jost function (6), one gets single particle bound states and resonant states.

Figure 1 shows solutions for PSS doublets with l = 6, i.e., $h_{11/2}$ with $\kappa = -6$ and $j_{13/2}$ with $\kappa' = 7$ for square-well potentials with D = 650 MeV and R = 7 fm. The conservation of the PSS for single particle resonant states is seen: When C = 0, the roots for $h_{11/2}$ and $j_{13/2}$ are the same. When C = -66 MeV, the PSS is broken.

By examining the Jost function, one can trace continuously the PSS partners from the PSS limit to the case with a finite potential depth. This has been done in Ref. [15] in which we found a threshold effect in the energy splitting and an anomaly in the width splitting of pseudospin partners when C varies from zero to a finite value. As the depth of the single particle potential becomes deeper, the PSS is broken more and a threshold effect in the energy splitting first increases then decreases

until the pseudospin doublets encounter the threshold where one of the levels becomes a bound state and the splitting takes a minimum value; When the potential becomes even deeper, the splitting increases again. When the depth of the single particle potential increases from zero, there also appears an anomaly in the width splitting of PS partners which is shown in Fig. 2: It first decreases from zero to a maximum value with negative sign, then increases and becomes zero again; after the inversion of the width splitting, the splitting increases and reaches a maximum and positive value, then it becomes smaller and finally reaches zero.

Finally we mention that there are still many open problems concerning the exact and breaking of the PSS in single particle resonant states; the readers are referred to Refs. [15, 37] for further discussions.

Acknowledgments

This work has been supported by Major State Basic Research Development Program of China (Grant No. 2013CB834400), National Natural Science Foundation of China (Grant Nos. 11121403, 11175252, 11120101005, 11211120152, and 11275248), the Knowledge Innovation Project of Chinese Academy of Sciences (Grant No. KJCX2-EW-N01).

References

- [1] Arima A, Harvey M and Shimizu K 1969 Phys. Lett. B 30 517-522
- [2] Hecht K T and Adler A 1969 Nucl. Phys. A 137 129-143
- [3] Bohr A, Hamamoto I and Mottelson B R 1982 Phys. Scr. 26 267–272
- [4] Dudek J, Nazarewicz W, Szymanski Z and Leander G A 1987 Phys. Rev. Lett. 59 1405–1408
- [5] Castanos O, Moshinsky M and Quesne C 1992 Phys. Lett. B 277 238-242
- [6] Bahri C, Draayer J P and Moszkowski S A 1992 Phys. Rev. Lett. 68 2133–2136
- [7] Blokhin A L, Bahri C and Draayer J P 1995 Phys. Rev. Lett. 74 4149–4152
- [8] Ginocchio J N 1997 Phys. Rev. Lett. 78 436–439
- [9] Meng J, Sugawara-Tanabe K, Yamaji S, Ring P and Arima A 1998 Phys. Rev. C 58 R628-631
- [10] Meng J, Sugawara-Tanabe K, Yamaji S and Arima A 1999 Phys. Rev. C 59 154–163
- [11] Arima A 1999 Dynamical symmetries and nuclear structure RIKEN-AF-NP-276
- [12] Alberto P, Fiolhais M, Malheiro M, Delfino A and Chiapparini M 2001 Phys. Rev. Lett. 86 5015–5018
- [13] Ginocchio J N 2005 Phys. Rep. 414 165-261
- [14] Liang H, Shen S, Zhao P and Meng J 2013 Phys. Rev. C 87 014334-13
- [15] Lu B N, Zhao E G and Zhou S G 2013 Phys. Rev. C 88 024323-11
- [16] Ginocchio J N 1999 Phys. Rep. 315 231-240
- [17] Zhou S G, Meng J and Ring P 2003 Phys. Rev. Lett. 91 262501-4
- [18] He X T, Zhou S G, Meng J, Zhao E G and Scheid W 2006 Eur. Phys. J. A 28 265-269
- [19] Liang H, Long W H, Meng J and Van Giai N 2010 Eur. Phys. J. A 44 119-124
- [20] Song C Y, Yao J M and Meng J 2009 Chin. Phys. Lett. 26 122102-3
- [21] Song C Y, Yao J M and Meng J 2011 Chin. Phys. Lett. 28 092101-4
- [22] Song C Y and Yao J M 2010 Chin. Phys. C 34 1425–1427
- [23] Lalazissis G A, Gambhir Y K, Maharana J P, Warke C S and Ring P 1998 Phys. Rev. C 58 R45–R48
- [24] Sugawara-Tanabe K and Arima A 1998 Phys. Rev. C 58 R3065–R3068; ibid, 1999 60 019901(Erratum)
- [25] Ginocchio J N, Leviatan A, Meng J and Zhou S G 2004 Phys. Rev. C 69 034303-11
- [26] Guo J Y, Chen S W, Niu Z M, Li D P and Liu Q arXiv:1309.1523 [nucl-th]
- [27] Yang S C, Meng J and Zhou S G 2001 Chin. Phys. Lett. 18 196–198
- $[28]\,$ Zhang S Q, Meng J and Zhou S G 2003 Sci. China G 46 632–658
- [29] Meng J, Toki H, Zhou S G, Zhang S Q, Long W H and Geng L S 2006 Prog. Part. Nucl. Phys. 57 470-563
- [30] Guo J Y, Wang R D and Fang X Z 2005 Phys. Rev. C 72 054319-8
- [31] Guo J Y and Fang X Z 2006 Phys. Rev. C 74 024320-8
- [32] Liu Q, Niu Z M and Guo J Y 2013 Phys. Rev. A 87 052122-7
- [33] Zhang S S, Zhang W, Sun B H, Guo J Y and Zhou S G 2006 High Ener. Phys. Nucl. Phys. 30(S2) 97-99
- [34] Zhang S S, Sun B H and Zhou S G 2007 Chin. Phys. Lett. 24 1199–1202
- [35] Xu Q and Guo J Y 2012 Int. J. Mod. Phys. E 21 1250096–12
- [36] Lu B N, Zhao E G and Zhou S G 2012 Phys. Rev. Lett. 109 072501-5
- [37] Lu B N, Zhao E G and Zhou S G 2013 AIP Conf. Proc. 1533 63-69