On higher spin extension of the Jackiw-Teitelboim gravity model

K.B. Alkalaev †

I.E. Tamm Department of Theoretical Physics, P.N. Lebedev Physical Institute, Leninsky ave. 53, 119991 Moscow, Russia

Abstract

We formulate AdS_2 higher spin gravity as BF theory with fields taking values in $sl(N,\mathbb{R})$ algebra treated as higher spin algebra. The theory is topological and naturally extends the Jackiw-Teitelboim gravity model so as to include higher spin fields. The BF equations linearized about AdS_2 background are interpreted as describing higher spin partially-massless fields of maximal depth along with dilaton fields. It is shown that there are dual metric-like formulations following from the original linearized BF higher spin theory. The duality establishes a dynamical equivalence of the metric-like field equations that can be given either as massive scalar field equations or as conservation conditions for higher spin currents.

1 Introduction

Two-dimensional gauge systems including generally covariant theories and those with higher rank fields strike a balance between descriptive simplicity and nontrivial content. Basically, this can be explained by that in two dimensions, unlike in higher dimensions, only scalar or spinor modes can propagate. It follows that some of gauge systems are topological while others may exhibit a quite complicated dynamical structure which encodes matter modes. Both cases are intrinsically related because the matter can interact via topological gauge fields, while matter modes themselves can be represented as particular components of gauge fields. This is most clearly illustrated by the massive Schwinger model of fermions interacting via topological Maxwell field, or by the Liouville field originating as the metric component in the conformal gauge.

 $^\dagger email:$ alkalaev@lpi.ru

A prominent example of topological gauge systems is the Jackiw-Teitelboim gravity model [1, 2] which replaces the standard gravitation theory. Recall that in two dimensions Einstein tensor vanishes identically so that the simplest alternative is given by diffeomorphism invariant equation of motion of the form ¹

$$R - \Lambda = 0 \,, \tag{1.1}$$

where R is the Ricci scalar curvature built of the metric field $g_{mn}(x)$, while Λ is the cosmological constant. ² The above theory is obviously non-Lagrangian since one has a single equation for three variables. Adding an auxiliary scalar field $\phi(x)$ allows one to introduce the variational principle so that the resulting local action describes a particular dilaton gravity. The theory is however topological and contains no local degrees of freedom. At the same time, AdS_2 spacetime and black hole type objects are solutions of the Jackiw-Teitelboim model [3, 4].

The main purpose of this paper is to elaborate a higher spin generalization of pure gravity system governed by equation (1.1). We introduce higher rank gauge fields and their equations of motion in such a way that the resulting gauge system contains the gravitational subsector. As a guiding principle, we employ the frame-like formulation instead of the metric-like one used to postulate equation (1.1). The idea behind the frame-like approach originates from the Cartan formulation of the Einstein gravity where the metric is the symmetric component of the vielbein, while the antisymmetric component is a pure gauge by local Lorentz transformations.

Indeed, when extending to higher spins, the Jackiw-Teitelboim model viewed as a gauge theory of the AdS_2 global symmetry algebra $o(2,1) \sim sl(2,\mathbb{R})$ with a BF local action [5] has conceptual advantages over alternative approaches. One considers BF action with fields taking values in $sl(N,\mathbb{R})$ algebra with a finite parameter N. We show that such a theory can be interpreted as two-dimensional topological AdS_2 higher spin gravity theory. Note that conventional higher-spin metric-like fields $g_{m_1...m_s}$ with s Lorentz indices appear as particular tensor components of $sl(N,\mathbb{R})$ gauge connection rewritten in $o(1,1) \subset o(2,1) \subset sl(N,\mathbb{R})$ manifestly Lorentz covariant basis. Other components are either auxiliary or pure gauges. We find that the set of local fields over AdS_2 background consists of gauge fields with spins s=2,3,...,N and masses $m_s^2=s(s-1)\Lambda$ interpreted as topological partially-massless fields of maximal depth. ³

On going from the original BF formulation linearized over AdS_2 space to the metric-like formulation one finds out that there are two different but dynamically equivalent forms

¹Manifold \mathcal{M}^2 is a general two-dimensional space-time with local coordinates x^m , Lorentz world indices run m, n = 0, 1, Lorentz fiber indices run a, b = 0, 1, o(2, 1) fiber indices run A, B, C = 0, 1, 2, o(2, 1) invariant metric is $\eta^{AB} = (+ - -)$. The Levi-Civita tensor ϵ_{ABC} is normalized as $\epsilon_{012} = +1$. Two-dimensional anti-de Sitter spacetime AdS_2 has a radius L and a signature (+-), so that the cosmological constant is $\Lambda = -1/L^2$. The Levi-Civita tensor ϵ_{mn} is normalized as $\epsilon_{01} = +1$.

²The Riemann tensor in two dimensions has only a scalar curvature component as one can see from the identity $\epsilon^{mn}\epsilon^{kl}R_{mn,kl}=2R$.

³The dilaton sector comprising 0-form fields of the BF formulation will be analyzed elsewhere [16].

of the metric-like theory. In particular, it turns out that the gauge sector of the quadratic theory can be dually represented either by massive scalar equations or by conservation conditions for higher spin currents. Both scalar/current equations are invariant with respect to on-shell symmetries/improvements that guarantee the absence of local degrees of freedom.

While the BF theories under consideration are topological, we have in mind to study AdS_2 models with local degrees of freedom interacting via higher spin topological fields. Such an interacting theory will contain at least cubic field-current interactions of the form $\sum_s \int d^2x \sqrt{g_{AdS_2}} \ \Omega_m^{a_1...a_s} J^m_{a_1...a_s}$, where Ω_m are some Lorentz spin-s gauge fields, while J_m are conserved Lorentz spin-s currents built of matter fields. ⁴ We suggest that the field-current duality mentioned above will play an important role in the analysis of the higher spin field-current couplings in AdS_2 space. It follows that one of the aims of the present paper is to classify possible higher spin gauge fields in AdS_2 space that may couple to conserved higher spin currents.

2 The BF higher spin extension

Gravitational fields in two dimensions can be identified with o(2,1) connection $W^A(x)$ $T_A = dx^m W_m^A(x) T_A$, where T_A are o(2,1) basis elements with the commutation relations $[T_A, T_B] = -\epsilon_{ABC} T^C$. ⁵ Adding 0-form fields $\Psi(x) = \Psi^A(x) T_A$ one arrives at the BF action [5]

$$S_{JT}[W, \Psi] = \int_{\mathcal{M}^2} \Psi_A \mathcal{R}^A , \qquad (2.1)$$

where the curvature is given by $\frac{1}{2}dx^m \wedge dx^n \mathcal{R}_{mn}^A \equiv \mathcal{R}^A = dW^A - \epsilon^{ABC}W_B \wedge W_C$. The equations of motion following from (2.1) are

$$\mathcal{R}_{mn}^A = 0 , \qquad D_m \Psi^A = 0 , \qquad (2.2)$$

where $D_m = \partial_m + W_m$ is the o(2,1) covariant derivative. The 1-form connection denoted as $W_0^A = (\sqrt{-\Lambda} h_m^a, w_m)$ corresponds to AdS_2 spacetime. The zero-curvature constraint $\mathcal{R}_{mn}^A(W_0) = 0$ expresses Lorentz spin connection w_m via the zweibein h_m^a , while the latter yields AdS_2 spacetime metric g_{mn} through the standard identification $g_{mn} = \eta_{ab}h_m^a h_n^b$, where the fiber Minkowski metric is $\eta_{ab} = (+-)$.

The BF frame-like equations of motion are dynamically equivalent to the metric-like equations of original Jackiw-Teitelboim model. In particular, the original equation (1.1)

⁴ A particular example of such a d=2 higher spin model has been proposed by Vasiliev [6]. Remarkably, it is formulated as BF theory with fields taking values in some infinite-dimensional algebra, and up to now this is the only example of fully non-linear higher spin gauge theory with local degrees of freedom formulated at the action level.

⁵Note that the basis elements can be equivalently represented in the dualized form as $T^A = \frac{1}{2} \epsilon^{ABC} T_{BC}$. The commutation relations for antisymmetric basis elements $T^{AB} = -T^{BA}$ take the standard form $[T^{AB}, T^{CD}] = \eta^{AD} T^{BC} + 3$ terms.

is identified with one component of the curvature (2.2), while other two components are constraints (see [5] for more details). Despite the general covariance, the theory describes a scalar Liouville mode that follows from equation (1.1) in the conformal gauge. However, the theory is topological because the resulting Liouville equation possesses a residual coordinate invariance that gauges away all functional freedom of the solutions, see, e.g., [7].

The higher spin generalization of the theory (2.1) is straightforward. Two-dimensional BF theory with \mathcal{A} -valued fields, where $\mathcal{A} = sl(N,\mathbb{R})$ can be considered as particular higher spin gauge theory. In order to introduce higher spin gauge fields one decomposes the adjoint of $sl(N,\mathbb{R})$ algebra into totally symmetric irreps of AdS_2 spacetime global symmetry algebra $sl(2,\mathbb{R}) \approx o(2,1)$. For instance, in the case of $\mathcal{A} = sl(3,\mathbb{R})$ the basis elements T^{α} labelled by $\alpha = 1, ..., 8$ can be equally rearranged as $T^A \oplus T^{(AB)}$ with A, B = 0, 1, 2, i.e., as a direct sum of rank-1 and rank-2 totally symmetric and traceless $sl(2,\mathbb{R})$ algebra tensors. In particular, respective one-form connections are identified with framelike fields of spins 2 and 3. The analogous construction has been used in the context of 3d Chern-Simons higher spin theory with algebra $\mathcal{A} \oplus \mathcal{A}$, where $\mathcal{A} = sl(3,\mathbb{R})$ [8, 9].

It follows that Lie algebra $sl(N,\mathbb{R})$ for N=2,3,... can be interpreted as a finite-dimensional higher spin algebra in two dimensions provided that its basis elements are organized as follows

$$T_{A_1} \oplus T_{A_1 A_2} \oplus \cdots \oplus T_{A_1 \dots A_{N-1}},$$
 (2.3)

where $T_{A_1...A_k}$ are rank-k totally symmetric and traceless $sl(2,\mathbb{R})$ algebra tensors,

$$T_{(A_1...A_k)}: \eta^{MN} T_{MNA_3...A_k} = 0.$$
 (2.4)

It is worth noting that the decomposition (2.3) comes from the principal embedding of $sl(2,\mathbb{R}) \subset sl(N,\mathbb{R})$.

Introduce differential 0-form and 1-form fields taking values in $\mathcal{A} = sl(N, \mathbb{R})$ with basis elements (2.3)

$$\Psi(x) = \sum_{s=2}^{N-1} \Psi^{A_1 \dots A_{s-1}}(x) T_{A_1 \dots A_{s-1}}, \qquad W(x) = \sum_{s=2}^{N-1} dx^m W_m^{A_1 \dots A_{s-1}}(x) T_{A_1 \dots A_{s-1}}. \quad (2.5)$$

Expansion coefficients in the higher spin algebra basis elements (2.3) are to be identified as higher spin fields. They satisfy o(2,1) irreducibility conditions of the form (2.4).

Let us define AdS_2 higher spin gravity by the following BF action

$$S[W, \Psi] = g \sum_{s=2}^{N-1} \int_{\mathcal{M}^2} \Psi_{A_1 \dots A_{s-1}} \mathcal{R}^{A_1 \dots A_{s-1}} , \qquad (2.6)$$

⁶A non-principal embedding gives rise to a different dynamical content, and, generally speaking, to a different theory. In the context of 3d Chern-Simons higher spin theory non-principal embeddings and respective spectra were discussed, e.g., in [10].

with curvature two-forms given by $\mathcal{R}_{mn}^{A_1...A_{s-1}} = \partial_{[m}W_{n]}^{A_1...A_{s-1}} + [W_m, W_n]^{A_1...A_{s-1}}$, where $[\cdot, \cdot]$ is the Lie bracket in \mathcal{A} evaluated in basis (2.3), and g is a dimensionless coupling constant. The action (2.6) is invariant under the $sl(N, \mathbb{R})$ gauge transformations

$$\delta W_m^{A_1...A_{s-1}} = \partial_m \xi^{A_1...A_{s-1}} + [\xi, W_m]^{A_1...A_{s-1}} , \qquad \delta \Psi^{A_1...A_{s-1}} = [\xi, \Psi]^{A_1...A_{s-1}} , \qquad (2.7)$$

where $\xi(x) = \sum_{s=2}^{N-1} \xi^{A_1...A_{s-1}}(x) T_{A_1...A_{s-1}}$ is an $\mathcal{A} = sl(N, \mathbb{R})$ -valued gauge parameter. Note

that *a-priori* there is no metric tensor as the theory is completely formulated via differential forms. It follows that the theory is diffeomorphism invariant. However, an on-shell diffeomorphism transformation is simply a gauge transformation with particular gauge parameter.

Variational equations of motion for $\Psi^{A_1...A_{s-1}}$ and $W_m^{A_1...A_{s-1}}$ are the zero-curvature equation and the covariance constancy equation

$$\mathcal{R}_{mn}^{A_1...A_{s-1}} = 0$$
, $D_m \Psi^{A_1...A_{s-1}} = 0$, $s = 2, ..., N-1$, (2.8)

where $D_m = \partial_m + W_m$ is a covariant derivative with 1-form connection W_m given by (2.5). The covariance constancy equation involves both 0-form and 1-form fields, while the zero-curvature equation forms an independent gauge sector of the theory that can be analyzed separately. Let us emphasize that solutions of the zero-curvature equation are given locally by pure gauge 1-form fields. For the case of finitely many gauge fields this observation guarantees that the theory is topological and therefore contains no local degrees of freedom. However, in the next section we explicitly examine the gauge sector in the free field approximation to provide a particular physical interpretation of topological modes.

3 Linearized higher spin dynamics in AdS_2 space

The higher spin theory (2.6) contains a gravitational sector described by the Jackiw-Teitelboim action. The gravitational fields are singled out by setting all higher spin fields to zero except for those taking values in $sl(2,\mathbb{R})$ subalgebra. This particular truncation is obviously consistent. At the same time, AdS_2 space is the ground state of the theory since it solves gravitational equations (1.1).

Since the action (2.6) is non-linear containing at most cubic terms $\sim \Psi W^2$, one can consider the perturbation theory over a particular solution. To this end introduce 1-form W_0 describing AdS_2 spacetime and 0-form $\Psi_0 = 0$ as the background of the theory, so that all fields are decomposed as $W = W_0 + \Omega$ and $\Psi = \Psi_0 + \Phi$, where Ω and Φ are fluctuations over the background. The background fields W_0 and Ψ_0 obviously solve BF higher spin equations (2.8). Similarly, the 2-form curvatures are expanded as $\mathcal{R} = \mathcal{R}_0 + R + ...$, where $\mathcal{R}_0 \equiv \mathcal{R}(W_0)$ is the zero-curvature gravitational constraint (2.2), while R is the linearized curvature and the ellipsis denote second order corrections.

In the one-form sector, gauge fields Ω and their linearized curvatures R have the component form

$$\Omega^{A_1...A_{s-1}} = dx^m \,\Omega_m^{A_1...A_{s-1}} \,, \qquad R^{A_1...A_{s-1}} = D_0 \Omega^{A_1...A_{s-1}} \,, \tag{3.1}$$

where $D_0 = d + W_0$ is the background covariant derivative. Representing the zerocurvature condition (2.2) as $\mathcal{R}(W_0) \equiv D_0^2 = 0$ one obtains that linearized curvatures are invariant under the gauge transformations (2.7) expanded over the background,

$$\delta\Omega^{A_1...A_{s-1}} = D_0 \xi^{A_1...A_{s-1}} , \qquad (3.2)$$

where $\xi^{A_1...A_{s-1}}$ are 0-form gauge parameters. The Bianchi identities $D_0 R^{A_1...A_{s-1}} \equiv 0$ following from (3.1) are trivial since any 3-form in two dimensions vanishes identically. Fluctuations in the zero-form sector are inert under the gauge transformations $\delta \Psi^{A_1...A_{s-1}} = 0$, at least to the lowest order in the fields, cf. (2.7).

Let us compare the resulting set of 1-form higher spin gauge fields (3.1) in d=2 dimensions with fields in $d \geq 3$ dimensions. By massless higher spin fields in AdS_d we call 1-form connections $\Omega_m^{A_1...A_{s-1},B_1...B_{s-1}}$ taking values in o(2,d-1) finite-dimensional irreps of the symmetry type described by two-row rectangular Young diagram [11]. In the metric-like formulation these fields reduce to Fronsdal spin-s massless fields. Obviously, for d=2 all 1-form connections vanish identically except for s=2 case of the gravity, $\Omega_m^{A_1,B_1}=-\Omega_m^{B_1,A_1}$. It follows that there are no spin s>2 Fronsdal fields in AdS_2 .

At the same time, in two dimensions fiber indices of spin-2 connection can be dualized by virtue of o(2,1) Levi-Civita tensor as $\Omega_m^{A_1} = \epsilon^{A_1MN} \Omega_{mM,N}$. The point is that the gravitational connection Ω_m^A starts an infinite sequence of 1-form connections $\Omega_m^{A_1...A_{s-1}}$, s=2,3,..., taking values in o(2,1) finite-dimensional irreps described by one-row Young diagrams. These are fields appearing via the gauging of $sl(N,\mathbb{R})$ algebra in the basis (2.3). Note that in the $d\geq 3$ case, the connections with totally symmetric o(d-1,2) fiber indices describe partially-massless fields of maximal depth [12, 13, 14]. It follows that in AdS_2 space all possible higher spin s>2 fields are exhausted by "topological partially-massless" fields of maximal depth. ⁷

The BF equations (2.8) linearized around AdS_2 background space decouple into a direct sum of N-1 independent subsectors

$$D_0 \Omega^{A_1 \dots A_{s-1}} = 0 , (3.3)$$

$$D_0 \Phi^{A_1 \dots A_{s-1}} = 0 , (3.4)$$

each of spin s = 2, 3, ..., N. In what follows, we analyze a set of topological metric-like fields encoded in the gauge sector described by equations (3.3).

⁷In higher dimensions, the depth is a number of derivatives in the gauge transformation law for a spin-s dynamical field, t = 1, ..., s. In two dimensions, a maximal value of the depth equals t = s - 1 due to the Hodge duality mentioned above.

To identify the dynamical content of equations (3.3) one decomposes o(2,1) covariant gauge fields and linearized curvatures (3.1) into Lorentz algebra $o(1,1) \subset o(2,1)$ irreducible components,

$$\Omega_m^{A_1 \dots A_{s-1}} = \bigoplus_{k=0}^{s-1} \omega_m^{a_1 \dots a_k} , \qquad R_{mn}^{A_1 \dots A_{s-1}} = \bigoplus_{k=0}^{s-1} R_{mn}^{a_1 \dots a_k} , \qquad (3.5)$$

i.e., totally symmetric and traceless with respect to the Minkowski tensor η^{ab} . Assuming that Lorentz indices a_i are symmetrized with a unit weight, we find that the component form of (3.1) is given by

$$R_{mn}^{a_1...a_k} = \nabla_{[m} \,\omega_{n]}^{a_1...a_k} - \Lambda \frac{(s-k-1)(s+k)}{2(k+1)} h_{[m,c} \wedge \omega_{n]}^{ca_1...a_k} + \left[h_{[m}^{a_1} \wedge \omega_{n]}^{a_2...a_k} - \frac{1}{k-1} \,\eta^{a_1 a_2} h_{[m,c} \wedge \omega_{n]}^{ca_3...a_k} \right],$$
(3.6)

where covariant derivative $\nabla_m = \partial_m + w_m$ is defined with respect to the background Lorentz spin connection w_m , while h_m^a is the background zweibein. Quite analogously, one finds the component form of the gauge transformations (3.2), where gauge parameters $\xi^{a_1...a_k}$ are o(1,1) irreducible components of the o(2,1) gauge parameter $\xi^{A_1...A_{s-1}}$ (3.2).

The Lorentz component form of BF equations (3.3) is

$$R_{mn}^{a_1...a_k} = 0$$
, $k = 0, 1, ..., s - 1$. (3.7)

In the spin s=1 case the only field ω_m is identified with potential A_m . The resulting BF equation $R_{mn} \equiv F_{mn} = 0$ is different form the Maxwell equation $\nabla_m F^{mn} = 0$. It is worth noting here that a similar difference is between Maxwell and Chern-Simons equations in three dimensions.

The analysis of equations (3.7) for spin s > 1 is more intricate because there are more different rank fields and their gauge transformations including both differential and algebraic symmetry parameters. All fields entering equations of motion (3.7) can be divided into three subsets consisting of dynamical fields, Stueckelberg fields, and auxiliary fields. ⁸ It turns out that such a triple decomposition of the original field space can be done in two different ways. In the following sections we explicitly demonstrate how it works in the spin-2 case, and then briefly generalize to arbitrary spins.

3.1 Spin-2 example

In the spin s=2 case fields ω_m^a and ω_m are identified with the zweibein and Lorentz spin connection (more precisely, with their fluctuations over AdS_2 values h_m^a and w_m). Field

⁸As usual, Stueckelberg fields are assumed to be set to zero by virtue of their algebraic gauge transformations, while auxiliary fields are expressed via dynamical fields. Despite the absence of local degrees of freedom we keep using the term "dynamical fields" to designate independent fields of the theory.

equations read off from (3.7) and (3.6) at s=2

$$R_{mn} \equiv \nabla_m \omega_n - \nabla_n \omega_m - \Lambda h_{m,b} \wedge \omega_n^b + \Lambda h_{n,b} \wedge \omega_m^b = 0 , \qquad (3.8)$$

and

$$R_{mn}^a \equiv \nabla_m \omega_n^a - \nabla_n \omega_m^a + h_m^a \wedge \omega_n - h_n^a \wedge \omega_m = 0 , \qquad (3.9)$$

are invariant under the gauge transformations

$$\delta\omega_m = \nabla_m \xi - \Lambda h_{m,c} \xi^c , \qquad \delta\omega_m^a = \nabla_m \xi^a + h_m^a \xi . \qquad (3.10)$$

Using dualized field redefinitions $\omega_m^a \equiv \epsilon^{ab} e_{m,b}$ and $\omega_m \equiv \epsilon_{ab} \omega_m^{[ab]}$, the equations (3.8) and (3.9) can be cast into the standard form for zweibein e_m^a and Lorentz spin connection ω_m^{ab} . In particular, the above field redefinition succeeded by setting $\Lambda = 0$ reproduces the known expressions for curvatures of the (1+1) Poincare algebra.

The standard choice of dynamical fields is achieved by using the gauge parameter ξ as the Stueckelberg one to gauge away the trace component of ω_m^a . ⁹ To this end, using the AdS_2 background frame h_m^a one converts world indices into fiber ones and then decomposes ω_m^a into three o(1,1) irreducible components as follows

$$\omega^{a|b} = \omega^{(ab)} + \eta^{ab}\hat{\omega} + \frac{1}{2}\epsilon^{ab}\tilde{\omega} , \qquad (3.11)$$

where $\eta_{mn}\omega^{(mn)} = 0$. Residual components of ω_m^a can be combined into a single traceful symmetric tensor $\varphi_{ab} = \epsilon_a{}^c\omega_{cb} + \frac{1}{2}\eta_{ab}\tilde{\omega}$, so that its traceless and trace parts are given by $\omega^{(ab)}$ and $\tilde{\omega}$. Field φ^{ab} is then identified as dynamical field with the gauge symmetry transformation $\delta\varphi_{ab} = \nabla_a(\epsilon_b{}^c\xi_c) + \nabla_b(\epsilon_a{}^c\xi_c)$ (cf. our comments below (3.10)). Simultaneously, one considers equation (3.9) as the constraint which allows one to express auxiliary field ω_m via dynamical field φ^{ab} .

Equation $\epsilon^{mn}R_{mn} = 0$ (3.8) is dynamical. In terms of φ_{ab} it reads $\nabla^a\nabla^b\varphi_{ab} - \frac{3}{2}\Box_{AdS_2}\varphi + 2\Lambda\varphi = 0$, where \Box_{AdS_2} is the wave operator for the AdS_2 metric, and $\varphi = \eta^{ab}\varphi_{ab}$. Obviously, this is the Jackiw-Teitelboim equation (1.1) linearized around AdS_2 background. Imposing the conformal gauge which in our case says that the traceless part of φ^{ab} is set to zero, one is left with φ subjected to the residual equation

$$\left(\Box_{AdS_2} - 2\Lambda\right)\varphi = 0. \tag{3.12}$$

The above equation is invariant under the residual gauge transformation $\delta \varphi = \epsilon_{ab} \nabla^a \xi^b$ with a parameter satisfying the Killing equation $\nabla^a \xi^b + \nabla^b \xi^a - \eta^{ab} \nabla_c \xi^c = 0$. Therefore, one is left with the scalar field which is massive gauge field. ¹⁰

It is crucial that for $\Lambda \neq 0$ the above choice of dynamical fields is not unique. Another option is to use the gauge parameter ξ^a as a Stueckelberg one to completely gauge away

⁹Scalar parameter ξ can be dualized into an antisymmetric parameter $\xi^{[ab]}$. Obviously, this choice corresponds to the standard gauge fixing of local Lorentz symmetry.

¹⁰Recall here the well-known example of a scalar gauge field given by the trace of the metric field.

the field ω^m (3.10). Then, from (3.8) and (3.11) one derives the algebraic constraint $\Lambda\tilde{\omega}=0$, which says that the antisymmetric component of ω_m^a (3.11) is set to zero. It follows that a dynamical field is again a traceful symmetric tensor $\phi^{ab}=\omega^{(ab)}+\eta^{ab}\hat{\omega}$, but now its traceless and trace components are identified with $\omega^{(ab)}$ and $\hat{\omega}$. The gauge transformations now read $\delta\phi^{ab}=\Lambda^{-1}\nabla^a\nabla^b\xi+\eta^{ab}\xi$. Note that it reproduces the gauge transformation of a spin-2 partially-massless field.

Equation $\epsilon^{mn}R^a_{mn}=0$ (3.9) is dynamical. Imposing the gauge $\eta_{ab}\phi^{ab}=0$ one is left with a traceless component subjected to the residual equation

$$\nabla^a J_{ab} = 0 (3.13)$$

where $J^{ab} \equiv \epsilon^{ac} \phi^b{}_c$ is symmetric and traceless tensor field. A residual gauge invariance of (3.13) is given by $\delta J_{ab} = \Lambda^{-1} \epsilon_{ac} \nabla^c \nabla_b \xi + \Lambda^{-1} \epsilon_{bc} \nabla^c \nabla_b \xi$ with the scalar parameter satisfying differential constraint $(\Box_{AdS_2} - 2\Lambda)\xi = 0$. It follows that equation (3.13) can be seen as the conservation condition for a rank-2 current J_{ab} on the AdS_2 space.

3.2 Standard realization

By the standard realization we mean using a set of dynamical fields that generalize the metric-like variable of the spin-2 case. A careful analysis shows that all higher rank equations in (3.7) are constraints except for the k=0 equation $\epsilon^{mn}R_{mn}=0$ treated as the dynamical equation. The constraints allow one to express all fields via derivatives of dynamical fields which are scalar and rank-s traceless tensors, φ and $\varphi^{a_1...a_s}$. As in the spin-2 case, the scalar component φ is identified with antisymmetric component of ω_m^a (3.11). The rank-s traceless component $\varphi^{a_1...a_s}$ is identified with a totally symmetrized highest rank k=s-1 connection $\varphi^{ma_1...a_{s-1}}=\omega^{(m|a_1...a_{s-1})}$. Note that the symmetrized combination is automatically traceless because all other components in $\omega_m^{a_1...a_{s-1}}$ are gauged away using the Stueckelberg symmetry transformations.

Quite analogously, one shows that the only independent gauge parameter is $\xi^{a_1...a_{s-1}}$, while all others are treated as Stueckelberg parameters or auxiliary parameters expressed via derivatives of $\xi^{a_1...a_{s-1}}$. The leftover gauge symmetry with parameter $\xi^{a_1...a_{s-1}}$ allows one to impose the higher spin gauge,

$$\varphi^{a_1 \dots a_s} = 0 , \qquad (3.14)$$

which is legitimate since any o(1,1) rank-k irreducible tensor for k > 0 has just two independent components. Then, the only dynamical field is given by the scalar φ . Imposing the higher spin gauge (3.14) and solving the constraints in (3.7) one finds that the leftover equation reduces to the massive scalar equation with a particular value of mass-like term

$$\left(\Box_{AdS_2} - m_s^2\right) \varphi = 0 , \quad \text{where} \quad m_s^2 = s(s-1)\Lambda , \quad s \ge 2 .$$
 (3.15)

This is invariant under the residual gauge transformations.

$$\delta \varphi = \epsilon_{mn_1} \nabla^m \nabla_{n_2} \cdots \nabla_{n_{s-1}} \xi^{n_1 \cdots n_{s-1}} , \qquad (3.16)$$

provided that the gauge parameter satisfies the generalized Killing equation $\nabla^{(a_1} \xi^{a_2...a_s)} - \frac{1}{s-1} \eta^{(a_1 a_2)} \nabla_c \xi^{a_3...a_{s-1})c} = 0$, which is simply the stability transformation for the higher spin gauge (3.14). The massive scalar field (3.15) endowed with higher order derivative transformations for the scalar parameter (3.16) can be treated as topological partially-massless field of spin s and the maximal depth.

3.3 Dual realization

By the dual realization we mean using a set of dynamical fields that generalize the current-like variable of the spin-2 case. Quite analogously to the standard realization one shows that all higher rank equations in (3.7) can be treated as constraints except for the k = s-1 equation $\epsilon^{mn}R_{mn}^{a_1...a_{s-1}} = 0$ which is dynamical. The constraints allow one to express all fields via derivatives of dynamical fields also given by scalar and rank-s symmetric traceless tensor, ϕ and $\phi^{a_1...a_s}$. But now the scalar φ is identified with a trace component of ω_m^a (3.11), while the rank-s traceless field $\phi^{a_1...a_s}$ is identified with the symmetric traceless component of k = s - 1 connection $\omega_{m|a_1...a_{s-1}}$, namely $\phi_{ma_1...a_{s-1}} = \omega_{(m|a_1...a_{s-1})} - traces$. An independent gauge parameter is the scalar ξ , while the gauge transformations read

$$\delta\phi = \Box_{AdS_2}\xi - s(s-1)\Lambda\xi , \qquad \delta\phi_{a_1...a_s} = \nabla_{a_1}\cdots\nabla_{a_s}\xi + ... , \qquad (3.17)$$

where the ellipsis refers to proper symmetrizations and trace terms. Using gauge symmetry (3.17) one can impose the scalar gauge $\phi = 0$ along with the residual gauge parameter equation $\Box_{AdS_2}\xi - s(s-1)\Lambda\xi = 0$.

In the scalar gauge, the only dynamical field is the rank-s component $\phi^{a_1...a_s}$. Then, solving the constraints in (3.7) one finds that the independent dynamical equation is given by

$$\nabla^n J_{na_1...a_{s-1}} = 0$$
 for $J_{na_1...a_{s-1}} = \epsilon_n^m \phi_{ma_1...a_{s-1}}$, (3.18)

where dualized tensor field $J_{na_1...a_{s-1}}$ is totally symmetric and traceless. One identifies $J_{a_1...a_s}$ with spin-s conserved current on the AdS_2 space, while higher order derivative transformations in (3.17) are "improvements". On the other hand, the "improvement" transformation can treated as the gauge transformation of a spin-s partially massless field of maximal depth.

4 Conclusions and outlooks

In this paper we have introduced AdS_2 higher spin gravity defined via topological BF action with fields taking values in $sl(N,\mathbb{R})$ algebra. We identified the spectrum of higher spin gauge field fluctuations over AdS_2 background, and for $\Lambda \neq 0$ found out their dual description in terms of higher rank conserved currents.

Note that the proposed duality between two metric-like formulations guarantees that the two realizations describe the same dynamical system. Indeed, in order to single out independent fields and parameters we used Stueckelberg algebraic symmetry and eliminated auxiliary fields of the frame-like formulation. Whence, two metric-like formulations (3.15) and (3.18) coming from the original (linearized) BF system are dynamically equivalent. This type of duality is similar to that one in the WZNW theory with the equations of motions for a group variable g(x) given by a second order field equation $\partial^m (g^{-1}\partial_m g) = 0$. At the same time, introducing the current $J_m = g^{-1}\partial_m g$ one equivalently arrives at the first order conservation condition $\partial^m J_m = 0$ (e.g., see [15]).

It is instructive to compare linearizations of AdS_2 higher spin theory for $\mathcal{A} = sl(N, \mathbb{R})$ algebra and 3d Chern-Simons higher spin theory for $\mathcal{A} \oplus \mathcal{A}$ algebra. In the 3d case, choosing AdS_3 spacetime as the background for higher spin fields yields Fronsdal equations of motion for massless gauge fields with spins s=2,...,N that describe, however, no local degrees of freedom [8, 9]. In the 2d case, the same analysis does not yield Fronsdal equations, although the system is also topological. Instead, in the higher spin gauge which generalizes the conformal gauge in the gravity, the dynamical equations in the standard realization are scalar field equations with higher order derivative gauge symmetries guaranteeing the absence of propagating modes. These gauge systems can be interpreted as two-dimensional topological partially-massless fields.

It is remarkable that BF higher spin theory considered in the present paper has been in fact formulated within the unfolded approach to higher spin dynamics. In [16] we consider the theory from the unfolded formulation perspective and use the cohomological technique to analyze BF equations of motion. In particular, independent parameters, fields and equations of the standard and the dual realizations elaborated in sections 3.2 and 3.3 have a streamlined interpretation as cohomology elements of the so-called σ_{-} and σ_{+} nilpotent operators acting in the field space. Also, using the cohomological machinery we analyze equations for 0-forms which were not considered in the present paper.

The proposed BF higher spin theory can be extended so as to include an infinite tower of higher spin fields provided a gauge algebra \mathcal{A} is chosen to be one of infinite-dimensional higher spin algebras [17, 16, 18]. The resulting theory is also topological. Recall that BF higher spin theory can however have local degrees of freedom provided that fields take values in particular infinite-dimensional associative algebra [6].

Acknowledgements. I am grateful to M. Grigoriev, E. Skvortsov and M.A. Vasiliev for useful comments and discussions. I would like to thank the Galileo Galilei Institute for Theoretical Physics, Florence, Italy, for hospitality during my staying at GGI Workshop on Higher Spins, Strings and Duality, and to the Dynasty foundation for the financial support of the visit. The work is supported by RFBR grant 12-02-31838.

Note added. After the present paper has been submitted to arXiv, we were informed by Soo-Jong Rey that 2d higher spin theories were partly discussed in his talk [18], mainly from the holographic perspective.

References

- [1] C. Teitelboim, "Gravitation and Hamiltonian Structure in Two Space-Time Dimensions," *Phys. Lett.* **B126** (1983) 41.
- [2] R. Jackiw, "Lower Dimensional Gravity", Nucl. Phys. **B252** (1985) 343.
- [3] M. Cadoni and S. Mignemi, "Nonsingular four-dimensional black holes and the Jackiw-Teitelboim theory," *Phys. Rev.* **D51** (1995) 4319–4329, hep-th/9410041.
- [4] M. Spradlin and A. Strominger, "Vacuum states for AdS(2) black holes," JHEP 9911 (1999) 021 hep-th/9904143.
- [5] T. Fukuyama and K. Kamimura, "Gauge Theory of Two-dimensional Gravity," Phys. Lett. B160 (1985) 259.
 - K. Isler and C. Trugenberger, "A Gauge Theory of Two-dimensional Quantum Gravity," *Phys. Rev. Lett.* **63** (1989) 834.
 - A. H. Chamseddine and D. Wyler, "Topological gravity in (1+1)-dimensions," *Nucl. Phys.* **B340** (1990) 595–616.
- [6] M. A. Vasiliev, "Higher-Spin Gauge Interactions for Matter Fields in Two Dimensions," *Phys. Lett.* **B363** (1995) 51–57, hep-th/9511063.
- [7] R. Jackiw, "Gauge theories for gravity on a line," *Theor.Math.Phys.* **92** (1992) 979–987, hep-th/9206093.
- [8] M. Henneaux and S.-J. Rey, "Nonlinear $W_{infinity}$ as Asymptotic Symmetry of Three-Dimensional Higher Spin Anti-de Sitter Gravity," *JHEP* **1012** (2010) 007, 1008.4579.
- [9] A. Campoleoni, S. Fredenhagen, S. Pfenninger, and S. Theisen, "Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields," *JHEP* **1011** (2010) 007, 1008.4744.
- [10] A. Castro, E. Hijano, and A. Lepage-Jutier, "Unitarity Bounds in AdS(3) Higher Spin Gravity," *JHEP* **1206** (2012) 001, 1202.4467.
- [11] M. A. Vasiliev, "Cubic interactions of bosonic higher spin gauge fields in AdS(5)," Nucl. Phys. **B616** (2001) 106–162, hep-th/0106200.
- [12] S. Deser and A. Waldron, "Partial masslessness of higher spins in (A)dS," Nucl. Phys. B607 (2001) 577–604, hep-th/0103198.
- [13] Y.M. Zinoviev, "On massive high spin particles in AdS," hep-th/0108192.
- [14] E. D. Skvortsov and M. A. Vasiliev, "Geometric formulation for partially massless fields," *Nucl. Phys.* **B756** (2006) 117–147, hep-th/0601095.
- [15] C. R. Nappi, "Some Properties of an Analog of the Nonlinear σ Model," *Phys.Rev.* **D21** (1980) 418; E. Fradkin and A. A. Tseytlin, "Quantum Equivalence of Dual Field Theories," *Annals Phys.* **162** (1985) 31.
- [16] K. Alkalaev, "Global and local properties of AdS_2 higher spin gravity," preprint FIAN-TD-2013-16, to be published.

- [17] E. Fradkin and V. Y. Linetsky, "Higher spin symmetry in one-dimension and two-dimensions, I," *Mod.Phys.Lett.* **A4** (1989) 2635–2647.
- [18] Soo-Jong Rey, "News from Higher Spins: W_{∞} , Black Holes and Entropy", talk given at the workshop "Higher spins and holography", Simons center, March 2011.