

An approach to the quantization of black hole quasi-normal modes

Soham Pal*

*Department of Physics,
Birla Institute of Technology and Science, Pilani - Hyderabad Campus
Hyderabad 500078, India*

Karthik Rajeev

S. Shankaranarayanan†

*School of Physics,
Indian Institute of Science Education and Research (IISER-TVM),
Thiruvananthapuram 695016, India*

In this work we derive the asymptotic quasi-normal modes of a BTZ black hole using a quantum field theoretic Lagrangian. The BTZ black hole is a very popular system in the context of $2+1$ -dimensional quantum gravity. However to our knowledge the quasi-normal modes of the BTZ black hole have been studied only in the classical domain. Here we show a way to quantize the quasi-normal modes of the BTZ black hole by mapping it to the Bateman-Feshbach-Tikochinsky oscillator and the Caldirola-Kanai oscillator. We have also discussed a couple of other black hole potentials to which this method can be applied.

Keywords: Black Holes; Quantum Dissipative Systems.

PACS numbers: 04.30.-w,04.62.+v,04.60.-m,04.70.-s,04.70.Dy

1. Introduction

Resonant modes play a central role in the phenomena of energy flow among coupled systems as they provide characteristic information about the physical system. In the case of gravitating systems, these resonant modes are commonly referred as *quasi-normal modes*. These are damped perturbations about a fixed background (black holes or Neutron stars) that propagate to spatial infinity.¹⁻³

To understand the importance of the black hole quasi-normal modes and why they have attracted attention over the last four decades, let us consider a non-spherical collapse of a gravitational object that results in a slightly perturbed black hole. The black hole will then reach it's quiescent state by radiating away the perturbations in the form of gravitational waves. The radiation form the quasi-normal modes spectrum of the black hole. One can therefore define the quasi-normal modes

*Currently at Iowa State University, Ames

†Communicating author. Email: shanki@iisertvm.ac.in

of a black hole as single-frequency oscillations of the black hole spacetime that satisfy ingoing boundary conditions at the black hole event horizon and outgoing boundary conditions at spatial infinity.⁴ The real part (that corresponds to the frequency of the oscillation) and complex part (that corresponds to the damping rate) of the frequencies are independent of the initial perturbations and depend only on the properties of the black holes.

In the above way of describing the dynamics and the modes emanating to infinity, quasi-normal modes are assumed to be purely classical and are not quantized. However, there have been indications that these carry some information about quantum gravity.⁵⁻⁸ Firstly, QNM has been shown to be an useful tool in understanding AdS/CFT correspondence. In other words, it has been shown that there is an one-to-one mapping of the damping time scales (evaluated via simple QNM techniques) of black holes in Anti-de Sitter spacetimes and the thermalization time scales of the corresponding conformal field theory (which are, in general, difficult to compute).^{5,6,9} Secondly, using Bohr's correspondence principle "the transition frequencies at high quantum numbers equate the classical oscillation frequencies" it was realised that one could possibly identify the transition frequency between different black hole mass with the black hole quasi-normal mode frequencies.^{7,8,10} This indeed has provided the correct black hole entropy spectrum like other more rigorous semiclassical quantization approaches.¹¹

It is important to note that although the quasi-normal frequencies are derived using quantum mechanical techniques, however, by definition these are purely classical (see, for instance,^{2,3}). The question which we address in this work is the following: If these modes are indeed quantum mechanical (which do not preserve the total probability at later times) can they yield directly the quasi-normal mode frequencies and whether the asymptotic frequencies match with the classical calculations? Also, can the quantization procedure provide the link between the quasi-normal frequencies and the black hole entropy spectrum.

To authors knowledge, the quantization of black hole quasi-normal modes was attempted by Kim.¹² Kim maps the black hole quasi-normal modes to a dissipative system, in particular, the Bateman-Feshbach-Tikochinsky (BFT) oscillator and showed that the two — the black hole quasi-normal modes and the BFT oscillator — systems have the same group structure and, hence, maps the quantum states of the BFT oscillator to that of the quasi-normal modes.

However, there are several drawbacks in Kim's approach: First, the BFT oscillator has two modes. One that decays in time and other that gets amplified in time. It is natural to map the decay modes to quasi-normal frequencies, however, as pointed by Kim, it is not clear what is the role of the amplifying modes? Second, unlike the normal modes, the quasi-normal modes do not form a complete set of basis vectors in the sense that the system cannot be described as a sum over its QNMs, unless the black hole potential satisfy certain conditions.¹³ Kim's analysis does not look into this fundamental problem. Last, as is well-known, it is *only* possible to write down an equivalent Lagrangian that leads to the equation of motion

for a dissipative harmonic oscillator.^{14–18} It is imperative to show that several of these explicitly time-dependent, equivalent Hamiltonians (Lagrangians) are related to each other by a canonical (point) transformations.

In this work, we systematically address each of the above points and show that using Feynman and Vernon’s approach one can quantize the quasi-normal modes. The Feynman-Vernon formulation involves integrating over the bath degrees of freedom. In the case of black holes, we show that naturally the bath degrees of freedom correspond to the “stretched horizon”. The concept of the stretched horizon was originally treated by ’t Hooft in his paper on the quantum nature of a black hole.¹⁹ This description of the bath degrees of freedom of a black hole is consistent with other descriptions used in literature. For non-degenerate horizon, the stretched horizon description provides a universal framework and leads to finite results.

Our main aim of this work is to obtain the quasi-normal mode frequencies from a completely quantum approach aka path integral quantization procedure. We start by introducing, in Sec. (2), the conditions necessary for the completeness of these mode. To illustrate these conditions we discuss two examples in Appendix A, namely, the Regge-Wheeler potential and the rectangular potential well. As it is well-known the equation of motion containing the Regge-Wheeler potential, corresponding to the Schwarzschild black hole, can be mapped to Huen equation whose solutions are yet unknown,^{20,21} we simplified the situation by modeling the Regge-Wheeler potential to Poschl-Teller potential whose solutions are known.²² In Appendix B we have explicitly shown the quantization procedure that we used to derive the quasi-normal modes of the 2+1 dimensional BTZ black hole in Sec. (3). We have also discussed the oscillator models, in Appendix C, that initiated the current quantization procedure. In Sec. (4) we discuss our result and present concluding remarks.

Throughout this work we use the Planck units $c = \hbar = G = k_B = 1$.

2. Quantization of quasi-normal modes

Before we go into the quantization, it may be useful to look at the crucial differences between the normal and quasi-normal modes and the issues involved in quantization of the quasi-normal modes: Let us consider a system whose motion is defined by an equation of the Sturm-Liouville type, for example a finite vibrating string, whose motion is given by the wave equation. Since we can write the Sturm-Liouville type equation as a self-adjoint operator and can use the concept of normal modes to describe the dynamics of the system. Each normal mode is an oscillation in which all the components of the system move with the same frequency and the same phase.²³ The normal modes form a complete set and we can write down the most general motion of the system as a superposition of the normal modes.²³

Now suppose we introduce dissipation to this system, for example we couple a semi-infinite string to the finite string via a spring.²⁴ Such a system can dissipate energy away to infinity via radiation. The dynamics of the system are described by quasi-normal modes. Quasi-normal modes, on the other hand, form complete sets

only under some special conditions.¹³ Therefore a system of quasi-normal modes can be quantized only if those conditions are satisfied. It should be noted that both canonical^{25,26} and path integral²⁷ quantizations of QNMs of leaky optical cavities have been attempted.

In the rest of this section, we discuss the conditions required for the completeness of the quasi-normal modes and definition of the inner product and normalization of the quasi-normal modes.

2.1. Completeness of QNMs

We use the Klein-Gordon (KG) equation to describe wave propagation in curved space

$$\tilde{D}\phi(x, t) \equiv [\partial_t^2 - \partial_x^2 + V(x)] \phi(x, t) = 0. \quad (1)$$

Assuming the time dependency of the form $e^{-i\omega t}$, the KG equation reduces to a Schrödinger-like equation

$$D\phi(x) \equiv [\partial_x^2 + \omega^2 - V(x)] \phi(x) = 0. \quad (2)$$

The QNMs are the eigenfunctions $f(x, t)$ of the operator \tilde{D} or equivalently the eigenfunctions $f(x)$ of the operator D , that satisfy the condition

$$f(x) \sim e^{i\omega|x|}, \quad |x| \rightarrow \infty. \quad (3)$$

If the potential $V(x)$ is finite everywhere and vanishes sufficiently rapidly as $|x| \rightarrow \infty$ and there are spatial discontinuities in $V(x)$ marking a “cavity” I , then for $x \in I$, $\{f_j\}$ is a complete set and we can express the time evolution of the wavefunction as a sum over the QNMs, $\phi(x, t) = \sum_j a_j f_j(x) e^{-i\omega_j t}$ or $\phi(x, t=0) = \sum_j a_j f_j$.^{4,13}

2.2. Inner product and normalization

Let $\phi(x, t)$ and $\psi(x, t)$ be two wavefunctions in the space spanned by $\{f_j(x, t)\}$ and let $\{f_j\}$ be complete. Therefore $\phi(x, t=0) = \sum_j a_j f_j(x)$ and $\psi(x, t=0) = \sum_j b_j f_j(x)$. Defining the two-component wavefunction $\phi = (\phi, \partial_t \phi)^T$ we give the inner product between the two wavefunctions as²⁹⁻³¹

$$(\phi|\psi) = i \left\{ \int_{-a}^a dx [\phi(\partial_t \psi) + (\partial_t \phi)\psi] + [\phi(-a)\psi(-a) + \phi(a)\psi(a)] \right\} \quad (4)$$

where $|a|$ is a finite value. The inner product gives the normalization relation between the QNMs

$$(\mathbf{f}_j|\mathbf{f}_j) = 2\omega_j \int_{-a}^a f_j^2(x) dx + i [f_j^2(-a) + f_j^2(a)] \quad (5)$$

where $\mathbf{f}_j = (f_j, -i\omega_j f_j)^T$ and the expansion coefficients are given by

$$a_j = \frac{(\phi_m|\mathbf{f}_j)}{(\mathbf{f}_j|\mathbf{f}_j)}, \quad b_j = \frac{(\psi_m|\mathbf{f}_j)}{(\mathbf{f}_j|\mathbf{f}_j)}. \quad (6)$$

Eqs. (4) and (5) give the orthogonality relation

$$\int_{-a}^a f_j(x) f_k(x) dx = \delta_{jk} - i \left(\frac{f_j(-a) f_k(-a) + f_j(a) f_k(a)}{\omega_j + \omega_k} \right). \quad (7)$$

3. Quasinormal modes of the BTZ black hole

We consider a massless scalar field in a BTZ black hole background, whose metric is given by

$$ds^2 = f(r) dt^2 - f(r)^{-1} dr^2 - r^2 d\varphi^2 \quad (8)$$

where $f(r) = -M + r^2 + J^2/4r^2$. In the remainder of the article we consider the black hole mass, M , to be 1 and the angular momentum, J , to be 0. The external black hole horizon is at $r = r_H = M^{1/2} = 1$ and the internal black hole horizon is at $r = 0$. Spatial infinity is at $r \rightarrow \infty$. To make the following calculations easier we now introduce the tortoise coordinate, $dx = dr/f(r)$. In the tortoise coordinate, the horizon is mapped to $x \rightarrow -\infty$ and the spatial infinity to $x \rightarrow 0$. The BTZ metric in the tortoise coordinate is given by

$$ds^2 = f(r)[dt^2 - dr^2] - r^2 d\varphi^2 \quad (9)$$

and the equation of motion of the massless scalar field is

$$[\partial_t^2 - \partial_x^2] \psi - \frac{f(r)}{r^2} \partial_\varphi^2 \psi = 0. \quad (10)$$

Writing the scalar field as a sum of partial waves, $\psi = \sum_l \psi_l(t, x) \exp(il\varphi)$, we get

$$[\partial_t^2 - \partial_x^2] \psi_l + V_l(x) \psi_l = 0 \quad (11)$$

where the potential $V_l(x) = l^2 f(r)/r^2$.

The BTZ black hole is asymptotically AdS. In the AdS geometry, fluctuations cannot escape to infinity and therefore dissipation of energy by time-localized fluctuations happen at the black hole horizon,⁴⁰ i.e. the scalar field satisfies completely ingoing boundary conditions at the black hole event horizon. The dissipation can be considered to be due to interaction between the scalar field and a bath. As discussed in Sec. (2) and Appendix A, the completeness of the QNMs requires us to keep the potential finite. While this looks artificial, in the case of black hole, this corresponds to the fact that the region close to the horizon that leads to large blue-shifting of modes needs to be removed. This naturally leads to the concept of a stretched horizon, which, in turn, helps us to define a bath. We consider a stretched horizon, $r = r_H + \varepsilon = 1 + \varepsilon$, and treat the variables between the stretched horizon and the horizon as the bath variables.⁴¹ The equations of motion of the bath variables are

$$\ddot{\psi}_l(t, x) = \partial_x^2 \psi_l(t, x) - V_l(x) \psi_l(t, x). \quad (12)$$

The value of the potential, $V_l(x)$ is vanishingly small near the horizon and can be neglected and the bath variables may be considered as free fields. Therefore the

total action for the scalar field and the bath variables in Euclidean time is

$$S = \frac{1}{2} \int_{\text{periodic}} d\mathbf{x}^2 \left[(\partial_\tau \psi_l)^2 + (\partial_x \psi_l)^2 + V_l(x) \psi_l^2 \right] \quad (13)$$

$$= S_{sys} + S_{bath} \quad (14)$$

where S_{sys} is the action over the fields in $x \in (-\infty, 0)$ and S_{bath} is the action over the fields in the region $x \in (-\infty, x(1 + \epsilon)]$ and $V_l(x) = 0$, for $x \in (-\infty, x(1 + \epsilon)]$. We consider the presence of a source $\chi = \sum_l \chi_l$ that interacts with the massless scalar field via

$$(\psi_l, \chi_l) = i \left\{ \int_{-\infty}^0 dx [\psi_l (\partial_\tau \chi_l) + (\partial_\tau \psi_l) \chi_l] + [\psi_l(-\infty) \chi_l(-\infty)] \right\}. \quad (15)$$

The global generating functional is

$$W[\chi_l] = \int \mathcal{D}\psi_l e^{-S(\psi_l, \partial_\mu \psi_l) + (\psi_l, \chi_l)}. \quad (16)$$

Separating the global generating functional into system and bath factors and integrating out the latter, leaves us with

$$W[\chi_l]^r = \int \mathcal{D}\psi_l^{(sys)} e^{-S_{sys} - S_{IF} + (\psi_l, \chi_l)} \quad (17)$$

where the influence functional, S_{IF} is defined by

$$e^{-S_{IF}} = \int \mathcal{D}\psi_l^{(bath)} e^{-S_{bath}}. \quad (18)$$

To make the following calculations easier we approximate the bath action by assuming that the horizon is at $x = -N$ for some very large N and the stretched horizon is at $x = -N + \delta$. The bath action under this approximation is

$$\begin{aligned} S_{bath} &= \frac{1}{2} \int_0^\beta d\tau \int_{-N}^{-N+\delta} dx [(\partial_\tau \psi_l)^2 + (\partial_x \psi_l)^2] \\ &= \frac{1}{2} \int_0^\beta d\tau \int_0^\delta d\xi [(\partial_\tau \psi_l)^2 + (\partial_x \psi_l)^2] \end{aligned} \quad (19)$$

where $\xi = x + N$. Following the procedure in Appendix B, we expand the bath fields,

$$\psi_l^{(bath)} = T \sum_k \left\{ \psi_{l,k}(-N) \frac{\delta - \xi}{\delta} + \sum_{u=1}^{\infty} \psi_{l,k,u} \sin\left(\frac{\pi u \xi}{\delta}\right) \right\} e^{-i\nu_k \tau}. \quad (20)$$

As defined earlier, T is the Hawking temperature and ν_k are the bosonic Matsubara frequencies. Therefore the influence functional is

$$e^{-S_{IF}} = \exp \left[\frac{T}{2} \sum_k |\psi_{l,k}(-N)|^2 \left[\frac{\delta \nu_k^2}{3} + \frac{1}{\delta} - \sum_{u=1}^{\infty} \frac{2(\delta \nu_k^2 / \pi u)}{\delta \nu_k^2 + \pi^2 u^2 / \delta} \right] \right] \quad (21)$$

$$= \exp \left[-\frac{T}{2} \sum_k |\psi_{l,k}(-N)|^2 c(\delta, \nu_k) \right]. \quad (22)$$

We expand the system fields as ψ_l as $\psi_l = \sum_k \psi_{l,k} = \sum_k \sum_m a_{lkm} f_m(x) \exp\{\omega_m \tau\} = \sum_k \sum_m a_{lkm}(\tau) f_m$. The modes $f_m(x)$ satisfy the ingoing boundary conditions at the horizon and if we impose the Dirichlet boundary conditions at $r \rightarrow \infty$, then these modes would correspond to the quasi-normal modes of the BTZ black hole. Similarly we expand the source, $\chi_l = \sum_k \sum_m b_{lkm}(\tau) f_m$. Now temporarily assuming that $V_l(x) = 0, \forall x$, we calculate the effective generating functional for the system to be

$$W[\chi_l]_0^r = \int \mathcal{D}\psi_l^{(sys)} \left[\exp \left\{ T \sum_k \left[- \sum_{mn} a_{lkm} a_{lkn} \right. \right. \right. \\ \left. \left. \left. \times \left(\omega_m \omega_n \delta_{mn} - \frac{i\omega_m \omega_n f_m(-N) f_n(-N)}{\omega_m + \omega_n} \right) + 2 \sum_m \omega_m a_{lkm} b_{lkm} \right] - S_{IF} \right\} \right] \quad (23)$$

where we have used the ‘‘orthogonality’’ condition

$$\int dx f_m(x) f_n(x) = \delta_{mn} - i \frac{f_m(-N) f_n(-N)}{\omega_m + \omega_n}.$$

Before proceeding further we introduce the following expressions to simplify the remaining mathematical equations:

$$\begin{aligned} \mathcal{F}_{mn} &= \frac{f_m(-N) f_n(-N)}{\omega_m + \omega_n}, \\ \mathcal{I} &= \int_{-N}^0 dx V_l(x) f_m(x) f_n(x), \\ \mathcal{O}_{kmn} &= \left(\theta(k) \frac{\omega_m}{\omega_m - ic(\delta, \nu_k)} + \theta(-k) \frac{\omega_n}{\omega_n + ic(\delta, \nu_k)} \right), \\ \mathcal{P}_{kmn} &= \left[1 - iT \frac{\mathcal{F}_{mn}}{2\omega_m \omega_n} \mathcal{O}_{kmn} (1 + iT \mathcal{F}_{mn} b_{lkm} b_{lkn}) \right]. \end{aligned}$$

Now for the more general case of $\omega_m \neq \omega_n$, we can simplify the generating functional to

$$\begin{aligned} W[\chi_l]_0^r &= \int \mathcal{D}\psi_l^{(sys)} \exp \left[T \sum_k \left\{ - \sum_{mn} a_{lkm} a_{lkn} f_m(-N) f_n(-N) \right. \right. \\ &\quad \left. \left. \times \left(\frac{c(\delta, \nu_k) (\theta(k) \omega_m - \theta(-k) \omega_n) + i\omega_m \omega_n}{\omega_m + \omega_n} \right) + \sum_m 2\omega_m a_{lkm} b_{lkm} \right\} \right] \\ &= \exp \left[iT \sum_{kmn} b_{lkm} b_{lkn} \mathcal{F}_{mn} \mathcal{O}_{kmn} \right]. \quad (24) \end{aligned}$$

Therefore the generating functional for the actual nonzero potential is

$$W[\chi_l]^r = \exp \left[-\beta \int_{-N}^0 dx V_l(x) f_m(x) f_n(x) \frac{1}{2\omega_m \omega_n} \frac{\partial}{\partial b_{lkm}} \frac{\partial}{\partial b_{lkn}} \right] W[\chi_l]_0^r. \quad (25)$$

The expression in the right-hand side of Eq. (25) can be calculated by perturbatively expanding the exponential. Taking only the first two terms of the perturbative expansion we get,

$$W[\chi_l]^r = \exp \left\{ iT \sum_{kmn} b_{lkm} b_{lkn} \mathcal{F}_{mn} \mathcal{O}_{kmn} \right\} \left[1 - iT \frac{\mathcal{F}_{mn} \mathcal{O}_{kmn}}{2\omega_m \omega_n} (1 + iT \mathcal{F}_{mn} b_{lkm} b_{lkn}) \right]. \quad (26)$$

The generating functional in Eq. (25) is related to the temperature Green's function via

$$\left. \frac{\partial^2 W[\chi_l]^r}{\partial b_{lkm} \partial b_{lkn}} \right|_{\chi=0} = -4\omega_m \omega_n \mathcal{G}_{mn}(\nu_k). \quad (27)$$

Double differentiating Eq. (26) we get,

$$\begin{aligned} \frac{\partial^2 W[\chi_l]^r}{\partial b_{lkm} \partial b_{lkn}} = \exp \left\{ iT \sum_{kmn} b_{lkm} b_{lkn} \mathcal{F}_{mn} \mathcal{O}_{kmn} \right\} & \left[(iT b_{lkm} \mathcal{F}_{mn} \mathcal{O}_{kmn})^2 \mathcal{P}_{kmn} \right. \\ & + iT \mathcal{F}_{mn} \mathcal{O}_{kmn} \mathcal{P}_{kmn} + iT T^2 b_{lkm} b_{lkn} \frac{\mathcal{F}_{mn}^3 \mathcal{O}_{kmn}^2}{2\omega_m \omega_n} \\ & \left. + iT T^2 \frac{\mathcal{F}_{mn}^3 \mathcal{O}_{kmn}^2}{2\omega_m \omega_n} b_{lkn}^2 + IT \frac{\mathcal{F}_{mn}^2 \mathcal{O}_{kmn}}{2\omega_m \omega_n} \right]. \quad (28) \end{aligned}$$

The condition that the source, $\chi = 0$, implies that the coefficients b_{klj} are all zero. Applying this condition to Eq. (28) we get

$$\left. \frac{\partial^2 W[\chi_l]^r}{\partial b_{lkm} \partial b_{lkn}} \right|_{\chi=0} = iT \mathcal{F}_{mn} \mathcal{O}_{kmn} + \frac{T^2 \mathcal{F}_{mn}^2 \mathcal{O}_{kmn}^2}{2\omega_m \omega_n} + \frac{IT \mathcal{F}_{mn}^2 \mathcal{O}_{kmn}}{2\omega_m \omega_n}. \quad (29)$$

The temperature Green's function therefore turns out to be

$$\begin{aligned} \mathcal{G}_{mn}(\nu_k) &= -\frac{1}{4\omega_m \omega_n} \left[iT \mathcal{F}_{mn} \mathcal{O}_{kmn} + \frac{T^2 \mathcal{F}_{mn}^2 \mathcal{O}_{kmn}^2}{2\omega_m \omega_n} + \frac{IT \mathcal{F}_{mn}^2 \mathcal{O}_{kmn}}{2\omega_m \omega_n} \right] \quad (30) \\ &= \frac{T \mathcal{F}_{mn}}{8\omega_m^2 \omega_n^2} \frac{[\omega_m \omega_n + ic(\delta, \nu_k) (\theta(k)\omega_m - \theta(-k)\omega_n)]}{(\omega_m - ic(\delta, \nu_k)) (\omega_n + ic(\delta, \nu_k))} \\ &\times \left[(i2\omega_m \omega_n + IT \mathcal{F}_{mn}) + \frac{T \mathcal{F}_{mn} (\omega_m \omega_n + ic(\delta, \nu_k) (\theta(k)\omega_m - \theta(-k)\omega_n))}{(\omega_m - ic(\delta, \nu_k)) (\omega_n + ic(\delta, \nu_k))} \right]. \quad (31) \end{aligned}$$

The temperature Green's function is related to the real-time retarded propagator as $\mathcal{G}_{mn}(\nu_k) = G_{mn}^R(i\nu_k)$ i.e., we make the substitution $c(\delta, \nu_k) \rightarrow c(\delta, i\nu_k)$ in Eq. (31) to get the real time retarded propagator. We can then compute the QNM frequencies as the poles of G_{mn}^R , i.e., $\omega_n = 0$ or $\omega_n = \pm ic(\delta, i\nu_k)$. We can express the function $c(\delta, i\nu_k)$ in terms of the digamma function, which is the logarithmic derivative of the gamma function,

$$c(\delta, i\nu_k) = \alpha_k - \frac{1}{\pi} \left(\psi^{(0)}(1 - s_k) + \psi^{(0)}(1 + s_k) \right) \quad (32)$$

where $\psi^{(0)}$ is the digamma function,

$$s_k = \frac{\delta \nu_k}{\pi}; \quad \alpha_k = -\frac{\delta \nu_k^2}{3} + \frac{1}{\delta} - \frac{2\gamma}{\pi} \quad (33)$$

and γ is the Euler-Mascheroni coefficient. The digamma function can be expressed as a converging rational zeta series if $|s_k| < 1$. We can ensure that this condition is satisfied by choosing a sufficiently small δ . The expression $\psi^{(0)}(1-s_k) + \psi^{(0)}(1+s_k)$ tends to infinity as $|s_k| \rightarrow 1$. Therefore to get the asymptotic QNM frequencies we expand $\psi^{(0)}(1-s_k) + \psi^{(0)}(1+s_k)$ at $s_k = 1$,

$$\psi^{(0)}(1-s_k) + \psi^{(0)}(1+s_k) = \frac{1}{1-s_k} + (1-2\gamma) + \frac{1}{3}(\pi^2-3)(s_k-1) + \mathcal{O}(s_k^2). \quad (34)$$

The asymptotic QNM frequencies are therefore

$$\omega_n^{\text{asym}} = \pm i \left\{ \alpha_k + \frac{1}{\pi} \left(\frac{1}{1-s_k} + (1-2\gamma) + \frac{1}{3}(\pi^2-3)(s_k-1) \right) \right\}. \quad (35)$$

As we have mentioned we have to choose a sufficiently small δ to derive the asymptotic QNM frequencies, one obvious point of curiosity would be the nature of the frequencies in the limit $\delta \rightarrow 0$. However this limit doesn't exist for the quantities α_k and s_k . To overcome this, we can consider δ to be of the order of k^{-1} for very large k (as defined in Appendix B, $k \in \mathbb{Z}$). Therefore under the limit $\delta \rightarrow 1/k$ the asymptotic QNM frequencies turn out to be

$$\omega_n^{\text{asym}} \sim \pm \frac{iT}{2} k. \quad (36)$$

The BTZ black hole metric had originally been obtained as a solution to the (2+1)-dimensional version of classical Einstein-Hilbert theory of gravitation with a negative cosmological constant. However this version of the Einstein-Hilbert theory does not allow propagating bulk degrees of freedom. To resolve this problem, alternative (2+1)-dimensional theories of gravitation, such as the topologically massive gravity (TMG) and new massive gravity (NMG), have been proposed. Both TMG and NMG admit the BTZ metric as a solution. In recent years, there have been numerous studies concerning the QNMs of BTZ black holes in all the three theories of gravity; many of which have dealt with massive scalar fields in more generalized BTZ backgrounds. All these classical calculations⁴²⁻⁴⁵ indicate that the scalar QNM frequencies of the BTZ black hole are of the form

$$\omega = l - i\mu T(k+1)$$

(l is the angular momentum of the scalar field and μ is a constant) which in the limit of large k becomes

$$\omega \sim -iT k.$$

This result is similar to our calculations. More general calculations,⁴⁶ in the context of NMG, show that QNM frequencies are of the form

$$\omega = -iT \frac{h(k^2)}{g(k)}$$

where h and g are functions of k , and in the asymptotic limit becomes

$$\omega \sim -i \frac{T}{2} k. \quad (37)$$

Though we have worked with a massless scalar field, the frequencies given by Eq. (35) are similar in structure to those obtained in the case of NMG and coincide in the limit of large k . Moreover, from Eq. (36), we can see that the QNM frequencies are proportional to the Hawking temperature, just like in the case of the classical models.

From the point of view of NMG, black holes with QNM frequencies of the form, $\omega = \omega_R + i\omega_I$, $\omega_R = 0$ and $\omega_I > 0$, are unstable.⁴⁷ This is because modes with positive ω_I will grow exponentially with time. Considering the classical QNM spectra, the BTZ black hole are stable in NMG. However, the quantum QNM spectra, that we obtained, has modes with either positive or negative imaginary parts and therefore doesn't confirm to this stability condition. One way to introduce stability would be to consider a finite time frame so that QNMs whose frequencies have positive imaginary part do not blow up. Another possible way would be consider a theory of (2+1)-dimensional gravity that naturally introduces a cutoff for the QNM frequencies. To authors knowledge, none of the competing classical theories of gravity do that.

4. Conclusion

In this paper we have presented two descriptions of the QNMs - one classical and the other quantum mechanical. In the classical picture (Appendix C) we have drawn analogy between the QNMs and the damped simple harmonic oscillator. We have shown that there's a connection between the system of QNMs and the Bateman oscillator and therefore the study of one system may give valuable information about the other. The damped harmonic oscillator is considered to be coupled to an amplified one and the system therefore follows the Bateman equations of motion.

We used the path integral formalism prescribed by Feynman and Vernon to develop our quantum theory. This is because the system-bath model of the Feynman-Vernon formalism is in spirit similar to Bateman's dual oscillator system and the physics can therefore be intuitively understood. Like in the case of the Bateman dual oscillator, the quantization of the system-bath model involves integrating over the bath degrees of freedom. We have given a consistent description of the bath degrees of freedom for a black hole using the concept of a stretched horizon, originally developed by 't Hooft.

In Sec. (3) we have given a detailed derivation of the QNM frequencies of a BTZ black hole, using our quantum theory. The asymptotic QNM frequencies in

this case matches with the classical results. However, our results do not show that the imaginary part of the asymptotic QNM frequencies are quantized.

Moreover the final forms of the generating functional obtained in Sec. (3) and Appendix B can be used to derive interesting physical quantities, such as the free energy, defined as, $F_c^R[\chi] \equiv \ln W_c^R[\chi]$ and entropy. We hope to present these applications of the generating functional in a future article.

Acknowledgments

The authors thank Sashideep Gutti for discussions. SP thanks IISER-TVM for hospitality for carrying out the initial part of the work and is supported by Junior Research Fellowship of CSIR, India. KR is supported by the INSPIRE fellowship, DST, India. The work is supported by Max Planck-India Partner Group on Gravity and Cosmology. SS is partially supported by Ramanujan Fellowship of DST, India.

Appendix A. Model Potentials

Taking into account the restrictions set on the potential in Sec. (2), we will use potentials that have support in the interval, $I \equiv [-a, a]$. In this section we will describe two such potentials and the associated QNMs.

The rectangular barrier is given below as an illustration of the method as it gives exact quasi-normal mode frequencies while the modified Pöschl-Teller potential has all the features of Regge-Wheeler potential.²²

Appendix A.1. Rectangular barrier

The rectangular barrier potential is given by

$$V(x) = \begin{cases} V_0 \neq 0, & x \in [-a, a] \\ 0, & \text{otherwise.} \end{cases} \quad (\text{A.1})$$

For this barrier potential the boundary conditions that should be satisfied by the QNMs are reduced to

$$\frac{\partial_x f(x)}{f(x)} = \pm i\omega, \quad x = \pm a$$

The eigenfunctions of D are given by Eq. (A.1)

$$f(x) = \begin{cases} P e^{i\omega x}, & x > a \\ Q e^{ikx} + R^{-ikx}, & a \geq x \geq -a \\ S e^{-i\omega x}, & -a > x, \end{cases} \quad (\text{A.2})$$

where $k = \sqrt{\omega^2 - V_0}$ and P , Q , R and S are constants. Maintaining the continuity of $f(x)$ and $\partial_x f(x)$ we obtain

$$\left(\frac{k - \omega}{k + \omega} \right)^2 e^{i4ka} = 1$$

which is simplified to

$$k = i\omega_0 \cos(ka) \quad \text{or} \quad k = -i\omega_0 \sin(ka), \quad (\text{A.3})$$

where $\omega_0 = V_0^{1/2}$. Depending on the sign of V_0 , Eq. (A.3) gives different sets of QNM frequencies.³²

Appendix A.1.1. *Negative* V_0

For $V_0 < 0$, ω_0 is purely imaginary, $\omega_0 = i|\omega_0|$. Therefore from Eq. (A.3)

$$k = -|\omega_0| \cos(ka) \quad \text{or} \quad k = |\omega_0| \sin(ka). \quad (\text{A.4})$$

Eq. (A.4) sometimes have solutions for real k if $|k| \leq |\omega_0|$. Since $\omega^2 = k^2 + \omega_0^2 = k^2 - |\omega_0|^2 \leq 0$, therefore the values of ω are purely imaginary corresponding to bound states of the potential and not quasi-normal modes.

However, if $\text{Im}(k) \gg 0$, then $\cos(ka) \approx \exp(-iak)/2$ and $\sin(ka) \approx -\exp(-iak)/2$. In such a situation the approximate values of the quasi-normal frequencies are given by the j -th solutions of

$$2k \approx -|\omega_0|e^{-ika}. \quad (\text{A.5})$$

Appendix A.1.2. *Positive* V_0

For $V_0 > 0$, ω_0 is real. If $\omega = i|\omega|$, corresponding to a purely damped mode, then from Eq. (A.3)

$$|k| = \omega_0 \cosh(|k|a) \quad \text{or} \quad k = 0. \quad (\text{A.6})$$

The $k = 0$ mode is not a true quasi-normal frequency, the other equation gives the physical quasi-normal frequencies

$$|k|a = \omega_0 a \cosh(|k|a). \quad (\text{A.7})$$

For small $\omega_0 a$ there are two quasi-normal frequencies, one of which is given by the perturbative expression

$$k = i\omega_0 \left\{ 1 + \frac{1}{2}(\omega_0 a)^2 + \frac{13}{24}(\omega_0 a)^4 + \mathcal{O}([\omega_0 a]^6) \right\} \quad (\text{A.8})$$

$$\text{or} \quad \omega = i\omega_0(\omega_0 a) \left\{ 1 + \frac{2}{3}(\omega_0 a)^2 + \frac{4}{5}(\omega_0 a)^4 + \mathcal{O}([\omega_0 a]^6) \right\}. \quad (\text{A.9})$$

No such representations exist for the other quasi-normal mode. However if $\omega_0 a \sim 0.663$, then the two quasi-normal modes merge and for $\omega_0 a \gtrsim 0.663$ there are no quasi-normal modes.

In this case, if $\text{Im}(k) \gg 0$ then the quasi-normal frequencies are the j -th solutions of

$$2k \approx -i\omega_0 e^{-ika}. \quad (\text{A.10})$$

Appendix A.2. Modified Pöschl-Teller potential

The Pöschl-Teller potential is a continuous function over the real line $(-\infty, \infty)$. The QNMs of this potential forms a complete set, in the sense that the solutions at late times can be represented by an infinite sum of the QNMs.³³ However the quantization procedure that we have used requires that spatial discontinuities be present in the potential. We therefore artificially introduce such discontinuities in the potential by making it go to zero for $x > a$ for a large a where the value of the potential is very small.¹³ Therefore the modified Pöschl-Teller potential is

$$V(x) = \begin{cases} \frac{1}{4\nu}(\nu^2 + c^2) \operatorname{sech}^2(\sqrt{\nu}x), & x \in [-a, a] \\ 0, & \text{otherwise.} \end{cases} \quad (\text{A.11})$$

where the factors ν and c can be adjusted to approximate actual black hole potentials,²² subject to the condition $(\nu^2 + c^2) > 0$. Here we have considered the quasi-exactly solvable form of the Pöschl-Teller potential,^{34,35} a special form of the more general Scarf II potential. The Schrödinger - like equation for the Pöschl-Teller potential is

$$\frac{d^2 f(x)}{dx^2} + \omega^2 f(x) = \begin{cases} \frac{1}{4\nu}(\nu^2 + c^2) \operatorname{sech}^2(\sqrt{\nu}x) f(x), & x \in [-a, a] \\ 0, & \text{otherwise.} \end{cases} \quad (\text{A.12})$$

We can approximate the QNM boundary condition as

$$f(x) = (\cosh(\sqrt{\nu}x))^{\nu+ic/2\nu}, \quad x = \pm a.$$

With these boundary conditions, we rewrite the wavefunction as $f = (\cosh(\sqrt{\nu}x)) g(x)$.³⁵ Introducing the new variable $y = \sinh(\sqrt{\nu}x)$ Eq. (A.12) becomes

$$\frac{d^2 g(y)}{dy^2} + \frac{iy(c - 2i\nu)}{(y^2 + 1)\nu} \frac{dg(y)}{dy} - \frac{c^2 - 2i\nu c - \nu(4\omega^2 + \nu)}{4(y^2 + 1)\nu^2} g(y) = 0. \quad (\text{A.13})$$

From Eq. (A.13), using the asymptotic iteration method we get the QNM frequencies as^{35,36}

$$\omega_n = \pm \left(\frac{c}{2\sqrt{\nu}} - i \frac{(2n+1)\sqrt{\nu}}{2} \right) \quad (\text{A.14})$$

and the QNMs as

$$f_n(x) \approx (\cosh(\sqrt{\nu}x))^{\nu+ic/2\nu} P_n^{(\alpha,\beta)}(i \sinh(\sqrt{\nu}x))$$

where $P_n^{(\alpha,\beta)}(x)$ are the Jacobi Polynomials.

Appendix B. Path Integral Quantization

Classical treatments of non-conservative or open system often use the cavity-bath model to describe such systems. The system of interest forms the cavity and the surroundings forms the bath with infinite degrees of freedom. Energy

is exchanged between the cavity and the bath via some interaction. The complete system is described by a Lagrangian of the form $L(q, \dot{q}, Q, \dot{Q}) = L_c(q, \dot{q}) + L_b(Q, \dot{Q}) + L_{int}(q, \dot{q}, Q, \dot{Q})$, where $L_c(q, \dot{q})$ describes the cavity, $L_b(Q, \dot{Q})$ the bath and $L_{int}(q, \dot{q}, Q, \dot{Q})$ gives the interaction between the two.³⁷ The structure of the Lagrangian shows that the quantization would involve both the cavity (q, \dot{q}) and the bath (Q, \dot{Q}) degrees of freedom. However we are not interested in the time evolution of the bath. It is therefore desirable to eliminate Q and \dot{Q} variables and compute the expectation value of any observable in terms of q and \dot{q} only. Feynman and Vernon showed that the path integral formulation is very effective in this regard.³⁸ The general idea is to write down the generating functional or the density matrix of the whole system and then integrate out the bath degrees of freedom. This leaves us a reduced density matrix which incorporates the effect of the bath on the cavity and is expressed in terms of the q and \dot{q} variables only.^{38,39} In the case of a black hole these bath degrees of freedom can be described as the fields populating the space between the black hole event horizon and a stretched horizon.

As mentioned in Sec. (2), QNMs are the exponentially damped eigensolutions of the operator \tilde{D} and form a complete set for the potentials with discontinuities, like those mentioned in Sec. (Appendix A). Moreover the spatial discontinuities in potentials allow us to treat the space around the source of the QNMs as an open system or cavity coupled to a bath, the interval I being the cavity and everything outside I forms the bath; the gravitational waves carry off energy from the cavity to the bath. The QNMs can therefore be used for exact eigenfunction expansions in the cavity which is analogous to the normal mode expansions in Hermitian systems. If we eliminate the bath modes then we can second quantize the open system in terms of the damped QNMs only. In this section we will use the Feynman-Vernon formalism as employed in²⁷ to quantize QNMs of the model potentials.

Appendix B.1. Integrating out the bath modes

To start with, we consider the universe to be restricted to $[-\lambda, \lambda]$ and the cavity to $[-a, a]$. The region $[-\lambda, -a] \cup [a, \lambda]$ forms the thermal bath. To get the equation of motion of the bath variables we divide the region between $[-\lambda, -a]$ into infinite segments as $x_n = x(1+\varepsilon) - nd$ (where $n = 0, 1, 2, \dots, \infty$ and d is the lattice spacing) and set harmonic oscillators q_n at these lattice points. The equation of motion of the j -th oscillator is given by

$$\ddot{q}_j = -\kappa(q_j - q_{j-1}) + \kappa(q_{j+1} - q_j) \quad (\text{B.1})$$

Using the forward and backward differentials

$$\Delta^+ q_j = \frac{q_{j+1} - q_j}{d}, \quad \Delta^- q_j = \frac{q_j - q_{j-1}}{d}$$

and the relation

$$(q_{j-1} - q_j) - (q_j - q_{j+1}) = d(\Delta^+ - \Delta^-) = d^2 \Delta^+ \Delta^- q_j$$

we can write the equation of motion of the j -th oscillator as

$$\ddot{q}_j = \kappa d(\Delta^+ - \Delta^-)q_j = \kappa d^2 \Delta^+ \Delta^- q_j$$

which in the continuum limit ($d \rightarrow 0$ and $kd^2 = 1$) gives

$$\ddot{\phi}(t, x) = \partial_x^2 \phi(t, x). \quad (\text{B.2})$$

The bath variables can therefore be considered as free fields. With this definition of the bath variables we can describe the universe by the Euclidean action

$$S = \frac{1}{2} \int_{\text{periodic}} d\mathbf{x}^2 \left[(\partial_\tau \phi)^2 + (\partial_x \phi)^2 + V(x)\phi^2 \right] \quad (\text{B.3})$$

$$= S_{\text{sys}} + S_{\text{bath}} \quad (\text{B.4})$$

where $V(x) = 0$, for $x \in [-\lambda, -a] \cup [a, \lambda]$. We also consider a source χ that interacts only with the cavity fields via

$$(\phi, \chi) = i \left\{ \int_a^{-a} dx [\phi(\partial_\tau \chi) + (\partial_\tau \phi)\chi] + [\phi(-a)\chi(-a) + \phi(a)\chi(a)] \right\}. \quad (\text{B.5})$$

Therefore the configuration space generating functional of the cavity-bath system is

$$W[\chi] = \frac{1}{Z} \int \mathcal{D}\phi e^{-S(\phi, \partial_\mu \phi) + (\phi, \chi)}. \quad (\text{B.6})$$

The real field ϕ satisfies the boundary conditions $\phi(-\lambda - a, 0) = \phi(a + \lambda, 0) = 0$ and $\phi(x, 0) = \phi(x, \beta)$, with β being the inverse of the Hawking temperature, T^{-1} . We now separate the generating functional into cavity and bath factors, $Z^{-1} \int \mathcal{D}\phi = Z_c^{-1} \int \mathcal{D}\phi_c \times Z_b^{-1} \int \mathcal{D}\phi_b$, the latter running over the fields in the region $[-\lambda - a, -a] \cup [a, a + \lambda]$, with given boundary values $\phi(a, \tau)$ and $\phi(-a, \tau)$. Z^{-1} , Z_b^{-1} and Z_c^{-1} are normalizing factors. The bath factor turns out to be

$$W_b = \frac{1}{Z_b} \int \mathcal{D}\phi_b \exp \left[\frac{1}{2} \int_0^\beta d\tau \int_{-a-\lambda}^{-a} dx \left[(\partial_\tau \phi)^2 + (\partial_x \phi)^2 \right] \right. \\ \left. + \frac{1}{2} \int_0^\beta d\tau \int_a^{a+\lambda} dx \left[(\partial_\tau \phi)^2 + (\partial_x \phi)^2 \right] \right]. \quad (\text{B.7})$$

Let $\xi = x + a$ for the first integral in Eq. (B.7) and $\eta = x - a$ for the second. Fourier-expansion of the bath modes in terms of the Matsubara frequencies²⁷ gives us

$$\phi(\eta, \tau) = \frac{1}{\beta} \sum_m \left[\phi_m(a) \frac{\lambda - \eta}{\lambda} + \sum_{u=1}^{\infty} \phi_{um} \sin \left(\frac{\pi u \eta}{\lambda} \right) \right] e^{-i\nu_k \tau}, \\ \phi(\xi, \tau) = \frac{1}{\beta} \sum_m \left[\phi_m(-a) \frac{\lambda + \xi}{\lambda} + \sum_{u=1}^{\infty} \phi_{um} \sin \left(\frac{\pi u \eta}{-\lambda} \right) \right] e^{-i\nu_k \tau}.$$

$\nu_k = 2\pi k/\beta$ ($k \in \mathbb{Z}$) are the bosonic Matsubara frequencies. Substituting these expansions in Eq. (B.7), we obtain

$$W_b = \exp \left[\frac{1}{2\beta} \sum_m \{ |\phi_m(a)|^2 - |\phi_m(-a)|^2 \} \left(-\frac{\lambda\nu_k^2}{3} - \frac{1}{\lambda} + \sum_{u=1}^{\infty} \frac{2(\lambda\nu_k^2/\pi u)^2}{\lambda\nu_k^2 + \pi^2 u^2/\lambda} \right) \right]$$

which in the limit $\lambda \rightarrow \infty$ gives

$$W_b = \exp \left[\frac{1}{2\beta} \sum_m |\nu_k| (|\phi_m(-a)|^2 - |\phi_m(a)|^2) \right]. \quad (\text{B.8})$$

Therefore the effective generating functional (or reduced density matrix) of the cavity fields is

$$W_c^R[\chi] = \frac{1}{Z_c} \int \mathcal{D}\phi_c \exp \left[(\phi_m, \chi_{-m}) + \frac{1}{2\beta} \sum_m \left[\int_{-a}^a dx [(\partial_\tau \phi_m)^2 + |(\partial_x \phi_m)|^2 + V(x)|\phi_m|^2] + |\nu_k| (|\phi_m(-a)|^2 - |\phi_m(a)|^2) \right] \right]. \quad (\text{B.9})$$

Appendix B.2. QNM expansion and path integral quantization

We expand the cavity and the source fields in terms of the QNMs, $\phi_m(x, 0) = \sum_m a_{jm} f_j$ and $\chi_{-m}(x, 0) = \sum_m b_{j,-m} f_j$. We temporarily assume that $V(x) = 0$, $x \in [-a, a]$. Therefore using the orthogonality condition given by Eq. (7) in Eq. (B.9) we obtain

$$W_c^R[\chi]_0 = \exp \left[\frac{1}{8\beta} \sum_{jkm} \frac{b_{jm} b_{k,-m}}{\omega_j + \omega_k} \left[f_j(-a) f_k(-a) \left(\frac{\theta(m)\omega_k}{i\omega_k + \nu_k} + \frac{\theta(-m)\omega_j}{i\omega_j - \nu_k} \right) + f_j(a) f_k(a) \left(\frac{\theta(m)\omega_j}{i\omega_j + \nu_k} + \frac{\theta(m)\omega_k}{i\omega_k - \nu_k} \right) \right] \right]. \quad (\text{B.10})$$

We calculated this path integral by temporarily assuming that $V(x) = 0$, $x \in [-a, a]$. Now for the actual case of $V(x) \neq 0$, $x \in [-a, a]$ we can calculate the cavity generating functional from Eq. (B.10) by perturbation techniques.²⁷ Let $a_{jm} \rightarrow \beta \partial / 2\omega_j \partial b_{j,-m}$. Then the generating functional for the cavity is

$$W_c^R[\chi] = \frac{1}{Z_c} \exp \left[-\frac{\beta}{2} \sum_{m_1 m_2} \sum_{j_1 j_2} \int_{-a}^a dx V(x) f_{j_1}(x) f_{j_2}(x) \frac{1}{2\omega_{j_1} \omega_{j_2}} \frac{\partial}{\partial b_{j_1 m_1}} \frac{\partial}{\partial b_{j_2 m_2}} \right] W_c^R[\chi]_0 \quad (\text{B.11})$$

where we can substitute $V(x) = V_0$ for the rectangular barrier and $V(x) = (1/4\nu)(\nu^2 + c^2) \text{sech}^2(\sqrt{\nu}x)$ for the modified Pöschl-Teller potential. Therefore Eq.

(B.11) gives the generating functional of the cavity in terms of the QNMs of the system only. As we have calculated the QNMs for our model potentials in Sec. (Appendix A), we can just replace them in Eq. (B.11) to obtain the final algebraic forms of the generating functional.

Appendix C. Dissipative oscillator models

The Bateman-Feshbach-Tikochinsky (BFT) oscillator has been the inspiration behind our quantization process. However there's another oscillator system, the Caldirola-Kanai(CK) oscillator, which could have been used to model the quasi-normal modes. Here, we show that the BFT and the CK oscillators are equivalent.

Appendix C.1. Bateman-Feshbach-Tikochinsky System

Just like in the case of curved spacetime, a free scalar field in a flat spacetime is also governed by the Klein-Gordon equation

$$[\square + m^2] \phi(\mathbf{x}, t) = 0 \quad (\text{C.1})$$

with m being the mass of the field. Fourier transforming Eq. (C.1), in the scalar variables, we obtain

$$\ddot{y}_{\mathbf{k}}(t) + (k^2 + m^2)y_{\mathbf{k}}(t) = 0 \quad (\text{C.2})$$

where

$$y_{\mathbf{k}}(t) \equiv \int \frac{d^n \mathbf{x}}{(2\pi)^{3/2}} e^{-i\mathbf{k}\cdot\mathbf{x}} \phi(\mathbf{x}, t).$$

Eq. (C.2) is the equation of motion of a simple harmonic oscillator with the frequency being $\omega_k \equiv \sqrt{k^2 + m^2}$. The scalar field can therefore be considered as a collection of an infinite number of simple harmonic oscillators with densely spaced frequencies. We now ask the question ‘‘What happens to the field if we include damping?’’, i.e., what would be nature of the field decomposed in terms of oscillators instead of satisfying Eq. (C.2):

$$\ddot{y}_{\mathbf{k}}(t) + \omega_k^2 y_{\mathbf{k}} + \alpha \dot{y}_{\mathbf{k}} = 0$$

where α is a spacetime resistance constant. To answer this question we start with a form invariant Lagrangian density. Using the Dirac matrices we construct a Lagrangian of the form

$$\mathcal{L} = \frac{1}{2} (\partial_\mu \Phi^D \partial^\mu \Phi + \partial^\mu \Phi^D \partial_\mu \Phi) + \alpha \Phi^D \gamma^\mu \partial_\mu \Phi \quad (\text{C.3})$$

where $\Phi^D = \Phi^\dagger \gamma^0$ and $\Phi = (\phi_1, \phi_2, \dots, \phi_n)^T$ is a n -component spinor object. The γ^μ 's are the Dirac matrices in n -dimensions. The Euler-Lagrange equation is

$$\partial^\mu \partial_\mu \Phi - \alpha \gamma^\mu \partial_\mu \Phi = 0. \quad (\text{C.4})$$

For simplicity we work in 1 + 1 dimensions, where the Euler-Lagrange equation reduces to a pair of coupled equations

$$\begin{aligned}\partial_x^2 \phi_1 - \partial_t^2 \phi_1 - \alpha[\partial_t \phi_2 + \partial_x \phi_1] &= 0 \\ \partial_x^2 \phi_2 - \partial_t^2 \phi_2 - \alpha[\partial_t \phi_1 - \partial_x \phi_2] &= 0.\end{aligned}\tag{C.5}$$

This is because the two dimensional Dirac matrices are

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \gamma^1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

If we naively assume a plane-wave like solution of the form $e^{i\omega x}(y_1(t), y_2(t))^T$, because of the first order position derivatives in the equation, we get a system of equations with complex parameters. But we can avoid this, thanks to the real 2×2 matrix representations of $\sqrt{-1}$, like for example $\sigma = \gamma^0 \gamma^1$. Putting $\Phi = e^{\sigma \omega x}(y_1(t), y_2(t))^T$ in Eq. (C.4) we obtain a coupled system of equations,

$$\begin{aligned}\ddot{y}_1 + \omega^2 y_1 + \alpha \dot{y}_2 - \alpha y_1 &= 0 \\ \ddot{y}_2 + \omega^2 y_2 + \alpha \dot{y}_1 - \alpha y_2 &= 0,\end{aligned}\tag{C.6}$$

which we can easily decouple using $y_+ = (y_1 + y_2)/2$ and $y_- = (y_1 - y_2)/2$. Thus, we obtain

$$\begin{aligned}\ddot{y}_+ + (\omega^2 - \alpha\omega)y_+ + \alpha \dot{y}_+ &= 0 \\ \ddot{y}_- + (\omega^2 - \alpha\omega)y_- - \alpha \dot{y}_- &= 0.\end{aligned}\tag{C.7}$$

For $\omega > \alpha$, eqs. (C.7) represent the Bateman dual system. Therefore we can consider the fields ϕ_1 and ϕ_2 to be infinite collections of coupled oscillators - one of which is dissipative and the other is amplified.

Appendix C.2. Caldirola-Kanai system from Galley's approach

Recently, Galley has come up with a new formulation for the nonconservative systems.¹⁸ One of the crucial feature of Galley's approach is the formal doubling of the variables. While this formal doubling of variables has been the feature for understanding dissipative oscillator, however, there is no physical meaning associated to these extra variables. Here we show that using the field theoretic concept it is possible to physically identify the doubling of modes in any dissipative system.

Consider a simple harmonic oscillator with variable coordinate, $q(t)$ and natural frequency, ω , coupled to N number of simple harmonic oscillators, $Q(\omega_n)$. The equations of motion are

$$M_n \ddot{Q}(\omega_n) + M_n \omega_n^2 Q(\omega_n) = \lambda(\omega_n) q \tag{C.8}$$

$$m \ddot{q} + m \omega^2 q = \sum_{n=1}^N \lambda(\omega_n) Q(\omega_n). \tag{C.9}$$

We double the degrees of freedom, $q \rightarrow (q_1, q_2)$ and $Q(\omega_n) \rightarrow (Q_1(\omega_n), Q_2(\omega_n))$ and define new coordinates

$$q_{\pm} = \frac{q_1 \pm q_2}{\sqrt{2}}, \quad Q_{\pm}(\omega_n) = \frac{Q_1(\omega_n) \pm Q_2(\omega_n)}{\sqrt{2}}.$$

Using the retarded and advanced Green's functions we write the equations of motion in terms of these new coordinates

$$Q_+(\omega_n, t) = Q^h(\omega_n, t) + \frac{\lambda(\omega_n)}{M_n} \int_{t_i}^{t_f} dt' G_{r,n}(t-t') q_+(t') \quad (\text{C.10})$$

$$Q_-(\omega_n, t) = \frac{\lambda(\omega_n)}{M_n} \int_{t_i}^{t_f} dt' G_{a,n}(t-t') q_-(t') \quad (\text{C.11})$$

where $G_{r,n}(G_{a,n})$ is the retarded(advanced)Green's function of the n^{th} oscillator and $Q^h(\omega_n)$ is a solution of the homogeneous equation. Using these solutions we can write the effective Lagrangian for the q -system as

$$L = m\dot{q}_+\dot{q}_- - m\omega^2 q_+ q_- + K(q_{\pm}, \dot{q}_{\pm}, t) \quad (\text{C.12})$$

where $K(q_{\pm}, \dot{q}_{\pm}, t) = q_- \sum_{n=1}^N \lambda(\omega_n) Q^h(\omega_n, t) + \int_{t_i}^{t_f} q_-(t) G(t-t') q_+(t') dt'$ and $G(t-t') = \sum_{n=1}^N (\lambda_n^2)/(M_n \omega_n) \sin \omega_n(t-t')$. Let $Q^h(\omega_n, t) = 0$, $M_n = M$ and $\lambda_n = \lambda(\omega_n) = \lambda \omega_n$. Then in the limit $N \rightarrow \infty$ and assuming that the frequencies are dense in R_+ we get

$$G(t-t') = -\frac{\lambda^2}{M} \sum_n \frac{d}{dt'} \cos \omega_n(t-t') = -\alpha \frac{d}{dt'} \delta(t-t')$$

where α is a constant. Therefore the effective Lagrangian becomes

$$L = m\dot{q}_+\dot{q}_- - m\omega^2 q_+ q_- - \alpha \dot{q}_+ q_-. \quad (\text{C.13})$$

We define $x = (q_+ - e^{\alpha t} q_-)/\sqrt{2}$ and $y = (q_- + e^{-\alpha t} q_+)/\sqrt{2}$. In terms of these new coordinates the effective Lagrangian becomes

$$L = m \frac{e^{\alpha t}}{2} (\dot{x}^2 - \omega^2 x^2) - m \frac{e^{-\alpha t}}{2} (\dot{y}^2 - \omega^2 y^2) + m \frac{d}{dt} \left(\frac{\lambda}{2} y^2 e^{-\alpha t} \right). \quad (\text{C.14})$$

The Lagrangian in Eq. (C.14) describes a dual-Caldirola-Kanai system. Therefore we see that the harmonic oscillator coupled to a bath of infinite number of oscillators (Eqs. (C.8) and (C.9)), the Bateman dual system (Eq. (C.12) and the Caldirola-Kanai system are all intimately connected.

References

1. K. D. Kokkotas and B. Schmidt, *Quasi-normal modes of stars and black holes*, *Living Reviews in Relativity* **2** (1999), no. 2.
2. E. Berti, V. Cardoso, and A. O. Starinets, *Quasinormal modes of black holes and black branes*, *Classical and Quantum Gravity* **26** (2009) 163001, [[arXiv:0905.2975](https://arxiv.org/abs/0905.2975)].
3. R. A. Konoplya and A. Zhidenko, *Quasinormal modes of black holes: From astrophysics to string theory*, *Rev. Mod. Phys.* **83** (Jul, 2011) 793–836.

4. R. H. Price and V. Husain, *Model for the completeness of quasi-normal modes of relativistic stellar oscillations*, *Phys. Rev. Lett.* **68** (Mar, 1992) 1973–1976.
5. S. Kalyana Rama and B. Sathiapalan, *On the role of chaos in the ads/cft connection*, *Mod. Phys. Lett.* **A14** (1999) 2635–2648, [[hep-th/9905219](#)].
6. G. T. Horowitz and V. E. Hubeny, *Quasinormal modes of ads black holes and the approach to thermal equilibrium*, *Phys. Rev. D* **62** (2000) 024027, [[hep-th/9909056](#)].
7. S. Hod, *Bohr’s correspondence principle and the area spectrum of quantum black holes*, *Phys. Rev. Lett.* **81** (1998) 4293, [[gr-qc/9812002](#)].
8. O. Dreyer, *Quasinormal modes, the area spectrum, and black hole entropy*, *Phys. Rev. Lett.* **90** (2003) 081301, [[gr-qc/0211076](#)].
9. J. S. F. Chan and R. B. Mann, *Scalar wave falloff in asymptotically antide sitter backgrounds*, *Phys. Rev. D* **55** (Jun, 1997) 7546–7562.
10. M. Maggiore, *The Physical interpretation of the spectrum of black hole quasi-normal modes*, *Phys. Rev. Lett.* **100** (2008) 141301, [[arXiv:0711.3145](#)].
11. J. Skakala and S. Shankaranarayanan, *Horizon spectroscopy in and beyond general relativity*, [arXiv:1311.4255](#).
12. S. P. Kim, *Quasinormal modes of black holes and dissipative open systems*, *J.Korean Phys.Soc.* **49** (2006) 764–772, [[gr-qc/0512005](#)].
13. E. S. C. Ching, P. T. Leung, W. M. Suen, and K. Young, *Wave propagation in gravitational systems: Completeness of quasi-normal modes*, *Phys. Rev. D* **54** (Sep, 1996) 3778–3791.
14. J. R. Ray, *Lagrangians and systems they describe-how not to treat dissipation in quantum mechanics*, *Am. J. Phys.* **47** (July, 1979) 626–629.
15. C.-I. Um, K.-H. Yeon, and T. F. George, *The quantum damped harmonic oscillator*, *Phys. Rept.* **362** (May, 2002) 63–192.
16. D. M. Gitman and V. G. Kupriyanov, *Canonical quantization of so-called non-Lagrangian systems*, *Eur. Phys. J. C* **50** (Apr., 2007) 691–700, [[hep-th/06](#)].
17. M. C. Baldiotti, R. Fresneda, and D. M. Gitman, *Quantization of the damped harmonic oscillator revisited*, *Phy. Lett. A* **375** (Apr., 2011) 1630–1636, [[arXiv:1005.4096](#)].
18. C. R. Galley, *Classical mechanics of nonconservative systems*, *Phys. Rev. Lett.* **110** (Apr, 2013) 174301.
19. G. ’t Hooft, *On the quantum structure of a black hole*, *Nuclear Physics B* **256** (1985), no. 0 727 – 745.
20. A. Ishkhanyan and K.-A. Suominen, *LETTER TO THE EDITOR: New solutions of Heun’s general equation*, *J. Phys. A.* **36** (Feb., 2003) L81–L85, [[arXiv:0909.1684](#)].
21. P. P. Fiziev, *Exact solutions of Regge-Wheeler equation*, *J. Phys. Conf. Ser.* **66** (May, 2007) 012016, [[gr-qc/070](#)].
22. V. Ferrari and B. Mashhoon, *New approach to the quasi-normal modes of a black hole*, *Phys. Rev. D* **30** (Jul, 1984) 295–304.
23. H. Goldstein, C. Poole, and J. Safko, *Classical Mechanics*. Pearson Education, 2002.
24. B. F. Schutz, *Asteroseismology of neutron stars and black holes*, *Journal of Physics: Conference Series* **118** (2008), no. 1 012005.
25. P. T. Leung, W. M. Suen, C. P. Sun, and K. Young, *Waves in open systems via a biorthogonal basis*, *Phys. Rev. E* **57** (May, 1998) 6101–6104.
26. K. C. Ho, P. T. Leung, A. Maassen van den Brink, and K. Young, *Second quantization of open systems using quasi-normal modes*, *Phys. Rev. E* **58** (Sep, 1998) 2965–2978.
27. A. M. van den Brink, *Exactly solvable path integral for open cavities in terms of quasi-normal modes*, *Phys. Rev. E* **61** (Mar, 2000) 2367–2375.

28. Chandrasekhar, S., *On the equations governing the perturbations of the Schwarzschild black hole*, *Proc. R. Soc. London, Ser. A* **343** (1975) 289298.
29. P. T. Leung, A. Maassen van den Brink, W. M. Suen, C. W. Wong, and K. Young, *Susy transformations for quasi-normal modes of open systems*, *Journal of Mathematical Physics* **42** (2001), no. 10 4802–4820.
30. P. T. Leung, S. Y. Liu, and K. Young, *Completeness and orthogonality of quasi-normal modes in leaky optical cavities*, *Phys. Rev. A* **49** (Apr, 1994) 3057–3067.
31. Z. Li, G. Yi-Bo, and W. Cheng, *Green function and perturbation method for dissipative systems based on birthogonal basis*, *Communications in Theoretical Physics* **51** (2009), no. 6 1017.
32. P. Boonserm and M. Visser, *Quasi-normal frequencies: key analytic results*, *Journal of High Energy Physics* **2011** (2011), no. 3 1–28.
33. H. R. Beyer, *On the completeness of the quasi-normal modes of the poschl-teller potential*, *Commun. Math. Phys.* **204** (1999) 397–423.
34. H.-T. Cho and C.-L. Ho, *Quasi-exactly solvable quasi-normal modes*, *Journal of Physics A: Mathematical and Theoretical* **40** (2007), no. 6 1325.
35. O. Özer and P. Roy, *The asymptotic iteration method applied to certain quasi-normal modes and non hermitian systems*, *Central European Journal of Physics* **7** (2009), no. 4 747–752.
36. H. Cho, A. Cornell, J. Doukas, T. Huang, and W. Naylor, *A New Approach to Black Hole Quasinormal Modes: A Review of the Asymptotic Iteration Method*, *Adv.Math.Phys.* **2012** (2012) 281705, [arXiv:1111.5024].
37. A. Caldeira and A. Leggett, *Quantum tunnelling in a dissipative system*, *Annals of Physics* **149** (1983), no. 2 374 – 456.
38. R. Feynman and F. V. Jr., *The theory of a general quantum system interacting with a linear dissipative system*, *Annals of Physics* **24** (1963), no. 0 118 – 173.
39. U. Weiss, *Quantum Dissipative Systems*. Series in modern condensed matter physics. World Scientific, 1999.
40. I. R. Amado, *Applications of Holography to Strongly Coupled Hydrodynamics*. Phd thesis, Universidad Autónoma de Madrid, 2010.
41. S. Iso and S. Okazawa, *Stochastic equations in black hole backgrounds and non-equilibrium fluctuation theorems*, *Nuclear Physics B* **851** (2011), no. 2 380 – 419.
42. T. R. Govindarajan and V. Suneeta, *Quasi-normal modes of ads black holes: a superpotential approach*, *Classical and Quantum Gravity* **18** (2001), no. 2 265.
43. V. Cardoso and J. P. S. Lemos, *Scalar, electromagnetic, and weyl perturbations of btz black holes: Quasinormal modes*, *Phys. Rev. D* **63** (May, 2001) 124015.
44. D. Birmingham, *Choptuik scaling and quasi-normal modes in the anti-desitter/conformal field theory correspondence*, *Phys. Rev. D* **64** (Aug, 2001) 064024.
45. I. Sachs and S. N. Solodukhin, *Quasi-normal modes in topologically massive gravity*, *Journal of High Energy Physics* **2008** (2008), no. 08 003.
46. Y. Kwon, S. Nam, J.-D. Park, and S.-H. Yi, *Quasi-normal modes for new type black holes in new massive gravity*, *Classical and Quantum Gravity* **28** (2011), no. 14 145006.
47. Y. S. Myung, Y.-W. Kim, T. Moon, and Y.-J. Park, *Classical stability of btz black holes in new massive gravity*, *Phys. Rev. D* **84** (Jul, 2011) 024044.
48. H.-P. Nollert, *Quasinormal modes of schwarzschild black holes: The determination of quasi-normal frequencies with very large imaginary parts*, *Phys. Rev. D* **47** (Jun, 1993) 5253–5258.
49. J. Skakala and M. Visser, *Highly damped quasi-normal frequencies for piecewise*

- eckart potentials*, *Phys. Rev. D* **81** (Jun, 2010) 125023.
50. H. T. Cho, A. S. Cornell, J. Doukas, and W. Naylor, *Black hole quasi-normal modes using the asymptotic iteration method*, *Classical and Quantum Gravity* **27** (2010), no. 15 155004.
 51. D. Birmingham, S. Carlip, and Y.-j. Chen, *Quasinormal modes and black hole quantum mechanics in (2+1)-dimensions*, *Classical and Quantum Gravity* **20** (2003) L239–L244, [[hep-th/0305113](#)].
 52. H. Dekker, *A note on the exact solution of the dynamics of an oscillator coupled to a finitely extended one-dimensional mechanical field and the ensuing quantum mechanical ultraviolet divergence*, *Physics Letters A* **104** (1984), no. 2 72 – 76.
 53. P. Ullersma, *An exactly solvable model for brownian motion: I. derivation of the langevin equation*, *Physica* **32** (1966), no. 1 27 – 55.
 54. L. H. Yu and C.-P. Sun, *Evolution of the wave function in a dissipative system*, *Phys. Rev. A* **49** (Jan, 1994) 592–595.
 55. H. Grabert, U. Weiss, and P. Talkner, *Quantum theory of the damped harmonic oscillator*, *Zeitschrift fr Physik B Condensed Matter* **55** (1984), no. 1 87–94.
 56. R. W. F. van der Plank and L. G. Suttorp, *Generalization of damping theory for cavities with mirrors of finite transmittivity*, *Phys. Rev. A* **53** (Mar, 1996) 1791–1800.
 57. H. Lai, P. Leung, and K. Young, *Thermal spectrum in leaky cavities: A string model*, *Physics Letters A* **119** (1987), no. 7 337 – 339.
 58. A. M. van den Brink and K. Young, *Jordan blocks and generalized bi-orthogonal bases: realizations in open wave systems*, *Journal of Physics A: Mathematical and General* **34** (2001), no. 12 2607.
 59. P. T. Leung and K. Young, *Time-independent perturbation theory for quasi-normal-mode solutions in quantum mechanics*, *Phys. Rev. A* **44** (Sep, 1991) 3152–3161.